

PRESCRIBING INFORMATION

Pr **SANDOZ MINOCYCLINE**

Minocycline Hydrochloride Capsules USP

50 mg and 100 mg minocycline

Antibiotic

USP

Sandoz Canada Inc.
110, de Lauzon Street
Boucherville, QC
J4B 1E6

Date of Revision
April 29, 2019

Submission Control Number: 226984

COMPLETE PRESCRIBING INFORMATION

Pr Sandoz Minocycline
minocycline hydrochloride capsules USP
50 mg and 100 mg minocycline

THERAPEUTIC CLASSIFICATION

Antibiotic

ACTION AND CLINICAL PHARMACOLOGY

Sandoz Minocycline (minocycline hydrochloride) is a tetracycline with antibacterial activity against some Gram-negative and Gram-positive organisms. The action of minocycline is primarily bacteriostatic and is thought to exert its antimicrobial effect by the inhibition of protein synthesis.

Comparative Bioavailability Studies

A comparative two-way cross-over bioavailability study between Sandoz Minocycline 100 mg capsules (Sandoz Canada Inc.) and Minocin 100 mg capsules (GlaxoSmithKline Inc.) was conducted, under fasting conditions, with eighteen (18) healthy male subjects who completed the study. The following results were obtained:

Minocycline (2 x 100 mg) capsules From measured data Oral administration Geometric Mean Arithmetic Mean (CV %)				
---	--	--	--	--

Parameter	Test*	Reference†	% Ratio of Geometric Means	Confidence Intervals (90%)
AUC _T ‡ (mcg·h/mL)	56.09 57.04 (-18.7)	55.54 56.88 (-20.1)	101.0	(91.9-111.0)
AUC _I (mcg·h/mL)	59.89 61.00 (-19.3)	59.31 60.76 (-20)	101.0	(91.6-111.3)
C _{MAX} (mcg·h/mL)	3.67 3.76 (-22.4)	3.75 3.88 (-25)	97.9	(88.1-108.7)
T _{MAX} § (h)	1.89 (0.58)	2.03 (-0.4)		
T _{1/2} ² (h)	15.8 (-3.24)	16.3 (-2.18)		

* Sandoz Minocycline, Sandoz Canada Inc.

† Minocin, 100 mg capsules, GlaxoSmithKline Inc., formerly known as, Lederle-Cyanamid Canada Inc.

§.2 For the T_{max} and T_{1/2el} parameters these are the arithmetic means with standard deviation in parenthesis.

INDICATIONS AND CLINICAL USE

Sandoz Minocycline (minocycline hydrochloride) may be indicated for the treatment of the following infections due to susceptible strains of the designated organisms:

Gall bladder infections caused by *Escherichia coli*.

Urinary Tract Infections: cystitis, gonorrhea, pyelonephritis caused by *Escherichia coli*, *Proteus* species, *Klebsiella* species, *Enterobacter aerogenes*, *Neisseria gonorrhoeae*.

When penicillin is contraindicated, Sandoz Minocycline may be employed as an alternative drug in the treatment of anal and pharyngeal gonorrhea and syphilis.

Skin and Soft Tissue Infections : abscess, cellulitis, furunculosis, impetigo and pyoderma caused by: *Staphylococcus epidermidis*, *Staphylococcus aureus*, *Streptococcus pyogenes*, *Proteus* species, *Escherichia coli*. Although tetracyclines are not the drugs of choice in any staphylococcal or streptococcal infection, Sandoz Minocycline could be useful in circumstances where these organisms are shown to be resistant to other agents but sensitive to minocycline hydrochloride. Bacterial evaluation of clinical cases involving *Proteus* suggests a relatively lower success rate may be expected where these organisms are concerned.

Respiratory Tract Infections : bronchitis, pharyngitis, pneumonia, bronchopneumonia, sinusitis and tonsillitis caused by: *Haemophilus influenzae*, *Klebsiella* species, *Enterobacter* species. Tetracyclines should not be prescribed for acute throat infections.

To reduce the development of drug-resistant bacteria and maintain the effectiveness of Sandoz Minocycline and other antibacterial drugs, Sandoz Minocycline should be used only to treat infections that are proven or strongly suspected to be caused by susceptible bacteria. When culture and susceptibility information are available, they should be considered in selecting or modifying antibacterial therapy. In the absence of such data, local epidemiology and susceptibility patterns may contribute to the empiric selection of therapy.

CONTRAINDICATIONS

- Patients who are hypersensitive to this drug or to any ingredient in the formulation. For a complete listing, see the Dosage Forms section.
- History of hypersensitivity to minocycline or any other tetracycline.
- Pregnancy and lactation (see WARNINGS, Pregnancy and Lactation)
- Children under 13 years (see WARNINGS, Newborns, Infants and Children)
- Complete renal failure
- Severe liver disease
- Myasthenia gravis

WARNINGS

Anaphylactic/Anaphylactoid Reactions

Rarely, anaphylactic/anaphylactoid reactions including shock and fatalities have been associated with the administration of minocycline hydrochloride.

Gastrointestinal

Clostridium difficile-associated disease: *Clostridium difficile*-associated disease (CDAD) has been reported with the use of many antibacterial agents, including minocycline (see ADVERSE REACTIONS). CDAD may range in severity from mild diarrhea to fatal colitis. It is important to consider this diagnosis in patients who present with diarrhea or symptoms of colitis, pseudomembranous colitis, toxic megacolon, or perforation of the colon subsequent to the administration of any antibacterial agent. CDAD has been reported to occur more than 2 months after the administration of antibacterial agents.

Treatment with antibacterial agents may alter the normal flora of the colon and may permit overgrowth of *Clostridium difficile*. *C. difficile* produces toxins A and B, which contribute to the development of CDAD. CDAD may cause significant morbidity and mortality. CDAD can be refractory to antimicrobial therapy.

If the diagnosis of CDAD is suspected or confirmed, appropriate therapeutic measures should be initiated. Mild cases of CDAD usually respond to discontinuation of antibacterial agents not directed against *Clostridium difficile*. In moderate to severe cases, consideration should be given to management with fluids and electrolytes, protein supplementation, and treatment with an antibacterial agent clinically effective against *Clostridium difficile*. Surgical evaluation should be instituted as clinically indicated since surgical intervention may be required in certain severe cases.

Newborns, Infants and Children

Minocycline hydrochloride is contraindicated in children under 13 years of age (see CONTRAINDICATIONS). The use of tetracyclines including minocycline hydrochloride, during tooth development (last half of pregnancy, infancy and childhood under the age of thirteen years) has been shown to cause permanent tooth discoloration (yellow-grey-brown). This is more common during long-term use, but has been observed following short-term courses. Enamel hypoplasia has also been reported. All tetracyclines including minocycline hydrochloride, administered during the last trimester form a stable calcium complex throughout the human fetal skeleton. A decrease in the fibula growth rate has been observed in premature human infants given oral tetracycline in doses of 25 mg/kg every 6 hours. This appeared to be reversible when the drug was discontinued.

Congenital anomalies including limb reductions have been reported in postmarketing experience.

Pregnancy and Lactation

Tetracyclines, including minocycline hydrochloride, are contraindicated during pregnancy and lactation (see CONTRAINDICATIONS) because of possible adverse effects on developing bones and teeth of the fetus and neonate. Results of animal studies indicate that tetracyclines cross the placenta, are found in fetal tissues and can have toxic effects on the developing fetus (often related to retardation of skeletal development). If minocycline hydrochloride is used during pregnancy or if the patient becomes pregnant while taking this drug, the patient should be apprised of the potential hazard to the fetus.

Evidence of embryotoxicity has also been noted in animals treated early in pregnancy. The safety of minocycline hydrochloride for use during pregnancy has not been established.

Tetracyclines, including minocycline hydrochloride, are excreted in the milk of lactating women; therefore, a decision should be made whether to discontinue breast-feeding or to discontinue minocycline.

Fertility

There are no relevant data available.

Elderly

Clinical studies of minocycline hydrochloride did not include sufficient numbers of subjects aged 65 and over to determine whether they respond differently than younger subjects. Dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal or cardiac function, and of concomitant disease or other drug therapy.

Penicillins

It is advisable to avoid giving minocycline hydrochloride in conjunction with penicillin since some bacteriostatic drugs may interfere with the bactericidal action of penicillin.

Treatment of Streptococcal Infections

Minocycline hydrochloride should not be used for the treatment of streptococcal diseases unless the organism is demonstrated to be sensitive, since most streptococci have been found to be resistant to tetracycline drugs. If it is deemed necessary that infection due to Group A beta-hemolytic streptococci be treated with minocycline, then such treatment should be continued for at least ten days.

Renal Impairment

In the presence of significant renal impairment, usual oral doses may lead to excessive systemic accumulations of minocycline hydrochloride and possible liver toxicity. Under such conditions, lower than usual doses may be indicated. After initial therapy, and if therapy is prolonged, serum level determinations of the drug are advisable.

The anti-anabolic action of tetracyclines can also produce dose-related increases in BUN, consequently, in patients with significant renal impairment, elevated serum minocycline hydrochloride levels can lead to azotemia, hypophosphatemia and acidosis.

Renal failure, including interstitial nephritis, has been reported rarely.

Auto-immune Disorders

Rare cases of auto-immune hepatotoxicity and isolated cases of systemic lupus erythematosus (SLE) have been reported (see ADVERSE REACTIONS). Also, minocycline hydrochloride is capable of aggravating the symptoms associated with lupus erythematosus. Therefore, caution should be taken when administering the drug to patients with this disease. If patients develop signs or symptoms of SLE or hepatotoxicity, or suffer exacerbation or pre-existing SLE, minocycline should be discontinued.

Anticoagulants

Minocycline hydrochloride has been shown to depress plasma prothrombin activity. Therefore, patients who are on anticoagulant therapy should be monitored regularly and may require downward adjustment of their anticoagulant dosage. Interference with vitamin K synthesis by microorganisms in the gut has been reported.

Myasthenia Gravis

Minocycline hydrochloride is contraindicated in patients with myasthenia gravis as tetracyclines can cause weak neuromuscular blockade (see CONTRAINDICATIONS).

Cross-sensitivities

Cross-resistance between tetracyclines may develop in micro-organisms and cross-sensitization among the various tetracyclines is extremely common. Minocycline hydrochloride should be discontinued if there are signs/symptoms of overgrowth of resistant organisms, enteritis, glossitis, stomatitis, vaginitis, pruritus ani or staphylococcal enteritis (see ADVERSE REACTIONS).

Hyperpigmentation

As with other tetracyclines, minocycline hydrochloride may cause hyperpigmentation at various body sites (see ADVERSE REACTIONS), including the skin, nails, teeth, oral mucosa, bones, thyroid, eyes (including sclera and conjunctiva), breast milk, lacrimal secretions and perspiration. The black/blue/grey or muddy-brown discolouration may be localized or diffuse. The most frequently reported site is in the skin (see ADVERSE REACTIONS).

Hyperpigmentation may present regardless of dose or duration of therapy but develops more commonly during long term treatment. Pigmentation is often reversible on discontinuation of the drug, although it may take several months or may persist in some cases. The generalised muddy-brown skin pigmentation may persist, particularly in areas exposed to the sun.

Patients should be advised to report any unusual pigmentation without delay and minocycline hydrochloride should be discontinued.

Oral Contraceptives

Reduced efficacy and increased incidence of breakthrough bleeding has been suggested with concomitant use of tetracycline and oral contraceptive preparations.

Patients taking oral contraceptives should be warned that if diarrhea or breakthrough bleeding occur there is a possibility of contraceptive failure.

Susceptibility/Resistance

Development of Drug Resistant Bacteria

Prescribing Sandoz Minocycline in the absence of a proven or strongly suspected bacterial infection is unlikely to provide benefit to the patient and risks the development of drug-resistant bacteria.

PRECAUTIONS

Children

The administration of minocycline hydrochloride to children under 13 years of age is contraindicated.

Skin and Subcutaneous Tissue Disorders

Very rare, serious events have occurred with minocycline hydrochloride including Stevens-Johnson Syndrome and toxic epidermal necrolysis. Minocycline hydrochloride should be discontinued if either of these serious skin reactions is suspected.

Intracranial Hypertension

Bulging fontanelles have been reported in young infants following full therapeutic dosage of tetracyclines including minocycline hydrochloride. Pseudotumor cerebri (benign intracranial hypertension) has been reported in juveniles and adults (see ADVERSE REACTIONS). The clinical manifestations were headache and visual disturbances including blurring of vision, scotoma and diplopia. While these conditions and related symptoms usually resolved after discontinuation of the tetracycline, permanent vision loss has been reported. Treatment should cease if evidence of raised intracranial pressure develops.

Photosensitivity

Patients should be warned to avoid exposure to direct sunlight and/or ultraviolet light while under treatment with minocycline hydrochloride or other tetracycline drugs, and treatment should be discontinued at the first evidence of skin erythema or discomfort. Photosensitivity manifested by an exaggerated sunburn reaction has been observed in some individuals taking tetracyclines. Studies to date indicate that photosensitivity is rarely reported with minocycline hydrochloride.

Ability to Perform Tasks that Require Judgement, Motor, or Cognitive Skills

Patients treated with minocycline hydrochloride may suffer from headaches, light-headedness, dizziness, tinnitus or vertigo (more common in women). Decreased hearing has been rarely reported in patients on minocycline hydrochloride. Administration of minocycline hydrochloride in excess of the recommended dosage can increase the frequency and severity of these CNS symptoms. Patients should be cautioned about driving vehicles or using hazardous machinery while on minocycline hydrochloride therapy. These symptoms may disappear during therapy and usually disappear rapidly when the drug is discontinued.

Overgrowth of Non-susceptible Organisms

As with other antibiotics, minocycline hydrochloride therapy may result in overgrowth of non-susceptible organisms (including fungi). If super infection occurs, minocycline hydrochloride should be discontinued and appropriate therapy instituted.

Cross-sensitivities

The development of cross-resistance to many antibiotics can develop rapidly in several species of micro-organisms. The clinician should bear this in mind if therapy with minocycline hydrochloride is not achieving expected results.

The frequency of resistance to minocycline hydrochloride in hemolytic streptococci is highest in strains from infections of the ear, wounds and skin. Culture and sensitivity studies should be performed whenever feasible and routinely in suspected streptococcal infections. Since sensitivity reactions are more likely to occur in persons with a history of allergy, asthma, hay fever, or urticaria, minocycline hydrochloride should be used with caution in such individuals.

Treatment of Gonorrhea

Before treating patients with gonorrhea, a darkfield examination should be made from any lesion suggestive of concurrent syphilis. Serological tests for syphilis should be repeated monthly for at least 4 months.

Hepatic Dysfunction

Hepatotoxicity has been reported with minocycline hydrochloride; therefore, minocycline hydrochloride should be used with caution in patients with mild to moderate hepatic dysfunction and in conjunction with alcohol or other hepatotoxic drugs.

Laboratory Monitoring

Periodic laboratory evaluation of organ system functions, including hematopoietic, renal and hepatic, should be performed.

Diuretics

Diuretics may aggravate nephrotoxicity by volume depletion.

Drugs Impairing Minocycline Absorption

Absorption of minocycline hydrochloride is impaired by antacids containing aluminum, calcium or magnesium, and oral iron preparations as well as bismuth and zinc salts - interactions with

specific salts and antacids, bismuth containing ulcer-healing drugs, quinapril which contains a magnesium carbonate excipient. These should not be given to patients taking oral minocycline hydrochloride.

Food Interactions

Food and/or milk reduce the absorption of tetracycline. Minocycline hydrochloride is not affected to the same extent.

In a study by Leyden, the absorption of a single 100 mg dose of minocycline was inhibited by the ingestion of solid food by 13% (as measured by a reduction in mean serum concentration), and the absorption of a single 250 mg dose of tetracycline was inhibited by 46% when that antibiotic was administered with solid food. When administered with milk, the mean serum concentration of minocycline was reduced by 27% and that of tetracycline, by 65%. The clinical significance of such declines in serum levels is not known.

Laboratory Tests

Interference with laboratory and other diagnostic tests: False evaluations of urinary catecholamine levels may occur due to interference with the fluorescence test.

Oral Contraceptives

The concurrent use of tetracyclines with oral contraceptives may render oral contraceptives less effective.

Retinoids

Administration of isotretinoin or other systemic retinoids or retinol should be avoided shortly before, during, and shortly after minocycline therapy. Each of these agents used alone has been associated with pseudotumor cerebri (benign intracranial hypertension).

Ergot Alkaloids

Increased risk of ergotism when ergot alkaloids or their derivatives are given with tetracyclines.

Adverse Reactions – Syndromes

The following syndromes have been reported. In some cases involving these syndromes, death has been reported (see ADVERSE REACTIONS). As with other serious adverse reactions, if any of these syndromes are recognized, the drug should be discontinued immediately:

- Hypersensitivity syndrome consisting of cutaneous reaction (such as rash or exfoliative dermatitis), eosinophilia, and one or more of the following: hepatitis, pneumonitis, nephritis, myocarditis, pericarditis. Fever and lymphadenopathy may be present.
- Lupus-like syndrome consisting of positive antinuclear antibody; arthralgia, arthritis, joint stiffness, or joint swelling; and one or more of the following: fever, myalgia, hepatitis, rash, vasculitis.
- Serum sickness-like syndrome consisting of fever; urticaria or rash; and arthralgia, arthritis, joint stiffness, or joint swelling. Eosinophilia may be present.

ADVERSE REACTIONS

The following adverse reactions have been reported with the tetracycline analogues including minocycline hydrochloride :

- a) **Central Nervous System:** increased intracranial pressure, light-headedness, dizziness or vertigo and, rarely, fainting spells have been reported with a variable but overall incidence of approximately 7% in patients treated with minocycline hydrochloride. These symptoms usually disappear rapidly when the drug is discontinued. Impaired hearing, tinnitus, headache, convulsions, sedation, hypesthesia or paresthesia have also been reported.
- b) **Gastrointestinal System:** anorexia, nausea, vomiting, diarrhea, stomatitis, glossitis, enterocolitis, pancreatitis, pruritis ani, constipation, dyspepsia, dysphagia, inflammatory lesions (with monilial overgrowth) in the anogenital region, increases in liver enzymes, and rarely hepatitis and acute liver failure have been reported. Rare instances of esophagitis and esophageal ulcerations have been reported in patients taking the tetracycline-class antibiotics in capsule and tablet form. Most of these patients took the medication immediately before going to bed. Very rare incidence of pseudomembranous colitis has been reported.
- c) **Teeth and Bone :** dental staining (yellow-gray-brown) has been reported in children of mothers given tetracyclines, including minocycline hydrochloride, during the latter half of pregnancy, and in children given the drug during the neonatal period, infancy and childhood to age of 13 years. Enamel hypoplasia has also been reported. Discolouration of bones and teeth has been documented to occur rarely in adolescents and adults upon extended treatment with minocycline hydrochloride. The effects may be irreversible. At present, the mechanism of staining, although not completely elucidated, appears to be mediated by the formation of a stable iron complex. Very rarely arthritis, joint stiffness and joint swelling have been reported.
- d) **Renal:** rise in BUN has been reported and is apparently dose-related. Increased excretion of nitrogen and sodium has also been reported. Acute renal failure, including interstitial nephritis has been reported rarely.
- e) **Skin:** maculopapular and erythematous rashes. Rarely reported - alopecia, fixed drug eruption, photosensitivity, pruritus, rash, urticaria, onycholysis, discolouration of the nails, tongue, gum and lip, pigmentation of the skin and mucous membrane, erythema multiforme, erythema nodosum. Lesions occurring on the glans penis have caused balanitis. Very rare, serious events have occurred with minocycline hydrochloride including angioedema, exfoliative dermatitis, hyperpigmentation of nails, Stevens-Johnson Syndrome, vasculitis and toxic epidermal necrolysis. Minocycline hydrochloride should be discontinued if either of these serious skin reactions is suspected.

- f) **Hypersensitivity Reactions:** urticaria, angioneurotic edema, polyarthralgia, anaphylaxis/anaphylactoid reactions (including shocks and fatalities), hypersensitivity, anaphylactoid purpura and pericarditis. Myalgia has also been reported.
- g) **Autoimmune:** autoimmune hepatotoxicity, lupus-like syndrome, cases of or exacerbation of systemic lupus erythematosus, and myocarditis.
- h) Pseudotumor cerebri (benign intracranial hypertension) in adults has been associated with the use of tetracyclines. The usual clinical manifestations are headache and blurred vision. Bulging fontanelles have been associated with the use of tetracyclines in infants. While both of these conditions and related symptoms usually resolve soon after discontinuation of the tetracycline, the possibility for permanent sequelae exists.
- i) **Respiratory:** rarely - cough and dyspnea, very rarely - bronchospasm, exacerbation of asthma and pulmonary eosinophilia and undetermined frequency of pneumonitis have been reported.
- j) **Other:** fever, elevated liver enzymes including SGOT or SGPT values, hepatic cholestasis, hepatic failure (including fatalities) hyperbilirubinemia, jaundice, autoimmune hepatitis, hemolytic anemia, leukopenia, neutropenia, thrombocytopenia and eosinophilia and pancytopenia and agranulocytosis. When given over prolonged periods, minocycline hydrochloride, like other tetracyclines, has been reported to produce brown-black microscopic discoloration of the thyroid gland. Very rarely, abnormalities of thyroid function have been reported. If adverse reactions or idiosyncrasy occur, the administration of minocycline hydrochloride should be discontinued and appropriate alternate therapy instituted. Very rare incidence of oral and anogenital candidiasis and vulvovaginitis have also been reported. Very rarely- Discoloration of secretions have been reported.

SYMPTOMS AND TREATMENT OF OVERDOSAGE

For management of a suspected drug overdose, contact your regional Poison Control Centre.

Symptoms and Signs

Dizziness, nausea, vomiting, abdominal pain, intestinal hemorrhage, hypotension, lethargy, coma, acidosis, azotemia without a concomitant rise in creatinine.

Treatment

There is no specific antidote. In cases of overdose, discontinue medication, treat symptomatically and with appropriate supportive measures. Minocycline is not removed in significant quantities by haemodialysis or peritoneal dialysis.

DOSAGE AND ADMINISTRATION

Children 13 years of age or older

The usual dosage of Sandoz Minocycline (minocycline hydrochloride) is 4 mg/kg initially followed by 2 mg/kg every 12 hours. Minocycline hydrochloride is contraindicated in children under 13 years of age (see CONTRAINDICATIONS).

Adults

The usual oral dosage of Sandoz Minocycline is 100 mg or 200 mg initially, followed by 100 mg every 12 hours. Alternatively, if more frequent doses are preferred, two or four 50 mg doses may be given initially, followed by one 50 mg dose every 6 hours. Therapy should be continued for 1 or 2 days beyond the time when characteristic symptoms or fever have subsided.

For treatment of syphilis, Sandoz Minocycline therapy should be administered over a period of 10 or 15 days. Close follow-up, including laboratory tests, is recommended.

Concomitant therapy: Antacids containing aluminum, calcium or magnesium and/or iron preparations impair absorption and should not be given to patients taking Sandoz Minocycline.

Dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or other drug therapy.

PHARMACEUTICAL INFORMATION


Drug Substance

Proper Name: Minocycline hydrochloride, USP

Chemical Name: 2-Naphthacencarboxamide,4,7bis(dimethylamino)-1,4,4a,5,5a,6,11,12a-octahydro-3,10,12,12a-tetrahydroxy-1,11-dioxo-, monohydrochloride, [4S(4 α ,4a α ,5a α ,12a α)]

4,7-Bis(dimethylamino)-1,4,4a,5,5a,6,11,12a-octahydro-3,10,12,12a-tetrahydroxy-1,11-dioxo-2-naphthacencarboxamide monohydrochloride [13614-98-7]

Structural Formula:



Molecular Formula: C₂₃H₂₇N₃O₇·HCl

Molecular Weight: 493.94 g/mol

Description: Minocycline hydrochloride is a yellow crystalline powder which is slightly hygroscopic and slightly sensitive to light and oxidation.

STORAGE AND STABILITY

Store at room temperature and protect from light.

DOSAGE FORMS, COMPOSITION AND PACKAGING

50 mg and 100 mg capsules packaged in bottles of 100 capsules.

Composition

Sandoz Minocycline 50 mg: Each orange hard gelatin capsule contains minocycline hydrochloride equivalent to 50 mg minocycline base. Nonmedicinal ingredients: cornstarch, gelatin, magnesium stearate and colouring agents: sunset yellow FCF and titanium dioxide. Tartrazine-free.

Sandoz Minocycline 100 mg: Each orange-purple hard gelatin capsule contains minocycline hydrochloride equivalent to 100 mg minocycline base. Nonmedicinal ingredients: cornstarch, gelatin, magnesium stearate and colouring agents: erythrosine, patent blue V, sunset yellow FCF, titanium dioxide. Tartrazine-free.

MICROBIOLOGY

This survey of the *in vitro* activity of minocycline against clinical isolates was compiled from data presented in 130 articles published from 1967 to 1980. The MICs of minocycline against clinical isolates representing gram-positive, gram-negative, actinomycetes, acid-fast and anaerobic bacteria and mycoplasma, were recorded and entered into a computer database file. The percent of clinical isolates inhibited at various antibiotic concentrations was determined directly from the total number of isolates tested by a computer-assisted statistical analysis system program.

BACTERIA	No. of Strains Tested	Cumulative Strains Inhibited at the Indicated Concentrations of Minocycline (mg/L)			
		≤1	≤4	≤8	≤16
GRAM-POSITIVE					
<i>Staphylococcus aureus</i>	3301	77	91	96	98
<i>Staphylococcus aureus</i> methicillin resistant	13	38	100		
<i>Staphylococcus aureus</i> penicillin resistant	100	100			
<i>Staphylococcus aureus</i> tetracycline resistant	736	50	75	84	93
<i>Staphylococcus epidermidis</i>	577	89	94	95	98
<i>Staphylococcus epidermidis</i> methicillin resistant	19	21	89	95	95
<i>Staphylococcus</i> species	775	82	89	96	99
<i>Staphylococcus</i> species tetracycline resistant	46	48	100		
<i>Staphylococcus</i> beta-hemolytic	654	73	83	95	99
<i>Streptococcus</i> Enterococcus group	844	18	23	28	46
<i>Streptococcus pneumoniae</i>	508	78	88	96	99
<i>Streptococcus pneumoniae</i> tetracycline resistant	70	27	57	96	100
GRAM-NEGATIVE					
<i>Acinetobacter calcoaceticus</i>	456	95	99	100	
<i>Acinetobacter</i> species	56	96	100		
<i>Bordetella pertussis</i>	23	100			
<i>Brucella</i> species	127	75	100		
<i>Citrobacter</i> species	37	8	81	81	84
<i>Enterobacter aerogenes</i>	130	0	13	35	61

BACTERIA	No. of Strains Tested	Cumulative Strains Inhibited at the Indicated Concentrations of Minocycline (mg/L)			
		≤1	≤4	≤8	≤16
<i>Enterobacter cloacae</i>	131	0	9	18	44
<i>Enterobacter</i> species	310	7	78	91	95
<i>Escherichia coli</i>	1538	33	56	69	78
<i>Haemophilus influenzae</i>	385	62	90	98	100
<i>Haemophilus</i> species	182	89	98	99	100
<i>Klebsiella-Enterobacter</i> group	309	30	48	59	68
<i>Klebsiella pneumoniae</i>	299	2	35	53	69
<i>Klebsiella</i> species	247	7	49	62	74
<i>Legionella pneumophila</i>	21	62	100		
<i>Neisseria gonorrhoeae</i>	1082	97	100		
<i>Neisseria gonorrhoeae</i> Beta-lactamase positive	50	90	100		
<i>Neisseria meningitidis</i>	613	94	100		
<i>Proteus</i> indole positive species	102	1	30	47	61
<i>Proteus mirabilis</i>	382	4	12	32	46
<i>Providencia</i> species	94	1	7	16	28
<i>Pseudomonas aeruginosa</i>	643	7	18	36	58
<i>Pseudomonas cepacia</i>	90	8	19	83	97
<i>Pseudomonas maltophilia</i>	81	89	99	99	99
<i>Pseudomonas pseudomallei</i>	157	10	77	89	9
<i>Pseudomonas</i> species	68	68	90	91	91
<i>Salmonella</i> species	128	2	59	76	80
<i>Salmonella</i> species Tetracycline-resistant	123	0	73	92	100
<i>Serratia</i> species	341	0	23	37	55
<i>Shigella</i> species	90	28	66	80	86
<i>Vibrio cholerae</i> type Eltor	203	61	100		
<i>Vibrio</i> species	367	53	100		
<i>Yersinia</i> species	212	94	100		
ACID-FAST BACTERIA					
<i>Mycobacterium tuberculosis</i>	5	0	0	80	100
<i>Mycobacterium</i> species	90	4	26	71	74
ACTINOMYCETES					
<i>Actinomyces israeli</i>	31	100			
<i>Actinomyces</i> species	110	89	95	100	
<i>Nocardia asteroides</i>	84	1	89	100	
<i>Nocardia</i> species	74	30	91	99	100
MYCOPLASMA					
<i>Mycoplasma pneumoniae</i>	14	100			
<i>Mycoplasma</i> species	223	85	91	92	93
CHLAMYDIA					
<i>Chlamydia trachomatis</i>	3	100			
ANAEROBIC					
<i>Bacteroides fragilis</i>	673	44	80	97	99

BACTERIA	No. of Strains Tested	Cumulative Strains Inhibited at the Indicated Concentrations of Minocycline (mg/L)			
		≤1	≤4	≤8	≤16
<i>Bacteroides</i> species	431	58	77	90	92
<i>Campylobacter fetus</i>	97	90	91	91	91
<i>Clostridium</i> species	297	69	81	91	98
<i>Eubacterium</i> species	144	53	87	99	100
<i>Fusobacterium</i> species	107	66	94	100	
<i>Peptococcus</i> species	375	46	81	97	99
<i>Peptostreptococcus</i> species	242	59	85	99	99
<i>Propionibacterium acnes</i>	102	89	95	100	
<i>Propionibacterium</i> species	70	94	97	99	100
<i>Veillonella</i> species	13	69	92	100	

Susceptibility Testing

Tube-Dilution Testing

Microorganisms may be considered susceptible (likely to respond to minocycline therapy), moderately susceptible (harbouring partial resistance) or resistant (not likely to respond to minocycline therapy) depending on the minimum inhibitory concentration (M.I.C.) as follows:

Minocycline M.I.C. Interpretive Standards (mg/L)

Susceptible	Moderately Susceptible	Resistant
≤4	8	≥16

Acceptable Quality Control Ranges of M.I.C. for Reference Strains

Reference Strain	ATCC NUMBER	mg/L
<i>Staphylococcus aureus</i>	29213	0.12-0.5
<i>Streptococcus faecalis</i>	29212	2.0-8.0
<i>Escherichia coli</i>	25922	0.5-2.0

Plate Testing

If the Kirby-Bauer method of susceptibility testing (using a 30 mcg tetracycline disc) gives a zone of 19 mm or greater, the bacterial strain is considered to be susceptible to any tetracycline. A zone of 14 mm or less is considered resistant.

Zone Diameter Interpretive Standards (30 mcg disc)

Susceptible	Moderately Susceptible	Resistant
≥19 mm	15-18 mm	≤14 mm

For *Staphylococcal* species, minocycline powder may be used for additional susceptibility testing.

Acceptable Quality Control Limits (Zone Diameter) for Disc Susceptibility Testing of Reference Strains

Reference Strain	ATCC Number	mg/L
<i>Escherichia coli</i>	25922	19-25
<i>Staphylococcus aureus</i>	25923	25-30

PHARMACOLOGY

Animal Pharmacology

Blood levels produced following oral dosing of minocycline hydrochloride to various animal species were: 21 mg/L at steady state in monkeys administered 30 mg/kg, and 6.5 mg/L at 3 hours post-dose in rats given a single 25 mg/kg dose. Minocycline hydrochloride was extensively distributed to all tissues examined in ¹⁴C-labelled drug studies in dogs.

Clinical Pharmacology

Minocycline hydrochloride pellet-filled capsules are rapidly absorbed from the gastrointestinal tract following oral administration. Following a single dose of two 100 mg pellet-filled capsules of minocycline hydrochloride administered to 18 normal fasting adult volunteers, maximum serum concentrations were attained in 1 to 4 hours (average 2.1 hours) and range from 2.1 to 5.1 mcg/mL (average 3.5 mcg/mL). The serum half-life in the normal volunteers ranged from 11.1 to 22.1 hours (average 15.5 hours).

When minocycline hydrochloride pellet-filled capsules were given concomitantly with a meal which included dairy products, the extent of absorption of minocycline hydrochloride pellet filled capsules was not noticeably influenced. The peak plasma concentrations were slightly decreased (11.2) and delayed by one hour when administered with food, compared to dosing under fasting conditions.

When minocycline hydrochloride tablets are administered with a meal including milk, the extent of absorption (AUC) is reduced by approximately 33% while the peak serum concentrations are reduced by approximately 32% and delayed one hour. In previous studies with other dosage forms, the minocycline half-life ranged from 11 to 16 hours in 7 patients with hepatic dysfunction, and from 18 to 69 hours in 5 patients with renal dysfunction. The urinary and faecal recovery of minocycline when administered to 12 normal volunteers is one-half to one-third that of other tetracyclines.

TOXICOLOGY

Minocycline hydrochloride has been tested in acute experiments in mice and rats, sub-chronic and chronic experiments in rats and dogs following oral and parenteral routes of administration.

Dietary administration of minocycline hydrochloride in long-term tumorigenicity studies in rats resulted in evidence of thyroid tumor production. In the rat, chronic treatment with minocycline hydrochloride has resulted in goiter accompanied by elevated radioactive iodine uptake and evidence of thyroid tumor production. Minocycline hydrochloride has been observed to cause a

dark discoloration of the thyroid in animals (rats, mice, dogs, and monkey). Minocycline hydrochloride has also been found to produce thyroid hyperplasia in rats and dogs.

In addition, there has been evidence of oncogenic activity in rats in studies with a related antibiotic, oxytetracycline (i.e., adrenal and pituitary tumors). Likewise, although mutagenicity studies of minocycline hydrochloride have not been conducted, positive results in *in vitro* mammalian cell assays (i.e., mouse lymphoma and Chinese hamster lung cells) have been reported for related antibiotics (tetracycline hydrochloride and oxytetracycline). Segment I (fertility and general reproduction) studies have provided evidence that minocycline hydrochloride impairs fertility in male rats.

The LD₅₀ of intravenous and intraperitoneal injections of minocycline in mice was 95 mg/kg and 280 mg/kg, respectively. The oral LD₅₀ in mice was 3100 mg/kg.

Minocycline hydrochloride has been given orally each day to dogs for six months at doses of 0, 4, 20 and 60 mg/kg/day (100 mg/kg/day for the first month) equally divided each day. At 20 mg/kg/day, there were no apparent drug-related findings except yellow discoloration of the skeleton and teeth in some animals, occasional emesis and black discoloration of the thyroid gland. At a dose of 4 mg/kg/day, there were no drug-related findings during the six month period, with the exception of discoloration of the thyroid gland and possibly some yellowing of the bones. Peak serum drug concentrations ranging from 8.5 to 100 mg/L were obtained with 60 and 100 mg/kg/day doses, 2.1 to 9.7 mg/L with the 20 mg/kg/day dose and 0.4 to 1.5 mg/L with the 4 mg/kg/day dose.

Minocycline hydrochloride was also given intravenously to dogs at doses of 5, 10, 20 and 40 mg/kg/day, a very similar dose range to that of the oral study, but administered for 1 month. Untoward findings such as body weight loss, reduced food consumption, erythema of the skin and of visible mucous membranes of varying duration, intensity and incidence, were associated primarily with the high dose (40 mg/kg/day). These findings were similar, except for erythema, to those obtained after the same dose of tetracycline. These drug-related findings with minocycline hydrochloride were associated with serum concentrations of 95 mg/L, three times those found with tetracycline (31 mg/L). Dogs that received 5, 10 and 20 mg/kg/day intravenously gave serum concentrations of 412 and 38 mg/L, respectively, and were found essentially to be without toxicity. These serum values are in considerable excess of those necessary for therapeutic effectiveness in man. In these experiments, minocycline hydrochloride appeared to be tolerated as well intravenously as it was orally.

Similar results were found following chronic oral administration of minocycline hydrochloride to rats for one year.

These animals were given a drug diet containing 0.008, 0.04, 0.2 and 1.0% minocycline hydrochloride, which corresponded to ranges of 4.4 to 8.5, 21.3 to 44.0, 108 to 122 and 593 to 812 mg/kg/day drug intake; these doses gave early morning plasma drug concentrations of 0.07 to 0.16, 0.36 to 0.51, 2.9 to 6.5 and 17 to 50 mg/L, respectively. With the exception of the

discoloration of the teeth (dose 0.04% drug diet or greater), femur and thyroid gland, there were no significant drug-related signs of toxicity at doses less than 1% drug diet.

As with other tetracyclines, minocycline hydrochloride has been found to produce discoloration of the thyroid gland in the rat, dog, monkey and human but not in the mouse. There was no evidence, however, from these investigations that thyroid function or bone growth was affected. A 23-month carcinogenicity study in the rat has shown that minocycline hydrochloride was not carcinogenic and that the black pigment in the thyroid gland did not cause neoplastic changes.

Biopsy specimens of thyroid tissue following the administration of minocycline hydrochloride and tetracycline to man revealed an intraepithelial lipofuscin deposition of both drugs, considered to be within normal variation. Thyroid function studies in man displayed a decrease within the normal range of thyroxine, indicating a tendency toward relative hypothyroidism.

Other than the tooth and bone discoloration that also occurs with other tetracyclines and the thyroid pigmentation seen in rats, dogs and monkeys, toxic effects of minocycline hydrochloride were observed only where serum concentrations were in excess of the therapeutic concentrations. It is concluded from the chronic safety evaluation studies that minocycline hydrochloride has a good margin of safety between therapeutic blood concentrations and concentrations producing toxic effects.

Reproduction studies performed in rats, rabbits and dogs have shown, as with other tetracyclines in animal studies that minocycline hydrochloride crosses the placenta, is found in fetal tissues and can produce toxic effects on the developing embryo, fetus or neonate when present in sufficient amounts.

The effects observed on the conceptus in rats and rabbits ranged from a low incidence of slight retardation of ossification and slight angulation of ribs at oral doses of 70 mg/kg/day in rats and 25 mg/kg/day in rabbits during pregnancy, to more extensive retardation of ossification and generalized morphologic changes and death at doses of 150 mg/kg/day and higher in the rat fetus. On other experiments, no deleterious effects were reported in rats or rabbits with oral doses as high as 100 and 75 mg/kg/day respectively. No adverse effects due to minocycline were seen in the newborn of 2 dogs given 20 mg/kg in 2 equally divided daily doses from days 35 to 62 of pregnancy.

REFERENCES

1. Anthony JR. Effect on deciduous and permanent teeth of tetracycline deposition in utero. *Postgrad Med* 1970;48(4):165-8.
2. Back MC, Zinner SH, Wilcox C, Finland M. Efficacy of standard disc diffusion test as applied to susceptibility of *Staphylococcus Aureus* to tetracycline and minocycline. *J Lab Clin Med* 77(2), 1972.
3. Bacon JF, Shenfield GM. Pregnancy attributable to interaction between tetracycline and oral contraceptives. *Br Med J* 1980;1:293.
4. Benitz KF, Roberts GKS, Yusa A. Morphologic effects of minocycline in laboratory animals. *Toxicol Appl Pharm* 1967;11:150-70.
5. Bernard B, Yin EJ, Simon HJ. Clinical pharmacologic studies with minocycline. *J Clin Pharm* 1971;332-48.
6. Bevelander G, Cohlan SQ. The effect on the rat fetus of transplacentally acquired tetracyclines. *Bio Neonat* 1962;4:365-70.
7. Bhattacharjee SB, Pal B, Bhaumik G. Further studies on tetracycline-induced mutation in V79 Chinese hamster cells. *Mutat Res* 1984;135:211-7.
8. Boucher D, Delast P. Développement post natal de la souris après traitement de la mère gestante et des descendants par les tétracyclines. *Compt Rend Soc Biol* 161:300-305,1967.
9. Boudreaux JP, Hayes DH, Mizrahi S, Hussey J, Regenstein F, Balart L. Fulminant hepatic failure, hepatorenal syndrome, and necrotizing pancreatitis after minocycline hepatotoxicity. *Transplant Proc* 1993;25(2):1873.
10. Burette A, Finet C, Prigogine T, De Roy G, Deltenre M. Acute hepatic injury associated with minocycline. *Arch Intern Med* 1984;144:1491-2.
11. Carney S, Butcher RA, Dawborn JK, Pattison G. Minocycline excretion and distribution in relation to renal function in man. *Clin Exp Pharm Physiol* 1974;1:299-308.
12. Cartwright AC, Hatfield HL, Yeadon A, London E. A comparison of the bioavailability of minocycline capsules and film-coated tablets. *J Antimicrob Chemother* 1975;1:317-22.
13. Chiu AM, Chuenkongkaew WL, Cornblath WT, Trobe JD, Digre KB, Dotan SA, et al. Minocycline treatment and pseudotumor cerebri syndrome. *Am J Ophthalmol* 1998;126(1):116-21.

14. Cohlman SQ, Bevelander G, Timsic T. Growth inhibition of prematures receiving tetracyclines. *Am J Dis Child* 1963;105:453-61.
15. Corcoran R, Castles JM. Tetracycline for acne vulgaris and possible teratogenesis. *Br Med J* 1977;2:807-8.
16. Cunha BA, Comer JB, Jonas M. The Tetracyclines. *Med Clin of North Am* 1982;66(1):293-302.
17. Danos EA. Apparent potentiation of warfarin activity by tetracycline. *Clin Pharm.* 1992;11:806-8.
18. Davies MG, Kersey PJW. Acute hepatitis and exfoliative dermatitis associated with minocycline. *BMJ* 1989;298(6686):1523-4.
19. Delaney RA, Wee D, Narayanaswamy TR. Pseudo-tumor cerebri and acne. *Milit Med* 1990;155(10):511.
20. Fedorke J, Katz S and Allnoch H. *In Vitro* activity of minocycline, a new tetracycline. *Amer J Med Sci* 255:252-258 Apr 1968.
21. Fields JP. Bulging fontanel: a complication of tetracycline therapy in infants. *J Pediatr* 1961;58(1):74-6.
22. Frisk AR, Tunevall G. Minocycline: clinical evaluation, abstract of papers. Eighth Interscience Conference on Antimicrobial Agents and Chemotherapy. New York (Oct.) 1968 p. 49
23. George CRP, Guinness MDG, Lark DJ, Evans RA. Minocycline toxicity in renal failure. *Med J Aust* 1973;1:640-1.
24. Graber CD, Jervey LP, Martin F, Boltjes BH. *In Vitro* and *In Vivo* sensitivity of staphylococci and selected bacteria to minocycline and doxycycline. Abstracts of Papers, Eighth Interscience Conference on Antimicrobial Agents and Chemotherapy. New York (Oct.) 1968 p. 49.
25. Grove Donald C, Randall WA. Assay methods of antibiotics: A laboratory manual. New York, 1955 *Med Encyclopedia*, (Follow the method for chlortetracycline.)
26. Gugler R, Allgayer H. Effects of antacids on the clinical pharmacokinetics of drugs. An update. *Clin Pharmacokinet* 1990;18(3):210-19.
27. Halme J, Aes J. Inhibition of collagen synthesis and bone calcification in foetal rat by tetracycline. *Scand J Clin Lab Invest* 21:4(Suppl 101)1968.
28. Jao RL, Finland M. Susceptibility of mycoplasma pneumoniae to 21 antibiotics *In Vitro*. *Amer J Med Sci* 253:639-650 June 1967.

29. Kasamaki A, Urasawa S. Induction of mutagenic activity by synergistic action of multiple chemicals. *J Toxicol. Sci* 1979;4:310-1.
30. Kelly RG, Kanegis LA. Metabolism and tissue distribution of radioisotopically labeled minocycline. *Toxicol Appl Pharmacol* 11:171-183,1967.
31. Kline AH, Blattner RJ, Lunin M. Transplacental effect of tetracyclines on teeth. *JAMA* 1964;188(2):178-80.
32. Kasamaki A, Urasawa S. Induction of mutagenic activity by synergistic action of multiple chemicals. *J Toxicol. Sci* 1979;4:310-1.
33. Krevsky S. The bulging fontanelle syndrome following tetracycline administration. *Mich Med* 1968;67(9):597-8.
34. Kuck NA, Redin GS, Forbes M. Activity of minocycline and other tetracyclines against tetracycline-sensitive and resistant staphylococci. *Pro Soc Exp Biol Med* 136:479-481,1971.
35. Lander CM. Minocycline-induced benign intracranial hypertension: *Clin Exp Neurol*. 1989;26:161-7.
36. Lee AG. Pseudotumor cerebri after treatment with tetracycline and isotretinoin for acne. *Cutis®* 1995;55:165-8.
37. Lepper MH, Dowling HF. Treatment of pneumococcal meningitis with penicillin compared with penicillin plus aureomycin. *Arch Intern Med* 1951;88:489-94.
38. Lester MR. Sulfite sensitivity: significance in human health. *J Amer Coll Nutr* 1995;14(3):229-32.
39. Leyden JJ. Absorption of minocycline hydrochloride and tetracycline hydrochloride. Effect of food, milk, and iron. *J Am Acad Dermatol* 1985;12:308-312.
40. Lewis PA, Kearney PJ. Pseudotumor cerebri induced by minocycline treatment for acne vulgaris. *Acta Dermato Venereologica* 1997;77(1):83.
41. Little PJ, Bailey RR. Tetracyclines and renal failure. *N Z Med J* 1970;72(460):183-4.
42. Macdonald H. Effect of food and milk on absorption of minocycline. *American Cyanamid Company Clinical Study* 66-37, 168-79, 1967.
43. Macdonald H, Kelly RG, Allen ES, Noble JF, Kanegis LA. Pharmacokinetic studies on minocycline in man. *Clin Pharm Ther* 1973;14(5):852-61.

44. Malcolm A, Heap TR, Eckstein RP, Lunzer MR. Minocycline-induced liver injury. *Amer J Gastroenterol* 1996;91(8):1641-3.
45. McGregor DB, Brown AG, Howgate S, McBride D, Riach C, Casprary WJ. Responses of the L5178Y mouse Lymphoma cell forward mutation assay. V:27 coded chemicals. *Environ Mol Mutagen* 1991;17:196-219.
46. McHenry MC, Gavan TL, Vidt DG, Jameson S, Wagner JG. Minocycline in renal failure. *Clinical Pharmacology and Therapeutics* 13:146,1972.
47. Min DI, Burke PA, Lewis WD, Jenkins RL. Acute Hepatic Failure Associated with Oral Minocycline: A case report. *Pharmacotherapy* 1992;12(1):68-71.
48. MINOCIN® Product Monograph, Professional Services Lederle Laboratories, 1990.
49. Minutello JS, Dimayuga RG, Carter J. Pseudotumor cerebri, a rare adverse reaction to tetracycline therapy: A case report. *J Periodontol* 1988;59(12):848-51.
50. Naline E, Sanceaume M, Toty L, Bakdach H, Pays M, Advenier C. Penetration of Minocycline into lung tissues. *Br J Clin Pharmacol* 1991;32:402-4.
51. Neuvonen PJ, Pentikainen PJ, Gothoni G. Inhibition of iron absorption by tetracycline {letter}. *Br J Clin Pharmacol* 1975;2(1):94-6.
52. Nishijima S, Namura S, Kawai S, Akamatsu H, Asada Y, Kawabata S. Sensitivity of *Staphylococcus aureus* and *Streptococcus pyogenes* isolated from skin infections in 1992 to antimicrobial agents. *J Dermatol* 1994;21:233-8.
53. Noble JF, Kanegis LA, Hallesy DW. Short-term toxicity and observations on certain aspects of the pharmacology of a unique tetracycline - minocycline. *Toxicol Appl Pharmacol* 11:128-149,1967.
54. Pellagali GV, d'Angelo A. Effetto della tetracycline cloridrato sulla comparsa dei nuclei di ossificazione di embrioni di ratto. *Bull Soc Ital Sper* 40:13211324,1964.
55. Posner AC, Prigot A, Konicoff NG. Further observations on the use of tetracycline hydrochloride in prophylaxis and treatment of obstetric infections. In Welch, H, and Morti-Ibanex, F, eds. *Antibiotics Annual 1954-1955*. New York Medical Encyclopedia, Inc. 1955:594-8.
56. Redin GS. Antibacterial activity in mice of minocycline, a new tetracycline. *Antimicrob Ag Chemother* p. 371-376,1966.
57. Renzini G, Bevilacqua RL, Boemi G, Ravagnan L. Antimicrobial *In Vitro* and *In Vivo* activity of a new tetracycline, CL 59, 806. *Antibiotica* 5:241-261. (Dec.) 1967.

58. Sadowski DC. Drug interactions with antacids: mechanisms and clinical significance. *Drug Saf* 1994;11:395-407.
59. Saito K., Jujio T, Hashizume I, Yamada T, Onaya T, Uehara T, et al. Studies on goitrogenic action of minocycline and related compounds. *Endocrinology* 1972;90(5):1192-201.
60. Saivin S, Houin G. Clinical pharmacokinetics of doxycycline and minocycline. *Clin Pharmacokinet* 1988;15:355-66.
61. Saxen L. Effect of tetracycline on osteogenesis *In Vitro*. *J Exper Zool* 162:269-294,1966.
62. Schlegel PN, Chang TSK, Marshall FF. Antibiotics: potential hazards to male fertility. *Fertil Steril* 1991;55(2):235-42.
63. Searcy RL, Craig RG, Foreman JA, Berquist LM. Blood clotting anomalies with intensive tetracycline therapy. *Clin Res* 1965;12:230.
64. Segal BM. Photosensitivity, nail discoloration, and onycholysis: side effects of tetracycline therapy. *Arch Intern Med* 1963;112:165-7.
65. Shils ME. Renal disease and the metabolic effects of tetracycline. *Ann Intern Med* 1963;58:389-408.
66. Sklenar I, Spring P, Dettli L. One-dose and multiple-dose kinetics of minocycline in patients with renal disease. *Agents Actions* 1977;7(3):369-77.
67. Steigbigel NH, McCall CE, Reed CW and Finland M. Antibacterial action of "Broad Spectrum" Penicillins, cephalosporins and other antibiotics against gram-negative bacilli isolated from bacteremic patients. *Ann N Y Acad Sci* 145:224-236(Sept. 27)1967.
68. Steigbigel NH, Reed CW, Finland M. Susceptibility of common pathogenic bacteria to seven tetracycline antibiotics *In Vitro*. *Amer J Med Sci* 225:179-195 Mar 1968.
69. Steigbigel NH, Reed CW, Finland M. Absorption and excretion of five tetracycline analogues in normal young men. *Amer J Med Sci* 255:296-312 May 1968.
70. Steiner G, Bradford W, Craig JM. Tetracycline-induced abortion in the rat. *Lab Invest* 1965;14(8):1456-63.
71. Traub WH, Leonhard B. Comparative susceptibility of clinical group A, B, C, F, and G β -hemolytic streptococcal isolates to 24 antimicrobial drugs. *Chemother* 1997;43:10-20.
72. Wallman IS, Hilton HB. Teeth pigmented by tetracycline. *Lancet* 1962;1:827-9.

73. Walters BNJ, Gubbay SS. Tetracycline and benign intracranial hypertension: report of five cases. *BMJ* 1981;282(6257):19-20.
74. Weller M, Klockgether T. Minocycline-induced benign intracranial hypertension. *J. Neurology* 1998;245(1):55.
75. Welling PG, Shaw WR, Uman SJ, Tse FLS, Craig WA. Pharmacokinetics of minocycline in renal failure. *Antimicrob Agents Chemother* 1975;8:532-7.
76. Williams DN, Laughlin LW, Lee YH. Minocycline: possible vestibular side-effects. *Lancet* 1974;2:744-6.
77. Zachariassen RD. Loss of oral contraceptive efficacy by concurrent antibiotic administration. *Women Health*. 1994;22(1):17-26.
78. PRESCRIBING INFORMATION, ^{Pf}Minocin®, minocycline hydrochloride, 50 mg and 100 mg Capsules, GlaxoSmithKline Inc., Submission Control No: 153630, Date of Revision: April 16, 2012
79. Apotex Inc. Prescribing Information: ^{Pf}Apo-Minocycline, Minocycline Hydrochloride Capsules USP, 50 mg and 100 mg Capsules. Control Number:210873, Date of Revision : November 14, 2018

READ THIS FOR SAFE AND EFFECTIVE USE OF YOUR MEDICINE

PATIENT MEDICATION INFORMATION

Sandoz Minocycline Minocycline Hydrochloride Capsules

Read this carefully before you start taking Sandoz Minocycline and each time you get a refill. This leaflet is a summary and will not tell you everything about this drug. Talk to your healthcare professional about your medical condition and treatment and ask if there is any new information about Sandoz Minocycline.

What is Sandoz Minocycline used for?

Sandoz Minocycline is used for the treatment of infections caused by certain bacteria in many different parts of the body.

Antibacterial drugs like Sandoz Minocycline treat only bacterial infections. They do not treat viral infections.

How does Sandoz Minocycline work?

Sandoz Minocycline is an antibiotic that belong to a class of drugs called tetracyclines. Sandoz Minocycline works by slowing the growth or reproduction of bacteria that causes the infection.

What are the ingredients in Sandoz Minocycline?

Medicinal Ingredient: minocycline hydrochloride

Nonmedicinal ingredients: cornstarch, gelatin, magnesium stearate The 50 mg capsule also contains the non-medicinal ingredients: sunset yellow FCF and titanium dioxide. Tartrazine-free.. The 100 mg capsule also contains the non-medicinal ingredients : erythrosine, patent blue V, sunset yellow FCF, titanium dioxide. Tartrazine-free.

Sandoz Minocycline comes in the following dosage forms:

Capsules: 50 mg and 100 mg.

Do not use Sandoz Minocycline if you:

- are allergic to minocycline or to any of the other ingredients in Sandoz Minocycline
- are allergic to other tetracycline antibiotics
- are pregnant or planning to become pregnant. Sandoz Minocycline can cause damage to your unborn babies bones and teeth. If you get pregnant while taking Sandoz Minocycline contact to your healthcare professional immediately.
- are breastfeeding or planning to breastfeed. Sandoz Minocycline passes into human milk and can cause damage to your babies bones and teeth. You should not breastfeed while taking Sandoz Minocycline.
- have liver or kidney problems
- have the autoimmune disease myasthenia gravis

- are lactose intolerant or have one of the following rare hereditary diseases:
 - Galactose intolerance
 - Fructose intolerance
 - Sucrose-isomaltase insufficiency
 - Lapp lactase deficiency
 - Glucose-galactose malabsorption

Because lactose is a non-medicinal ingredient in Sandoz Minocycline.

Sandoz Minocycline is not recommended for children under 13 years old.

To help avoid side effects and ensure proper use, talk to your healthcare professional before you take Sandoz Minocycline. Talk about any health conditions or problems you may have, including if you:

- have liver or kidney problems
- have the autoimmune disease systemic lupus erythematosus
- suffer from allergies, asthma, hay fever or itchy skin rashes. You may be more likely to experience side effects while taking Sandoz Minocycline.
- are also taking a penicillin antibiotic
- are taking birth control pills. Sandoz Minocycline may make your birth control pills less effective. Talk to your healthcare professional about using a back-up method of birth control, such as condoms, while you are taking Sandoz Minocycline.

Other warnings you should know about:

Sensitivity to Sunlight

Sandoz Minocycline can make your skin more sensitive to the sun. While you are taking Sandoz Minocycline avoid direct sunlight, sunlamps and tanning beds. If you develop any skin redness or discomfort while taking Sandoz Minocycline, contact your healthcare professional immediately.

Driving and Using Machines

Sandoz Minocycline can cause headaches, light-headedness, dizziness, ringing in the ears and vertigo (feeling like you're spinning). Use caution when driving or using dangerous machinery while taking Sandoz Minocycline.

Serious Side Effects

Serious side effects, in some cases causing death, have been seen in people taking Sandoz Minocycline. If you have any of the following symptoms stop taking Sandoz Minocycline and seek immediate medical help.

- **Allergic reaction (hypersensitivity) with skin rashes:** rash, skin redness, peeling skin, blisters on your nose, mouth, eyes or genitals, hives, swelling of the face, lips, tongue or throat, difficulty swallowing or breathing, fever, swollen lymph nodes combined with any of the following: liver problems (yellowing of the skin or eyes, dark urine, abdominal pain, nausea, vomiting, loss of appetite), lung problems (flu-like symptoms, cough, shortness of breath), kidney problems (decreased urination, nausea,

vomiting, swelling of extremities, fatigue), heart problems (chest pain, fast or irregular heartbeat, shortness of breath, swelling in the legs, ankles or feet, fatigue)

- **Lupus-like syndrome:** joint and muscle pain, fatigue, fever, swollen lymph nodes, inflammation around the lungs or heart that causes pain or discomfort
- **Serum-sickness syndrome:** fever, generally feeling unwell, hives, itching, rash, joint pain, swollen lymph nodes

Tell your healthcare professional about all the medicines you take, including any drugs, vitamins, minerals, natural supplements or alternative medicines.

The following may interact with Sandoz Minocycline:

- penicillin antibiotics, used to treat infections
- acne medicines known as retinoids
- quinapril, used to treat heart problems and high blood pressure
- blood thinners, used to prevent blood clots
- birth control pills, Sandoz Minocycline may make your birth control pills less effective.
- diuretics or “water pills” used to treat high blood pressure
- medicines used to treat migraines called ergot alkaloids
- antacids containing aluminum calcium, magnesium, bismuth or iron containing products
- magnesium or zinc salts

How to take Sandoz Minocycline:

- Although you may feel better early in treatment, Sandoz Minocycline should be used exactly as directed.
- Misuse or overuse of Sandoz Minocycline could lead to the growth of bacteria that will not be killed by Sandoz Minocycline (resistance). This means that Sandoz Minocycline may not work for you in the future.
- Do not share your medicine.

Usual Dose:

Children 13 years of age and older: Your healthcare professional will tell you how much Sandoz Minocycline to give your child and how often based on their body weight.

Adults: 100 mg or 200 mg to start followed by 100 mg every 12 hours.

Overdose:

If you think you have taken too much Sandoz Minocycline, contact your healthcare professional, hospital emergency department or regional poison control centre immediately, even if there are no symptoms.

Missed Dose:

If you forget to take a dose of Sandoz Minocycline take it as soon as you remember.

What are possible side effects from using Sandoz Minocycline?

These are not all the possible side effects you may feel when taking Sandoz Minocycline. If you experience any side effects not listed here, contact your healthcare professional.

Side effects may include:

- light-headedness, dizziness, vertigo (feeling like you’re spinning or losing your balance)
- headache
- loss of appetite, constipation
- nausea, vomiting, diarrhea
- indigestion
- trouble swallowing
- inflammation or sores in the mouth, lips and/or tongue
- joint swelling, pain and stiffness, aching muscles
- rash, itching
- cough
- sleepiness
- fever

Sandoz Minocycline can cause abnormal blood test results. Your healthcare professional will decide when to perform blood tests and will interpret the results.

Serious side effects and what to do about them			
Symptom / effect	Talk to your healthcare professional		Stop taking drug and get immediate medical help
	Only if severe	In all cases	
RARE			
Fainting		√	
Liver problems: yellowing of the skin or eyes, dark urine, abdominal pain, nausea, vomiting, loss of appetite			√
Inflammation or ulcers of the food pipe (esophagus): trouble swallowing, pain when swallowing, sore throat, hoarse voice, heartburn, chest pain that is worse with eating, nausea			√
Kidney problems: decreased urination, nausea, vomiting, swelling of extremities, fatigue			√
Unusual hair loss or thinning	√		
Increased sensitivity of the skin to sun		√	

Discolouration of the nails, skin, tongue, gums, lips or teeth	√		
Shortness of breath		√	
VERY RARE			
Pseudomembranous colitis: watery, bloody diarrhea, mucus in the stool, abdominal cramps and pain, fever			√
Angioedema: swelling of the face, eyes, lips, tongue, throat, arms or legs, trouble breathing or swallowing, hoarseness, itching, hives, rash on the hands, arms and feet, fever, abdominal cramps			√
Allergic reaction (hypersensitivity) with skin rashes: rash, skin redness, peeling skin, blisters on your nose, mouth, eyes or genitals, hives, swelling of the face, lips, tongue or throat, difficulty swallowing or breathing, fever, swollen lymph nodes combined with any of the following: liver problems (yellowing of the skin or eyes, dark urine, abdominal pain, nausea, vomiting, loss of appetite), lung problems (flu-like symptoms, cough, shortness of breath), kidney problems (decreased urination, nausea, vomiting, swelling of extremities, fatigue), heart problems (chest pain, fast or irregular heartbeat, shortness of breath, swelling in the legs, ankles or feet, fatigue)			√
Thyroid problems: weight changes, fatigue, heart palpitations, constipation, dry skin, joint or muscle pain		√	
Yeast infections: Oral: creamy white bumps on the tongue, cheeks, gums or throat that bleed when scraped, pain, trouble swallowing, bad taste in the mouth Genital and Anal: genital (vagina or penis) or anal itching, burning during intercourse or urination, pain, redness, swelling, discharge		√	
Bronchospasm: pain, tightness and a feeling of constriction in the chest and back, trouble breathing, wheezing, coughing, dizziness			√
NOT KNOWN			
Hearing problems: buzzing, ringing or other persistent noise in ear, loss of hearing	√		
Tingling or numbness of the hands or feet	√		

Lupus-like syndrome: joint and muscle pain, fatigue, fever, swollen lymph nodes, inflammation around the lungs or heart that causes pain or discomfort			√
High blood pressure in the brain: headache, blurred vision, nausea, vomiting, confusion			√
Respiratory disorder:			
Pneumonitis: flu-like symptoms, cough, shortness of breath, fever, chills, fatigue			√
Nervous system disorder:			
Seizures or fits			√
Gastrointestinal disorder:			
Inflammation of the pancreas: abdominal pain that lasts and gets worse when you lie down, nausea, vomiting		√	
Immune system disorder:			
Serum-sickness syndrome: fever, generally feeling unwell, hives, itching, rash, joint pain, swollen lymph nodes	√		
Blood disorder:			
Hemolytic anemia: pale skin, yellow skin, eyes and mouth (jaundice), darkcoloured urine, fever, weakness, dizziness, confusion, reduced ability to exercise		√	
Low levels of white blood cells: infection, fatigue, fever, aches, pain, flu-like symptoms		√	
Low levels of blood platelets: bruising, bleeding, fatigue, weakness		√	

If you have a troublesome symptom or side effect that is not listed here or becomes bad enough to interfere with your daily activities, talk to your healthcare professional.

Reporting Side Effects

You can report any suspected side effects associated with the use of health products to Health Canada by:

- Visiting the Web page on Adverse Reaction Reporting (<https://www.canada.ca/en/healthcanada/services/drugs-health-products/medeffect-canada/adverse-reaction-reporting.html>) for information on how to report online, by mail or by fax; or
- Calling toll-free at 1-866-234-2345.

NOTE: Contact your health professional if you need information about how to manage your side effects. The Canada Vigilance Program does not provide medical advice.

Storage :

Store at room temperature and protect from light.

Keep out of reach and sight of children.

If you want more information about Sandoz Minocycline :

- Talk to your healthcare professional
- Find the full product monograph that is prepared for healthcare professionals and includes this Patient Medication Information by visiting the Health Canada website (<https://www.canada.ca/en/health-canada/services/drugs-health-products/drug-products/drug-product-database.html>); the manufacturer's website <http://www.sandoz.ca> or by calling 1-800-361-3062.

This leaflet was prepared by Sandoz Canada Inc.

Last Revised: April 29, 2019