

recommend no print on short lap (except full coverage color)

2019-05-08

DIN 02321920

# Avatec<sup>®</sup> 20

## Lasalocid Sodium Medicated Premix

### Veterinary Use Only

For Feed Manufacturing Use Only

Active Ingredient:  
Each kg contains 200 g (20%) of lasalocid sodium in a carrier suitable for incorporation in feed.

PRODUCT CODE: 710154

NET 25 KG

Zoetis is a trademark and Avatec is a registered trademark of Zoetis or its licensors, used under license by Zoetis Canada Inc.

zoetis

Avatec<sup>®</sup> 20

Lot See bag seal  
EXP See bag seal



Zoetis Canada Inc.  
Kirkland QC H9H 4M7

DIN 02321920

# Avatec<sup>®</sup> 20

## Prémélange médicamenteux de lasalocide sodique

### Usage vétérinaire seulement

Réservé à la fabrication de moulées

Ingrédient actif :  
Chaque kg renferme 200 g (20 %) de lasalocide sodique dans un support propice à l'incorporation dans les aliments

CODE DU PRODUIT : 710154

NET 25 KG

Zoetis est une marque de commerce et Avatec est une marque déposée de Zoetis ou de ses concédants de licence, utilisées sous licence par Zoetis Canada Inc.

#### FOR USE IN BROILER CHICKEN AND TURKEY FEEDS

**Indications**  
A. As an aid in the prevention of coccidiosis caused by *Eimeria tenella*, *E. tenella*, *E. scrovaria*, *E. brunetti*, *E. mitis* and *E. necatrix* in broiler chickens.  
B. As an aid in the prevention of coccidiosis caused by *E. adenovaridis*, *E. meleagridis* and *E. gallopavonis* in growing chickens.

**Mixing Directions**  
Thoroughly mix 0.55 (0.52 kg) of Avatec<sup>®</sup> 20 Medicated Premix in one tonne (1000 kg) of feed to provide 100 mg of lasalocid sodium per kg of complete feed. It is recommended that Avatec<sup>®</sup> 20 be mixed with a small quantity of the feed to ensure uniform distribution throughout the batch.

**Feeding Directions**  
A. As an aid in the prevention of coccidiosis caused by *E. adenovaridis*, *E. meleagridis* and *E. gallopavonis* in growing chickens.  
B. As an aid in the prevention of coccidiosis caused by *E. tenella*, *E. scrovaria*, *E. brunetti*, *E. mitis* and *E. necatrix* in broiler chickens.

**Caution**  
1. Do not use this medicated feed for treatment of coccidiosis.  
2. Consult a veterinarian or poultry pathologist if losses exceed 0.5% in a broiler period.

**Warnings**  
1. No withdrawal period is required for broiler chickens and turkeys when treated according to the label.  
2. Do not feed to pregnant, laying and brooding chickens and turkeys.  
3. Keep out of reach of children.

#### FOR USE IN CATTLE AND LAMB FEEDS

**Indications**  
A. For improved feed efficiency and increased rate of weight gain in feeder cattle being fed in confinement for slaughter.  
B. For increased rate of weight gain in pasture cattle (steers, heifer cattle, and beef and dairy replacement heifers).  
C. As an aid in the prevention of coccidiosis caused by *Eimeria bovis* and *Eimeria zuernii* in calves up to 360 kg of body weight being fed in confinement.

**Mixing Directions**  
Thoroughly mix the supplement in the total daily diet or in the complete feed (gran portion of the ration) before use. Feed continuously during periods of exposure to coccidiosis or when coccidiosis is likely to be a hazard.

**Level of Drug in Supplement**  
Supplement must provide an average daily intake as follows:

| Weight of cattle (kg) | Recommended lasalocid dosage (mg/head/day) | 0.5 kg | 1.0 kg | 1.5 kg | 2.0 kg |
|-----------------------|--|--------|--------|--------|--------|
| 200-250               | 200  | 2.00   | 1.25   | 0.80   | 0.63   |
| 300                   | 300  | 3.00   | 1.75   | 1.17   | 0.87   |

**Caution**  
1. Do not allow horses or other equine access to lasalocid sodium as ingestion may be fatal (equine only).  
2. Do not feed to pregnant, laying and brooding chickens and turkeys.

**Warnings**  
1. Treated sheep must not be slaughtered for use in food for at least 2 days after the latest treatment with this drug.  
2. No withdrawal period is required for cattle when treated according to the label.  
3. Keep out of reach of children.

Thoroughly mix the supplement in the total daily diet or in the complete feed (gran portion of the ration) before use.

**Mixing Directions**  
Thoroughly mix 0.55 (0.52 kg) of Avatec<sup>®</sup> 20 Medicated Premix in one tonne (1000 kg) of feed to provide 100 mg of lasalocid sodium per kg of complete feed.

**Feeding Directions**  
A. As an aid in the prevention of coccidiosis caused by *Eimeria tenella*, *E. tenella*, *E. scrovaria*, *E. brunetti*, *E. mitis* and *E. necatrix* in broiler chickens.  
B. As an aid in the prevention of coccidiosis caused by *E. adenovaridis*, *E. meleagridis* and *E. gallopavonis* in growing chickens.

**Indications**  
A. As an aid in the prevention of coccidiosis caused by *Eimeria tenella*, *E. tenella*, *E. scrovaria*, *E. brunetti*, *E. mitis* and *E. necatrix* in broiler chickens.  
B. As an aid in the prevention of coccidiosis caused by *E. adenovaridis*, *E. meleagridis* and *E. gallopavonis* in growing chickens.

**Mixing Directions**  
Thoroughly mix 0.55 (0.52 kg) of Avatec<sup>®</sup> 20 Medicated Premix in one tonne (1000 kg) of feed to provide 100 mg of lasalocid sodium per kg of complete feed. It is recommended that Avatec<sup>®</sup> 20 be mixed with a small quantity of the feed to ensure uniform distribution throughout the batch.

**Feeding Directions**  
A. As an aid in the prevention of coccidiosis caused by *E. adenovaridis*, *E. meleagridis* and *E. gallopavonis* in growing chickens.  
B. As an aid in the prevention of coccidiosis caused by *E. tenella*, *E. scrovaria*, *E. brunetti*, *E. mitis* and *E. necatrix* in broiler chickens.

**Caution**  
1. Do not use this medicated feed for treatment of coccidiosis.  
2. Consult a veterinarian or poultry pathologist if losses exceed 0.5% in a broiler period.

**Warnings**  
1. No withdrawal period is required for broiler chickens and turkeys when treated according to the label.  
2. Do not feed to pregnant, laying and brooding chickens and turkeys.  
3. Keep out of reach of children.

Thoroughly mix the supplement in the total daily diet or in the complete feed (gran portion of the ration) before use. Feed continuously during periods of exposure to coccidiosis or when coccidiosis is likely to be a hazard.

| Weight of cattle (kg) | Recommended lasalocid dosage (mg/head/day) | 0.5 kg | 1.0 kg | 1.5 kg | 2.0 kg |
|-----------------------|--|--------|--------|--------|--------|
| 100-150               | 100  | 1.00   | 0.50   | 0.33   | 0.25   |
| 100-140               | 100  | 1.50   | 0.75   | 0.50   | 0.37   |
| 100-130               | 200  | 2.00   | 1.00   | 0.67   | 0.50   |
| 200-250               | 200  | 2.00   | 1.25   | 0.80   | 0.63   |
| 200-200               | 300  | 3.00   | 1.50   | 1.00   | 0.75   |
| 300-300               | 300  | 3.00   | 1.75   | 1.17   | 0.87   |

**Level of Drug in Supplement**  
Supplement must provide an average daily intake as follows:

| Weight of cattle (kg) | Recommended lasalocid dosage (mg/head/day) | 0.5 kg | 1.0 kg | 1.5 kg | 2.0 kg |
|-----------------------|--|--------|--------|--------|--------|
| 10                    | 10   | 0.10   | 0.10   | 0.10   | 0.10   |
| 20                    | 20   | 0.20   | 0.20   | 0.20   | 0.20   |
| 30                    | 30   | 0.30   | 0.30   | 0.30   | 0.30   |
| 40                    | 40   | 0.40   | 0.40   | 0.40   | 0.40   |
| 50                    | 50   | 0.50   | 0.50   | 0.50   | 0.50   |
| 75                    | 75   | 0.75   | 0.75   | 0.75   | 0.75   |

**Caution**  
1. Do not allow horses or other equine access to lasalocid sodium as ingestion may be fatal (equine only).  
2. Do not feed to pregnant, laying and brooding chickens and turkeys.

**Warnings**  
1. Treated sheep must not be slaughtered for use in food for at least 2 days after the latest treatment with this drug.  
2. No withdrawal period is required for cattle when treated according to the label.  
3. Keep out of reach of children.

Thoroughly mix the supplement in the total daily diet or in the complete feed (gran portion of the ration) before use. Feed continuously during periods of exposure to coccidiosis or when coccidiosis is likely to be a hazard.

| Weight of cattle (kg) | Recommended lasalocid dosage (mg/head/day) | 0.5 kg | 1.0 kg | 1.5 kg | 2.0 kg |
|-----------------------|--|--------|--------|--------|--------|
| 200-250               | 200  | 2.00   | 1.25   | 0.80   | 0.63   |
| 300                   | 300  | 3.00   | 1.75   | 1.17   | 0.87   |

**Level of Drug in Supplement**  
Supplement must provide an average daily intake as follows:

| Weight of cattle (kg) | Recommended lasalocid dosage (mg/head/day) | 0.5 kg | 1.0 kg | 1.5 kg | 2.0 kg |
|-----------------------|--|--------|--------|--------|--------|
| 200-250               | 200  | 2.00   | 1.25   | 0.80   | 0.63   |
| 300                   | 300  | 3.00   | 1.75   | 1.17   | 0.87   |

**Caution**  
1. Do not allow horses or other equine access to lasalocid sodium as ingestion may be fatal (equine only).  
2. Do not feed to pregnant, laying and brooding chickens and turkeys.

**Warnings**  
1. Treated sheep must not be slaughtered for use in food for at least 2 days after the latest treatment with this drug.  
2. No withdrawal period is required for cattle when treated according to the label.  
3. Keep out of reach of children.

Thoroughly mix the supplement in the total daily diet or in the complete feed (gran portion of the ration) before use.

**Mixing Directions**  
Thoroughly mix 0.55 (0.52 kg) of Avatec<sup>®</sup> 20 Medicated Premix in one tonne (1000 kg) of feed to provide 100 mg of lasalocid sodium per kg of complete feed.

**Feeding Directions**  
A. As an aid in the prevention of coccidiosis caused by *Eimeria tenella*, *E. tenella*, *E. scrovaria*, *E. brunetti*, *E. mitis* and *E. necatrix* in broiler chickens.  
B. As an aid in the prevention of coccidiosis caused by *E. adenovaridis*, *E. meleagridis* and *E. gallopavonis* in growing chickens.

**Caution**  
1. Do not use this medicated feed for treatment of coccidiosis.  
2. Consult a veterinarian or poultry pathologist if losses exceed 0.5% in a broiler period.

**Warnings**  
1. No withdrawal period is required for broiler chickens and turkeys when treated according to the label.  
2. Do not feed to pregnant, laying and brooding chickens and turkeys.  
3. Keep out of reach of children.

Avatec<sup>®</sup> 20

zoetis

710154-20

| Project No. | Batch Number | Description | Country |
|-------------|--------------|-------------|---------|
| 12907       | 3025200      | Avatec 20%  | Canada  |

  

| Part No.         | Part Description | Part No. | Part Description |
|------------------|------------------|----------|------------------|
| 38" x 16.5" x 6" | DWG-300175-00    | 710154   | Bag              |

  

| Address Info                | Colours | Part No. | Part Description |
|-----------------------------|---------|----------|------------------|
| JDE Component Code - 710254 | PSI 308 | Diète    | EDITOR'S COPY    |

  

| Rev | By        | Date     | Rev | By        | Date     |
|-----|-----------|----------|-----|-----------|----------|
| 1   | J. Hicks  | 02/07/14 | 1   | J. Hicks  | 02/07/14 |
| 2   | L. Nelson | 04/07/14 | 2   | L. Nelson | 04/07/14 |

710154-20  
3025200