

## **PRODUCT MONOGRAPH**

**<sup>Pr</sup> APO-QUINAPRIL/HCTZ**

**Quinapril and Hydrochlorothiazide Tablets  
Apotex Standard**

**10/12.5, 20/12.5 and 20/25 mg**

**Angiotensin Converting Enzyme Inhibitor/Diuretic**

APOTEX INC.  
150 Signet Drive  
Toronto, Ontario  
M9L 1T9

**Date of Revision:**  
June 10, 2019

**Submission Control No.: 228262**

## Table of Contents

<b>PART I: HEALTH PROFESSIONAL INFORMATION.....</b>	<b>3</b>
SUMMARY PRODUCT INFORMATION .....	3
INDICATIONS AND CLINICAL USE .....	3
CONTRAINDICATIONS .....	3
WARNINGS AND PRECAUTIONS .....	4
ADVERSE REACTIONS .....	13
DRUG INTERACTIONS .....	16
DOSAGE AND ADMINISTRATION .....	24
OVERDOSAGE .....	26
ACTION AND CLINICAL PHARMACOLOGY .....	26
STORAGE AND STABILITY .....	28
DOSAGE FORMS, COMPOSITION AND PACKAGING .....	28
COMPOSITION .....	29
<b>PART II: SCIENTIFIC INFORMATION.....</b>	<b>30</b>
PHARMACEUTICAL INFORMATION.....	30
CLINICAL TRIALS .....	33
DETAILED PHARMACOLOGY .....	34
TOXICOLOGY .....	35
REFERENCES .....	44
<b>PART III: CONSUMER INFORMATION .....</b>	<b>46</b>

**Pr APO-QUINAPRIL/HCTZ**  
**Quinapril and Hydrochlorothiazide Tablets**  
**Apotex Standard**

**PART I: HEALTH PROFESSIONAL INFORMATION**

**SUMMARY PRODUCT INFORMATION**

<b>Route of Administration</b>	<b>Dosage Form/ Strength</b>	<b>All Nonmedicinal Ingredients</b>
Oral	Tablet: 10/12.5, 20/12.5 and 20/25 mg	copovidone, crospovidone, ferric oxide red, ferric oxide yellow, hydroxypropyl cellulose, hypromellose, magnesium carbonate, polyethylene glycol, titanium dioxide, zinc stearate

**INDICATIONS AND CLINICAL USE**

APO-QUINAPRIL/HCTZ Tablets (quinapril hydrochloride and hydrochlorothiazide) are indicated for the treatment of essential hypertension in patients for whom combination therapy is appropriate.

APO-QUINAPRIL/HCTZ Tablets are not indicated for initial therapy. Patients in whom quinapril and hydrochlorothiazide are initiated simultaneously can develop symptomatic hypotension (see WARNINGS AND PRECAUTIONS, **Cardiovascular**, - **Hypotension** and DRUG INTERACTIONS).

Patients should be titrated on the individual drugs. If the fixed combination represents the dosage determined by this titration, the use of APO-QUINAPRIL/HCTZ Tablets may be more convenient in the management of patients. If during maintenance therapy dosage adjustment is necessary, it is advisable to use individual drugs.

**Pediatrics (<18 years of age)**

The safety and effectiveness of APO-QUINAPRIL/HCTZ in children have not been established, therefore use in this age group is not recommended.

**CONTRAINDICATIONS**

APO-QUINAPRIL/HCTZ Tablets are contraindicated in:

- Patients who are hypersensitive to the drug or to any ingredient in the formulation. For a complete listing, see the Dosage Forms, Composition and Packaging section of the product monograph.

- Patients with a history of angioedema related to previous treatment with an angiotensin converting enzyme (ACE) inhibitor (see WARNINGS AND PRECAUTIONS, **General, Angioedema**).
- Combination with sacubitril/valsartan due to increased risk of angioedema.
- Patients hypersensitive to other sulfonamide-derived drugs because of the hydrochlorothiazide component.
- Patients with anuria.
- Women who are pregnant, intend to become pregnant, or of childbearing potential who are not using adequate contraception (see WARNINGS AND PRECAUTIONS, **Special Populations, Pregnant Women** and ADVERSE REACTIONS).
- Nursing women (see WARNINGS AND PRECAUTIONS, **Special Populations, Nursing Women**).
- Combination with aliskiren-containing medicines in patients with
  - diabetes mellitus (type 1 or type 2),
  - moderate to severe kidney insufficiency ( $\text{GFR} < 60 \text{ mL/min/1.73 m}^2$ ),
  - hyperkalemia ( $> 5 \text{ mMol/L}$ ) or
  - congestive heart failure who are hypotensive (see WARNINGS AND PRECAUTIONS, **Dual blockade of the Renin-Angiotensin System (RAS)** and **Renal Impairment**, and DRUG INTERACTIONS, **Aliskiren-containing medicines and Angiotensin receptor blockers (ARBs) or other ACE inhibitors**).
- Combination with angiotensin receptor blockers (ARBs) or other ACE inhibitors in patients with
  - diabetes with end organ damage,
  - moderate to severe kidney insufficiency ( $\text{GFR} < 60 \text{ mL/min/1.73m}^2$ ),
  - hyperkalemia ( $> 5\text{mMol/L}$ ) or
  - congestive heart failure who are hypotensive (see DRUG INTERACTIONS, **Angiotensin receptor blockers (ARBs) or other ACE inhibitors**).

## WARNINGS AND PRECAUTIONS

### Serious Warnings and Precautions

**When used in pregnancy, angiotensin converting enzyme (ACE) inhibitors can cause injury or even death of the developing fetus. When pregnancy is detected, APO-QUINAPRIL/HCTZ Tablets should be discontinued as soon as possible.**

## **General**

### **Angioedema**

Angioedema has been reported in patients treated with ACE inhibitors, including quinapril. Angioedema associated with laryngeal involvement may be fatal. If laryngeal stridor or angioedema of the face, tongue, or glottis occurs, treatment with APO-QUINAPRIL/HCTZ (quinapril hydrochloride and hydrochlorothiazide) should be discontinued immediately, the patient treated appropriately in accordance with accepted medical care, and carefully observed until the swelling disappears. In instances where swelling is confined to the face and lips, the condition generally resolves without treatment, although antihistamines may be useful in relieving symptoms. Where there is involvement of the tongue, glottis or larynx, likely to cause airway obstruction, appropriate therapy (including but not limited to 0.3 to 0.5 mL of subcutaneous epinephrine solution 1:1000) should be administered promptly (see ADVERSE REACTIONS).

The incidence of angioedema during ACE inhibitor therapy has been reported to be higher in black than in non-black patients.

Patients taking a concomitant mTOR inhibitor (e.g. temsirolimus), DPP-4 inhibitor (e.g. sitagliptin) or neutral endopeptidase (NEP) inhibitor may be at increased risk for angioedema. Caution should be used when either initiating ACE inhibitor therapy in patients already taking a mTOR inhibitor, DPP-4 inhibitor or NEP inhibitor or vice versa (see DRUG INTERACTIONS).

Patients with a history of angioedema related or unrelated to ACE inhibitor therapy may be at increased risk of angioedema while receiving an ACE inhibitor (see CONTRAINDICATIONS).

### **Carcinogenesis and Mutagenesis**

#### **Non-melanoma skin cancer**

An increased risk of non-melanoma skin cancer (NMSC) [basal cell carcinoma (BCC) and squamous cell carcinoma (SCC) of the skin] after hydrochlorothiazide therapy was reported in some epidemiological studies. The risk may be higher with increasing cumulative use (see ADVERSE REACTIONS, Post Market Adverse Drug Reactions). The photosensitizing action of hydrochlorothiazide may be a possible mechanism for NMSC (see TOXICOLOGY, Carcinogenicity – Hydrochlorothiazide).

Patients taking hydrochlorothiazide should be informed of the potential risk of NMSC. They should be advised to regularly check their skin for new lesions as well as changes to existing ones, and to promptly report any suspicious skin lesions. Patients should also be advised to limit exposure to sunlight, to avoid the use of indoor tanning equipment, and to use adequate protection (e.g. a broad spectrum sunscreen with a SPF of 30 or higher, clothing, and a hat) when exposed to sunlight or UV light to minimize the risk of skin cancer.

Alternatives to hydrochlorothiazide may be considered for patients who are at a particularly high risk for NMSC (e.g., light coloured skin, known personal or family history of skin cancer, ongoing immunosuppressive therapy, etc.) (see ADVERSE REACTIONS, Post Market Adverse Drug Reactions).

## **Cardiovascular**

### **Dual blockade of the Renin-Angiotensin System (RAS)**

Dual blockade of the Renin-Angiotensin System (RAS): There is evidence that co-administration of angiotensin converting enzyme (ACE) inhibitors, such as APO-QUINAPRIL/HCTZ, or of angiotensin receptor antagonists (ARBs) with aliskiren increases the risk of hypotension, syncope, stroke, hyperkalemia and deterioration of renal function, including acute renal failure, in patients with diabetes mellitus (type 1 or type 2) and/or moderate to severe kidney insufficiency (GFR < 60 mL/min/1.73 m<sup>2</sup>). Therefore, the use of APO-QUINAPRIL/HCTZ in combination with aliskiren-containing drugs is contraindicated in these patients (see CONTRAINDICATIONS).

Further, co-administration of ACE inhibitors, including APO-QUINAPRIL/HCTZ, with other agents blocking the RAS, such as ARBs or aliskiren-containing drugs, is generally not recommended in other patients, since such treatment has been associated with an increased incidence of severe hypotension, acute renal failure, and hyperkalemia. Administration should be limited to individually defined cases with close monitoring of renal function and blood potassium levels (see CONTRAINDICATIONS).

### **Hypotension**

Symptomatic hypotension has occurred after administration of quinapril, usually after the first or second dose or when the dose was increased. It is more likely to occur in patients who are volume depleted by diuretic therapy, dietary salt restriction, dialysis, diarrhea, or vomiting. In patients with ischemic heart or cerebrovascular disease, an excessive fall in blood pressure could result in a myocardial infarction or cerebrovascular accident (See ADVERSE REACTIONS). Because of the potential fall in blood pressure in these patients, therapy with APO-QUINAPRIL/HCTZ Tablets should be started under close medical supervision. Such patients should be followed closely for the first weeks of treatment and whenever the dose of APO-QUINAPRIL/HCTZ Tablets is increased. In patients with severe congestive heart failure, with or without associated renal insufficiency, excessive hypotension has been observed and may be associated with oliguria and/or progressive azotemia, and rarely with acute renal failure and/or death.

If hypotension occurs, the patient should be placed in supine position and, if necessary, receive an intravenous infusion of 0.9% sodium chloride. A transient hypotensive response is not a contraindication to further doses which usually can be given without difficulty once the blood pressure has increased after volume expansion. If symptoms persist, the dosage should be reduced or the drug discontinued.

### **Valvular Stenosis**

There is concern on theoretical grounds that patients with aortic stenosis might be at particular risk of decreased coronary perfusion when treated with vasodilators because they do not develop as much afterload reduction.

### **Endocrine and Metabolism**

Initial and periodic determination of serum electrolytes should be performed at appropriate intervals to detect possible electrolyte imbalance.

### **Hyperkalemia/Hypokalemia**

Quinapril: Elevated serum potassium ( $>5.7$  mMol/L) was observed in approximately 2% of patients receiving quinapril. In most cases these were isolated values which resolved despite continued therapy. Hyperkalemia was a cause of discontinuation of therapy in  $<0.1\%$  of hypertensive patients. Risk factors for the development of hyperkalemia include renal insufficiency, diabetes mellitus, and the concomitant use of agents to treat hypokalemia, potassium-sparing diuretics, potassium supplements, potassium-containing salt substitutes, or other drugs known to raise serum potassium levels (see WARNINGS AND PRECAUTIONS, **Monitoring and Laboratory Tests**, **Serum Electrolytes**, **ADVERSE REACTIONS**, and **DRUG INTERACTIONS, Agents Increasing Serum Potassium, Trimethoprim-containing products**). The addition of a potassium-sparing diuretic to APO-QUINAPRIL/HCTZ Tablets, which contains a diuretic, is not recommended.

Hydrochlorothiazide: Treatment with thiazide diuretics has been associated with hypokalemia. Hypokalemia can also sensitize or exaggerate the response of the heart to the toxic effects of digitalis. The risk of hypokalemia is greatest in patients with cirrhosis of the liver, in patients experiencing a brisk diuresis, in patients who are receiving inadequate oral intake of electrolytes, and in patients receiving concomitant therapy with corticosteroids or ACTH or with other drugs known to increase the risk of hypokalemia induced by thiazide diuretics (e.g. aminoglycoside antibiotics, cisplatin, foscarnet, amphotericin B and loop diuretics (furosemide)).

Quinapril/Hydrochlorothiazide: The opposite effects of hydrochlorothiazide and quinapril on serum potassium may approximately balance each other in many patients so that no net effect will be seen. In other patients, one or the other effect may be dominant.

### **Other electrolytes imbalances**

Hydrochlorothiazide: In addition to hypokalemia, treatment with thiazide diuretics has also been associated with hyponatremia and hypochloremic alkalosis. These disturbances have sometimes been manifest as one or more of the following: dryness of mouth, thirst, weakness, lethargy, drowsiness, restlessness, muscle pain or cramps, muscular fatigue, hypotension, oliguria, tachycardia, nausea, confusion, seizures and vomiting.

Chloride deficits secondary to thiazide therapy are generally mild and require specific treatment only under extraordinary circumstances (e.g. in liver disease or renal disease). Dilutional hyponatremia may occur in edematous patients, especially in hot weather; appropriate therapy is water restriction rather than administration of salt, except when the hyponatremia is life threatening. In actual salt depletion, replacement of salt is the therapy of choice.

Thiazides may decrease calcium excretion. Thiazides may cause intermittent and slight elevation of serum calcium in the absence of known disorders of calcium metabolism. Marked hypercalcemia may be evidence of hidden hypoparathyroidism. In a few patients on prolonged thiazide therapy, pathological changes in the parathyroid gland have been observed, with hypercalcemia and hypophosphatemia. More serious complications of hyperparathyroidism (renal lithiasis, bone resorption, and peptic ulceration) have not been seen. Thiazides should be discontinued before performing tests for parathyroid function.

Thiazides increase the urinary excretion of magnesium, and hypomagnesaemia may result.

### **Hypoglycemia/Hyperglycemia and Diabetes**

Quinapril: ACE inhibitors may reduce insulin resistance and may lead to hypoglycemia in diabetic patients on insulin or oral hypoglycemic agents; closer monitoring of diabetic patients may be required.

Hydrochlorothiazide: Thiazide-induced hyperglycemia may compromise blood sugar control. Depletion of serum potassium augments glucose intolerance. Monitor glycemic control, supplement potassium, if necessary, to maintain appropriate serum potassium levels, and adjust diabetes medications as required (see DRUG INTERACTIONS, **Antidiabetic agents (e.g. insulin, oral hypoglycemic agents, sitagliptin)**).

Overt diabetes may be precipitated in susceptible individuals.

### **Other metabolic parameters**

Hyperuricemia may occur, or acute hyperuricemia may be precipitated, in certain patients receiving thiazide therapy.

Increase in cholesterol, triglyceride and glucose levels may be associated with thiazide diuretic therapy.

Thiazides may decrease serum PBI levels without signs of thyroid disturbance.

### **Hematologic**

#### **Neutropenia/Agranulocytosis**

Agranulocytosis and bone marrow depression have been caused by ACE inhibitors. Agranulocytosis did occur during quinapril treatment in one patient with a history of neutropenia during previous captopril therapy. Periodic monitoring of white blood cell counts should be considered, especially in patients with collagen vascular disease and/or renal disease.

### **Hepatic**

#### **Impairment of Liver Function**

Hepatitis (hepatocellular and/or cholestatic), elevations of liver enzymes and/or serum bilirubin have occurred during therapy with other ACE inhibitors in patients with or without pre-existing liver abnormalities. In most cases the changes were reversed on discontinuation of the drug.

Elevations of liver enzymes and/or serum bilirubin have been reported for quinapril and hydrochlorothiazide tablets (see ADVERSE REACTIONS). Should the patient receiving APO-QUINAPRIL/HCTZ Tablets experience any unexplained symptoms particularly during the first weeks or months of treatment, it is recommended that a full set of liver function tests and any other necessary investigation be carried out. Discontinuation of APO-QUINAPRIL/HCTZ Tablets should be considered when appropriate.

There are no adequate studies in patients with cirrhosis and/or liver dysfunction. APO-QUINAPRIL/HCTZ Tablets should be used with particular caution in patients with pre-existing liver



abnormalities. In such patients baseline liver function tests should be obtained before administration of the drug and close monitoring of response and metabolic effects should apply.

### **Patients with Impaired Liver Function**

APO-QUINAPRIL/HCTZ Tablets should be used with caution in patients with impaired hepatic function or progressive liver disease, since minor alterations of fluid and electrolyte balance may precipitate hepatic coma. Also, since the metabolism of quinapril to quinaprilat is normally dependent upon hepatic esterase, patients with impaired liver function could develop markedly elevated plasma levels of quinapril.

### **Immune**

#### **Anaphylactoid Reactions during Membrane Exposure**

Anaphylactoid reactions have been reported in patients dialysed with high-flux membranes (e.g.: polyacrylonitrile [PAN]) and treated concomitantly with an ACE inhibitor. Dialysis should be stopped immediately if symptoms such as nausea, abdominal cramps, burning, angioedema, shortness of breath and severe hypotension occur. Symptoms are not relieved by antihistamines. In these patients consideration should be given to using a different type of dialysis membrane or a different class of antihypertensive agent.

#### **Anaphylactoid Reactions during LDL Apheresis**

Rarely, patients receiving ACE inhibitors during low density lipoprotein apheresis with dextran sulfate have experienced life-threatening anaphylactoid reactions. These reactions were avoided by temporarily withholding the ACE inhibitor therapy prior to each apheresis.

#### **Anaphylactoid Reactions during Desensitization**

There have been isolated reports of patients experiencing sustained life threatening anaphylactoid reactions while receiving ACE inhibitors during desensitizing treatment with hymenoptera (bees, wasps) venom. In the same patients, these reactions have been avoided when ACE inhibitors were temporarily withheld for at least 24 hours, but they have reappeared upon inadvertent rechallenge to an ACE inhibitor.

#### **Hypersensitivity to Hydrochlorothiazide**

Sensitivity reactions to hydrochlorothiazide may occur in patients with or without a history of allergy or bronchial asthma.

#### **Nitritoid Reactions-Gold**

Nitritoid reactions (symptoms include facial flushing, nausea, vomiting and symptomatic hypotension) have been reported rarely in patients on therapy with injectable gold (sodium aurothiomalate) and concomitant ACE inhibitor therapy including quinapril and hydrochlorothiazide tablets (see DRUG INTERACTIONS).

#### **Systemic Lupus Erythematosus**

Thiazide diuretics have been reported to cause exacerbation or activation of systemic lupus erythematosus.

### **Ophthalmologic**

### **Acute Myopia and Secondary Angle-Closure Glaucoma related to Hydrochlorothiazide**

Hydrochlorothiazide, a sulfonamide, can cause an idiosyncratic reaction, resulting in acute transient myopia and acute angle-closure glaucoma. Symptoms include acute onset of decreased visual acuity or ocular pain and typically occur within hours to weeks of drug initiation. Untreated acute angle-closure glaucoma can lead to permanent vision loss.

The primary treatment is to discontinue hydrochlorothiazide as rapidly as possible. Prompt medical or surgical treatments may need to be considered if the intraocular pressure remains uncontrolled. Risk factors for developing acute angle-closure glaucoma may include a history of sulfonamide or penicillin allergy.

### **Peri-Operative Considerations**

#### **Surgery/Anaesthesia**

In patients undergoing major surgery or during anaesthesia with agents that produce hypotension, ACE inhibitors will block angiotensin II formation secondary to compensatory renin release. If hypotension occurs and is considered to be due to this mechanism, it can be corrected by volume expansion.

### **Renal**

#### **Azotemia**

Azotemia may be precipitated or increased by hydrochlorothiazide. Cumulative effects of the drug may develop in patients with impaired renal function. If increasing azotemia and oliguria occur, APO-QUINAPRIL/HCTZ Tablets should be discontinued.

#### **Renal Impairment**

The use of ACE inhibitors, including APO-QUINAPRIL/HCTZ Tablets, with ARBs or aliskiren-containing drugs is contraindicated in patients with moderate to severe kidney insufficiency (GFR < 60 mL/min/1.73m<sup>2</sup>) (see CONTRAINDICATIONS and DRUG INTERACTIONS, **Aliskiren-containing medicines and Angiotensin receptor blockers (ARBs) or other ACE inhibitors**).

As a consequence of inhibiting the renin-angiotensin-aldosterone system (RAAS), changes in renal function have been seen in susceptible individuals. In patients whose renal function may depend on the activity of the RAAS, such as patients with bilateral renal artery stenosis, unilateral renal artery stenosis to a solitary kidney, or severe congestive heart failure, treatment with agents that inhibit this system has been associated with oliguria, progressive azotemia, and rarely, acute renal failure and/or death. In susceptible patients, concomitant diuretic use may further increase risk (see ADVERSE REACTIONS and DOSAGE ADMINISTRATION).

Use of APO-QUINAPRIL/HCTZ Tablets should be followed by the appropriate assessment of renal function (see ADVERSE REACTIONS and DOSAGE AND ADMINISTRATION).

Thiazides may not be appropriate diuretics for use in patients with renal impairment and are ineffective at creatinine clearance values of ≤30 mL/min (i.e. moderate or severe renal insufficiency) (see ADVERSE REACTIONS and DOSAGE ADMINISTRATION).

## **Respiratory**

### **Cough**

A dry, persistent cough, which usually disappears only after withdrawal or lowering of the dose of quinapril has been reported. Such possibility should be considered as part of the differential diagnosis of the cough.

## **Skin**

### **Photosensitivity**

Photosensitivity reactions have been reported with the use of thiazide diuretics.

If photosensitivity reactions occur during treatment with hydrochlorothiazide-containing drugs, treatment should be stopped.

## **Special Populations**

### **Pregnant Women**

Quinapril is contraindicated in pregnancy (see CONTRAINDICATIONS and ADVERSE REACTIONS). ACE inhibitors can cause fetal and neonatal morbidity and mortality when administered to pregnant women. Several dozen cases have been reported in the world literature. When pregnancy is detected, APO-QUINAPRIL/HCTZ Tablets should be discontinued as soon as possible.

The use of ACE inhibitors during the second and third trimesters of pregnancy has been associated with fetal and neonatal injury including hypotension, neonatal skull hypoplasia, anuria, reversible or irreversible renal failure, and death. Oligohydramnios has also been reported, presumably resulting from decreased fetal renal function; oligohydramnios in this setting has been associated with fetal limb contractures, craniofacial deformation, and hypoplastic lung development.

Prematurity, and patent ductus arteriosus and other structural cardiac malformations, as well as neurological malformations, have also been reported following exposure in the first trimester of pregnancy.

Infants with a history of *in utero* exposure to ACE inhibitors should be closely observed for hypotension, oliguria, and hyperkalemia. If oliguria occurs, attention should be directed toward support of blood pressure and renal perfusion. Exchange transfusion or dialysis may be required as a means of reversing hypotension and/or substituting for impaired renal function; however, limited experience with those procedures has not been associated with significant clinical benefit.

If oligohydramnios is observed, a non-stress test (NST), and/or a biophysical profiling (BPP) may be appropriate, depending upon the week of pregnancy. If concerns regarding fetal well-being still persist, a contraction stress test (CST) should be considered. Patients and physicians should be aware, however, that oligohydramnios may not appear until after the fetus has sustained irreversible injury.

Thiazides cross the placental barrier and appear in cord blood. Although studies in humans have not been done, effects to the fetus may include fetal or neonatal jaundice, thrombocytopenia, and possibly other adverse reactions which have occurred in the adult.

**Animal Data:** No fetotoxic or teratogenic effects were observed in rats at quinapril doses as high as 300 mg/kg/day (180x maximum daily human dose), despite maternal toxicity at 150 mg/kg/day. Offspring body weights were reduced in rats treated late in gestation and during lactation with doses of  $\geq 25$  mg/kg/day. Quinapril hydrochloride was not teratogenic in rabbits; however, maternal and embryo toxicity were seen in some rabbits at doses as low as 0.5 mg/kg/day and 1 mg/kg/day, respectively.

No adverse effects on fertility or reproduction were observed in rats at quinapril dose levels  $\leq 100$  mg/kg/day (60x maximum daily human dose).

### **Nursing Women**

The presence of concentrations of ACE inhibitor has been reported in human milk. Thiazides also appear in human milk. The use of APO-QUINAPRIL/HCTZ Tablets is contraindicated during breast-feeding (see CONTRAINDICATIONS).

### **Pediatrics (<18 years of age)**

The safety and effectiveness of APO-QUINAPRIL/HCTZ Tablets in children have not been established, therefore use in this age group is not recommended.

### **Monitoring and Laboratory Tests**

**Serum Electrolytes:** Variations of serum electrolytes levels have been observed with quinapril and hydrochlorothiazide. Initial and periodic determination of serum electrolytes should be performed at appropriate intervals to detect possible electrolyte imbalance (See CONTRAINDICATIONS and WARNINGS and PRECAUTIONS).

**Creatinine and Blood Nitrogen:** Increases ( $>1.25$ x ULN) in serum creatinine and blood urea nitrogen (BUN) were observed in 3% and 4% respectively, of patients treated with quinapril and hydrochlorothiazide (see WARNINGS AND PRECAUTIONS and DOSAGE AND ADMINISTRATION).

**Hepatic:** Elevations of liver enzymes and/or serum bilirubin have occurred in patients receiving quinapril and hydrochlorothiazide. If a patient receiving quinapril and hydrochlorothiazide experience any unexplained symptoms, particularly during the first weeks or months of treatment, a full set of liver function tests and any other investigation should be carried out. Discontinuation of quinapril and hydrochlorothiazide should be considered when appropriate. In patients with pre-existing liver abnormalities, baseline liver function tests should be obtained before administration of the drug. The response and metabolic effects should be closely monitored (see WARNINGS AND PRECAUTIONS).

**Glucose:** Elevations in glucose values have occurred. Monitor glycemic control, supplement potassium, if necessary, to maintain appropriate serum potassium levels, and adjust diabetes medications as required (see WARNINGS AND PRECAUTIONS, **Endocrine and Metabolism**, **Hypoglycemia/Hyperglycemia and Diabetes** and DRUG INTERACTIONS, **Antidiabetic agents** (e.g. insulin, oral hypoglycemic agents, sitagliptin)).

**Triglyceride:** Elevations in triglyceride values have occurred (see WARNINGS AND PRECAUTIONS, Endocrine and Metabolism, Other metabolic parameters).

**Serum Uric Acid:** Elevations in serum uric acid values have occurred (see WARNINGS AND PRECAUTIONS, Endocrine and Metabolism, Other metabolic parameters).

**Hematology:** Possibly clinically important increases and decreases in hematology parameters have occurred. Periodic monitoring of white blood cell counts should be considered, especially in patients with collagen vascular disease and/or renal disease (see WARNINGS AND PRECAUTIONS, Hematologic, Neutropenia/Agranulocytosis).

Other laboratory test values with clinically important deviations during controlled and uncontrolled trials included: Magnesium, Cholesterol, PBI, Parathyroid Function Tests and Calcium (see WARNINGS AND PRECAUTIONS, Endocrine and Metabolism, Other electrolytes imbalances and Other metabolic parameters).

## ADVERSE REACTIONS

### Clinical Trial Adverse Drug Reactions

*Because clinical trials are conducted under very specific conditions the adverse reaction rates observed in the clinical trials may not reflect the rates observed in practice and should not be compared to the rates in the clinical trials of another drug. Adverse drug reaction information from clinical trials is useful for identifying drug-related adverse events and for approximating rates.*

Quinapril hydrochloride and hydrochlorothiazide was evaluated for safety in 1571 patients with essential hypertension, including 943 patients in controlled studies (see Table 1), 345 patients in placebo-controlled trials, and 517 patients who were treated with quinapril and hydrochlorothiazide  $\geq 1$  year. Adverse reactions were limited to those reported previously with quinapril or hydrochlorothiazide (HCTZ) when used separately for the treatment of hypertension.

Serious or clinically significant adverse reactions observed in  $<0.2\%$  of patients treated with quinapril and HCTZ were: hematemesis, gout, syncope and angioedema. Therapy was discontinued in 2.1% of patients due to an adverse event (AE). Headache (0.5%) and dizziness (0.3%) were the most frequent reasons for withdrawal.

The most frequent adverse experiences in controlled trials were headache (6.7%), dizziness (4.8%), cough (3.2%) and fatigue (2.9%). The cough is characteristically non-productive, persistent and resolves after discontinuation of therapy (see WARNINGS AND PRECAUTIONS, Respiratory, Cough).

**Table 1. Adverse Events in  $\geq 1\%$  of Quinapril/Hydrochlorothiazide Patients in Controlled Clinical Studies.**

---

<b>Adverse Events</b>	<b><u>Quinapril / HCTZ</u></b> <b>N = 943</b> <b>(% Patients)</b>	<b><u>Quinapril</u></b> <b>N = 799</b> <b>(% Patients)</b>
<b>Body as a Whole</b>		
Asthenia	1.1	1.2
Fatigue	2.9	2.0
Headache	6.7	4.8
Back pain	1.5	0.7
Chest pain	1.0	1.2
Viral infection	1.9	2.0
<b>Cardiovascular System</b>		
Vasodilatation	1.0	0.4
<b>Digestive System</b>		
Dyspepsia	1.2	1.9
Nausea and/or vomiting	1.8	2.0
Diarrhea	1.4	1.7
Abdominal pain	1.7	1.6
<b>Musculoskeletal System</b>		
Myalgia	2.4	0.9
<b>Nervous System</b>		
Dizziness	4.8	2.7
Insomnia	1.2	1.5
Somnolence	1.2	0.9
Vertigo	1.0	0.3
<b>Respiratory System</b>		
Pharyngitis	1.1	1.4
Rhinitis	2.0	3.0
Bronchitis	1.2	1.3
Coughing	3.2	2.7
Upper respiratory infection	1.3	1.1

Clinical AEs regardless of relationship to therapy, occurring in  $\geq 0.5\%$  to  $< 1.0\%$  of patients treated with quinapril plus HCTZ in controlled and uncontrolled trials and less frequent clinically significant events seen in clinical trials or in post marketing experience included:

**Cardiovascular:** Hypotension, palpitations, tachycardia  
**Gastrointestinal:** Dry mouth or throat, flatulence, pancreatitis  
**Respiratory:** Dyspnea, sinusitis

***Integumentary:*** Alopecia, erythema multiforme, exfoliative dermatitis, pemphigus, pruritus, rash  
***Nervous/Psychiatric:*** Nervousness, paresthesia  
***Urogenital:*** Impotence, urinary tract infection  
***Other:*** Arthralgia, hemolytic anemia, peripheral edema

Rare AEs, not listed above, which have been reported with either HCTZ, quinapril, or the combination include:

***Cardiovascular:*** Atrial flutter, cerebrovascular accident, heart arrest, heart failure, myocardial ischemia, necrotizing angitis, transient ischemic attack, vasodilation. Orthostatic hypotension may occur, especially in elderly patients with reduced plasma volume, and may be potentiated by alcohol, barbiturates, or narcotics

***Gastrointestinal:*** Anorexia, bloody stools, constipation, cramping, gastric irritation, GI hemorrhage, jaundice (intrahepatic cholestatic), pancreatitis, sialadenitis

***Eye Disorders:*** Acute myopia and acute angle closure glaucoma (see WARNINGS AND PRECAUTIONS, **Ophthalmologic**, **Acute myopia and acute angle closure glaucoma**)

***Respiratory:*** Respiratory distress including pneumonitis, asthma, hoarseness

***Integumentary:*** Eczema, photosensitivity, rash, Stevens Johnson syndrome, urticaria

***Nervous System:*** Amnesia, anxiety, confusion, facial paralysis, paresthesias, polyneuritis, xanthopsia

***Hematological:*** Agranulocytosis, aplastic anemia, hemolytic anemia, leukopenia, purpura, thrombocytopenia

***Urogenital:*** Dysuria, glycosuria, hematuria, impaired renal function, polyuria

***Special Senses:*** Taste disturbance, tinnitus, transient blurred vision

***Congenital and familial/genetic disorders:***  
 Fetal/neonatal injury including: anuria, hypotension, oligohydramnios, skull hypoplasia, reversible or irreversible renal failure, and death (See CONTRAINDICATIONS, WARNINGS AND PRECAUTIONS, **Pregnant Women**)

***Other:*** Allergy, anaphylactic reactions, arthritis, chill, dehydration, face edema, fever, fracture, muscle spasm, restlessness, weakness, weight increase

### **Laboratory Deviations:**

Azotemia, hyperglycemia, hyperuricemia, transient hyperlipidemia, WBC decreased

### **Post-Market Adverse Drug Reactions**

**Non-melanoma skin cancer:** Some pharmacoepidemiological studies have suggested a higher risk of squamous cell carcinoma (SCC) and basal cell carcinoma (BCC) of the skin with increasing use of hydrochlorothiazide. A systematic review and meta-analysis undertaken by Health Canada suggested, with important uncertainty, that the use of hydrochlorothiazide for several years (>3 years) could lead to:

- 122 additional cases (95% CI, from 112 to 133 additional cases) of SCC per 1000 treated patients compared with non-use of hydrochlorothiazide (meta-analysis of 3 observational studies);
- 31 additional cases (95% CI, from 24 to 37 additional cases) of BCC per 1000 treated patients compared with non-use of hydrochlorothiazide (meta-analysis of 2 observational studies).

## **DRUG INTERACTIONS**

### **Drug-Drug Interactions**

<b>Proper name</b>	<b>Reference</b>	<b>Effect</b>	<b>Clinical comment</b>
<b>Agents Affecting Sympathetic Activity</b>	---		Agents affecting sympathetic activity (e.g. ganglionic blocking agents or adrenergic neuron blocking agents) may be used with caution. Beta- adrenergic blocking drugs add some further antihypertensive effect to quinapril.
<b>Agents Increasing Serum Potassium</b>	---	Since quinapril decreases aldosterone production, elevation of serum potassium may occur.	Potassium sparing diuretics such as spironolactone, triamterene or amiloride, potassium supplements or other drugs known to raise serum potassium levels should be given with caution and with frequent monitoring of serum potassium, since they may lead to a significant increase in serum potassium. Salt substitutes which contain potassium should also be used with caution.  Since quinapril/HCTZ contains a diuretic, the addition of a potassium-sparing diuretic is not recommended.



Proper name	Reference	Effect	Clinical comment
<b>Alcohol, barbiturates or narcotics</b>	C	Potential of orthostatic hypotension may occur in the presence of hydrochlorothiazide.	Avoid alcohol, barbiturates or narcotics, especially with initiation of therapy.
<b>Aliskiren-containing medicines</b>	CT	Dual blockade of the renin-angiotensin-aldosterone system by combining an ACE inhibitor with aliskiren-containing medicines is not recommended since there is an increased risk of hypotension, syncope, stroke, hyperkalemia and changes in renal function, including renal failure.	<p>The use of APO-QUINAPRIL/HCTZ in combination with aliskiren-containing medicines is contraindicated in patients with</p> <ul style="list-style-type: none"> <li>• diabetes mellitus (type 1 or type 2),</li> <li>• moderate to severe kidney insufficiency (GFR &lt; 60 mL/min/1.73 m<sup>2</sup>),</li> <li>• hyperkalemia (&gt; 5mMol/L) or</li> <li>• congestive heart failure who are hypotensive.</li> </ul> <p>It is not recommended in other patients (see CONTRAINDICATIONS; WARNINGS AND PRECAUTIONS, <b><u>Dual blockade of the Renin-Angiotensin System (RAS)</u></b>)</p>
<b>Amphotericin B</b>	T	Amphotericin B increases the risk of hypokalemia induced by thiazide diuretics.	Monitor serum potassium level.
<b>Angiotensin receptor blockers (ARBs) or other ACE inhibitors</b>	CT	Dual blockade of the renin-angiotensin-aldosterone system by combining an ACE inhibitor with ARBs or other ACEs inhibitors is not recommended since there is an increased risk of hypotension, syncope, stroke, hyperkalemia and changes in renal function, including renal failure.	<p>The use of APO-QUINAPRIL/HCTZ in combination with ARBs or other ACE inhibitors is contraindicated in patients with</p> <ul style="list-style-type: none"> <li>• diabetes with end organ damage ,</li> <li>• moderate to severe kidney insufficiency (GFR &lt; 60 mL/min/1.73 m<sup>2</sup>),</li> <li>• hyperkalemia (&gt; 5mMol/L) or</li> <li>• congestive heart failure who are hypotensive.</li> </ul> <p>It is not recommended in other patients (see CONTRAINDICATIONS; WARNINGS AND PRECAUTIONS, <b>Cardiovascular, Dual blockade of the</b></p>

Proper name	Reference	Effect	Clinical comment
			<b>Renin-Angiotensin System (RAS)</b>
<b>Anion Exchange Resins - Bile acid sequestrants (e.g. cholestyramine)</b>	CT	Bile acid sequestrants bind thiazide diuretics in the gut and impair gastrointestinal absorption by 43-85%. Administration of thiazide 4 hours after a bile acid sequestrant reduced absorption of hydrochlorothiazide by 30 to 35%.	Give thiazide 2-4 hours before or 6 hours after the bile acid sequestrant. Maintain a consistent sequence of administration. Monitor blood pressure, and increase dose for thiazide, if necessary.
<b>Antidiabetic agents (e.g. insulin oral hypoglycemic agents, sitagliptin)</b>	CT	ACE inhibitors may reduce insulin resistance and may lead to hypoglycemia in diabetic patients on insulin or oral hypoglycemic agents.	Monitor closely diabetic patients (see WARNINGS AND PRECAUTIONS, <b><u>Endocrine and Metabolism</u></b> , <b>Hypoglycemia and Diabetes</b> ).
		Thiazide-induced hyperglycemia may compromise blood sugar control. Depletion of serum potassium augments glucose intolerance.	Monitor glycemic control, supplement potassium if necessary, to maintain appropriate serum potassium levels, and adjust diabetes medications as required (see WARNINGS AND PRECAUTIONS, <b><u>Endocrine and Metabolism</u></b> , <b>Hypoglycemia and Diabetes</b> ).
		Patients taking concomitant DPP-4 inhibitor therapy may be at increased risk for angioedema.	Caution should be used when either initiating ACE inhibitor therapy in patients already taking a DPP-4 inhibitor or <i>vice versa</i> (see WARNINGS AND PRECAUTIONS, <b><u>General</u></b> , <b>Angioedema</b> ).
<b>Anti-neoplastic</b>	C, CT	Concomitant use of thiazide diuretics may	Hematological status should be closely monitored in patients receiving this

Proper name	Reference	Effect	Clinical comment
<b>drugs, including cyclophosphamide, methotrexate and mTOR inhibitors (e.g. temsirolimus, everolimus)</b>		reduce renal excretion of cytotoxic agents and enhance their myelosuppressive effects.	combination. Dose adjustment of cytotoxic agents may be required.
		Patients taking concomitant mTOR inhibitor therapy may be at increased risk for angioedema.	Caution should be used when either initiating ACE inhibitor therapy in patients already taking an mTOR inhibitor or <i>vice versa</i> (see WARNINGS AND PRECAUTIONS, <b><u>General</u></b> , <b>Angioedema</b> ).
<b>Calcium and vitamin D supplements</b>	C	Thiazides decrease renal excretion of calcium and increase calcium release from bone.	Monitor serum calcium, especially with concomitant use of high doses of calcium supplements. Dose reduction or withdrawal of calcium and/or vitamin D supplements may be necessary.
<b>Carbamazepine</b>	C	Carbamazepine may cause clinically significant hyponatremia. Concomitant use with thiazide diuretics may potentiate hyponatremia.	Monitor serum sodium levels. Use with caution.
<b>Concomitant Diuretic Therapy</b>	---	Patients concomitantly taking ACE inhibitors and diuretics, and especially those in whom diuretic therapy was recently instituted, may occasionally experience an excessive reduction of blood pressure after initiation of therapy.	The possibility of hypotensive effects after the first dose of quinapril can be minimized by either discontinuing the diuretic or increasing the salt intake (except in patients with heart failure) prior to initiation of treatment with quinapril. If it is not possible to discontinue the diuretic, the starting dose of quinapril should be reduced and the patient should be closely observed for several hours following initial dose and until blood pressure has stabilized. (See WARNINGS, and DOSAGE AND ADMINISTRATION).

Proper name	Reference	Effect	Clinical comment
<b>Corticosteroids, and adrenocorticotrophic hormone (ACTH)</b>	T	Intensified electrolyte depletion, particularly hypokalemia, may occur when administered with hydrochlorothiazide	Monitor serum potassium, and adjust medications, as required.
<b>Digoxin</b>	CT	Thiazide-induced electrolyte disturbances, i.e. hypokalemia, hypomagnesemia, increase the risk of digoxin toxicity, which may lead to fatal arrhythmic events.	Concomitant administration of hydrochlorothiazide and digoxin requires caution. Monitor electrolytes and digoxin levels closely. Supplement potassium or adjust doses of digoxin or thiazides, as required.
<b>Drugs that alter GI motility, i.e., anti-cholinergic agents, such as atropine and prokinetic agents, such as metoclopramide, domperidone</b>	CT, T	Bioavailability of thiazide diuretics may be increased by anticholinergic agents due to a decrease in gastrointestinal motility and gastric emptying. Conversely, prokinetic drugs may decrease the bioavailability of thiazide diuretics.	Dose adjustment of thiazide may be required.
<b>Gold</b>	C	Nitritoid reactions (symptoms include facial flushing, nausea, vomiting and hypotension) have been reported rarely in patients on therapy with injectable gold (sodium aurothiomalate) and concomitant ACE inhibitor therapy including quinapril and hydrochlorothiazide	
<b>Gout medications (allopurinol, uricosurics, xanthine oxidase inhibitors)</b>	T, RC	Thiazide-induced hyperuricemia may compromise control of gout by allopurinol and probenecid. The co-	Dose adjustment of gout medications may be required.

Proper name	Reference	Effect	Clinical comment
		administration of hydrochlorothiazide and allopurinol may increase the incidence of hypersensitivity reactions to allopurinol.	
<b>Lithium</b>	CT	Thiazide diuretic agents and ACE inhibitors reduce the renal clearance of lithium and increase the risk of lithium toxicity.	Concomitant use of thiazide diuretics or ACE inhibitors with lithium is generally not recommended. If such use is deemed necessary, reduce lithium dose by 50% and monitor lithium levels closely.
<b>Neutral endopeptidase inhibitor</b>	---	Patients taking concomitant neutral endopeptidase inhibitor may be at increased risk for angioedema.	Caution should be used when either initiating ACE inhibitor therapy in patients already taking a neutral endopeptidase inhibitor or vice versa (see WARNINGS AND PRECAUTIONS, General, Angioedema).
<b>Non-Steroidal Anti-Inflammatory Drugs (NSAID) including selective cyclooxygenase-2 inhibitors (COX-2 inhibitors)</b>	CT	<p>There are two types of interaction between APO-QUINAPRIL/HCTZ and NSAIDs:</p> <p>Interaction with ACE-Inhibitor component of APO-QUINAPRIL/HCTZ: In patients who are elderly, volume-depleted (including those on diuretic therapy), or with compromised renal function, co-administration of NSAIDs, including selective COX-2 inhibitors, with ACE inhibitors, including quinapril, may result in deterioration of renal function, including</p>	If combination use is necessary, monitor renal function, serum potassium, and blood pressure closely. Dose adjustment may be required.

Proper name	Reference	Effect	Clinical comment
		<p>possible acute renal failure. These effects are usually reversible.</p> <p>The antihypertensive effect of ACE inhibitors, including quinapril may be attenuated by NSAIDs.</p> <p>Interaction with Diuretic component of APO-QUINAPRIL/HCTZ: In some patients, the administration of a NSAID agent can reduce the diuretic, natriuretic, and antihypertensive effects of loop, potassium-sparing, and thiazide diuretics.</p> <p>NSAID-induced inhibition of renal prostaglandins leading to decreases of renal blood flow, along with thiazide-induced decreases in GFR may lead to acute renal failure. Patients with heart failure may be at particular risk.</p>	
<b>Other Antihypertensive Agents</b>	CT	Hydrochlorothiazide may potentiate the action of other antihypertensive drugs (e.g. guanethidine, methyldopa, beta-blockers, vasodilators, calcium channel blockers, ACEI, ARB, ARB, and direct renin	

Proper name	Reference	Effect	Clinical comment
		inhibitors)	
<b>Pressor Amines (e.g. noradrenaline)</b>	---	Possible decreased response to pressor amines may occur in the presence of a thiazide diuretic, but is not sufficient to preclude their use.	
<b>Selective serotonin reuptake inhibitors (SSRIs, e.g. citalopram, escitalopram, sertraline)</b>	T, C	Concomitant use with thiazide diuretics may potentiate hyponatremia.	Monitor serum sodium levels. Use with caution.
<b>Sirolimus (immune-suppressant mTOR inhibitor)</b>	CT	Organ transplant patients taking concomitant sirolimus therapy may be at increased risk for angioedema.	Caution should be used when either initiating ACE inhibitor therapy in patients already taking sirolimus or <i>vice versa</i> (see WARNINGS AND PRECAUTIONS, <b>General, Angioedema</b> ).
<b>Skeletal muscle relaxants of the curare family, e.g., tubocurare</b>	C	Thiazide drugs may increase responsiveness of some skeletal muscle relaxants, such as curare derivatives.	
<b>Tetracycline</b>	---	Concomitant administration of tetracycline with quinapril reduced the absorption of tetracycline in healthy volunteers (by 28-37%) due to the presence of magnesium carbonate as an excipient in the formulation.	This interaction should be considered with concomitant use of APO-QUINAPRIL/HCTZ and tetracycline or other drugs which interact with magnesium.
<b>Topiramate</b>	CT	Additive hypokalemia. Possible thiazide-induced increase in	Monitor serum potassium and topiramate levels. Use potassium supplements, or adjust

Proper name	Reference	Effect	Clinical comment
		topiramate serum concentrations.	topiramate dose as necessary.
<b>Trimethoprim-containing products</b> (sulfamethoxazole/trimethoprim)	C	In patients who are elderly or have compromised renal function, co-administration of an ACE inhibitor with sulfamethoxazole/trimethoprim has been associated with severe hyperkalemia, likely due to the hyperkalemic effects of trimethoprim.	Quinapril/HCTZ and trimethoprim-containing products should only be co-administered with caution and with appropriate monitoring of serum potassium.

Legend: C=Case Study; RCS=Retrospective Cohort Study; CT=Clinical Trial; T=Theoretical

### **Drug-Food Interactions**

The rate and extent of quinapril absorption are diminished moderately (approximately 25 to 30%) when administered during a high-fat meal. However, no effect on quinapril absorption occurs when taken during a regular meal.

### **Drug-Herb Interactions**

Interactions with herbal products have not been established.

### **Drug-Laboratory Test Interactions**

Interactions with laboratory products/methods have not been established.

### **Drug-Lifestyle Interactions**

Lifestyle interactions have not been established.

## **DOSAGE AND ADMINISTRATION**

### **Dosing Considerations**

**Dosage must be individualized. The fixed combination is not for initial therapy. The dose of APO-QUINAPRIL/HCTZ Tablets (quinapril hydrochloride and hydrochlorothiazide) should be determined by titration of the individual components.**

Once the patient has been successfully titrated with the individual components as described below, APO-QUINAPRIL/HCTZ Tablets may be substituted if the titrated doses and dosing schedule can be achieved by the fixed combination (see INDICATIONS AND CLINICAL USE and WARNINGS AND PRECAUTIONS). In some patients a twice daily administration may be required.

Patients do not generally require HCTZ in excess of 50 mg daily, particularly when combined with other antihypertensive agents.



### **Recommended Dose and Dosage Adjustment**

***Monotherapy:*** The recommended initial dose of quinapril in patients not on diuretics is 10 mg once daily. An initial dose of 20 mg once daily can be considered for patients without advanced age, renal impairment, or concomitant heart failure and who are not volume depleted (see WARNINGS AND PRECAUTIONS, **Cardiovascular** - **Hypotension**). Dosage should be adjusted according to blood pressure (BP) response, generally at intervals of 2 to 4 weeks. A dose of 40 mg daily should not be exceeded.

In some patients treated once daily, the antihypertensive effect may diminish towards the end of the dosing interval. This can be evaluated by measuring BP just prior to dosing to determine whether satisfactory control is being maintained for 24 hours. If it is not, either 2x daily administration with the same total daily dose, or an increase in dose should be considered. If BP is not controlled with quinapril alone, a diuretic may be added. After the addition of a diuretic, it may be possible to reduce the dose of quinapril.

***Concomitant Diuretic Therapy:*** Symptomatic hypotension occasionally may occur following the initial dose of quinapril and is more likely in patients who are currently being treated with a diuretic. The diuretic should, if possible, be discontinued for 2 to 3 days before beginning therapy with quinapril to reduce the likelihood of hypotension (see WARNINGS AND PRECAUTIONS, **Cardiovascular**, **Hypotension**). If the diuretic cannot be discontinued, an initial dose of 5 mg of quinapril should be used with careful medical supervision for several hours and until BP has stabilized. The dosage of quinapril should subsequently be titrated (as described above) to the optimal response.

***Dosage Adjustment in Renal Impairment:*** For use in hemodialysis patients, see WARNINGS AND PRECAUTIONS and ADVERSE REACTIONS-Anaphylactoid Reactions during Membrane Exposure. Quinapril should be administered on days when dialysis is not performed.

Starting doses should be reduced according to the following guidelines:

<b>Creatinine Clearance (mL/min)</b>	<b>Maximum Recommended Initial Dose (mg)</b>
>60	10
30-60	5
10-30	2.5
<10	Insufficient data for dosage recommendation

Patients should subsequently have dosage titrated (as described above) to the optimal response as described under Monotherapy.

When concomitant diuretic therapy is required in patients with severe renal impairment a loop diuretic rather than a thiazide is preferred for use with quinapril. Therefore, for patients with severe renal dysfunction APO-QUINAPRIL/HCTZ Tablets are not recommended.

***Dosage in the Elderly:*** The recommended initial dosage of quinapril is 10 mg once daily (depending on renal function), followed by titration to the optimal response as described above under Monotherapy.

## OVERDOSAGE

No data are available regarding overdosage with quinapril and hydrochlorothiazide tablets or quinapril. The most likely clinical manifestation would be symptoms attributable to severe hypotension, which should be normally treated by intravenous volume expansion with 0.9% sodium chloride. Hemodialysis and peritoneal dialysis have little effect on the elimination of quinapril and quinaprilat.

The most common signs and symptoms observed for HCTZ monotherapy overdosage are those caused by electrolyte depletion (hypokalemia, hypochloremia, hyponatremia) and dehydration resulting from excessive diuresis. If digitalis has also been administered, hypokalemia may accentuate cardiac arrhythmias.

For management of a suspected drug overdose, contact your regional Poison Control Centre.

## ACTION AND CLINICAL PHARMACOLOGY

### **Mechanism of Action**

APO-QUINAPRIL/HCTZ (quinapril hydrochloride and hydrochlorothiazide) is a fixed-combination tablet which combines the antihypertensive actions of an angiotension-converting enzyme (ACE) inhibitor, quinapril hydrochloride and a diuretic, hydrochlorothiazide (HCTZ). In clinical studies, administration of this combination produced greater reductions in blood pressure (BP) than the single agents given alone.

### **Pharmacodynamics**

***Quinapril:*** Quinapril is a nonpeptide, nonsulphydryl ACE inhibitor. ACE is a peptidyl dipeptidase that catalyzes the conversion of angiotensin I to the vasoconstrictor angiotensin II. After absorption, quinapril is rapidly de-esterified to quinaprilat (quinapril diacid), its principal active metabolite. Its primary mode of action is to inhibit circulating and tissue ACE, thereby decreasing vasopressor activity and aldosterone secretion. Although the decrease in aldosterone is small, it results in a small increase in serum K<sup>+</sup> (see WARNINGS AND PRECAUTIONS). Removal of angiotensin II negative feedback on renin secretion leads to increased plasma renin activity.

ACE is identical to kininase II. Thus, quinapril may interfere with the degradation of bradykinin, a potent peptide vasodilator. However, it is not known whether this system contributes to the therapeutic effects of quinapril.

In animal studies, the antihypertensive effect of quinapril outlasts its inhibitory effect on circulating ACE. Tissue ACE inhibition more closely correlates with the duration of antihypertensive effects and this may be related to enzyme binding characteristics.

Administration of 10 to 40 mg of quinapril to patients with essential hypertension results in a reduction of both sitting and standing BP with minimal effect on heart rate. Antihypertensive activity commences within 1 hour with peak effects usually achieved by 2 to 4 hours after dosing. Achievement of maximum BP lowering effects may require 2 to 4 weeks of therapy in some patients. At the recommended doses, antihypertensive effects are maintained throughout the 24-hour dosing interval in most patients. While the dose response relationship is relatively flat, a dose of 40 mg was somewhat more effective at trough than 10 to 20 mg, and 2x daily dosing tended to give a somewhat lower BP than 1x daily dosing with the same total daily dose. The antihypertensive effect of quinapril was maintained during long-term therapy with no evidence of loss of effectiveness.

Hemodynamic assessments in patients with essential hypertension indicate that BP reduction produced by quinapril is accompanied by a reduction in total peripheral resistance and renal vascular resistance with little or no change in heart rate and cardiac index. There was an increase in renal blood flow which was not significant. Little or no change in glomerular filtration rate (GFR) or filtration fraction was observed.

***Hydrochlorothiazide:*** HCTZ acts directly on the kidney to increase excretion of sodium and chloride, and an accompanying volume of water. HCTZ also increases the excretion of potassium and bicarbonate and decreases calcium excretion.

As a result of its diuretic effect, HCTZ increases plasma renin activity, increases aldosterone secretion, decreases serum potassium, and increases urinary potassium loss. Administration of quinapril inhibits the renin-angiotensin-aldosterone axis and tends to attenuate the potassium decrease associated with HCTZ.

The mechanism underlying the antihypertensive activity of diuretics is unknown. During chronic administration peripheral vascular resistance is reduced; however, this may be secondary to changes in sodium balance.

***Quinapril/Hydrochlorothiazide:*** When quinapril and HCTZ are given together, the antihypertensive effects are approximately additive.

### **Pharmacokinetics**

***Quinapril:*** Following oral administration of quinapril, peak plasma concentrations of quinapril occur within 1 hour. Based on the recovery of quinapril and its metabolites in urine, the extent of absorption is  $\geq 60\%$ . Following absorption, quinapril is de-esterified to its major active metabolite, quinaprilat (quinapril diacid), a potent ACE inhibitor, and to minor inactive metabolites. Quinapril has an apparent half-life in plasma of approximately 1 hour. Peak plasma quinaprilat concentrations occur approximately 2 hours after an oral dose of quinapril. Quinaprilat is eliminated primarily by renal excretion and has an effective accumulation half-life of approximately 3 hours. Quinaprilat has an elimination half-life in plasma of approximately 2 hours with a prolonged terminal phase of 25 hours. Approximately 97% of either quinapril or quinaprilat circulating in plasma is bound to proteins.

The rate and extent of quinapril absorption are diminished moderately (approximately 25 to 30%) when administered during a high-fat meal. However, no effect on quinapril absorption occurs when taken during a regular meal.

Studies in rats indicate that quinapril and its metabolites do not cross the blood-brain barrier.

**Hydrochlorothiazide:** After oral administration of HCTZ, diuresis begins within 2 hours, peaks in about 4 hours, and lasts about 6 to 12 hours; the extent of absorption is approximately 50 to 80%. HCTZ is excreted unchanged by the kidney. When plasma levels have been followed for  $\geq 24$  hours, the plasma half-life has been observed to vary between 4 to 15 hours. At least 61% of the oral dose is eliminated unchanged within 24 hours. HCTZ crosses the placental but not the blood-brain barrier.

**Quinapril/Hydrochlorothiazide:** Concomitant administration of quinapril and HCTZ has little or no effect on the bioavailability or the pharmacokinetics of either drug.

### **Special Populations and Conditions**

#### **Geriatrics:**

**Quinapril:** Therapeutic effects appear to be the same for elderly ( $>65$  years of age) and younger adult patients given the same daily dosages, with no increase in AEs in elderly patients.

#### **Race:**

**Quinapril:** The antihypertensive effect of ACE inhibitors is generally lower in black than in non-black patients.

#### **Hepatic Insufficiency:**

**Quinapril:** Pharmacokinetic studies in patients with end-stage renal disease or chronic hemodialysis or continuous ambulatory peritoneal dialysis indicate that dialysis has little effect on the elimination of quinapril and quinaprilat.

The disposition of quinapril and quinaprilat in patients with renal insufficiency is similar to that in patients with normal renal function until creatinine clearance is  $\leq 60$  mL/min. With creatinine clearance  $<60$  mL/min, peak and trough quinaprilat concentrations increase, apparent half-life increases, and time to steady state may be delayed. The elimination of quinaprilat may be reduced in elderly patients ( $>65$  years) and in those with heart failure; this reduction is attributable to decrease in renal function (see DOSAGE and ADMINISTRATION). Quinaprilat concentrations are reduced in patients with alcoholic cirrhosis due to impaired de-esterification of quinapril.

### **STORAGE AND STABILITY**

Protect from moisture. Keep container tightly closed and store between 15°C and 25°C.

### **DOSAGE FORMS, COMPOSITION AND PACKAGING**

APO-QUINAPRIL/HCTZ is available as fixed combination tablets in 3 strengths of quinapril (as quinapril hydrochloride) with hydrochlorothiazide:

- 10 mg/12.5 mg: Peach, oval, scored, biconvex, film coated tablets engraved “APO” on one side and “10” bisect “12.5” on the other side.
- 20 mg/12.5 mg: Peach, round, scored, biconvex, film coated tablets engraved “APO” on one side and “20” bisect “12.5” on the other side.
- 20 mg/25 mg: Peach, round, biconvex, film coated tablets engraved “APO” on one side and “20” over “25” on the other side.

Available in bottles of 100 tablets.

Available in blister pack of 100 (10 x 10) tablets

## COMPOSITION

**Medicinal Ingredients:** Quinapril (as quinapril hydrochloride) and hydrochlorothiazide in ratios of 10 mg: 12.5 mg; 20 mg: 12.5 mg and 20 mg: 25 mg respectively.

**Nonmedicinal Ingredients:** copovidone, crospovidone, ferric oxide red, ferric oxide yellow, hydroxypropyl cellulose, hypromellose, magnesium carbonate, polyethylene glycol, titanium dioxide, zinc stearate.


## PHARMACEUTICAL INFORMATION

**Proper Name:** Quinapril Hydrochloride Acetone Solvate

**Chemical Name:**

- 1) 3-Isoquinolinecarboxylic acid, 2-[2-[[1-(ethoxycarbonyl)-3-phenylpropyl]amino]-1-oxopropyl]-1,2,3,4-tetrahydro-, monohydrochloride, [3S[2[R\*(R\*)],3R\*]], acetone solvate;
- 2) (S)-2-[(S)-N-[(S)-1-Carboxy-3-phenylpropyl]alanyl]-1,2,3,4-tetrahydro-3-isoquinolinecarboxylic acid, 1-ethyl ester, monohydrochloride, acetone solvate.

**Structural Formula:**



***Solubility:***

<u>Solvent</u>	<u>% w/v solubility at RT</u>
Water	>10
0.1N HCl	>10
Methanol	>5
95% Ethanol	>5

<u>Solvent</u>	<u>% w/v solubility at RT (cont'd)</u>
Acetone	>5
Chloroform	>5
PEG 400	>10
Propylene Glycol	>10

**Dissociation Constants:**  $pK_{a1} = 2.8$ ,  $pK_{a2} = 5.4$

**Partition Coefficients:**

<u>Medium</u>	<u>Log-P</u>
0.1N HCl	0.86
0.05M phosphate buffer, pH 2.5	0.68
0.05M phosphate buffer, pH 4.0	1.35
0.05M phosphate buffer, pH 7.4	0.33

**Melting Range:** Melts with decomposition, 108-115°C


### **Hydrochlorothiazide**

**Chemical Name:**

- 1) 2H-1,2,4-Benzothiadiazine-7-sulfonamide, 6-chloro-3,4-dihydro-1,1-dioxide
- 2) 6-chloro-3,4-dihydro-2H-1,2,4-benzothiadiazine-7-sulfonamide 1,1-dioxide
- 3) 6-chloro-3,4-dihydro-7-sulfamoyl-2H-1,2,4-benzothiadiazine,1,1-dioxide
- 4) 6-chloro-7-sulfamoyl-3,4-dihydro-1,2,4-benzothiadiazine,1,1-dioxide

**Molecular Formula:**  $C_7H_8ClN_3O_4S_2$  (M.W. = 297.74 g/mol)

**Molecular Structure:**



**Description:** Hydrochlorothiazide is a white or almost white, crystalline powder which is soluble in acetone, sparingly soluble in alcohol and methanol and dissolves in dilute solutions of alkali hydroxides.

**Dissociation Constants:**  $pK_{a1} = 7.0$

$$\text{pK}_{\text{a}2} = 9.2$$

**Melting Range:** 273-275°C

**Solubility:**

Solvent	Solubility (mg/ml) at ambient temperature
pH 7.3 Water	0.59038
pH 1.2 0.1 N HCl	0.62059
pH 2.1 0.01N HCl	0.61728
pH 1.2 SGF	0.61744
pH 2.5 Phosphate Buffer (0.05 M)	0.60475
pH 3.5 Phosphate Buffer (0.05 M)	0.60012
pH 4.5 Phosphate Buffer (0.05 M)	0.59595
pH 5.5 Phosphate Buffer (0.05 M)	0.57382
pH 6.0 Phosphate Buffer (0.05 M)	0.57753
pH 6.8 Phosphate Buffer (0.05 M)	0.58204
pH 7.2 Phosphate Buffer (0.05 M)	0.60272
pH 7.5 Phosphate Buffer (0.05 M)	0.59700

**Partition Coefficient (octanol - water):** -0.1



## CLINICAL TRIALS

### Comparative Bioavailability Studies

A randomized, single dose, double-blinded, 2-way crossover comparative bioavailability study, conducted under fasting conditions, was performed on healthy male volunteers. The rate and extent of absorption of Quinapril and Hydrochlorothiazide compared following a single oral dose (1x 20/25 mg tablet) of APO-QUINAPRIL/HCTZ tablets and Accuretic<sup>®</sup> tablets in 17 volunteers. The results from measured data of both the analytes (Quinapril and Hydrochlorothiazide) are summarized in the following tables:

<b>Summary Table of the Comparative Bioavailability Data</b> <b>Quinapril</b> <b>(A single 20 mg of Quinapril /25 mg of hydrochlorothiazide dose:</b> <b>1 x 20/25 mg tablet)</b> <b>From Measured Data/Fasting Conditions</b> <b>Geometric Mean</b> <b>Arithmetic Mean (CV%)</b>				
Parameter	APO- QUINAPRIL/HCTZ tablets (Apotex Inc.)	Accuretic <sup>®</sup> tablets, (Pfizer Canada Inc.)†	% Ratio of Geometric Means	Confidence Interval
AUC <sub>t</sub> (ng•h/mL)	312.120 325.241 (36)	308.785 315.987 (27)	101.1	90.4 – 113.0
AUC <sub>inf</sub> (ng•h/mL)	316.908 330.774 (36)	314.010 321.162 (27)	100.9	90.0 – 113.2
C <sub>max</sub> (ng/mL)	290.758 312.694(41)	260.063 270.096 (28)	111.8	92.0 - 135.9
T <sub>max</sub> § (h)	0.69 (73)	0.71 (27)		
T <sub>1/2</sub> § (h)	1.45 (21)	1.57 (31)		
§ Arithmetic means (CV %) only.				
† Accuretic <sup>®</sup> Tablets (Pfizer Canada Inc.) were purchased in Canada.				

<b>Summary Table of the Comparative Bioavailability Data</b> <b>Hydrochlorothiazide</b> <b>(A single 20 mg of Quinapril /25 mg of hydrochlorothiazide dose:</b> <b>1 x 20/25 mg tablet)</b> <b>From Measured Data/Fasting Conditions</b> <b>Geometric Mean</b> <b>Arithmetic Mean (CV%)</b>				
Parameter	APO- QUINAPRIL/HCTZ tablets (Apotex Inc.)	Accuretic <sup>®</sup> tablets, (Pfizer Canada Inc.)†	% Ratio of Geometric Means	Confidence Interval
AUC <sub>t</sub> (ng•h/mL)	989.83 1024.47 (26)	1002.98 1030.10 (25)	98.7	93.1 – 104.6
AUC <sub>inf</sub> (ng•h/mL)	1049.16 1088.26 (27)	1059.66 1088.82 (25)	99.0	93.3 - 105.1

C <sub>max</sub> (ng/mL)	151.39 156.95 (27)	141.30 145.66 (27)	107.1	98.3 – 116.7
T <sub>max</sub> § (h)	1.81 (29)	2.28 (36)		
T <sub>½</sub> § (h)	9.76 (13)	9.42 (8)		
§ Arithmetic means (CV %) only.				
†Accuretic® Tablets (Pfizer Canada Inc.) were purchased in Canada.				

## DETAILED PHARMACOLOGY

### Quinapril: Mechanism of Action

***In Vitro Studies:*** Quinapril was shown to be an inhibitor of angiotensin converting enzyme (ACE) in both plasma and tissue. In assays utilizing human plasma as sources of ACE, the diacid form of quinapril (quinaprilat) exhibited greater inhibition of ACE activity than quinapril ( $6.4 \times 10^{-10}$  M and  $8.4 \times 10^{-8}$  M, respectively). In rabbit and rat aortic strips, quinapril ( $10^{-7}$  M,  $10^{-5}$  M) specifically suppressed the contractile responses elicited by angiotensin I (50% contraction at approximately  $10^{-7}$  M and  $10^{-6}$  M angiotensin I, respectively), but had no effect on contractions induced by angiotensin II and potassium chloride.

***In Vivo Studies:*** Following oral dosing of quinapril, captopril or enalapril (0.1 to 3 mg/kg) to conscious normotensive rats, plasma ACE inhibition was assessed *in vivo* by the decrease in pressor response to intravenous (IV) angiotensin I, angiotensin II, norepinephrine and bradykinin. Quinapril produced a dose-dependent reduction (44% at 0.1 mg/kg, 81% at 0.3 mg/kg) of angiotensin I (0.32 mcg/kg IV) pressor response and potentiated the response to bradykinin (154% after 0.3 mg/kg quinapril), but had no effect on angiotensin II and norepinephrine responses. Quinapril was equipotent to captopril and enalapril, but had a longer duration of action than captopril. In the conscious dog, oral administration of quinapril (0.1 to 3 mg/kg) resulted in plasma ACE inhibition comparable to that of enalapril and captopril.

In human subjects, quinapril at single oral doses of 10 to 20 mg/day produced 95 to 100% inhibition of plasma ACE activity at 0.5 hour postdose, with >80% inhibition persisting at 24 hours postdose. Multiple oral doses of quinapril to humans for 12-weeks (20 to 80 mg/day) confirmed the inhibitory effect on plasma ACE and showed that it produces corresponding decreases in angiotensin II with significant increases in plasma renin activity. Once or 2x daily dosing did not alter the results.

### Hydrochlorothiazide

For a discussion of the pharmacology of HCTZ, consult the corresponding product monograph.

### Quinapril Hydrochloride and Hydrochlorothiazide: Antihypertensive Activity

***Animal Studies:*** In spontaneously hypertensive rats the combination of quinapril/ HCTZ administered for 3 days was more efficacious than quinapril or HCTZ monotherapy. Quinapril alone (0.3 mg/kg PO) and HCTZ (30 mg/kg PO) alone reduced blood pressure (BP) by  $\leq 17\%$  and 16%, respectively ( $p > 0.05$ ). Plasma renin activity was elevated 3x by HCTZ. The combination of quinapril/HCTZ (0.3

and 30 mg/kg PO) produced significant reductions BP on days 1 and 2 and a sustained reduction (28 to 32%) in BP on day 3.

In chronic two kidney, perinephritic hypertensive dogs neither quinapril nor enalapril had antihypertensive activity at oral doses  $\leq 10$  mg/kg. Plasma renin activity was only transiently elevated in this model. Coadministration of HCTZ (10 mg/kg PO) and quinapril (10 mg/kg PO) in HCTZ pretreated dogs (10 mg/kg for 1 day) reduced mean arterial BP by 35 mmHg. In this model, HCTZ monotherapy had no significant effect on BP over a 24-hour period.

**Clinical Studies:** Two controlled studies evaluated the efficacy and safety of quinapril and HCTZ combination therapy compared with each drug given as monotherapy in patients with essential hypertension. The combination therapy caused a statistically significant greater fall in diastolic blood pressure (DBP) than each drug given as monotherapy. In a placebo controlled study, when quinapril hydrochloride (10 mg, 40 mg) and HCTZ (12.5 mg, 25 mg) were administered alone or in combination, mean reductions in DBP (at trough) produced by quinapril monotherapy ranged from 7.3 to 10.3 mmHg, by HCTZ monotherapy from 7.2 to 11.4 mmHg, and by combination therapy from 8.2 to 14.9 mmHg. Placebo produced a mean reduction in DBP of 2.2 mmHg.

## TOXICOLOGY

### Quinapril Hydrochloride

**Acute Toxicity:** The acute oral (PO) and intravenous (IV) toxicity of quinapril are summarized in Table 2.

**Table 2: Acute Toxicity of Quinapril**

Species	Sex	Route of Administration	Median Lethal Dose(mg/kg)
Mouse	Male	PO	1492 - 2150
	Female	PO	1440 - 2005
	Male	IV	504
	Female	IV	523
Rat	Male	PO	4280
	Female	PO	3541
	Male	IV	158 - 300
	Female	IV	108 - 273
Dog	Male & Female	PO	> 400

Quinapril showed a low order of acute toxicity. Clinical signs of toxicity in both mice and rats were depression or hypoactivity, prostration and ataxia. Peak mortality occurred within 24 hours in oral studies and within 15 minutes in IV studies. Asymptomatic oral dose levels were 500 mg/kg in mice and 1000 mg/kg in rats.

In the dog study, escalating oral doses of 50 to 400 mg/kg were given over 13-consecutive days. Vomiting occurred after doses of >150 mg/kg. BPs decreased with increasing dose. At 400 mg/kg, the female had elevated creatinine and blood urea nitrogen (BUN) levels, decreased sodium and chloride levels, and granular casts in the urine. Gastric erosions and ulcers were seen in both animals and renal tubular dilatation was noted in the female.

The results of quinapril toxicity from subacute, chronic, reproductive, genetic, and carcinogenicity studies are given in Tables 3 to 7, respectively.

Table 8 summarizes the results of toxicity studies with quinaprilat, the major active metabolite of quinapril.

**Table 3: Subacute Toxicity Studies of Quinapril**

Species	Duration (Week)	No. of Animals/ Sex/Group	Route	Doses (mg/kg/day)	Results
Mouse	2	10	PO	VC <sup>1</sup> , 125, 250, 500, 750	One drug-related death at 750 mg/kg; reduced food consumption and body weight gain. MTD <sup>2</sup> about 500 mg/kg.
Mouse	13	10	PO	VC, 50, 125, 250, 500	Body weight gain suppression, decreases in heart weight, hyperplasia of juxtaglomerular apparatus (JGA). MTD between 50 and 125 mg/kg.
Rat	2	5	PO	VC, 200, 400, 800, 1200	Deaths at 400, 800, and 1200 mg/kg; salivation, reduced food consumption, body weight gain suppression, pulmonary, renal, and gastric lesions.
Rat	2	10	PO	UC <sup>1</sup> , VC, 100, 400, 800	Deaths at 400 and 800 mg/kg; respiratory signs, salivation, increased BUN, decreased RBC, Hgb, and Hct; increased liver weights, decreased heart weights; pulmonary edema and focal gastric erosions. MTD was between 400 and 800 mg/kg
Rat	13	12	PO	UC, VC, 50, 250, 500	Deaths at 250 and 500 mg/kg; salivation; slightly increased BUN, CPK, and LDH; decreased RBC, Hct, and Hgb; decreased heart weight, pulmonary and gastric lesions at ≥250 mg/kg; increased renin granules in JG cells. MTD between 50 and 250 mg/kg.
Dog	2	2	PO	VC, 25, 125, 250 (125 b.i.d.)	No deaths; emesis, mild focal erosions and inflammation of the stomach at 125 mg/kg. MTD estimated as 250 mg/kg.
Dog	13	3	PO	VC, 25, 125, 250 (125 b.i.d.)	Sporadic emesis and anorexia; mild to moderate reversible increase in BUN and mild depression of RBC, Hct, Hgb at 250 mg/kg; focal gastric erosions at ≥125 mg/kg, increases in renin granules in JG cells; hypertrophy and hyperplasia of JGA. MTD was between 25 and 125 mg/kg.

<sup>1</sup> VC = Vehicle Control; UC = Untreated Control.

<sup>2</sup> MTD = Maximum Tolerated Dose; RBC = red blood cell count; Hgb = hemoglobin; Hct = hematocrit; CPK = creatine phosphokinase; LDH = lactate dehydrogenase; JG = juxtaglomerular apparatus; BUN = blood urea nitrogen.

**Table 4: Chronic Toxicity Studies of Quinapril**

<b>Species</b>	<b>Duration (Week)</b>	<b>No. of Animals/ Sex/Group</b>	<b>Route</b>	<b>Doses (mg/kg/day)</b>	<b>Results</b>
Rat	57 <sup>1</sup>	30	PO	UC <sup>2</sup> , VC <sup>2</sup> , 10, 50, 100	No drug-related deaths; transient post-dose salivation, body weight gain suppression, increased BUN, decreased glucose, increased plasma renin level, decreased heart weight, JGA hypertrophy and hyperplasia with increased granules; degenerative changes in kidneys.
Dog	52	4	PO	VC, 10, 50, 100	No deaths; elevation of plasma renin and liver enzyme levels, focal areas of chronic active inflammation in the liver at 100 mg/kg; gastric erosion at 50 mg/kg, and hypertrophy/hyperplasia of renal JGA.

<sup>1</sup> 52 weeks treatment plus 4 weeks without treatment for some animals

<sup>2</sup> UC = Untreated Control; VC = Vehicle Control; BUN = blood urea nitrogen; JGA = juxtaglomerular apparatus.

**Table 5: Reproductive Toxicology Studies of Quinapril**

Species	No. of Animals/ Sex/Group	Route	Doses (mg/kg/day)	Duration of Dosing	Results
<u>Fertility:</u>					
Rat	12 Male 24 Female	PO	VC <sup>1</sup> , 10, 50, 100	<u>Males</u> -60 days prior to mating <u>Females</u> -14 days prior to mating until weaning of offspring	No effects on fertility, no adverse effects on F <sub>1</sub> offspring parameters, and no teratogenic effects.
<u>Teratology:</u>					
Rat	5 Female	PO	100, 200, 400, 600, 800	Days 6 to 15 of gestation	No teratogenicity. Maternal deaths at 600 and 800 mg/kg; decreased fetal body weights at ≥200 mg/kg.
Rat	20 Female	PO	Uco, VC, 50, 150, 300	Days 6 to 15 of gestation	No fetotoxic or teratogenic effects. Reversible maternal toxicity.
Rabbit	5-7 Female	PO	10, 15, 25, 50, 100, 200, 400	Days 6 to 18 of gestation	Severe materno- and fetotoxicity.
Rabbit	5 Female	PO	VC, 1, 2, 4, 6, 8	Days 6 to 18 of gestation	Abortions and maternal deaths at 4, 6, and 8 mg/kg; materno- and fetotoxicity at doses >1 mg/kg.
Rabbit	14 Female	PO	VC, 0.5, 1.0, 1.5	Days 6 to 18 of gestation	Not teratogenic. Maternal weight loss; increased incidence of postimplantation loss (embryotoxicity) at 1.0 and 1.5 mg/kg.
<u>Perinatal/ Postnatal:</u>					
Rat	20 Female	PO	VC, 25, 75, 150	Day 15 of gestation to Day 20 of lactation	Reduction in offspring body weights from birth to Day 21 postnatally at 25, 75, and 150 mg/kg.

<sup>1</sup> UC = Untreated Control; VC = Vehicle Control

**Table 6: Genetic Toxicology Studies of Quinapril**

Test		Dosage Range	Results	
<u>Mutagenicity</u>				
1) <i>In Vitro</i>	a)	Initial cytotoxicity in <i>Salmonella</i> strain	≤10,000 mcg/plate	Non-cytotoxic.
	b)	Mutagenesis assay in <i>Salmonella</i>	625 to 10,000 mcg/plate	Negative-with or without metabolic activation.
2) <i>In Vitro</i>	a)	Initial cytotoxicity assay	≤44,300 mcg/mL	Cytotoxic at ≥1400 mcg/mL.
	b)	Point mutation assay in Chinese hamster lung cells	175 - 1400 mcg/mL	Negative - did not manifest direct acting or promutagen activity.
<u>Cytogenetics</u>				
1) <i>In Vitro</i>	a)	Initial cytotoxicity assay	≤44,300 mcg/mL	Cytotoxic at concentrations >700 mcg/mL.
	b)	Sister chromatid exchange (SCE) assay in Chinese hamster ovary cells	10.94 - 1400 mcg/mL	No increase in SCE at toxicity-limited doses ≤700 mcg/mL in the presence of metabolic activation or ≤1400 mcg/mL in the absence of metabolic activation.
2) <i>In Vitro</i>	a)	Initial cytotoxicity assay	≤2700 mcg/mL	Cytotoxic at ≥1200 mcg/mL.
	b)	Structural chromosomal aberration (SCA) assay in Chinese hamster lung cells	800 -1800 mcg/mL	Slight, statistically significant increase in SCA with metabolic activation; not considered biologically significant.
3) <i>In Vivo</i>	a)	Mouse micronucleus assay	1 - 1430 mcg/kg	Not clastogenic; no increased frequency of micronuclei.



**Table 7: Carcinogenicity Studies of Quinapril**

<b>Species</b>	<b>Duration (Week)</b>	<b>No. of Animals/ Sex/Group</b>	<b>Route</b>	<b>Doses (mg/kg/day)</b>	<b>Results</b>
Mouse	104	50	PO	UC <sup>1</sup> , VC <sup>1</sup> , 5, 35, 75	No evidence of tumorigenic potential. Reduced heart weight, nephritis, and JGA hypertrophy/hyperplasia.
Rat	104	65	PO	UC, VC, 10, 50, 100	No evidence of tumorigenic potential. Reduced RBC, JGA hypertrophy/hyperplasia and renal degenerative changes.

<sup>1</sup> UC = Untreated Control; VC= Vehicle Control; JGA = juxtaglomerular apparatus; RBC = red blood cell count

**Table 8: Toxicity Studies of Quinaprilat**

Species	Duration (Week)	No. of Animals/ Sex/Group	Route	Doses (mg/kg/day)	Results
<b>A. Acute Studies:</b>					
Mouse	Single-dose	10	IV	VC <sup>1</sup> , 250, 500, 1000	No deaths; MLD >1000 mg/kg. No clinical or gross pathological changes.
Rat	Single-dose	10	IV	VC, 50, 100, 200, 300, 400	No deaths; MLD >400 mg/kg. No clinical or gross pathological changes.
Dog	Escalating doses	1	IV	Escalating; 1-240	No deaths; MLD >240 mg/kg. Reduced food consumption, weight loss, and slight increase in myeloid to erythroid ratio.
<b>B. Subacute Studies:</b>					
Rat	2	5	IV	VC, 25, 50, 100, 200	No deaths, clinical signs or adverse pathological findings.
Rat	4	10	IV	VC, 20, 100, 200	No drug-related deaths or clinical signs; reduced heart weights.
Dog	2	1	IV	VC, 10, 50, 100	Sporadic increases in heart rate.
Dog	4	3	IV	VC, 10, 50, 100	No clinical or gross pathologic findings; JGA hypertrophy/hyperplasia.

<sup>1</sup> VC = Vehicle Control; MLD = median lethal dose; JGA = juxtaglomerular apparatus

**C. Genotoxicity Studies:**

Test		Dosage Range		Results
<u>Mutagenicity:</u>				
<i>In Vitro</i>	a)	Initial cytotoxicity in	<i>Salmonella</i> ≤1200 mcg/plate	Non-cytotoxic.
	b)	Mutagenesis assay in	<i>Salmonella</i> 75 - 1200 mcg/plate	Negative-with or without metabolic activation.

## **Quinapril Hydrochloride and Hydrochlorothiazide**

The 14 day Median Lethal Dose (MLD) in mice was 1068/667 mg/kg quinapril/HCTZ; in rats it was 4640/2896 mg/kg. For quinapril alone, the oral MLD ranged from 1440 to 2150 mg/kg for mice, and from 3531 to 4280 mg/kg in rats. In dogs, no drug-related clinical signs of toxicity were observed at doses of 125 to 250x the maximum human dose of quinapril given in combination with hydrochlorothiazide, and 60 to 120x the maximum human dose of hydrochlorothiazide in combination (100/60 mg/kg in males, 200/120 mg/kg in females).

## **Hydrochlorothiazide**

According to the experimental data available, hydrochlorothiazide revealed inconsistent evidence of carcinogenic activity in rats and mice, with conflicting evidence of hepatic adenoma in male mice at the highest dose and adrenal pheochromocytoma in one rat study but not in another. Current evidence is inadequate to draw a clear conclusion for a carcinogenic effect of hydrochlorothiazide in animals.

The mutagenic potential was assessed in a series of in vitro and in vivo test systems. While some positive results were obtained in vitro, all in vivo studies provided negative results. Hydrochlorothiazide enhanced the UVA-induced formation of pyrimidine dimers in vitro and in the skin of mice following oral treatment. It is therefore concluded that although there is no relevant mutagenic potential in vivo, hydrochlorothiazide could enhance the genotoxic effects of UVA light. This mechanism of photosensitization could be associated with a higher risk for non-melanoma skin cancer.

## REFERENCES

1. Ferry J, Horvath A, Sedman A, et al. Influence of food on the pharmacokinetics of quinapril and its active diacid metabolite, CI-928. *J Clin Pharmacol* 1987;27:397-399.
2. Ferry J, Cetnarowski A, Sedman A, et al. Multiple-Dose Cimetidine Administration Does Not Influence the Single-Dose Pharmacokinetics of Quinapril and Its Active Metabolite (CI-928). *J Clin Pharmacol* 1988;28:48-51.
3. Fabris B, Chen B, Pupic V, et al. Inhibition of angiotensin-converting enzyme (ACE) in plasma and tissue. *J Cardiovasc Pharmacol* 1990;15(Suppl 2):S6-S13.
4. Frank G. Overview of the clinical development of quinapril. *Clin Cardiol* 1990; 13(Suppl 7):13-18.
5. Frishman W. The safety and efficacy of quinapril in the treatment of mild to moderate essential hypertension. *Clin Cardiol* 1990;13(Suppl 7):19-25.
6. Goldstein R. The treatment of moderate to severe hypertension with ACE inhibitors. *J Cardiovasc Pharmacol* 1990;15(Suppl 2):S29-S35.
7. Gupta R, Kjeldsen S, Krause L, et al. Hemodynamic effects of quinapril a novel angiotensin-converting enzyme inhibitor. *Clin Pharmacol Ther* 1990;48:41-49.
8. Kaplan H, Taylor D, Olson S. Quinapril: Overview of preclinical data. *Clin Cardiol* 1990;13(Suppl 7):4-12.
9. Knapp L, Frank G, McLain R, et al. The safety and tolerability of quinapril. *J Cardiovasc Pharmacol* 1990;15(Suppl 2); S47-S55.
10. Schnaper H. The management of hypertension in older patients. *J Cardiovasc Pharmacol* 1990;15(Suppl 2):S56-S61.
11. Swartz R, Stermann B, Horvath A, et al. Pharmacokinetics of Quinapril and its active metabolite quinaprilat during continuous ambulatory peritoneal dialysis. *J Clin Pharmacol* 1990;30:1136-41.
12. Taylor S. The treatment of mild to moderate hypertension with ACE inhibitors. *J Cardiovasc Pharmacol* 1990;15(Suppl 2):S24-S28.
13. Verresen L, Waer M, Vanrenterhem Y, et al. Angiotensin converting enzyme inhibitors and anaphylactoid reactions to high-flux membrane dialysis. *Lancet* 1990;336:1360-1362.
14. Wadworth A, Brogden R. Quinapril: A review of its pharmacological properties and therapeutic efficacy in cardiovascular disorders. *Drugs* 1991; 41:378-399.

15. Halstenson C, Opsahl J, Rachael K, et al. The pharmacokinetics of quinapril and its active metabolite, quinaprilat, in patients with various degrees of renal function. *J Clin Pharmacol* 1992;32:344-350.
16. Canter D, Frank G, et al. Quinapril and hydrochlorothiazide combination for control of hypertension: assessment by factorial design. *J Human Hypertension* 1994;8:155-162.
17. Lenz T, Schulte K, et al. Quinapril, hydrochlorothiazide, and combination in patients with moderate to severe hypertension. *Eur Heart J* 1994;15:940-946.
18. Product Monograph – ACCURETIC\* (Quinapril Hydrochloride and Hydrochlorothiazide tablets) film coated scored tablets. Pfizer Canada ULC, Licensee. Submission control no. 224313, Date of revision: March 20, 2019.

## PART III: CONSUMER INFORMATION

### <sup>Pr</sup>APO-QUINAPRIL/HCTZ Quinapril and Hydrochlorothiazide Tablets Apotex Standard

Read this carefully before you start taking APO-QUINAPRIL/HCTZ and each time you get a refill. This leaflet is a summary and will not tell you everything about APO-QUINAPRIL/HCTZ. Talk to your doctor, nurse or pharmacist about your medical condition and treatment and ask if there is any new information about APO-QUINAPRIL/HCTZ.

#### ABOUT THIS MEDICATION

##### What the medication is used for:

APO-QUINAPRIL/HCTZ lowers high blood pressure.

##### What it does:

APO-QUINAPRIL/HCTZ contains a combination of 2 drugs, quinapril hydrochloride and hydrochlorothiazide:

- Quinapril hydrochloride is an angiotensin converting enzyme (ACE) inhibitor. You can recognize ACE inhibitors because their medicinal ingredient ends in “PRIL”. It lowers blood pressure.
- Hydrochlorothiazide is a diuretic or “water pill” that increases urination. This lowers blood pressure.

This medicine does not cure high blood pressure. It helps to control it. Therefore, it is important to continue taking APO-QUINAPRIL/HCTZ regularly even if you feel fine.

##### When it should not be used:

Do not take APO-QUINAPRIL/HCTZ if you:

- Are allergic to quinapril hydrochloride or hydrochlorothiazide or to any non-medicinal ingredients in the formulation
- Are allergic to any sulfonamide-derived drugs (sulfa drugs); most of them have a medicinal ingredient that ends in “-MIDE”
- Have experienced an allergic reaction (angioedema) with swelling of the hands, feet, or ankles, face, lips, tongue, throat, or sudden difficulty breathing or swallowing, to any ACE inhibitor or without a known cause. Be sure to tell your doctor, nurse, or pharmacist that this has happened to you
- Have been diagnosed with hereditary angioedema: an increased risk of getting an allergic reaction that is passed down through families. This can be triggered by different factors, such as surgery, flu, or dental procedures
- Are taking Entresto (sacubitril/valsartan), due to the increased risk of serious allergic reaction which causes swelling of the face or throat (angioedema) when taken with APO-QUINAPRIL/HCTZ.

- Have difficulty urinating or produce no urine
- Are pregnant or intend to become pregnant. Taking APO-QUINAPRIL/HCTZ during pregnancy can cause injury and even death to your baby
- Are breastfeeding. APO-QUINAPRIL/HCTZ passes into breast milk.
- Are taking aliskiren-containing medicines, such as Rasilez, and have one of the following conditions:
  - Diabetes
  - Kidney disease
  - High levels of potassium
  - Congestive heart failure combined with hypotension.
- Are taking an angiotensin receptor blocker (ARB), another medicine to treat your high blood pressure, or another ACE inhibitor and have one of the following conditions:
  - Diabetes with end organ damage
  - Kidney disease
  - High levels of potassium
  - Congestive heart failure combined with hypotension.

You can recognize ARBs because their medicinal ingredient ends in “-SARTAN”.

##### What the medicinal ingredients are:

- Quinapril (as quinapril hydrochloride).  
Hydrochlorothiazide

##### What the non-medicinal ingredients are:

Copovidone, crospovidone, ferric oxide red, ferric oxide yellow, hydroxypropyl cellulose, hypromellose, magnesium carbonate, polyethylene glycol, titanium dioxide, zinc stearate

##### What dosage forms it comes in:

Tablets:

10 mg quinapril (as quinapril hydrochloride) and 12.5 mg hydrochlorothiazide

20 mg quinapril (as quinapril hydrochloride) and 12.5 mg hydrochlorothiazide

20 mg quinapril (as quinapril hydrochloride) and 25 mg hydrochlorothiazide

#### WARNINGS AND PRECAUTIONS

##### **Serious Warnings and Precautions - Pregnancy**

APO-QUINAPRIL/HCTZ should not be used during pregnancy. If you discover that you are pregnant while taking APO-QUINAPRIL/HCTZ, stop the medication and contact your doctor, nurse, or pharmacist as soon as possible.

**Before you use APO-QUINAPRIL/HCTZ, talk to your doctor, nurse or pharmacist if you:**

- Are allergic to any drug used to lower blood pressure or penicillin
- Have recently received or are planning to get allergy shots for bee or wasp stings
- Have narrowing of an artery or a heart valve
- Have had a heart attack or stroke
- Have heart failure
- Have diabetes, liver or kidney disease
- Have lupus or gout
- Are on dialysis or receiving LDL apheresis (treatment to remove “bad cholesterol” from the blood)
- Are dehydrated or suffer from excessive vomiting, diarrhea, or sweating
- Are taking a salt substitute that contains potassium, potassium supplements, or a potassium-sparing diuretic (a specific kind of “water pill”)
- Are taking an antibiotic containing trimethoprim
- Are on a low-salt diet
- Are receiving gold (sodium aurothiomalate) injections
- Are less than 18 years old
- Are taking a neutral endopeptidase inhibitor. The combination with APO-QUINAPRIL/HCTZ is not recommended.
- Are taking an aliskiren-containing medicine such as Rasilez, used to lower high blood pressure. The combination with APO-QUINAPRIL/HCTZ is not recommended.
- Are taking an angiotensin receptor blocker (ARB). You can recognize an ARB because its medicinal ingredient ends in “-SARTAN”
- Are taking a medicine that contains aliskiren, such as Rasilez, an angiotensin receptor blocker (ARB), or another ACE inhibitor (in addition to APO-QUINAPRIL/HCTZ. The combination with APO-QUINAPRIL/HCTZ is not recommended.
- Are currently taking anti-cancer (temsirolimus, everolimus), anti-rejection (sirolimus) or anti-diabetic (gliptins) drugs. Use of ACE inhibitors, such as APO-QUINAPRIL/HCTZ, with these drugs may increase the chance of having an allergic reaction.
- Have had skin cancer or have a family history of skin cancer.
- Have a greater chance of developing skin cancer because you have light-coloured skin, get sunburned easily, or are taking drugs to suppress your immune system.

**Risk of skin cancer:**

- APO-QUINAPRIL/HCTZ contains hydrochlorothiazide. Treatment with hydrochlorothiazide may increase the risk of developing non-melanoma skin cancer. The risk is higher if you have been taking APO-QUINAPRIL/HCTZ for many years (more than 3) or at a high dose.
- While taking APO-QUINAPRIL/HCTZ:
  - Make sure to regularly check your skin for any new lesions. Check areas that are most exposed to the sun, such as the face, ears, hands, shoulders, upper chest and back.
  - Limit your exposure to the sun and to indoor tanning. Always use sunscreen (SPF-30 or higher) and wear protective clothing when going outside.
  - Talk to your doctor immediately if you get more sensitive to the sun or UV light or if you develop an unexpected skin lesion (such as a lump, bump, sore, or patch) during the treatment.

**Hydrochlorothiazide in APO-QUINAPRIL/HCTZ can cause sudden eye disorders:**

- **Myopia:** sudden nearsightedness or blurred vision
- **Glaucoma:** an increased pressure in your eye, eye pain. Untreated, it may lead to permanent vision loss

These eye disorders are related and can develop within hours to weeks of starting APO-QUINAPRIL/HCTZ.

You may become sensitive to the sun while taking APO-QUINAPRIL/HCTZ. Exposure to sunlight should be minimized until you know how you respond.

If you are going to have surgery and will be given an anesthetic. Be sure to tell your doctor or dentist that you are taking APO-QUINAPRIL/HCTZ.

**Driving and using machines:** before you perform tasks which may require special attention, wait until you know how you respond to APO-QUINAPRIL/HCTZ. Dizziness, lightheadedness, or fainting can especially occur after the first dose and when the dose is increased.

**INTERACTIONS WITH THIS MEDICATION**

As with most medicines, interactions with other drugs are possible. Tell your doctor, nurse, or pharmacist about the medicines you take, including drugs prescribed by other doctors, vitamins, minerals, natural supplements, or alternative medicines.

The following may interact with APO-QUINAPRIL/HCTZ:

- Adrenocorticotrophic hormone (ACTH) used to treat West Syndrome
- Alcohol, narcotics (strong pain medications) or barbiturates (sleeping pills). They may cause low blood pressure and dizziness when you go from lying down or sitting to standing up.
- Amphotericin B, an antifungal drug
- Anti-cancer drugs, including cyclophosphamide, methotrexate, temsirolimus and everolimus.
- Anti-rejection drugs, such as sirolimus (Rapamune)
- Antidepressants, in particular selective serotonin reuptake inhibitors (SSRIs), including citalopram, escitalopram, and sertraline
- Anti-diabetic drugs including insulin and oral medicines (e.g. metformin, gliptins, sulfonylureas)
- Bile acid resins used to lower cholesterol
- Calcium or vitamin D supplements
- Corticosteroids used to treat joint pain and swelling
- Digoxin, a heart medication
- Drugs known to increase the potassium level in the blood such as a salt substitute that contains potassium, potassium supplements, potassium-sparing diuretic (a specific kind of “water pill”) (e.g. spironolactone, triamterene, amiloride, sulfamethoxazole/trimethoprim).
- Drugs that slow down or speed up bowel function, including atropine, metoclopramide, and domperidone
- Drugs used to treat epilepsy, including carbamazepine and topiramate
- Gold for the treatment of rheumatoid arthritis
- Gout medications, including allopurinol and probenecid
- Lithium used to treat bipolar disease
- Non-steroidal anti-inflammatory drugs (NSAIDs) used to reduce pain and swelling. Examples include ibuprofen, naproxen, and celecoxib
- Blood pressure lowering drugs, including diuretics (“water pills”), aliskiren-containing products (e.g. Rasilez), angiotensin receptor blockers (ARBs) or other ACE inhibitors (in addition to APO-QUINAPRIL/HCTZ).
- Pressor amines (drugs which increase blood pressure, such as adrenaline)
- Skeletal muscle relaxants used to relieve muscle spasms, including tubocurane
- Tetracycline (a type of antibiotic)

#### PROPER USE OF THIS MEDICATION

APO-QUINAPRIL/HCTZ is not for initial therapy. You must first be stabilized on the individual medicinal ingredients (quinapril hydrochloride and

hydrochlorothiazide) of APO-QUINAPRIL/HCTZ. If your dosage matches the dosages in APO-QUINAPRIL/HCTZ, your doctor may prescribe APO-QUINAPRIL/HCTZ taken once a day (instead of each medicinal ingredient as a separate pill).

Take APO-QUINAPRIL/HCTZ exactly as prescribed. It is recommended to take your dose at about the same time every day.

APO-QUINAPRIL/HCTZ can be taken with or without food. If APO-QUINAPRIL/HCTZ causes upset stomach, take it with food or milk.

#### Usual Adult dose:

The recommended starting dose is one 10 mg/12.5 mg tablet daily.

#### Overdose:

If you think you have taken too much APO-QUINAPRIL/HCTZ, contact your healthcare professional, hospital emergency department or regional Poison Control Centre immediately, even if there are no symptoms.

#### Missed Dose:

If you have forgotten to take your dose during the day, carry on with the next one at the usual time. Do NOT double dose.

#### SIDE EFFECTS AND WHAT TO DO ABOUT THEM

Side effects may include:

- Dizziness, headache, trouble sleeping
- Drowsiness, fatigue, weakness
- Cough, stuffy and runny nose
- Rash, itching
- Abdominal pain, upset stomach, decreased appetite, constipation,
- Muscle pain, spasms, back pain, restlessness
- Pins and needles in your fingers
- Nausea, vomiting, diarrhea
- Sore throat
- Stuffy, runny nose
- Reduced libido
- Rash, red patches on the skin

**If any of these affects you severely, tell your doctor, nurse or pharmacist.**

APO-QUINAPRIL/HCTZ can cause abnormal blood test results. Your doctor will decide when to perform blood tests and will interpret the results.



SERIOUS SIDE EFFECTS, HOW OFTEN THEY HAPPEN AND WHAT TO DO ABOUT THEM				
Symptom / effect		Talk to your healthcare professional		Stop taking drug and get immediate medical help
		Only if severe	In all cases	
Common	<b>Low blood pressure:</b> dizziness, fainting, lightheadedness May occur when you go from lying or sitting to standing up	✓		
	<b>Decreased or increased levels of potassium in the blood:</b> irregular heartbeats, muscle weakness and generally feeling unwell		✓	
Common	<b>Non-melanoma skin cancer:</b> lump or discoloured patch on the skin that stays after a few weeks and slowly changes. Cancerous lumps are red/pink and firm and sometimes turn into ulcers. Cancerous patches are usually flat		✓	

SERIOUS SIDE EFFECTS, HOW OFTEN THEY HAPPEN AND WHAT TO DO ABOUT THEM				
Symptom / effect		Talk to your healthcare professional		Stop taking drug and get immediate medical help
		Only if severe	In all cases	
	and scaly.			
Uncommon	<b>Allergic reaction including angioedema:</b> rash, hives. Swelling of the face, lips, tongue or throat, difficulty swallowing or breathing			✓
	<b>Kidney disorder:</b> decreased urination, nausea, vomiting, swelling of extremities, fatigue		✓	
	<b>Liver disorder:</b> yellowing of the skin or eyes, dark urine, abdominal pain, nausea, vomiting, loss of appetite		✓	
Uncommon	<b>Increased blood sugar:</b> frequent urination, thirst and hunger	✓		
	<b>Electrolyte imbalance:</b> Weakness, drowsiness, muscle pain or cramps,		✓	

SERIOUS SIDE EFFECTS, HOW OFTEN THEY HAPPEN AND WHAT TO DO ABOUT THEM				
Symptom / effect		Talk to your healthcare professional		Stop taking drug and get immediate medical help
		Only if severe	In all cases	
	irregular heartbeat			
Rare	<b>Decreased platelets:</b> Bruising, bleeding, fatigue and weakness		✓	
	<b>Decreased white blood cells:</b> Infections, fatigue, fever, aches, pains and flu-like symptoms		✓	
	<b>Edema:</b> Swelling of the hands, ankles or feet		✓	
	<b>Vomiting blood</b>			✓
	<b>High nitrogen compound found in blood (Azotemia):</b> rapid heart rate, high blood pressure, fatigue, confusion, lightheadedness, dizziness, decreased urine production			✓
	<b>Chest Pain</b>			✓
	<b>Heart Attack</b>			✓

SERIOUS SIDE EFFECTS, HOW OFTEN THEY HAPPEN AND WHAT TO DO ABOUT THEM				
Symptom / effect		Talk to your healthcare professional		Stop taking drug and get immediate medical help
		Only if severe	In all cases	
Very rare	<b>Toxic epidermal necrolysis:</b> Severe skin peeling, especially in the mouth and eyes			✓
Unknown	<b>Eye disorders:</b> <b>Myopia:</b> sudden near sightedness or blurred vision <b>Glaucoma:</b> Increased pressure in your eyes, eye pain			✓
	<b>Anemia:</b> fatigue, loss of energy, weakness, shortness of breath		✓	
	<b>Inflammation of the pancreas:</b> Abdominal pain that lasts and gets worse when you lie down, nausea, vomiting		✓	
	<b>Tachycardia:</b> Fast heart beats		✓	

This is not a complete list of side effects. For any unexpected effects while taking APO-QUINAPRIL/HCTZ, contact your doctor, nurse or pharmacist.

## HOW TO STORE IT

Protect from moisture. Keep container tightly closed and store between 15°C and 25°C. Keep APO-QUINAPRIL/HCTZ out of the reach and sight of children

## Reporting Side Effects

You can report any suspected side effects associated with the use of health products to Health Canada by:

- Visiting the Web page on Adverse Reaction Reporting (<https://www.canada.ca/en/health-canada/services/drugs-health-products/medeffect-canada/adverse-reaction-reporting.html>) for information on how to report online, by mail or by fax; or By calling 1-866-234-2345 (toll-free);
- Calling toll-free at 1-866-234-2345.

*NOTE: Contact your health professional if you need information about how to manage your side effects. The Canada Vigilance Program does not provide medical advice.*

## MORE INFORMATION

### If you want more information about APO-QUINAPRIL/HCTZ:

- Talk to your healthcare professional
- Find the full product monograph that is prepared for healthcare professionals and includes this Consumer Information by visiting the Health Canada website (<https://health-products.canada.ca/dpd-bdpp/index-eng.jsp>). Find the Consumer Information on the manufacturer's website (<http://www.apotex.ca/products>), or by calling 1-800-667-4708.

This leaflet was prepared by Apotex Inc., Toronto, Ontario, M9L 1T9

Last revised: June 10, 2019