

PRODUCT MONOGRAPH
INCLUDING PATIENT MEDICATION INFORMATION

^NACT OXYCODONE CR

Oxycodone Hydrochloride Controlled Release Tablets

5 mg, 10 mg, 20 mg, 40 mg and 80 mg oxycodone hydrochloride

Teva Standard

Opioid Analgesic

Teva Canada Limited
30 Novopharm Court,
Toronto, Ontario
Canada M1B 2K9
www.tevacanada.com

Date of Revision:
October 1, 2019

Submission Control No: 230130

Table of Contents

PART I: HEALTH PROFESSIONAL INFORMATION.....	3
SUMMARY PRODUCT INFORMATION	3
INDICATIONS AND CLINICAL USE.....	3
CONTRAINDICATIONS	4
WARNINGS AND PRECAUTIONS.....	4
ADVERSE REACTIONS.....	14
DRUG INTERACTIONS	17
DOSAGE AND ADMINISTRATION	19
OVERDOSAGE	23
ACTION AND CLINICAL PHARMACOLOGY	24
STORAGE AND STABILITY	29
DOSAGE FORMS, COMPOSITION AND PACKAGING	29
PART II: SCIENTIFIC INFORMATION	30
PHARMACEUTICAL INFORMATION.....	30
CLINICAL TRIALS.....	31
DETAILED PHARMACOLOGY	40
TOXICOLOGY	40
REFERENCES	42
PATIENT MEDICATION INFORMATION	44

^NACT OXYCODONE CR

Oxycodone Hydrochloride Controlled Release Tablets

5 mg, 10 mg, 20 mg, 40 mg & 80 mg

An important Risk Management Plan (RMP) for ACT OXYCODONE CR has been implemented to minimize potential risks and problems associated with possible abuse, dependence and diversion. For more information, please visit the following web site: <http://soar-essor.ca/> or call toll free at 1-866-254-6111.

PART I: HEALTH PROFESSIONAL INFORMATION

SUMMARY PRODUCT INFORMATION

Route of Administration	Dosage Form / Strength	All Nonmedicinal Ingredients
Oral	Controlled Release Tablet / 5 mg, 10 mg, 20 mg, 40 mg and 80 mg	Colloidal silicon dioxide, hydroxypropyl methyl cellulose, magnesium stearate, microcrystalline cellulose, polyvinyl acetate and povidone. Coating ingredients include: hypromellose, hydroxypropyl cellulose, polyethylene glycol, titanium dioxide, iron oxide yellow (20 mg, 40 mg and 80 mg), iron oxide red (20 mg and 40 mg), FD&C Blue Aluminium Lake (5 mg and 80 mg)

INDICATIONS AND CLINICAL USE

ACT OXYCODONE CR (oxycodone hydrochloride) controlled release tablets is indicated for the management of pain severe enough to require daily, continuous, long-term opioid treatment, and:

- that is opioid-responsive; and
- for which alternative options are inadequate.

ACT OXYCODONE CR is not indicated as an as-needed (prn) analgesic.

Geriatrics (> 65 years of age):

In general, dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal, or cardiac function, concomitant disease or other drug therapy (see DOSAGE AND ADMINISTRATION).

Pediatrics (< 18 years of age):

The safety and efficacy of ACT OXYCODONE CR has not been studied in the pediatric

population. Therefore the use of ACT OXYCODONE CR is not recommended in patients under 18 years of age.

CONTRAINDICATIONS

ACT OXYCODONE CR (oxycodone hydrochloride) controlled release tablets are contraindicated in:

- Patients who are hypersensitive to the active substance (oxycodone) or other opioid analgesics or to any ingredient in the formulation. For a complete listing, see the **DOSAGE FORMS, COMPOSITION AND PACKAGING** section of the Product Monograph.
- In patients with known or suspected mechanical gastrointestinal obstruction (e.g., bowel obstruction, strictures) or any diseases/conditions that affect bowel transit (e.g., ileus of any type).
- Patients with suspected surgical abdomen (e.g., acute appendicitis or pancreatitis).
- Patients with mild, intermittent or short duration pain that can be managed with other pain medications.
- The management of acute pain.
- Patients with acute or severe bronchial asthma, chronic obstructive airway or status asthmaticus.
- Patients with acute respiratory depression, elevated carbon dioxide levels in the blood, and cor pulmonale.
- Patients with acute alcoholism, delirium tremens, and convulsive disorders.
- Patients with severe CNS depression, increased cerebrospinal or intracranial pressure, and head injury.
- Patients taking monoamine oxidase (MAO) inhibitors (or within 14 days of such therapy).
- Women who are breast-feeding, pregnant, or during labour and delivery (see **SERIOUS WARNINGS AND PRECAUTIONS** and **WARNINGS AND PRECAUTIONS**).

WARNINGS AND PRECAUTIONS

SERIOUS WARNINGS AND PRECAUTIONS

- **Limitations of Use**
Because of the risks of addiction, abuse, and misuse with opioids, even at recommended doses, and because of the greater risks of overdose and death with controlled-release opioid formulations, ACT OXYCODONE CR (oxycodone hydrochloride) controlled release tablets should only be used in patients for whom alternative treatment options (eg. non-opioid analgesic) are ineffective, not tolerated, or would be otherwise inadequate to provide appropriate management of pain (see **DOSAGE AND ADMINISTRATION**).
- **Addiction, Abuse, and Misuse**
ACT OXYCODONE CR poses risks of opioid addiction, abuse, and misuse, which can lead to overdose and death. Each patient's risk should be assessed prior to prescribing ACT OXYCODONE CR and all patients should be monitored regularly for the development of these behaviours or conditions (see **WARNINGS AND PRECAUTIONS**). ACT OXYCODONE CR should be stored securely to avoid theft or misuse.

- **Life-threatening Respiratory Depression: OVERDOSE**
Serious, life-threatening, or fatal respiratory depression may occur with use of ACT OXYCODONE CR. Infants exposed in-utero or through breast milk are at risk of life-threatening respiratory depression upon delivery or when nursed. Patients should be monitored for respiratory depression, especially during initiation of ACT OXYCODONE CR or following a dose increase. ACT OXYCODONE CR must be swallowed whole. Cutting, breaking, crushing, chewing, or dissolving ACT OXYCODONE CR tablets can lead to rapid release and absorption of a potentially fatal dose of oxycodone (see WARNINGS AND PRECAUTIONS). Further, instruct patients of the hazards related to taking opioids including fatal overdose.
- **Accidental Exposure**
Accidental ingestion of even one dose of ACT OXYCODONE CR, especially by children, can result in a fatal overdose of oxycodone (see DOSAGE AND ADMINISTRATION, Disposal, for instructions on proper disposal).
- **Neonatal Opioid Withdrawal Syndrome**
Prolonged maternal use of ACT OXYCODONE CR during pregnancy can result in neonatal opioid withdrawal syndrome, which may be life-threatening (see WARNINGS AND PRECAUTIONS).
- **Interaction with Alcohol**
The co-ingestion of alcohol with ACT OXYCODONE CR should be avoided as it may result in dangerous additive effects, causing serious injury or death (see WARNINGS AND PRECAUTIONS and DRUG INTERACTIONS).
- **Risks from Concomitant Use with Benzodiazepines or Other CNS Depressants**
Concomitant use of opioids with benzodiazepines or other CNS depressants, including alcohol, may result in profound sedation, respiratory depression, coma, and death (see WARNINGS AND PRECAUTIONS, Neurologic and DRUG INTERACTIONS).
 - Reserve concomitant prescribing of ACT OXYCODONE CR and benzodiazepines or other CNS depressants for use in patients for whom alternative treatment options are inadequate.
 - Limit dosages and durations to the minimum required.
 - Follow patients for signs and symptoms of respiratory depression and sedation.

General

ACT OXYCODONE CR must be swallowed whole. Taking cut, broken, chewed, dissolved or crushed ACT OXYCODONE CR tablets could lead to the rapid release and absorption of a potentially fatal dose of oxycodone.

ACT OXYCODONE CR 80 mg tablets, or a single dose greater than 40 mg, are for use in opioid tolerant patients only (see also DOSAGE AND ADMINISTRATION). A single dose greater than 40 mg of oxycodone, or total daily doses greater than 80 mg of oxycodone, may cause fatal respiratory depression when administered to patients who are not tolerant to the respiratory depressant effects of opioids (see WARNINGS AND PRECAUTIONS and DRUG INTERACTIONS).

Patients should be instructed not to give ACT OXYCODONE CR to anyone other than the patient for whom it was prescribed as such inappropriate use may have severe medical consequences, including death.

Patients should be cautioned not to consume alcohol while taking ACT OXYCODONE CR, as it may increase the chance of experiencing dangerous side effects.

Hyperalgesia that will not respond to a further dose increase of oxycodone may occur at particularly high doses. An oxycodone dose reduction or change in opioid may be required.

Addiction, Abuse and Misuse

Like all opioids, ACT OXYCODONE CR is a potential drug of abuse and misuse, which can lead to overdose and death. Therefore, ACT OXYCODONE CR should be prescribed and handled with caution.

Patients should be assessed for their clinical risks for opioid abuse or addiction prior to being prescribed opioids. All patients receiving opioids should be routinely monitored for signs of misuse and abuse.

Opioids, such as ACT OXYCODONE CR, should be used with particular care in patients with a history of alcohol and illicit/prescription drug abuse. However, concerns about abuse, addiction, and diversion should not prevent the proper management of pain.

With parenteral abuse, the tablet excipients can be expected to result in local tissue necrosis, infection, pulmonary granulomas, and increased risk of endocarditis and valvular heart injury.

Carcinogenesis and Mutagenesis

See **TOXICOLOGY** section.

Cardiovascular

Oxycodone administration may result in severe hypotension in patients whose ability to maintain adequate blood pressure is compromised by reduced blood volume, or concurrent administration of such drugs as phenothiazines or certain anesthetics.

Dependence/Tolerance

As with other opioids, tolerance and physical dependence may develop upon repeated administration of ACT OXYCODONE CR and there is a potential for development of psychological dependence.

Physical dependence and tolerance reflect the neuroadaptation of the opioid receptors to chronic exposure to an opioid, and are separate and distinct from abuse and addiction. Tolerance, as well as physical dependence, may develop upon repeated administration of opioids, and are not by themselves evidence of an addictive disorder or abuse.

Patients on prolonged therapy should be tapered gradually from the drug if it is no longer required for pain control. Withdrawal symptoms may occur following abrupt discontinuation of therapy or upon administration of an opioid antagonist. Some of the symptoms that may be associated with abrupt withdrawal of an opioid analgesic include body aches, diarrhea, gooseflesh, loss of appetite, nausea, nervousness or restlessness, anxiety, runny nose, sneezing, tremors or shivering, stomach cramps, tachycardia, trouble with sleeping, unusual increase in sweating, palpitations, unexplained fever, weakness and yawning).

Use in Drug and Alcohol Addiction

ACT OXYCODONE CR is an opioid with no approved use in the management of addictive disorders. Its proper usage in individuals with drug or alcohol dependence, either active or in remission, is for the management of pain requiring opioid analgesia. Patients with a history of addiction to drugs or alcohol may be at higher risk of becoming addicted to ACT OXYCODONE CR unless used under extreme caution and awareness.

Endocrine

Adrenal Insufficiency

Cases of adrenal insufficiency have been reported with opioid use, more often following greater than one month of use. Presentation of adrenal insufficiency may include non-specific symptoms and signs including nausea, vomiting, anorexia, fatigue, weakness, dizziness, and low blood pressure. If adrenal insufficiency is suspected, confirm the diagnosis with diagnostic testing as soon as possible. If adrenal insufficiency is diagnosed, treat with physiologic replacement doses of corticosteroids. Wean the patient off of the opioid to allow adrenal function to recover and continue corticosteroid treatment until adrenal function recovers. Other opioids may be tried as some cases reported use of a different opioid without recurrence of adrenal insufficiency. The information available does not identify any particular opioids as being more likely to be associated with adrenal insufficiency.

Gastrointestinal Effects

Oxycodone and other morphine-like opioids have been shown to decrease bowel motility. Oxycodone may obscure the diagnosis or clinical course of patients with acute abdominal conditions.

Neonatal Opioid Withdrawal Syndrome (NOWS)

Prolonged maternal use of opioids during pregnancy can result in withdrawal signs in the neonate. Neonatal opioid withdrawal syndrome, unlike opioid withdrawal syndrome in adults, may be life-threatening.

Neonatal opioid withdrawal syndrome presents as irritability, hyperactivity and abnormal sleep pattern, high pitched cry, tremor, vomiting, diarrhea and failure to gain weight. The onset, duration, and severity of neonatal opioid withdrawal syndrome vary based on the specific opioid used, duration of use, timing and amount of last maternal use, and rate of elimination of the drug by the newborn.

Use of ACT OXYCODONE CR is contraindicated in pregnant women (see **CONTRAINDICATIONS**).

Neurologic

Interactions with Central Nervous System Depressants (including benzodiazepines and alcohol): ACT OXYCODONE CR should be used with caution and in a reduced dosage during concomitant administration of other opioid analgesics, general anesthetics, phenothiazines and other tranquilizers, sedatives, hypnotics, tricyclic antidepressants, antipsychotics, antihistamines, benzodiazepines, centrally-active anti-emetics and other CNS depressants. Respiratory depression, hypotension and profound sedation, coma or death may result. When such combination therapy is contemplated, a substantial reduction in the dose of one or both agents should be considered and patients should be carefully monitored. ACT OXYCODONE CR should not be consumed with alcohol as it may increase the chance of experiencing dangerous side effects (see **DRUG INTERACTIONS**).

Observational studies have demonstrated that concomitant use of opioid analgesics and benzodiazepines increases the risk of drug-related mortality compared to use of opioid analgesics alone. Because of similar pharmacological properties, it is reasonable to expect similar risk with the concomitant use of other CNS depressant drugs with opioid analgesics (see **DRUG INTERACTIONS**). If the decision is made to prescribe a benzodiazepine or other CNS depressant concomitantly with an opioid analgesic, prescribe the lowest effective dosages and minimum durations of concomitant use. In patients already receiving an opioid analgesic, prescribe a lower initial dose of the benzodiazepine or other CNS depressant than indicated in the absence of an opioid, and titrate based on clinical response. If an opioid analgesic is initiated in a patient already taking a benzodiazepine or other CNS depressant, prescribe a lower initial dose of the opioid analgesic, and titrate based on clinical response. Follow patients closely for signs and symptoms of respiratory depression and sedation.

Advise both patients and caregivers about the risks of respiratory depression and sedation when

ACT OXYCODONE CR is used with benzodiazepines or other CNS depressants (including alcohol and illicit drugs). Advise patients not to drive or operate heavy machinery until the effects of concomitant use of the benzodiazepine or other CNS depressant have been determined. Screen patients for risk of substance use disorders, including opioid abuse and misuse, and warn them of the risk for overdose and death associated with the use of additional CNS depressants including alcohol and illicit drugs (see DRUG INTERACTIONS).

ACT OXYCODONE CR should not be consumed with alcohol as it may increase the chance of experiencing dangerous side effects, including death (see CONTRAINDICATIONS and ADVERSE REACTIONS, Sedation, and DRUG INTERACTIONS).

Severe pain antagonizes the subjective and respiratory depressant actions of opioid analgesics. Should pain suddenly subside, these effects may rapidly become manifest.

Use in Patients with Convulsive or Seizure Disorders: The oxycodone hydrochloride in ACT OXYCODONE CR may aggravate convulsions in patients with convulsive disorders, and may induce or aggravate seizures in some clinical settings. Therefore, ACT OXYCODONE CR should not be used in these patients (see CONTRAINDICATIONS).

Serotonin Syndrome: ACT OXYCODONE CR could cause a rare but potentially life-threatening condition resulting from concomitant administration of serotonergic drugs (e.g., antidepressants, migraine medications). Treatment with the serotonergic drug should be discontinued if such events (characterized by clusters of symptoms such as hyperthermia, rigidity, myoclonus, autonomic instability with possible rapid fluctuations of vital signs, mental status changes including confusion, irritability, extreme agitation progressing to delirium and coma) occur and supportive symptomatic treatment should be initiated. ACT OXYCODONE CR should not be used in combination with MAO inhibitors or serotonin-precursors (such as L-tryptophan, oxitriptan) and should be used with caution in combination with other serotonergic drugs (triptans, certain tricyclic antidepressants, lithium, tramadol, St. John's Wort) due to the risk of serotonin syndrome (see DRUG INTERACTIONS).

Head Injury: The respiratory depressant effects of oxycodone and the capacity to elevate cerebrospinal fluid pressure, may be greatly increased in the presence of an already elevated intracranial pressure produced by trauma. Also, oxycodone may produce confusion, miosis, vomiting and other side effects which obscure the clinical course of patients with head injury. In such patients, oxycodone must be used with extreme caution and only if it is judged essential.

Peri-Operative Considerations:

ACT OXYCODONE CR is not indicated for pre-emptive analgesia (administration pre-operatively for the management of post-operative pain).

In the case of planned chordotomy or other pain-relieving operations, patients should not be treated with ACT OXYCODONE CR for at least 24 hours before the operation and ACT OXYCODONE CR should not be used in the immediate post-operative period.

Physicians should individualize treatment, moving from parenteral to oral analgesics as appropriate. Thereafter, if ACT OXYCODONE CR is to be continued after the patient recovers from the postoperative period, a new dosage should be administered in accordance with the changed need for pain relief. The risk of withdrawal in opioid-tolerant patients should be addressed as clinically indicated.

The administration of analgesics in the peri-operative period should be managed by healthcare providers with adequate training and experience (e.g., by an anesthesiologist).

Oxycodone and other morphine-like opioids have been shown to decrease bowel motility. Ileus is a common post-operative complication, especially after intra-abdominal surgery with opioid analgesia. Caution should be taken to monitor for decreased bowel motility in post-operative patients receiving opioids. Standard supportive therapy should be implemented.

ACT OXYCODONE CR should not be used in the early post-operative period (12 to 24 hours post-surgery) unless the patient is ambulatory and gastrointestinal function is normal.

Psychomotor Impairment

Oxycodone may impair the mental and/or physical abilities needed for certain potentially hazardous activities such as driving a car or operating machinery. Patients should be cautioned accordingly. Patients should also be cautioned about the combined effects of oxycodone with other CNS depressants, including other opioids, phenothiazines, sedatives, hypnotics and alcohol.

Respiratory

Respiratory Depression: Serious, life-threatening, or fatal respiratory depression has been reported with the use of opioids, even when used as recommended. Respiratory depression from opioid use, if not immediately recognized and treated, may lead to respiratory arrest and death. Management of respiratory depression may include close observation, supportive measures, and use of opioid antagonists, depending on the patient's clinical status. Carbon dioxide (CO₂) retention from opioid-induced respiratory depression can exacerbate the sedating effects of opioids.

While serious, life-threatening, or fatal respiratory depression can occur at any time during the use of ACT OXYCODONE CR, the risk is greatest during the initiation of therapy or following a dose increase. Patients should be closely monitored for respiratory depression when initiating therapy with ACT OXYCODONE CR and following dose increases.

To reduce the risk of respiratory depression, proper dosing and titration of ACT OXYCODONE CR are essential (see **DOSAGE AND ADMINISTRATION**). Overestimating the ACT OXYCODONE CR dose when converting patients from another opioid product can result in a fatal overdose with the first dose.

Patient Counselling Information:

A patient information sheet should be provided to patients when ACT OXYCODONE CR tablets are dispensed to them.

Patients receiving ACT OXYCODONE CR should be given the following instructions by the physician:

1. Patients should be informed that accidental ingestion or use by individuals (including children) other than the patient for whom it was originally prescribed, may lead to severe, even fatal consequences.
2. Patients should be advised that ACT OXYCODONE CR contains oxycodone, an opioid pain medicine.
3. Patients should be advised that ACT OXYCODONE CR should only be taken as directed. The dose of ACT OXYCODONE CR should not be adjusted without consulting with a physician.
4. ACT OXYCODONE CR must be swallowed whole (not cut, broken, chewed, dissolved or crushed) due to the risk of fatal oxycodone overdose.
5. Patients should be advised to report episodes of pain and adverse experiences occurring during therapy. Individualization of dosage is essential to make optimal use of this medication.
6. Patients should not combine ACT OXYCODONE CR with alcohol or other central nervous system depressants (sleep aids, tranquilizers) because dangerous additive effects may occur, resulting in serious injury or death.
7. Patients should be advised to consult their physician or pharmacist if other medications are being used or will be used with ACT OXYCODONE CR.
8. Patients should be advised that if they have been receiving treatment with ACT OXYCODONE CR and cessation of therapy is indicated, it may be appropriate to taper ACT OXYCODONE CR dose, rather than abruptly discontinue it, due to the risk of precipitating withdrawal symptoms.

9. Patients should be advised that the most common adverse reactions that may occur while taking ACT OXYCODONE CR are asthenia, constipation, dizziness, dry mouth, headache, nausea, pruritus, somnolence, sweating and vomiting.
10. Patients should be advised that ACT OXYCODONE CR may cause drowsiness, dizziness or light-headedness and may impair mental and/or physical ability required for the performance of potentially hazardous tasks (e.g., driving, operating machinery). Patients started on ACT OXYCODONE CR or patients whose dose has been adjusted should be advised not to drive a car or operate machinery unless they are tolerant to the effects of ACT OXYCODONE CR.
11. Patients should be advised that ACT OXYCODONE CR is a potential drug of abuse. They should protect it from theft or misuse.
12. Patients should be advised that ACT OXYCODONE CR should never be given to anyone other than the individual for whom it was prescribed.
13. Patients should be advised that ACT OXYCODONE CR 80 mg tablets or a single dose greater than 40 mg are for use only in individuals tolerant to the effect of opioids.
14. Women of childbearing potential who become or are planning to become pregnant should be advised to consult a physician prior to initiating or continuing therapy with ACT OXYCODONE CR. Women who are breast-feeding or pregnant should not use ACT OXYCODONE CR.

Sexual Function/Reproduction

Long term use of opioids may be associated with decreased sex hormone levels and symptoms such as low libido, erectile dysfunction, or infertility (see ADVERSE REACTIONS, Post-Market Adverse Drug Reactions).

Special Populations

Special Risk Groups: Oxycodone should be administered with caution and in a reduced dosage to debilitated patients, and in patients with severely impaired pulmonary function, Addison's disease, hypothyroidism, myxedema, toxic psychosis, prostatic hypertrophy or urethral stricture.

Female subjects have, on average, plasma oxycodone concentrations up to 25% higher than males on a body weight adjusted basis. Plasma concentrations of oxycodone are increased by approximately 15% in elderly subjects receiving oxycodone hydrochloride; by 50 to 60% in patients with moderate degrees of renal impairment; and by approximately two-fold in patients with hepatic cirrhosis.

Pregnant Women: Animal reproduction studies have revealed no evidence of harm to the fetus due to oxycodone, however, as studies in humans have not been conducted, ACT OXYCODONE CR is contraindicated in patients who are pregnant (see CONTRAINDICATIONS).

Prolonged maternal use of opioids during pregnancy can result in withdrawal signs in the neonate. Neonatal opioid withdrawal syndrome, unlike opioid withdrawal syndrome in adults, may be life-threatening (see **WARNINGS AND PRECAUTIONS – Neonatal Opioid Withdrawal Syndrome**).

Labour, Delivery and Nursing Women:

ACT OXYCODONE CR is contraindicated during labour, delivery, pregnancy and in nursing mothers. Oxycodone can cross the placental barrier and is also excreted in breast milk. Life-threatening respiratory depression may occur in the infant if opioids are administered to the mother. Naloxone, a drug that counters the effect of opioids, should be readily available if ACT OXYCODONE CR is used in this population.

Pediatrics (< 18 years of age): The safety and efficacy of ACT OXYCODONE CR have not been studied in the pediatric population. Therefore, use of ACT OXYCODONE CR is not recommended in patients under 18 years of age.

Geriatrics (> 65 years of age): In general, dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or other drug therapy (see **DOSAGE AND ADMINISTRATION**).

Patients with Hepatic Impairment: When compared to normal subjects, patients with mild to severe hepatic dysfunction may have higher plasma concentrations of oxycodone and noroxycodone, and lower plasma concentrations of oxymorphone. There may be an increase in the elimination half-life of oxycodone, and this may be accompanied by an increase in drug effects.

Patients with Renal Impairment: When compared to normal subjects, patients with mild to severe renal dysfunction (creatinine clearance < 60 mL/min) may have higher plasma concentrations of oxycodone and its metabolites. There may be an increase in the elimination half-life of oxycodone, and this may be accompanied by an increase in drug effects.

In Vitro Dissolution Studies of Interaction with Alcohol

Among readily available drugs with the established potential to pharmacologically augment the CNS depressant effect of opioids, ethanol also has the potential to chemically interact with the pharmaceutical formulation to accelerate the release of opioids from the dosage form. Given the larger doses of opioids in controlled release opioid formulations on average, the occurrence of

such a formulation effect can further augment the risk of serious and unintended respiratory depression. A method to assess the potential for ethanol to accelerate the release of opioids from a pharmaceutical formulation requires the use of *in vitro* dissolution studies using simulated gastric fluid and 40% ethanol.

With oxycodone hydrochloride controlled release tablets, increasing concentrations of alcohol in the dissolution medium (from 0% to 40% v/v), resulted in a slight decrease in the rate of release of oxycodone from intact tablets. Additional *in vitro* dissolution testing in ethanol (40% v/v), conducted with oxycodone hydrochloride controlled release tablet fragments over a range of particles sizes, showed that dose dumping did not occur with the particle sizes tested.

ADVERSE REACTIONS

Adverse Drug Reaction Overview

Adverse effects of oxycodone hydrochloride controlled release tablets are similar to those of other opioid analgesics, and represent an extension of pharmacological effects of the drug class. The major hazards of opioids include respiratory and central nervous system depression and to a lesser degree, circulatory depression, respiratory arrest, shock and cardiac arrest.

The most frequently observed adverse effects of oxycodone hydrochloride controlled release tablets are asthenia, constipation, dizziness, dry mouth, headache, hyperhidrosis, nausea, pruritus, somnolence, and vomiting.

Sedation: Sedation is a common side effect of opioid analgesics, especially in opioid naïve individuals. Sedation may also occur partly because patients often recuperate from prolonged fatigue after the relief of persistent pain. Most patients develop tolerance to the sedative effects of opioids within three to five days and, if the sedation is not severe, will not require any treatment except reassurance. If excessive sedation persists beyond a few days, the dose of the opioid should be reduced and alternate causes investigated. Some of these are: concurrent CNS depressant medication, hepatic or renal dysfunction, brain metastases, hypercalcemia and respiratory failure. If it is necessary to reduce the dose, it can be carefully increased again after three or four days if it is obvious that the pain is not being well controlled. Dizziness and unsteadiness may be caused by postural hypotension, particularly in elderly or debilitated patients, and may be alleviated if the patient lies down.

Nausea and Vomiting: Nausea is a common side effect on initiation of therapy with opioid analgesics and is thought to occur by activation of the chemoreceptor trigger zone, stimulation of the vestibular apparatus and through delayed gastric emptying. The prevalence of nausea declines following continued treatment with opioid analgesics. When instituting therapy with an opioid for chronic pain, the routine prescription of an antiemetic should be considered. In the cancer patient, investigation of nausea should include such causes as constipation, bowel obstruction, uremia, hypercalcemia, hepatomegaly, tumour invasion of celiac plexus and

concurrent use of drugs with emetogenic properties. Persistent nausea which does not respond to dosage reduction may be caused by opioid-induced gastric stasis and may be accompanied by other symptoms including anorexia, early satiety, vomiting and abdominal fullness. These symptoms respond to chronic treatment with gastrointestinal prokinetic agents.

Constipation: Practically all patients become constipated while taking opioids on a persistent basis. In some patients, particularly the elderly or bedridden, fecal impaction may result. It is essential to caution the patients in this regard and to institute an appropriate regimen of bowel management at the start of prolonged opioid therapy. Stimulant laxatives, stool softeners, and other appropriate measures should be used as required. As fecal impaction may present as overflow diarrhea, the presence of constipation should be excluded in patients on opioid therapy prior to initiating treatment for diarrhea.

The following adverse effects occur with opioid analgesics and include those reported in oxycodone hydrochloride controlled release tablet clinical trials. The reactions are categorized by body system and frequency according to the following definitions: Very common ($\geq 1/10$); Common ($\geq 1/100$ to $< 1/10$); Uncommon ($\geq 1/1,000$ to $< 1/100$); Rare ($\geq 1/10,000$ to $< 1/1,000$); Very rare ($< 1/10,000$), Not known (cannot be estimated from the available data).

Blood and Lymphatic System Disorders:

Not known: lymphadenopathy

Cardiac Disorders:

Uncommon: palpitations, tachycardia

Not known: ST depression

Ear and Labyrinth Disorders:

Uncommon: vertigo, tinnitus

Eye Disorders:

Uncommon: miosis, visual impairment

Gastrointestinal Disorders:

Very common: constipation, nausea, vomiting

Common: abdominal pain, diarrhea, dry mouth, dyspepsia

Uncommon: dysphagia, eructation, flatulence, gastritis, hiccups, ileus, stomatitis

Not known: biliary spasm, dental caries

General Disorders and Administration Site Conditions:

Common: asthenia, fatigue, fever, hypotonia

Uncommon: abnormal gait, chest pain, chills, drug withdrawal syndrome, edema, edema peripheral, malaise, thirst, drug tolerance

Not known: drug withdrawal syndrome neonatal

Hepatobiliary Disorders:

Uncommon: increased hepatic enzyme

Not known: cholestasis

Immune System Disorders:

Uncommon: hypersensitivity

Not known: anaphylactic reaction, anaphylactoid reaction

Investigations:

Uncommon: weight loss

Metabolism and Nutrition Disorders:

Common: decreased appetite

Uncommon: dehydration, hypoglycemia

Rare: increased appetite

Nervous System Disorders:

Very common: dizziness, headache, somnolence

Common: tremor, lethargy

Uncommon: amnesia, convulsion, dysgeusia, hypertonia, hypoaesthesia, migraine, muscle contractions involuntary, paresthesia, speech disorder, syncope

Psychiatric Disorders:

Common: abnormal dreams, anxiety, confusional state, depression, insomnia, nervousness, thinking abnormal

Uncommon: affect lability, agitation, depersonalization, euphoric mood, hallucination, libido decreased, drug dependence

Rare: dysphoria

Not known: aggression, delirium

Renal and Urinary Disorders:

Uncommon: dysuria, hematuria, polyuria, urinary retention or hesitancy

Reproductive System and Breast Disorders:

Uncommon: erectile dysfunction

Not known: amenorrhea

Respiratory, Thoracic and Mediastinal Disorders:

Common: dyspnea

Uncommon: bronchitis, cough, pharyngitis, respiratory depression, yawning

Rare: sinusitis

Not known: bronchospasm, pneumonia

Skin and Subcutaneous Tissue Disorders:

Very common: pruritus

Common: hyperhidrosis, rash

Uncommon: dry skin, exfoliative dermatitis

Rare: urticaria

Vascular Disorders:

Uncommon: vasodilatation

Rare: hypotension, orthostatic hypotension

Post-marketing Experience

Hyperalgesia and hypogonadism have also been reported during post-marketing experience with oxycodone.

Androgen deficiency: Chronic use of opioids may influence the hypothalamic-pituitary-gonadal axis, leading to androgen deficiency that may manifest as low libido, impotence, erectile dysfunction, amenorrhea, or infertility. The causal role of opioids in the clinical syndrome of hypogonadism is unknown because the various medical, physical, lifestyle, and psychological stressors that may influence gonadal hormone levels have not been adequately controlled for in studies conducted to date. Patients presenting with symptoms of androgen deficiency should undergo laboratory evaluation.

DRUG INTERACTIONS

Overview

Interactions with Central Nervous System (CNS) Depressants (including benzodiazepines and alcohol): ACT OXYCODONE CR (oxycodone hydrochloride controlled release tablets) should be dosed with caution and started in a reduced dosage (1/3 to 1/2 of the usual dosage) in patients who are currently taking other central nervous system depressants (e.g., other opioids, sedatives, hypnotics, anti-depressants, phenothiazines, neuroleptics, anti-histamines, and anti-emetics) and beta-blockers, as they may enhance the CNS-depressant effect (e.g., respiratory depression) of ACT OXYCODONE CR. ACT OXYCODONE CR should not be consumed with alcohol as it may increase the chance of experiencing dangerous side effects.

Interactions with Anticholinergics: Concomitant administration of oxycodone with anticholinergics or medications with anticholinergic activity (e.g. tricyclic antidepressants,

antihistamines, antipsychotics, muscle relaxants, anti-Parkinson drugs) may result in increased anticholinergic adverse effects.

Drug-Drug Interactions

Drugs Metabolized by Cytochrome P450 Isozymes: Oxycodone is metabolized in part by cytochrome P450 2D6 and cytochrome P450 3A4 pathways. The activities of these metabolic pathways may be inhibited or induced by various co-administered drugs or dietary elements, which may alter plasma oxycodone concentrations. Oxycodone doses may need to be adjusted accordingly.

Inhibitors of CYP3A4:

Since the CYP3A4 isoenzyme plays a major role in the metabolism of oxycodone hydrochloride, drugs that inhibit CYP3A4 activity, such as macrolide antibiotics (e.g., erythromycin, clarithromycin), azole-antifungal agents (e.g., ketoconazole), and protease inhibitors (e.g., ritonavir) and grapefruit juice may cause decreased clearance of oxycodone which could lead to an increase in oxycodone plasma concentrations. A published study showed that the co-administration of the antifungal drug, voriconazole, increased oxycodone AUC and C_{max} by 3.6- and 1.7-fold, respectively. Although clinical studies have not been conducted with other CYP3A4 inhibitors, the expected clinical results would be increased or prolonged opioid effects. If co-administration with ACT OXYCODONE CR is necessary, caution is advised when initiating therapy with, currently taking, or discontinuing CYP450 inhibitors. Evaluate these patients at frequent intervals and consider dose adjustments until stable drug effects are achieved.

Inducers of CYP3A4:

CYP450 inducers, such as rifampin, carbamazepine, phenytoin, and St. John's wort, may induce the metabolism of oxycodone and, therefore, may cause increased clearance of the drug which could lead to a decrease in oxycodone plasma concentrations, lack of efficacy or possibly the development of an abstinence syndrome in a patient who had developed physical dependence to oxycodone. A published study showed that the co-administration of rifampin, a drug metabolizing enzyme inducer, decreased oxycodone (oral) AUC and C_{max} by 86% and 63% respectively. If co-administration with ACT OXYCODONE CR is necessary, caution is advised when initiating therapy with, currently taking or discontinuing CYP3A4 inducers. Evaluate these patients at frequent intervals and consider dose adjustments until stable drug effects are achieved.

Inhibitors of CYP2D6:

Oxycodone is metabolized in part to oxymorphone via cytochrome CYP2D6. While this pathway may be blocked by a variety of drugs (e.g., certain cardiovascular drugs including amiodarone and quinidine as well as polycyclic antidepressants), such blockade has not been shown to be of clinical significance during oxycodone treatment.

Administration with Mixed Activity Agonist/Antagonist Opioids:

Mixed agonist/antagonist opioid analgesics (i.e., pentazocine, nalbuphine, butorphanol, and buprenorphine) should be administered with caution to a patient who has received or is receiving a course of therapy with a pure opioid agonist analgesic such as oxycodone. In this situation, mixed agonist/antagonist analgesics may reduce the analgesic effect of oxycodone and/or may precipitate withdrawal symptoms in these patients.

MAO Inhibitors:

MAO inhibitors intensify the effects of opioid drugs which can cause anxiety, confusion and decreased respiration. ACT OXYCODONE CR is contraindicated in patients receiving MAO inhibitors or who have used them within the previous 14 days (see **CONTRAINDICATIONS**).

Warfarin and Other Coumarin Anticoagulants: Clinically relevant changes in International Normalized Ratio (INR or Quick-value) in both directions have been observed in individuals when oxycodone and coumarin anticoagulants are co-administered.

Serotonergic Agents: Co-administration of oxycodone hydrochloride with a serotonergic agent, such as a selective serotonin re-uptake inhibitor or a serotonin norepinephrine re-uptake inhibitor, may increase the risk of serotonin syndrome, a potentially life-threatening condition (see **WARNINGS AND PRECAUTIONS, Neurologic**).

Drug-Food Interactions

Administration of oxycodone hydrochloride controlled release tablets with food results in an increase in peak plasma oxycodone concentration of up to 1.5-fold but has no significant effect on the extent of absorption of oxycodone.

Drug-Herb Interactions

Interactions with herbal products have not been established.

Drug-Laboratory Interactions

Interactions with laboratory tests have not been established.

Drug-Lifestyle Interactions

The concomitant use of alcohol should be avoided (see **WARNINGS AND PRECAUTIONS, General**).

DOSAGE AND ADMINISTRATION

All doses of opioids carry an inherent risk of fatal or non-fatal adverse events. This risk is increased with higher doses. For the management of chronic non-cancer, non-palliative

pain, it is recommended that 60 mg (90 morphine milligram equivalent) of oxycodone hydrochloride controlled release tablets not be exceeded. Each patient should be assessed for their risk prior to prescribing ACT OXYCODONE CR, as the likelihood of experiencing serious adverse events can depend upon the type of opioid, duration of treatment, level of pain as well as the patient's own level of tolerance. In addition, the level of pain should be assessed routinely to confirm the most appropriate dose and the need for further use of ACT OXYCODONE CR (see DOSAGE AND ADMINISTRATION - Adjustment or Reduction of Dosage).

Dosing Considerations

ACT OXYCODONE CR should only be used in patients for whom alternative treatment options (e.g., non-opioid analgesics) are ineffective, or not tolerated, or would be otherwise inadequate to provide appropriate management of pain.

ACT OXYCODONE CR must be swallowed whole. Cutting, breaking, crushing, chewing, or dissolving ACT OXYCODONE CR tablets can lead to rapid release and absorption of a potentially fatal dose of oxycodone (see WARNINGS AND PRECAUTIONS).

ACT OXYCODONE CR 80 mg tablets, or a single dose greater than 40 mg, are for use in opioid tolerant patients only. A single dose greater than 40 mg, or total daily doses greater than 80 mg, may cause fatal respiratory depression when administered to patients who are not tolerant to the respiratory depressant effects of opioids.

ACT OXYCODONE CR should not be used in the early post-operative period (12 to 24 hours post-surgery) unless the patient is ambulatory and gastrointestinal function is normal.

ACT OXYCODONE CR is not indicated for rectal administration.

The controlled release tablets may be taken with or without food, with a glass of water.

In general, dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal, or cardiac function, concomitant disease or other drug therapy. The dosage should be adjusted to the lowest ACT OXYCODONE CR dose that will achieve the goal of satisfactory pain relief with acceptable side effects.

Recommended Dose and Dosage Adjustment

Adults: Individual dosing requirements vary considerably based on each patient's age, weight, severity and cause of pain, and medical and analgesic history. Monitor patients closely for respiratory depression, especially within the first 24 to 72 hours of initiating therapy with ACT OXYCODONE CR.

Patients Not Receiving Opioids at the Time of Initiation of ACT OXYCODONE CR

Treatment: The usual initial adult dose of ACT OXYCODONE CR for patients who have not previously received opioid analgesics is 10 mg every 12 hours.

Use of higher starting doses in patients who are not opioid tolerant may cause fatal respiratory depression.

Patients Currently Receiving Opioids: Patients currently receiving other oral oxycodone formulations may be transferred to ACT OXYCODONE CR tablets at the same total daily oxycodone dosage, equally divided into two 12 hourly ACT OXYCODONE CR doses.

For patients who are receiving an alternate opioid, the "oral oxycodone equivalent" of the analgesic presently being used should be determined. Having determined the total daily dosage of the present analgesic, Table 1 can be used to calculate the approximate daily oral oxycodone dosage that should provide equivalent analgesia. This total daily oral oxycodone dose should then be equally divided into two 12 hourly ACT OXYCODONE CR doses. It is usually appropriate to treat a patient with only one opioid at a time. Further dose reductions should be considered due to incomplete cross-tolerance between opioids.

Patients with Hepatic and Renal Impairment: The dose initiation should follow a conservative approach in these patients. The recommended adult starting dose in these patients should be at 1/3 to 1/2 the usual starting dose followed by careful dose titration to adequate pain control according to their clinical situation.

Opioid Rotation: Conversion ratios for opioids are subject to variations in kinetics governed by genetics and other factors. When switching from one opioid to another, consider **reducing the calculated dose by 25 to 50%** to minimize the risk of overdose. Subsequently, up-titrate the dose, as required, to reach the appropriate maintenance dose.

Table 1: Opioid Conversion Table^a

Opioids	To convert to oral morphine equivalent	To convert from oral morphine multiply by	Daily 90 mg MED ^b
Morphine	1	1	90 mg
Codeine	0.15	6.67	600 mg
Hydromorphone	5	0.2	18 mg
Oxycodone	1.5	0.667	60 mg
Tapentadol	0.3 to 0.4	2.5 to 3.33	300 mg
Tramadol	0.1 to 0.2	6	***
Methadone	Morphine dose equivalence is not reliably established		

*** The maximum recommended daily dose of tramadol is 300 mg – 400 mg depending on the formulation.

a. Adapted from the 2017 Canadian guideline for opioids for chronic non-cancer pain. McMaster University; 2017

b. MED. Morphine Equivalent Dose

Patients who are receiving 1 to 5 tablets/capsules per day of a fixed-dose combination opioid/non-opioid containing 5 mg of oxycodone or 30 mg codeine should be started on 10 mg to 20 mg ACT OXYCODONE CR q12h. For patients receiving 6 to 9 tablets/capsules per day of a fixed-dose combination opioid/non-opioid containing 5 mg of oxycodone or 30 mg codeine, a starting dose of 20 mg to 30 mg* q12h should be used and for patients receiving 10 to 12 tablets/capsules per day of a fixed-dose combination opioid/non-opioid containing 5 mg of oxycodone or 30 mg codeine, a starting dose of 30 mg* to 40 mg q12h is suggested. For those receiving > 12 tablets/capsules per day of a fixed-dose combination opioid/non-opioid containing 5 mg of oxycodone or 30 mg codeine, conversions should be based on the total daily opioid dose.

* ACT OXYCODONE CR is NOT available as 30 mg oxycodone hydrochloride controlled release tablets.

Use with Non-Opioid Medications: If a non-opioid analgesic is being provided, it may be continued. If the non-opioid is discontinued, consideration should be given to increasing the opioid dose to compensate for the non-opioid analgesic. ACT OXYCODONE CR can be safely used concomitantly with usual doses of other non-opioid analgesics.

Dose Titration: Dose titration is the key to success with opioid analgesic therapy. **Proper optimization of doses scaled to the relief of the individual's pain should aim at regular administration of the lowest dose of controlled release oxycodone (ACT OXYCODONE CR) which will achieve the overall treatment goal of satisfactory pain relief with acceptable side effects.**

Dosage adjustments should be based on the patient's clinical response. In patients receiving ACT OXYCODONE CR, the dose may be titrated at intervals of 24 to 36 hours to that which provides satisfactory pain relief without unmanageable side effects. ACT OXYCODONE CR is designed to allow 12 hourly dosing.

If pain repeatedly occurs at the end of the dosing interval it is generally an indication for a dosage increase rather than more frequent administration of controlled release oxycodone (ACT OXYCODONE CR).

Adjustment or Reduction of Dosage: Physical dependence with or without psychological dependence tends to occur with chronic administration of opioids, including ACT OXYCODONE CR. Withdrawal (abstinence) symptoms may occur following abrupt discontinuation of therapy. These symptoms may include body aches, diarrhea, gooseflesh, loss of appetite, nausea, nervousness or restlessness, runny nose, sneezing, tremors or shivering, stomach cramps, tachycardia, trouble with sleeping, unusual increase in sweating, palpitations, unexplained fever, weakness and yawning.

Following successful relief of pain, periodic attempts to reduce the opioid dose should be made. Patients on prolonged therapy should be withdrawn gradually from the drug if it is no longer required for pain control. In patients who are appropriately treated with opioid analgesics and who undergo gradual withdrawal for the drug, these symptoms are usually mild (see **WARNINGS AND PRECAUTIONS**). Tapering should be individualized and carried out under medical supervision.

Patient should be informed that reducing and/or discontinuing opioids decreases their tolerance to these drugs. If treatment needs to be re-initiated, the patient must start at the lowest dose and titrate up to avoid overdose.

Opioid analgesics may only be partially effective in relieving dysesthetic pain, stabbing pains, activity-related pain and some forms of headache. That is not to say that patients with these types of pain should not be given an adequate trial of opioid analgesics, but it may be necessary to refer such patients at an early time to other forms of pain therapy.

Missed Dose

If the patient forgets to take one or more doses, they should take their next dose at the next scheduled time and in the normal amount.

Disposal

ACT OXYCODONE CR should be kept in a safe place, such as under lock and out of the sight and reach of children before, during and after use. ACT OXYCODONE CR should not be used in front of children, since they may copy these actions.

Unused or expired ACT OXYCODONE CR should be properly disposed of as soon as it is no longer needed to prevent accidental exposure to others, including children or pets. ACT OXYCODONE CR should not be shared with others and steps should be taken to protect it from theft or misuse. The patient should speak to their pharmacist about temporary storage options, if required, until the medication can be returned to the pharmacy for safe disposal.

ACT OXYCODONE CR should never be disposed of in household trash. Disposal via a pharmacy take back program is recommended.

OVERDOSAGE

For management of a suspected drug overdose, contact your regional Poison Control Centre.

Symptoms: Serious overdose with oxycodone may be characterized by respiratory depression (a decrease in respiratory rate and/or tidal volume, Cheyne-Stokes respiration, cyanosis), extreme somnolence progressing to stupor or coma, miosis, hypotonia, cold and clammy skin, and sometimes bradycardia and hypotension. Severe overdose may result in apnea, circulatory collapse, cardiac arrest, pulmonary edema and death.

Treatment: Primary attention should be given to the establishment of adequate respiratory exchange through the provision of a patent airway and controlled or assisted ventilation. The opioid antagonist naloxone hydrochloride is a specific antidote against respiratory depression due to overdose or as a result of unusual sensitivity to oxycodone. An appropriate dose of an opioid antagonist should therefore be administered, preferably by the intravenous route. The usual initial i.v. adult dose of naloxone is 0.4 mg or higher. Concomitant efforts at respiratory resuscitation should be carried out. Since the duration of action of oxycodone, particularly sustained release formulations, may exceed that of the antagonist, the patient should be under continued surveillance and doses of the antagonist should be repeated as needed to maintain adequate respiration.

An antagonist should not be administered in the absence of clinically significant respiratory or cardiovascular depression. Oxygen, intravenous fluids, vasopressors and other supportive measures should be used as indicated.

In individuals physically dependent on opioids, the administration of the usual dose of narcotic antagonist will precipitate an acute withdrawal syndrome. The severity of this syndrome will depend on the degree of physical dependence and the dose of antagonist administered. The use of narcotic antagonists in such individuals should be avoided if possible. If a narcotic antagonist must be used to treat serious respiratory depression in the physically dependent patient, the antagonist should be administered with extreme care by using dosage titration, commencing with 10% to 20% of the usual recommended initial dose.

Evacuation of gastric contents may be useful in removing unabsorbed drug, particularly when a sustained release formulation has been taken.

ACTION AND CLINICAL PHARMACOLOGY

Mechanism of Action

Oxycodone is a semi-synthetic opioid analgesic which exerts an agonist effect at specific, saturable opioid receptors in the CNS and other tissues. In man, oxycodone produces a variety of effects including analgesia, constipation from decreased gastrointestinal motility, suppression of the cough reflex, respiratory depression from reduced responsiveness of the respiratory center to CO₂, nausea and vomiting via stimulation of the chemoreceptor trigger zone, changes in mood including euphoria and dysphoria, sedation, mental clouding, and alterations of the endocrine and autonomic nervous systems.

Pharmacodynamics

Oxycodone retains at least one-half of its analgesic activity when administered orally and with acute dosing is approximately twice as potent as orally administered morphine.

There is no intrinsic limit to the analgesic effect of oxycodone; like morphine, adequate doses will relieve even the most severe pain. Clinically however, dosage limitations are imposed by the adverse effects, primarily respiratory depression, nausea and vomiting, which can result from high doses.

Studies with oxycodone hydrochloride controlled release tablets and immediate release oxycodone hydrochloride tablets (IR) in normal volunteers and patients demonstrate a consistent relationship between oxycodone dosage and plasma oxycodone concentrations as well as between concentration and pharmacodynamic effects. In a single dose analgesic assay, the peak effect of oxycodone hydrochloride controlled release tablets (20 and 30 mg) was greater than that of 10 mg oxycodone hydrochloride controlled release tablet and was equivalent to that of two tablets of oxycodone (5 mg) plus acetaminophen (325 mg), or 15 mg of immediate release oxycodone but with a longer duration of action. In patients with pain due to osteoarthritis, oxycodone hydrochloride q12h was more effective than placebo in decreasing pain and in improving quality of life, mood and sleep. In patients with cancer pain, oxycodone hydrochloride controlled release tablets administered q12h produced equivalent analgesia to immediate release oxycodone hydrochloride tablets administered four times per day. In patients with low back pain, oxycodone hydrochloride controlled release tablets q12h and immediate release oxycodone hydrochloride given four times per day, were equally effective. Titration to analgesic effect was achieved as easily with Oxycodone hydrochloride controlled release tablets as with immediate release oxycodone hydrochloride tablets.

Central Nervous System: Oxycodone produces respiratory depression by direct action on brain stem respiratory centres. The respiratory depression involves both a reduction in the responsiveness of the brain stem centres to increases in CO₂ tension and to electrical stimulation.

Oxycodone depresses the cough reflex by direct effect on the cough centre in the medulla. Antitussive effects may occur with doses lower than those usually required for analgesia.

Oxycodone causes miosis, even in total darkness. Pinpoint pupils are a sign of opioid overdose but are not pathognomonic (e.g., pontine lesions of hemorrhagic or ischemic origin may produce similar findings). Marked mydriasis rather than miosis may be seen with hypoxia in the setting of oxycodone overdose.

Gastrointestinal Tract and Other Smooth Muscle: Oxycodone causes a reduction in motility associated with an increase in smooth muscle tone in the antrum of the stomach and duodenum. Digestion of food in the small intestine is delayed and propulsive contractions are decreased.

Propulsive peristaltic waves in the colon are decreased, while tone may be increased to the point of spasm resulting in constipation. Other opioid-induced effects may include a reduction in gastric, biliary and pancreatic secretions, spasm of the sphincter of Oddi, and transient elevations in serum amylase.

Cardiovascular System: Oxycodone may produce release of histamine with or without associated peripheral vasodilatation. Manifestations of histamine release and/or peripheral vasodilatation may include pruritus, flushing, red eyes, hyperhidrosis and/or orthostatic hypotension.

Immune System: *In vitro* and animal studies indicate that opioids have a variety of effects on immune functions, depending on the context in which they are used. The clinical significance of these findings is unknown.

Concentration – Efficacy Relationships

Studies in normal volunteers and patients reveal predictable relationships between oxycodone dosage and plasma oxycodone concentrations, as well as between concentration and certain expected opioid effects, such as papillary constriction, sedation, overall subjective “drug effect”, analgesia and feelings of “relaxation”.

The minimum effective analgesic concentration will vary widely among patients, especially among patients who have been previously treated with potent agonist opioids. As a result, patients must be treated with individualized titration of dosage to the desired effect. The minimum effective analgesic concentration of oxycodone for any individual patient may increase over time due to an increase in pain, the development of a new pain syndrome and/or the development of analgesic tolerance.

Concentration – Adverse Reaction Relationship

There is a significant relationship between increasing oxycodone plasma concentrations and increasing frequency of dose-related opioid adverse reactions such as nausea, vomiting, CNS effects, and respiratory depression. In opioid-tolerant patients, the situation may be altered by the development of tolerance to opioid-related side effects.

The dose of ACT OXYCODONE CR must be individualized (see **DOSAGE AND ADMINISTRATION**) because the effective analgesic dose for some patients will be too high to be tolerated by other patients.

Pharmacokinetics

The activity of ACT OXYCODONE CR is primarily due to the parent drug oxycodone. ACT OXYCODONE CR is designed to provide delivery of oxycodone over 12 hours.

Cutting, breaking, chewing, crushing or dissolving ACT OXYCODONE CR impairs the controlled release delivery mechanism and could lead to the rapid release and absorption of a potentially fatal dose of oxycodone.

Absorption:

About 60% to 87% of an oral dose of oxycodone reaches the central compartment in comparison to a parenteral dose. The high oral bioavailability is due to low pre-systemic and/or first-pass metabolism.

Distribution: Following intravenous administration, the steady-state volume of distribution (V_{ss}) for oxycodone was 2.6 L/kg. Oxycodone binding to plasma protein at 37°C and a pH of 7.4 was about 45%. Once absorbed, oxycodone is distributed to skeletal muscle, liver, intestinal tract, lungs, spleen, and brain. Oxycodone has been found in breast milk.

Metabolism: Oxycodone is extensively metabolized by multiple metabolic pathways to produce noroxycodone, oxymorphone and noroxymorphone, which are subsequently glucuronidated. Noroxycodone and noroxymorphone are the major circulating metabolites. CYP3A mediated N-demethylation to noroxycodone is the primary metabolic pathway of oxycodone with a lower contribution from CYP2D6 mediated O-demethylation to oxymorphone. Therefore, the formation of these and related metabolites can, in theory, be affected by other drugs (see **Drug-Drug Interactions**).

Noroxycodone exhibits very weak anti-nociceptive potency compared to oxycodone, however, it undergoes further oxidation to produce noroxymorphone, which is active at opioid receptors. Although noroxymorphone is an active metabolite and is present at relatively high concentrations in circulation, it does not appear to cross the blood-brain barrier to a significant extent.

Oxymorphone has been shown to be active and possessing analgesic activity but its contribution to analgesia following oxycodone administration is thought to be clinically insignificant. Other metabolites (α - and β -oxycodol, noroxycodol and oxymorphol) may be present at very low concentrations and demonstrate limited penetration in to the brain as compared to oxycodone. The enzymes responsible for keto-reduction and glucuronidation pathways in oxycodone metabolism have not been established.

Excretion: Oxycodone and its metabolites are excreted in both urine and feces. The amounts measured in the urine have been reported as follows: free and conjugated oxycodone 8.9%, free noroxycodone 23%, free oxymorphone less than 1%, conjugated oxymorphone 10%, free and conjugated noroxymorphone 14%, reduced free and conjugated metabolites up to 18%. The total plasma clearance was approximately 1.4 L/min in adults.

Pharmacokinetic studies of oxycodone hydrochloride controlled release tablets in normal volunteers demonstrate that both AUC and C_{max} increase in a dose proportional manner and that

the six tablet strengths are bioequivalent. In single dose studies, oxycodone hydrochloride controlled release tablet was absorbed to an equivalent extent as immediate release oxycodone but with a reduced maximum concentration (C_{max} ratio approximately 50%), a prolonged (2.4x) time to maximum concentration (t_{max} approximately 2.8 hours), with a biphasic absorption pattern, with two apparent absorption half-times of 0.6 and 6.9 hours, which describe the initial release of oxycodone from the tablet, followed by a prolonged release. Release in vitro is pH-independent.

In steady state pharmacokinetic studies of oxycodone hydrochloride controlled release tablets q12h, maximum plasma concentrations (C_{max}) of oxycodone were equivalent to those obtained with q6h administration of oral immediate release preparations and was achieved approximately 3 hours after administration of oxycodone hydrochloride controlled release tablets. Steady-state was achieved within 24-36 hours of initiation of dosing. The absorption of oxycodone from oxycodone hydrochloride controlled release tablets is not significantly influenced when administered in the presence of food.

Special Populations and Conditions

Pediatrics: Oxycodone hydrochloride controlled release tablets has not been studied in children and is not indicated for patients less than 18 years of age.

Geriatrics: Plasma concentrations of oxycodone are increased by approximately 15% in elderly subjects receiving oxycodone hydrochloride controlled release tablets.

Race: No data available.

Hepatic Insufficiency: Patients with mild to moderate hepatic dysfunction showed peak plasma oxycodone and noroxycodone concentrations approximately 50% and 20% higher, respectively, than normal subjects. AUC values were approximately 95% and 75% higher, respectively. Oxymorphone peak plasma concentrations and AUC values were lower by 15% to 50%. The $t_{1/2}$ elimination for oxycodone increased by 2.3 hours.

Plasma concentrations of oxycodone are increased by approximately 2-fold in patients with hepatic cirrhosis.

Renal Insufficiency: Patients with mild to severe renal dysfunction showed peak plasma oxycodone and noroxycodone concentrations 50% and 20% higher, respectively, and AUC values for oxycodone, noroxycodone, and oxymorphone 60%, 50%, and 40% higher than normal subjects, respectively. This was accompanied by an increase in sedation but not by differences in respiratory rate, pupillary constriction, or several other measures of drug effect. There was an increase in mean elimination $t_{1/2}$ for oxycodone of 1 hour.

Genetic Polymorphism: No data available.

STORAGE AND STABILITY

Store ACT OXYCODONE CR Tablets between 15°C and 30°C. Keep container tightly closed. Protect from moisture.

DOSAGE FORMS, COMPOSITION AND PACKAGING

Each ACT OXYCODONE CR tablet contains 5 mg, 10 mg, 20 mg, 40 mg or 80 mg of oxycodone hydrochloride, as well as the following inactive ingredients; colloidal silicon dioxide, hydroxy propyl methyl cellulose, magnesium stearate, microcrystalline cellulose, polyvinyl acetate and povidone. Coating ingredients include: hypromellose, hydroxypropyl cellulose, polyethylene glycol, titanium dioxide, iron oxide yellow (20 mg, 40 mg and 80 mg), iron oxide red (20 mg and 40 mg), FD&C Blue Aluminium Lake (5 mg and 80 mg).

ACT OXYCODONE CR 5 mg are blue, round, unscored, biconvex tablets engraved with “OXY” over “CR” on one side and “W 5” on the other side. Available in bottles of 100 tablets.

ACT OXYCODONE CR 10 mg are white, round, unscored, biconvex tablets engraved with “OXY” over “CR” on one side and “W 10” on the other side. Available in bottles of 100 tablets.

ACT OXYCODONE CR 20 mg are pink, round, unscored, biconvex tablets engraved with “OXY” over “CR” on one side and “W 20” on the other side. Available in bottles of 100 and 500 tablets.

ACT OXYCODONE CR 40 mg are yellow, round, unscored, biconvex tablets engraved with “OXY” over “CR” on one side and “W 40” on the other side. Available in bottles of 100 and 500 tablets.

ACT OXYCODONE CR 80 mg are green, round, unscored, biconvex tablets engraved with “OXY” over “CR” on one side and “W 80” on the other side. Available in bottles of 100 tablets.

PART II: SCIENTIFIC INFORMATION

PHARMACEUTICAL INFORMATION

Drug Substance

Oxycodone hydrochloride is a semi-synthetic derivative of the naturally occurring opium alkaloid, thebaine.


Proper name: Oxycodone Hydrochloride

Chemical name: 4,5 α epoxy-14-hydroxy-3-methoxy-17-methylmorphinan-6-one hydrochloride

Molecular formula: $C_{18}H_{21}NO_4 \cdot HCl$

Molecular mass: 351.9 g/mol

Structural formula:



*Chiral Centres

Physicochemical properties: Oxycodone hydrochloride is a white to off-white, odourless, crystalline powder. It is soluble in water and slightly soluble in alcohol and has a melting point of 218°C to 223°C.

CLINICAL TRIALS

Comparative Bioavailability Studies

A blind, randomized, 2-way crossover, bioequivalence study of ACT OXYCODONE CR and OxyContin[®] administered as 1 x 10 mg tablet in healthy males under fasting conditions (n=29) was conducted and demonstrates the bioequivalence of ACT OXYCODONE CR and OxyContin[®].

Oxycodone (1 x 10 mg) From measured data Geometric LS Mean Arithmetic Mean (CV %)				
Parameter	ACT OXYCODONE CR 10 mg Tablet	OxyContin [†] 10 mg Tablet	% Ratio of Geometric LS Means	90% Confidence Interval
AUC _T (ng·h/mL)	91.17 93.79 (21.59)	91.72 94.78 (25.74)	99.40%	94.34% - 104.73%
AUC _∞ (ng·h/mL)	10.02 10.36 (23.70)	10.90 11.40 (28.06)	91.87%	85.57% - 98.64%
C _{max} (ng/mL)	80.78 83.92 (25.20)	79.74 81.92 (23.34)	101.30%	96.13% - 106.75%
T _{max} [§] (h)	4.00 (1.00 – 6.00)	2.00 (1.00 – 5.00)		
t _{1/2} [€] (h)	6.38	8.87		

[†] Manufacturer: Purdue Pharma; origin (country of purchase): Canada.

[§] Expressed as the median (range) only

[€] Expressed as the arithmetic mean only

A blind, randomized, 2-way crossover, bioequivalence study of ACT OXYCODONE CR and OxyContin[®] administered as 1 x 10 mg tablet in healthy males under fed conditions (n=27) was conducted and demonstrates the bioequivalence of ACT OXYCODONE CR and OxyContin[®].

Oxycodone (1 x 10 mg) From measured data Geometric LS Mean Arithmetic Mean (CV %)				
Parameter	ACT OXYCODONE CR 10 mg Tablet	OxyContin [†] 10 mg Tablet	% Ratio of Geometric LS Means	90% Confidence Interval
AUC _T (ng·h/mL)	112.42 114.795 (20.2)	112.05 114.399 (20.8)	100.33	98.13 – 102.58
AUC _∞ (ng·h/mL)	117.84 120.450 (20.7)	121.87 124.553 (21.4)	96.69	94.42 – 99.02
C _{max} (ng/mL)	12.85 13.221 (24.3)	11.71 11.864 (16.7)	109.78	105.25 – 114.49
T _{max} [§] (h)	4.50 (1.00 – 7.00)	3.00 (2.00 – 7.00)		
t _{1/2} [€] (h)	4.94 (11.9)	6.12 (21.1)		

[†] Manufacturer: Purdue Pharma; origin (country of purchase): Canada.

[§] Expressed as the median (range) only

[€] Expressed as the arithmetic mean only

A blind, randomized, 2-way crossover, bioequivalence study of ACT OXYCODONE CR and OxyContin[®] administered as 1 x 10 mg tablet every 12 hours for 4 consecutive days in healthy subjects (n=26) was conducted and demonstrates the bioequivalence of ACT OXYCODONE CR and OxyContin[®].

Oxycodone (1 x 10 mg) From measured data Geometric LS Mean Arithmetic Mean (CV %)				
Parameter	ACT OXYCODONE CR 10 mg Tablet	OxyContin [†] 10 mg Tablet	% Ratio of Geometric LS Means	90% Confidence Interval
AUC _{TSS} (ng·h/mL)	10.04 10.40 (25.77)	10.43 10.84 (24.81)	96.21 %	92.92% - 99.61%
C _{max} SS (ng/mL)	11.88 12.29 (23.05)	12.91 13.37 (21.84)	92.04%	88.03% - 96.24%
C _{min} SS (ng/mL)	4.73 5.02 (35.19)	4.62 4.89 (33.50)	102.46%	94.62% - 110.95%
T _{max} [§] (h)	2.25 (1.00 – 4.50)	1.50 (0.50 – 4.00)		
F1 (%)	86.06	95.81		

[†] Manufacturer: Purdue Pharma; origin (country of purchase): Canada.

[§] Expressed as the median (range) only

A blind, randomized, 2-way crossover, bioequivalence study of ACT OXYCODONE CR and OxyContin[®] administered as 1 x 20 mg tablet in healthy males under fasting conditions (n=33) was conducted and demonstrates the bioequivalence of ACT OXYCODONE CR and OxyContin[®].

Oxycodone (1 x 20 mg) From measured data Geometric LS Mean Arithmetic Mean (CV %)				
Parameter	ACT OXYCODONE CR 20 mg Tablet	OxyContin [†] 20 mg Tablet	% Ratio of Geometric LS Means ¹	90% Confidence Interval
AUC _T (ng·h/mL)	213.05 221.35 (29.12)	226.76 239.44 (33.31)	93.96%	89.48% - 98.65%
AUC _∞ (ng·h/mL)	223.02 233.28 (31.91)	237.72 251.96 (34.39)	93.81%	89.13% - 98.74%
C _{max} (ng/mL)	16.99 17.47 (23.75)	19.84 20.51 (26.33)	85.66%	82.18% - 89.28%
T _{max} [§] (h)	3.00 (1.00 – 6.00)	2.00 (1.00 – 6.00)		
t _{1/2} ^ε (h)	6.12 (21.25)	6.33 (19.81)		

[†] Manufacturer: Purdue Pharma; origin (country of purchase): Canada.

[§] Expressed as the median (range) only

¹ Based on least-square means due to unbalanced study

A blind, randomized, 2-way crossover, bioequivalence study of ACT OXYCODONE CR and OxyContin[®] administered as 1 x 20 mg tablet in healthy males under fed conditions (n=28) was conducted and demonstrates the bioequivalence of ACT OXYCODONE CR and OxyContin[®].

Oxycodone (1 x 20 mg) From measured data Geometric LS Mean Arithmetic Mean (CV %)				
Parameter	ACT OXYCODONE CR 20 mg Tablet	OxyContin [†] 20 mg Tablet	% Ratio of Geometric LS Means	90% Confidence Interval
AUC _T (ng·h/mL)	226.29 232.52 (23.7)	237.00 243.91 (24.4)	95.48	92.92 – 98.11
AUC _∞ (ng·h/mL)	232.40 239.03 (24.2)	243.63 251.01 (24.9)	95.39	92.80 – 98.06
C _{max} (ng/mL)	22.1 22.61 (20.5)	23.41 24.01 (22.4)	94.44	90.79 – 98.23
T _{max} [§] (h)	4.50 (1.00 – 7.00)	3.50 (1.50 – 6.00)		
t _{1/2} (h)	5.28 (15.4)	5.28 (12.6)		

[†] Manufacturer: Purdue Pharma; origin (country of purchase): Canada.

[§] Expressed as the median (range) only

A blind, randomized, 2-way crossover, bioequivalence study of ACT OXYCODONE CR and OxyContin[®] administered as 1 x 20 mg tablet every 12 hours for 4 consecutive days in healthy subjects (n=28) was conducted and demonstrates the bioequivalence of ACT OXYCODONE CR and OxyContin[®].

Oxycodone (1 x 20 mg) From measured data Geometric LS Mean Arithmetic Mean (CV %)				
Parameter	ACT OXYCODONE CR 20 mg Tablet	OxyContin [†] 20 mg Tablet	% Ratio of Geometric LS Means	90% Confidence Interval
AUC _{TSS} (ng·h/mL)	18.91 19.93 (32.49)	19.91 20.94 (32.47)	95.00%	91.84% - 98.27%
C _{max SS} (ng/mL)	22.30 23.36 (31.36)	23.97 25.34 (38.61)	93.01%	88.96% - 97.25%
C _{min SS} (ng/mL)	8.54 9.38 (43.85)	8.93 9.74 (43.33)	95.66%	88.49% - 103.41%
T _{max} [§] (h)	2.50 (1.00 – 4.00)	2.50 (1.00 – 4.50)		
F1 (%)	87.56	90.99		

[†] Manufacturer: Purdue Pharma; origin (country of purchase): Canada.

[§] Expressed as the median (range) only

A blind, randomized, 2-way crossover, bioequivalence study of ACT OXYCODONE CR and OxyContin[®] administered as 1 x 40 mg tablet in healthy males under fasting conditions (n=34) was conducted and demonstrates the bioequivalence of ACT OXYCODONE CR and OxyContin[®].

Oxycodone (1 x 40 mg) From measured data Geometric LS Mean Arithmetic Mean (CV %)				
Parameter	ACT OXYCODONE CR 40 mg Tablet	OxyContin [†] 40 mg Tablet	% Ratio of Geometric LS Means	90% Confidence Interval
AUC _T (ng·h/mL)	415.86 421.34 (16.2)	412.89 417.50 (14.8)	100.72	97.39 - 104.16
AUC _∞ (ng·h/mL)	431.26 437.29 (16.7)	434.20 439.19 (15.1)	99.32	95.58 - 103.21
C _{max} (ng/mL)	38.00 38.48 (15.9)	37.24 37.55 (12.9)	102.04	98.46 - 105.76
T _{max} [§] (h)	2.77 (1.00 - 2.97)	2.00 (1.00 - 2.24)		
t _{1/2} (h)	5.77 (17.7)	6.51 (27.1)		

[†] Manufacturer: Purdue Pharma; origin (country of purchase): Canada.

[§] Expressed as the median (range) only

A blind, randomized, 2-way crossover, bioequivalence study of ACT OXYCODONE CR and OxyContin[®] administered as 1 x 40 mg tablet in healthy males under fed conditions (n=20) was conducted and demonstrates the bioequivalence of ACT OXYCODONE CR and OxyContin[®].

Oxycodone (1 x 40 mg) From measured data Geometric LS Mean Arithmetic Mean (CV %)				
Parameter	ACT OXYCODONE CR 40 mg Tablet	OxyContin [†] 40 mg Tablet	% Ratio of Geometric LS Means	90% Confidence Interval
AUC _T (ng·h/mL)	501.45 512.51 (20.8)	521.09 536.15 (24.4)	96.23	92.30 – 100.33
AUC _∞ (ng·h/mL)	510.99 522.56 (21.2)	536.04 552.42 (25.3)	95.33	91.39 – 99.43
C _{max} (ng/mL)	50.33 51.19 (19.4)	51.51 52.68 (20.9)	97.70	91.81 – 103.97
T _{max} (h)	4.75 (2.00 – 10.00)	3.25 (1.50 – 6.00)		
t _{1/2} [€] (h)	4.71 (14.5)	5.30 (13.8)		

[†] Manufacturer: Purdue Pharma; origin (country of purchase): Canada.

[§] Expressed as the median (range) only

A blind, randomized, 2-way crossover, bioequivalence study of ACT OXYCODONE CR and OxyContin[®] administered as 1 x 40 mg tablet every 12 hours for 4 consecutive days in healthy subjects (n=29) was conducted and demonstrates the bioequivalence of ACT OXYCODONE CR and OxyContin[®].

Oxycodone (1 x 40 mg) From measured data Geometric LS Mean Arithmetic Mean (CV %)				
Parameter	ACT OXYCODONE CR 40 mg Tablet	OxyContin [†] 40 mg Tablet	% Ratio of Geometric LS Means	90% Confidence Interval
AUC _{TSS} (ng·h/mL)	396.23 404.78 (19.93)	409.10 421.67 (24.01)	96.85%	93.21% - 100.63%
C _{max SS} (ng/mL)	47.19 47.93 (16.76)	49.68 50.94 (21.71)	94.99%	91.04% - 99.11%
C _{min SS} (ng/mL)	18.58 19.29 (25.89)	17.32 18.76 (36.95)	107.25%	97.83% - 117.59%
T _{max} [§] (h)	2.00 (1.00 – 4.50)	2.00 (1.00 – 4.50)		
F1 (%)	86.57	94.35		

[†] Manufacturer: Purdue Pharma; origin (country of purchase): Canada.

[§] Expressed as the median (range) only

A series of single-dose, randomized, crossover comparative bioavailability studies were conducted in healthy male and female volunteers to compare ACT Oxycodone CR tablets, 10, 20 and 40 mg strengths to the respective strengths of OxyNEO[®] (oxycodone hydrochloride) controlled release tablets, manufactured by Purdue Pharma, Canada, under both fasting and fed conditions. The results demonstrate that ACT Oxycodone CR tablets are bioequivalent to the respective strengths of OxyNEO[®] tablets. The results of these studies should not be interpreted or construed as equivalence, or Declaration of Equivalence of the products, as OxyNEO[®] tablets are formulated with inactive ingredients intended to make the tablet more difficult to manipulate for misuse and abuse.

DETAILED PHARMACOLOGY

Pharmacodynamics:

Oxycodone and related μ -agonist opioids produce their major effects on the CNS and the bowel by acting at specific saturable opioid receptors in the CNS and other tissues. The effects include analgesia, drowsiness, changes in mood, respiratory depression, cough suppression, decreased gastrointestinal motility, nausea, vomiting, and alterations of the endocrine and autonomic nervous systems.

Oxycodone receptor selectivity has not been extensively studied or characterized, and there appears to be a discrepancy between its weak affinity for opioid receptors and its potent antinociceptive activity.

Oxycodone has been shown to be 2 to 4 times more potent than morphine after both subcutaneous and intraperitoneal administration in rats. In clinical studies in patients with acute post-operative pain, oxycodone has been demonstrated to be twice as potent as morphine.

TOXICOLOGY

The LD₅₀ after subcutaneous administration of oxycodone in mice was 275 to 340 mg/kg. The lowest lethal dose has been reported to be 200 mg/kg after subcutaneous administration in mice. These values are similar to those obtained for morphine. In a preliminary 12 day study in rabbits, no drug related toxic effects were discernable at 5 mg/kg. Doses of 25, 75 and 150 mg/kg were associated with variable and transient pharmacotoxic effects typical of high dose opioid treatment in animals (decreased activity, decreased or absent defecation and convulsions).

Teratogenicity: Oxycodone had no effect on fertility or early embryonic development in male and female rats at doses as high as 8 mg/kg/day. Also, oxycodone did not induce any malformations in rats at doses as high as 8 mg/kg/day or in rabbits at doses as high as 125 mg/kg/day. Dose-related increases in developmental variations (increased incidences of extra presacral vertebrae and extra pairs of ribs) were observed in rabbits when the data for individual fetuses were analyzed. However, when the same data were analyzed using litters as opposed to individual fetuses, there was no dose-related increase in developmental variations although the incidence of extra presacral vertebrae remained significantly higher in the 125 mg/kg/day group compared to the control group. Since this dose level was associated with severe pharmacotoxic effects in the pregnant animals, the fetal findings may have been a secondary consequence of severe maternal toxicity.

In a study of peri- and post-natal development in rats, maternal body weight and food intake parameters were reduced for doses ≥ 2 mg/kg/day compared to the control group. Body weights were lower in the F1 generation from maternal rats in the 6 mg/kg/day dosing group. There were

no effects on physical, reflexological, or sensory developmental parameters or on behavioural and reproductive indices in the F1 pups (the NOEL for F1 pups was 2 mg/kg/day based on body weight effects seen at 6 mg/kg/day). There were no effects on the F2 generation at any dose in the study.

There are no adequate and well-controlled studies in pregnant women, and no studies on fertility or the post-natal effects of intrauterine exposure have been carried out.

Mutagenicity: Oxycodone was not mutagenic in the following assays: Ames *Salmonella* and *E. coli* test with and without metabolic activation at doses of up to 5,000 mcg, chromosomal aberration test in human lymphocytes in the absence of metabolic activation at doses of up to 1,500 mcg/mL and with activation 48 hours after exposure at doses of up to 5,000 mcg/mL, and in the *in vivo* bone marrow micronucleus test in mice at plasma levels of up to 48 mcg/mL.

Mutagenic results occurred in the presence of metabolic activation in the human chromosomal aberration test (at greater than or equal to 1,250 mcg/mL) at 24 but not 48 hours of exposure and in the mouse lymphoma assay at doses of 50 mcg/mL or greater with metabolic activation and at 400 mcg/mL or greater without metabolic activation. The data from these tests indicate that the genotoxic risk to humans may be considered low.

Carcinogenicity: Studies of oxycodone in animals to evaluate its carcinogenic potential have not been conducted owing to the length of clinical experience with the drug substance.

REFERENCES

1. Beaver WT, Wallenstein SL, Rogers A, Houde RW. Analgesic studies of codeine and oxycodone in patients with cancer. I. Comparisons of oral with intramuscular codeine and of oral with intramuscular oxycodone. *J Pharmacol Exp Ther* 1978; 207:92-100.
2. Beaver WT, Wallenstein SL, Rogers A, Houde RW. Analgesic studies of codeine and oxycodone in patients with cancer. II. Comparisons of intramuscular oxycodone with intramuscular morphine and codeine. *J Pharmacol Exp Ther* 1978; 207:101-8.
3. Chen ZR, Irvine RJ, Somogyi AA, Bochner F. Mu receptor binding of some commonly used opioids and their metabolites. *Life Sciences* 1991;48:2165-71.
4. Dickson PH, Lind A, Studts P, Nipper HC, Makoid M, Makoid M, et al. The routine analysis of breast milk for drugs of abuse in a clinical toxicology laboratory. *J Forensic Sci* 1994; 39(1):207-14.
5. Gimbel JS, Richards P, Portenoy RK. Controlled-release oxycodone for pain in diabetic neuropathy. A randomized controlled trial. *Neurology* 2003;60:927-34.
6. Glare PA, Walsh TC. Dose-ranging study of oxycodone for chronic pain in advanced cancer. *J Clin Oncology* 1993; 11:973-8.
7. Hagelberg NM, Nieminen TH, Saari TI, Neuvonen M, Neuvonen PJ, Laine K, et al. Voriconazole drastically increases exposure to oral oxycodone. *Eur J Clin Pharmacol* 2009;65(3):263-71.
8. Heiskanen T, Olkkola KT and Kalso E. Effects of blocking CYP2D6 on the pharmacokinetics and pharmacodynamics of oxycodone. *Clin Pharmacol Ther* 1998;64:603-11.
9. Kalso E, Vainio A. Morphine and oxycodone hydrochloride in the management of cancer pain. *Clin Pharmacol Ther* 1990; 47:639-46.
10. Leow KP, Smith MT, Williams B, Cramond T. Single-dose and steady-state pharmacokinetics and pharmacodynamics of oxycodone in patients with cancer. *Clin Pharmacol Ther* 1992; 52:487-95.
11. Leow KP, Wright AWE, Cramond T, Smith MT. Determination of the serum protein binding of oxycodone and morphine using ultrafiltration. *Ther Drug Monit* 1993; 15:440-7.

12. Mandema JW, Kaiko RF, Oshlack B, Reder RF, Stanski DR. Characterization and validation of a pharmacokinetic model for controlled-release oxycodone. *Br J Clin Pharmacol* 1996; 42:747-56.
13. Niemeinen TH, Hagelberg NM, Saari TI, Neuvonen M, Neuvonen PJ, Laine K, et al. Grapefruit juice enhances the exposure to oral oxycodone. *Basic Clin Pharmacol Toxicol* 2010;107:782-8.
14. Nieminen TH, Hagelberg NM, Saari TI, Neuvonen M, Laine K, Neuvonen PJ et al. St. John's wort greatly reduces the concentrations of oral oxycodone. *Eur J Pain* 2010;14: 854-9.
15. Pöyhiä R, Seppälä T, Olkkola KT, Kalso E. The pharmacokinetics and metabolism of oxycodone after intramuscular and oral administration to healthy subjects. *Br J Clin Pharmacol* 1992; 33:617-21.
16. Pöyhiä R, Kalso E, Seppälä T. Pharmacodynamic interactions of oxycodone and amitriptyline in healthy volunteers. *Current Ther Res* 1992; 51:739-49.
17. Watson CPN, Babul N. Efficacy of oxycodone in neuropathic pain. A randomized trial in postherpetic neuralgia. *Neurology* 1998;50:1837-41.
18. Watson CPN, Moulin D, Watt-Watson J, Gordon A, Eisenhoffer J. Controlled-release oxycodone relieves neuropathic pain: a randomized controlled trial in painful diabetic neuropathy. *Pain* 2003;105:71-8.
19. Weinstein SH, Gaylord JC. Determination of oxycodone in plasma and identification of a major metabolite. *J Pharm Sciences* 1979; 68:527-8.
20. ^NOXYNEO[®] (Oxycodone Hydrochloride Controlled Release Tablets) Product Monograph. Purdue Pharma. Control No. 209991, Date of Revision: February 7, 2018. NOTE: Since the Canadian Reference Product for ACT OXYCODONE CR is no longer marketed, the safety changes to the PM for ^NOXYNEO[®] is applicable, except for any tamper resistance claims as they do not apply to ACT OXYCODONE CR.

READ THIS FOR SAFE AND EFFECTIVE USE OF YOUR MEDICINE

PATIENT MEDICATION INFORMATION

^N ACT OXYCODONE CR
Oxycodone Hydrochloride
Controlled Release Tablets
5 mg, 10 mg, 20 mg, 40 mg, and 80 mg

Read this carefully before you start taking **ACT OXYCODONE CR** and each time you get a refill. This leaflet is a summary and will not tell you everything about this drug. Talk to your healthcare professional about your medical condition and treatment and ask if there is any new information about **ACT OXYCODONE CR**.

An important Risk Management Plan (RMP) for ACT OXYCODONE CR has been implemented to minimize potential risks and problems associated with possible abuse, dependence and diversion. For more information, please visit the following web site: <http://soar-essor.ca/> or call toll free at 1-866-254-6111.

Serious Warnings and Precautions

- **Even if you take ACT OXYCODONE CR as prescribed you are at risk for opioid addiction, abuse, and misuse that can lead to overdose and death. To understand your risk of opioid addiction, abuse, and misuse, you should speak to your prescriber (e.g., doctor).**
- **Life-threatening breathing problems can happen while taking ACT OXYCODONE CR, especially if not taken as directed. Babies are at risk of life-threatening breathing problems if their mothers take opioids while pregnant or nursing.**
- **Never give anyone your ACT OXYCODONE CR. They could die from taking it. If a person has not been prescribed ACT OXYCODONE CR, taking even one dose can cause a fatal overdose. This is especially true for children.**
- **If you took ACT OXYCODONE CR while you were pregnant, whether for short or long periods of time or in small or large doses, your baby can suffer life-threatening withdrawal symptoms after birth. This can occur in the days after birth and for up to 4 weeks after delivery. If your baby has any of the following symptoms:**
 - **has changes in their breathing (such as weak, difficult or fast breathing)**
 - **is unusually difficult to comfort**
 - **has tremors (shakiness)**
 - **has increased stools, sneezing, yawning, vomiting, or fever****Seek immediate medical help for your baby.**
- **Taking ACT OXYCODONE CR with other opioid medicines, benzodiazepines, alcohol, or other central nervous system depressants (including street drugs) can cause severe drowsiness, decreased awareness, breathing problems, coma, and death.**
-

What is ACT OXYCODONE CR used for?

ACT OXYCODONE CR is used for the long-term management of pain, when:

- the pain is severe enough to require daily, around-the-clock pain medication
- the doctor determines that other treatment options are not able to effectively manage your pain

ACT OXYCODONE CR is NOT used (“as needed”) to treat pain that you only have once in a while.

How does ACT OXYCODONE CR work?

ACT OXYCODONE CR is an oral controlled release tablet that slowly releases oxycodone hydrochloride over a 12 hour period.

ACT OXYCODONE CR contains oxycodone which is a pain medication belonging to the class of medicines known as opioids which includes codeine, fentanyl and morphine. It relieves pain by acting on specific nerve cells of the spinal cord and brain.

What are the ingredients in ACT OXYCODONE CR?

Medicinal ingredients: oxycodone hydrochloride

Non-medicinal ingredients: colloidal silicon dioxide, hydroxyl propyl methyl cellulose, magnesium stearate, microcrystalline cellulose, polyvinyl acetate and povidone. Coating ingredients include: hypromellose, hydroxypropyl cellulose, polyethylene glycol, titanium dioxide, iron oxide yellow (20 mg, 40 mg and 80 mg), iron oxide red (20 mg and 40 mg) and FD&C Blue Aluminium Lake (5 mg and 80 mg).

ACT OXYCODONE CR comes in the following dosage forms:

Controlled release tablets, 5 mg, 10 mg, 20 mg, 40 mg and 80 mg.

Do not use ACT OXYCODONE CR if:

- your doctor did not prescribe it for you;
- you are allergic to oxycodone hydrochloride, other opioids, or any of the other ingredients of ACT OXYCODONE CR;
- you have mild or short term pain that can be controlled by the occasional use of pain medications, including those available without a prescription;
- **you have difficulty in swallowing or have been diagnosed with narrowing of the esophagus;**
- you have severe asthma, trouble breathing, or other lung problems;
- you have a condition where the small bowel does not work properly (paralytic ileus) or
- you have severe pain in your abdomen ;you have a head injury;
- you are at risk for seizures;
- you suffer from alcoholism;
- you are taking, or have taken within the past 2 weeks, a monoamine oxidase inhibitor medication (e.g., phenelzine sulphate, tranylcypromine sulphate, moclobemide or selegiline);
- you are pregnant or plan to become pregnant, breast-feeding, or in labour;
- you are under 18 years of age;
- you are going to have, or recently had, a planned surgery.

To help avoid side effects and ensure proper use, talk to your healthcare professional before you take ACT OXYCODONE CR. Talk about any health conditions or problems you may have, including if you:

- have a history of illicit or prescription drug or alcohol abuse;
- have severe kidney, liver or lung disease;
- have heart disease;
- have low blood pressure;
- have past or current depression;
- have problems with your thyroid, adrenal or prostate gland;

- suffer from chronic or severe constipation;
- have, or had in the past hallucinations or other severe mental problems;
- suffer from migraines.

Other warnings you should know about:

Opioid dependence and addiction: There are important differences between physical dependence and addiction. It is important that you talk to your doctor if you have questions or concerns about abuse, addiction or physical dependence.

Pregnancy, nursing, labour and delivery: Do not use ACT OXYCODONE CR while pregnant, nursing, during labour or delivery. Opioids can be transferred to your baby through breast milk, or while still in the womb. ACT OXYCODONE CR can then cause life-threatening breathing problems in your unborn baby or nursing infant.

Driving and using machines: Before you do tasks which may require special attention, you should wait until you know how you react to ACT OXYCODONE CR. ACT OXYCODONE CR can cause:

- drowsiness;
- dizziness;
- light-headedness.

This can usually occur after you take your first dose and when dose is increased.

Disorder of the adrenal gland: You may develop a disorder of the adrenal gland called adrenal insufficiency. This means that your adrenal gland is not making enough certain hormones. You may experience symptoms such as:

- nausea, vomiting;
- feeling tired, weak or dizzy;
- decreased appetite.

You may be more likely to have problems with your adrenal gland if you have been taking opioids for longer than one month. Your doctor may do tests, give you another medication, and slowly take you off ACT OXYCODONE CR.

Serotonin Syndrome: ACT OXYCODONE CR can cause serotonin syndrome, a rare but potentially life-threatening condition. It can cause serious changes in how your brain, muscles and digestive system work. You may develop serotonin syndrome if you take ACT OXYCODONE CR with certain antidepressants or migraine medications.

Serotonin syndrome symptoms include:

- fever, sweating, shivering, diarrhea, nausea, vomiting;
- muscle shakes, jerks, twitches or stiffness, overactive reflexes, loss of coordination;
- fast heartbeat, changes in blood pressure;
- confusion, agitation, restlessness, hallucinations, mood changes, unconsciousness, and coma.

Sexual Function/Reproduction: Long term use of opioids may lead to a decrease in sex hormone levels. It may also lead to low libido (desire to have sex), erectile dysfunction or being infertile.

Tell your healthcare professional about all the medicines you take, including any drugs, vitamins, minerals, natural supplements or alternative medicines.

The following may interact with ACT OXYCODONE CR:

- alcohol, including prescription and non-prescription medications containing alcohol. Do not drink alcohol while taking ACT OXYCODONE CR. This can lead to drowsiness, depressed breathing, unusually slow or weak breathing, serious side effects or a fatal overdose;
- other sedative drugs which may enhance the drowsiness caused by ACT OXYCODONE CR;
- other opioid analgesics (for pain);
- general anesthetics (used during surgery);
- drugs used to help you sleep or to reduce anxiety;
- antidepressants (for depression and mood disorders). Do not take ACT OXYCODONE CR with monoamine oxidase (MAO) inhibitors or if you have taken MAO inhibitors in the last 14 days before treatment with ACT OXYCODONE CR;
- drugs used to treat serious mental or emotional disorders, such as schizophrenia;
- antihistamines (for allergies);
- anti-emetics (for prevention of vomiting);
- drugs used to treat muscle spasms and back pain;
- warfarin and other coumarin anticoagulants (for prevention/treatment of blood clots);
- some heart medications (beta-blockers);
- anti-retroviral,azole-anti-fungal and macrolide-antibiotic drugs;
- grapefruit juice;
- St. John's Wort.

How to take ACT OXYCODONE CR:

Swallow whole. Do not cut, break, chew, dissolve or crush ACT OXYCODONE CR tablets before swallowing since this can lead to the release and absorption of an excessive dose of oxycodone which can seriously harm you.

Usual Adult Starting Dose:

Dosage is individualized. Be sure to follow your doctor's dosing instructions exactly. Taking higher doses can lead to more side effects and a greater chance of overdose.

Take the dose prescribed by your doctor. ACT OXYCODONE CR tablets should be taken every 12 hours with a glass of water.

Your dose of ACT OXYCODONE CR will be clearly labelled on the medication bottle. Be sure to follow the directions on the label exactly; this is very important. Do not increase or decrease your dose without consulting your doctor. If your dosage is changed by your doctor, be sure to write it down at the time your doctor calls or sees you, and follow the new directions exactly. Review your pain regularly with your doctor to determine if you still need ACT OXYCODONE CR. Be sure to use ACT OXYCODONE CR only for the condition for which it was prescribed.

Stopping your Medication:

You should not stop taking ACT OXYCODONE CR all at once if you have been taking it for more than a few days.

Your doctor will monitor and guide you on how to slowly stop taking ACT OXYCODONE CR. You should do it slowly to avoid uncomfortable symptoms such as having:

- body aches
- diarrhea
- goosebumps
- loss of appetite
- nausea
- feeling nervous or restless
- runny nose
- sneezing
- tremors or shivering
- stomach cramps
- rapid heart rate (tachycardia)
- having trouble with sleeping
- an unusual increase in sweating
- heart palpitations
- an unexplained fever
- weakness
- yawning

By reducing or stopping your opioid treatment, your body will become less used to opioids. If you start treatment again, you will need to start at the lowest dose. You may overdose if you restart at the last dose you took before you slowly stopped taking ACT OXYCODONE CR.

Refilling Prescriptions for ACT OXYCODONE CR:

A new written prescription is required from your doctor each time you need more ACT OXYCODONE CR. Therefore, it is important that you contact your doctor before your current supply runs out.

Only obtain prescriptions for this medicine from the doctor in charge of your treatment. Do not seek prescriptions from other doctors unless you switch to another doctor for your pain management.

Overdose:

If you think you have taken too much ACT OXYCODONE CR, contact your healthcare professional, hospital emergency department or regional Poison Control Centre immediately, even if there are no symptoms.

Signs of overdose may include:

- unusually slow or weak breathing
- dizziness
- confusion
- extreme drowsiness

Missed Dose:

It is important that you do not miss any doses. If you miss a dose, take your next dose at your usual time. You should always try to get back on track with your regular dosing schedule (e.g., 8 o'clock in the morning and 8 o'clock in the evening). If you miss several doses in a row, talk to your doctor before restarting your medication.

What are possible side effects from using ACT OXYCODONE CR?

These are not all the possible side effects you may feel when taking ACT OXYCODONE CR. If you experience any side effects not listed here, contact your healthcare professional.

Side effects may include:

- Constipation
- Dizziness
- Drowsiness

- Dry mouth
- Headache
- Itching
- Weakness, uncoordinated muscle movement
- Nausea and/or vomiting, or poor appetite
- Sweating
- Insomnia
- Abdominal pain
- Fever
- Diarrhea
- Indigestion
- Tremor

- Abnormal dreams or thoughts
- Anxiety
- Confusion
- Depression
- Nervousness
- Rash
- Difficulty breathing
- Low sex drive, impotence (erectile dysfunction), infertility

Talk with your doctor or pharmacist about ways to prevent constipation when you start using ACT OXYCODONE CR.

If nausea and vomiting become troublesome during prolonged therapy with ACT OXYCODONE CR, talk to your doctor or pharmacist.

Serious side effects and what to do about them				
Symptom / effect		Talk to your healthcare professional		Stop taking drug and get immediate medical help
		Only if severe	In all cases	
RARE	Overdose: hallucinations, confusion, inability to walk normally, slow or weak breathing, extreme sleepiness, sedation, or dizziness, floppy muscles/low muscle tone, cold and clammy skin.			✓
	Respiratory Depression: slow, shallow or weak breathing.			✓
	Allergic Reaction: rash, hives, swelling of the face, lips, tongue or throat, difficulty swallowing or breathing.			✓
	Bowel Blockage (Impaction): abdominal pain, severe constipation, nausea.			✓
	Withdrawal: nausea, vomiting, diarrhea, anxiety, shivering, cold		✓	

Serious side effects and what to do about them			
Symptom / effect	Talk to your healthcare professional		Stop taking drug and get immediate medical help
	Only if severe	In all cases	
and clammy skin, body aches, loss of appetite, sweating.			
Fast, Slow or Irregular heartbeat: Heart palpitations.		✓	
Low Blood Pressure: dizziness, fainting, light-headedness.	✓		
Serotonin Syndrome: agitation or restlessness, loss of muscle control or muscle twitching, tremor, diarrhea.			✓

If you have a troublesome symptom or side effect that is not listed here or becomes bad enough to interfere with your daily activities, talk to your healthcare professional.

Reporting Side Effects
<p>You can report any suspected side effects associated with the use of health products to Health Canada by:</p> <ul style="list-style-type: none"> • Visiting the Web page on Adverse Reaction Reporting (https://www.canada.ca/en/health-canada/services/drugs-health-products/medeffect-canada/adverse-reaction-reporting.html) for information on how to report online, by mail or by fax; or • Calling toll-free at 1-866-234-2345. <p><i>NOTE: Contact your health professional if you need information about how to manage your side effects. The Canada Vigilance Program does not provide medical advice.</i></p>

Storage:

- **Keep unused or expired ACT OXYCODONE CR in a secure place to prevent theft, misuse or accidental exposure.**
- Store ACT OXYCODONE CR tablets between 15°C and 30°C. Keep container tightly closed. Protect from moisture.
- **Keep ACT OXYCODONE CR under lock, out of sight and reach of children and pets.**
- **Never take medicine in front of small children as they will want to copy you.**

Accidental ingestion by a child is dangerous and may result in death. If a child accidentally takes ACT OXYCODONE CR, get emergency help right away.

Disposal:

ACT OXYCODONE CR should never be thrown into household trash, where children and pets may find it. It should be returned to a pharmacy for proper disposal.

If you want more information about ACT OXYCODONE CR:

- Talk to your healthcare professional
- Find the full Product Monograph that is prepared for healthcare professionals and includes this Patient Medication Information by visiting the <https://www.canada.ca/en/health-canada/services/drugs-health-products/drug-products/drug-product-database.html> ; the manufacturer's website <http://www.tevacanada.com>; or by calling 1-800-268-4127 ext. 3; or email druginfo@tevacanada.com.

This leaflet was prepared by Teva Canada Limited, Toronto, Ontario, Canada M1B 2K9.

Last Revised: October 1, 2019