

PRODUCT MONOGRAPH
INCLUDING PATIENT MEDICATION INFORMATION

Pr **JORVEZA™**

Budesonide Orodispersible Tablets

1 mg, Oral

Corticosteroids acting locally

AVIR Pharma Inc.
660 Boulevard Industriel
Blainville, Québec
J7C 3V4

www.avirpharma.com

Submission Control No: 227232

Date of Initial Approval:
November 5, 2019

JORVEZA™ is a trademark of Dr. Falk Pharma GmbH, used under license by AVIR Pharma Inc.

RECENT MAJOR LABEL CHANGES

Not applicable.

TABLE OF CONTENTS

PART I: HEALTH PROFESSIONAL INFORMATION	4
1 INDICATIONS	4
1.1 Pediatrics (< 18 years of age).....	4
2 CONTRAINDICATIONS	4
3 DOSAGE AND ADMINISTRATION	4
3.1 Dosing Considerations	4
3.2 Recommended Dose and Dosage Adjustment	4
3.3 Administration.....	5
3.4 Missed Dose	5
4 OVERDOSAGE.....	5
5 DOSAGE FORMS, STRENGTHS, COMPOSITION AND PACKAGING	5
6 WARNINGS AND PRECAUTIONS	6
6.1 Special Populations	7
6.1.1 Pregnant Women	7
6.1.2 Breast-feeding.....	8
6.1.3 Pediatrics	8
7 ADVERSE REACTIONS	8
7.1 Adverse Reaction Overview	8
7.2 Clinical Trial Adverse Reactions	8
7.3 Post-Market Adverse Reactions	9
8 DRUG INTERACTIONS	10
8.1 Drug-Drug Interactions	10
8.2 Drug-Food Interactions.....	11
8.3 Drug-Herb Interactions	11
8.4 Drug-Laboratory Test Interactions	11
9 ACTION AND CLINICAL PHARMACOLOGY.....	11
9.1 Mechanism of Action	11
9.2 Pharmacodynamics.....	11
9.3 Pharmacokinetics	12
10 STORAGE, STABILITY AND DISPOSAL.....	13

PART II: SCIENTIFIC INFORMATION	14
11 PHARMACEUTICAL INFORMATION	14
12 CLINICAL TRIALS	15
12.1 Trial Design and Study Demographics	15
12.2 Study Results	15
13 NON-CLINICAL TOXICOLOGY	16
PATIENT MEDICATION INFORMATION	19

PART I: HEALTH PROFESSIONAL INFORMATION

1 INDICATIONS

JORVEZA (budesonide) is indicated for:

- Induction of clinico-pathological remission in adults with eosinophilic esophagitis (EoE).

1.1 Pediatrics (< 18 years of age)

No data are available. Therefore, Health Canada has not authorised an indication for pediatric use.

1.2 Geriatrics (≥65 years of age)

JORVEZA has not been adequately studied in elderly ≥ 65 years of age. See **WARNINGS AND PRECAUTIONS, Special Populations**.

2 CONTRAINDICATIONS

JORVEZA is contraindicated in patients who

- Are hypersensitive to this drug or to any ingredient in the formulation, including any non-medicinal ingredient, or component of the container. For a complete listing, see **DOSAGE FORMS, STRENGTHS, COMPOSITION AND PACKAGING**.
- Have uncontrolled infections
- Have active tuberculosis

3 DOSAGE AND ADMINISTRATION

3.1 Dosing Considerations

- **Renal impairment:** There are no data available for patients with renal impairment. Because budesonide is not excreted via the kidneys, patients with mild to moderate impairment may be treated with caution with the same doses as patients without renal impairment. JORVEZA is not recommended for use in patients with severe renal impairment.
- **Hepatic impairment:** During treatment of patients with hepatic impairment with other budesonide containing products, budesonide levels were increased. However, no systematic study investigating different levels of hepatic impairment is available. Patients with hepatic impairment should not be treated with JORVEZA (see **WARNINGS AND PRECAUTIONS, Special Populations** and **ACTION AND CLINICAL PHARMACOLOGY, Pharmacokinetics, Special Populations and Conditions, Hepatic Impairment**).

3.2 Recommended Dose and Dosage Adjustment

The treatment with JORVEZA should be initiated by a physician experienced in the diagnosis and treatment of eosinophilic esophagitis.

The recommended daily dose is 2 mg budesonide as one 1-mg-tablet in the morning and one in the evening.

The usual duration of treatment is 6 weeks.

Health Canada has not authorized an indication for pediatric use.

3.3 Administration

- The orodispersible tablet should be taken after a meal.
- The tablet should not be taken with liquid or food.
- There should be at least 30 minutes before eating or drinking or performing oral hygiene. Any oral solutions, sprays or chewable tablets should be used at least 30 minutes before or after administration of JORVEZA.
- It should be placed on the tip of the tongue and gently pressed against the top of the mouth, where it will dissolve. This will usually take about two minutes. The effervescence process of the tablet starts after JORVEZA comes into contact with saliva and stimulates the production of further saliva. The dissolved material should be swallowed with saliva little by little while the orodispersible tablet disintegrates.
- The orodispersible tablet should not be chewed or swallowed undissolved. These measures ensure optimal exposure of the esophageal mucosa to the active substance.

3.4 Missed Dose

If a dose is missed, treatment should be continued at the prescribed dosage. A double dose should not be used to make up for a forgotten dose.

4 OVERDOSAGE

In case of short-term overdose, no emergency medical treatment is required. There is no specific antidote. Subsequent treatment should be symptomatic and supportive.

For management of a suspected drug overdose, contact your regional poison control centre.

5 DOSAGE FORMS, STRENGTHS, COMPOSITION AND PACKAGING

Table 1: Dosage forms, strengths, composition and packaging.

Route of Administration	Dosage Form / Strength/Composition	Non-medicinal Ingredients
Oral	orodispersible tablet 1 mg	Anhydrous monosodium citrate, Disodium hydrogen citrate, Docusate sodium, Macrogol 6000, Magnesium stearate, Mannitol, Povidone K25, Sodium hydrogen carbonate and Sucralose.

JORVEZA is supplied as a white or almost white, round, biplane orodispersible tablet, with diameter of 6.9-7.3 mm and height of 1.8-2.6 mm.

Alu / Alu-blister. Pack sizes: 20, 30 or 90 orodispersible tablets. Some pack sizes may not be marketed.

6 WARNINGS AND PRECAUTIONS

General

Concomitant treatment with ketoconazole or other CYP3A4 inhibitors should be avoided (see **Drug-Drug Interactions, CYP3A4 inhibitors**).

Patients with tuberculosis, hypertension, diabetes mellitus, osteoporosis, peptic ulcer, glaucoma, cataract, family history of diabetes or family history of glaucoma condition may be at higher risk of experiencing systemic glucocorticosteroid adverse reactions (see **ADVERSE REACTIONS**) and should therefore be monitored for the occurrence of such effects. In these patients, caution should be exercised and the benefits of an oral glucocorticosteroid must be weighed against its risks. JORVEZA should not be used in patients with active tuberculosis, or uncontrolled infection.

Driving and Operating Machinery

JORVEZA has no or negligible influence on the ability to drive and use machines.

Endocrine and Metabolism

Glucocorticoids may cause suppression of the hypothalamus-pituitary-adrenal (HPA) axis and reduce the stress response. Where patients are subject to surgery or other stresses, supplementary systemic glucocorticoid treatment is recommended.

Systemic effects of steroids may occur, particularly when prescribed at high doses and for prolonged periods. Such effects may include Cushing's syndrome, adrenal suppression, growth retardation, decreased bone mineral density, cataract, glaucoma and very rarely psychiatric/behavioral effects (see **ADVERSE REACTIONS**).

Particular care is needed in patients who are transferred from a glucocorticosteroid treatment with higher systemic effect (e.g., prednisolone). Tapering of the dose of such conventional therapy when treatment with JORVEZA is initiated and monitoring of adrenocortical function may be needed in these patients. Some patients feel unwell during withdrawal (e.g., pain in muscles and joints), or experience flare up of allergies previously controlled by the conventional systemic corticosteroid drug. A general insufficient glucocorticosteroid effect should be suspected if, in rare cases, symptoms such as tiredness, headache, nausea and vomiting occur. In these cases a temporary adjustment in the dose of systemic glucocorticosteroids may sometimes be necessary.

Hepatic/Biliary/Pancreatic

Reduced liver function may affect the elimination of budesonide, causing higher systemic exposure. The risk of adverse reactions (systemic glucocorticosteroid effects) will be increased. However, no systematic data are available. Patients with hepatic impairment should therefore not be treated with JORVEZA.

Immune

Infections

Suppression of the inflammatory response and immune function increases the susceptibility to infections and their severity. Symptoms of infections can be atypical or masked. JORVEZA should not be used in patients with uncontrolled infections, or with active tuberculosis.

In clinical studies conducted with JORVEZA, oral, oropharyngeal and esophageal candida infections have been observed with a high frequency (see **ADVERSE REACTIONS**).

If indicated, symptomatic candidiasis of the mouth and throat can be treated with topical or systemic anti-fungal therapy whilst still continuing treatment with JORVEZA.

Chickenpox, herpes zoster and measles can have a more serious or even fatal course in patients treated with glucocorticosteroids. In patients who have not had these diseases, the vaccination status should be checked, and particular care should be taken to avoid exposure. If patients are infected or suspected of being infected, consider reduction or discontinuation of glucocorticoid treatment.

Vaccines

The co-administration of live vaccines and glucocorticosteroids should be avoided as this is likely to reduce the immune response to vaccines. The antibody response to other vaccines may be diminished.

Ophthalmologic

Visual disturbance may be reported with systemic and topical corticosteroid use. If a patient presents with symptoms such as blurred vision or other visual disturbances, the patient should be considered for referral to an ophthalmologist for evaluation of possible causes which may include cataract, glaucoma or rare diseases such as central serous chorioretinopathy (CSCR) which have been reported after use of systemic and topical corticosteroids.

Psychiatric

Particular care is required when considering the use of systemic corticosteroids in patients with current or previous history of severe affective disorders, or such history in any of the first-degree relatives. Systemic effects of steroids may occur, particularly when prescribed at high doses and for prolonged periods. Such effects may include psychiatric/behavioral effects.

Renal impairment:

There are currently no data available for patients with renal impairment. Because budesonide is not excreted via the kidneys, patients with mild to moderate impairment may be treated with caution with the same doses as patients without renal impairment. JORVEZA is not recommended for use in patients with severe renal impairment.

Sexual Health

Fertility

There are no data on the effect of budesonide on human fertility. Fertility was unaffected following budesonide treatment in animal studies (see **NON-CLINICAL TOXICOLOGY, Reproductive and Developmental Toxicity**).

6.1 Special Populations

6.1.1 Pregnant Women

Administration during pregnancy should be avoided unless there are compelling reasons for therapy with JORVEZA. In the absence of adequate studies in pregnant women, JORVEZA

should be used during pregnancy only if the potential benefits to the mother clearly outweigh the risks to the fetus.

In pregnant animals, budesonide, like other glucocorticosteroids, has been shown to cause fetal malformations and abnormalities (see **NON-CLINICAL TOXICOLOGY, Reproductive and Developmental Toxicity**). The relevance of these findings to humans has not been established.

In animal studies, budesonide was found to cross the placental barrier, therefore, infants born of mothers who have received glucocorticosteroids during pregnancy should be carefully observed for hypoadrenalism. Note that the maximal concentration of budesonide in plasma is expected to be higher in the treatment with JORVEZA compared to inhaled budesonide.

6.1.2 Breast-feeding

Budesonide is excreted in human milk. However, only minor effects on the breast-fed child are anticipated after oral use of JORVEZA within the therapeutic range. A decision must be made whether to discontinue breast-feeding or to discontinue/abstain from budesonide therapy taking into account the benefit and potential risks of breast feeding for the child and the benefit of therapy for the woman.

6.1.3 Pediatrics

The safety and efficacy of JORVEZA have not been studied in pediatric populations. JORVEZA should not be used in children and adolescents under the age of 18 years. Corticosteroids, including JORVEZA, may reduce growth velocity in children.

6.1.4 Geriatrics (≥ 65 years of age):

There is no sufficient and adequate data in subjects ≥65 years of age. Caution should be exercised in elderly patients due to the potential for decreased hepatic, renal, or cardiac function, or due to concomitant disease or therapies.

7 ADVERSE REACTIONS

7.1 Adverse Reaction Overview

The nature and frequency of adverse events observed in the JORVEZA 1 mg BID group were consistent with the known safety profile of budesonide. In the double-blind treatment phase, the majority of adverse events were of mild or moderate intensity. Fungal infections (Candidiasis) of the mouth, pharynx and the esophagus were the most frequently observed adverse reactions in clinical studies with JORVEZA. In the clinical study BUL-1/EEA, a total of 10 out of 87 patients (11.5%) exposed to JORVEZA (double-blind and open-label extension phase) experienced cases of suspected fungal infections associated with clinical symptoms, which were all of mild intensity, except for one adverse event. The total number of infections (including those diagnosed by endoscopy and histology without symptoms) was 33, occurring in 27 out of 87 patients (31%).

There was no adverse event leading to discontinuation in the JORVEZA treatment group.

7.2 Clinical Trial Adverse Reactions

Because clinical trials are conducted under very specific conditions, the adverse reaction rates observed in the clinical trials may not reflect the rates observed in practice and should not be compared to the rates in the clinical trials of another drug. Adverse reaction information from

clinical trials is useful for identifying drug-related adverse events and for approximating rates.

Adverse reactions observed in the pivotal study (induction of remission) are listed in the **Table 2** below.

Table 2: Adverse drug reactions by System Organ Class and Preferred Term – 6-week Double-Blind Phase (Safety Analysis Set)

System Organ Class	JORVEZA n = 59 (%)	Placebo n = 29 (%)
Gastrointestinal disorders		
Gastroesophageal reflux disease	2 (3.4%)	0
Dyspepsia	1 (1.7%)	0
Feces soft	1 (1.7%)	0
Nausea	1 (1.7%)	0
Infections and infestations		
Candida infection	2 (3.4%)	0
Esophageal candidiasis	10 (16.9%)	0
Oral candidiasis	2 (3.4%)	0
Oropharyngeal candidiasis	3 (5.1%)	0
Investigations		
Blood cortisol decreased	3 (5.1%)	0
Psychiatric disorders		
Insomnia	0	1 (3.4%)
Renal and urinary disorders		
Polyuria	1 (1.7%)	0
Vascular disorders		
Hypertension	1 (1.7%)	0

A similar adverse event profile was reported during the 6-week open-label induction phase for eligible clinico-pathological non-remitters (n=51, of which 23 were treated with JORVEZA for a total of 12 weeks). However, it is not possible to conclude on the safety or efficacy of JORVEZA beyond 6 weeks of treatment. Additional adverse drug reactions reported (1 patient each) include lymphocytosis, abdominal pain, gastric ulcer, lip edema, melaena and oral paresthesia.

Other adverse reaction observed in clinical studies with JORVEZA, with a frequency of common ($\geq 1/100$ to $< 1/10$): fatigue, headache.

7.3 Post-Market Adverse Reactions

Adverse Events Seen with JORVEZA During Post-Marketing Surveillance:

Nervous system disorders: Dysgeusia

The following known undesirable effects of the therapeutic class (corticosteroids, budesonide) could also occur with JORVEZA (frequency = not known):

Endocrine disorders:	Cushing's syndrome, moon-face, truncal obesity, adrenal suppression, growth retardation in children, disturbance of sex hormone secretion (e.g. amenorrhea, hirsutism, impotence)
Eye disorders:	Glaucoma, cataract (including subcapsular cataract), blurred vision, central serous chorioretinopathy (CSCR), see WARNINGS AND PRECAUTIONS, Ophthalmologic
Gastrointestinal disorders:	Dyspepsia, duodenal or gastric ulcers, pancreatitis, constipation
General disorders and administration site conditions:	Malaise
Immune system disorders:	Immune suppression (e.g. increased risk of infection)
Metabolism and nutrition disorders:	Hypokalaemia, hyperglycaemia, reduced glucose tolerance, diabetes mellitus, sodium retention with edema
Musculoskeletal and connective tissue disorders:	Muscle and joint pain, muscle weakness and twitching, osteoporosis, osteonecrosis
Nervous system disorders:	Dysgeusia, pseudotumor cerebri including papilloedema in adolescents
Psychiatric disorders:	Depression, irritability, euphoria, psychomotor hyperactivity, anxiety, aggression
Skin and subcutaneous tissue disorders:	Allergic exanthema, petechiae, delayed wound healing, contact dermatitis, ecchymosis, steroid acne, red striae
Vascular disorders:	Increased risk of thrombosis, vasculitis (withdrawal syndrome after long-term therapy), hypertension

8 DRUG INTERACTIONS

8.1 Drug-Drug Interactions

CYP3A4 inhibitors

Co-treatment with potent CYP3A inhibitors such as ketoconazole, ritonavir, itraconazole, clarithromycin and cobicistat may cause a marked increase of the plasma concentration of budesonide and is expected to increase the risk of systemic adverse reactions. Therefore, concomitant use should be avoided unless the benefit outweighs the increased risk of systemic corticosteroid side effects, in which case patients should be monitored for systemic corticosteroid side effects.

Ketoconazole 200 mg once daily orally increased the plasma concentration of budesonide (3 mg single dose) approximately 6-fold during concomitant administration. When ketoconazole was administered approximately 12 hours after budesonide, the plasma concentration of budesonide increased approximately 3-fold.

Oestrogens, oral contraceptives

Elevated plasma concentrations and enhanced effects of glucocorticosteroids have been reported in women also receiving oestrogens or oral contraceptives. No such effect has been observed with budesonide and concomitant intake of low-dose combination oral contraceptives.

Cardiac glycosides

The action of glycoside can be potentiated by potassium deficiency which is a potential and known adverse reaction of glucocorticoids.

Saluretics

Concomitant use of glucocorticoids may result in enhanced potassium excretion and aggravated hypokalaemia.

8.2 Drug-Food Interactions

Inhibitors of CYP3A4 such as grapefruit juice may cause a marked increase of the plasma concentration of budesonide. Therefore, concomitant use should be avoided.

8.3 Drug-Herb Interactions

Interactions with herbal products have not been established.

8.4 Drug-Laboratory Test Interactions

Interference with serological testing

Because adrenal function may be suppressed by treatment with budesonide, an ACTH stimulation test for diagnosing pituitary insufficiency might show false results (low values).

9 ACTION AND CLINICAL PHARMACOLOGY

9.1 Mechanism of Action

Budesonide is a non-halogenated glucocorticosteroid, which acts primarily as anti-inflammatory via binding to the glucocorticoid receptor. JORVEZA is a partly locally acting corticosteroid. The exact mechanism of action in the treatment of EoE is not fully understood. JORVEZA, as a corticosteroid, may inhibit antigen-stimulated secretion of many pro-inflammatory signal molecules, which may result in a significant reduction of the esophageal eosinophilic inflammatory infiltrate.

9.2 Pharmacodynamics

The primary pharmacodynamic effect of budesonide is its anti-inflammatory activity. Budesonide has a high glucocorticoid effect and a weak mineralocorticoid effect, and the affinity of budesonide to GCS receptors, which reflects the intrinsic potency of the drug, is about 200-fold that of cortisol and 15-fold that of prednisolone.

Effect on hypothalamus-pituitary-adrenal and endogenous cortisol levels

Treatment with systemically active glucocorticosteroid is associated with a suppression of endogenous cortisol concentrations and impairment of the hypothalamus-pituitary-adrenal (HPA) axis function.

Following 6 weeks of JORVEZA 1 mg BID treatment in patients with EoE, the rate of patients with adverse events of decreased plasma cortisol levels was 5% with JORVEZA and 0% with placebo.

Pharmacokinetics data in healthy subjects following one week of treatment with 4 mg daily of budesonide orodispersible tablets showed a decrease in plasma and urinary cortisol levels.

9.3 Pharmacokinetics

Absorption: Following administration of JORVEZA, budesonide is rapidly absorbed. Pharmacokinetic data following administration of single doses of 1 mg budesonide to fasted healthy subjects show a median lag time of 0.17 hours (range 0.00 – 0.33 hours) and a median time to peak plasma concentration of 1.00 hour (range 0.50 – 2.00 hours). The mean peak plasma concentration (\pm standard deviation) was 0.44 ± 0.31 ng/mL, the area under the plasma-concentration–time curve (AUC_{0-12}) was 1.44 ± 0.31 hr*ng/mL.

Single dose pharmacokinetic data in fasted patients with EoE are available with 4 mg budesonide: Median lag-time was 0.00 hours (range 0.00 – 0.17), median time to peak plasma concentration was 1.00 hour (range 0.67 – 2.00 hours); peak plasma concentration was 2.56 ± 1.36 ng/mL, and AUC_{0-12} was 8.96 ± 4.21 hr*ng/mL.

Patients showed a 35% increase in peak plasma concentrations and a 60% increase in AUC_{0-12} compared to healthy subjects.

Distribution: The volume of distribution following administration of 1 mg budesonide to healthy subjects was 35.52 ± 14.94 L/kg and 42.46 ± 23.90 L/kg following administration of 4 mg budesonide to patients with EoE. Plasma protein binding is on average 85-90%.

Metabolism: Metabolism of budesonide is decreased in EoE patients compared to healthy subjects resulting in increased plasma concentrations of budesonide.

Budesonide undergoes extensive biotransformation by CYP3A4 in the mucosa of the small intestine and in the liver to metabolites of low glucocorticosteroid activity. The glucocorticosteroid activity of the major metabolites, 6 β -hydroxybudesonide and 16 α -hydroxyprednisolone, is less than 1% of that of budesonide. CYP3A5 does not contribute significantly to the metabolism of budesonide.

Elimination: The median elimination half-life is 2.13 hours in healthy subjects (receiving 1 mg budesonide) and 4.56 hours in EoE patients (receiving 4 mg budesonide).

Clearance of budesonide is about 14.61 ± 9.86 L/hour/kg in healthy subjects and 6.54 ± 4.4 L/hour/kg in EoE patients. Budesonide is eliminated only in marginal (if any) amounts by the kidney. No budesonide was detected in urine, only budesonide metabolites.

Special Populations and Conditions

Hepatic Impairment

A relevant proportion of budesonide is metabolised in the liver by CYP3A4. The systemic exposure of budesonide is considerably increased in patients with severely impaired hepatic function. No studies have been conducted with JORVEZA in patients with impaired liver function. JORVEZA should not be used in patients with hepatic impairment.

10 STORAGE, STABILITY AND DISPOSAL


Store at room temperature (15°C to 25°C). Store in the original package in order to protect from light and moisture.

PART II: SCIENTIFIC INFORMATION

11 PHARMACEUTICAL INFORMATION

Drug Substance

Proper/Common name:	Budesonide
Chemical name:	16 α ,17-[(1RS)-butylidenebis(oxy)]-11 β ,21-dihydroxypregna-1,4-diene-3,20-dione
Molecular formula:	C ₂₅ H ₃₄ O ₆
Molecular mass:	430.5
Structural formula:	



Physicochemical properties:	Budesonide is a white or almost-white crystalline powder, with a pKa of 12.85 \pm 0.10. Budesonide is practically insoluble in water, freely soluble in methylene chloride, sparingly soluble in ethanol (96%).
-----------------------------	--

12 CLINICAL TRIALS

12.1 Trial Design and Study Demographics

The efficacy and safety of JORVEZA were evaluated for the induction of clinico-pathological remission of eosinophilic esophagitis (EoE) in adults in one pivotal clinical study (BUL-1/EEA), as described in **Table 3**.

Table 3 Summary of trial design and patient demographics

Trial Design	Dosage, Route of Administration and Duration	Study Subjects (n)	Mean Age (Range)	Sex
Phase III, pivotal, randomized, double-blind, placebo-controlled, multicenter study	Test: JORVEZA (budesonide 1 mg orodispersible tablets BID) Treatment duration: 6 weeks double-blind followed by 6 weeks open-label for non-remitters	Adult patients with active EoE (88)	37.0 (18 – 69)	Male (83%) Female (17%)

BID = twice daily; EoE = eosinophilic esophagitis

The patients had to have active symptomatic and histological EoE based on the presence of (a) symptoms on at least 1 day in the last 7 days prior to baseline with severity of ≥ 4 points on a 11-point Numerical Rating Scale (NRS) of *Dysphagia*, OR *Pain at swallowing*, and (b) Peak eosinophils count $\geq 65/\text{mm}^2$ high power field (hpf) in at least one hpf out of a total of 6 hpfs derived from 6 biopsies.

Patients with PPI-responsive esophageal eosinophilia were excluded. Patients with conditions for which corticosteroids therapy is not recommended were also excluded.

The primary efficacy endpoint was clinico-pathological remission defined as peak of < 16 eosinophils/ mm^2 high power field in esophageal biopsies, and resolution of symptoms was defined as severity of ≤ 2 points on 0 to 10-point Numerical Rating Scale [NRS] for dysphagia AND a severity of ≤ 2 points on 0-10 NRS for pain during swallowing [odynophagia] on each day in the week 6.

12.2 Study Results

Study BUL-1/EEA

A total of 59 patients were treated with JORVEZA 1 mg twice daily, and 29 patients were treated with placebo for 6 weeks during the double-blind placebo phase. Among the 59 patients, 23 continued JORVEZA in an open-label extension for another 6 weeks (total of 12 weeks).

JORVEZA (budesonide orodispersible tablets) was highly statistically significantly superior to placebo on the primary efficacy endpoint. In the final primary confirmatory analysis, 0/29 (0%) placebo patients and 34/59 (57.6%) JORVEZA patients were in clinico-pathological remission at DB week 6 (LOCF) (1-sided Fisher's exact test $p = 0.00000001$) (see **Table 4**). The results were confirmed in the PP analysis.

Table 4: Clinico-pathological remission rates at double-blind week 6 (LOCF) (double-blind phase, confirmatory) – FAS population

	Number (%) of patients with clinico-pathological remission at DB Week 6 (LOCF)		Difference between proportions ^a [95% RCI]	Testing of H ₀ ^b 1-sided p-value from Fisher's exact test
	JORVEZA 1 mg BID	Placebo		
Final FAS-DB	34/59 (57.6%)	0/29 (0.0%)	57.63% [38.22%; 71.97%]	0.00000001

BID: twice daily; DB: double-blind; FAS: full analysis set; LOCF: last observation carried forward; RCI: repeated confidence interval.

^a Difference between proportions ($\pi_{Eff} - \pi_{Pla}$)

^b Testing of H₀ ($\pi_{Pla} \geq \pi_{Eff}$) by means of the Fisher's exact test

In addition, key secondary efficacy endpoints (individual components of the primary endpoint) proved superiority of JORVEZA 1 mg BID vs placebo (see **Table 5**).

Table 5: Results of key secondary endpoints

Key secondary endpoints		Placebo (n = 29)	JORVEZA 1 mg BID (n = 59)	Difference between proportions [95% RCI]	p-value (one-sided)
1. Rate of patients with histological remission at week 6 (LOCF) ^a	n (%)	0/29 (0.0%)	55/59 (93.2%)	93.2% [86.8%; 99.6%]	< 0.0001 ^b
2. Rate of patients with resolution of symptoms on each day in the week prior to week 6 (LOCF)	n (%)	4/29 (13.8%)	35/59 (59.3%)	45.5% [27.8%; 63.3%]	< 0.0001 ^b

DB: double-blind; BID: twice daily; LOCF: last observation carried forward.

^a For this analysis not evaluable results were set to 'No'.

^b Fisher's exact test was used for testing.

During the open-label extension, 23 patients received JORVEZA 1 mg twice daily for an additional 6 weeks in patients without remission at the end of the double-blind phase. The results suggested that the rate of patients with clinico-pathological remission reached 84.7%. However, no definitive conclusion can be drawn regarding the benefit of JORVEZA beyond 6 weeks given the open-label design and the very low number of patients receiving 12 weeks of treatment.

13 NON-CLINICAL TOXICOLOGY

Acute Toxicity

The acute toxicity of budesonide has been studied in mice, rats, and dogs. Published data indicate that the acute toxicity of budesonide depends on the route of administration. Mortality was seen at substantially lower doses after i.v., i.p. and s.c. administration compared to oral administration. The LD₅₀ values ranged between 53.6 and 173 mg/kg for s.c. administration, between 98.9 and 320 mg/kg for i.v. administration and between 138 and 300 mg/kg for the i.p.

route. In contrast, after oral administration, LD₅₀ values between >3200 mg/kg and >10,000 mg/kg were obtained. This likely reflects low systemic bioavailability due to extensive first-pass effect in the liver.

Toxicity After Repeated Administration

In repeated dose toxicity studies, budesonide induced expected systemic effects typical for glucocorticoids. Notable effects included atrophies of skin and adrenal cortex and immune system effects (decreased WBC, lymphocyte and platelet counts, lymphoid depletion, and atrophy of the thymus and spleen). The glucocorticosteroid effects were also partly present at low doses (comparable to or lower than expected exposures at the recommended human therapeutic dose), and a NOEL could not be established in most of the repeated dose toxicity studies.

Repeated dose toxicity of budesonide orodispersible formulation was assessed as part of local tolerance study in hamsters. In this study, budesonide was administered to hamsters at oral (into the cheek pouches) doses of 0, 0.125 and 0.250 mg/day for 4 weeks. Systemic toxicity (premature deaths, marked reduction of body weight, and marked changes in hematology and clinical biochemistry) was noted with both doses, which is probably not unexpected at these doses. The systemic exposure at the lowest tested dose was 10-fold (C_{max}) and 15-fold (AUC) higher than expected systemic exposures (at the recommended therapeutic dose). For local tolerance effects, see section **Local Tolerance of Budesonide Orodispersible Tablets**.

Carcinogenesis and Genotoxicity

Carcinogenicity studies with budesonide were conducted in rats and mice. In a 91-week study in mice, budesonide caused no treatment-related carcinogenicity at oral doses up to 200 µg/kg/day.

In a two-year study in Sprague-Dawley (SD) rats, budesonide caused a small but statistically significant increase in the incidence of gliomas in male rats at an oral dose of 50 µg/kg/day (approximately 0.24 times the maximum recommended human dose on a body surface area basis). In addition, there were increased incidences of primary hepatocellular tumors in male rats at doses ≥25 µg/kg/day (approximately 0.12 times the maximum recommended human dose on a body surface area basis). In female rats, there was a statistically significant increase in combined primary neoplasms of the mammary gland at a dose of 50 µg/kg/day. The results regarding gliomas were considered equivocal since the SD rat is very variable with regard to spontaneous glioma incidence.

To elucidate these results, two further 104-week carcinogenicity studies with budesonide 50 µg/kg/day were performed, one using male SD rats, and one using male Fischer rats (which have a lower and less variable incidence of gliomas). Prednisolone and triamcinolone acetonide were used as reference glucocorticoids in both studies. The results from these new carcinogenicity studies in male rats did not demonstrate an increased glioma incidence in budesonide treated animals, as compared to concurrent controls or reference glucocorticosteroid treated groups. However, consistent with the original study, a statistically significant increase in the incidence of hepatocellular tumors was noted in the repeat study in male SD rats. The concurrent reference corticosteroids (prednisolone and triamcinolone acetonide) showed similar findings thus indicating a class effect of glucocorticosteroids.

Budesonide had no mutagenic and chromosome damaging properties in *in vitro* and *in vivo* tests demonstrating no genotoxic potential.

Reproductive and Developmental Toxicity

Effects on fertility and general reproductive performance:

In a fertility and early embryonic development study in rats, budesonide had no effect on fertility but caused early embryonic death, reduced fetal viability and delayed ossification at a dose of 20 µg/kg/day (approximately 0.09 times the maximum recommended human dose on a body surface area basis). Maternal effects consisting of reduced body weight and food consumption were also noted at a dose of 20 µg/kg/day.

In a study that evaluated fertility and general reproductive performance, reduced peri- and post-natal viability was noted at a dose of 80 µg/kg/day (approximately 0.38 times the maximum recommended human dose on a body surface area basis).

Embryo-Fetal development:

Budesonide, like other glucocorticosteroids, caused fetal death and abnormalities of fetal development in rats and rabbits.

In embryo-fetal development studies in rats, budesonide (doses ranging from 4 to 500 µg/kg/day) administered subcutaneously to pregnant rats during the fetal organogenesis period resulted in inhibition of weight gain or weight loss and systemic corticosteroid effects in dams at doses \geq 20 µg/kg/day. Reduced fetal weights and delayed sternebral or vertebral ossification were noted at doses \geq 20 µg/kg/day (approximately 0.1 times the maximum recommended human dose on a body surface area basis) and reduced fetal viability was noted at doses \geq 100 µg/kg/day (approximately 0.5 times the maximum recommended human dose on a body surface area basis). Significant maternal toxicity (deteriorated general condition including piloerection, drowsiness, decreased food consumption and reduced body weight gain) and fetal abnormalities (fetal loss, reduced weight and skeletal abnormalities) occurred at doses \geq 500 µg/kg/day (approximately 2.4 times the maximum recommended human dose on a body area basis).

In pregnant rabbits, doses of 5, 25, and 125 µg/kg/day (approximately 0.05, 0.24, and 1.2 times the maximum recommended human dose on a body surface area basis) were administered subcutaneously during days 6-18 of gestation. In the low and medium dose groups, food consumption and body weight gain were decreased during the fourth gestational week. Some also showed signs of diarrhea and vaginal bleeding. In the high dose group, all rabbits aborted at the end of the gestation period. In the medium dose group, a marked increase in the frequency of abnormalities, mainly skeletal defects (mainly skull and vertebral abnormalities), was observed.

Local Tolerance of Budesonide Orodispersible Tablets

Potential irritating effects of budesonide orodispersible tablets on mucous membranes of the oral cavity and the esophagus were assessed by administration into the cheek pouch of female hamsters. 0.5 mg and 1 mg budesonide orodispersible tablets as well as placebo tablets were tested.

Budesonide orodispersible tablets showed a favorable local tolerance as demonstrated by the daily macroscopic inspections of the cheek pouches and by histopathological examination of the mucosa at necropsy. The irritation of the epithelium of the cheek pouches, tongue and esophagus was classified as minimal according to the irritation index. Epithelial atrophy was the main histopathological finding in cheek pouch, tongue and esophagus. The thinning of the epithelium was considered as a typical glucocorticoid effect and assumed to be at least partly caused by systemically available budesonide.

READ THIS FOR SAFE AND EFFECTIVE USE OF YOUR MEDICINE

PATIENT MEDICATION INFORMATION

Pr **JORVEZA™**

Budesonide Orodispersible Tablets

Read this carefully before you start taking JORVEZA and each time you get a refill. This leaflet is a summary and will not tell you everything about this drug. Talk to your healthcare professional about your medical condition and treatment and ask if there is any new information about JORVEZA.

What is JORVEZA used for?

- For the treatment of eosinophilic esophagitis. This is an inflammatory condition of the throat (esophagus) that causes problems swallowing solid food.

How does JORVEZA work?

JORVEZA contains budesonide, which is used to decrease inflammation. As the tablet dissolves in your mouth, it mixes with your saliva, which is slowly swallowed and coats your throat (esophagus).

What are the ingredients in JORVEZA?

Medicinal ingredient: Budesonide

Non-medicinal ingredients: Anhydrous monosodium citrate, Disodium hydrogen citrate, Docusate sodium, Macrogol 6000, Magnesium stearate, Mannitol, Povidone K25, Sodium hydrogen carbonate and Sucralose.

JORVEZA comes in the following dosage forms:

Blister. Pack sizes: 20, 30 or 90 orodispersible tablets.

Do not use JORVEZA if you:

- Are allergic to budesonide or any of the other ingredients of this medicine
- Have active tuberculosis
- Have uncontrolled infections

To help avoid side effects and ensure proper use, talk to your healthcare professional before you take JORVEZA. Talk about any health conditions or problems you may have, including if you:

- Have high blood pressure
- Have diabetes, or if someone in your family has diabetes
- Have weakening of the bones (osteoporosis)
- Have ulcers in the stomach or first part of the small intestine (peptic ulcer)
- Have increased pressure in your eye (glaucoma) or eye problems such as clouding of the lens (cataracts) or if somebody in your family has glaucoma
- Have liver or kidney disease

- Are pregnant or breast-feeding, think you may be pregnant or planning to have a baby
- Are about to have an operation
- Are on other steroid treatment.

Other warnings you should know about:

Your immune system may be weakened while you are taking JORVEZA. Take the following steps to help avoid getting an infection:

- Keep away from people who have chicken pox or herpes zoster (shingles), if you have never had them. These illnesses may get worse during treatment with JORVEZA. If you do come into contact with chickenpox or shingles, see your doctor right away.
- Tell your doctor if you have not yet had measles.
- If you need to be vaccinated, please speak to your doctor first.
- If you know that you have a planned operation, please tell your doctor that you are using JORVEZA.

Tell your doctor if you have white spots in your mouth and throat or any signs of an infection. The symptoms of some infections can be unusual or less pronounced.

Children and adolescents

JORVEZA should not be used in children and adolescents under 18 years of age. The use of this medicine in children younger than 18 years of age has not yet been studied.

Tell your healthcare professional about all the medicines you take, including any drugs, vitamins, minerals, natural supplements or alternative medicines.

The following may interact with JORVEZA:

- Ketoconazole or itraconazole (to treat fungal infection)
- Clarithromycin or erythromycin (antibiotic medicine used to treat infections)
- Ritonavir and cobicistat (to treat HIV infections)
- Oestrogens (used for hormone replacement therapy or birth control)
- Cardiac glycoside such as digoxin (medicines used to treat heart conditions)
- Diuretics (to remove excess fluid from the body)

You should not drink grapefruit juice while you are taking this medicine as this can worsen its side effects.

JORVEZA could affect the results of tests (ACTH stimulation test) performed by your doctor or in hospital. Tell your doctor that you are taking JORVEZA before any tests are carried out.

How to take JORVEZA:

Always use this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

- Take the tablet after a meal
- Do NOT take any liquid or food with the tablet
- Do not eat, drink, brush your teeth or rinse your mouth for at least 30 minutes after you have taken the tablet. Do not use any oral solutions, sprays or chewable tablets at least 30

minutes before or after administration of JORVEZA. This will ensure that your medicine works properly.

- Place the tablet on the tip of your tongue and close your mouth. Press the tablet gently against the roof of your mouth until it has dissolved completely (this usually takes about two minutes). Swallow the dissolved material with saliva little by little while the tablet disintegrates
- Do not chew or swallow the undissolved tablet

Usual adult dose:

The recommended dose is two tablets per day, one in the morning and one in the evening after a meal (total daily dose: 2 mg budesonide).

Your treatment should last about 6 weeks.

Speak to your doctor if you want to interrupt or end your treatment early. It is important that you do not stop taking your medicine without talking to your doctor. Keep taking your medicine until your doctor tells you to stop, even if you start to feel better.

Overdose:

If you think you have taken too much JORVEZA, contact your healthcare professional, hospital emergency department or regional poison control centre immediately, even if there are no symptoms.

Missed Dose:

If you miss a dose, continue your treatment at the prescribed dosage. Do not use a double dose to make up for a forgotten dose.

What are possible side effects from using JORVEZA?

These are not all the possible side effects you may feel when taking JORVEZA. If you experience any side effects that are not listed here, contact your healthcare professional.

Typical side effects of cortisone preparations such as JORVEZA may occur which may affect all parts of the body, particularly when you use JORVEZA at high doses and for prolonged periods.

The following side effects have been reported during the use of JORVEZA:

Common (may affect up to 1 in 10 people)

- Headache
- High blood pressure
- Pain in the upper part of the stomach
- Heartburn
- Swelling of the lips
- Feeling sick (nausea)
- Tingling or numbness in the mouth
- Tiredness

Not known (cannot be estimated based on available data)

- Dysgeusia (taste disorder)

The following side effects have been reported with medicines similar to JORVEZA, and can therefore also occur with JORVEZA:

- Restlessness with increased physical activity, anxiety, aggression
- Increased risk of blood clotting, inflammation of the blood vessels (associated with stopping cortisone use after long-term therapy)
- Indigestion, irritable stomach (dyspepsia), constipation, ulcers in the stomach or small intestine
- Rash from hypersensitivity reactions, red spots from bleeding in the skin, delayed wound healing, local skin reactions such as contact dermatitis, bruising
- General feeling of being ill

Contact your doctor if you experience blurred vision or other visual disturbances.

Serious side effects and what to do about them			
Symptom / effect	Talk to your healthcare professional		Stop taking drug and get immediate medical help
	Only if severe	In all cases	
VERY COMMON			
Fungal infections in the throat/esophagus: white spots in mouth or throat		✓	
REPORTED WITH MEDICINES SIMILAR TO JORVEZA			
Changes in vision that may be signs of glaucoma (increased pressure in your eye), cataract (clouding of the lens of your eye), central serous chorioretinopathy (CSCR)		✓	
Brittle bones (osteoporosis), bone loss due to poor circulation of blood (osteonecrosis)	✓		
Cushing's syndrome: roundness of the face, weight gain, reduced glucose tolerance, high blood sugar, fluid retention in the tissues (e.g. swollen legs), increased excretion of potassium (hypokalemia), irregular periods in women, unwanted body hair in women, impotence, abnormal laboratory findings (reduced adrenal function), red stripes on the skin (stretch marks), acne	✓		

Serious side effects and what to do about them			
Symptom / effect	Talk to your healthcare professional		Stop taking drug and get immediate medical help
	Only if severe	In all cases	
Hyperglycemia (high levels of sugar in the blood): increased thirst, headache, blurred vision, fatigue		✓	
Increased risk of infection (such as chicken pox and shingles)	✓		
Pancreatitis (inflammation of the pancreas): severe pain in the abdomen and back	✓		
Mood changes, such as depression, irritability or euphoria	✓		
Muscle and joint pain, muscle weakness, muscle twitching	✓		

If you have a troublesome symptom or side effect that is not listed here or becomes bad enough to interfere with your daily activities, talk to your healthcare professional.

Reporting Side Effects

You can report any suspected side effects associated with the use of health products to Health Canada by:

- Visiting the Web page on Adverse Reaction Reporting (<https://www.canada.ca/en/health-canada/services/drugs-health-products/medeffect-canada/adverse-reaction-reporting.html>) for information on how to report online, by mail or by fax; or
- Calling toll-free at 1-866-234-2345.

NOTE: Contact your health professional if you need information about how to manage your side effects. The Canada Vigilance Program does not provide medical advice.

Storage:

Store at room temperature (15°C to 25°C). Store in the original package in order to protect from light and moisture.

Keep out of reach and sight of children.

If you want more information about JORVEZA:

- Talk to your healthcare professional
- Find the full product monograph that is prepared for healthcare professionals and includes this Patient Medication Information by visiting the Health Canada website (<https://health-products.canada.ca/dpd-bdpp/index-eng.jsp>); the manufacturer's website www.avirpharma.com, or by calling 1-800-363-7988.

This leaflet was prepared by:

AVIR Pharma Inc.

660 Boul. Industriel

Blainville, Québec

J7C 3V4

www.avirpharma.com

Last Revised November 5, 2019

JORVEZA™ is a trademark of Dr. Falk Pharma GmbH, used under license by AVIR Pharma Inc.