

PRODUCT MONOGRAPH
INCLUDING PATIENT MEDICATION INFORMATION

PrTALTZ[®]

ixekizumab

Solution for Injection
80 mg / 1.0 mL

Immunomodulator

Eli Lilly Canada Inc.,
Exchange Tower
130 King Street West, Suite 900,
PO Box 73
Toronto, Ontario
M5X 1B1
1-888-545-5972
www.lilly.ca

Date of Initial Approval:
May 25, 2016

Date of Revision:
February 4, 2020

Submission Control No: 225178

TALTZ is a registered trademark owned or licensed to Eli Lilly and Company, its subsidiaries or affiliates.

Table of Contents

PART I: HEALTH PROFESSIONAL INFORMATION..... 3
SUMMARY PRODUCT INFORMATION 3
DESCRIPTION..... 3
INDICATIONS AND CLINICAL USE..... 3
CONTRAINDICATIONS 4
WARNINGS AND PRECAUTIONS..... 4
ADVERSE REACTIONS..... 6
DRUG INTERACTIONS 13
DOSAGE AND ADMINISTRATION 14
OVERDOSAGE 16
ACTION AND CLINICAL PHARMACOLOGY 16
STORAGE AND STABILITY 17
SPECIAL HANDLING INSTRUCTIONS 18
DOSAGE FORMS, COMPOSITION AND PACKAGING 18

PART II: SCIENTIFIC INFORMATION 19
PHARMACEUTICAL INFORMATION..... 19
CLINICAL TRIALS 19
TOXICOLOGY 37
REFERENCES 38

PART III: PATIENT MEDICATION INFORMATION 40

INSTRUCTIONS FOR USE – PREFILLED AUTOINJECTOR..... 47

INSTRUCTIONS FOR USE – PREFILLED SYRINGE..... 55

PrTALTZ®

ixekizumab

Solution for Injection
80 mg / 1.0 mL

Immunomodulator

PART I: HEALTH PROFESSIONAL INFORMATION

SUMMARY PRODUCT INFORMATION

Route of Administration	Dosage Form / Strength	Clinically Relevant Nonmedicinal Ingredients
Subcutaneous Injection (S.C.)	Sterile solution for injection / 80 mg/1 mL (prefilled auto injector or prefilled syringe)	None <i>For a complete listing see DOSAGE FORMS, COMPOSITION and PACKAGING section.</i>

DESCRIPTION

TALTZ® (ixekizumab) is a humanized immunoglobulin G subclass 4 (IgG4) monoclonal antibody (mAb) comprised of two identical light chain polypeptides of 219 amino acids each and two identical heavy chain polypeptides of 445 amino acids each, and has a molecular weight of 146,158 Daltons for the protein backbone of the molecule. Ixekizumab selectively binds and neutralizes the proinflammatory cytokine interleukin-17A (IL-17A). Ixekizumab works by targeting IL-17A and inhibiting its interaction with the IL-17 receptor, which is expressed on various cell types including keratinocytes.

TALTZ, for subcutaneous use, is available as: 80 mg of ixekizumab in a 1 mL single-dose prefilled autoinjector or a single-dose prefilled syringe, manufactured to deliver 80 mg of ixekizumab. TALTZ prefilled autoinjector and prefilled syringe do not contain latex. See DOSAGE FORMS, COMPOSITION AND PACKAGING.

TALTZ is for single use and, therefore, contains no antimicrobial preservative.

INDICATIONS AND CLINICAL USE

Plaque Psoriasis

TALTZ (ixekizumab) is indicated for the treatment of adult patients with moderate-to-severe

plaque psoriasis who are candidates for systemic therapy or phototherapy.

Psoriatic Arthritis

TALTZ is indicated for the treatment of adult patients with active psoriatic arthritis who have responded inadequately to, or are intolerant to one or more disease-modifying antirheumatic drugs (DMARD). TALTZ can be used alone or in combination with a conventional DMARD (cDMARD) (e.g., methotrexate).

Ankylosing Spondylitis

TALTZ is indicated for the treatment of adult patients with active ankylosing spondylitis who have responded inadequately to, or are intolerant to conventional therapy (see CLINICAL TRIALS, Ankylosing spondylitis).

CONTRAINDICATIONS

TALTZ is contraindicated in patients with known serious hypersensitivity to ixekizumab or to any of the excipients. For a complete listing of excipients, see the Dosage Forms, Composition and Packaging section of the product monograph. See WARNINGS AND PRECAUTIONS, Hypersensitivity.

WARNINGS AND PRECAUTIONS

General

Infections

TALTZ has a potential to increase the risk of infection. In clinical trials, infections have been observed in TALTZ-treated patients. Most of the infections were mild or moderate in severity and did not lead to early discontinuation. See ADVERSE REACTIONS.

TALTZ should be used with caution in patients with clinically important chronic or active infection.

Patients should be instructed to seek medical advice if signs or symptoms suggestive of an infection occur. If a patient develops a serious infection or is not responding to standard therapy, the patient should be closely monitored and TALTZ should not be administered until the infection resolves.

TALTZ should not be given to patients with active tuberculosis (TB). Prior to initiating treatment with TALTZ, patients should be evaluated for TB infection. Treatment of latent TB infection should be initiated prior to administering TALTZ. Anti-tuberculosis therapy should also be considered prior to initiation of TALTZ in patients with past history of latent or active TB in whom an adequate course of treatment cannot be confirmed. Patients receiving TALTZ should be monitored closely for signs and symptoms of active TB during and after treatment.

Hypersensitivity

Serious hypersensitivity reactions, including anaphylaxis, angioedema and urticaria, have been reported in TALTZ-treated patients. See ADVERSE REACTIONS. If a serious hypersensitivity reaction occurs, administration of TALTZ should be discontinued immediately and appropriate therapy initiated.

Gastrointestinal

Inflammatory Bowel Disease

Caution should be exercised when prescribing TALTZ to patients with inflammatory bowel disease, including Crohn's disease and ulcerative colitis, as new cases or exacerbations were reported in TALTZ-treated patients. Patients who are treated with TALTZ and have inflammatory bowel disease should be monitored closely.

Immune

Vaccination

Prior to initiating therapy with TALTZ, consider completion of all age appropriate immunizations according to current immunization guidelines. Patients treated with TALTZ should not receive live vaccines. See DRUG INTERACTIONS. No data are available on the response to live vaccines.

Special Populations

Pregnant Women:

No clinical studies have been conducted with TALTZ in pregnant women to establish the safety of TALTZ during pregnancy. Studies in cynomolgus monkeys showed that ixekizumab crosses the placental barrier. No effects on embryo-fetal development were observed in fetuses from pregnant monkeys administered ixekizumab by subcutaneous injection during organogenesis to near parturition up to 19 times the maximum human recommended dose (MRHD). Neonatal deaths occurred in the offspring of pregnant monkeys administered weekly subcutaneous injections of ixekizumab from the beginning of organogenesis until parturition at 1.9 times the MRHD. Animal studies are not always predictive of human response, therefore the clinical significance of these findings is not known. See Part II: TOXICOLOGY.

Nursing Women:

There is no information regarding the presence of TALTZ in human breast milk, the effects on the breastfed infant, or the effects on milk production. Because many drugs and immunoglobulins are excreted in human milk, caution should be exercised when TALTZ is administered to a nursing woman.

Fertility:

No data are available on the effect of TALTZ on human fertility. Animal studies did not show any effects on fertility endpoints. See Part II: TOXICOLOGY.

Pediatrics (<18 years of age):

Safety and effectiveness of TALTZ in pediatric patients (<18 years of age) have not been evaluated.

Geriatrics (≥65 years of age):

Of the 4204 plaque psoriasis patients exposed to TALTZ in clinical trials, a total of 301 were 65 years or older, and 36 patients were 75 years or older. Of the 454 psoriatic arthritis patients exposed to TALTZ during the 24-week placebo-controlled period of the 2 pivotal clinical trials, a total of 69 were 65 years or older, and 6 patients were 75 years or older. Of the 376 ankylosing spondylitis patients exposed to TALTZ during the 16-week placebo-controlled period of the 2 pivotal clinical trials, a total of 19 were 65 years or older, and 3 patients were 75 years or older. Although no differences in safety or efficacy were observed between older and younger patients, the number of patients aged 65 and over is not sufficient to determine whether they respond differently from younger patients. See ACTION AND CLINICAL PHARMACOLOGY, *Special Populations and Conditions*, Geriatrics.

ADVERSE REACTIONS**Adverse Drug Reaction Overview**

The most frequently reported adverse drug reactions were injection site reactions and upper respiratory tract infections (most frequently nasopharyngitis). Most of the reactions were mild or moderate in severity and did not lead to discontinuation of TALTZ.

In the placebo-controlled period of the phase 3 studies, the proportion of patients who discontinued treatment due to adverse events was 2% in TALTZ-treated patients and 1.1% in placebo-treated patients in plaque psoriasis studies, 4.2% in the TALTZ-treated patients and 3.6% in the placebo-treated patients in psoriatic arthritis studies, and 4.3% in the TALTZ-treated patients and 1.1% in the placebo-treated patients in ankylosing spondylitis studies.

Clinical Trial Adverse Drug Reactions

Because clinical trials are conducted under very specific conditions the adverse reaction rates observed in the clinical trials may not reflect the rates observed in practice and should not be compared to the rates in the clinical trials of another drug. Adverse drug reaction information from clinical trials is useful for identifying drug-related adverse events and for approximating rates.

Adverse Drug Reactions in Plaque Psoriasis Trials

A total of 4204 plaque psoriasis patients were treated with TALTZ in blinded and open-label clinical trials. Of these, 2190 patients were exposed for at least 1 year.

Three randomized, double-blind, placebo-controlled phase 3 trials (UNCOVER-1, UNCOVER-2, and UNCOVER-3) in plaque psoriasis patients were integrated to evaluate the safety of TALTZ in comparison to placebo up to 12 weeks. In two of the trials (UNCOVER-2 and UNCOVER-3), the safety of TALTZ included a comparison to an active comparator, etanercept, up to 12 weeks. In total, 3858 patients were evaluated (1167 to TALTZ 80 mg Q2W group, 1161 to TALTZ 80 mg Q4W group, 739 to etanercept 50 mg twice weekly group, and 791 to placebo group). See CLINICAL TRIALS.

Table 1 summarizes the adverse drug reactions that occurred at a frequency of ≥1% in patients treated with TALTZ during the placebo-controlled 12-week period of UNCOVER-1,

UNCOVER-2, and UNCOVER-3.

Table 1: Adverse Drug Reactions Reported by Greater Than or Equal to 1% of Patients with Plaque Psoriasis Through Week 12 in the 3 UNCOVER Trials

Adverse Drug Reactions	TALTZ		PLACEBO	Etanercept ^a
	80 mg Q2W (N = 1167) n (%)	80 mg Q4W (N = 1161) n (%)	(N = 791) n (%)	(N = 739) n (%)
General disorders and administration site conditions				
Injection site reactions	196 (16.8%)	150 (12.9%)	26 (3.3%)	121 (16.4%)
Infections and Infestations				
Upper respiratory tract infection ^b	163 (14.0%)	155 (13.4%)	101 (12.8%)	92 (12.4%)
Tinea infections	17 (1.5%)	10 (0.9%)	1 (0.1%)	1 (0.1%)
Gastrointestinal disorders				
Nausea	23 (2.0%)	15 (1.3%)	5 (0.6%)	3 (0.4%)
Respiratory, thoracic and mediastinal disorders				
Oropharyngeal pain	16 (1.4%)	20 (1.7%)	4 (0.5%)	7 (0.9%)

^a Etanercept data from UNCOVER-2 and UNCOVER-3 studies only

^b Upper respiratory tract infection includes: nasopharyngitis and upper respiratory tract infection.

Adverse Drug Reactions in Psoriatic Arthritis Trials

A total of 1118 psoriatic arthritis patients were treated with TALTZ in blinded and open-label clinical trials. Of these, 365 patients were exposed for at least 1 year.

Two randomized, double-blind, placebo-controlled phase 3 trials (SPIRIT-P1 and SPIRIT-P2) in psoriatic arthritis patients were integrated to evaluate the safety of TALTZ in comparison to placebo up to 24 weeks. A total of 678 patients were studied (225 to TALTZ 80 mg Q2W, 229 to TALTZ 80 mg Q4W and 224 to placebo). See CLINICAL TRIALS.

During the 24-week placebo-controlled period of these trials, the proportion of patients with adverse events was higher in the TALTZ Q4W group compared to the placebo group (67% and 57%, respectively). The safety profile observed in patients with psoriatic arthritis treated with TALTZ Q4W is consistent with the safety profile in patients with plaque psoriasis with the exception of the frequencies of influenza (1.3%) and conjunctivitis (1.3%).

Table 2 summarizes the adverse drug reactions that occurred at a frequency of $\geq 1\%$ in patients treated with TALTZ during the 24-week placebo-controlled period of the SPIRIT-P1 and SPIRIT-P2 trials.

Table 2: Adverse Drug Reactions Reported by Greater Than or Equal to 1% of Patients with Psoriatic Arthritis Through Week 24 in the 2 SPIRIT Trials

Adverse Drug Reactions	TALTZ		PLACEBO
	80 mg Q4W (N = 229) n (%)	80 mg Q2W (N = 225) n (%)	(N = 224) n (%)
General disorders and administration site conditions			
Injection site reactions ^a	40 (17.5%)	57 (25.3%)	10 (4.5%)
Infections and Infestations			
Upper respiratory tract infection ^b	33 (14.4%)	23 (10.2%)	25 (11.2%)
Conjunctivitis	3 (1.3%)	3 (1.3%)	0
Influenza	3 (1.3%)	1 (0.4%)	1 (0.4%)
Oral candidiasis	1 (0.4%)	4 (1.8%)	0
Rhinitis	1 (0.4%)	4 (1.8%)	0
Gastrointestinal disorders			
Nausea	1 (0.4%)	5 (2.2%)	3 (1.3%)
Respiratory, thoracic and mediastinal disorders			
Oropharyngeal pain	7 (3.1%)	2 (0.9%)	1 (0.4%)

^a Injection site reactions includes: injection site reaction, injection site erythema, injection site hypersensitivity, injection site pruritis, injection site swelling, injection site pain, injection site rash, injection site haematoma, injection site bruising, injection site induration, injection site urticaria, injection site discolouration, injection site inflammation, injection site papule, and injection site haemorrhage.

^b Upper respiratory tract infection includes: nasopharyngitis, upper respiratory tract infection, and viral respiratory tract infection.

Adverse Drug Reactions in Ankylosing Spondylitis Trials

A total of 639 ankylosing spondylitis patients were treated with TALTZ in blinded and open-label clinical trials. Of these, 341 patients were exposed for at least 1 year.

Two randomized, double-blind, placebo-controlled phase 3 trials (COAST-V and COAST-W) in ankylosing spondylitis patients were integrated to evaluate the safety of TALTZ in comparison to placebo up to 16 weeks. A total of 566 patients were studied (181 on TALTZ 80 mg Q2W, 195 on TALTZ 80 mg Q4W and 190 on placebo). See CLINICAL TRIALS.

During the 16-week placebo-controlled period of these trials, the proportion of patients with adverse events was higher in the TALTZ Q4W group compared to the placebo group (56.4% and 44.7%, respectively). The safety profile observed in patients with ankylosing spondylitis treated with TALTZ Q4W is consistent with the safety profile in patients with plaque psoriasis with the exception of the frequency of inflammatory bowel disease (1.5%) and rhinitis (1.0%). Additional treatment-emergent adverse events reported by greater than or equal to 1% of patients treated with TALTZ Q4W and occurring more frequently than in the placebo group included diarrhea

(3.1%), headache (1.5%), iridocyclitis (1.5%), eczema (1.0%), and pruritis (1.0%). At baseline, 22.3% of patients in the COAST-V and COAST-W studies had a history of iridocyclitis.

Table 3 summarizes the adverse drug reactions that occurred at a frequency of $\geq 1\%$ in patients treated with TALTZ during the 16-week placebo-controlled period of the COAST-V and COAST-W trials.

Table 3: Adverse Drug Reactions Reported by Greater Than or Equal to 1% of Patients with Ankylosing Spondylitis Through Week 16 in the 2 COAST Trials

Adverse Drug Reactions	TALTZ		PLACEBO
	80 mg Q4W (N = 195) n (%)	80 mg Q2W (N = 181) n (%)	(N = 190) n (%)
General disorders and administration site conditions			
Injection site reactions ^a	12 (6.2%)	27 (14.9%)	10 (5.3%)
Infections and Infestations			
Upper respiratory tract infection ^b	32 (16.4%)	25 (13.8%)	15 (7.9%)
Tinea infection	2 (1.0%)	1 (0.6%)	0
Rhinitis	2 (1.0%)	1 (0.6%)	0
Gastrointestinal disorders			
Inflammatory bowel disease ^c	3 (1.5%)	1 (0.6%)	1 (0.5%)
Respiratory, thoracic and mediastinal disorders			
Oropharyngeal pain	5 (2.6%)	0	0

^a Injection site reactions includes: injection site reaction, injection site pain, injection site erythema, injection site rash, injection site pruritus, injection site dermatitis, injection site hypersensitivity, injection site mass, injection site oedema, injection site bruising, injection site paraesthesia.

^b Upper respiratory tract infection includes: nasopharyngitis, pharyngitis, and upper respiratory tract infection.

^c Inflammatory bowel disease includes adjudicated cases of Crohn's disease and ulcerative colitis.

Injection Site Reactions

The most frequent injection site reactions observed in psoriasis patients were injection site erythema and injection site pain.

In psoriatic arthritis patients, the most frequent injection site reactions observed were injection site erythema, injection site hypersensitivity and injection site pruritus.

In ankylosing spondylitis patients, the most frequent injection site reactions observed were injection site pain and injection site erythema.

Injection site reactions across all indications were predominantly mild-to-moderate in severity and rarely led to discontinuation of TALTZ.

Infections

In the placebo-controlled period of the phase 3 clinical trials in plaque psoriasis (a total of 2328 patients treated with TALTZ and 791 patients treated with placebo up to 12 weeks), infections were reported in 27.2% of patients treated with TALTZ (Q2W and Q4W) compared with 22.9% of patients treated with placebo. Most of these were mild or moderate. Serious infections occurred in 0.4% and 0.7% of patients treated with TALTZ Q2W and Q4W, respectively, and in 0.4% of patients treated with placebo. See WARNINGS AND PRECAUTIONS.

During the maintenance treatment period, the exposure-adjusted incidence rate of infections was 0.71 per patient year (56.0%) in patients treated with TALTZ with the recommended dosing regimen (Q4W), compared with 0.78 per patient year (35.6%) in patients treated with placebo. The exposure-adjusted incidence rate of serious infections was 0.02 per patient year (1.4%) in patients treated with TALTZ (Q4W) and 0.02 per patient year (0.7%) in patients treated with placebo.

Over the entire treatment period (a total of 4204 plaque psoriasis patients treated with TALTZ for up to 60 weeks for the majority of patients), the exposure-adjusted incidence rate of infections was 0.47 per patient year (52.8%) in patients treated with TALTZ. The exposure-adjusted incidence rate of serious infections was 0.02 per patient year (1.6%) in patients treated with TALTZ.

In the placebo-controlled period of the psoriatic arthritis clinical trials, infections were reported in 32.8% of patients treated with TALTZ (Q2W and Q4W) compared with 27.7% of patients treated with placebo. Most of these were mild or moderate. Patients treated with TALTZ reported serious infections (1.3%) and opportunistic infections (3.3%) more commonly than patients treated with placebo (0% serious and 0.4% opportunistic infections). In patients treated with TALTZ, 9.3% had multiple or recurrent infections compared to 4.9% in placebo-treated patients. No increase in risk of infections has been observed with increasing duration of exposure to TALTZ. See WARNINGS AND PRECAUTIONS.

In the placebo-controlled period of the ankylosing spondylitis clinical trials, infections were reported in 23.9% of patients treated with TALTZ (Q2W and Q4W) compared with 12.1% of patients treated with placebo. Most of these were mild or moderate. Patients treated with TALTZ reported serious infections (1.1%) and opportunistic infections (0.5%) more commonly than patients treated with placebo (0% serious and 0% opportunistic infections). No increase in risk of infections has been observed with increasing duration of exposure to TALTZ. See WARNINGS AND PRECAUTIONS.

Laboratory Assessment of Cytopenia

Neutropenia was observed in clinical trials. In general, neutropenia was transient and did not require discontinuation of TALTZ and was not associated with an increased frequency of infections.

In the placebo-controlled and active-controlled period of the clinical trials in patients with plaque psoriasis (UNCOVER-2 and UNCOVER-3), neutropenia \geq Grade 3 ($<1,000$ cells/mm³) was observed in 0.3% of patients receiving TALTZ Q2W, compared to 0.5% of patients treated with etanercept and 0.3% of patients treated with placebo. The remaining cases of neutropenia were

low grade, either Grade 2 (2.6% for TALTZ Q2W vs 3.3% for etanercept; 0.3% for placebo; $\geq 1,000$ to $< 1,500$ cells/mm³) or Grade 1 (7.0% for TALTZ Q2W vs 9.9% for etanercept; 3.4% for placebo; $\geq 1,500$ cells/mm³ up to normal).

In the placebo-controlled and active-controlled period of the plaque psoriasis clinical trials (UNCOVER-2 and UNCOVER-3), thrombocytopenia (Grade 1) was reported in 3.0% of patients treated with TALTZ Q2W compared to 4.5% of patients treated with etanercept and 1.7% of patients treated with placebo.

The cytopenia profile in psoriatic arthritis and ankylosing spondylitis clinical studies was similar to that observed in the psoriasis studies. Compared to placebo-treated patients, more patients treated with ixekizumab had a decrease in neutrophil counts. Most of the TALTZ-treated patients with decreased neutrophil counts had treatment-emergent decreases to Grade 1 or 2. In ankylosing spondylitis clinical studies, neutropenia was not associated with an increased frequency of infections.

Active Comparator Studies

In the 2 psoriasis clinical studies that included an active comparator (UNCOVER-2 and UNCOVER-3), the frequency of serious adverse events was 1.9% for both etanercept and for TALTZ, and the frequency of discontinuation due to adverse events was 1.2% for etanercept and 2.0% for TALTZ. The frequency of infections was 21.5% for etanercept and 26.0% for TALTZ, with the majority of the events mild to moderate in severity. The frequency of serious infections was 0.4% for etanercept and 0.5% for TALTZ.

Immunogenicity

As with all therapeutic proteins there is a potential for immunogenicity with TALTZ.

Plaque psoriasis

By week 12, approximately 9% of patients treated with TALTZ at the recommended dosing regimen developed anti-drug antibodies.

Approximately 22% of patients treated with TALTZ at the recommended dosing regimen developed antibodies to ixekizumab during the 60-week treatment period. The clinical effects of antibodies to ixekizumab are dependent on the antibody titer; higher antibody titers were associated with decreasing drug concentration and clinical response.

Of the patients who developed anti-drug antibodies to ixekizumab during the 60-week treatment period, approximately 9%, which equates to 2% of subjects treated with TALTZ at the recommended dosing regimen, had antibodies that were classified as neutralizing. Neutralizing antibodies were associated with low drug concentrations and reduced clinical response.

Psoriatic arthritis

In patients treated with TALTZ 80 mg every 4 weeks for up to 52 weeks (SPIRIT-P1), 11% (n=12) developed anti-drug antibodies. Of the 12 patients who developed anti-drug antibodies, 8 had confirmed neutralizing antibodies. An association between immunogenicity and efficacy has not been established.

Ankylosing spondylitis

In patients treated with TALTZ 80 mg every 4 weeks for up to 16 weeks, 5.2% (n=10) developed anti-drug antibodies. Of the 10 patients who developed anti-drug antibodies, 3 had confirmed neutralizing antibodies. An association between immunogenicity and efficacy has not been established.

Across all indications an association between immunogenicity and treatment emergent adverse events has not been established. The percentages of patients that had allergic reactions/hypersensitivity adverse events were similar between patients with and without antibodies to ixekizumab. An association between injection site reactions and the presence of antibodies to ixekizumab has not been established.

The detection of antibody formation is highly dependent on the sensitivity and specificity of the assay. Additionally, the observed incidence of antibody (including neutralizing antibody) positivity in an assay may be influenced by several factors including assay methodology, sample handling, timing of sample collection, concomitant medications, and underlying disease. For these reasons, comparison of incidence of antibodies to TALTZ across indications or with the incidences of antibodies to other products may be misleading.

Allergic reactions/hypersensitivity

Plaque psoriasis

In the placebo-controlled period of the UNCOVER-1, UNCOVER-2 and UNCOVER-3 clinical trials through Week 12, 3.7% of patients treated with TALTZ reported one or more allergic reaction or hypersensitivity events compared with 2.1% of patients treated with placebo. Serious reactions and events identified as potentially anaphylactic occurred in 0.2% and 0.3% of patients treated with TALTZ, respectively and in 0.1% and 0.3% of patients treated with placebo, respectively.

Psoriatic arthritis

In the placebo-controlled period of the SPIRIT-P1 and SPIRIT-P2 clinical trials through Week 24, patients treated with TALTZ had a higher incidence of non-anaphylactic allergic or hypersensitivity reactions (5.3%) compared to those treated with placebo (1.8%). Serious reactions were rare but have occurred after TALTZ administration.

Ankylosing spondylitis

In the placebo-controlled period of the COAST-V and COAST-W clinical trials through Week 16, patients treated with TALTZ had a higher incidence of non-anaphylactic allergic or hypersensitivity reactions (4.0%) compared to those treated with placebo (1.1%). Serious non-anaphylactic reactions occurred in 1 (0.3%) patient treated with TALTZ, 0 patients treated with placebo, respectively.

Less Common Clinical Trial Adverse Drug Reactions (<1%)

Plaque psoriasis

Adverse reactions that occurred at a frequency less than 1% in the placebo-controlled period of UNCOVER-1, UNCOVER-2, and UNCOVER-3 through Week 12 included: influenza, rhinitis, conjunctivitis, urticaria, oral candidiasis, and inflammatory bowel disease.

Psoriatic arthritis

Adverse reactions that occurred at a frequency less than 1% in the placebo-controlled period of SPIRIT-P1 and SPIRIT-P2 through Week 24 included: tinea infection and urticaria.

Ankylosing spondylitis

Adverse reactions that occurred at a frequency less than 1% in the placebo-controlled period of COAST-V and COAST-W through Week 16 included: urticaria.

Post-Market Adverse Drug Reactions

The following undesirable effect (adverse drug reaction) is based on postmarketing spontaneous reports. Because these reactions are reported voluntarily from a population of uncertain size, it is not possible to reliably estimate their frequency.

Immune system disorders

Anaphylaxis

DRUG INTERACTIONS

Drug-Drug Interactions

The safety of TALTZ in combination with other immunomodulatory agents or phototherapy has not been evaluated.

The pharmacokinetics of ixekizumab were not significantly different when ixekizumab was administered alone compared with concomitant administration of methotrexate in patients with psoriatic arthritis.

The clearance of ixekizumab was not significantly different when ixekizumab was administered alone compared with concomitant administration of oral corticosteroids, non-steroidal anti-inflammatory drugs (NSAIDs), or cDMARDs (sulfasalazine and methotrexate) in patients with ankylosing spondylitis.

Live vaccinations should not be given concurrently with TALTZ. See WARNINGS AND PRECAUTIONS.

Healthy individuals who received ixekizumab had similar antibody responses 1 month after vaccination with tetanus and pneumococcal vaccines compared to individuals who did not receive ixekizumab. The safety and clinical effectiveness of vaccines have not been assessed in patients undergoing treatment with ixekizumab.

Results from a drug-drug interaction study in patients with moderate-to-severe psoriasis determined that administration of TALTZ with drugs metabolized by CYP3A4 (i.e., midazolam), CYP2C9 (i.e., warfarin), CYP2C19 (i.e., omeprazole), CYP1A2 (i.e., caffeine) or CYP2D6 (i.e., dextromethorphan) does not have a clinically significant impact on the pharmacokinetics of these drugs.

Drug-Food Interactions

Interactions with food have not been established.

Drug-Herb Interactions

Interactions with herbal products have not been established.

Drug-Laboratory Interactions

Interactions with laboratory tests have not been established.

DOSAGE AND ADMINISTRATION

Dosing Considerations

TALTZ is intended for use under the guidance and supervision of a health care professional. Patients may self-inject after training in subcutaneous injection technique using the prefilled autoinjector or prefilled syringe.

Recommended Dose and Dosage Adjustment

Plaque psoriasis

The recommended dose is 160 mg by subcutaneous injection (two 80 mg injections) at Week 0, followed by 80 mg (one injection) at Weeks 2, 4, 6, 8, 10, and 12, then 80 mg (one injection) every 4 weeks.

Psoriatic arthritis

The recommended dose is 160 mg by subcutaneous injection (two 80 mg injections) at Week 0, followed by 80 mg every 4 weeks. For psoriatic arthritis patients with coexistent moderate-to-severe plaque psoriasis, use the dosing regimen for plaque psoriasis (see DOSAGE AND ADMINISTRATION, Plaque psoriasis). For psoriatic arthritis patients with coexistent mild plaque psoriasis, use the dosing regimen for psoriatic arthritis: 160 mg at Week 0, followed by 80 mg every 4 weeks (see Part II, CLINICAL TRIALS, Psoriatic arthritis).

Ankylosing spondylitis

The recommended dose is 80 mg by subcutaneous injection every 4 weeks. Limited data suggests that some TNF inhibitor experienced patients with ankylosing spondylitis may benefit from a 160 mg starting dose. Conventional disease-modifying antirheumatic drugs (cDMARD) (e.g., sulfasalazine), corticosteroids, non-steroidal anti-inflammatory drugs (NSAIDs), and/or analgesics may be continued during treatment with ixekizumab.

Renal impairment/Hepatic impairment

TALTZ has not been studied specifically in these patient populations.

Pediatric patients

Safety and effectiveness of TALTZ in pediatric patients (<18 years of age) have not been evaluated.

Geriatric patients

Of the 4204 plaque psoriasis patients exposed to TALTZ in clinical trials, a total of 301 were 65 years or older, and 36 patients were 75 years or older. Of the 454 patients with psoriatic arthritis exposed to TALTZ during the 24-week placebo-controlled period of the 2 pivotal clinical trials, a total of 69 were 65 years or older, and 6 patients were 75 years or older. Of the 376 patients with ankylosing spondylitis exposed to TALTZ during the 16-week placebo-controlled period of the 2 pivotal clinical trials, a total of 19 were 65 years or older, and 3 patients were 75 years or older. Although no differences in safety or efficacy were observed between older and younger patients, the number of patients aged 65 and over is not sufficient to determine whether they respond differently from younger patients.

Missed Dose

Patients who miss a dose of TALTZ should be advised to inject this missed dose as soon as they become aware of it, and then follow with their next scheduled dose.

Administration

There are two presentations for TALTZ (prefilled autoinjector and prefilled syringe). See the TALTZ Instructions for Use for each presentation for more detailed instructions on the preparation and administration of TALTZ.

Preparation for Use of TALTZ Prefilled Autoinjector and Prefilled Syringe

Before injection, remove TALTZ prefilled autoinjector or TALTZ prefilled syringe from the refrigerator and allow TALTZ to reach room temperature (30 minutes) without removing the needle cap.

Inspect TALTZ visually for particulate matter and discoloration prior to administration. The TALTZ solution is clear and colourless to slightly yellow. Do not use if the liquid contains visible particles, is discoloured or cloudy. TALTZ does not contain preservatives therefore discard any unused product remaining in the prefilled autoinjector or prefilled syringe after injection.

TALTZ is for subcutaneous administration. Administer each injection at a different anatomic location (such as upper arms, thighs or any quadrant of abdomen) than the previous injection, and not into areas where the skin is tender, bruised, erythematous, indurated or affected by psoriasis. Administration of TALTZ in the upper, outer arm may be performed by a caregiver or healthcare provider.

Instruct patients using the prefilled autoinjector or prefilled syringe to inject the full amount (1 mL), which provides 80 mg of TALTZ, according to the directions provided in the Instructions for Use.

OVERDOSAGE

Doses up to 180 mg have been administered subcutaneously in clinical trials without dose-limiting toxicity. Overdoses up to 240 mg, subcutaneously, have been reported without any serious adverse events. In the event of overdose, it is recommended that the patient be monitored for any signs or symptoms of adverse reactions and appropriate symptomatic treatment be instituted immediately.

For management of a suspected drug overdose, contact your regional Poison Control Centre.

ACTION AND CLINICAL PHARMACOLOGY

Mechanism of Action

Ixekizumab is an IgG4 monoclonal antibody that has a binding affinity of <3 pM to interleukin 17A (IL-17A), a naturally occurring proinflammatory cytokine. Elevated levels of IL-17A have been implicated in the pathogenesis of a variety of autoimmune diseases. Ixekizumab inhibits the release of proinflammatory cytokines and chemokines.

Pharmacodynamics

No formal pharmacodynamic studies have been conducted with TALTZ.

Pharmacokinetics

The pharmacokinetic properties of ixekizumab were similar in the plaque psoriasis, psoriatic arthritis and ankylosing spondylitis indications. Unless otherwise indicated, the data referenced in this section refer to analyses conducted in plaque psoriasis patients.

Absorption:

Following a single subcutaneous dose of 160 mg ixekizumab in patients with plaque psoriasis, the mean peak concentration (C_{max}) of 16.2 ± 6.57 $\mu\text{g/mL}$ was achieved within approximately 5 days.

After the 160 mg starting dose and the 80 mg Q2W dosing regimen, steady state was achieved by Week 5 (range 2 – 10 weeks). The mean (\pm SD) steady-state maximum concentration ($C_{max,ss}$) and trough concentration ($C_{trough,ss}$) estimates were 15.6 ± 6.23 $\mu\text{g/mL}$, and 9.74 ± 4.81 $\mu\text{g/mL}$, respectively.

After switching from the 80 mg Q2W dosing regimen to the 80 mg Q4W dosing regimen at Week 12, steady state was achieved by approximately 12 weeks. The mean (\pm SD) $C_{max,ss}$ and $C_{trough,ss}$ estimates were 10.5 ± 4.16 $\mu\text{g/mL}$, and 3.63 ± 2.33 $\mu\text{g/mL}$, respectively. The average subcutaneous bioavailability of ixekizumab was estimated in the range of 60% to 81% by population pharmacokinetic analysis in subjects with plaque psoriasis. Administration of ixekizumab via injection in the thigh achieved a higher bioavailability relative to that achieved using other injection sites including the arm and abdomen.

Distribution:

From population pharmacokinetic analyses, the mean total volume of distribution at steady-state was 7.11 L in patients with plaque psoriasis.

Metabolism:

The metabolic pathway of ixekizumab has not been characterized. Ixekizumab is a monoclonal antibody and is expected to be degraded into small peptides and amino acids via catabolic pathways in the same manner as endogenous IgGs.

Excretion:

In a population pharmacokinetic analysis, mean serum clearance was 0.0161 L/hr (0.39L/day). The estimated mean elimination half-life is 13 days in plaque psoriasis patients. Ixekizumab clearance and volume of distribution increase as body weight increases.

Dose Linearity:

Ixekizumab exhibited dose-proportional pharmacokinetics in plaque psoriasis patients over a dose range of 5 to 160 mg following subcutaneous administration.

Special Populations and Conditions**Pediatrics (<18 years of age):**

Safety and effectiveness of TALTZ in pediatric patients (<18 years of age) have not yet been evaluated.

Geriatrics (≥65 years of age):

Of the 4204 plaque psoriasis patients exposed to TALTZ in clinical trials, a total of 301 were 65 years or older, and 36 patients were 75 years or older. Based on population pharmacokinetic analysis, the clearance of ixekizumab in elderly patients was similar to patients less than 65 years of age.

Hepatic or Renal Insufficiency:

Specific clinical pharmacology studies to evaluate the effects of hepatic or renal impairment on the pharmacokinetic of ixekizumab have not been conducted.

STORAGE AND STABILITY

TALTZ is sterile and preservative-free. Discard any unused portion after injection.

TALTZ must be protected from light until use. Store refrigerated at 2°C to 8°C (36°F to 46°F). TALTZ may be stored unrefrigerated for up to 5 days at a temperature not above 30°C (86°F). Do not freeze. Do not use TALTZ if it has been frozen. Do not shake.

Keep in safe place out of the reach and sight of children.

SPECIAL HANDLING INSTRUCTIONS

Discard the TALTZ single-dose prefilled autoinjector or syringe after use in a puncture-resistant container. Patients or caregivers should be instructed in the technique as well as proper syringe and needle disposal, and not to reuse these items.

DOSAGE FORMS, COMPOSITION AND PACKAGING

TALTZ is supplied as:

- 80 mg single-dose prefilled autoinjector (pack sizes: carton of 1, 2 or 3)*
- 80 mg single-dose prefilled syringe (pack sizes: carton of 1, 2 or 3)*

*Not all pack sizes and presentations may be marketed.

The TALTZ prefilled autoinjector and prefilled syringe each contain a 1 mL glass syringe with a fixed 27 gauge ½ inch needle and are manufactured to deliver 80 mg ixekizumab.

Each prefilled autoinjector or prefilled syringe is composed of ixekizumab (80 mg/mL) and the inactive ingredients, citric acid anhydrous, polysorbate 80, sodium chloride; sodium citrate dihydrate, and water for injection.

TALTZ is for single use and, therefore, contains no antimicrobial preservative.

TALTZ prefilled autoinjector and prefilled syringe do not contain latex.

PART II: SCIENTIFIC INFORMATION

PHARMACEUTICAL INFORMATION

Drug Substance

Proper name:	ixekizumab
Chemical name:	ixekizumab is a humanized IgG4 monoclonal antibody, anti-(human interleukin 17A)
Molecular mass:	146,158 Daltons
Physicochemical properties:	ixekizumab solution is clear to opalescent, colourless to slightly yellow to slightly brown with a pH of 5.4 - 6.0

Product Characteristics

TALTZ is available in a single-dose prefilled autoinjector or a single-dose prefilled syringe to deliver 80 mg ixekizumab.

TALTZ is supplied as:

- 80 mg single-dose prefilled autoinjector (pack sizes: carton of 1, 2 or 3)*
- 80 mg single-dose prefilled syringe (pack sizes: carton of 1, 2 or 3)*

*Not all pack sizes and presentations may be marketed.

The prefilled autoinjector and prefilled syringe each contain a 1 mL glass syringe and are manufactured to deliver 80 mg of ixekizumab.

CLINICAL TRIALS

Plaque psoriasis

The safety and efficacy of TALTZ were assessed in three multicentre, randomized, double-blind, placebo-controlled studies (UNCOVER-1, UNCOVER-2, and UNCOVER-3) in a total of 3866 patients 18 years of age and older with plaque psoriasis who had a minimum body surface area involvement of 10%, a static Physician Global Assessment (sPGA) score of ≥ 3 and Psoriasis Area and Severity Index (PASI) score ≥ 12 , and who were candidates for phototherapy or systemic therapy. Patients with guttate, erythrodermic, or pustular psoriasis were excluded from the studies.

Each pivotal study evaluated short-term efficacy (up to 12 weeks, the “Induction Dosing Period”) of TALTZ versus placebo; 2 of the 3 studies (UNCOVER-2 and UNCOVER-3) included etanercept as an active comparator treatment. Studies UNCOVER-1 and UNCOVER-2 evaluated maintenance of efficacy for an additional 48 weeks after induction treatment (up to Week 60, the “Maintenance Dosing Period”) using a randomized withdrawal design for TALTZ-

treated patients who met response criteria (defined as static Physician Global Assessment [sPGA] (0,1) at Week 12). In the induction dosing period, subjects were randomized to either placebo or TALTZ (80 mg every two weeks [Q2W]) for 12 weeks, following a 160 mg starting dose, or etanercept 50 mg twice weekly for 12 weeks in studies UNCOVER-2 and UNCOVER-3.

In all three studies, the co-primary endpoints were the proportion of patients who achieved at least a 75% reduction in PASI score (PASI 75) from baseline to Week 12 and the proportion of patients with an sPGA (0,1) (clear or minimal) with at least a 2-point improvement from baseline. PASI is a composite score that takes into consideration both the percentage of body surface area affected and the nature and severity of psoriatic plaques (induration, erythema and scaling) within the affected regions. The sPGA is a 6 category scale ranging from 0 (clear) to 5 (very severe) that indicates the physician's overall assessment of psoriasis based on plaque thickness/induration, erythema, and scaling.

Other evaluated outcomes included the proportion of patients with an sPGA score of 0 (clear), a reduction of at least 90% in PASI (PASI 90), and a reduction of 100% in PASI (PASI 100).

Patient demographics, baseline characteristics, or disease severity were consistent across all 3 studies. The majority of patients were enrolled in North America (51.3%) and Europe (42.8%). Patients' baseline illness was moderate to severe as indicated by a mean baseline sPGA score of 3.6, a mean baseline PASI score of 20.2, and a mean baseline percentage of body surface area involvement (%BSA) of 27.3. Baseline sPGA score was severe or very severe in 51% of subjects in UNCOVER-1, 50% in UNCOVER-2, and 48% in UNCOVER-3.

Approximately 43.5% of patients had received phototherapy prior to enrollment; 49.3% had received prior conventional systemic therapy; 26.4% had received prior biologic therapy for the treatment of psoriasis. Of the patients who had received prior biologic therapy, 14.9% had received at least one anti-TNF- α agent, and 8.7% had received an anti-IL-12/IL-23 and 10.3% had an inadequate response to biologic therapy. A total of 23.4% of study patients had a history of psoriatic arthritis.

Table 4: Summary of study design and patient demographics for UNCOVER-1, UNCOVER-2, and UNCOVER-3

Study #	Trial design	Dosage, route of administration and duration	Study subjects (n = number)	Mean age (Range)	Gender
UNCOVER-1 (RHAZ)	multicentre, randomized, double-blind, placebo-controlled	<u>Induction (Week 0 to Week 12):</u> 160 mg starting dose, then 80 mg Q2W (n = 433); 160 mg starting dose, then 80 mg Q4W (n = 432); Placebo (n = 431)	N = 1296	45.7 (18 - 88)	M = 883 (68.1%) F = 413 (31.9%)
		<u>Maintenance (Week 12 to Week 60):</u> 80 mg Q4W (n = 229); 80 mg Q12W (n = 227); Placebo (n = 226)	682	44.8 (18-88)	M = 460 (67.4%) F = 222 (32.6%)
UNCOVER-2 (RHBA)	multicentre, randomized, double-blind, placebo-controlled, active-comparator	<u>Induction (Week 0 to Week 12):</u> 160 mg starting dose, then 80 mg Q2W (n = 351); 160 mg starting dose, then 80 mg Q4W (n = 347); Placebo (n = 168); Etanercept (n = 358)	1224	45.0 (18 - 84)	M = 821 (67.1%) F = 403 (32.9%)
		<u>Maintenance (Week 12 to Week 60):</u> 80 mg Q4W (n = 187); 80 mg Q12W (n = 181); Placebo (n = 176)	544	44.0 (18 - 84)	M = 361 (66.4%) F = 183 (33.6%)
UNCOVER-3 (RHBC)	multicentre, randomized, double-blind, placebo-controlled, active-comparator	<u>Induction (Week 0 to Week 12):</u> 160 mg starting dose, then 80 mg Q2W (n = 385); 160 mg starting dose, then 80 mg Q4W; (n = 386) Placebo (n = 193); Etanercept (n = 382)	1346	45.8 (17 - 88)	M = 918 (68.2%) F = 428 (31.8%)

Abbreviations: Q2W = every 2 weeks; Q4W = every 4 weeks; Q12W = every 12 weeks

Study Results

Clinical Response at 12 Weeks

The results for UNCOVER-1, UNCOVER-2 and UNCOVER-3 are presented in Tables 5, 6, and 7, respectively. In addition, see Figure 1 for PASI 75 response rates.

Table 5: Efficacy Results at Week 12 (NRI)^a in UNCOVER-1

	Placebo N = 431	TALTZ	
		80 mg Q2W N = 433	80 mg Q4W N = 432
sPGA of “0” (clear) or “1” (minimal)^c, n (%)	14 (3.2%)	354 (81.8%)	330 (76.4%)
Difference from Placebo (97.5% CI)	--	78.5% (73.9%, 83.1%) ^b	73.1% (68.2, 78.1) ^b
PASI 75^c, n (%)	17 (3.9%)	386 (89.1%)	357 (82.6%)
Difference from Placebo (97.5% CI)	--	85.2% (81.2%, 89.2%) ^b	78.7% (74.1%, 83.3%) ^b
sPGA of “0” (clear), n (%)	0	160 (37.0%)	149 (34.5%)
Difference from Placebo (97.5% CI)	--	37.0% (31.7%, 42.2%) ^b	34.5% (29.4%, 39.6%) ^b
PASI 90, n (%)	2 (0.5%)	307 (70.9%)	279 (64.6%)
Difference from Placebo (97.5% CI)	--	70.4% (65.5%, 75.4%) ^b	64.1% (58.9%, 69.3%) ^b
PASI 100, n (%)	0	153 (35.3%)	145 (33.6%)
Difference from Placebo (97.5% CI)	--	35.3% (30.2%, 40.5%) ^b	33.6 (28.5%, 38.7%) ^b

Abbreviations: N = number of patients in the intent-to-treat population; NRI = Non-Responder Imputation

^a patients with missing data were counted as non-responders

^b p-value <0.001 compared with placebo

p-value is based on a logistic regression analysis with treatment, geographic region, previous non-biologic systemic therapy, and baseline weight category included as factors (where placebo response rate is zero, p-value is based on Fisher's exact test)

p-value is adjusted for multiplicity of testing based on pre-defined hierarchy.

Confidence interval corresponds to an alpha level of 0.025

^c co-primary objectives

Table 6: Efficacy Results at Week 12 (NRI)^a in UNCOVER-2

	Placebo (N = 168)	TALTZ		Etanercept (N = 358)
		80 mg Q2W (N = 351)	80 mg Q4W (N = 347)	
sPGA of “0” clear) or “1” (minimal)^d, n (%)	4 (2.4%)	292 (83.2%)	253 (72.9%)	129 (36.0%)
Difference from Placebo (97.5% CI)	--	80.8% (75.6%, 86.0%) ^b	70.5% (64.6%, 76.5%) ^b	--
Difference from Etanercept (97.5% CI)	--	47.2 (39.9%, 54.4%) ^{b,c}	36.9% (29.1%, 44.7%) ^{b,c}	--
PASI 75^d, n (%)	4 (2.4%)	315 (89.7%)	269 (77.5%)	149 (41.6%)
Difference from Placebo (97.5% CI)	--	87.4% (82.9%, 91.9%) ^b	75.1 (69.5%, 80.8%) ^b	--
Difference from Etanercept (97.5% CI)	--	48.1% (41.2%, 55.0%) ^{b,c}	35.9% (28.2%, 43.6%) ^{b,c}	--
sPGA of “0” clear), n (%)	1 (0.6%)	147 (41.9%)	112 (32.3%)	21 (5.9%)
Difference from Placebo (95% CI)	--	41.3% (35.2%, 47.3%) ^b	31.7% (25.9%, 37.5%) ^b	--
Difference from Etanercept (97.5% CI)	--	36.0% (29.5%, 42.5%) ^b	26.4% (20.1%, 32.7%) ^b	--
PASI 90, n (%)	1 (0.6%)	248 (70.7%)	207 (59.7%)	67 (18.7%)
Difference from Placebo (97.5% CI)	--	70.1% (64.4%, 75.7%) ^b	59.1% (53.0%, 65.1%) ^b	--
Difference from Etanercept (97.5% CI)	--	51.9% (44.8%, 59.1%) ^b	40.9% (33.4%, 48.4%) ^b	--
PASI 100, n (%)	1 (0.6%)	142 (40.5%)	107 (30.8%)	19 (5.3%)
Difference from Placebo (97.5% CI)	--	39.9% (33.8%, 45.9%) ^b	30.2% (24.5%, 36.0%) ^b	--
Difference from Etanercept (97.5% CI)	--	35.1% (28.7%, 41.6%) ^b	25.5% (19.4%, 31.7%) ^b	--

Abbreviations: N = number of patients in the intent-to-treat population; NRI = Non-Responder Imputation

^a patients with missing data were counted as non-responders

^b p-value <0.001

p-value is based on Cochran-Mantel-Haenszel test stratified by pooled center and adjusted for multiplicity of testing based on pre-defined hierarchy.

Confidence interval corresponds to an alpha level of 0.025

^c Non-inferiority was demonstrated using a fixed margin approach before proceeding with superiority testing of ixekizumab compared to etanercept.

^d co-primary objectives

Table 7: Efficacy Results at Week 12 (NRI)^a in UNCOVER-3

	Placebo (N = 193)	TALTZ		Etanercept (N = 382)
		80 mg Q2W (N = 385)	80 mg Q4W (N = 386)	
sPGA of “0” (clear) or “1” (minimal)^d, n (%)	13 (6.7%)	310 (80.5%)	291 (75.4%)	159 (41.6%)
Difference from Placebo (97.5% CI)	--	73.8% (67.7%, 79.9%) ^b	68.7% (62.3%, 75.0%) ^b	--
Difference from Etanercept (97.5% CI)	--	38.9% (31.7%, 46.1%) ^{b,c}	33.8% (26.3%, 41.3%) ^{b,c}	--
PASI 75^d, n (%)	14 (7.3%)	336 (87.3%)	325 (84.2%)	204 (53.4%)
Difference from Placebo (97.5% CI)	--	80.0% (74.4%, 85.7%) ^b	76.9% (71.0%, 82.8%) ^b	--
Difference from Etanercept (97.5% CI)	--	33.9% (27.0%, 40.7%) ^{b,c}	30.8% (23.7%, 37.9%) ^{b,c}	--
sPGA of “0” (clear), n (%)	0	155 (40.3%)	139 (36.0%)	33 (8.6%)
Difference from Placebo (97.5% CI)	--	40.3% (34.7%, 45.9%) ^b	36.0% (30.5%, 41.5%) ^b	--
Difference from Etanercept (97.5% CI)	--	31.6% (25.2%, 38.1%) ^b	27.4% (21.0%, 33.7%) ^b	--
PASI 90, n (%)	6 (3.1%)	262 (68.1%)	252 (65.3%)	98 (25.7%)
Difference from Placebo (97.5% CI)	--	64.9% (58.9%, 71.0%) ^b	62.2% (56.1%, 68.3%) ^b	--
Difference from Etanercept (97.5% CI)	--	42.4% (35.1%, 49.7%) ^b	39.6% (32.2%, 47.0%) ^b	--
PASI 100, n (%)	0	145 (37.7%)	135 (35.0%)	28 (7.3%)
Difference from Placebo (97.5% CI)	--	37.7% (32.1%, 43.2%) ^b	35.0% (29.5%, 40.4%) ^b	--
Difference from Etanercept (97.5% CI)	--	30.3% (24.0%, 36.6%) ^b	27.6% (21.4%, 33.9%) ^b	--

Abbreviations: N = number of patients in the intent-to-treat population; NRI = Non-Responder Imputation

^a patients with missing data were counted as non-responders

^b p-value <0.001


p-value is based on Cochran-Mantel-Haenszel test stratified by pooled center and adjusted for multiplicity of testing based on pre-defined hierarchy.

Confidence interval corresponds to an alpha level of 0.025

^c Non-inferiority was demonstrated using a fixed margin approach before proceeding with superiority testing of ixekizumab compare to etanercept.

^d co-primary objectives

Figure 1: PASI 75 response rates at each post-baseline visit (NRI), Induction Dosing Period (UNCOVER-2 and UNCOVER-3)



Abbreviations: PBO = Placebo; ETN = Etanercept; IXE80 Q2W = Ixekizumab 80 mg Q2W; IXE80 Q4W = Ixekizumab 80 mg Q4W; N = number of patients in the analysis population; NRI = non-responder imputation; PASI = psoriasis area and severity index.

Examination of results stratified by previous treatment with a biologic agent did not identify differences in response to TALTZ in biologic-naïve and biologic-experienced patients.

Maintenance of Response at 60 Weeks

To evaluate the maintenance and durability of response, patients originally randomized to TALTZ and who were responders at Week 12 (i.e., sPGA score of 0,1) in UNCOVER-1 and UNCOVER-2 were re-randomized to an additional 48 weeks of one of the following treatment regimens: TALTZ 80 mg every 4 weeks (Q4W), TALTZ 80 mg every 12 weeks (Q12W), or placebo. Non-responders (sPGA >1) at Week 12 and patients who relapsed (sPGA ≥3) during the maintenance period were placed on TALTZ 80 mg Q4W.

The response rates for those patients re-randomized to the recommended maintenance dose of TALTZ 80 mg Q4W based on the recommended induction dose of TALTZ 80 mg Q2W are provided in Table 8.

Table 8: Maintenance of Response at Week 60 (NRI)^a in UNCOVER-1

Endpoint at Week 60	TALTZ 80 mg Q2W (induction)/ placebo (maintenance) (N = 110)	TALTZ 80 mg Q2W (induction)/80 mg Q4W (maintenance) (N=119)
Maintained sPGA of “0” (clear) or “1” (minimal) ^c ; n (%)	9 (7.7%)	89 (74.8%)
Difference from Placebo (98.75% CI)	--	67.1% (55.4%, 78.8%) ^b

Abbreviations: N = number of patients in the analysis population; NRI = non-responder imputation

^a patients with missing data were counted as non-responders

^b p-value <0.001

p-value is based on a logistic regression analysis with treatment and baseline weight category as factors

Confidence interval corresponds to an alpha level of 0.0125

^c p-value is adjusted for multiplicity of testing based on pre-defined hierarchy

Among the patients who achieved sPGA 0 or 1 at Week 12, the response rates were sPGA 0 = 55% and PASI 100 = 52% at Week 60.

Improvements at Week 12 from baseline compared to placebo (UNCOVER-1 and UNCOVER-3) were demonstrated in Dermatology Life Quality Index (DLQI); these improvements were maintained for 60 weeks (UNCOVER-1).

Clinical Response in Genital Psoriasis

A randomized, double-blind, placebo-controlled study (IXORA-Q) was conducted in 149 adult patients with plaque psoriasis who had a minimum BSA involvement of 1%, a sPGA score of ≥ 3 (at least moderate psoriasis), a sPGA of Genitalia score of ≥ 3 (at least moderate genital psoriasis), who failed to respond to or were intolerant of at least one topical therapy used for treatment of psoriasis affecting the genital area, and who were candidates for systemic therapy or phototherapy. Patients were randomized to placebo or TALTZ (initial dose of 160 mg followed by 80 mg every 2 weeks for 12 weeks). Randomization was stratified by BSA involvements of 1% to <10% and $\geq 10\%$. The primary endpoint was the proportion of patients who achieved a “0” (clear) or “1” (minimal) response on the sPGA of Genitalia at Week 12. Baseline BSA involvement was $\geq 10\%$ for approximately 60% of patients. Median baseline PASI score was 12.2 and approximately 42% of patients had a sPGA of Genitalia score at baseline indicating severe or very severe genital psoriasis. Approximately 24% of study participants were female.

Results for the primary endpoint in Study IXORA-Q are presented in Table 9.

Table 9: Efficacy Results at Week 12 (NRI)^a in Adults with Genital Psoriasis in IXORA-Q

Endpoint	Placebo	TALTZ 80 mg Q2W ^b
Number of patients randomized	N=74	N=75
sPGA of Genitalia “0” (clear) or “1” (minimal), n (%)	6 (8.1%)	55 (73.3%)
Difference from placebo (95% CI) p-value	--	65.2% (53.4%, 77.0%) ^c <0.001

Abbreviations: NRI = Non-Responder Imputation

^a Patients with missing data were counted as non-responders.

^b At Week 0, patients received 160 mg of TALTZ, followed by 80 mg every 2 weeks for 12 weeks.

^c p-value is based on the logistic regression analysis with treatment and baseline BSA category as factors. Confidence intervals are constructed using the simple asymptotic method, without continuity correction.

For patients with a baseline Genital Psoriasis Symptoms Scale (GPSS) Itch numeric rating scale (NRS) score of ≥ 4 (GPSS Itch NRS is an 11-point scale), the proportion of patients that reported a ≥ 4 -point reduction in GPSS Itch NRS score at Week 12, was 55.4% for patients receiving TALTZ and 5.9% for patients receiving placebo. Patient-perceived impact of genital psoriasis on limiting frequency of sexual activity (intercourse or other activities) was assessed by the Genital Psoriasis Sexual Frequency Questionnaire (GenPs-SFQ) Item 2 (In the past week how often did your genital psoriasis limit the frequency of your sexual activity?). Scores for the GenPs-SFQ Item 2 range from 0 to 4 (0=never, 1=rarely, 2=sometimes, 3=often, 4=always). For patients who reported a GenPs-SFQ Item 2 score ≥ 2 at baseline, the proportion of patients that reported a score of 0 or 1 at Week 12, was 78.4% for patients receiving TALTZ and 21.4% for patients receiving placebo.

Psoriatic arthritis

The safety and efficacy of TALTZ were assessed in 780 patients, in two randomized, double-blind, placebo-controlled studies (SPIRIT-P1 and SPIRIT-P2) in adult patients, age 18 years and older with active psoriatic arthritis (at least 3 swollen and at least 3 tender joints) despite non-steroidal anti-inflammatory drug (NSAID), corticosteroid or DMARD therapy. In SPIRIT-P1, patients also had at least 1 disease-related definite joint erosion on hand or foot x-rays as determined by the central reader or had a C-reactive protein (CRP) >6 mg/L at screening. In SPIRIT-P2, patients were required to have prior treatment with 1 or more cDMARDs and prior treatment with at least 1 but not more than 2 anti-TNF- α agents, and to have discontinued the anti-TNF- α agent due to either an inadequate response (based on a minimum of 12 weeks on therapy) or documented intolerance. In both studies, patients were required to have active psoriatic skin lesions (plaque) or a documented history of plaque psoriasis.

The SPIRIT-P1 study evaluated patients with psoriatic arthritis who were biologic-naïve, whereas the SPIRIT-P2 study evaluated patients with psoriatic arthritis who were anti-TNF- α agent-experienced. In both studies, patients were treated with either TALTZ 160 mg at Week 0 followed by 80 mg every 2 weeks (Q2W) or 4 weeks (Q4W), or placebo. SPIRIT-P1 also included adalimumab 40 mg Q2W as an active reference arm for sensitivity analysis. In both studies, patients who received placebo (or adalimumab in SPIRIT-P1) were re-randomized to TALTZ (80 mg Q2W or Q4W) at Week 16 if they were inadequate responders (defined as

patients with <20% improvement from baseline in both tender joint count and swollen joint count) or at Week 24 (see Table 10).

In SPIRIT-P1, 69.5% had psoriasis involving $\geq 3\%$ of body surface area (BSA), 58.0% had enthesitis, and 37.6% had dactylitis at baseline. The majority of patients (64.0%) were receiving a cDMARD at baseline and 14.6% of patients were cDMARD naïve. A total of 54.2% of patients were using methotrexate (MTX) at baseline. The majority of patients were Caucasian (94%) and were enrolled in Europe (73%) and the United States (20%).

In SPIRIT-P2, 62.5% of patients had psoriasis involving $\geq 3\%$ of BSA, 75.2% had enthesitis, and 23.7% had dactylitis at baseline. A total of 56.2% and 35.3% of patients were inadequate responders to 1 and 2 anti-TNF- α agents, respectively, and 8.5% were intolerant to an anti-TNF- α agent. More than half of the patients (51.0%) were receiving a cDMARD at baseline. A total of 41.0% of patients were using MTX at baseline. The majority of patients were Caucasian (92%) and were enrolled in the United States (52%) and Europe (41%).

The primary endpoint for both studies was the percentage of patients achieving at least a 20% improvement in the American College of Rheumatology (ACR20) criteria at Week 24 (see Table 12 for components of ACR20). Secondary endpoints evaluated included ACR50, ACR70; PASI 75; Health Assessment Questionnaire - Disability Index (HAQ-DI); Leeds Enthesitis Index (LEI) scores, and Minimal Disease Activity (MDA). Structural damage was also followed radiographically in the SPIRIT-P1 study by measuring the mean change from baseline to Week 24 in the modified total Sharp score (mTSS).

Table 10: Summary of study design and patient demographics for SPIRIT-P1 and SPIRIT-P2

Study #	Trial design	Dosage, route of administration and duration ^a	Study subjects (N = number)	Mean age (Range)	Gender
SPIRIT-P1 (RHAP)	multicentre, randomized, double-blind, placebo-controlled, active-reference	<u>TALTZ</u> ^b 160 mg SC starting dose at Week 0, followed by 80 mg SC Q2W (n = 103) or 80 mg SC Q4W (n = 107); Placebo ^c SC Q2W (n = 106); Adalimumab ^d 40 mg SC Q2W (n=101)	417	49.5 (19.0 – 76.0)	M = 192 (46.0%) F = 225 (54.0%)
SPIRIT-P2 (RHBE)	multicentre, randomized, double-blind, placebo-controlled	<u>TALTZ</u> ^b 160 mg SC starting dose at Week 0, followed by 80 mg SC Q2W (n = 123) or 80 mg SC Q4W (n = 122); Placebo ^c SC Q2W (n = 118)	363	51.9 (18.0 – 86.0)	M = 169 (46.6%) F = 194 (53.4%)

^a Patients in all treatment groups with an inadequate response at Week 16 received rescue therapy (modification to background therapy).

^b At Week 16, TALTZ inadequate responders received rescue therapy but remained on their assigned treatment.

^c At Week 16, placebo inadequate responders were re-randomized (1:1) to one of the TALTZ regimens. They received a 160 mg SC starting dose of TALTZ at Week 16 and their assigned TALTZ regimen thereafter. At Week 24, all patients still receiving placebo were re-randomized (1:1) to one of the TALTZ regimens. They received a 160 mg starting dose of TALTZ at Week 24 and their assigned TALTZ thereafter.

^d At Week 16, adalimumab inadequate responders were re-randomized (1:1) to one of the TALTZ regimens, but first underwent an 8-week washout period. They received a 160 mg SC starting dose of TALTZ at Week 24 and their assigned TALTZ regimen thereafter. At Week 24, all patients still receiving adalimumab were re-randomized (1:1) to one of the TALTZ regimens, but first underwent an 8-week washout period. They received an 80 mg dose of TALTZ at Week 32 and their assigned TALTZ regimen thereafter. The study was not powered to test equivalence or non-inferiority of TALTZ versus adalimumab.

Study Results

Clinical Response

In both studies, patients treated with TALTZ 80 mg Q2W or 80 mg Q4W demonstrated a greater clinical response compared to placebo at Week 12 and Week 24 (Table 11). Responses were similar in patients regardless of concomitant methotrexate or cDMARD treatment. In SPIRIT-P2, responses were seen regardless of prior inadequate response or intolerance to anti-TNF- α therapy.

There was no clear evidence of improved ACR response with the more frequent TALTZ 80 mg Q2W dose group compared to the 80 mg Q4W dose group. Responses were similar regardless of age, gender, race, and body weight. In patients with psoriatic arthritis and coexistent moderate-to-severe plaque psoriasis, TALTZ 80 mg Q2W demonstrated additional clinical benefit on skin clearance, whereas in patients with psoriatic arthritis and coexistent mild plaque psoriasis there was no added benefit to receiving TALTZ 80 mg Q2W compared to Q4W.

Table 11: Efficacy Results at Week 12 and 24 (NRI)^{a,b} in SPIRIT-P1 and SPIRIT-P2

	SPIRIT-P1 – biologic-naïve		SPIRIT-P2 – anti-TNF- α – experienced ^c	
	TALTZ 80 mg Q4W (N=107)	Placebo (N=106)	TALTZ 80 mg Q4W (N=122)	Placebo (N=118)
ACR20 response				
Week 12 (%)	57	31	50	22
Difference from placebo (95% CI)	26 (13, 39) ^d		28 (16, 40) ^d	
Week 24 (%)	58	30	53	20
Difference from placebo (95% CI)	28 (15, 41) ^d		34 (22, 45) ^d	
ACR50 response				
Week 12 (%)	34	5	31	3
Difference from placebo (95% CI)	29 (19, 39) ^e		28 (19, 37) ^e	
Week 24 (%)	40	15	35	5
Difference from placebo (95% CI)	25 (14, 37) ^e		30 (21, 40) ^e	
ACR70 response				
Week 12 (%)	15	0	15	2
Difference from placebo (95% CI)	15 (8, 22) ^e		13 (6, 20) ^e	
Week 24 (%)	23	6	22	0

Difference from placebo (95% CI)	18 (9, 27) ^c	22 (15, 30) ^e
----------------------------------	-------------------------	--------------------------

Abbreviations: ACR = American College of Rheumatology; CI = Confidence Interval; N = number of patients in the intent-to-treat population; NRI = Non-Responder Imputation.

^a Patients with missing data were counted as non-responders

^b Patients who met inadequate responder criteria (less than 20% improvement in tender and swollen joint counts) at Week 16 were considered non-responders at Week 24.


^c Included patients who had prior treatment with at least 1 but not more than 2 anti-TNF- α agents, and discontinued the anti-TNF- α agent due to either an inadequate response or documented intolerance

^d p value <0.001.

^e ACR50 and ACR70 are not adjusted for multiplicity testing.

The percentage of patients achieving ACR20 response by visit, up to Week 24, is shown in Figure 2. In both studies, clinical response occurred as early as Week 1.

Figure 2: Percent of Patients Achieving ACR20 Response^a in SPIRIT-P1 and SPIRIT-P2 Through Week 24



^a Patients who met inadequate responder criteria (less than 20% improvement in tender and swollen joint counts) at Week 16 were considered non-responders at Week 24.

The improvements in the components of the ACR response and Minimal Disease Activity (MDA) criteria are shown in Table 12.

Table 12: Efficacy Results in ACR Components and MDA Criteria at Week 12 and 24

	SPIRIT-P1 – biologic-naïve		SPIRIT-P2 – anti-TNF- α – experienced ^a	
	Placebo (N=106)	TALTZ 80 mg Q4W (N=107)	Placebo (N=118)	TALTZ 80 mg Q4W (N=122)
Number of Swollen Joints				
Mean Baseline	10.6	11.4	10.3	13.1
Mean Change at Week 12	-3.2	-6.2	-2.6	-5.8
Mean Change at Week 24	-3.5	-7.0	-5.0	-8.5

Number of Tender Joints				
Mean Baseline	19.2	20.5	23.0	22.0
Mean Change at Week 12	-3.5	-10.3	-5.4	-9.4
Mean Change at Week 24	-4.7	-11.9	-6.2	-12.7
Patient's Assessment of Pain^b				
Mean Baseline	58.5	60.1	63.9	63.9
Mean Change at Week 12	-9.1	-26.6	-11.9	-29.8
Mean Change at Week 24	-14.0	-29.6	-21.4	-36.9
Patient's Global Assessment of Disease Activity^b				
Mean Baseline	61.1	62.7	64.1	66.4
Mean Change at Week 12	-11.1	-29.7	-10.7	-34.5
Mean Change at Week 24	-14.8	-33.8	-19.0	-40.7
Physician's Global Assessment of Disease Activity^b				
Mean Baseline	55.9	57.6	58.9	60.3
Mean Change at Week 12	-16.6	-34.0	-15.9	-34.4
Mean Change at Week 24	-24.2	-38.5	-18.3	-40.0
Disability Index (HAQ-DI)^c				
Mean Baseline	1.2	1.2	1.2	1.2
Mean Change at Week 12	-0.1	-0.4	-0.1	-0.4
Mean Change at Week 24	-0.2	-0.4 ^c	-0.2	-0.6 ^c
CRP (mg/L)				
Mean Baseline	15.1	12.8	12.1	17.0
Mean Change at Week 12	-3.2	-8.8	-4.3	-11.4
Mean Change at Week 24	-3.9	-8.8	-3.6	-11.8
Minimal Disease Activity (MDA)^d				
Response at Week 12 (%)	5	21	5	25
Response at Week 24 (%)	15	30	3	28 ^c

Abbreviations: N = number of patients in the intent-to-treat population

^a Included patients who had prior treatment with at least 1 but not more than 2 anti-TNF- α agents, and discontinued the anti-TNF- α agent due to either an inadequate response or documented intolerance

^b Visual analog scale (mm); 0=best, 100=worst

^c Health Assessment Questionnaire - Disability Index (HAQ-DI): 0 = best, 3 = worst; measures the patient's ability to perform the following: dress/groom, arise, eat, walk, reach, grip, maintain hygiene, and maintain daily activity. HAQ-DI at Week 24 was adjusted for multiplicity in both studies.

^d Minimal Disease Activity (MDA): patients are classified as achieving MDA if they fulfill 5 of 7 outcome measures: tender joint count (TJC) ≤ 1 ; swollen joint count (SJC) ≤ 1 ; Psoriasis Activity and Severity Index (PASI total score) ≤ 1 or body surface area (BSA) ≤ 3 ; patient pain visual analog scale (VAS) score of ≤ 15 ; patient global disease activity VAS score of ≤ 20 ; HAQ-DI score ≤ 0.5 ; and tender enthesal points ≤ 1 . MDA is based on Non-Responder Imputation (patients with missing data were counted as non-responders). MDA at Week 24 was adjusted for multiplicity in SPIRIT-P2.

^e p value < 0.001

In patients with pre-existing enthesitis, numerical improvements in LEI mean change from baseline at Week 12 in SPIRIT-P1 and in resolution of enthesitis (LEI=0) at Week 24 in SPIRIT-P2 were seen in TALTZ-treated patients compared to those treated with placebo.

In patients with coexistent plaque psoriasis ($\geq 3\%$ BSA psoriasis skin involvement at baseline), the skin lesions of psoriasis improved with TALTZ treatment, relative to placebo, as measured by PASI 75 at Week 12 in the TALTZ 80 mg Q4W compared to the placebo groups. The proportion of patients who achieved PASI 75 responses were 75.3% (55/73) vs 7.5% (5/67) in SPIRIT-P1 and 57.4% (39/68) vs 10.4% (7/67) in SPIRIT-P2. In SPIRIT-P1, patients experienced improvement in itch severity, when compared to placebo at Week 12.

Radiographic Response

Radiographic changes were assessed in SPIRIT-P1. Inhibition of progression of structural damage was assessed radiographically and expressed as the change in mTSS and its components, the erosion score and joint space narrowing score at Week 24, compared to baseline. The mTSS was modified for psoriatic arthritis by addition of hand distal interphalangeal joints.

TALTZ 80 mg Q4W inhibited the progression of structural joint damage (mTSS) compared to placebo at Week 24. The adjusted mean change from baseline in the mTSS was 0.17 for TALTZ 80 mg Q4W and 0.49 for placebo (95% CI for mean differences for TALTZ minus placebo; -0.55, -0.10).

Patient Reported Outcomes

General health status was assessed by the Short Form Health Survey (SF-36). At Weeks 12 and 24 in SPIRIT-P1 and SPIRIT-P2, patients treated with TALTZ showed greater improvement from baseline in the SF-36 Physical Component Summary (PCS) score and all 8 domains compared to patients treated with placebo, and improved or did not worsen in the SF-36 Mental Component Summary score.

Maintenance of Response at 52 Weeks

To evaluate the maintenance of response, patients originally randomized to TALTZ who completed Week 24 of the SPIRIT-P1 trial continued in an extension phase on either TALTZ 80 mg Q2W or Q4W treatment. At Week 52, 80.6% of patients on Q4W dosing who had achieved ACR20 at Week 24 maintained this response. For patients on Q4W dosing for 52 weeks, 74.2% had a change from baseline of ≤ 0.5 in their mTSS score, indicating minimal progression of structural damage.

Ankylosing spondylitis

The efficacy and safety of TALTZ were assessed in 657 patients, in two randomized, double-blind, placebo-controlled studies (COAST-V and COAST-W) in adult patients, age 18 years and older with active ankylosing spondylitis. Patients had active disease as defined by the Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) ≥ 4 despite non-steroidal anti-inflammatory drug (NSAID), corticosteroid, or disease modifying anti-rheumatic drug (DMARD) therapy.

COAST-V evaluated 341 biologic-naive patients, who were treated with either TALTZ 80 mg or 160 mg at Week 0 followed by 80 mg every 2 weeks (Q2W) or 4 weeks (Q4W), adalimumab 40 mg every 2 weeks (active reference arm), or with placebo. Patients receiving placebo were re-randomized at Week 16 to receive TALTZ (160 mg initial dose, followed by 80 mg Q2W or Q4W). Patients receiving adalimumab were re-randomized at Week 16 to receive TALTZ at Week 20 (80 mg initial dose, followed by 80 mg Q2W or Q4W).

COAST-W evaluated 316 TNF inhibitor experienced patients (90% were inadequate responders and 10% were intolerant to TNF inhibitors). All patients were treated with TALTZ 80 mg or 160 mg at Week 0 followed by 80 mg Q2W or Q4W, or with placebo. Patients receiving placebo were re-randomized at Week 16 to receive TALTZ (160 mg initial dose, followed by 80 mg Q2W or Q4W).

At baseline, patients had symptoms of ankylosing spondylitis for an average of 17 years across both studies. At baseline, approximately 32% of the patients were on a concomitant cDMARD. In COAST-W, all patients discontinued previous treatment with 1 (63%) or 2 (37%) TNF inhibitors due to either inadequate response or intolerance. Across both studies the majority of patients were Caucasian (73%) and were enrolled in Europe (46%).

The primary efficacy endpoint in both studies was the percentage of patients achieving an Assessment of Spondyloarthritis International Society 40 (ASAS40) response at Week 16. Secondary endpoints evaluated included ASAS20, ASDAS, BASDAI Score, BASMI, hsCRP, MRI Spine SPARCC, BASDAI50, ASDAS <2.1, ASAS HI, and SF-36.

Table 13: Summary of study design and patient demographics for COAST-V and COAST-W

Study #	Trial design	Dosage, route of administration and duration ^a	Study subjects (N = number)	Mean age (Range)	Gender
COAST-V (RHBV)	Multicentre, randomized, double-blind, placebo- and active-controlled period (0 to 16 weeks) followed by extended treatment period (16-52 weeks)	<u>TALTZ</u> 80 mg or 160 mg SC starting dose at Week 0, followed by 80 mg SC Q2W (n = 83) or 80 mg SC Q4W (n = 81); Placebo SC Q2W (n = 87); Adalimumab ^a 40 mg SC Q2W (n = 90)	341	41.7 (19 – 78)	M = 276 (81.2%) F = 64 (18.8%)
COAST-W (RHBW)	Multicentre, randomized, double-blind, placebo-controlled period (0 to 16 weeks) followed by extended treatment period (16-52 weeks)	<u>TALTZ</u> 80 mg or 160 mg SC starting dose at Week 0, followed by 80 mg SC Q2W (n = 98) or 80 mg SC Q4W (n = 114); Placebo SC Q2W (n = 104)	316	46.1 (18 – 76)	M = 253 (80.1%) F = 63 (19.9%)

^a At Week 16, patients on adalimumab were re-randomized (1:1 ratio) to TALTZ 80 mg Q2W or 80 mg Q4W with an 80 mg starting dose. The last dose of adalimumab was administered at Week 14 followed by a 6-week washout period, and the first TALTZ 80 mg starting dose was given at Week 20.

Study Results

Clinical Response

In both studies, patients treated with TALTZ 80 mg Q2W or 80 mg Q4W demonstrated a greater clinical response compared to placebo at Week 16 (Table 14). Responses were similar in patients regardless of concomitant therapies (i.e., NSAIDs and cDMARDs).

There was no clear evidence of improved ASAS40 response with the more frequent TALTZ 80 mg Q2W dose group compared to the TALTZ 80 mg Q4W dose group. Responses were similar regardless of gender and body weight. In COAST-W responses were seen regardless of the number of prior TNF inhibitors.

Table 14: ASAS20 and ASAS40 Efficacy Results at Week 16 (NRI)^a in COAST-V and COAST-W

	COAST-V – biologic-naive		COAST-W – TNF inhibitor experienced ^b	
	TALTZ 80 mg Q4W ^c (N=81)	Placebo (N=87)	TALTZ 80 mg Q4W ^c (N=114)	Placebo (N=104)
ASAS40 response ^{d, e} , %	48.1	18.4	25.4	12.5
Difference from placebo (95% CI)	29.8 (16.2, 43.3) ^f	--	12.9 (2.7, 23.2) ^h	--
ASAS20 response ^d , %	64.2	40.2	48.2	29.8
Difference from placebo (95% CI)	24.0 (9.3, 38.6) ^g	--	18.4 (5.7, 31.1) ^g	--

Abbreviations: N = number of patients in the intent-to-treat population; NRI = Non-Responder Imputation.

^a Patients with missing data were counted as non-responders.

^b TNF inhibitor experienced includes patients who discontinued previous treatment with 1 or 2 TNF inhibitors due to either inadequate response or intolerance.

^c At Week 0, patients received 80 mg or 160 mg of TALTZ.

^d An ASAS20 response is defined as a $\geq 20\%$ improvement and an absolute improvement from baseline of ≥ 1 unit (range 0 to 10) in ≥ 3 of 4 domains (Patient Global Assessment, Spinal Pain, Function [BASFI], and Inflammation), and no worsening of $\geq 20\%$ and ≥ 1 unit (range 0 to 10) in the remaining domain. An ASAS40 response is defined as a $\geq 40\%$ improvement and an absolute improvement from baseline of ≥ 2 units in ≥ 3 of 4 domains without any worsening in the remaining domain.

^e Primary endpoint.


^f p value < 0.001 .

^g p value < 0.01 .

^h p value < 0.05 .

In COAST-V and COAST-W, statistically significant differences from placebo for ASAS20 were observed as early as Week 1 in the TALTZ 80 mg Q4W group, compared to the placebo group. For ASAS40, TALTZ 80 mg Q4W achieved a statistically significant difference from placebo by Week 1 in COAST-W and by Week 2 in COAST-V (see Figure 3).

Figure 3: Percent of Patients Achieving ASAS40 Response in COAST-V and COAST-W through Week 16, NRI^a



^a Patients with missing data were counted as non-responders.

The improvement in the main components of the ASAS40 response criteria and other measures of disease activity are shown in Table 15.

Table 15: ASAS Components and Other Measures of Disease Activity at Week 16

	COAST-V – biologic-naive		COAST-W – TNF inhibitor experienced ^a	
	TALTZ 80 mg Q4W ^b (N=81)	Placebo (N=87)	TALTZ 80 mg Q4W ^b (N=114)	Placebo (N=104)
ASAS Components				
Patient Global Assessment (0-10)				
Baseline	6.9	7.1	8.0	7.8
Mean Change from Baseline	-2.5	-1.4	-2.4	-0.7
Total Spinal Pain (0-10)				
Baseline	7.2	7.4	7.9	7.8
Mean Change from Baseline	-3.2	-1.7	-2.4	-1.0
BASFI (0-10)				
Baseline	6.06	6.35	7.35	7.01
Mean Change from Baseline	-2.39 ^g	-1.16	-1.69 ^g	-0.64
Inflammation (0-10) ^c				
Baseline	6.51	6.76	7.21	7.20
Mean Change from Baseline	-3.18	-1.27	-2.42	-0.70
ASDAS				
Baseline	3.71	3.89	4.15	4.05
Mean Change from Baseline	-1.43 ^g	-0.46	-1.16 ^g	-0.11
BASDAI Score				
Baseline	6.75	6.81	7.54	7.32
Mean Change from Baseline	-2.92	-1.39	-2.17 ^g	-0.92
BASMI				
Baseline	3.87	4.51	4.68	4.88
Mean Change from Baseline	-0.50	-0.08	-0.35	-0.05

hsCRP (mg/L)				
Baseline	12.19	15.97	20.16	16.02
Mean Change from Baseline	-5.21	1.43	-11.10	9.72
MRI Spine SPARCC ^d				
Baseline	14.53	15.80	8.30	6.37
Mean Change from Baseline	-11.02 ^g	-1.51	-2.99 ^h	3.29
BASDAI50 ^e (%), NRI ^f	42 ^g	17	22	10
ASDAS <2.1 (%) (Low Disease Activity), NRI ^f	43	13	18 ^h	5

Abbreviations: ASDAS = Ankylosing Spondylitis Disease Activity Score; BASDAI = Bath Ankylosing Spondylitis Disease Activity Index; BASFI = Bath Ankylosing Spondylitis Functional Index; BASMI = Bath Ankylosing Spondylitis Metrology Index; hsCRP = High sensitivity C-reactive protein; ITT = Intent-to-Treat; MRI Spine SPARCC = Spondyloarthritis Research Consortium of Canada Magnetic Resonance Imaging Scoring of the Spine (23 discovertebral unit scale); NRI = Non-Responder Imputation.

^a TNF inhibitor experienced includes patients who discontinued previous treatment with 1 or 2 TNF inhibitors due to either inadequate response or intolerance.

^b At Week 0, patients received 80 mg or 160 mg of TALTZ.

^c Inflammation is the mean of patient-reported stiffness self-assessments (questions 5 and 6) in BASDAI.

^d The numbers of ITT patients with MRI data at baseline are as follows: COAST-V: TALTZ, n = 81; PBO, n = 82. COAST-W: TALTZ, n = 58; PBO, n = 51.

^e BASDAI50 response defined as an improvement of $\geq 50\%$ of the BASDAI score from baseline.

^f Patients with missing data were counted as non-responders.

^g p value <0.001 (p values are only shown for endpoints with type I error controlled in pre-defined multiple testing scheme).

^h p value <0.01 (p values are only shown for endpoints with type I error controlled in pre-defined multiple testing scheme).

Patient Reported Outcomes

General health status and quality of life was assessed by the Short Form Health Survey (SF-36). At Week 16, in COAST-V and COAST-W, compared to placebo, patients treated with TALTZ 80 mg Q4W showed significantly greater improvement from baseline in the SF-36 Physical Component Summary (PCS) score (p<0.001 for both studies).

TOXICOLOGY

No significant adverse effects, including any organ toxicity or undesirable effects on immune function (e.g. T-cell dependent antibody response or NK cell activity) were observed in cynomolgus monkeys when administered ixekizumab by intravenous and subcutaneous injection up to dose levels of 50 mg/kg QW (19 times the MRHD on a mg/kg basis) for 8 and 39 weeks respectively.

Carcinogenicity, mutagenicity and genotoxicity studies have not been conducted with ixekizumab. There were no ixekizumab related effects on surrogate markers of fertility (e.g. estrous cyclicity, sperm parameters and reproductive organ weights) or histopathological findings in reproductive tissues when administered to sexually mature cynomolgus monkeys by subcutaneous injection up to doses of 50 mg/kg QW for 13 weeks. The monkeys were not mated to evaluate fertility.

No effects on embryo-fetal development were observed in fetuses from pregnant cynomolgus monkeys administered ixekizumab by subcutaneous injection during organogenesis to near parturition up to doses of 50 mg/kg QW. Neonatal deaths occurred in the offspring of pregnant monkeys administered ixekizumab by subcutaneous injection from the beginning of organogenesis until parturition. Infant losses (death or forced euthanasia) occurred in 0/14, 2/12 and 5/14 successful pregnancies in control, 5 mg/kg and 50 mg/kg QW treatment groups,

respectively and were attributable to maternal neglect (3), early delivery (2), congenital defect (1) and trauma (1). Surviving animals (14, 10 and 9 at 0, 5 and 50 mg/kg, respectively) did not display any treatment related functional or immunological developmental effects. Ixekizumab was shown to cross the placenta and was present in the blood of offspring for up to 6 months of age. Ixekizumab was detected at low levels in the breast milk of cynomolgus monkeys. The clinical significance of these findings is not known.

REFERENCES

1. Blauvelt, A., et al. (2017). Efficacy and safety of ixekizumab for the treatment of moderate-to-severe plaque psoriasis: results through 108 weeks of a randomized, controlled phase III clinical trial (UNCOVER-3). *J Am Acad Dermatol* published online Sept 13. <https://doi.org/10.1016/j.jaad.2017.06.153>.
2. Deodhar, A., et al. (2019). Efficacy and safety of ixekizumab in the treatment of radiographic axial spondyloarthritis: 16-week results from a phase 3 randomized, double-blind, placebo-controlled trial in patients with prior inadequate response or intolerance to tumor necrosis factor inhibitors. *Arthritis Rheumatol* 71(4): 599-611.
3. Gordon, K. B., et al. (2016). Phase 3 trials of ixekizumab in moderate-to-severe plaque psoriasis. *N Engl J Med* 375(4): 345-356.
4. Supplement to: Gordon, K. B., et al. (2016). Phase 3 trials of ixekizumab in moderate-to-severe plaque psoriasis. *N Engl J Med* 375(4): 345-356.
5. Griffiths, C. E., et al. (2015). Comparison of ixekizumab with etanercept or placebo in moderate-to-severe psoriasis (UNCOVER-2 and UNCOVER-3): results from two phase 3 randomised trials. *Lancet* 386(9993): 541-551.
6. Supplement to: Griffiths CEM, Reich K, Lebwohl M, et al, for the UNCOVER-2 and UNCOVER-3 investigators. Comparison of ixekizumab with etanercept or placebo in moderate-to-severe psoriasis (UNCOVER-2 and UNCOVER-3): results from two phase 3 randomised trials. *Lancet* 2015; published online June 10. [http://dx.doi.org/10.1016/S0140-6736\(15\)60125-8](http://dx.doi.org/10.1016/S0140-6736(15)60125-8).
7. Mease, P. J., et al. (2017). Ixekizumab, an interleukin-17A specific monoclonal antibody, for the treatment of biologic-naïve patients with active psoriatic arthritis: results from the 24-week randomized, double-blind, placebo-controlled and active (adalimumab)-controlled period of the phase III trial SPIRIT-P1. *Ann Rheum Dis* 76:79-87.
8. Nash, P., et al. (2017). Ixekizumab for the treatment of patients with active psoriatic arthritis and an inadequate response to tumour necrosis factor inhibitors: results from the 24-week randomised, double-blind, placebo-controlled period of the SPIRIT-P2 phase 3 trial. *Lancet* 389(10086): 2317-2327.
9. Ryan, C., et al. (2018). Efficacy and safety of ixekizumab in a randomized, double-blinded, placebo-controlled phase 3b study of patients with moderate-to-severe genital psoriasis. *Br J Dermatol* published online. <http://dx.doi.org/10.1111/bjd.16736>.
10. Strober, B., et al. (2017). Short and long-term safety outcomes with ixekizumab from 7 clinical trials in psoriasis: Etanercept comparisons and integrated data. *J Am Acad Dermatol* 76(3): 432-440.

11. Van der Heijde, D., et al. (2018). Efficacy and safety of ixekizumab in patients with active psoriatic arthritis: 52-week results from a phase III study (SPIRIT-P1). *J Rheumatol* 45(3): 367-377.
12. Van der Heijde, D., et al. (2018). Ixekizumab, an interleukin-17A antagonist in the treatment of ankylosing spondylitis or radiographic axial spondyloarthritis in patients previously untreated with biological disease-modifying anti-rheumatic drugs (COAST-V): 16 week results of a phase 3 randomised, double-blind, active-controlled and placebo-controlled trial. *Lancet* 392(10163): 2441-2451.

READ THIS FOR SAFE AND EFFECTIVE USE OF YOUR MEDICINE

PART III: PATIENT MEDICATION INFORMATION

PrTALTZ®
ixekizumab

www.lilly.ca



Read this carefully before you start taking TALTZ® and each time you get a refill. This leaflet is a summary and will not tell you everything about this drug. Talk to your healthcare professional about your medical condition and treatment and ask if there is any new information about TALTZ.

What is TALTZ used for?

Plaque Psoriasis

TALTZ is a prescription medicine used to treat adults with a skin condition called “plaque psoriasis”. Plaque psoriasis causes inflammation of the skin, appearing as raised, thick, red and scaly patches (“psoriatic lesions”) that can appear anywhere on your body. TALTZ reduces the inflammation and other symptoms of the disease.

Psoriatic Arthritis

TALTZ is used to treat adults with active psoriatic arthritis and can be used alone or with another medicine called a conventional disease modifying anti-rheumatic drug (cDMARD), for example methotrexate. You may first be given other medicines for this disease. If you do not respond well enough to these medicines, you will be given TALTZ. Psoriatic arthritis is an inflammatory disease of the joints, usually accompanied by psoriasis. TALTZ reduces signs and symptoms of your arthritis and improves your psoriasis.

Ankylosing Spondylitis

TALTZ is used to treat a condition called ankylosing spondylitis. The condition is an inflammatory disease primarily affecting the spine, which causes inflammation of the spinal joints. TALTZ is used in adults with active ankylosing spondylitis. You may first be given other medicines for this disease. If you do not respond well enough to these medicines, you will be given TALTZ.

How does TALTZ work?

TALTZ contains the active substance ixekizumab. Ixekizumab is a monoclonal antibody. Monoclonal antibodies are proteins that recognize and bind specifically to certain proteins in the body.

TALTZ belongs to a group of medicines called interleukin (IL) inhibitors. This medicine works by neutralizing the activity of a protein called IL-17A, which is present at increased levels in

diseases such as plaque psoriasis, psoriatic arthritis, and ankylosing spondylitis. By blocking IL-17A, TALTZ helps reduce the signs and symptoms of psoriasis such as itching, pain, and scaling; psoriatic arthritis such as painful and swollen joints; and ankylosing spondylitis such as back pain and morning stiffness.

Using TALTZ will benefit you by leading to fast and sustained improvements of skin clearance and reducing your symptoms such as scaling, itching and pain. If you have any questions about how TALTZ works or why this medicine has been prescribed for you, ask your healthcare professional.

What are the ingredients in TALTZ?

Medicinal ingredients: ixekizumab

Non-medicinal ingredients: citric acid anhydrous, polysorbate 80, sodium chloride, sodium citrate dihydrate, and water for injection

TALTZ comes in the following dosage forms:

- 80 mg solution for injection in prefilled autoinjector*
- 80 mg solution for injection in prefilled syringe*

*not all presentations may be marketed.

Do not use TALTZ if:

- you are allergic to ixekizumab or any of the other ingredients in TALTZ. See **What are the ingredients in TALTZ.**

If you think you may be allergic, ask your healthcare professional for advice before using TALTZ.

To help avoid side effects and ensure proper use, talk to your healthcare professional before you take TALTZ. Talk about any health conditions or problems you may have, including if you:

- currently have an infection or if you have long-term or repeated infections
- have tuberculosis or have been in close contact with someone with tuberculosis
- have inflammatory bowel disease (Crohn's disease or ulcerative colitis)
- have recently had a vaccination or if you are due to have a vaccination during treatment with TALTZ
- are receiving any other treatment for psoriasis, such as another immunosuppressant or phototherapy with ultraviolet (UV) light
- are pregnant, think that you may be pregnant or are planning to have a baby
- are breast-feeding or plan to breast-feed

Other warnings you should know about:

TALTZ can potentially cause serious side effects, including infections and allergic reactions. You must look out for signs of these conditions while you are taking TALTZ.

Stop using TALTZ and tell your healthcare professional or seek medical help immediately if you notice any signs indicating a possible serious infection or a serious allergic reaction.

Signs or symptoms of a potentially serious infection may include:

- fever, sweats, or chills
- muscle aches
- cough
- shortness of breath
- blood in your phlegm (mucus)
- weight loss
- warm, red, or painful skin or sores on your body
- diarrhea or stomach pain
- burning when you urinate or urinate more often than normal

Signs or symptoms of a serious allergic reaction may include:

- feeling faint
- swelling of your face, eyelids, lips, mouth, tongue, or throat
- trouble breathing or throat tightness
- chest tightness
- skin rash

Children and adolescents

TALTZ is not recommended for children and adolescents under 18 years of age because it has not been studied in this age group.

Tell your healthcare professional about all the medicines you take, including any drugs, vitamins, minerals, natural supplements or alternative medicines.

How to take TALTZ:

Always use this medicine exactly as your healthcare professional has told you. Check with your healthcare professional if you are not sure.

Each prefilled autoinjector and each prefilled syringe contains one dose of TALTZ (80 mg). Each prefilled autoinjector and each prefilled syringe delivers only one dose.

TALTZ is given via injection under your skin (known as a subcutaneous injection). You and your healthcare professional should decide if you should inject TALTZ yourself.

It is important not to try to inject yourself until you have been trained by your healthcare professional. A caregiver may also give you your TALTZ injection after proper training.

See the detailed Instructions for Use provided for information on how to prepare and inject a dose of TALTZ, and how to properly throw away used TALTZ prefilled autoinjectors and prefilled syringes.

Usual Dose:

Your doctor will decide how much TALTZ you need and for how long.

Plaque Psoriasis

- The initial dose is 160 mg (two injections) by subcutaneous injection.
- After the first dose you will receive further injections at Weeks 2, 4, 6, 8, 10, and 12. From Week 12, you will receive injections every 4 weeks. At each time point, after the initial dose, you will receive an 80 mg dose (one injection).

Psoriatic Arthritis

- The initial dose is 160 mg (two injections) by subcutaneous injection.
- After the first dose, you will receive 80 mg (one injection) every 4 weeks.
- For patients with psoriatic arthritis and with moderate-to-severe plaque psoriasis, use the dosing and administration recommendations for plaque psoriasis.
- For patients with psoriatic arthritis and with mild plaque psoriasis, use the dosing and administration recommendations for psoriatic arthritis.

Ankylosing Spondylitis

- You will receive 80 mg (one injection) by subcutaneous injection every 4 weeks.

TALTZ is for long-term treatment. Your healthcare professional will regularly monitor your condition to check that the treatment is having the desired effect.

Overdose:

If you accidentally inject more TALTZ than you should or the dose has been administered sooner than according to your doctor's prescription, inform your healthcare professional.

If you think you have taken too much TALTZ, contact your healthcare professional, hospital emergency department or regional Poison Control Centre immediately, even if there are no symptoms.

Missed Dose:

If you have forgotten to inject a dose of TALTZ, inject the next dose as soon as you remember. Then talk to your healthcare professional to discuss when you should inject the next dose.

What are possible side effects from using TALTZ?

Like all medicines, this medicine can cause side effects, although not everybody gets them.

Most of the following side effects are mild to moderate. If any of these side effects becomes severe, tell your healthcare professional.

Some side effects are very common (may affect more than 1 in 10 people):

- upper respiratory tract infections with symptoms such as sore throat and stuffy nose (nasopharyngitis)
- injections site reactions (e.g. rash, pain, itch or swelling)

Some side effects are common (may affect up to 1 in 10 people):

- feeling sick to your stomach (nausea)
- athlete's foot (tinea pedis)
- sore throat

- diarrhea
- headache
- inflammation of the eye with redness, pain, and blurred vision (iridocyclitis)

Some side effects are uncommon (may affect up to 1 in 100 people):

- oral thrush (oral candidiasis)
- fever, flu-like symptoms
- runny nose
- hives
- signs of low levels of white blood cells, such as fever, sore throat or mouth ulcers due to infections (neutropenia)
- discharge from the eye with itching, redness and swelling (conjunctivitis)
- stomach pain, diarrhea with or without blood, weight loss (inflammatory bowel disease)
- eczema
- itching

These are not all the possible side effects you may feel when taking TALTZ. If you experience any side effects not listed here, contact your healthcare professional.

Serious side effects and what to do about them			
Symptom / effect	Talk to your healthcare professional		Stop taking drug and get immediate medical help
	Only if severe	In all cases	
RARE Serious allergic reactions (such as feeling faint; swelling of your face eyelids, lips, mouth, tongue, or throat; trouble breathing or throat tightness; chest tightness; or skin rash)		✓	✓

If you have a troublesome symptom or side effect that is not listed here or becomes bad enough to interfere with your daily activities, talk to your healthcare professional.

Reporting Side Effects

You can help improve the safe use of health products for Canadians by reporting serious and unexpected side effects to Health Canada. Your report may help to identify new side effects and change the product safety information.

3 ways to report:

- Online at [MedEffect](#);
- By calling 1-866-234-2345 (toll-free);
- By completing a Patient Side Effect Reporting Form and sending it by:
 - Fax to 1-866-678-6789 (toll-free), or
 - Mail to: Canada Vigilance Program

Health Canada, Postal Locator 1908C
Ottawa, ON
K1A 0K9

Postage paid labels and the Patient Side Effect Reporting Form are available at [MedEffect](#).

NOTE: Contact your health professional if you need information about how to manage your side effects. The Canada Vigilance Program does not provide medical advice.

Storage:

Store in a refrigerator (2°C - 8°C). Do not freeze or shake. Do not use if TALTZ has been frozen. If needed, TALTZ may be left out of the refrigerator at a temperature up to 30°C (86°F) for up to 5 days. TALTZ should be discarded if not used within this 5-day period.

Store in the original packaging in order to protect from light.

Do not use TALTZ prefilled syringe or prefilled autoinjector:

- if you notice that it is damaged, or the medicine is cloudy, distinctly brown or yellow or has particles in it, or
- after the expiry date which is stated on the label and on the outer carton after “EXP”.

This medicine is for single use only. Ask your healthcare professional how to throw away medicines no longer required.

Keep out of reach and sight of children.

If you want more information about TALTZ:

- Talk to your healthcare professional
- Find the full product monograph that is prepared for healthcare professionals and includes this Patient Medication Information by visiting the Health Canada website; the manufacturer’s website www.lilly.ca, or by calling 1-888-545-5972.
- Instructions for Use can be found on www.lilly.ca.

For more information, please contact your healthcare professionals or pharmacist first, or Eli Lilly Canada Inc. at: 1-888-545-5972 or visit the website at www.lilly.ca.

The information in this document is current as of the last revision date shown below. For the most current information please visit our website or contact us directly.

TALTZ is a registered trademark owned by or licensed to Eli Lilly and Company, its subsidiaries or affiliates.

You may need to read this package insert again. Please do not throw it away until you have finished your medicine.

This leaflet was prepared by Eli Lilly Canada Inc.

Last Revised:

A4.0-TAL-PMI-0005-YYYYMMDD

INSTRUCTIONS FOR USE

How to use the TALTZ prefilled autoinjector

PrTALTZ®
(ixekizumab) injection
80 mg/mL



www.lilly.ca

PLEASE READ THESE INSTRUCTIONS BEFORE USE

Before you use the TALTZ prefilled autoinjector, read and carefully follow all the step-by-step instructions.

BEFORE USING YOUR PREFILLED AUTOINJECTOR:

IMPORTANT POINTS TO KNOW

- Read and carefully follow all the instructions. Keep the Instructions for Use and refer to them as needed.
- The prefilled autoinjector contains 1 dose of TALTZ. The prefilled autoinjector is for ONE-TIME USE ONLY.
- The prefilled autoinjector contains glass parts. Handle it carefully. If you drop it on a hard surface, do not use it. Use a new prefilled autoinjector for your injection.
- Your healthcare professional may help you decide where on your body to inject your dose. You can also read the GET READY section of these instructions to help you choose which area can work best for you.
- Read the TALTZ Patient Medication Information inside this box to learn more about your medicine.


IMPORTANT SAFETY INFORMATION

- If you have questions or need help with your prefilled autoinjector, call your healthcare professional or visit the manufacturer's website at www.lilly.ca or call 1-888-545-5972.
- If you have vision problems, **DO NOT** use the prefilled autoinjector, without help from a person trained to use it.
- Keep the prefilled autoinjector out of the reach and sight of children.
- If you do not have a sharps container, ask your healthcare professional where you can get one.

INSTRUCTIONS FOR USE

Before you use the TALTZ prefilled autoinjector, read and carefully follow all the step-by-step instructions.

Guide to the parts



1 GET READY

- 1a Take the prefilled autoinjector from the refrigerator.** Leave the base cap on until you are ready to inject. **Wait 30 minutes** to let the prefilled autoinjector warm to room temperature before you use it. This will make the medicine easier to inject.




DO NOT microwave the prefilled autoinjector, run hot water over it, or leave it in direct sunlight.

- 1b Gather the supplies for your injection:**

- 1 alcohol wipe
- 1 cotton ball or piece of gauze
- 1 sharps container for disposal of prefilled autoinjector

- 1c**



Inspect the prefilled autoinjector. Check the label. Make sure the name TALTZ appears on the label. Also make sure the medicine has not expired.


The medicine inside should be clear. Its color may vary from colorless to slightly yellow.

If you see any of the following, **DO NOT USE** the prefilled autoinjector, and dispose of it as directed:

- It is past the expiration date
- It looks damaged
- The medicine is cloudy, is distinctly brown, or has small specks

- 1d Wash your hands before you inject your medicine.**

1e



Talk with your healthcare professional about where on your body may be best to inject your dose.

Choose your injection site.

You may inject in your abdomen (stomach area), in your thigh, or in the back of your arm. To inject in your arm, you will need someone to help you.


DO NOT inject into areas where the skin is tender, bruised, red, scaly, or hard or where you have scars or stretch marks. **DO NOT** inject within 1 inch (2.5 centimeters) around of the navel (belly button).

Alternate your injection sites. DO NOT inject in the exact same spot every time. For example, if your last injection was in your left thigh, your next injection should be in your right thigh, your abdomen, or the back of either arm.

1f Prepare your skin. Clean your skin with an alcohol wipe. Let the injection site dry before you inject your medicine.

2 INJECT

2a




Make sure the lock ring is in the lock position.

Leave the base cap on until you are ready to inject. **DO NOT** touch the needle.

Twist off the base cap.


Throw the base cap in the garbage. You will not need to put the base cap back on—doing so could damage the needle or cause you to stick yourself by accident.

2b




Place the clear base flat and firmly against your skin.

2c



Keep the base on your skin, and then turn the lock ring to the unlock position. You are now ready to inject.

2d



Press the green injection button. There will be a loud click.

Keep holding the clear base firmly against your skin. You will hear a second click in about 5 to 10 seconds after the first one. **The second click tells you that your injection is complete.**


You will also see the gray plunger at the top of the clear base.

Remove the prefilled autoinjector from your skin.

Press a cotton ball or gauze over the injection site. **DO NOT** rub the injection site, as this may cause bruising. You may have slight bleeding. This is normal.

3 FINISH

3a



Dispose of the prefilled autoinjector.

DO NOT put the base cap back on. Dispose of the used prefilled autoinjector in a sharps container such as a closeable, puncture-resistant container.

When you dispose of the prefilled autoinjector and the sharps container:

- Put the prefilled autoinjector in a sharps container (like a biohazard container) or a hard plastic container with a secure lid. Do not throw the prefilled autoinjector directly into your household garbage.
- Do not recycle the filled container. The full container must be disposed of according to your provincial and local laws.
- For information on how to dispose of the container properly, ask your healthcare professional about options available in your area.

Commonly asked questions

Q. What if I see bubbles in the prefilled autoinjector?

A. It is normal to have air bubbles in the prefilled autoinjector. TALTZ is injected under the skin (subcutaneous injection). Air bubbles are not a problem in this type of injection. They will not harm you or affect your dose.

Q. What if there is a drop of liquid on the tip of the needle when I remove the base cap?

A. It is okay to see a drop of liquid on the tip of the needle. This will not harm you or affect your dose.

Q. What if I unlocked the prefilled autoinjector and pressed the green injection button before I twisted off the base cap?

A. Do not remove the base cap. Contact 1-888-545-5972.

Q. Do I need to hold the injection button down until the injection is complete?

A. This is not necessary, but it may help you keep the prefilled autoinjector steady and firm against your skin.

Q. What if the needle did not retract after my injection?

A. Do not touch the needle or replace the base cap. Store in a safe place and contact 1-888-545-5972.

Q. What if I heard more than 2 clicks during my injection—2 loud clicks and a soft one. Did I get my complete injection?

A. Some patients may hear a soft click right before the second loud click. That is the normal operation of the prefilled autoinjector. Do not remove the prefilled autoinjector from your skin until you hear the second loud click.

Q. How can I tell if my injection is complete?

A. After you press the green injection button, you will hear 2 loud clicks. The second click tells you that your injection is complete. You will also see the gray plunger at the top of the clear base.

For questions or more information about TALTZ

It's important to know how to inject your medicine correctly and safely. If you have questions about TALTZ prefilled autoinjector:

- Talk to your healthcare professional
- Call Lilly at 1-888-545-5972
- Visit www.lilly.ca

How to store your TALTZ prefilled autoinjector

DO



DO store the prefilled autoinjector in the refrigerator between 2°C to 8°C (36°F to 46°F) until you are ready to use it.

- If needed, a prefilled autoinjector may be left out of the refrigerator at a temperature up to 30°C (86°F) for up to 5 days. TALTZ should be discarded if not used within this 5-day period.



DO wait 30 minutes to let the prefilled autoinjector warm to room temperature before you use it.

DO NOT



DO NOT freeze the prefilled autoinjector. If the prefilled autoinjector has been frozen, **do not use it**.



DO NOT microwave the prefilled autoinjector, run hot water over it, or leave it in direct sunlight.

DO NOT shake the prefilled autoinjector.

Read the Patient Medication Information for TALTZ inside this box to learn more about your medicine.

TALTZ is a registered trademark owned or licensed to Eli Lilly and Company, its subsidiaries or affiliates.

Copyright © 2016, 2019, Eli Lilly and Company. All rights reserved.

Document Revision Date: December 4, 2019


The TALTZ prefilled autoinjector meets the current dose accuracy and functional requirements of ISO 11608-1:2012 and 11608-5:2012

TAL-AI-0004-CA-IFU-20191204

INSTRUCTIONS FOR USE

How to use the TALTZ prefilled syringe

PrTALTZ®
(ixekizumab) injection
80 mg/mL



www.lilly.ca

Lilly

PLEASE READ THESE INSTRUCTIONS BEFORE USE

Before you use the TALTZ prefilled syringe, read and carefully follow all the step-by-step instructions.

BEFORE USING YOUR PREFILLED SYRINGE:

IMPORTANT POINTS TO KNOW

- Read and carefully follow all the instructions. Keep the Instructions for Use and refer to them as needed.
- The prefilled syringe contains 1 dose of TALTZ. The syringe is for ONE-TIME USE ONLY.
- Your healthcare professional may help you decide where on your body to inject your dose. You can also read the GET READY section of these instructions to help you choose which area can work best for you.
- Read the TALTZ Patient Medication Information inside this box to learn more about your medicine.

IMPORTANT SAFETY INFORMATION


- If you have questions or need help with your prefilled syringe, call your healthcare professional or visit the manufacturer's website at www.lilly.ca or call 1-888-545-5972.
- If you have vision problems, **DO NOT** use the prefilled syringe, without help from a person trained to use it.
- **DO NOT** share or reuse your TALTZ prefilled syringe. You may give or get an infection.

- Keep the syringe out of the reach and sight of children.
- If you do not have a sharps container, ask your healthcare professional where you can get one.

INSTRUCTIONS FOR USE

Before you use the TALTZ prefilled syringe, read and carefully follow all the step-by-step instructions.

Guide to the parts



1 GET READY

- 1a Take the syringe from the refrigerator.** Leave the needle cap on the syringe until you are ready to inject. **Wait 30 minutes** to let the syringe warm to room temperature before you use it. This will make the medicine easier to inject.




DO NOT microwave the syringe, run hot water over it, or leave it in direct sunlight.

- 1b Gather the supplies for your injection:**

- 1 alcohol wipe
- 1 cotton ball or piece of gauze
- 1 sharps container for disposal of syringe

1c



Inspect the syringe for damage to the outside. Leave the needle cap on the syringe until you are ready to inject. Check the label. Make sure the name TALTZ appears on the label. Also make sure the medicine has not expired.


The medicine inside should be clear. Its color may vary from colorless to slightly yellow.

If you see any of the following, **DO NOT USE** the syringe, and dispose of it as directed:

- It is past the expiration date
- It looks damaged
- The medicine is cloudy, is distinctly brown, or has small specks

1d **Wash your hands before you inject your medicine.**

1e



Choose your injection site.

You may inject in your abdomen (stomach area), in your thigh, or in the back of your arm. To inject in your arm, you will need someone to help you.

DO NOT inject into areas where the skin is tender, bruised, red, scaly, or hard or where you have scars or stretch marks. **DO NOT** inject within 1 inch (2.5 centimeters) of the navel (belly button).


Alternate your injection sites. DO NOT inject in the exact same spot every time. For example, if your last injection was in your left thigh, your next injection should be in your right thigh, your abdomen, or the back of either arm.

Talk with your healthcare professional about where on your body may be best to inject your dose.

1f **Prepare your skin.** Clean your skin with an alcohol wipe. Let the injection site dry before you inject your medicine.

2 INJECT

2a




Pull the needle cap off and throw it away.

DO NOT put the cap back on—you could damage the needle or stick yourself by accident.


DO NOT touch the needle.

2b




Gently pinch and hold a fold of skin where you will inject.

2c




Insert the needle at a 45-degree angle. Then gently let go of your skin. Make sure to keep the needle in place.



Let go of your skin before you push the plunger in.

2d



Push in the plunger.

Slowly push the plunger all the way in until all the medicine is injected. The gray syringe plunger should be pushed all the way to the needle end of the syringe. Gently remove the needle from your skin.

Press a cotton ball or gauze over the injection site. **DO NOT** rub the injection site, as this may cause bruising. You may have slight bleeding. This is normal.

You should see the green plunger rod show through the syringe body when the injection is complete.

3 FINISH

3a



Dispose of the syringe.

DO NOT put the needle cap back on. Dispose of the used syringe in a sharps container such as a closeable, puncture-resistant container.

When you dispose of syringes and the sharps container:

- Put the syringe in a sharps container (like a biohazard container) or a hard plastic container with a secure lid. Do not throw the syringe directly into your household garbage.
- Do not recycle the filled container. The full container must be disposed of according to your provincial and local laws.
- For information on how to dispose of the container properly, ask your healthcare professional about options available in your area.

Commonly asked questions

Q. What if I see air bubbles in my syringe?

A. It is normal to sometimes have air bubbles in the syringe. TALTZ is injected under your skin (subcutaneous injection). Air bubbles are not a problem in this type of injection. They will not harm you or affect your dose.

Q. What if there is a drop of liquid on the tip of the needle when I remove the needle cap?

A. It is okay to see a drop of liquid on the tip of the needle. This will not harm you or affect your dose.

Q. What if I cannot push in the plunger?

A. If the plunger is stuck or damaged:

- **DO NOT** continue to use the syringe
- Remove the needle from your skin
- Contact 1-888-545-5972

Q. How can I tell if my injection is complete?

A. When your injection is complete:

- The green plunger rod should show through the body of the syringe
- The gray syringe plunger should be pushed all the way to the needle end of the syringe

For questions or more information about TALTZ

It's important to know how to inject your medicine correctly and safely. If you have questions about TALTZ prefilled syringe:

- Talk to your healthcare professional
- Call Lilly at 1-888-545-5972
- Visit www.lilly.ca

How to store your TALTZ prefilled syringe

DO



DO store the syringe in the refrigerator between 2°C to 8°C (36°F to 46°F) until you are ready to use it.

- If needed, a prefilled syringe may be left out of the refrigerator at a temperature up to 30°C (86°F) for up to 5 days. TALTZ should be discarded if not used within this 5-day period.



DO wait 30 minutes to let the syringe warm to room temperature before you use it.

DO NOT



DO NOT freeze the syringe. If the syringe has been frozen, **do not use it**.



DO NOT microwave the syringe, run hot water over it, or leave it in direct sunlight.

DO NOT shake the syringe.

Read the Patient Medication Information for TALTZ inside this box to learn more about your medicine.

TALTZ is a registered trademark owned or licensed to Eli Lilly and Company, its subsidiaries or affiliates.

Copyright © 2016, 2019, Eli Lilly and Company. All rights reserved.

Document Revision Date:

TAL-PFS-0003-CA-IFU-20191204