

PRODUCT MONOGRAPH

^{Pr} Sodium Nitroprusside for Injection

25 mg / mL Sodium Nitroprusside Dihydrate

Antihypertensive Agent

Manufactured By:
Dr. Reddy's Laboratories Limited
Bachupally – 500090
India.

Date of Preparation:
Oct 29, 2020

Imported and Distributed By:
Dr. Reddy's Laboratories Canada Inc.
Mississauga, ON L4W 4Y1
Canada

Submission Control Number: 234455

Table of Contents

PART I: HEALTH PROFESSIONAL INFORMATION.....3
SUMMARY PRODUCT INFORMATION3
INDICATIONS AND CLINICAL USE.....3
CONTRAINDICATIONS3
WARNINGS AND PRECAUTIONS.....4
ADVERSE REACTIONS.....7
DRUG INTERACTIONS7
DOSAGE AND ADMINISTRATION7
OVERDOSAGE9
ACTION AND CLINICAL PHARMACOLOGY10
STORAGE AND STABILITY11
DOSAGE FORMS, COMPOSITION AND PACKAGING11

PART II: SCIENTIFIC INFORMATION12
PHARMACEUTICAL INFORMATION.....12
DETAILED PHARMACOLOGY12
TOXICOLOGY15
REFERENCES17

PART III: CONSUMER INFORMATION.....19

Pr Sodium Nitroprusside for Injection

PART I: HEALTH PROFESSIONAL INFORMATION

SUMMARY PRODUCT INFORMATION

Route of Administration	Dosage Form / Strength	Nonmedicinal Ingredients
Intravenous	Solution for injection; 25 mg / mL	Water for Injection

INDICATIONS AND CLINICAL USE

Sodium Nitroprusside for Injection is indicated for the treatment of acute hypertension, refractory to standard therapeutic measures.

Sodium Nitroprusside for Injection is also indicated for producing controlled hypotension during anesthesia in order to reduce bleeding in surgical procedures, where surgeon and anesthesiologist deem it appropriate. In each case, the benefit-risk ratio should be carefully considered on an individual basis.

CONTRAINDICATIONS

- Patients who are hypersensitive to sodium nitroprusside or to any ingredient in the formulation or component of the container. For a complete listing, see the **DOSAGE FORMS, COMPOSITION AND PACKAGING**.
- Sodium Nitroprusside for Injection should not be used in the treatment of compensatory hypertension (e.g., arteriovenous shunt or coarctation of the aorta). It is also contraindicated in physically poor-risk patients (A.S.A. Risk 5), in patients with uncorrected anemia or hypovolemia, or in those with known inadequate cerebral circulation.
- Controlled hypotension during anaesthesia induced by sodium nitroprusside is contraindicated in patients with liver disease, severe renal disease, Leber's optic atrophy, tobacco amblyopia and disease states associated with vitamin B₁₂ deficiency.

WARNINGS AND PRECAUTIONS

Serious Warnings and Precautions

Sodium Nitroprusside for Injection is only to be used as an infusion with 5% Dextrose Injection. Not for direct injection.

Sodium Nitroprusside for Injection can cause precipitous decreases in blood pressure. In patients not properly monitored, these decreases can lead to irreversible ischemic injuries or death. Sodium Nitroprusside for Injection should be used only when available equipment and personnel allow blood pressure to be continuously monitored.

Except when used briefly or at low (< 2 mcg/kg/min) infusion rates, Sodium Nitroprusside for Injection gives rise to important quantities of cyanide ion, which can reach toxic, potentially lethal levels. The usual dose rate is 0.5 to 8 mcg/kg/min, but infusion at the upper dose rate should never last more than 10 minutes. If blood pressure has not been adequately controlled after 10 minutes of infusion at 8 mcg/kg/min, administration of Sodium Nitroprusside for Injection should be terminated immediately. Infusion rates greater than 8 mcg/kg/min are virtually never required.

Although acid-base balance and venous oxygen concentration should be monitored and may indicate cyanide toxicity, these laboratory tests provide imperfect guidance.

Fatalities due to cyanide poisoning have occurred following sodium nitroprusside administration. One factor is common to all known cases, namely that large amounts of nitroprusside were infused at high rates, since detoxification relies upon enzymatic action, the rare possibility of deficient or atypical enzymes occurring in humans should always be considered. Patients most apt to run into difficulty are those who are resistant to the hypotensive effect or those in whom maintenance at the selected blood pressure level is difficult or impossible.

Constant attention to the patient's dose-response characteristics is mandatory. If infusion rates are in excess of 8 mcg/kg/min, determine the nature of the response (effective constant response at higher dose; tachyphylactic; resistant- none or less than the expected). As soon as either tachyphylaxis or resistance is determined, the infusion of Sodium Nitroprusside for Injection should be discontinued immediately. In abnormal responders, acid-base balance should be monitored since metabolic acidosis is evidence of cyanide toxicity.

Cardiovascular

Adequate facilities, equipment and trained personnel should be available for frequent and vigilant monitoring of blood pressure. The hypotensive effects of Sodium Nitroprusside for Injection occur rapidly, and blood pressure usually begins to rise immediately and returns to pre-treatment levels within one to ten minutes when the infusion is slowed or stopped. It should be used with caution and initially in low doses in elderly patients, since they may be more sensitive to the hypotensive effects of the drug.

Because of the rapid onset of action and potency of Sodium Nitroprusside for Injection, it should be administered with the use of an infusion pump, micro-drip regulator, or any similar device that would allow precise measurement of the flow rate.

If in the clinical situation, stress, induced by pain or manipulation, is reduced or eliminated during sodium nitroprusside infusion, the patient could experience a greater than expected reduction in blood pressure unless the rate of infusion is adjusted downward as required.

Endocrine and Metabolism

Sodium Nitroprusside for Injection is metabolized to cyanide and then to thiocyanate which in turn is excreted by the kidney (see **ACTION AND CLINICAL PHARMACOLOGY**). If excessive amounts of Sodium Nitroprusside for Injection are used, thiocyanate toxicity (e.g., tinnitus, blurred vision (miosis), delirium hyperreflexia) may occur. Estimating the thiocyanate blood levels, will help to determine thiocyanate toxicity, but may not reflect cyanide toxicity. Thiocyanate is mildly neurotoxic at serum levels of 60 mg/L (1 mmol/L). Thiocyanate toxicity is life-threatening when levels are three or four times higher (200 mg/L).

Since thiocyanate inhibits both the uptake and binding of iodine, caution should be exercised in using Sodium Nitroprusside for Injection in patients with hypothyroidism.

Hematologic

Sodium nitroprusside infusions at rates above 2 mcg/kg/min generate cyanide ion (CN⁻) faster than the body can normally dispose of it. When sodium thiosulfate is given, the body's capacity for CN⁻ elimination is greatly increased. Methemoglobin, normally present in the body, can buffer a certain amount of CN⁻, but the capacity of this system is exhausted by the CN⁻ produced from about 500 mcg/kg of sodium nitroprusside (see **ACTION AND CLINICAL PHARMACOLOGY**). This amount of sodium nitroprusside is administered in approximately one hour when the drug is administered at 8 mcg/kg/min.

Neurologic

Signs of cerebral underperfusion, such as confusion and somnolence, may occur if blood pressure is reduced too rapidly, especially in hypertensive patients with encephalopathy.

Peri-Operative Considerations

The following warnings apply to the use of Sodium Nitroprusside for Injection for controlled hypotension during anaesthesia:

1. Extreme caution should be exercised in patients who are especially poor surgical risks (A.S.A class 4 and 4E).
2. Tolerance to blood loss, anemia and hypovolemia may be diminished. If possible, pre-existing anemia and hypovolemia should be corrected prior to employing controlled hypotension.

3. Hypotensive anaesthetic techniques may alter pulmonary ventilation perfusion ratio. Patients intolerant of additional dead air space at ordinary oxygen partial pressure may benefit from higher oxygen partial pressure.
4. Resistance and tachyphylaxis occur more frequently in normotensive patients infused with sodium nitroprusside. Induction of deliberate hypotension in healthy young individuals may prove to be more difficult than in other segments of the population.
5. Upon discontinuation of the sodium nitroprusside infusion, for the purpose of controlled hypotension during anaesthesia, a rebound hypertension has been observed on rare occasions.

Renal

Since thiocyanate inhibits both the uptake and binding of iodine, caution should be exercised in using Sodium Nitroprusside for Injection in patients with severe renal impairment.

Blood levels of thiocyanate should be determined if treatment is to be extended. This monitoring is critical in patients with severe renal dysfunction. Although intravenous infusions of Sodium Nitroprusside for Injection are not intended for long-term use, as long as blood thiocyanate levels are measured daily and do not exceed 10 mg percent (100 mg/L), it is probably safe to continue with the infusion until the patient can be safely treated with oral antihypertensive medications. Peritoneal dialysis can be helpful if too high levels of thiocyanate are found.

Sensitivity/Resistance

There is marked variation in individual sensitivity to the antihypertensive action of sodium nitroprusside.

Hypertensive patients are more sensitive to the intravenous effect of sodium nitroprusside than are normotensive subjects.

Several authors have reported tachyphylaxis in young male patients during hypotensive anaesthesia. However, tachyphylaxis has not been reported to date with sodium nitroprusside in the treatment of hypertensive emergencies.

Special Populations

Pregnant Women: The safety of sodium nitroprusside in women who are or who may become pregnant has not been established. Hence, it should be given only when the potential benefits have been weighed against possible hazard to the mother and child.

Nursing Mothers: It is not known whether sodium nitroprusside and its metabolites are excreted in human milk. Because many drugs are excreted in human milk and because of the potential for serious adverse reactions in nursing infants from sodium nitroprusside, a decision should be made whether to discontinue nursing or to discontinue the drug, taking into account the importance of the drug to the mother.

Geriatrics: Sodium Nitroprusside for Injection should be used with caution and initially in low doses in elderly patients, since they may be more sensitive to the hypotensive effects of the drug.

ADVERSE REACTIONS

Adverse Drug Reaction Overview

Fatalities due to cyanide poisoning have occurred following sodium nitroprusside administration (see **WARNINGS AND PRECAUTIONS**).

Nausea, retching, emesis, diaphoresis, apprehension, headache, restlessness, agitation, muscle twitching, retrosternal discomfort and chest pain, palpitations, dizziness, faintness, weakness, rash, abdominal pain, confusion and somnolence have been noted with too rapid reduction in blood pressure. However, these symptoms rapidly disappeared with slowing of the rate of the infusion or temporary discontinuation of infusion, and did not reappear with continued slower rate of administration.

Irritation of the injection site may occur.

One case of hypothyroidism following prolonged therapy with intravenous sodium nitroprusside has been reported. Thiocyanate blood levels were 9.5 mg/100 mL after 21 days of administration of sodium nitroprusside to this patient with severe hypertension and renal disease.

Methemoglobinemia has been reported in the literature. Sodium nitroprusside infusions can cause sequestration of hemoglobin as methemoglobin (see **ACTION AND CLINICAL PHARMACOLOGY**). Clinically significant methemoglobinemia (> 10%) is seen only rarely in patients receiving sodium nitroprusside. Methemoglobinemia should be suspected in patients who have received > 10 mg/kg of sodium nitroprusside and who exhibit signs of impaired oxygen delivery.

DRUG INTERACTIONS

Drug-Drug Interactions

Patients who are also receiving concomitant antihypertensive medications (specifically, hydralazine or hexamethonium) are more sensitive to the hypotensive effect of sodium nitroprusside, and the dosage of Sodium Nitroprusside for Injection should be adjusted downward accordingly.

DOSAGE AND ADMINISTRATION

Dosing Considerations

Use of Sodium Nitroprusside for Injection in anaesthetized normotensive patients undergoing deliberate hypotensive surgery must be restricted to carefully selected cases. There is a possibility of an abnormal response occurring in normotensive patients. In the event of an abnormal response, the infusion of Sodium Nitroprusside for Injection should be discontinued immediately (see WARNINGS AND PRECAUTIONS).

Recommended Dose and Dosage Adjustment

Depending on the desired concentration, 2 mL of Sodium Nitroprusside for Injection (50 mg of Sodium Nitroprusside Dihydrate) must be further diluted in 500 to 1000 mL of 5% Dextrose Injection. **No other diluents should be used.** The diluted solution should be protected from light, using the supplied opaque sleeve, aluminum foil or other opaque materials, the infusion solution should be freshly prepared and any unused portion discarded. The freshly prepared solution for infusion has a very faint brownish tint. If it is highly coloured, it should be discarded (see **WARNINGS AND PRECAUTIONS**).

Storage period from the time of reconstitution to the completion of intravenous administration should not exceed 24 hours. The infusion fluid used for administration of Sodium Nitroprusside for Injection should not be employed as a vehicle for simultaneous administration of any other drug.

As with all parenteral drug products, intravenous admixtures should be inspected visually for clarity, particulate matter, precipitate, discolouration and leakage prior to administration whenever solution and container permit. Solutions showing haziness, particulate matter, precipitate, discolouration or leakage should not be used. Discard unused portion.

The intravenous infusion of Sodium Nitroprusside for Injection should be administered by an infusion pump, micro-drip regulator or any similar device that will allow precise measurement of the flow rate. Care should be taken to avoid extravasation.

2 mL of Sodium Nitroprusside for Injection (50 mg of Sodium Nitroprusside Dihydrate) in 1000 mL D5W (50 mcg/mL)

<u>Dose/kg</u>	<u>mcg/kg/min</u>	<u>mL/kg/min</u>
Average	3	0.06
Range	0.5 to 8	0.01 to 0.16

2 mL of Sodium Nitroprusside for Injection (50 mg of Sodium Nitroprusside Dihydrate) in 500 mL D5W (100 mcg/mL)

<u>Dose/kg</u>	<u>mcg/kg/min</u>	<u>mL/kg/min</u>
Average	3	0.03
Range	0.5 to 8	0.005 to 0.08

In patients who are not receiving antihypertensive drugs, the average dose of Sodium Nitroprusside for Injection for both adults and children is 3 mcg/kg/minute (range of 0.5 to 8 mcg/kg/minute). Usually, at 3 mcg/kg/minute, blood pressure can be lowered by about 30 to 40% below the pre-treatment diastolic levels and maintained.

In hypertensive patients receiving concomitant antihypertensive medications, smaller doses are required.

In order to avoid excessive levels of thiocyanate and to lessen the possibility of a precipitous drop in blood pressure, infusion rates greater than 8 mcg/kg/minute should rarely be used. If an adequate reduction of blood pressure is not obtained within ten minutes at this rate, administration of Sodium Nitroprusside for Injection should be stopped.

Sodium Nitroprusside for Injection dosage varies considerably from patient to patient, hence the need for individual titration. The infusion should be started at the lower dosage range (0.5 mcg/kg/minute) and adjusted in a stepwise fashion (e.g. increments of 0.2 mcg/kg/minute usually every five minutes) until the desired reduction in blood pressure is obtained. Continuous careful monitoring of the blood pressure on a minute to minute basis is necessary. Adjustments in the rate of infusion may be required to keep the blood pressure smoothly controlled and prevent extremes of hypotension and hypertension.

The blood pressure usually starts to drop immediately or at least within a few minutes. It is recommended that the blood pressure should not be allowed to drop at a too rapid rate and that the systolic pressure not be lowered below 60 mm Hg. Too rapid a reduction in blood pressure may result in retching or vomiting, muscular twitching, diaphoresis and agitation. These symptoms subside promptly when the rate of infusion is slowed or temporarily stopped.

In hypertensive emergencies, Sodium Nitroprusside for Injection infusion may be continued until the patient can safely be treated with oral antihypertensive medications alone.

OVERDOSAGE

For management of a suspected drug overdose, contact your regional Poison Control Centre.

Overdosage of nitroprusside can be manifested as excessive hypotension or cyanide toxicity or as thiocyanate toxicity. In moderate cases, the signs are dyspnea, headache, vomiting, dizziness, ataxia and loss of consciousness. Massive overdosage produces coma with imperceptible pulse, absent reflexes, widely dilated pupils, pink colour, distant heart sounds, hypotension and very shallow breathing. Relief with oxygen only is not seen. Death may result. High overdosage also results in the occurrence of hyperkalemia and metabolic acidosis which will require appropriate measures for correction.

Treatment of cyanide toxicity: Cyanide levels can be measured by many laboratories, and blood-gas studies that can detect venous hyperoxemia or acidosis are widely available. Acidosis may not appear until more than an hour after the appearance of dangerous cyanide levels, and laboratory tests should not be awaited. Reasonable suspicion of cyanide toxicity is adequate grounds for initiation of treatment.

The treatment of cyanide toxicity consists of: discontinuing the administration of Sodium Nitroprusside for Injection; providing a buffer for cyanide by using sodium nitrite to convert as much hemoglobin into methemoglobin as the patient can safely tolerate; and then infusing sodium thiosulfate in sufficient quantity to convert the cyanide into thiocyanate.

Treatment:

- a. Immediate discontinuation of Sodium Nitroprusside for Injection infusion or any other medication.
- b. Should the patient be conscious, amyl nitrite ampoules should be administered by inhalation immediately, one for 30 seconds every two minutes, unless pressure is below 80 mm Hg. If there is a delay in obtaining 3% sodium nitrite solution, amyl nitrite may be repeated as indicated.
- c. Follow as soon as possible (but not together with amyl nitrite) by intravenous injection of 10 mL of 3% sodium nitrite over a three-minute period (i.v. infusion of noradrenaline may be necessary to maintain blood pressure during this injection).
- d. 50 mL of 25% sodium thiosulfate should be administered intravenously over a ten-minute period following the administration of sodium nitrite.
- e. Supportive measures should be instituted as soon as possible (e.g. artificial respiration with 100% oxygen).
- f. Administration of sodium nitrite and sodium thiosulfate may have to be repeated if symptoms re-appear, but doses should be reduced by 50%.
- g. The institution of peritoneal dialysis may be helpful in the reduction of thiocyanate levels.
- h. Constant monitoring for cyanide and thiocyanate blood levels should be undertaken.
- i. If a severe and prolonged hypoxemia results due to excessive methemoglobinemia, inhalation of pure oxygen or a blood transfusion may be required.
- j. Further treatment should be symptomatic.

A cyanide antidote should be used if needed.

ACTION AND CLINICAL PHARMACOLOGY

Mechanism of Action

Sodium nitroprusside is a potent, rapid-acting, intravenous antihypertensive agent. The antihypertensive action of sodium nitroprusside is probably due to the nitroso (NO) group. Its effect is almost immediate and usually ends when the intravenous infusion is stopped. The brief duration of the drug's action is due to its rapid biotransformation. The hypotensive effect is augmented by ganglionic blocking agents. The hypotensive effects of sodium nitroprusside are caused by peripheral vasodilation as a result of a direct action on the blood vessels, independent of autonomic innervation. No relaxation is seen in the smooth muscle of the uterus or duodenum *in situ* in animals.

Sodium nitroprusside administered intravenously to hypertensive and normotensive patients produced a marked lowering of the arterial blood pressure, a slight increase in heart rate, a mild decrease in cardiac output and a moderate diminution in calculated total peripheral vascular resistance.

The decrease in calculated total peripheral vascular resistance suggests arteriolar vasodilation. The decreases in cardiac and stroke index noted may be due to the peripheral vascular pooling of blood.

STORAGE AND STABILITY

Sodium Nitroprusside for Injection should be stored at controlled room temperature (15 to 30°C). It should be protected from light and freezing.

Once diluted in 5% Dextrose Injection, Sodium Nitroprusside for Injection tends to deteriorate in the presence of light. The diluted solution should also be protected from light, using the supplied opaque sleeve, aluminium foil, or other opaque material. It is not necessary to cover the infusion drip chamber or the tubing.

Storage period of solutions of Sodium Nitroprusside for Injection from the time of reconstitution to the completion of intravenous administration should not exceed 24 hours. Sodium Nitroprusside for Injection in aqueous solution yields the nitroprusside ion, which reacts with even minute quantities of a wide variety of inorganic and organic substances to form usually highly coloured reaction products (blue, green or dark red). If this occurs, the infusion should be replaced as quickly as possible.

DOSAGE FORMS, COMPOSITION AND PACKAGING

Composition:

Sodium Nitroprusside for Injection is a solution available in single-use amber glass vials (with latex-free stopper) in 50 mg / 2 mL format (25 mg / mL), containing sodium nitroprusside dihydrate, in water for injection.

PART II: SCIENTIFIC INFORMATION

PHARMACEUTICAL INFORMATION


Drug Substance

Proper name: Sodium nitroprusside

Chemical name: Ferrate(2-), pentakis(cyano-C)nitrosyl-, disodium, hydrate, (OC-6-22)-

Molecular formula and molecular mass: $[\text{Na}_2\text{Fe}(\text{CN})_5\text{NO}] \cdot 2\text{H}_2\text{O}$ (297.95 g/mol)

Structural formula:



Physicochemical properties:

Physical description: Reddish-brown, crystals or powder

Solubility: Freely soluble in water

pH: 5.0 – 7.0 (25 mg / mL solution in water)

DETAILED PHARMACOLOGY


Human Pharmacology

The hypotensive effects of intravenously administered sodium nitroprusside are due to peripheral vasodilatation and reduction in peripheral resistance as a result of a direct action on the blood vessel walls, independent of autonomic innervation. The active component in sodium nitroprusside is the free nitroprusside radical. The evanescent nature of the drug's hypotensive effect is due to the destruction of the active radical. Infused sodium nitroprusside is rapidly distributed to a volume that is approximately coextensive with the extracellular space. The drug is cleared from this volume by intraerythrocytic reaction with hemoglobin (Hgb), and sodium nitroprusside's resulting circulatory half-life is about two minutes.

The products of the nitroprusside/hemoglobin reaction are cyanmethemoglobin (cyanmetHgb) and cyanide ion (CN^-). Safe use of Sodium Nitroprusside for Injection must be guided by knowledge of the further metabolism of these products.

As shown in the diagram below, the essential features of nitroprusside metabolism are:

- One molecule of sodium nitroprusside is metabolized by combination with hemoglobin to produce one molecule of cyanmethemoglobin and four CN⁻ ions;
- Methemoglobin, obtained from hemoglobin, can sequester cyanide as cyanmethemoglobin;
- Thiosulfate reacts with cyanide to produce thiocyanate;
- Thiocyanate is eliminated in the urine.
- Cyanide not otherwise removed binds to cytochromes; and
- Cyanide is much more toxic than methemoglobin or thiocyanate.



Cyanide ion is normally found in serum; it is derived from dietary substrates and from tobacco smoke.

Cyanide binds avidly (but reversibly) to ferric ion (Fe⁺⁺⁺), most body stores of which are found in erythrocyte methemoglobin (metHgb) and in mitochondrial cytochromes. When CN⁻ is infused or generated within the bloodstream, essentially all of it is bound to methemoglobin until intraerythrocytic methemoglobin has been saturated.

When the Fe⁺⁺⁺ of cytochromes is bound to cyanide, the cytochromes are unable to participate in oxidative metabolism. In this situation, cells may be able to provide for their energy needs by utilizing anaerobic pathways, but they thereby generate an increasing body burden of lactic acid. Other cells may be unable to utilize these alternate pathways, and they may die hypoxic deaths. CN⁻ levels in packed erythrocytes are typically less than 1 mmol/L (less than 25 mcg/L); levels are roughly doubled in heavy smokers.

At healthy steady-state, most people have less than 1% of their hemoglobin in the form of methemoglobin. Nitroprusside metabolism can lead to methemoglobin formation (a) through dissociation of cyanmethemoglobin formed in the original reaction of sodium nitroprusside with Hgb and (b) by direct oxidation of Hgb by the released nitroso group. Relatively large quantities of sodium nitroprusside, however, are required to produce significant methemoglobinemia.

At physiologic methemoglobin levels, the CN^- binding capacity of packed red cells is a little less than 200 $\mu\text{mol/L}$ (5 mg/L). Cytochrome toxicity is seen at levels only slightly higher, and death has been reported at levels from 300 to 3,000 $\mu\text{mol/L}$ (8 to 80 mg/L). Put another way, a patient with a normal red-cell mass (35 mL/kg) and normal methemoglobin levels can buffer about 175 mcg/kg of CN^- , corresponding to little less than 500 mcg/kg of infused sodium nitroprusside.

Some cyanide is eliminated from the body as expired hydrogen cyanide, but most is enzymatically converted to thiocyanate (SCN^-) by thiosulfate-cyanide sulfur transferase (rhodanase, EC 2.8.1.1), a mitochondrial enzyme. The enzyme is normally present in great excess, so the reaction is rate-limited by the availability of sulfur donors, especially thiosulfate, cystine and cysteine.

Thiosulfate is a normal constituent of serum, produced from cysteine by way of β -mercaptopyruvate. Physiological levels of thiosulfate are typically about 0.1 mmol/L (11 mg/L), but they are approximately twice this level in children and in adults who are not eating. Infused thiosulfate is cleared from the body (primarily by the kidneys) with a $t_{1/2}$ of about 20 minutes.

When thiosulfate is being supplied only by normal physiologic mechanisms, conversion of CN^- to SCN^- generally proceeds at about 1 mcg/kg/min. The rate of CN^- clearance corresponds to steady-state processing of a sodium nitroprusside infusion of slightly more than 2 mcg/kg/min. CN^- begins to accumulate when sodium nitroprusside infusions exceed this rate.

Thiocyanate (SCN^-) is also a normal physiological constituent of serum, with normal levels typically in the range of 50 to 250 $\mu\text{mol/L}$ (3 to 15 mg/L). Clearance of SCN^- is primarily renal, with a $t_{1/2}$ of about three days. In renal failure, the $t_{1/2}$ can be doubled or tripled.

Oral administration of sodium nitroprusside does not produce the dramatic decrease in blood pressure seen with intravenous administration. The effects of chronic oral administration are similar to those obtained with oral potassium thiocyanate.

In hypertensive patients, moderate depressor doses of sodium nitroprusside induce renal vasodilation roughly equivalent to the decrease in pressure, without an appreciable increase in renal blood flow or a decrease in glomerular filtration.

In normotensive subjects, acute reduction of mean arterial pressure to 60 -75 mm Hg by infusion of sodium nitroprusside caused significant increase in renin activity of renal venous plasma in correlation with a degree of reduction in pressure. Renal response to reduction in pressure was more striking in renovascular-hypertensive patients, with significant increase in renin release occurring from the involved kidney at mean arterial pressures ranging from 90 -137 mm Hg. Furthermore, the magnitude of renin release from the involved kidney was significantly greater when compared to that in normotensive subjects; while in the contralateral, uninvolved kidney, no significant release of renin was detected during the reduction of pressure.

Animal Pharmacology

Intravenous administration of sodium nitroprusside in doses of 0.125, 0.5 and 2.0 mg/kg/day (as 1 mL injections given in a two-minute period) was tolerated in dogs for a two-week period. The results noted were consistent with the transient vasodilation and lowering of blood pressure known to occur with intravenous sodium nitroprusside.

In dogs, the myocardial contractile force measured by a strain gauge arch showed no definite change with the administration of smaller doses of sodium nitroprusside (1 to 4 mcg/kg), while with high doses (8 to 16 mcg/kg) there was a decrease in the contractile force in about 60 to 90 seconds. The decrease was preceded by a fall in blood pressure occurring about 15 seconds post- infusion, suggesting that decrease in myocardial contractile force is not a primary effect of sodium nitroprusside but perhaps is due to the decrease in cardiac filling from relative peripheral pooling of blood. Adrenergic blocking agents failed to abolish its action, and autonomic ganglionic blocking agents markedly augmented the hypotensive response.

In dogs, intravenous infusion of sodium nitroprusside caused a slight increase in renal blood flow and decrease in vascular resistance, with no effect on the maximal tubular excretory capacity for PAH or the maximal tubular capacity for reabsorption of glucose.

TOXICOLOGY

The LD₅₀'s of intravenously administered sodium nitroprusside determined according to the method of Miller and Tainter (*Proc Soc Exp Biol Med*, 1944;57:261) in four different species of animals are as follows: mice, 8.4 ± 0.3 mg/kg; rats, 11.2 ± 1.1 mg/kg; rabbits, 2.8 ± 1.1 mg/kg; and dogs, 5 mg/kg (approximate).

Sodium nitroprusside was injected intravenously to 12 beagle dogs of both sexes over a period of two minutes, at doses of 0.125, 0.5 and 2.0 mg/kg/day for two weeks. All dogs survived the two weeks. After each injection, the dogs demonstrated signs consistent with the transient vasodilatation and lowering of blood pressure known to occur with sodium nitroprusside administered intravenously. Blood glucose was elevated temporarily after injections at the high and mid doses. Slight increases in the relative weights of the liver and adrenals at the high dose may have been related to vasodilation. Microscopic examination of the tissues showed congestion (vascular dilatation) of the liver, kidney and spleen at the high- and mid-doses, and of the liver at the low dose. There were no adverse cellular changes related to the vasodilation caused by sodium nitroprusside. No other changes were noted.

McDowall and co-workers (1974) have studied induced hypotension by intravenous infusion of sodium nitroprusside in baboons and reported that in four of eight animals receiving doses considered to be four to six times higher than "normal" in this species there was failure to recover normal arterial pressure after discontinuation of the infusion. These animals were in a state of metabolic acidosis and exhibited a markedly depressed cerebral oxygen uptake. In the four animals receiving doses considered "normal" for this species it was found that arterial pressure recovered rapidly. The early appearance of metabolic acidosis appeared to be coincident in animals resistant to nitroprusside induced hypotension.

Another study conducted in dogs (Stoyka and Schultz, 1975), reported that physiologic cerebral oxygenation was maintained at low perfusion pressures achieved with sodium nitroprusside infusion.

REFERENCES

Pharmacology and Toxicology

1. Bhatia SK, Frohlich ED. Hemodynamic comparison of agents useful in hypertensive emergencies. *Am Heart J* 1973;85:367-73.
2. Boxer GE, Rickards JC. Studies on the metabolism of the carbon of cyanide and thiocyanate. *Arch Biochem* 1952;39:7-26.
3. Goldstein F, Rieders F. Formation of cyanide in dog and man following administration of thiocyanate. *Am J Physiol* 1951;167:47-51.
4. Goldstein F, Rieders F. Conversion of thiocyanate to cyanide by an erythrocytic enzyme. *Am J Physiol* 1953;173:287-90.
5. Greiss L, Tremblay NAG, Davies DW. The toxicity of sodium nitroprusside. *Can Anaesth Soc J* 1976;23:480-5.
6. Hill HE. A contribution to the toxicology of sodium nitroprusside. I. The decomposition and determination of sodium nitroprusside. *Austr Chem Inst J Proc* 1942;9:89-93.
7. Johnson CC. The actions and toxicity of sodium nitroprusside. *Arch Int Pharmacodyn Ther* 1929;35:480-96.
8. McDowall DG, et al. The toxicity of sodium nitroprusside. *Br J Anaesth* 1974;46:327-32.
9. Michenfelder JD, Tinker JH. Cyanide toxicity and thiosulfate protection during chronic administration of sodium nitroprusside in the dog. *Anesthesiology* 1977;47:441-8.
10. Page IH. Treatment of essential and malignant hypertension. *JAMA* 1951;147:1311-8.
11. Page IH, et al. Cardiovascular actions of sodium nitroprusside in animals and hypertensive patients. *Circulation* 1955;11:188-98.
12. Pines KL, Crymble MM. In vitro conversion of thiocyanate to cyanide in the presence of erythrocytes. *Proc Soc Exp Biol Med* 1952;81:160-3.
13. Schlant RC, Tsagaris TS, Robertson RJ. Studies on the acute cardiovascular effects of intravenous sodium nitroprusside. *Am J Cardiol* 1962;9:51-9.
14. Stoyka WW, Schutz H. The cerebral response to sodium nitroprusside and trimethaphan controlled hypotension. *Can Anaesth Soc J* 1975;22:275-83.

Clinical

15. Amaranath L, Kellermeyer WF. Tachyphylaxis to sodium nitroprusside. *Anesthesiology* 1976;44:345-8.
16. Eppens H. Sodium nitroprusside in hypotensive anaesthesia. *Br J Anaesth* 1973;45:124.
17. Gifford RW, Jr. Current practices in general medicine. 7. Treatment of hypertensive emergencies including use of sodium nitroprusside. *Proc Staff Meet Mayo Clin* 1959;34:387-94.

18. Katz RL, Wolf CE. Sodium nitroprusside for controlled hypotension and hypertensive emergencies. In: Mark LC, Ngari SH, eds. *Highlights of clinical anesthesiology*. New York, Harper and Row, 1971:48-54.
19. Lowson JA. Sodium nitroprusside in hypotensive anaesthesia. *Br J Anaesth* 1972;44:908.
20. Nourok DS, et al. Hypothyroidism following prolonged sodium nitroprusside therapy. *Am J Med Sci* 1964;248:129-38.
21. Palmer RF, Lasseter KC. Sodium nitroprusside. *N Engl J Med* 1975;292:294-7.
22. Posner MA, Tobey RE, McElroy H. Hydroxocobalamin therapy of cyanide intoxication in guinea pigs. *Anesthesiology* 1976;44:157-60.
23. Posner MA, Rodkey FL, Tobey RE. Nitroprusside-induced cyanide poisoning: antidotal effect of hydroxocobalamin. *Anesthesiology* 1976;44:330-5.
24. Siegel P, et al. Sodium nitroprusside in the surgical treatment of cerebral aneurysms and arteriovenous malformations. *Br J Anaesth* 1971;43:790-5.
25. Styles M, Coleman AJ, Leary WP. Some hemodynamic effects of sodium nitroprusside. *Anesthesiology* 1973;38:173-6.
26. Tuzel IH. Sodium nitroprusside: a review of its clinical effectiveness as a hypotensive agent. *J Clin Pharmacol* 1974;14:494-503.
27. Wildsmith JAW, et al. Haemodynamic effects of sodium nitroprusside during nitrous oxide/halothane anaesthesia. *Br J Anaesth* 1973;45:71-4.

Overdosage

28. Lovejoy FH, Linden CH., Acute Poison and Drug Overdosage, *Harrison's Principles of Internal Medicine*, 12th ed., McGraw-Hill, 1991:2172.
29. Gerber JG, Nies AS. *Goodman and Gilman's The Pharmacological Basis of Therapeutics*. 8th ed., Pergamon Press, (New York), 1991:803-4.
30. PRODUCT MONOGRAPH ^{Pr}NIPRIDE[®] sodium nitroprusside injection
25 mg / mL sodium nitroprusside dihydrate Last revised: July 27, 2017 Control
Number.206712.

PART III: CONSUMER INFORMATION

^{Pr}Sodium Nitroprusside for Injection

This leaflet is part III of a three-part "Product Monograph" published when Sodium Nitroprusside for Injection was approved for sale in Canada and is designed specifically for Consumers. This leaflet is a summary and will not tell you everything about Sodium Nitroprusside for Injection. Contact your doctor or pharmacist if you have any questions about the drug.

ABOUT THIS MEDICATION

What the medication is used for:

- The immediate reduction of blood pressure of patients in hypertensive crises who have not responded adequately to standard treatment.
- Producing controlled hypotension (low blood pressure) in order to reduce bleeding during surgery when both the surgeon and anaesthesiologist decide it is necessary.

What it does:

Sodium Nitroprusside for Injection is a vasodilator that works by relaxing the muscles in your blood vessels to help them dilate (widen). This lowers blood pressure.

When it should not be used:

Sodium Nitroprusside for Injection should not be used if you:

- Are allergic to sodium nitroprusside;
- Are a physically poor risk patient being treated for compensatory hypertension (mechanism to correct high blood pressure);
- Have low brain circulation;
- Have anemia (a reduction of red blood cells)

If you are going to receive this drug in surgery, please tell your doctor if you:

- Have liver disease
- Have severe kidney disease
- Have hereditary vision loss (Leber's disease)
- Have vision problems caused by smoking
- Have a history of blood clot in your brain
- Have vitamin B₁₂ deficiency.

What the nonmedicinal ingredient is:

Water for Injection

What the medicinal ingredient is:

Sodium Nitroprusside dihydrate

What dosage forms it comes in:

Sodium Nitroprusside for Injection is a solution available in single-use amber glass vials (with latex-free stopper) in 50 mg / 2 mL format (25 mg / mL), containing sodium nitroprusside dihydrate, in water for injection.

WARNINGS AND PRECAUTIONS

Serious Warnings and Precautions

- Sodium Nitroprusside for Injection is only to be used as an intravenous infusion with 5% Dextrose. Not for direct injection.
- Sodium Nitroprusside for Injection can cause rapid decreases in blood pressure. In patients not properly monitored, these decreases can lead to irreversible injuries or death related to poor blood supply. Sodium Nitroprusside for Injection should be used only when available equipment and personnel allow blood pressure to be continuously monitored.
- Except when used briefly or at low infusion rates, Sodium Nitroprusside for Injection gives rise to important quantities of cyanide ion which can cause death.

BEFORE receiving Sodium Nitroprusside for Injection tell your doctor if:

- You have liver disease, kidney disease, anemia (a reduction of red blood cells), a seizure disorder, or a history of head injury or brain tumor, hypothyroidism (you do not produce enough thyroid hormone), difficulty breathing;
- You have an existing severe disease that may make you a poor candidate to undergo surgery;
- You are pregnant, plan to become pregnant or are breastfeeding;
- Geriatrics: you may experience a stronger lower blood pressure when you receive Sodium Nitroprusside for Injection, please tell your doctor your age.

INTERACTIONS WITH THIS MEDICATION

Drugs that may interact with Sodium Nitroprusside for Injection include: medications to help lower your blood pressure (specifically, hydralazine or hexamethonium).

PROPER USE OF THIS MEDICATION

Sodium Nitroprusside for Injection is only given in hospital with adequate equipment and trained personnel who can monitor you frequently for changes to your blood pressure. It is only given using an infusion pump, micro-drip regulator or any similar device that would allow precise measurement of the flow rate of the intravenous.

Sodium Nitroprusside for Injection is only to be used as an intravenous infusion with 5% Dextrose. Sodium Nitroprusside for Injection is not for direct injection.

The infusion fluid used for administration of Sodium Nitroprusside for Injection should not be employed as a vehicle for simultaneous administration of any other drug.

Usual dose:

Your health care professionals will decide the best dose for you. Sodium Nitroprusside for Injection must be diluted in 500 to 1000 ml of 5% Dextrose Injection. No other diluents should be used. The diluted solution should be protected from light, using the supplied opaque sleeve, aluminum foil or other opaque materials. The infusion solution should be freshly prepared and any unused portion discarded. The freshly prepared solution for infusion has a very faint brownish tint. If it is highly coloured, it should be discarded.

Storage period from the time of reconstitution to the completion of intravenous administration should not exceed 24 hours.

As with all parenteral drug products, intravenous admixtures should be inspected visually for clarity, particulate matter, precipitate, discoloration and leakage prior to administration whenever solution and container permit. Solutions showing haziness, particulate matter, precipitate, discoloration or leakage should not be used. Discard unused portion.

Tell your health care practitioner right away if you think that your intravenous has come out of the vein, or if it becomes sore, red or swollen.

Overdose:

If you think you have taken too much Sodium Nitroprusside for Injection, contact your healthcare professional, hospital emergency department or regional poison control centre immediately, even if there are no symptoms.

Seek emergency medical attention if you think you have received too much of this medicine. Symptoms of Sodium Nitroprusside for Injection overdose may include low blood pressure, shortness of breath, shallow breathing, nausea, vomiting, stomach pain, sweating, severe dizziness, headache, muscle twitching, fast or pounding heartbeat, restless feeling, loss of consciousness, and chest or back pain.

SIDE EFFECTS AND WHAT TO DO ABOUT THEM

Along with its needed effects, a medicine may cause some unwanted effects. These are referred to as “side effects.” Although not all of these side effects may occur, if they do occur they may need medical attention.

Nausea, retching, vomiting, sweating, apprehension, headache, restlessness, agitation, muscle twitching, abdominal pain, have been noted with too rapid reduction in blood pressure. However, these symptoms rapidly disappear with slowing of the rate of the infusion or temporary discontinuation of the infusion. Your doctor will decide the best rate of infusion for you.

Irritation of the injection site may occur.

SERIOUS SIDE EFFECTS AND WHAT TO DO ABOUT THEM				
Symptom / effect		Talk to your healthcare professional		Stop taking drug and get immediate medical help
		Only if severe	In all cases	
Common	Feeling dizzy and faint or your blood pressure is too low.		√	
	Difficulty breathing		√	
	Swelling in the arms or legs		√	
Uncommon	Feeling an irregular heart beat.		√	
	Rash or other skin irritation.		√	
	Weakness		√	
	Chest pain		√	
	Confusion and somnolence (a strong desire for sleep)		√	
	Methemoglobinemia, Symptoms include: Shortness of breath, blueness at mouth and finger tips, headache, fatigue, dizziness and loss of consciousness		√	

This is not a complete list of side effects. For any unexpected effects while taking Sodium Nitroprusside for Injection, contact your doctor or pharmacist.

HOW TO STORE IT

Sodium Nitroprusside for Injection should be stored at room temperature (15 to 30°C); protect from light and freezing.

Keep out of reach and sight of children

Following dilution in 5% Dextrose Injection, the diluted solution should be protected from light, using the supplied opaque sleeve, aluminum foil, or other opaque material. It is not necessary to cover the infusion drip chamber or the tubing.

The storage period from the time of reconstitution to the completion of intravenous administration should not exceed 24 hours.

REPORTING SIDE EFFECTS

You can report any suspected side effects associated with the use of health products to Health Canada by:

- Visiting the Web page on Adverse Reaction Reporting (<https://www.canada.ca/en/health-canada/services/drugs-health-products/medeffect-canada/adverse-reaction-reporting.html>) for information on how to report online, by mail or by fax; or
- Calling toll-free at 1-866-234-2345.

NOTE: Contact your health professional if you need information about how to manage your side effects. The Canada Vigilance Program does not provide medical advice.

MORE INFORMATION

If you want more information about Sodium Nitroprusside for Injection:

- Talk to your healthcare professional
- Find the full product monograph that is prepared for healthcare professionals and includes this Consumer Information by visiting the Health Canada website (<https://health-products.canada.ca/dpd-bdpp/index-eng.jsp>); the manufacturer's website www.drreddys.com, or by calling 1-855-845-1739

This leaflet was prepared by

Dr. Reddy's Laboratories Limited

Bachupally – 500090 India

Imported and Distributed by:

Dr. Reddy's Laboratories Canada Inc.

Mississauga, ON L4W 4Y1

Canada

Date Prepared: Oct 29, 2020