

# PRODUCT MONOGRAPH

## **Pr TAMSULOSIN CR**

Tamsulosin Hydrochloride Controlled-Release Tablets

0.4 mg Tablet

Manufacturer Standard

SELECTIVE ANTAGONIST OF  
ALPHA<sub>1A/1D</sub> ADRENORECEPTOR SUBTYPES  
IN THE PROSTATE AND BLADDER

**PRO DOC LTÉE**  
2925, boul. Industriel  
Laval, Québec  
H7L 3W9

Date of Revision:  
January 21, 2021

Submission Control No: 247291

**Table of Contents**

**PART I: HEALTH PROFESSIONAL INFORMATION..... 3**

SUMMARY PRODUCT INFORMATION ..... 3

INDICATIONS AND CLINICAL USE ..... 3

CONTRAINDICATIONS ..... 3

WARNINGS AND PRECAUTIONS ..... 4

ADVERSE REACTIONS..... 7

DRUG INTERACTIONS ..... 11

DOSAGE AND ADMINISTRATION ..... 13

OVERDOSAGE..... 13

ACTION AND CLINICAL PHARMACOLOGY ..... 14

STORAGE AND STABILITY ..... 18

DOSAGE FORMS, COMPOSITION AND PACKAGING ..... 18

**PART II: SCIENTIFIC INFORMATION ..... 19**

PHARMACEUTICAL INFORMATION..... 19

CLINICAL TRIALS ..... 20

DETAILED PHARMACOLOGY ..... 25

TOXICOLOGY ..... 26

REFERENCES..... 27

**PART III: CONSUMER INFORMATION..... 29**

**PrTAMSULOSIN CR**  
Tamsulosin Hydrochloride Controlled-Release Tablets

Manufacturer Standard

**PART I: HEALTH PROFESSIONAL INFORMATION**

**SUMMARY PRODUCT INFORMATION**

<b>Route of Administration</b>	<b>Dosage Form / Strength</b>	<b>All Nonmedicinal Ingredients</b>
Oral	Controlled release tablet 0.4 mg	Citric acid, colloidal silicon dioxide, hydroxypropyl cellulose, hydroxypropyl methylcellulose 2208 and 2910, magnesium stearate, polyethylene glycol, sodium alginate and yellow ferric oxide.

**INDICATIONS AND CLINICAL USE**

TAMSULOSIN CR (tamsulosin hydrochloride) is indicated for the treatment of Lower Urinary Tract Symptoms (LUTS) associated with benign prostatic hyperplasia (BPH).

**Geriatrics (> 65 years of age):**

Tamsulosin hydrochloride has been found to be a safe and effective alpha<sub>1</sub> adrenoceptor antagonist when administered at therapeutic doses (0.4 mg once daily) to patients over the age of 65 years.

**Pediatrics:**

TAMSULOSIN CR is not indicated for use in children.

The effectiveness of tamsulosin in 161 pediatric patients (ages 2 to 16 years) with neuropathic bladder was not demonstrated (see WARNINGS AND PRECAUTIONS, Special Populations, Pediatrics).

**CONTRAINDICATIONS**

- TAMSULOSIN CR (tamsulosin hydrochloride) is contraindicated in patients known to have hypersensitivity including drug induced angioedema to tamsulosin or any component of the TAMSULOSIN CR controlled release formulation. For a complete listing, see the DOSAGE FORMS, COMPOSITION AND PACKAGING section of the Product Monograph.
- TAMSULOSIN CR (tamsulosin hydrochloride) should not be administered to patients using concomitant strong CYP3A4 inhibitors (e.g. ketoconazole) (see section DRUG INTERACTIONS).

## **WARNINGS AND PRECAUTIONS**

**As with all  $\alpha_1$ -adrenoceptor antagonists, a reduction in blood pressure can occur in individual cases during treatment with tamsulosin hydrochloride, as a result of which, rarely, syncope can occur. At the first signs of orthostatic hypotension (dizziness, weakness), the patient should sit or lie down until the symptoms have disappeared.**

**Patients beginning treatment with tamsulosin hydrochloride should be cautioned to avoid situations where injury could result should syncope occur (see ADVERSE REACTIONS).**

### **General**

Tamsulosin hydrochloride is not indicated for the treatment of hypertension.

### **Drug-Drug Interactions**

- Tamsulosin is extensively metabolized, mainly by CYP3A4 and CYP2D6. TAMSULOSIN CR should not be used in combination with strong inhibitors of CYP3A4 (e.g., ketoconazole). TAMSULOSIN CR should be used with caution in combination with moderate inhibitors of CYP3A4 (e.g., erythromycin), in combination with strong (e.g., paroxetine) or moderate (e.g., terbinafine) inhibitors of CYP2D6, in patients known to be CYP2D6 poor metabolizers.
- TAMSULOSIN CR should be used with caution in combination with cimetidine.
- TAMSULOSIN CR should not be used in combination with other alpha adrenergic blocking agents.
- Caution is advised when alpha adrenergic blocking agents including TAMSULOSIN CR are coadministered with PDE5 inhibitors. Alpha-adrenergic blockers and PDE5 inhibitors are both vasodilators that can lower blood pressure. Concomitant use of these two drug classes can potentially cause symptomatic hypotension.
- Caution should be exercised with concomitant administration of warfarin and TAMSULOSIN CR

See DRUG INTERACTIONS

### **Carcinoma of the Prostate**

Carcinoma of the prostate and BPH cause many of the same symptoms. These two diseases frequently co-exist. Patients should be evaluated to rule out the presence of carcinoma of the prostate.

### **Orthostatic Hypotension**

While syncope is the most severe orthostatic symptom of  $\alpha_1$ -adrenoceptor antagonists, other symptoms can occur (dizziness and postural hypotension). In a phase III, randomized, double-blind, placebo-controlled trial involving male patients treated once daily with either 0.4 mg tamsulosin hydrochloride (n=350) or placebo (n=356), both supine and standing blood pressure were monitored over the course of the 12 week treatment period. There was a small, clinically insignificant decrease from baseline in mean supine and standing systolic/diastolic BP in both treatment groups; the decrease in BP from baseline in the tamsulosin hydrochloride group (< 2 mmHg) was comparable to the placebo group

(< 1.5 mmHg). There were no cases of orthostatic hypotension or syncope reported in either treatment group.

Patients in occupations in which orthostatic hypotension could be dangerous should be treated with caution.

If hypotension occurs, the patient should be placed in the supine position and if this measure is inadequate, volume expansion with intravenous fluids or vasopressor therapy may be used. A transient hypotensive response is not a contraindication to further therapy with tamsulosin hydrochloride.

### **Hepatic**

The treatment of patients with severe hepatic impairment should be approached with caution as no studies have been conducted in this patient population. No dose adjustment is warranted in hepatic insufficiency.

### **Renal**

The treatment of patients with severe renal impairment (creatinine clearance of <10mL/min) should be approached with caution, as these patients have not been studied.

### **Intraoperative Floppy Iris Syndrome**

During cataract and/or glaucoma surgery, a variant of small pupil syndrome known as Intraoperative Floppy Iris Syndrome (IFIS) has been reported during post-marketing surveillance in association with alpha-1 blocker therapy, including tamsulosin hydrochloride. Most reports to date were in patients taking tamsulosin hydrochloride when IFIS occurred, but in some cases, tamsulosin hydrochloride had been stopped prior to surgery. In most of these cases, tamsulosin hydrochloride had been stopped recently prior to surgery (2 to 14 days), but in a few cases, IFIS was reported after the patient had been off tamsulosin hydrochloride for a longer period. This variant of small pupil syndrome is characterized by the combination of a flaccid iris that billows in response to intraoperative irrigation currents, progressive intraoperative miosis despite preoperative dilation with standard mydriatic drugs and potential prolapse of the iris toward the phacoemulsification incisions. The patient's ophthalmologist should be prepared for possible modifications to their surgical technique, such as the utilization of iris hooks, iris dilator rings, or viscoelastic substances. IFIS may increase the risk of eye complications during and after the operation. The benefit of stopping alpha-1 blocker therapy, including tamsulosin hydrochloride prior to cataract and/or glaucoma surgery has not been established. IFIS has also been reported in patients who had discontinued tamsulosin for a longer than 2 week period prior to the surgery. The initiation of therapy with tamsulosin hydrochloride in patients for whom cataract and/or glaucoma surgery is scheduled is not recommended.

### **Reproduction**

Ejaculation disorders have been observed in short and long term clinical studies with tamsulosin (see ADVERSE REACTIONS, Clinical Trial Adverse Drug Reactions). Events of ejaculation disorder, retrograde ejaculation and ejaculation failure have been reported in post marketing.

### **Sulfa Allergy**

In patients with sulfa allergy, allergic reaction to tamsulosin hydrochloride capsules has been rarely reported. If a patient reports a serious or life-threatening sulfa allergy, caution is warranted when administering tamsulosin hydrochloride.

### **Special Populations**

**Pregnant Women:** Tamsulosin hydrochloride is not indicated for use in women. Studies in pregnant rats and rabbits at daily doses of 300 and 50 mg/kg, respectively (30,000 and 5,000 times the anticipated human dose), revealed no evidence of harm to the fetus. There are no adequate data on the use of tamsulosin in pregnant women; therefore the potential risk from the use of tamsulosin during pregnancy in humans is unknown.

**Nursing Women:** Tamsulosin hydrochloride is not indicated for use in women.

**Pediatrics:** Tamsulosin hydrochloride is not indicated for use in children. Tamsulosin hydrochloride has been studied in 161 pediatric patients (ages 2 to 16 years) with an elevated detrusor leak point pressure associated with a known neurological disorder (e.g., spina bifida). The effectiveness of tamsulosin in this pediatric population was not demonstrated. The most frequently reported adverse events ( $\geq 5\%$ ) were urinary tract infection, vomiting, nasopharyngitis, influenza, headache, and abdominal pain.

**Geriatrics (> 65 years of age):** There were no pharmacokinetic studies conducted in geriatric patients with tamsulosin hydrochloride controlled-release tablets. Cross-study comparisons of overall exposure (AUC) and half-life of tamsulosin hydrochloride capsules indicate that the pharmacokinetic disposition of tamsulosin may be slightly prolonged in geriatric males compared to young healthy male volunteers. However, tamsulosin hydrochloride capsules have been found to be a safe and effective  $\alpha_1$  adrenoreceptor antagonist when administered at therapeutic doses to patients over the age of 65 years.

**Gender Effects:** Tamsulosin hydrochloride is not indicated for use in women. Safety, effectiveness, and pharmacokinetics have not been evaluated in women.

### **Monitoring and Laboratory Tests**

No laboratory test interactions with tamsulosin hydrochloride are known. Treatment with tamsulosin hydrochloride for up to 3 months had no significant effect on prostate specific antigen (PSA).

### **Information for the patient (See PART III: CONSUMER INFORMATION)**

Patients should be advised not to crush or chew tamsulosin hydrochloride controlled-release tablets. These tablets are specially formulated to control the delivery of tamsulosin HCl to the blood stream.

There are no specific studies conducted with tamsulosin hydrochloride and the ability to drive vehicles or use machinery. However patients should be advised that dizziness can occur with tamsulosin hydrochloride, requiring caution in people who must drive, operate machinery, or perform hazardous tasks.

Patients should be advised about the possibility of priapism as a result of treatment with tamsulosin hydrochloride and other similar medications. Patients should be informed that this reaction is extremely rare, but if not brought to immediate medical attention, can lead to permanent erectile dysfunction (impotence).

## **ADVERSE REACTIONS**

### **Adverse Drug Reaction Overview**

Information on the safety profile of tamsulosin hydrochloride was derived from two, 3-month placebo-controlled clinical trials involving 1840 male subjects. Of these, 563 were treated with tamsulosin hydrochloride controlled-release tablets 0.4 mg, 709 with tamsulosin hydrochloride capsules 0.4 mg and 568 with placebo. The results suggest that tamsulosin hydrochloride controlled-release tablets 0.4 mg and tamsulosin hydrochloride capsules 0.4 mg were very well tolerated with the AE profile of tamsulosin hydrochloride controlled-release tablets 0.4 mg tending to be more favourable than that of tamsulosin hydrochloride capsules.

In these studies, 3.6% of patients taking tamsulosin hydrochloride controlled-release tablets (0.4 mg) discontinued from the study due to adverse events compared with 1.2% in the placebo group. The most frequently reported Treatment Emergent Adverse Events (TEAE) in the tamsulosin hydrochloride controlled-release tablets 0.4 mg group was dizziness and those related to abnormal ejaculation, although the incidence of both was comparable to placebo.

Impotence and other events related to sexual function are commonly associated with other alpha<sub>1</sub>-blockers, however in the 3-month studies with tamsulosin hydrochloride controlled-release tablets there were minimal effects on sexual function and ejaculatory disorders/abnormalities with no reports of priapism. The difference in incidence of ejaculatory disorders/abnormalities between tamsulosin hydrochloride controlled-release tablets and placebo was not statistically significant. No patient discontinued treatment with tamsulosin hydrochloride controlled-release tablets 0.4 mg due to ejaculatory disorders/abnormalities.

### **Clinical Trial Adverse Drug Reactions**

*Because clinical trials are conducted under very specific conditions the adverse reaction rates observed in the clinical trials may not reflect the rates observed in practice and should not be compared to the rates in the clinical trials of another drug. Adverse drug reaction information from clinical trials is useful for identifying drug-related adverse events and for approximating rates.*

**TABLE 1: TREATMENT-EMERGENT ADVERSE EVENTS IN  $\geq$  2% OF PATIENTS RECEIVING EITHER TAMSULOSIN OR PLACEBO DURING THE 3 MONTH PLACEBO AND ACTIVE-CONTROLLED STUDY.**

<b>SOC/Preferred term</b>	<b>Placebo N=356</b>	<b>Tamsulosin Hydrochloride Controlled-Release Tablets 0.4 mg N=360</b>	<b>Tamsulosin Capsules 0.4 mg N=709</b>
Any TEAE	71 (19.9%)	93 (25.8%)	168 (23.7%)
Cardiac disorders	8 (2.2%)	8 (2.2%)	16 (2.3%)
Gastrointestinal disorders	7 (2.0%)	14 (3.9%)	34 (4.8%)
General Disorders and administration site conditions	2 (0.6%)	8 (2.2%)	11 (1.6%)
Infections and infestations	16 (4.5%)	20 (5.6%)	32 (4.5%)
Investigations	10 (2.8%)	6 (1.7%)	10 (1.4%)
Musculoskeletal and connective tissue disorders	7 (2.0%)	9 (2.5%)	12* (1.7%)
Nervous system disorders	9 (2.5%)	11 (3.1%)	29 (4.1%)
Reproductive system and breast disorders	2 (0.6%)	12 (3.3%)	28 (3.9%)
Respiratory, thoracic and mediastinal disorders	3 (0.8%)	10 (2.8%)	20 (2.8%)
Vascular disorders	8 (2.2%)	6 <sup>#</sup> (1.7%)	15 (2.1%)

Number (%) of patients

A patient may experience an AE more than once or may experience more than one AE within the same SOC.

\*Post database lock: deletion of 1 AE

<sup>#</sup>Post database lock: addition of 1 AE

**TABLE 2: NUMBER (%) OF PATIENTS WITH TEAES COMMONLY ASSOCIATED WITH A<sub>1</sub>-AR ANTAGONISTS DURING THE 3 MONTH PLACEBO AND ACTIVE-CONTROLLED STUDY.**

SOC/Preferred term	Placebo N=356	Tamsulosin Hydrochloride Controlled-Release Tablets 0.4 mg N=360	Tamsulosin Capsules 0.4 mg N=709
<b>Non-cardiovascular class effects</b>			
Retrograde ejaculation	1 (0.3%)	6 (1.7%)	10 (1.4%)
Ejaculation Failure	0 (0.0%)	0 (0.0%)	2 (0.3%)
Semen volume reduced	0 (0.0%)	1 (0.3%)	2 (0.3%)
Ejaculation delayed	0 (0.0%)	1 (0.3%)	2 (0.3%)
Ejaculation disorder NOS	0 (0.0%)	0 (0.0%)	6 (0.8%)
ABNORMAL EJACULATION POOLED	1 (0.3%)	7 (1.9%)	22 (3.1%)
Headache NOS	4 (1.1%)	3 (0.8%)	10 (1.4%)
Asthenia	1 (0.3%)	1 (0.3%)	1 (0.1%)
Fatigue	1 (0.3%)	3 (0.8%)	2 (0.3%)
Somnolence	0 (0.0%)	0 (0.0%)	2 (0.3%)
Rhinitis NOS	0 (0.0%)	1 (0.3%)	2 (0.3%)
Nasal congestion	0 (0.0%)	1 (0.3%)	1 (0.1%)
Nasal obstruction	0 (0.0%)	0 (0.0%)	0 (0.0%)
SUB-TOTAL	7 (2.0%)	16 (4.4%)	36 (5.1%)
<b>Cardiovascular class effects</b>			
Dizziness	5 (1.4%)	5 (1.4%)	9 (1.3%)
Dizziness aggravated	0 (0.0%)	0 (0.0%)	2 (0.3%)
Dizzy spell	0 (0.0%)	0 (0.0%)	1 (0.1%)
DIZZINESS POOLED	5 (1.4%)	5 (1.4%)	12 (1.7%)
Palpitations	2 (0.6%)	2 (0.6%)	1 (0.1%)
Tachycardia NOS	0 (0.0%)	1 (0.3%)	2 (0.3%)
Hypotension NOS	1 (0.3%)	0 (0.0%)	2 (0.3%)
Orthostatic hypotension	0 (0.0%)	0 (0.0%)	3 (0.4%)
Dizziness postural	0 (0.0%)	0 (0.0%)	2 (0.3%)
Syncope	0 (0.0%)	0 (0.0%)	1 (0.1%)
Orthostatic/circulatory collapse	0 (0.0%)	0 (0.0%)	0 (0.0%)
Depressed level of/loss of consciousness	0 (0.0%)	1 (0.3%)	1 (0.1%)
SUB-TOTAL	8 (2.2%)	9 (2.5%)	23 (3.2%)
TOTAL	13 (3.7%)	25 (6.9%)	55 (7.8%)

A patient may experience an AE more than once or may experience more than one AE within the same SOC.

Angioedema or priapism was not reported in the phase 2 or 3 studies.

### **Post-Market Adverse Drug Reactions**

The following adverse reactions have been reported during the use of tamsulosin hydrochloride at a frequency of:

#### >1% AND < 10%:

Nervous System Disorders: dizziness

Reproductive system and breast disorders: ejaculation disorders including retrograde ejaculation and ejaculation failure

#### > 0.1% AND < 1%:

Cardiac disorders: palpitations

Gastrointestinal Disorders: constipation, diarrhea, nausea, and vomiting

General disorders and administration site conditions: asthenia

Nervous systems disorders: headache

Respiratory, thoracic and mediastinal disorders: rhinitis

Skin and subcutaneous tissue disorders: rash, pruritus, urticaria

Vascular disorders: Orthostatic hypotension

#### > 0.01% AND < 0.1%:

Nervous system disorders: syncope

Skin and subcutaneous tissue disorders: angioedema

#### < 0.01%:

Reproductive systems and breast disorders: priapism

Skin and subcutaneous tissue disorders: Stevens-Johnson syndrome

### **Not Known (cannot be estimated from the available data)**

Eye disorders: vision blurred, visual impairment

Respiratory, thoracic and mediastinal disorders: epistaxis

Skin and subcutaneous tissue disorders: erythema multiforme, dermatitis exfoliative, tamsulosin-induced photosensitivity reaction

Gastrointestinal Disorders: dry mouth

In addition to the adverse events listed above, atrial fibrillation, arrhythmia, tachycardia and dyspnoea have been reported in association with tamsulosin use. Because these spontaneously reported events are from the worldwide post marketing experience, the frequency of events and the role of tamsulosin in their causation cannot be reliably determined.

During cataract and glaucoma surgery, a variant of small pupil syndrome known as Intraoperative Floppy Iris Syndrome (IFIS) has been reported during post-marketing surveillance in association with alpha-1 blocker therapy, including tamsulosin hydrochloride (see **WARNINGS AND PRECAUTIONS**).

An open label extension study involving 609 male patients with lower urinary tract symptoms (LUTS) associated with BPH demonstrated sustained efficacy, safety and long-term tolerability of tamsulosin for up to 6 years.

## **DRUG INTERACTIONS**

### **Overview**

There were no drug interaction studies conducted specifically with tamsulosin hydrochloride controlled-release tablets and it is expected that the interaction profile would not be any different than that of tamsulosin hydrochloride capsules. As with tamsulosin hydrochloride capsules, caution should be exercised with concomitant administration of tamsulosin hydrochloride controlled-release tablets and other alpha-adrenergic blocking agents.

No clinically significant drug-drug interactions were observed when tamsulosin hydrochloride capsules 0.4 mg or 0.8 mg were administered with one of the following therapeutic agents: nifedipine, atenolol, enalapril, digoxin, furosemide or theophylline.

### **Drug-Drug Interactions**

#### **Strong and Moderate Inhibitors of CYP3A4 or CYP2D6**

Tamsulosin is extensively metabolized, mainly by CYP3A4 and CYP2D6.

The effects of ketoconazole (a strong inhibitor of CYP3A4) at 400 mg once daily for 5 days on the pharmacokinetics of a single tamsulosin capsule 0.4 mg dose was investigated in 24 healthy volunteers (age range 23 to 47 years). Concomitant treatment with ketoconazole resulted in an increase in the C<sub>max</sub> and AUC of tamsulosin by a factor of 2.2 and 2.8, respectively. The effects of concomitant administration of a moderate CYP3A4 inhibitor (e.g., erythromycin) on the pharmacokinetics of tamsulosin have not been evaluated.

The effects of paroxetine (a strong inhibitor of CYP2D6) at 20 mg once daily for 9 days on the pharmacokinetics of a single tamsulosin capsule 0.4 mg dose was investigated in 24 healthy volunteers (age range 23 to 47 years). Concomitant treatment with paroxetine resulted in an increase in the C<sub>max</sub> and AUC of tamsulosin by a factor of 1.3 and 1.6, respectively. A similar increase in exposure is expected in CYP2D6 poor metabolizers (PM) as compared to extensive metabolizers (EM). A fraction of the population (about 7% of Caucasians and 2% of African Americans) is CYP2D6 PMs. Since CYP2D6 PMs cannot be readily identified and the potential for significant increase in tamsulosin exposure exists when tamsulosin is co-administered with strong CYP3A4 inhibitors in CYP2D6 PMs, tamsulosin should not be used in combination with strong inhibitors of CYP3A4 (e.g., ketoconazole). Tamsulosin should be given with caution in combination with moderate inhibitors of CYP3A4.

The effects of concomitant administration of a moderate CYP2D6 inhibitor (e.g., terbinafine) on the pharmacokinetics of tamsulosin have not been evaluated.

The effects of co-administration of both a CYP3A4 and a CYP2D6 inhibitor with tamsulosin have not been evaluated. However, there is a potential for significant increase in tamsulosin exposure when tamsulosin hydrochloride is co-administered with a combination of both CYP3A4 and CYP2D6 inhibitors.

**Nifedipine, Atenolol, Enalapril:** No dosage adjustments are necessary when tamsulosin hydrochloride controlled release tablets are administered concomitantly with nifedipine extended release tablets, atenolol, or enalapril. In three studies in hypertensive subjects (age range 47 to 79 years) whose blood pressure was controlled with stable doses of nifedipine extended release tablets, atenolol or enalapril for at least three months, tamsulosin hydrochloride 0.4 mg capsules for seven days followed by tamsulosin hydrochloride 0.8 mg capsules for another seven days (n=8 per study) resulted in no clinically significant effects on blood pressure and pulse rate compared to placebo (n=4 per study).

**Warfarin:** A definitive drug-drug interaction study between tamsulosin and warfarin was not conducted. Results from limited in-vitro and in-vivo studies are inconclusive. Therefore, caution should be exercised with concomitant administration of warfarin and tamsulosin hydrochloride.

**Digoxin and Theophylline:** No dosage adjustments are necessary when tamsulosin hydrochloride is administered concomitantly with digoxin or theophylline. In two studies in healthy volunteers (n=10 per study; age range 19 to 39 years), receiving tamsulosin hydrochloride capsules 0.4 mg/day for two days, followed by tamsulosin hydrochloride capsules 0.8 mg/day for five to eight days, single intravenous doses of digoxin 0.5 mg or theophylline 5 mg/kg resulted in no change in the pharmacokinetics of digoxin or theophylline.

**Furosemide:** No dosage adjustments are necessary when tamsulosin hydrochloride is administered concomitantly with furosemide. The pharmacokinetic and pharmacodynamic interaction between tamsulosin hydrochloride capsules 0.8 mg/day (steady-state) and furosemide 20 mg intravenously (single dose) was evaluated in ten healthy volunteers (age range 21 to 40 years). Tamsulosin hydrochloride capsules had no effect on the pharmacodynamics (excretion of electrolytes) of furosemide. While furosemide produced an 11% to 12% reduction in tamsulosin  $C_{max}$  and AUC, these changes are expected to be clinically insignificant and do not require adjustment of the tamsulosin hydrochloride controlled-release tablets dosage.

**Cimetidine:** The effects of cimetidine at the highest recommended dose (400 mg every six hours for six days) on the pharmacokinetics of a single tamsulosin hydrochloride capsules 0.4 mg dose was investigated in ten healthy volunteers (age range 21 to 38 years). Treatment with cimetidine resulted in a moderate increase in tamsulosin AUC (44%) due to a significant decrease (26%) in the clearance of tamsulosin. Therefore, tamsulosin hydrochloride controlled-release tablets should be used with caution in combination with cimetidine.

## **PDE5 Inhibitors**

Alpha-adrenergic blockers and PDE5 inhibitors are both vasodilators that can lower blood pressure. Concomitant use of these two drug classes can potentially cause symptomatic hypotension. Therefore, caution is advised when alpha adrenergic blocking agents including tamsulosin are co-administered with PDE5 inhibitors.

## **Other Alpha Adrenergic Blocking Agents**

The pharmacokinetic and pharmacodynamic interactions between tamsulosine and other alpha adrenergic blocking agents have not been determined; however, interactions between tamsulosine and other alpha adrenergic blocking agents may be expected.

## **Drug-Laboratory Test Interactions**

No laboratory test interactions with tamsulosin hydrochloride are known. Treatment with tamsulosin hydrochloride for up to 3 months had no significant effect on prostate specific antigen (PSA).

## **DOSAGE AND ADMINISTRATION**

### **Dosing Considerations**

TAMSULOSIN CR (tamsulosin hydrochloride) 0.4 mg once daily is recommended as the dose for the treatment of lower urinary tract symptoms (LUTS) associated with Benign Prostatic Hyperplasia (BPH).

### **Missed Dose**

If a dose of TAMSULOSIN CR is missed, the missed dose can be taken later the same day. If a day is missed, the missed dose should be skipped and the regular dosing schedule should be resumed. Doses must not be doubled.

### **Administration**

TAMSULOSIN CR should be taken at the same time each day with or without food. TAMSULOSIN CR tablets **must be swallowed whole, as crushing or chewing will interfere with the controlled release of the active ingredient.**

Taking TAMSULOSIN CR with a high fat meal increase exposure to tamsulosin (see ACTION AND CLINICAL PHARMACOLOGY – Pharmacokinetics section)

## **OVERDOSAGE**

Overdosage with tamsulosin hydrochloride can potentially result in severe hypotensive effects. Severe hypotensive effects have been observed at different levels of overdosage.

Should overdosage of tamsulosin hydrochloride lead to hypotensive effects (see **WARNINGS AND PRECAUTIONS**), support of the cardiovascular system is of first importance. Restoration of blood pressure and normalization of heart rate may be accomplished by keeping the patient in the supine position. If this measure is inadequate, then administration of intravenous fluids should be considered. If necessary, vasopressors should then be used and renal function should be monitored and supported as

needed. Laboratory data indicate that tamsulosin is 94% to 99% protein bound: therefore dialysis is unlikely to be of benefit.

Measures such as emesis can be taken to impede absorption. When large quantities are involved, gastric lavage can be applied and activated charcoal and an osmotic laxative, such as sodium sulphate can be administered.

Acute overdose with 5 mg of tamsulosin hydrochloride has been reported. Acute hypotension (systolic blood pressure 70 mm Hg), vomiting and diarrhoea were observed, which were treated with fluid replacement and the patient could be discharged the same day. One patient reported an overdose of 30 X 0.4 mg tamsulosin hydrochloride capsules. Following the ingestion of the capsules, the patient reported a headache judged to be severe and probably drug-related that resolved the same day.

For management of a suspected drug overdose, contact your regional Poison Control Centre.
---

## **ACTION AND CLINICAL PHARMACOLOGY**

### **Mechanism of Action**

Tamsulosin hydrochloride is an  $\alpha_1$  adrenoreceptor (AR) blocking agent used for the treatment of lower urinary tract symptoms (LUTS) associated with benign prostatic hyperplasia (BPH). It exhibits selectivity for both  $\alpha_{1A}$  and  $\alpha_{1D}$  receptors over the  $\alpha_{1B}$  AR subtype. These three AR subtypes have a distinct distribution pattern in human tissue. Whereas approximately 70% of the  $\alpha_1$ -receptors in human prostate are of the  $\alpha_{1A}$  subtype, the human bladder contains predominantly the  $\alpha_{1D}$  subtype while blood vessels express predominantly  $\alpha_{1B}$  subtype.

Stimulation/antagonism of each of the receptor subtypes gives rise to a distinct pharmacological effect. Lower Urinary Tract Symptoms (LUTS) suggestive of benign prostatic obstruction (BPO) formerly referred to as symptomatic benign prostatic hyperplasia (BPH) are very common in men > 50 years old; the prevalence increases with age. The symptoms associated with LUTS/BPH are comprised of two underlying components: the static and dynamic. The static component is related to an increase in prostate size caused, in part, by a proliferation of smooth muscle cells in the prostatic stroma. However, the severity of BPH symptoms and the degree of urethral obstruction do not correlate well with the size of the prostate. The dynamic component is a function of an increase in smooth muscle tone in the prostate and bladder neck leading to constriction of the bladder outlet. Smooth muscle tone is mediated by the sympathetic nervous stimulation of  $\alpha_1$  adrenoreceptors, which are abundant in the prostate, prostatic capsule, prostatic urethra, and bladder neck. Blockade of these adrenoreceptors can cause smooth muscles in the bladder neck and prostate to relax, resulting in an improvement in urine flow rate and a reduction in symptoms of BPH.

It is further believed that blockade of  $\alpha_{1D}$  subtypes in the human obstructed bladder may be responsible for reducing detrusor overactivity and subsequent relief of storage symptoms.

Tamsulosin hydrochloride is not intended for use as an antihypertensive drug.

## Pharmacodynamics

The pharmacokinetics of tamsulosin have been evaluated in adult healthy volunteers with doses ranging from 0.4 mg to 1.6 mg.

## Pharmacokinetics

Absorption: After a single oral dose of 0.4 mg of tamsulosin hydrochloride controlled-release tablets in the fasted state, the plasma concentration of tamsulosin gradually increased reaching  $C_{max}$  at a median time of 6 hours. At steady state, which is reached by day 4 of multiple dosing, plasma concentrations of tamsulosin peak at 4 to 6 hours in the fasted and fed state. Peak plasma concentrations increase from approximately 6 ng/mL after the first dose to 11 ng/mL in steady state. After  $C_{max}$  is reached, the plasma concentration decreases, but at approximately 16 to 24 hours post-dose, a small increase or second plateau is observed. Under fasted conditions the absolute bioavailability of tamsulosin from tamsulosin hydrochloride controlled-release tablets was estimated to be 57%.


A study conducted at steady state with 0.4 mg tamsulosin hydrochloride controlled-release tablets demonstrated that the plasma concentration-time profile in the fed state was bioequivalent to the fasted state, indicating the absence of a food effect, by a low fat meal (Table 3). After a single oral dose of 0.4mg tamsulosin hydrochloride controlled-release tablets, the extent of absorption is increased by 64% and 149% (AUC and  $C_{max}$  respectively) by a high-fat meal compared to fasted.

**Table 3: MEAN PHARMACOKINETIC PARAMETERS OF TAMSULOSIN AT STEADY STATE FOLLOWING ADMINISTRATION OF ONCE DAILY DOSES OF 0.4 MG TAMSULOSIN HYDROCHLORIDE CONTROLLED-RELEASE TABLETS IN BOTH THE FED AND FASTED STATE.**

Parameter	Tamsulosin Hydrochloride Controlled-Release Tablets 0.4 mg (Fed) (n=24)	Tamsulosin Hydrochloride Controlled-Release Tablets 0.4 mg (Fasted) (n=24)
AUC <sub>0-inf</sub> (ng·h/mL)	291.1	278.7
$C_{max}$ (ng/mL)	11.1	10.7
$C_{24}$ (ng/mL)	4.8	4.6
$T_{max}$ (h)	4.16	4.75
$T_{1/2}$ (h)	14.6	15.6

The 0.4 mg tamsulosin hydrochloride controlled-release tablet is not bioequivalent to the 0.4 mg tamsulosin hydrochloride capsule, as the test/reference ratio for  $C_{max}$  and AUC did not fall within the predefined limits of 80 to 125%. The plasma concentration-time profile presented in Figure 1. shows the lack of a pronounced spike in  $C_{max}$  with tamsulosin hydrochloride controlled-release tablets compared with capsules which may be consistent with a more favourable safety profile.

**FIGURE 1: MEAN TAMSULOSIN PLASMA VS. TIME PROFILES OF TAMSULOSIN HYDROCHLORIDE CONTROLLED-RELEASE TABLETS 0.4 MG AND TAMSULOSIN CAPSULES, 0.4 MG (N=12)**



**Distribution:** The mean steady-state apparent volume of distribution of tamsulosin after intravenous administration to ten healthy male adults was 16 litres, which is suggestive of distribution into extracellular fluids in the body. Additionally, whole body autoradiographic studies in mice, rats and dogs indicate that tamsulosin is widely distributed to most tissues including kidney, prostate, liver, gall bladder, heart, aorta, and brown fat, and minimally distributed to the brain, spinal cord, and testes.

Tamsulosin is extensively bound to human plasma proteins (94% to 99%), primarily alpha-1-acid glycoprotein (AAG) in humans, with linear binding over a wide concentration range (20 to 600 ng/mL). The results of two-way *in vitro* studies indicate that the binding of tamsulosin to human plasma proteins is not affected by amitriptyline, diclofenac, glyburide, simvastatin plus simvastatin-hydroxy acid metabolite, warfarin, diazepam, propranolol, trichlormethiazide, or chlormadinone. Likewise, tamsulosin had no effect on the extent of binding of these drugs.

**Metabolism:** Tamsulosin is extensively metabolized by cytochrome P450 enzymes (CYP3A4 and CYP2D6) in the liver, followed by extensive glucuronide or sulfate conjugation of metabolites. On administration of a dose of radiolabelled tamsulosin to four healthy volunteers, 97% of the administered radioactivity was recovered, with urine (76%) representing the primary route of excretion compared to feces (21%) over 168 hours. Less than 10% of the dose was recovered as unchanged (parent) compound in the urine.

Metabolites of tamsulosin do not contribute significantly to tamsulosin adrenoreceptor antagonist activity. Furthermore, there is no enantiomeric bioconversion from tamsulosin [R(-) isomer] to the S(+) isomer in studies with mice, rats, dogs, and humans.

Incubations with human liver microsomes showed no evidence of clinically significant interactions between tamsulosin and drugs which are known to interact or be metabolized by hepatic enzymes, such as amitriptyline, diclofenac, albuterol (beta agonist), glyburide (glibenclamide), finasteride (5 alpha-

reductase inhibitor for treatment of BPH), and warfarin. No dose adjustment is warranted in hepatic insufficiency.

**Excretion:** Tamsulosin undergoes restrictive clearance in humans, with a relatively low systemic clearance (2.88 L/h). Tamsulosin exhibits linear pharmacokinetics following single or multiple dosing of tamsulosin hydrochloride controlled-release tablets resulting in a proportional increase in  $C_{max}$  and AUC with increasing doses. Intrinsic clearance is independent of tamsulosin binding to AAG, but diminishes with age, resulting in a 40% overall higher exposure (AUC) in subjects of age 55 to 75 years compared to subjects of age 20 to 32 years.

Following intravenous or oral administration of an immediate-release formulation, the elimination half-life of tamsulosin in plasma ranged from five to seven hours. Because of absorption rate-controlled pharmacokinetics with the tamsulosin hydrochloride controlled-release tablets formulation, the apparent half-life of tamsulosin increases to approximately 12 to 15 hours in healthy volunteers.

### **Special Populations and Conditions**

**Pediatrics:** Tamsulosin hydrochloride is not indicated for use in children. The effectiveness of tamsulosin in children (ages 2 to 16 years) with neuropathic bladder was not demonstrated (see WARNINGS AND PRECAUTIONS, Special Populations, Pediatrics). Pharmacokinetics have not been evaluated in pediatrics.

**Geriatrics:** There were no pharmacokinetic studies conducted in geriatric patients with tamsulosin hydrochloride controlled-release tablets. Cross-study comparisons of overall exposure (AUC) and half-life of tamsulosin hydrochloride capsules indicate that the pharmacokinetic disposition of tamsulosin may be slightly prolonged in geriatric males compared to young healthy male volunteers. However, tamsulosin hydrochloride capsules have been found to be a safe and effective  $\alpha_1$  adrenoreceptor antagonist when administered at therapeutic doses to patients over the age of 65 years.

**Gender Effects:** Tamsulosin hydrochloride is not indicated for use in women. Pharmacokinetics has not been evaluated in women.

**Hepatic Insufficiency:** The pharmacokinetics of tamsulosin have been compared in subjects with hepatic dysfunction (n=8) and in normal subjects (n=8). While a change in the overall plasma concentration of tamsulosin was observed as the result of altered binding to AAG, the unbound (active) concentration of tamsulosin does not change significantly with only a modest (32%) change in intrinsic clearance of unbound tamsulosin. Therefore, patients with mild to moderate hepatic dysfunction do not require an adjustment in tamsulosin hydrochloride dosage.

**Renal Insufficiency:** The pharmacokinetics of tamsulosin have been compared in subjects with moderate (n=6) or severe (n=6) renal impairment and in normal subjects (n=6). While a change in the overall plasma concentration of tamsulosin was observed as the result of altered binding to AAG, the unbound (active) concentration of tamsulosin, as well as the intrinsic clearance, remained relatively constant. Therefore, patients with such renal impairment do not require an adjustment in tamsulosin hydrochloride dosing. Patients with end stage renal disease ( $Cl_{cr} < 10 \text{ mL/min}$ ) have not been studied.

## **STORAGE AND STABILITY**

Store at room temperature (15°C to 30°C).

## **DOSAGE FORMS, COMPOSITION AND PACKAGING**

TAMSULOSIN CR Tablets 0.4 mg: Each yellowish-brown, round, biconvex film-coated tablet, engraved “TA” over “0.4” on one side, and plain on the other side contains 0.4 mg of tamsulosin hydrochloride. Available in HDPE bottles of 100 and 500 tablets and blister packs of 30 and 100 tablets.

In addition to tamsulosin hydrochloride, each tablet also contains the following **non-medicinal ingredients** (in alphabetical order): citric acid, colloidal silicon dioxide, hydroxypropyl cellulose, hydroxypropyl methylcellulose 2208 and 2910, magnesium stearate, polyethylene glycol, sodium alginate and yellow ferric oxide.

## PART II: SCIENTIFIC INFORMATION

### PHARMACEUTICAL INFORMATION


#### Drug Substance

Proper name: Tamsulosin hydrochloride

Chemical name: 5-[(2R)-2-[[2-(2-ethoxyphenoxy) ethyl]amino]propyl]-2-methoxybenzenesulfonamide hydrochloride.

Molecular formula and molecular mass:  $C_{20}H_{28}N_2O_5S \cdot HCl$ ; 444.97 g/mol

Structural Formula:



Physicochemical properties: Tamsulosin HCl is a white to almost white solid. Slightly soluble in water, freely soluble in formic acid, slightly soluble in anhydrous ethanol.

pH: 5.34 (1% aqueous solution)

pKa: 8.4 (secondary amine); 10.2 (sulfonamide)

## CLINICAL TRIALS

### Comparative Bioavailability Studies

A randomized, single dose, double-blinded, 2-way crossover comparative bioavailability study, conducted under fasting conditions was performed on healthy male volunteers. The rate and extent of absorption of tamsulosin was measured and compared following a single oral dose (1 x 0.4 mg tablet) of Flomax<sup>®</sup> CR (tamsulosin hydrochloride) and Tamsulosin (tamsulosin hydrochloride) in 30 volunteers. The results from measured data are summarized in the following table:

Summary Table of the Comparative Bioavailability Data Tamsulosin (A single 0.4 mg dose: 1 x 0.4 mg) From Measured Data/Fasting Conditions Geometric Mean Arithmetic Mean (CV%)				
Parameter	Tamsulosin (Pro Doc Ltée) (Canada)	Flomax <sup>®</sup> CR† (Boehringer Ingelheim (Canada) Ltd.) (Canada)	Ratio of Geometric Means (%)	90% Confidence Interval (%)
AUC <sub>t</sub> (ng•h/mL)	96.471 110.617 (50)	94.411 110.257 (56)	102.2	91.9 – 113.6
AUC <sub>inf</sub> (ng•h/mL)	103.299 119.182 (47)	102.177 124.844 (59)	101.1	88.9 – 114.9
C <sub>max</sub> (ng/mL)	5.273 5.586 (33)	4.786 5.061 (35)	110.2	99.4 – 122.1
T <sub>max</sub> § (h)	5.34 (60)	4.60 (30)		
T <sub>half</sub> § (h)	12.97 (28)	14.60 (45)		
For balanced treatment sequence, results are based on Geometric means. For unbalanced treatment sequence, results are based on Least Squares Means (LSM).				
§ Arithmetic means (CV%) only.				
† Flomax <sup>®</sup> CR is manufactured by Boehringer Ingelheim (Canada) Ltd. and was purchased in Canada.				

A randomized, single dose, double-blinded, 2-way crossover comparative bioavailability study, conducted under fed conditions was performed on healthy male volunteers. The rate and extent of absorption of tamsulosin was measured and compared following a single oral dose (1 x 0.4 mg tablet) of Flomax<sup>®</sup> CR (tamsulosin hydrochloride) and Tamsulosin (tamsulosin hydrochloride) in 18 volunteers. The results from measured data are summarized in the following table:

Summary Table of the Comparative Bioavailability Data				
Tamsulosin				
(A single 0.4 mg dose: 1 x 0.4 mg tablet)				
From Measured Data/Fed Conditions				
Geometric Mean				
Arithmetic Mean (CV%)				
Parameter	Tamsulosin (Pro Doc Ltée) (Canada)	Flomax <sup>®</sup> CR † (Boehringer Ingelheim (Canada) Ltd.) (Canada)	Ratio of Geometric Means (%)	90% Confidence Interval (%)
AUC <sub>t</sub> (ng•h/mL)	140.116 148.673 (36)	145.742 158.189 (42)	96.1	83.9 – 110.2
AUC <sub>inf</sub> (ng•h/mL)	147.466 156.057 (35)	152.511 164.811 (41)	96.7	84.9 – 110.1
C <sub>max</sub> (ng/mL)	7.266 7.550 (28)	8.576 9.198 (41)	84.7	75.4 – 95.2
T <sub>max</sub> § (h)	6.44 (46)	6.78 (33)		
T <sub>half</sub> § (h)	13.41 (35)	13.28 (33)		
For balanced treatment sequence, results are based on Geometric means. For unbalanced treatment sequence, results are based on Least Squares Means (LSM).				
§ Arithmetic means (CV%) only.				
† Flomax <sup>®</sup> CR is manufactured by Boehringer Ingelheim (Canada) Ltd. and was purchased in Canada.				

A randomized, multiple dose, double-blinded, 2-way crossover comparative bioavailability study, conducted under fasting conditions at steady state was performed on healthy male volunteers. The rate and extent of absorption of tamsulosin was measured and compared following a multiple oral dose (7 x 0.4 mg tablet) of Flomax<sup>®</sup> CR (tamsulosin hydrochloride) and Tamsulosin (tamsulosin hydrochloride) in 33 volunteers. The results from measured data are summarized in the following table:

Summary Table of the Comparative Bioavailability Data Tamsulosin (A Multiple 0.4 mg dose: 7 x 0.4 mg) From Measured Data/Fasting Conditions at Steady State Geometric Mean Arithmetic Mean (CV%)				
Parameter	Tamsulosin (Pro Doc Ltée) (Canada)	Flomax <sup>®</sup> CR <sup>†</sup> (Boehringer Ingelheim (Canada) Ltd.) (Canada)	Ratio of Geometric Means (%)	90% Confidence Interval (%)
AUC <sub>tau</sub> (ng•h/mL)	101.810 109.924 (41)	110.651 128.335 (61)	92.0	83.2 – 101.8
C <sub>max</sub> (ng/mL)	7.988 8.382 (32)	8.111 9.019 (51)	98.5	90.3 – 107.4
C <sub>min</sub> (ng/mL)	2.048 2.417 (61)	2.474 3.142 (75)	82.8	70.9 – 96.7
T <sub>max</sub> <sup>§</sup> (h)	3.82 (25)	4.52 (24)		
Fluc <sup>§</sup> (%)	140.78 (30)	123.83 (38)		
For balanced treatment sequence, results are based on Geometric means. For unbalanced treatment sequence, results are based on Least Squares Means (LSM). § Arithmetic means (CV%) only. † Flomax <sup>®</sup> CR is manufactured by Boehringer Ingelheim (Canada) Ltd. and was purchased in Canada.				

## Other Studies

### Study demographics and trial design

Efficacy of tamsulosin hydrochloride controlled-release tablets has been evaluated in two double-blind, randomized, placebo-controlled studies of 12-weeks duration involving a total of 1840 male subjects. Of these, 563 were treated with tamsulosin hydrochloride controlled-release tablets 0.4 mg, 709 with tamsulosin hydrochloride capsules 0.4 mg and 568 with placebo. The main inclusion criteria in both trials were: male patients aged  $\geq 45$  years with symptoms diagnosed as LUTS suggestive of BPH. These patients had to have a total International Prostate Symptom Score (I-PSS) of  $\geq 13$  at enrollment and after 2 week placebo run-in. In both studies, tamsulosin (or placebo) was orally administered at the specified dosage once daily.

The primary efficacy parameter for both studies was the change from baseline to endpoint in Total I-PSS for the full analysis set (FAS). The I-PSS consists of questions that assess the severity of both irritative and obstructive symptoms, with possible scores ranging from 0 to 35. The secondary efficacy analysis contained the changes from baseline in voiding and storage I-PSS subscores, I-PSS QoL score and the individual I-PSS items.


**TABLE 4: EFFECT ON TOTAL I-PSS IN THE 3- MONTH STUDIES**

Study	Treatment Arm	No. Baseline/Endpoint	Baseline Mean (SD)	Endpoint Mean (SD)	Change at Endpoint	Difference vs. Placebo	P-value Vs. Placebo
					Mean (SD) [%]	Mean (95% CI)	
617-CL-303	Placebo	210 <sup>a</sup> /211	17.8 (4.0)	11.7 (6.1)	-6.0 (5.4) [-34.5]	-	0.0016
	Tamsulosin Hydrochloride Controlled-Release tablets 0.4 mg	203/203	18.0 (4.3)	10.4 (5.5)	-7.6 (5.3) [-42.4]	-1.6 (-2.5, -0.6)	


617-CL-307	Placebo	350/350	18.3 (4.5)	12.4 (6.4)	-5.8 (5.6) [-32.0]	-	-
	Tamsulosin Hydrochloride Controlled-Release tablets 0.4 mg	354/354	18.5 (4.4)	10.8 (6.2)	-7.7 (5.8) [41.7]	-1.7 (-2.5, -1.0)	<0.0001
	Tamsulosin Capsules 0.4 mg	700/700	18.5 (4.5)	10.6 (5.9)	-8.0 (5.6) [-43.2]	-2.0 (-2.6,-1.3)	<0.0001

<sup>a</sup> Patient 1607 in the placebo group did not have an I-PSS at baseline (Visit 2) and the Visit 1 I-PSS of this patient was not included in the mean (SD) at baseline

**FIGURE 2: MEAN CHANGE FROM BASELINE IN TOTAL I-PSS OVER TIME IN THE PLACEBO CONTROLLED STUDY**



**FIGURE 3: MEAN CHANGE FROM BASELINE IN TOTAL I-PSS OVER TIME IN THE PLACEBO AND ACTIVE-CONTROLLED STUDY**



In both studies, tamsulosin hydrochloride controlled-release tablets 0.4 mg had a fast onset of action with decrease in I-PSS at 2 to 4 weeks. As evident from Table 4 and Figures 2 and 3, there was a statistically significant reduction ( $p < 0.001$ ) in the I-PSS vs. placebo in both studies indicating a reduction in symptom severity. This was due to a statistically significant improvement in both the irritative and obstructive subscores. Tamsulosin hydrochloride controlled-release tablets 0.4 mg was an efficacious dose and provided a response which was equivalent to that of tamsulosin hydrochloride 0.4 mg capsules confirming the recommendation of once daily dosing of 0.4 mg.

**DETAILED PHARMACOLOGY**

See ACTION AND CLINICAL PHARMACOLOGY section.

## TOXICOLOGY

### **Carcinogenesis, Mutagenesis, and Impairment of Fertility**

Rats administered doses up to 43 mg/kg/day in males and 52 mg/kg/day in females had no increases in tumour incidence with the exception of a modest increase in the frequency of mammary gland fibroadenomas in female rats receiving doses  $\geq 5.4$  mg/kg ( $P < 0.015$ ). The highest doses of tamsulosin evaluated in the rat carcinogenicity study produced systemic exposures (AUC) in rats 3 times the exposures in men receiving doses of 0.8 mg/day.

Mice were administered doses up to 127 mg/kg/day in males and 158 mg/kg/day in females. There were no significant tumour findings in male mice. Female mice treated for 2 years with the two highest doses of 45 and 158 mg/kg/day had statistically significant increases in the incidence of mammary gland fibroadenomas ( $P < 0.0001$ ) and adenocarcinomas ( $P < 0.0075$ ). The highest dose levels of tamsulosin evaluated in the mice carcinogenicity study produced systemic exposures (AUC) in mice 8 times the exposures in men receiving doses of 0.8 mg/day.

The increased incidences of mammary gland neoplasms in female rats and mice were considered secondary to tamsulosin-induced hyperprolactinemia. It is not known if tamsulosin hydrochloride elevates prolactin in humans. The relevance for human risk of the findings of prolactin-mediated endocrine tumours in rodents is not known.

Tamsulosin produced no evidence of mutagenic potential *in vitro* in the Ames reverse mutation test, mouse lymphoma thymidine kinase assay, and chromosomal aberration assays in Chinese hamster ovary cells or human lymphocytes. There were no mutagenic effects in the *in vivo* sister chromatid exchange and mouse micronucleus assay.

Studies in rats revealed significantly reduced fertility in males dosed with single or multiple daily doses of 300 mg/kg/day of tamsulosin (AUC exposure in rats about 50 times the human exposure at a dose of 0.8 mg/day). The mechanism of decreased fertility in male rats is considered to be an effect of the compound on the vaginal plug formation possibly due to changes of semen content or impairment of ejaculation. The effects on fertility were reversible showing improvement by 3 days after a single dose and 4 weeks after multiple dosing. Effects on fertility in males were completely reversed within nine weeks of discontinuation of multiple dosing. Multiple doses of 10 and 100 mg/kg/day tamsulosin (1/5 and 16 times the anticipated human AUC exposure) did not significantly alter fertility in male rats. Effects of tamsulosin on sperm counts or sperm function have not been evaluated.

Studies in female rats revealed significant reductions in fertility after single or multiple dosing with 300 mg/kg/day of the R-isomer or racemic mixture of tamsulosin, respectively. In female rats, the reductions in fertility after single doses were considered to be associated with impairments in fertilization. Multiple dosing with 10 or 100 mg/kg/day of the racemic mixture did not significantly alter fertility in female rats.

## REFERENCES

1. McGrath, J.C., Lepor H., Wyllit M.G. Report of a unique meeting between the alphablocker subcommittee and pharmaceutical industry. *Urol* 48(5), 1996.
2. Price D.T., Lomasney J.W., Allen L.F., Caron M.G., Lefkowitz. Identification, quantification, and localization of mRNA for three distinct  $\alpha_1$  adrenergic receptor subtypes in human prostate. *J Urol* 1993; 150:546-551.
3. Hatano A., Takahashi H., Tamaki M., Komeyama T., Koizumi T., Takeda M. Pharmacological evidence of distinct  $\alpha_1$  - adrenoceptor subtypes mediating the contraction of human prostatic urethra and peripheral artery. *Br J Pharmacol* 1994;113:723-728.
4. Moriyama N., Hamada K., Takanashi M., Kurimoto S., Kimura K., Inagaki O. Evaluation of  $\alpha_1$  - adrenoceptor subtypes in human hypertrophied prostate using [<sup>3</sup>H]YM617, an  $\alpha_1$ -selective antagonist. *Acta Histochem Cytochem* 1994; 27 (3):219-225.
5. Faure C., Pimoule C., Vallancien G., Langer S.Z., Graham D. Identification of  $\alpha_1$  -adrenoceptor subtypes present in the human prostate. *Life Sci* 1994;54(21):1595-1605.
6. Michel M.C., Insel P.A. Comparison of cloned and pharmacologically defined rat tissue  $\alpha_1$  - adrenoceptor subtypes. *Naunyn-Schmiedebergs Arch Pharmacol* 1994;350(2):136-142.
7. Yamada S., Suzuki M., Tanaka C., Mori R., Kimura R., Inagaki O. Comparative study on  $\alpha_1$  - adrenoceptor antagonist binding in human prostate and aorta. *Clinical and Experimental Pharmacol and Physiol* 1994;21:405-411.
8. Chapple, CR. The total approach in lower urinary symptoms/ benign prostatic hyperplasia (LUTS/BPH) management: introduction and conclusions. *Eur Urol* 2003; 2:1-5.
9. Dunn C.J., Matheson A., Faulds D.M. Tamsulosin: A review of its pharmacology and therapeutic efficacy in the management of lower urinary tract symptoms. *Drugs Aging* 2002;19(2):135-161.
10. Narayan P., Evans C.P., Moon T., Long-term safety and efficacy of tamsulosin for the treatment of lower urinary tract symptoms associated with benign prostatic hyperplasia. *J Urol* 2003;170:498-502.
11. Foglar R., Shibata K., Horie K., Hirasawa A., Tsujimoto G. Use of recombinant  $\alpha_1$ -adrenoceptors to characterize subtype selectivity of drugs for the treatment of prostatic hypertrophy. *Eur J Pharmacol* 1995; 288:201-207.
12. Kim H.L., Kim J.C., Benson D..A., Bales G., Gerber G.S. Results of treatment with tamsulosin in men with acute urinary retention. *Tech Urol* 2001; 7(4):256-60.

13. Patel R., Fiske J., Lepor H. Tamsulosin reduces the incidence of acute urinary retention following early removal of the urinary catheter after radical retropubic prostatectomy. *Urology* 2003; 62(2):287-91.
14. Kloner R.A., Jackson G., Emmick J.T., *et al.* Interaction between the phosphodiesterase 5 inhibitor, tadalafil, and two alpha-blockers: doxazosin and tamsulosin. *J Urol* 2004; 4(Suppl):abstr 1195.
15. Romic I., Kiss T., Kisbenedek L., *et al.* Tamsulosin drug ratio in prostate versus free fraction in plasma supports pharmacokinetic (pk) contribution to its uroselectivity. Ann Mtg of the American Urological Association, Chicago, 26 April -1 May 2003.
16. Davis B.J., Noble A., Chapple C.R, Chess-Williams R. Human prostate-artery selectivity of tamsulosin, doxazosin and alfuzosin. *Br J Pharmacol* 2000; 129(Suppl):abstr 52p.
17. Michel MC., Korstanje C, Krauwinkel W, Shear M, Davies J, Quartel A. Comparison of vascular  $\alpha_1$ -adrenoceptor antagonism of tamsulosin in oral controlled absorption system (OCAS) and modified release (MR) formulations. *European Urology Supplements* 4 (2005) 45–52.
18. Michel MC., Korstanje C, Krauwinkel W, Shear M, Davies J, Quartel A. Cardiovascular safety of the oral controlled absorption system (OCAS) formulation compared to the modified release (MR) formulation. *European Urology Supplements* 4 (2005) 53-60.
19. Product Monograph - Flomax<sup>®</sup> CR (tamsulosin hydrochloride) Controlled-Release Tablets 0.4 mg. Boehringer Ingelheim (Canada) Ltd. Control # 229213 Date of Revision: August 28, 2019.

**PART III: CONSUMER INFORMATION**

**P<sup>r</sup>TAMSULOSIN CR**  
**Tamsulosin Hydrochloride Controlled-Release**  
**Tablets**  
**Manufacturer Standard**

**This leaflet is part III of a three-part "Product Monograph" published when TAMSULOSIN CR was approved for sale in Canada and is designed specifically for Consumers. This leaflet is a summary and will not tell you everything about TAMSULOSIN CR. Contact your doctor or pharmacist if you have any questions about the drug.**

Please read this leaflet before you start taking TAMSULOSIN CR (tamsulosin hydrochloride). Also, read it each time you renew your prescription, just in case new information has been added.

**ABOUT THIS MEDICATION****What the medication is used for:**

Your doctor has prescribed TAMSULOSIN CR because you have lower urinary tract symptoms (LUTS) associated with a medical condition called benign prostatic hyperplasia or BPH.

**What it does:**

TAMSULOSIN CR acts by relaxing muscles in the prostate and bladder neck at the site of obstruction, resulting in improved urine flow, and reduced BPH symptoms.

BPH is an enlargement of the prostate gland. BPH is the most common cause of lower urinary tract symptoms (LUTS) in elderly males. After age 50, most men develop enlarged prostates. The prostate is located below the bladder. As the prostate enlarges, it may slowly restrict the flow of urine. This can lead to symptoms such as:

- frequent sleep interruption caused by a need to urinate;
- having a weak urinary stream;
- a sensation of not emptying your bladder completely after you finish urinating;
- pushing or straining to begin urination;
- stopping and starting again several times when urinating;
- urinating again less than 2 hours after you finish urinating;
- finding it difficult to postpone urination.

**When it should not be used:**

- You should not use TAMSULOSIN CR if you are allergic (hypersensitive) to tamsulosin or any of the other ingredients in TAMSULOSIN CR. Hypersensitivity may present as sudden local swelling of the soft tissues of the body (e.g. the throat or tongue), difficult breathing and/or itching and rash (angioedema).;
- You should not use TAMSULOSIN CR if you are taking ketoconazole (an antifungal used to treat fungal skin infections)

**What the medicinal ingredient is:**

Tamsulosin Hydrochloride

**What the important non-medicinal ingredients are:**

citric acid, colloidal silicon dioxide, hydroxypropyl cellulose, hydroxypropyl methylcellulose 2208 and 2910, magnesium stearate, polyethylene glycol, sodium alginate and yellow ferric oxide.

**What dosage forms it comes in:**

Controlled-release Tablets. Each yellowish-brown, round, biconvex film-coated tablet, engraved "TA" over "0.4" on one side, and plain on the other side, contains 0.4 mg tamsulosin hydrochloride.

**WARNINGS AND PRECAUTIONS**

**Rarely, fainting can occur during the use of TAMSULOSIN CR, as with other medicinal products of this type. At the first signs of dizziness or weakness you should sit or lie down until they have disappeared.**

**TAMSULOSIN CR should not be used in women or children.**

**BEFORE you use TAMSULOSIN CR talk to your doctor or pharmacist:**

- if you suffer from severe liver problems;
- if you suffer from fainting due to reduced blood pressure when changing posture (going to sit or stand up);
- if you are allergic to sulfa drugs;
- if you have kidney problems;
- if you have previously taken tamsulosin hydrochloride and became unwell;

- if you are going to have eye surgery for cataract (cloudiness of the lens of the eye) and/or glaucoma (high pressure inside the eye), ask your doctor if you need to temporarily stop taking this medicine before your surgery. A condition known as Intraoperative Floppy Iris Syndrome (IFIS) has been reported in association with tamsulosin hydrochloride during cataract and glaucoma surgery.

TAMSULOSIN CR may cause dizziness. Do not drive or operate machinery or perform a hazardous task for 12 hours after taking the initial dose of TAMSULOSIN CR, or until you know how the medication affects you.

**You must see your doctor regularly.** While taking TAMSULOSIN CR, you must have regular checkups. Follow your doctor's advice about when to have these checkups.

### **INTERACTIONS WITH THIS MEDICATION**

Taking TAMSULOSIN CR with other medicines from the same class (alpha<sub>1</sub>- adrenoceptor blockers) may cause an unwanted decrease in blood pressure.

Please tell your doctor or pharmacist if you are taking or have recently taken cimetidine, ketoconazole, warfarin or any other medicines, including medicines obtained without a prescription.

### **PROPER USE OF THIS MEDICATION**

#### **Usual dose:**

Follow your doctor's advice about how to take TAMSULOSIN CR. You should take one tablet (0.4 mg) once daily at the same time each day with or without food.

Do not crush or chew TAMSULOSIN CR tablets. These tablets are specially formulated to control the delivery of tamsulosin hydrochloride to the blood stream.

Do not share TAMSULOSIN CR with anyone else; it is prescribed only for you.

If you interrupt your treatment for several days or more, resume treatment after consulting with your physician.

#### **Overdose:**

If you think you have taken too much TAMSULOSIN CR, contact your healthcare professional, hospital emergency department or regional poison control centre immediately, even if there are no symptoms.

#### **Missed Dose:**

You may take your daily TAMSULOSIN CR tablet later the same day if you have forgotten to take it as recommended. If you have missed a day, just continue to take your daily tablet as prescribed. Never take a double dose to make up for the forgotten tablet.

### **SIDE EFFECTS AND WHAT TO DO ABOUT THEM**

Like all prescription drugs, TAMSULOSIN CR may cause side effects. Side effects due to TAMSULOSIN CR may include:

#### **Common (1 to 10%):**

Abnormal ejaculation (ejaculation disorder). This means that semen does not leave the body via the urethra, but instead goes into the bladder (retrograde ejaculation) or the ejaculation volume is reduced or absent (ejaculation failure). This phenomenon is harmless.

#### **Uncommon (0.1 to 1%):**

Headache, reduced blood pressure (orthostatic hypotension) e.g. when getting up quickly from a seated or lying position sometimes associated with dizziness; runny or blocked nose (rhinitis), diarrhea, feeling sick and vomiting, constipation, weakness (asthenia), rashes, itching and hives (urticaria).

#### **Rare (0.01 to 0.1%):**

Fainting - at the first signs of dizziness or weakness you should sit or lie down until they have disappeared.

Allergic reaction (angioedema) – the symptoms include sudden local swelling of the soft tissues of the body (e.g. the throat or tongue), difficulty breathing and/or itching and rash.

#### **Very rare (<0.01%):**

Priapism (the painful prolonged unwanted erection of the penis), which is unrelieved by sexual intercourse or masturbation. This condition, if untreated, can lead to permanent inability to have an erection. Stevens-Johnson syndrome (a life-threatening skin disorder which can cause: rash, sores in the mouth, and sores on the body).

#### **Not Known (frequency cannot be estimated from the available data):**

Blurred vision or impaired vision, nose bleeds (epistaxis), serious skin rashes (erythema multiform, dermatitis exfoliative), dry mouth, increased sensitivity of the skin to light caused by TAMSULOSIN CR.

**IMPORTANT: PLEASE READ**

If you are undergoing eye surgery because of cloudiness of the lens (cataract) or high pressure inside the eye (glaucoma) and are already taking or have previously taken tamsulosin hydrochloride, the pupil may dilate poorly and the iris (the coloured circular part of the eye) may become floppy during the procedure.

*This is not a complete list of side effects. For any unexpected effects while taking TAMSULOSIN CR, contact your doctor or pharmacist immediately, so that these effects may be properly addressed.*

Serious side effects and what to do about them				
Symptom/effect		Talk to your healthcare professional		Stop taking drug and get immediate medical help
		Only if severe	In all cases	
Common	Dizziness, particularly when getting up from a seated or lying position		✓	
Uncommon	Palpitations (feeling of rapid beating of the heart that may be more forceful)		✓	
	Rashes, itching and hives (urticaria)			✓
	Reduced blood pressure e.g. when getting up quickly from a seated or lying position, sometimes associated with		✓	

Serious side effects and what to do about them				
	dizziness			
Rare	Fainting			✓
	*Sudden local swelling of soft tissues of the body (e.g. the throat or tongue), difficulty breathing and/or itching and rash (angioedema)			✓
Very rare	*Priapism (painful prolonged unwanted erection)			✓

\* If you suspect such symptoms, call your doctor or go to an Emergency Room as soon as possible.

**HOW TO STORE IT**

Store at room temperature (15°C to 30°C).

Keep TAMSULOSIN CR and all medicines out of reach of children

### **Reporting Side Effects**

You can report any suspected side effects associated with the use of health products to Health Canada by:

- Visiting the Web page on Adverse Reaction Reporting (<https://www.canada.ca/en/health-canada/services/drugs-health-products/medeffect-canada/adverse-reaction-reporting.html>) for information on how to report online, by mail or by fax; or
- Calling toll-free at 1-866-234-2345.

*NOTE: Contact your health professional if you need information about how to manage your side effects. The Canada Vigilance Program does not provide medical advice.*

### **MORE INFORMATION**

#### **If you want more information about TAMSULOSIN CR:**

- Talk to your healthcare professional
- Find the full product monograph that is prepared for healthcare professionals and includes this Consumer Information by visiting the Health Canada website (<https://health-products.canada.ca/dpd-bdpp/index-eng.jsp>). This leaflet, plus the full product monograph prepared for health professionals, can be obtained by contacting Pro Doc Ltée at 1-800-361-8559, [www.prodoc.qc.ca](http://www.prodoc.qc.ca) or [info@prodoc.qc.ca](mailto:info@prodoc.qc.ca).

This leaflet was prepared by  
Pro Doc Ltée, Laval, Québec, H7L 3W9.

Last revised: January 21, 2021