

PRODUCT MONOGRAPH

Pr JAMP-METOPROLOL-L

Metoprolol Tartrate Tablets USP

25mg, 50 mg and 100 mg film coated tablets,
USP

β -Adrenergic Receptor Blocking Agent

JAMP Pharma Corporation
1310 rue Nobel
Boucherville, Québec
J4B 5H3, Canada

Date of Revision:
January 25, 2021

Submission Control No: 247050

Table of Contents

PART I: HEALTH PROFESSIONAL INFORMATION..... 3

SUMMARY PRODUCT INFORMATION.....3

INDICATIONS AND CLINICAL USE.....3

CONTRAINDICATIONS.....4

WARNINGS AND PRECAUTIONS5

ADVERSE REACTIONS..... 10

DRUG INTERACTIONS 12

DOSAGE AND ADMINISTRATION..... 18

OVERDOSAGE.....20

ACTION AND CLINICAL PHARMACOLOGY 21

STORAGE AND STABILITY24

DOSAGE FORMS, COMPOSITION AND PACKAGING.....25

PART II: SCIENTIFIC INFORMATION..... 26

PHARMACEUTICAL INFORMATION.....26

CLINICAL TRIALS.....27

DETAILED PHARMACOLOGY28

TOXICOLOGY29

REFERENCES32

PART I: HEALTH PROFESSIONAL INFORMATION..... 36

^{Pr}**JAMP-METOPROLOL-L**

Metoprolol Tartrate Tablets USP

PART I: HEALTH PROFESSIONAL INFORMATION SUMMARY

PRODUCT INFORMATION

| Route of Administration | Dosage Form/ Strength | All Nonmedicinal Ingredients |
|--------------------------------|--|---|
| Oral | Film coated tablets, 25mg, 50mg and 100 mg | Microcrystalline cellulose, lactose, Sodium starch glycolate, povidone, colloidal silicone dioxide, magnesium stearate. In addition, these dosage strengths also contain 25mg tablet: Opadry 03F580000 White (HPMC 2910/Hypromellose, titanium dioxide, macrogol/Polyethylene glycol, talc). 50mg tablet: Opadry 03F540232 Pink (HPMC 2910/Hypromellose, titanium dioxide, macrogol/Polyethylene glycol, talc, FD&C Red #40, D&C Yellow #10, D&C Red #30). 100mg tablet: Opadry 03F505098 Blue (HPMC 2910/Hypromellose, titanium dioxide, macrogol/Polyethylene glycol, talc, FD&C Blue #2, FD&C Blue #1). |

INDICATIONS AND CLINICAL USE

Hypertension

JAMP-METOPROLOL-L (Metoprolol Tartrate Tablets USP) is indicated for mild or moderate hypertension. Usually combined with other antihypertensive agents (thiazide diuretics), it may be tried alone when the physician judges that a beta-blocker, rather than a diuretic, should be the initial treatment.

Combining JAMP-METOPROLOL-L with a diuretic or peripheral vasodilator has been found to be compatible and generally more effective than metoprolol tartrate alone. Limited experience with other antihypertensive agents has not shown evidence of incompatibility with JAMP-METOPROLOL-L.

JAMP-METOPROLOL-L is not recommended for the emergency treatment of hypertensive crises.

Angina Pectoris

JAMP-METOPROLOL-L is indicated for the long-term treatment of angina pectoris due to ischemic heart disease.

Myocardial Infarction

JAMP-METOPROLOL-L is indicated in the treatment of hemodynamically stable patients with definite or suspected acute myocardial infarction, to reduce cardiovascular mortality.

In patients with proven myocardial infarction, oral treatment can begin within 3 to 10 days of the acute event (*see Dosage and Administration*). Data are not available as to whether benefit would ensue if the treatment is initiated later.

Clinical trials have shown that patients with unconfirmed myocardial infarction received no benefit from early metoprolol tartrate therapy.

Geriatrics:

Caution is indicated when using JAMP-METOPROLOL-L in elderly patients. An excessively pronounced decrease in blood pressure or pulse rate may cause the blood supply to vital organs to fall to inadequate levels.

Pediatrics:

No pediatric studies have been performed. The safety and efficacy of JAMP-METOPROLOL-L in pediatric patients have not been established.

CONTRAINDICATIONS

JAMP-METOPROLOL-L (Metoprolol Tartrate Tablets USP) should not be used in the presence of:

- Known hypersensitivity to metoprolol and derivatives, JAMP-METOPROLOL-L components, or hypersensitivity to other beta-blockers (cross-sensitivity between beta-blockers can occur)
- Sinus bradycardia
- Sick sinus syndrome
- Second and third degree A-V block
- Right ventricular failure secondary to pulmonary hypertension
- Overt heart failure
- Cardiogenic shock
- Severe peripheral arterial circulatory disorders
- Anesthesia with agents that produce myocardial depression, (e.g., ether)
- Untreated pheochromocytoma

Myocardial Infarction Patients - Additional Contraindications

JAMP-METOPROLOL-L is contraindicated in patients with a heart rate < 45 beats/min; significant heart block greater than first degree (PR interval \geq 0.24 s); systolic blood pressure < 100 mmHg; or moderate to severe cardiac failure (see *Warnings and Precautions*).

WARNINGS AND PRECAUTIONS

General

Cardiovascular system: Special caution should be exercised when administering Jamp-Metoprolol-L to patients with a history of heart failure. Sympathetic stimulation is a vital component supporting circulatory function in congestive heart failure, and inhibition with β -blockade always carries the potential hazard of further depressing myocardial contractility and precipitating cardiac failure. The positive inotropic action of digitalis may be reduced by the negative inotropic effect of metoprolol tartrate when the two drugs are used concomitantly. The effects of β -blockers and digitalis are additive in depressing A-V conduction. This also applies to combinations with calcium-antagonists of the verapamil type or some antiarrhythmics (see **Drug Interactions**).

In patients without a history of cardiac failure, continued depression of the myocardium over a period of time can, in some cases, lead to cardiac failure and/or hypotension (systolic blood pressure ≤ 90 mmHg). Therefore, at the first sign or symptom of impending cardiac failure, patients should be fully digitalized and/or given a diuretic and the response observed closely. If cardiac failure continues, despite adequate digitalization and diuretic therapy, Jamp-Metoprolol-L therapy should be reduced or withdrawn.

Cardiovascular

Severe Sinus Bradycardia: Severe sinus bradycardia may occur after β_1 -adrenergic receptor blockade with metoprolol tartrate because of unopposed vagal activity. Very rarely a pre-existing A-V conduction disorder of moderate degree may become aggravated, possibly leading to A-V block. In such cases, dosage should be reduced or gradually withdrawn. Atropine, isoproterenol or dobutamine should be considered in patients with acute myocardial infarction.

Prinzmetal's angina: Beta-blockers may increase the number and duration of angina attacks in patients with Prinzmetal's angina (variant angina pectoris).

Peripheral Circulatory Disorders: Metoprolol may aggravate the symptoms of peripheral arterial circulatory disorders, mainly due to its blood pressure lowering effect.

Myocardial Infarction - Additional Warnings

Cardiac Failure: Depression of the myocardium with metoprolol tartrate may lead to cardiac failure (see general **Warnings** above). Special caution should be exercised when administering JAMP-METOPROLOL-L to patients with a history of cardiac failure or those with minimal cardiac reserve. Should failure occur, treatment should be as described in **WARNINGS**.

Severe Sinus Bradycardia: Severe sinus bradycardia may occur with JAMP-METOPROLOL-L use (see general **Warnings** above). Acute myocardial infarction (particularly inferior infarcts) may significantly decrease sinus rate. If the rate falls below 40 beats/min, especially with signs of decreased cardiac output, administer atropine (0.25-0.5 mg) intravenously. If atropine treatment is unsuccessful, discontinue JAMP-METOPROLOL-L and consider cautious administration of isoproterenol or installation of a cardiac pacemaker.

A-V Conduction: Metoprolol Tartrate Tablets USP slows A-V conduction and may produce

significant first- (PR interval \geq 0.24 sec), second-, or third-degree heart block. Acute myocardial infarction may also produce heart block. If heart block occurs, discontinue JAMP-METOPROLOL-L and administer atropine (0.25-0.5 mg) intravenously. If atropine treatment is unsuccessful, consider cautious administration of isoproterenol or installation of a cardiac pacemaker. Because of their negative effect on atrioventricular conduction, beta-blockers, including Jamp- Metoprolol-L, should only be given with caution to patients with first degree atrioventricular block.

Hypotension: If hypotension (systolic blood pressure \leq 90 mmHg) occurs, JAMP-METOPROLOL-L should be discontinued, and the hemodynamic status of the patient and the extent of myocardial ischemia carefully assessed. Invasive monitoring of central venous, pulmonary capillary wedge, and arterial pressures may be required. Appropriate therapy with fluids, positive inotropic agents, balloon counterpulsation, or other treatment modalities should be instituted. If hypotension is associated with sinus bradycardia or A-V block, treatment should be directed at reversing these (see above).

Abrupt withdrawal

Patients with angina or hypertension should be warned against abrupt discontinuation of JAMP-METOPROLOL-L. There have been reports of severe exacerbation of angina, and of myocardial infarction or ventricular arrhythmias occurring in patients with angina pectoris, following abrupt discontinuation of β -blocker therapy. The last two complications may occur with or without preceding exacerbation of angina pectoris. Therefore, when discontinuation of JAMP-METOPROLOL-L is planned in patients with angina pectoris or previous myocardial infarction, the dosage should be gradually reduced over a period of about two weeks. The patient should be carefully observed. The same frequency of administration should be maintained. In situations of greater urgency, JAMP-METOPROLOL-L therapy should be discontinued stepwise and with closer observation. If angina markedly worsens or acute coronary insufficiency develops, it is recommended that treatment with JAMP-METOPROLOL-L be reinstated promptly, at least temporarily.

Patients should be warned against interruption or discontinuation of therapy without the physician's advice. Because coronary artery disease is common and may be unrecognized, it is prudent not to discontinue JAMP-METOPROLOL-L therapy abruptly even in patients treated only for hypertension.

Endocrine and Metabolism

Thyrototoxicosis: Although metoprolol has been used successfully for the symptomatic (adjuvant) therapy of thyrototoxicosis, possible deleterious effects from long-term use of JAMP-METOPROLOL-L have not been adequately appraised. β -blockade may mask the clinical signs of continuing hyperthyroidism or its complications, and give a false impression of improvement. Therefore, abrupt withdrawal of metoprolol tartrate may be followed by an exacerbation of the symptoms of hyperthyroidism, including thyroid storm.

Diabetic patients: JAMP-METOPROLOL-L should be administered cautiously to patients spontaneously hypoglycemic or diabetic patients who are receiving insulin or oral hypoglycemic agents. β -adrenergic receptor blockers, including metoprolol tartrate, affect glucose metabolism and may mask the premonitory signs and symptoms of acute hypoglycemia, such as tachycardia. In patients with insulin or non-insulin dependent diabetes, especially labile diabetes, or with a history of spontaneous hypoglycaemia, beta-blockade may result in the loss of diabetic control and delayed recovery from hypoglycaemia. The dose of insulin or oral hypoglycaemic agent may need adjustment. Diabetic patients receiving JAMP-METOPROLOL-L should be monitored to ensure

that diabetes control is maintained.

Pheochromocytoma: Where a β -blocker is prescribed for a patient known to be suffering from a pheochromocytoma, an alpha-blocker should be given concomitantly and only after the alpha-blocker has been initiated.

Hepatic/Biliary/Pancreatic

Metoprolol Tartrate Tablets USP is mainly eliminated by means of hepatic metabolism (see ***Actions and Clinical Pharmacology - Pharmacokinetics***).

Hepatic impairment: may increase the systemic bioavailability of metoprolol and reduce its total clearance, leading to increased plasma concentrations. Therefore, JAMP-METOPROLOL-L should be used with caution in patients with impaired liver function. Liver function tests should be performed at regular intervals during long-term treatment (see ***Actions and Clinical Pharmacology - Pharmacokinetics***). Therefore, hepatic impairment may increase the systemic bioavailability of metoprolol and reduce its total clearance, leading to increased plasma concentrations. Therefore, dose adjustment and regular monitoring of hepatic function are advised in patients with mild to moderate hepatic impairment. Patients with severe hepatic impairment should be treated with caution i.e. lower initial and maintenance doses as well as regular monitoring of hepatic function, as they are more sensitive to therapeutic effects/adverse effects of drugs.

Interactions

Calcium channel blocker of the verapamil (phenylalkylamine) type should not be given intravenously to patients receiving JAMP-METOPROLOL-L because there is a risk of cardiac arrest in this situation (see ***Drug Interactions***). Patients taking an oral calcium channel blocker of the verapamil type in combination with JAMP-METOPROLOL-L should be closely monitored. See the complete list of observed and potential drug-drug and other drug interactions with Jamp-Metoprolol-L in ***Drug Interactions*** section.

Peripheral vascular disease:

Beta-blockade may impair the peripheral circulation and exacerbate the symptoms of peripheral vascular disease (see ***Contraindications***).

Peri-Operative Considerations

Anesthesia and Surgery: The necessity or desirability of withdrawing beta-blocking agents prior to major surgery is controversial. The impaired ability of the heart to respond to reflex adrenergic stimuli may augment the risks of general anaesthesia and surgical procedures. The benefits of continuing a treatment with a beta-blocker should be balanced against the risk of withdrawing it in each patient. However, care should be taken to avoid using anesthetic agents that may depress the myocardium. Vagal dominance, if it occurs, may be corrected with atropine (1-2 mg i.v.).

In patients receiving beta-blocker therapy, inhalation anaesthetics may enhance the cardiodepressant effect. Beta-blockade may have beneficial effects in decreasing the incidence of arrhythmias and myocardial ischaemia during anaesthesia and the postoperative period. It is currently recommended that maintenance beta-blockade be continued peri-operatively. The anaesthetist must be made aware of beta-blockade because of the potential for interactions with other drugs, resulting in severe bradyarrhythmias and hypotension, the decreased reflex ability to

compensate for blood loss, hypovolaemia and regional sympathetic blockade, and the increased propensity for vagal-induced bradycardia. Incidents of protracted severe hypotension or difficulty restoring normal cardiac rhythm during anaesthesia have been reported.

Modern inhalational anaesthetic agents are generally well tolerated, although older agents (ether, cyclopropane, methoxyflurane, trichlorethylene) were sometimes associated with severe circulatory depression in the presence of beta-blockade. If it is thought necessary to withdraw beta-blocker therapy before surgery, this should be done gradually and completed about 48 hours before anaesthesia. Since metoprolol is a competitive inhibitor of β -adrenoceptor agonists, its effects may be reversed, if necessary, by sufficient doses of such agonists as isoproterenol or dobutamine.

Renal:

Renal impairment: In patients with severe renal disease, haemodynamic changes following beta-blockade may impair renal function further. Beta-blockers which are excreted mainly by the kidney may require dose adjustment and safety monitoring in patients with severe renal impairment, including renal failure.

Respiratory

Bronchospastic Diseases: In general, patients with bronchospastic diseases should not receive β -blockers, including JAMP-METOPROLOL-L. However, because of its relative β_1 -selectivity, JAMP-METOPROLOL-L (Metoprolol Tartrate Tablets USP) may be used with caution in patients with asymptomatic bronchospastic disease who do not respond to, or cannot tolerate, other antihypertensive treatment. Since β_1 -selectivity is not absolute, a β_2 -stimulating agent should preferably be administered concomitantly, and the lowest possible dose of Jamp- Metoprolol-L should be used. In these circumstances it would be prudent initially to administer JAMP-METOPROLOL-L in smaller doses three times daily, instead of larger doses two times daily, to avoid the higher plasma levels associated with the longer dosing interval (see *Dosage and Administration*).

Because it is unknown to what extent β_2 -stimulating agents may exacerbate myocardial ischemia and the extent of infarction, these agents should not be used prophylactically in patients with proven or suspected acute myocardial infarction. If bronchospasm not related to congestive heart failure occurs, JAMP-METOPROLOL-L should be discontinued. A theophylline derivative or a β_2 -agonist may be administered cautiously, depending on the clinical condition of the patient. Both theophylline derivatives and β_2 -agonists may produce serious cardiac arrhythmias.

Sensitivity/Resistance

Anaphylactic reactions: There may be increased difficulty in treating an allergic type reaction in patients on β -blockers. Whenever possible, β -blockers, including JAMP-METOPROLOL-L, should be avoided in patients who are at risk of anaphylaxis. In these patients, the reaction may be more severe due to pharmacologic effects of the β -blockers and problems with fluid changes. Epinephrine should be administered with caution since it may not have its usual effects in the treatment of anaphylaxis. On the one hand, larger doses of epinephrine may be needed to overcome the bronchospasm, while on the other these doses can be associated with excessive alpha-adrenergic stimulation with consequent hypertension, reflex bradycardia and heart block and possible potentiation of bronchospasm. Alternatives to the use of large doses of epinephrine include vigorous supportive care such as fluids and the use of β -agonists including parenteral salbutamol or isoproterenol, to overcome bronchospasm and norepinephrine to overcome hypotension.

Skin

Oculomucocutaneous Syndrome: Various skin rashes and conjunctival xerosis have been reported with β -blockers, including metoprolol tartrate. Oculomucocutaneous syndrome, a severe syndrome whose signs include conjunctivitis sicca and psoriasiform rashes, otitis, and sclerosing serositis has occurred with the chronic use of one β -adrenergic receptor-blocking agent (practolol). This syndrome has not been observed with metoprolol tartrate or any other such agent. However, physicians should be alert to the possibility of such reactions and should discontinue treatment in the event that they occur (see ***Warnings and Precautions-Abrupt withdrawal***).

Special Populations

Women of child-bearing potential: Upon confirming the diagnosis of pregnancy, women should immediately inform the doctor and stop gradually taking the drug. The use of any drug in patients of child-bearing potential requires that the anticipated benefit be weighed against the possible hazards.

Pregnant Women: There is a limited amount of data on the use of metoprolol in pregnant women. Metoprolol crosses the placental barrier. Since JAMP-METOPROLOL-L has not been studied in human pregnancy, the drug should not be given to pregnant women.

Nursing Women: Metoprolol is excreted in breast milk. If drug use is essential, patients should stop nursing.

Pediatrics: No pediatric studies have been performed. The safety and efficacy of Jamp-Metoprolol-L in pediatric patients have not been established.

Geriatrics: Caution is indicated when using JAMP-METOPROLOL-L in elderly patients. An excessively pronounced decrease in blood pressure or pulse rate may cause the blood supply to vital organs to fall to inadequate levels. Lower starting and maintenance doses and safety monitoring are advised in these patients (see ***Indications and Clinical use, Dosage and Administration*** and ***Action and Clinical Pharmacology-Pharmacokinetics-Special populations***).

Driving and using machines

Dizziness, fatigue or visual impairment may occur during treatment with JAMP-METOPROLOL-L (see ***Adverse Drug reactions***) and may adversely affect the patient's ability to drive or use machines.

Patients should be advised to avoid operating automobiles and machinery or engaging in other tasks requiring alertness until the patient's response to JAMP-METOPROLOL-L therapy has been determined.

ADVERSE REACTIONS

Adverse Drug Reaction Overview

The most common adverse events reported are exertional tiredness, gastrointestinal disorders,

and disturbances of sleep patterns. The most serious adverse events reported are congestive heart failure, bronchospasm and hypotension.

Reported adverse events according to organ systems are:

| | |
|---|---|
| Cardiac disorders | Secondary effects of decreased cardiac output which include: syncope, vertigo, light-headedness and postural hypotension; Significant Conduction disorders (First, Second and third degree A-V block) (<i>see Contraindications</i>); Congestive heart failure (<i>see Warnings</i>); Severe bradycardia; Hot flushes; Arrhythmias; Lengthening of PR interval; Palpitations; Sinus arrest; Cold extremities; Claudication; Chest pain |
| Vascular disorders | Raynaud's phenomenon; Gangrene in patients with pre-existing severe peripheral circulatory disorders; Oedema |
| Psychiatric disorders | Mental depression; Vivid dreams/ nightmares; Hallucination; Personality disorder |
| Nervous System disorders | Headache, Weakness, Dizziness, Sedation, Light-headedness, Somnolence, insomnia, Vertigo, Paresthesia, Anxiety, Depressed level of consciousness |
| Gastrointestinal disorders | Diarrhea, Abdominal pain, Constipation, Heartburn, Flatulence, Dry mouth, Nausea and vomiting, Retroperitoneal fibrosis |
| Hepatobiliary disorders | Hepatitis |
| Respiratory disorders | Shortness of breath; Wheezing; Bronchospasm; Rhinitis; Status asthmaticus; Exertional dyspnea |
| Skin and subcutaneous tissue disorders (<i>see Warnings</i>) | Rash (exanthema, urticaria, psoriasiform and dystrophic skin lesions); Hyperhidrosis; Pruritus; Photosensitivity reaction; Alopecia; Worsening of psoriasis |
| Musculoskeletal and connective tissue disorders | Muscle spasms; Arthritis |
| Reproductive system and breast disorders | Erectile dysfunction; Libido disorder; Peyronie's disease |

| | |
|---|---|
| Ear and labyrinth disorders | Tinnitus; Hearing disorders (e.g. hypoacusis or deafness) when doses exceed those recommended |
| Eye disorders | Dry eyes, eye irritation; Visual impairment (e.g. blurred vision); Conjunctivitis |
| General disorders and administration site conditions | Fatigue; Exertional tiredness |
| Metabolism | Weight increase |

Adverse reactions in clinical trials

In a placebo-controlled study in patients with acute myocardial infarction the incidence of the following cardiovascular reactions were:

| | Metoprolol | Placebo |
|---|------------|---------|
| Orthostatic hypotension (systolic BP < 90 mmHg) | 27.4% | 23.2% |
| Bradycardia (heart rate < 40 beats/min) | 15.9% | 6.7% |
| Second- or third-degree heart block | 4.7% | 4.7% |
| First-degree heart block (PR ≥ 0.24 s) | 5.3% | 1.9% |
| Cardiac failure | 27.5% | 29.6% |

Abnormal Hematologic and Clinical Chemistry Findings

Clinical Laboratory

The following laboratory parameters have been elevated on rare occasions: transaminases, BUN, alkaline phosphatase and bilirubin.

Hematology

Isolated cases of thrombocytopenia and leucopenia.

Post-Market Adverse Drug Reactions

The following adverse reactions have been derived from post-marketing experience with metoprolol tartrate via spontaneous case reports and literature cases. Because these reactions are reported voluntarily from a population of uncertain size and are subject to confounding factors, it is not possible to reliably estimate their frequency which is therefore categorized as not known. Adverse drug reactions are listed according to system organ classes in MedDRA. Within each system organ class, ADRs are presented in order of decreasing seriousness.

Nervous system disorders

Confusional state

Investigations

Blood triglycerides increased, High Density Lipoprotein (HDL) decreased

DRUG INTERACTIONS

Overview

Established or Potential Drug-Drug Interactions

(Legend: CT = Clinical Trial; C=Postmarket (Case Study); T = Theoretical)

| Metoprolol | Ref | Effect | Clinical comment |
|-------------------------------|-----|---|--|
| Alcohol | C | Increased concentration of metoprolol in blood | Metoprolol modifies the pharmacokinetics (decreases the elimination rate) of alcohol. Which may increase certain side effects of metoprolol |
| Anti-adrenergic agents | C | Potentiate Antihypertensive effect of alpha-adrenergic blockers | Antihypertensive effect of alpha-adrenergic blockers such as guanethidine, betanidine, reserpine, alpha-methyldopa or clonidine may be potentiated by β -blockers. β -adrenergic blockers may also potentiate the postural hypotensive effect of the first dose of prazosin, probably by preventing reflex tachycardia. On the contrary, β -adrenergic blockers may also potentiate the hypertensive response to withdrawal of clonidine as patients receiving concomitant clonidine and β -adrenergic blocker. Withdrawing the β -blocker several days before the clonidine may reduce the danger of rebound effects. |

| Metoprolol | Ref | Effect | Clinical comment |
|-------------------------------------|------------|--|---|
| Antiarrhythmic Agents | C | Potentiate the negative inotropic effect of antiarrhythmic agents and their effect on atrial-conduction time | β -blockers may potentiate the negative inotropic effect of antiarrhythmic agents and their effect on atrial-conduction time. Particularly, in patients with preexisting sinus node dysfunction, concomitant administration of amiodarone may result in additive electrophysiologic effects including bradycardia, sinus arrest, and atrioventricular block. Antiarrhythmic agents such as quinidine, tocainide, procainamide, ajmaline, amiodarone, flecainide, and disopyramide may potentiate the effects of metoprolol tartrate on heart rate and atrioventricular conduction. |
| Other Antihypertensive drugs | CT | Hypertension | JAMP-METOPROLOL-L dosage should be adjusted to the individual requirements of the patient especially when used concomitantly with other antihypertensive agents (see <i>Dosage and Administration</i>). Patients receiving concurrent treatment with catecholamine depleting drugs, other beta-blockers (including those in form of eye drops, such as timolol), or monoamine oxidase (MAO) inhibitors, should be carefully monitored. In addition, possibly significant hypertension may theoretically occur up to 14 days following discontinuation of the concomitant administration with an irreversible MAO inhibitor. |

| Metoprolol | Ref | Effect | Clinical comment |
|--|------------|--|--|
| Calcium Channel Blockers (IV use) | CT | Potentiate the depressant effects of β -blockers | Calcium channel blockers such as verapamil and diltiazem may potentiate the depressant effects of beta-blockers on blood pressure, heart rate, cardiac contractility and atrioventricular conduction. A calcium channel blocker of the verapamil (phenylalkylamine) type should not be given intravenously to patients receiving JAMP-METOPROLOL-L because there is a risk of cardiac arrest in this situation. However, in exceptional cases, when the physician considers concomitant use essential, such use should be instituted gradually in a hospital setting under careful supervision. Negative inotropic, dromotropic and chronotropic effects may occur when metoprolol is given together with calcium antagonists. Verapamil and diltiazem reduce metoprolol clearance (see <i>Warnings and Precautions</i>). |
| Calcium channel blockers (oral use) | CT | Additive reduction in myocardial contractility | Concomitant administration of a beta-adrenergic antagonist with a calcium channel blocker may produce an additive reduction in myocardial contractility due to negative chronotropic and inotropic effects. Patients taking an oral calcium channel blocker of the verapamil type in combination with JAMP-METOPROLOL-L should be closely monitored. |
| CYP2D6 inhibitors | CT | \uparrow plasma concentration of metoprolol | Potent inhibitors of this enzyme may increase the plasma concentration of metoprolol. Strong inhibition of CYP2D6 would result in the change of phenotype into poor metabolizer (see <i>Clinical Pharmacology</i>). Caution should therefore be exercised when coadministering potent CYP2D6 inhibitors with metoprolol. |

| Metoprolol | Ref | Effect | Clinical comment |
|--------------------------------|------------|--|---|
| | | | Known clinically significant potent inhibitors of CYP2D6 are antidepressants such as fluvoxamine, fluoxetine, paroxetine, sertraline, bupropion, clomipramine, desipramine antipsychotics such as chlorpromazine, fluphenazine, haloperidol, thioridazine, antiarrhythmics such as quinidine or propafenone, antiretrovirals such as ritonavir, antihistamines such as diphenhydramine, antimalarials such as hydroxychloroquine or quinine, antifungals such as terbinafine. |
| Digitalis glycosides | C | Excessive bradycardia and/or ↑ in atrioventricular conduction time | Concurrent use of digitalis glycosides may result in excessive bradycardia and/or increase in atrioventricular conduction time. Monitoring heart rate and PR interval is recommended. |
| Dipyridamole | C | Careful monitoring of heart rate | In general, administration of a β-blocker should be withheld before dipyridamole testing, with careful monitoring of heart rate following the dipyridamole injection. |
| Ergot alkaloid | C | ↑vasoconstrictive action of ergot alkaloids | Concomitant administration with β-blockers may enhance the vasoconstrictive action of ergot alkaloids. |
| Fingolimod | CT | bradycardia | Concomitant administration of beta blockers with other drugs known to decrease heart rate such as sphingosine-1-phosphate receptor modulators (e.g. fingolimod) may result in additive heart rate lowering effects and is not recommended. Where such coadministration is considered necessary, appropriate monitoring at treatment initiation, i.e. at least overnight monitoring, is recommended. |
| Hepatic Enzyme-Inducers | CT | Influence plasma level of metoprolol | Hepatic enzyme-inducing substances may exert an influence on the plasma level of metoprolol. The plasma concentration of metoprolol is lowered by rifampicin. |
| Hydralazine | C | ↑ concentrations of metoprolol | Concomitant administration of hydralazine may inhibit presystemic metabolism of metoprolol leading to increased concentrations of metoprolol. |

| Metoprolol | Ref | Effect | Clinical comment |
|--|------------|---|--|
| Inhalation anesthetics | C | ↑cardiodepression of certain anesthetics | β-blockers enhance the cardiodepression produced by certain anesthetics (see <i>Warnings and Precautions-Patients Undergoing Surgery</i>). |
| Lidocaine | C | ↓ clearance of lidocaine | Metoprolol may reduce the clearance of lidocaine. |
| MAO Inhibitors and Adrenergic Neuron Blockers | C | ↓ sympathetic activity | Closely monitor patients receiving MAO inhibitors or catecholamine-depleting drugs (such as reserpine or guanethidine). The added β-adrenergic- blockade of metoprolol may excessively reduce sympathetic activity. JAMP-METOPROLOL-L should not be combined with other β-blockers. |
| Nitroglycerin | C | ↑ hypotensive effect of metoprolol tartrate | Nitroglycerin may enhance the hypotensive effect of metoprolol tartrate. |
| NSAIDs | C | ↓ antihypertensive effect of β-blockers | Concomitant administration of non-steroidal anti-inflammatory drugs including COX-2 inhibitors with a beta-blocker may decrease the antihypertensive effect of β-blockers, possibly as a result of the inhibition of renal prostaglandin synthesis and sodium and fluid retention caused by non-steroidal anti-inflammatory drugs. |
| Oral Antidiabetics drugs and insulin | C | ↑ blood pressure associated with severe bradycardia | β-blockers may interfere with the usual hemodynamic response to hypoglycemia and produce a rise in blood pressure associated with severe bradycardia. The dosage of oral antidiabetics may have to be readjusted in patients receiving β-blockers (see <i>Warnings and Precautions</i>). |

| Metoprolol | Ref | Effect | Clinical comment |
|--|-----|---|--|
| Prazosin (selective alpha-1-adrenergic antagonist) | C | ↑ acute postural hypotension | The acute postural hypotension that can follow the first dose of prazosin may be increased in patients already taking a β blocker, including JAMP-METOPROLOL-L. |
| Sympathomimetics | C | Hypertension | Concomitant administration of sympathomimetic drugs such as adrenaline, noradrenaline, isoprenaline, ephedrine, phenylephrine, phenylpropanolamine, and xanthine derivatives (including antitussives or nose and eye drops) with a β-blocker may enhance the pressor response resulting in hypertension due to mutual inhibition of therapeutic effects. |
| Aldesleukin or other drugs known to decrease blood pressure | T | ↑ hypotensive effect of metoprolol tartrate | Concomitant administration of beta-blockers with other drugs known to decrease blood pressure such as aldesleukin may result in an enhanced hypotensive effect. |

Drug-Food Interactions

Food enhances the bioavailability of an oral dose of metoprolol by approximately 20-40%. Indeed, food intake affects the pharmacokinetics of metoprolol leading to increased exposure (AUC) and a higher maximum plasma concentration (C_{max}) (see *Action and Clinical Pharmacology*). Hence, in order to minimize the effect variations within the individual, it is recommended that the drug should be administered with or immediately following meals.

In one clinical study with metoprolol immediate release formulation, it was found that C_{max} and AUC were higher by about 32% and 38%, respectively, when administered after standard breakfast as compared to fasting condition. The study recommended that the drug should be administered with or immediately following meals to minimize the variations within an individual.

Drug-Herb Interactions

The interaction of metoprolol with herbal medications or supplements has not been studied.

Drug-Laboratory Interactions

No data suggest that metoprolol interferes with laboratory tests.

Drug-Lifestyle Interactions

Dizziness, fatigue or visual impairment may occur during treatment with JAMP-METOPROLOL-L (see *Adverse Drug reactions*) and may adversely affect the patient's ability to drive or use machines.

DOSAGE AND ADMINISTRATION

Dosing Considerations

- **Special Note to Pharmacist:** Splitting JAMP-METOPROLOL-L tablets is not recommended since the film coating will be damaged.

Recommended Dose and Dosage Adjustment

Hypertension

JAMP-METOPROLOL-L (Metoprolol Tartrate Tablets USP) is usually used in conjunction with other antihypertensive agents, particularly a thiazide diuretic, but may be used alone (see *Indications*).

The dose must always be adjusted to the individual requirements of the patient, in accordance with the following guidelines.

JAMP-METOPROLOL-L treatment should be initiated with doses of 50 mg b.i.d. If an adequate response is not seen after one week, dosage should be increased to 100 mg b.i.d. In some cases the daily dosage may need to be increased by further 100 mg increments at intervals of not less than two weeks up to a maximum of 200 mg b.i.d., which should not be exceeded. The usual maintenance dose is within the range of 100-200 mg daily.

When JAMP-METOPROLOL-L is combined with another antihypertensive agent which is already being administered, JAMP-METOPROLOL-L should be added initially at a dose of 50 mg b.i.d. After one or two weeks the daily dosage may be increased if required, in increments of 100 mg, at intervals of not less than two weeks, until adequate blood pressure control is obtained.

Given the interactions of JAMP-METOPROLOL-L with food, it is recommended that the drug should be administered with or immediately following meals (see *Action and Clinical Pharmacology- Pharmacokinetics, Drug Interactions-Drug-Food interactions*).

Angina Pectoris

The recommended dosage range for JAMP-METOPROLOL-L in angina pectoris is 100-400 mg per day in divided doses. Treatment should be initiated with 50 mg b.i.d. for the first week. If response is not adequate, the daily dosage should be increased by 100 mg for the next week. The usual maintenance dose is 200 mg/day. The need for further increases should be closely monitored at weekly intervals and the dosage increased in 100 mg increments to a maximum of 400 mg/day in two or three divided doses. JAMP-METOPROLOL-L dose of 400 mg/day should not be exceeded.

Myocardial Infarction

In addition to the usual contraindications:

ONLY PATIENTS WITH SUSPECTED ACUTE MYOCARDIAL INFARCTION WHO MEET THE FOLLOWING CRITERIA ARE SUITABLE FOR THERAPY AS DESCRIBED BELOW:

| | |
|---------------------------------|-----------------------|
| Systolic Blood Pressure | ≥100 mmHg |
| Heart Rate * | ≥ 45 beats per minute |
| PR Interval | < 0.24 seconds |
| Rales* | < 10 cm |
| Adequate peripheral circulation | |

*Extreme caution should be exercised when giving intravenous metoprolol to patients with heart rate between 45 and 60 and/or pulmonary rales less than 10 cm.

Therapy should be discontinued in patients if the heart rate drops below 45 or the systolic blood pressure drops below 100 mmHg.

Early Treatment

During the early phase of definite or suspected acute myocardial infarction, treatment with metoprolol can be initiated as soon as possible after the patient's arrival in the hospital. Such treatment should be initiated in a coronary care or similar unit immediately after the patient's hemodynamic condition has stabilized.

Treatment in this early phase should begin with the intravenous administration of three bolus injections of 5 mg of metoprolol each. The injections should be given at approximately 2- minute intervals. During the intravenous administration of metoprolol, blood pressure, heart rate, and electrocardiogram should be carefully monitored. If any of the injections are associated with adverse cardiovascular effects, intravenous administration should be stopped immediately and the patient should be observed carefully and appropriate therapy instituted.

In patients who tolerate the full intravenous dose (15 mg), JAMP-METOPROLOL-L tablets, 50 mg every 6 hours, should be initiated 15 minutes after the last intravenous dose and continued for 48 hours. Thereafter, patients should receive a maintenance dosage of 100 mg twice daily (see *Late Treatment* below).

Patients who appear not to tolerate the full intravenous dose should be started on either 25 mg or 50 mg every 6 hours (depending on the degree of intolerance) 15 minutes after the last intravenous dose or as soon as their clinical condition allows. In patients with severe intolerance, treatment with metoprolol should be discontinued (see *Warnings and Precautions*).

Late Treatment (For proven myocardial infarction patients only)

Patients with contraindications to treatment during the early phase of myocardial infarction, patients who appear not to tolerate the full early treatment, and patients in whom the physician wishes to delay therapy for any other reason should be started on JAMP-METOPROLOL-L tablets, 100 mg twice daily, as soon as their clinical condition allows. Treatment can begin within 3-10 days of the acute event. Therapy should be continued for at least 3 months. Although the efficacy of treatment with JAMP-METOPROLOL-L beyond 6 months has not been conclusively established data from studies with other β -blockers suggest that the treatment should be continued for 1-3 years.

Special populations

Pediatric patients

No pediatric studies have been performed. The safety and efficacy of metoprolol in pediatric patients have not been established.

Renal impairment

No dose adjustment of JAMP-METOPROLOL-L is required in patients mild to moderate renal impairment. Caution and regular monitoring of renal function are required in patients with severe renal impairment (see *Action and Clinical Pharmacology-Pharmacokinetics-Special populations*).

Hepatic impairment

JAMP-METOPROLOL-L blood levels are likely to increase substantially in patients with mild to moderate hepatic impairment. Therefore, JAMP-METOPROLOL-L should be initiated at low doses with cautious gradual dose titration according to clinical response and safety monitoring. Patients with severe hepatic impairment should be treated with caution i.e. lower initial and maintenance doses as well as regular monitoring of hepatic function, as they are more sensitive to therapeutic effects/adverse effects of drugs (see *Action and Clinical Pharmacology-Pharmacokinetics-Special populations*).

Geriatric patients (>65 years)

JAMP-METOPROLOL-L should be given with caution in geriatric patients due to increased likelihood of adverse events. Lower starting and maintenance doses and safety monitoring are recommended (see *Action and Clinical Pharmacology-Pharmacokinetics-Special populations*).

OVERDOSAGE

Symptoms

The most common signs to be expected with overdosage of a β -adrenoreceptor agent are hypotension, bradycardia, congestive heart failure, myocardial infarction, bronchospasm and hypoglycemia. Atrioventricular block, cardiogenic shock and cardiac arrest may develop. In addition, impairment of consciousness (or even coma), convulsions, nausea, vomiting and cyanosis and death may occur.

Concomitant ingestion of alcohol, antihypertensives, quinidine, or barbiturates aggravate the signs and symptoms.

The first manifestations of overdosage set in 20 minutes to 2 hours after drug administration.

Management

If overdosage occurs, in all cases therapy with JAMP-METOPROLOL-L (Metoprolol Tartrate Tablets USP) should be discontinued, the patient hospitalized and observed closely. Remove any drug remaining in the stomach by induction of emesis or gastric lavage.

Other clinical manifestations of overdose should be managed symptomatically based on modern methods of intensive care.

For management of a suspected drug overdose, contact your regional Poison Control Centre.

Bradycardia and Hypotension: Initially 1-2 mg of atropine sulfate should be given intravenously. If a satisfactory effect is not achieved, norepinephrine or dopamine may be administered after preceding treatment with atropine. (*See Precaution* concerning the use of epinephrine in β -blocked patients.) In case of hypoglycemia glucagon (1-10 mg) can be administered.

Heart Block (second- or third- degree): Isoproterenol or transvenous cardiac pacemaker.

1. Congestive Heart Failure: Conventional therapy.
2. Bronchospasm: Intravenous aminophylline or a β_2 -agonist.
3. Hypoglycemia: Intravenous glucose.

It should be remembered that JAMP-METOPROLOL-L is a competitive antagonist of isoproterenol and hence large doses of isoproterenol can be expected to reverse many of the effects of excessive doses of JAMP-METOPROLOL-L. However, the complications of excess isoproterenol, e.g. hypotension and tachycardia, should not be overlooked.

ACTION AND CLINICAL PHARMACOLOGY

Mechanism of Action

Metoprolol tartrate is a β -adrenergic receptor-blocking agent. *In vitro and in vivo* animal studies have shown that it has a preferential effect on the β_1 -adrenoreceptors, chiefly located in cardiac muscle. This preferential effect is not absolute, however, and at higher doses, metoprolol also inhibits β_2 -adrenoreceptors, chiefly located in the bronchial and vascular musculature. Metoprolol has no membrane-stabilizing or partial agonism (intrinsic sympathomimetic) activities. It is used in the treatment of hypertension, angina pectoris and to reduce mortality in patients with myocardial infarction.

The mechanism of the antihypertensive effect has not been established. Among the factors that may be involved are:

- a) competitive ability to antagonize catecholamine-induced tachycardia at the β -receptor sites in the heart, thus decreasing heart rate, cardiac contractility and cardiac output;
- b) inhibition of renin release by the kidneys;
- c) inhibition of the vasomotor centres.

By blocking catecholamine-induced increases in heart rate, in velocity and extent of myocardial contraction, and in blood pressure, metoprolol reduces the oxygen requirements of the heart at any given level of effort, thus making it useful in the long-term management of angina pectoris. However, in patients with heart failure, β -adrenergic receptor blockade may increase oxygen requirements by increasing left ventricular fiber length and end-diastolic pressure (preload).

The mechanisms involved in reducing mortality in patients with acute myocardial infarction are

not fully understood.

Pharmacodynamics

Significant β -blocking effect (as measured by reduction of exercise heart rate) occurs within one hour after oral administration, and its duration is dose-related. For example, a 50% reduction of the maximum effect after single oral doses of 20, 50 and 100 mg occurred at 3.3, 5.0 and 6.4 hours, respectively, in normal subjects. After repeated oral dosages of 100 mg twice daily, a significant reduction in exercise systolic blood pressure was evident at 12 hours.

Pharmacokinetic and pharmacodynamic relationship

Following intravenous administration of metoprolol, the half-life of the distribution phase is approximately 12 minutes; the urinary recovery of unchanged drug is approximately 10%. When the drug was infused over a 10-minute period, in normal volunteers, maximum β -blockade was achieved at approximately 20 minutes. Doses of 5 mg and 15 mg yielded a maximal reduction in exercise-induced heart rate of approximately 10% and 15%, respectively. The effect on exercise heart rate decreased linearly with time at the same rate for both doses, and disappeared at approximately 5 hours and 8 hours for the 5 mg and 15 mg doses, respectively.

Equivalent maximal β -blocking effect is achieved with oral and intravenous doses in the ratio of approximately 2.5:1.

There is a linear relationship between the log of plasma levels and reduction of exercise heart rate. However, antihypertensive activity does not appear to be related to plasma levels. Because of variable plasma levels attained with a given dose and lack of a consistent relationship of antihypertensive activity to dose, selection of proper dosage requires individual titration.

In several studies of patients with acute myocardial infarction, intravenous followed by oral administration of metoprolol caused a reduction in heart rate, systolic blood pressure, and cardiac output. Stroke volume, diastolic blood pressure, and pulmonary artery end-diastolic pressure remained unchanged.

The SR formulation produced lower peak metoprolol plasma concentrations than the regular tablets in studies with volunteers. Between 4 to 6 hours, both concentration curves were similar. During the 8 to 24 hour period concentrations were higher with the SR tablets.

Pharmacokinetics

The drug is available in racemic form and it exhibits stereo-specific pharmacokinetics.

Absorption: In humans, following oral administration of conventional tablet, metoprolol is rapidly and almost completely absorbed from the gastrointestinal tract. The drug is absorbed evenly throughout gastrointestinal tract. Plasma levels following oral administration, however, approximate 50% of levels following intravenous administration, indicating about 50% first-pass metabolism.

Inter-subject plasma levels achieved are highly variable after oral administration, although they show good reproducibility within each individual. Peak plasma concentrations are attained after

approximately 1.5-2 hours with conventional metoprolol formulations, and after approximately 4-5 hours with slow-release formulations. Following repeated oral administration, the percentage of the dose systemically available is higher than after a single dose and also increases dose dependently. Only a small fraction of the drug (about 12%) is bound to human serum albumin.

Distribution: Metoprolol is rapidly and extensively distributed to the extra-vascular tissue. The mean volume of distribution is 3.2 to 5.6 L/kg. The apparent volume of distribution at steady-state (V_{ss}) in extensive metabolizers (4.84 L/kg) is almost 2-fold higher than that of poor metabolizers (2.83 L/kg). At therapeutic concentrations, approximately 12 % of the active ingredient in JAMP-METOPROLOL-L tablets (metoprolol tartrate tablets USP) is bound to human serum proteins. Metoprolol crosses the placenta and is found in breast milk (see **Warnings and Precautions- Nursing Women**).

Biotransformation / Metabolism: Metoprolol is not a significant P-glycoprotein substrate but is extensively metabolised by enzymes of the cytochrome P450 system in the liver. The oxidative metabolism of metoprolol is under genetic control with a major contribution of the polymorphic cytochrome P450 isoform 2D6 (CYP2D6), which causes inter-individual variability in pharmacokinetics and pharmacodynamics of metoprolol.

Indeed, the accumulation of metoprolol leads to high levels of the drug in plasma in poor metabolizers (PMs), which are associated with higher intensity of therapeutic effects, an increase in duration of action and an increase in the occurrence and severity of AEs as compared to extensive metabolizers (EMs).

Metabolism & Dose-proportionality: Metoprolol exhibits saturable pre-systemic metabolism leading to non-proportionate increase in exposure with increased dose. However, dose proportionate pharmacokinetics is expected with extended release formulations.

Food enhances the bioavailability of an oral dose of metoprolol by approximately 20-40%. Indeed, food intake affects the pharmacokinetics of metoprolol leading to increased exposure (AUC) and a higher maximum plasma concentration (C_{max}) (see **Drug Interactions - Drug Food Interactions**).

In one clinical study with metoprolol immediate release formulation, it was found that C_{max} and AUC were higher by about 32% and 38%, respectively, when administered after standard breakfast as compared to fasting condition. The study recommended that the drug should be administered with or immediately following meals to minimize the variations within an individual.

Elimination: Elimination is mainly by biotransformation in the liver, and the plasma half-life averages 3.5 hours (range: 1 [in EMs] to 9 hours [in PMs]). The total clearance rate of an intravenous dose is approximately 1L/min and the protein binding rate is approximately 10%. Less than 5% of an oral dose of metoprolol is recovered unchanged in the urine; the rest is excreted by the kidneys as metabolites that appear to have no clinical significance. Following single oral administration of 100 mg metoprolol the median clearance were 31, 168, and 367 L/h in poor metabolizers, extensive metabolizers, and ultra-rapid metabolizers, respectively.

Special Populations and Conditions

Elderly: The elderly population show higher plasma concentrations of metoprolol (up to 28%AUC increase in elderly patients as compared to young healthy volunteers) as a combined result of a decreased elimination of metoprolol and the metabolite α -hydroxy-metoprolol and a decreased hepatic blood flow due to age-related physiological changes. In addition, time to reach peak concentration, T_{max} , was significantly longer in the elderly population. Hence, it is recommended to initiate therapy with lower doses in this group and safety monitoring may be recommended.

Ethnic sensitivity: The oxidative metabolism of metoprolol is under genetic control with a major contribution of the polymorphic cytochrome P450 isoform 2D6 (CYP2D6). There are marked ethnic differences in the prevalence of the poor metabolizers (PM) phenotype. Approximately 7% of Caucasians and less than 1% Orientals are PMs. CYP2D6 poor metabolizers exhibit 5-fold higher plasma concentrations of metoprolol than extensive metabolizers with normal CYP2D6 activity.

Hepatic impairment: Since the drug is primarily eliminated by hepatic metabolism, hepatic impairment impacts the pharmacokinetics of metoprolol. The elimination half-life of metoprolol is considerably prolonged, depending on severity (up to 7.2 h), in patients with liver impairment. Hence, dose adjustment and safety monitoring are advised in patients with mild to moderate hepatic impairment. Patients with severe hepatic impairment should be treated with caution, i.e. lower initial and maintenance doses as well as regular monitoring of hepatic function, as they are more sensitive to therapeutic effects/adverse effects of drugs.

Renal impairment: Pharmacokinetics of metoprolol in patient with renal impairment did not differ to a clinically significant degree from normal subjects. However, there is accumulation of one of its less active metabolite in patients with a creatinine clearance below 5 mL/min. Since the resulting metabolite accumulation has no significant effect on the β -blocking effects, metoprolol dosing does not need to be altered in patient with mild to moderate renal impairment. Caution is advised in the use of a beta-blocker in patients with severe renal impairment and safety monitoring is advised in these patients.

STORAGE AND STABILITY

JAMP-METOPROLOL-L tablets must be stored between 15°-30°C. Protected from moisture and light.

DOSAGE FORMS, COMPOSITION AND PACKAGING

JAMP-METOPROLOL-L (metoprolol) tablets

| | 25mg Tablets | 50mg Tablets | 100mg Tablets |
|---------------------------------------|--|--|--|
| Description | White colored caplet shaped film coated biconvex tablet with "ML" and "25" engraved on either side of break-line and "JP" engraved on other side. | Pink colored caplet shaped film coated biconvex tablet with "ML" and "50" engraved on either side of break-line and "JP" engraved on other side. | Light blue colored caplet shaped film coated biconvex tablet with "ML" and "100" engraved on either side of break-line and "JP" engraved on other side |
| Colour | White | Pink | Blue |
| Metoprolol tartrate content | 25mg | 50 mg | 100mg |
| Non-medicinal ingredients | Microcrystalline cellulose, lactose, Sodium, starch glycolate, povidone, colloidal silicone dioxide, magnesium stearate. In addition to these ingredients the 25mg tablet contains Opadry 03F580000 White (HPMC 2910/Hypromellose, titanium dioxide, macrogol/Polyethylene glycol, talc), the 50mg tablet contains Opadry 03F540232 Pink (HPMC 2910/Hypromellose, titanium dioxide, macrogol/Polyethylene glycol, talc, FD&C Red #40, D&C Yellow #10, D&C Red #30) and the 100mg tablet contains Opadry 03F505098 Blue (HPMC 2910/Hypromellose, titanium dioxide, macrogol/Polyethylene glycol, talc, FD&C Blue #2, FD&C Blue #1). | | |
| Availability | HDPE Bottles of 100 or 500 tablets | HDPE Bottles of 100 or 500 or 1000 tablets | HDPE Bottles of 100 or 500 or 1000 tablets |
| Stability and Storage Recommendations | Stored between 15° to 30°C. Protect from moisture and light. | | |

PART II: SCIENTIFIC INFORMATION

PHARMACEUTICAL INFORMATION


Drug Substance

Proper name: Metoprolol Tartrate

Chemical name: 2-propanol,1-[4-(2-methoxyethyl)phenoxy]-3-[(1-methylethyl)amino]-, (±)-, [R-(R*,R*)]-2,3-dihydroxybutanedioate (2:1) (salt).

Molecular formula and molecular mass: $(C_{15}H_{25}NO_3)_2 \cdot C_4H_6O_6$; 684.83 g/mol

Structural formula:



Physicochemical properties: Description: Colorless, odorless, crystalline powder with a bitter taste.

Solubility: Very soluble in water at 20°C pH: 6.7 (2% water solution)

CLINICAL TRIALS

Comparative Bioavailability Studies

A double-blind, randomized, two-period, two-treatment, two-sequence, crossover, balanced, single (1 x 100 mg) dose, oral bioequivalence study of JAMP-METOPROLOL-L (Metoprolol Tartrate Tablets), 100 mg (by JAMP Pharma Corporation) and LOPRESOR® (metoprolol tartrate) Tablets, 100 mg by Novartis Pharmaceuticals Canada, Inc.) was conducted in healthy, adult, Asian male subjects under fasting conditions. The results from the 29 subjects who completed the study are summarized below.

SUMMARY TABLE OF COMPARATIVE BIOAVAILABILITY DATA

| Metoprolol (1 x 100 mg) From measured data Geometric Mean Arithmetic Mean (CV %) | | | | |
|--|--------------------------|--------------------------|----------------------------|-------------------------|
| Parameter | Test* | Reference [†] | % Ratio of Geometric Means | 90% Confidence Interval |
| AUC _T (ng·h/mL) | 687.12 769.8 (57.81) | 646.26 720.98 (56.98) | 106.30 | 102.20-110.60 |
| AUC _I (ng·h/mL) | 692.10 774.79 (57.67) | 651.10 726.05 (56.86) | 106.30 | 102.20-110.50 |
| C _{max} (ng/mL) | 119.66 128.33 (40.12) | 115.74 122.41 (34.67) | 103.40 | 97.50-109.60 |
| T _{max} [§] (h) | 1.50 (0.75-3.00) | 2.00 (0.75–3.00) | | |
| T _½ (h) ^ε | 3.71 (21.50) | 3.77 (21.93) | | |

* JAMP-METOPROLOL-L (metoprolol tartrate) Tablets, 100 mg (by JAMP Pharma Corporation)

† LOPRESOR® (metoprolol tartrate) Tablets, 100 mg (by Novartis Pharmaceuticals Canada Inc.) were purchased in Canada

§ T_{max} is expressed as the median (range) only

ε T_½ is expressed as the arithmetic mean (CV%) only

DETAILED PHARMACOLOGY

Effect on the Cardiovascular System

Metoprolol produced dose-dependent reductions in heart rate and contractile force responses to sympathetic nerve stimulation in the anaesthetized cat. The ED₅₀ value for blockade of the chronotropic response to nerve stimulation was approximately 7 times less than that for isoproterenol stimulation. Metoprolol reduced the heart rate in conscious dogs at rest and during exercise. With the exception of PR interval prolongation, ECG complexes were not changed.

In anaesthetized cats, intravenous doses up to 2.0 mg/kg did not significantly influence the pressor response to intravenous epinephrine.

In anaesthetized cats, intravenously-administered metoprolol antagonized the hind limb vasodilating response to intra-arterial isoproterenol in much higher doses (ED₅₀ 5 mg/kg) than required to block the increase in chronotropic response (ED₅₀ 0.4 mg/kg) or increase in contractile force (ED₅₀ 0.2 mg/kg).

In 5 healthy volunteers, intravenously-administered 10 mg doses of metoprolol reduced exercise-induced tachycardia by 13% and systolic blood pressure during exercise by 13%. The decrease in mean blood pressure after epinephrine was abolished by metoprolol, whereas the increase in systolic blood pressure was reduced by 50%; vascular resistance in the forearm was unchanged after metoprolol.

In healthy volunteers, intravenous metoprolol 0.15 mg/kg significantly lowered cardiac output by 1.3 litre/min. at rest, and 3.6 litre/min. during exercise. The mean decreases in heart rate were 9 and 16 beats/min. during rest and exercise, respectively. Right atrial pressure was significantly increased during rest and exercise. Oxygen consumption was not significantly influenced by drug administration. Significant increases in the calculated arteriovenous oxygen differences were observed (6 and 20 mL/litre at rest and during exercise, respectively).

Studies in reserpinized cats showed that metoprolol was devoid of β -receptor stimulating (intrinsic) activity. In cumulative doses up to 0.85 mg/kg, the drug did not significantly influence heart rate or contractile force.

A single oral dose of 40 mg of metoprolol administered to 17 anginal patients 90 minutes before testing, increased total work performed from 5994 to 8462 k.p.m. (40%). Times to onset of pain and appearance of ST depression were similarly increased from 11.8 to 16.9 minutes and 9.9 to 13.9 minutes respectively.

Effects on Pulmonary Function

The effects on specific airways resistance (S_{Raw}) of single oral doses of 100 mg of metoprolol were assessed in 6 healthy volunteers and in 12 patients with bronchial asthma. No bronchodilator was used. Metoprolol did not have a significant effect on S_{Raw} in the normal subjects, but in the asthmatic patients, S_{Raw} was significantly increased. Similar findings were observed with an 80 mg dose of propranolol.

In a controlled study, 17 patients with bronchial asthma received concomitantly a bronchodilator (terbutaline) with 50 or 100 mg b.i.d. of metoprolol. The FEV1 values fell only in the high dose group, indicating some b2-blocking effect.

Other Effects

Metoprolol showed a negligible local anaesthetic effect on the isolated sciatic nerve of the frog and in the intracutaneous wheal test in guinea pigs. The cardiostimulant effects of ouabain, glucagon and theophylline were not affected by doses of 2-3 mg/kg in the anaesthetized cat. The same dosage of metoprolol was found to be devoid of anticholinergic, ganglionic blocking, antihistaminic and alpha-receptor blocking properties in cats.

Metoprolol inhibited the increase in plasma renin activity induced by furosemide.

The effects of metoprolol on isoproterenol-stimulated metabolic effects showed inhibition of the increase in liberation of glycerol, glucose, insulin, and free fatty acids.

TOXICOLOGY

Acute Toxicity

| Species | Sex | Route | Solutions | LD₅₀(mg/kg) |
|----------------|------------|--------------|------------------|-------------------------------|
| Mouse | male | I.V. | 1% | 69.4 ± 5.1 |
| Mouse | female | I.V. | 1% | 79.9 ± 4.5 |
| Mouse | male | P.O. | 23% | 2460 ± 210 |
| Mouse | female | P.O. | 25% | 2300 ± 200 |
| Rat | male | I.V. | 5% | 71.9 ± 4.1 |
| Rat | female | I.V. | 5% | 74.3 ± 4.4 |
| Rat | male | P.O. | 50% | 4670 ± 1210 |
| Rat | female | P.O. | 50% | 3470 ± 580 |

The toxic symptoms in rats include: sedation, ataxia, piloerection, irritation, spasm, and lacrimation. Rats were unconscious before death, which occurred within 5-10 minutes after intravenous injection and 6-20 hours after oral administration.

In mice the most pronounced symptoms were: sedation, hypersensitivity, irritation, spasms, and ptosis. Convulsions were seen before death, which occurred within 5 minutes after intravenous injection. No symptoms of toxicity were detectable 24 hours after administration in surviving animals.

Long-Term Toxicity (Subacute)

| Strain Species | No. of Groups | N per Group | Dose (mg/kg) | Route | Duration | Toxic Effects |
|---------------------|---------------|--------------|--|-------|----------|---|
| Sprague-Dawley Rats | 4 | 10 M 10 F | Saline, 10, 50, 100/day (after 14 days, high dose increased to 200/day). | P.O. | 5 Wks | Slight increase in hematocrit and slight decrease in blood sugar in high-dose females. |
| Beagle Dogs | 1 | 1 M 1 F | 40 x 3 days, increased by 20/day to 140 x 6 days to 160/day. | P.O. | 3 Wks | Disturbance of balance; increased abdominal muscular tone, mydriasis, hyperemia in visible mucous membranes. One dog died at dose level of 140 mg/kg/day. |
| Beagle Dogs | 2 | 1 M | 80 b.i.d. one day; 2 days later, single dose of 100. | P.O. | 3 Days | Disturbance of balance; vomiting, prostration, dyspnea, loss of consciousness, death. |
| | | 2 F | 20 b.i.d. increased every 5 days by 20 b.i.d. up to 120 b.i.d. | P.O. | 4 Wks | Vomiting; increased salivation, tremor, ataxia. One dog died at highest dose. |
| Beagle Dogs | 4 | 1 M 1 F | 0, 5, 20, 40/day | P.O. | 4 Wks | None. |
| Beagle Dogs | 3 | 1 M 1 F | Saline, 0.5, 5 /day | I.V. | 2 Wks | Prolonged PR interval in ECG. |
| Beagle Dogs | 2 | 1 M 1 F | Saline, 5 /day | I.V. | 2 Wks | Prolonged PR interval in ECG. |

Long-Term Toxicity (Chronic)

| Strain Species | No. Of Groups | N per Group | Dose (mg/kg) | Route | Duration | Toxic Effects |
|--------------------|---------------|--------------|---|-------|----------|--|
| Sprague-Dawley Rat | 4 | 15 M 15 F | Saline, 10, 100, 200/day. High dose increased to 200/day after 13 Weeks | P.O. | 6 Months | None. |
| Beagle Dogs | One Control | 2 M 2 F | 0, 5, 20, 40 b.i.d. After 7 weeks, high dose increased to 50/b.i.d. After 3 months, intermediate dose increased to 30 b.i.d. and high dose to 80 b.i.d. | P.O. | 6 Months | Bradycardia, increased PR interval and QT interval in ECG. |
| | Three Active | 3 M 3 F | | | | |
| Beagle Dog | One Control | 6 M 6 F | 0, 10, 60 day. High level dogs received 120 on day 1, 60 on days 3 to 8; 90/day on days 9 to 22 and 105/day for balance. | P.O. | 1 Year | 2 high-dose dogs died on day 1, otherwise, none. |
| | Three Active | 6 M 6F | | | | |

Teratology and Reproduction Studies

Rat: (Sprague-Dawley strain) Doses of 10, 50 and 200 mg/kg were administered orally to groups of 20 pregnant rats on days 6-15 of gestation. Treatment with metoprolol did not adversely affect any of the parameters studied.

Rabbit: (New Zealand White strain) Doses of 5, 12.5 and 25 mg/kg were administered orally to groups of 20 pregnant rabbits on days 6-18 of gestation. Parameters studied were not significantly affected, although litter size was lower and fetal loss higher in the high dose group. The incidence of fetal abnormality was unaffected by treatment.

Rat: (Sprague-Dawley strain) Doses of 10, 50 and 200 mg/kg were administered orally to groups of 50 rats from day 15 of gestation, through lactation to 21 days postpartum. Parameters studied in litter and parent animals were not adversely affected.

Rat: (Charles River CD strain) Doses of 50 and 500 mg/kg were administered orally to groups of 10 male and 20 female rats. Males were treated for 63 days prior to mating and during the mating period. The females were treated for 14 days prior to mating, during mating and throughout the gestation and lactation periods to 21 days postpartum, with an interim sacrifice at day 13 of gestation. The only significant finding in this study was a slight reduction of intrauterine growth in rats at 50 and 500 mg/kg/day and a higher frequency of stillbirths in the high dose group.

Carcinogenicity Studies

Metoprolol was administered to 3 groups of 60 male and 60 female Charles River Sprague-Dawley rats at dietary levels of 50, 200 and 800 mg/kg per day for 78 weeks. A fourth group received 2-AAF (positive control) and the fifth was the negative control group. The incidence of nodules and masses observed at necropsy were comparable between the treated and control groups. The only histopathological changes noted were an increased incidence of impaction of pulmonary alveoli by septal cells in the high and intermediate metoprolol-treated groups. The strain of rats was susceptible to the known carcinogen 2-AAF; a statistically higher incidence of neoplasms, primarily hepatomas, was present.

A similar study in Swiss albino mice at doses of 75, 150 and 750 mg/kg per day for 78 weeks showed that the tumors were distributed with equal frequency in the treated and control groups. The strain was susceptible to the known carcinogen.

REFERENCES

1. Ablad B, Borg KO, Carlsson E, Ek L, Johnsson G, Malmfors T, and Regardh C-G. A survey of the pharmacological properties of metoprolol in animals and man. *Acta Pharmacol Toxicol* 1975; 36 (suppl V): 7-23.
2. Ablad B, Borjesson I, Carlsson E, and Johnsson G. Effects of metoprolol and propranolol on some metabolic responses to catecholamines in the anaesthetized dog. *Acta Pharmacol Toxicol* 1975; 36 (suppl V): 85-95.
3. Adolfsson L, Areskog N-H, Furburg C, and Johnsson G. Effects of single doses of alprenolol and two cardio-selective β -blockers (H 87/07 and H 93/26) on exercise-induced angina pectoris. *Eur J Clin Pharmacol* 1974; 7 (2): 111-118.
4. Biemann P, Leduc G, Jequier J-C, Davignon J, and Cartwright K. Changes in the lipoprotein composition after chronic administration of metoprolol and propranolol in hypertriglyceridemic-hypertensive subjects. *Curr Ther Res* 1981; 30 (6): 956-967.
5. Bodin N-O, Flodh H, Magnussen G, Malmfors T, and Nyberg J-A. Toxicological studies on metoprolol. *Acta Pharmacol Toxicol* 1975; 36 (suppl V): 96-103.
6. Borer JS, Comerford MB, and Sowton E. Assessment of metoprolol, a cardioselective β -blocking agent, during chronic therapy in patients with angina pectoris. *J Int Med Res* 1976; 4: 15-22.
7. Borg KO, Carlsson E, Hoffmann KJ, Jönsson TE, Thorin H, Wallin B. Metabolism of metoprolol-(3H) in man, the dog and the rat. *Acta Pharmacol Toxicol* 1975;36 Suppl V:125-35.
8. Boyle DMcC, Barber JM, McIlmoyle EL, Salathia KS, Evans AE, Cran G, Elwood JH, and Shanks RG. Effect of very early intervention with metoprolol on myocardial infarct size. *Br Heart J* 1983; 49: 229-233.
9. Comerford MB, and Besterman EMM. An eighteen months' study of the clinical response to metoprolol, a selective β_1 -receptor blocking agent, in patients with anginal pectoris. *Postgrad Med J* 1976; 52 (610): 481-486.
10. Ekelund L-G, Johnsson G, Melcher A, and Oro L. Effects of cedilanid-D in combination with metoprolol on exercise tolerance and systolic time intervals in angina pectoris. *Am J Cardiol* 1976; 37 (4): 630-634.
11. Epstein SE, and Braunwald E. β -adrenergic receptor blocking drugs. Mechanisms of action and clinical applications. *New Eng J Med* 1966; 275: 1106-1112, 1175-1183
12. Fitzgerald JD. The role of β -adrenergic blockade in acute myocardial ischaemia. IN: Oliver MF, Julian DG, and Donald KW (eds). *Effect of acute ischaemia on myocardial function*. Churchill Livingstone, Edinburgh and London, 1972, pp. 321-353.
13. Formgren H. The effect of metoprolol and practolol on lung function and blood pressure in hypertensive asthmatics. *Br J Clin Pharmacol* 1976; 3 (6): 1007-1014 The Göteborg Metoprolol Trial in Acute Myocardial Infarction. *Am J Cardiol* 1984; 53 (13): 1D-50D.

14. Hansten PD, Horn JR, editors. Principles of beta-adrenergic blocker drug interactions 79-82. Anti- arrhythmic drug interactions 153-8, 162-3, 170-1, 188-9. Antidiabetic drug interactions 373-5. Beta-adrenergic blocker drug interactions 435-57. Drug interactions & updates. Vancouver: Applied Therapeutics Inc., 1993.
15. Herlitz J, Elmfeldt D, Hjalmarson A, Holmberg S, Malek I, Nyberg G, Ryden L, Swedberg K, Vedin A, Waagstein F, Waldenstrom A, Waldenstrom J, Wedel H, Wilhelmsen L, and Wilhelmsson C. Effect of metoprolol on indirect signs of the size and severity of acute myocardial infarction. *Am J Cardiol* 1983; 51: 1282-1288.
16. Hjalmarson A, Ariniego R, Herlitz J, et al. Limitation of infarct size in man by the β_1 -blocker metoprolol. *Circulation* 1979; 59 and 60 (Suppl II): 164 (Abstr.)
17. Hjalmarson A, Elmfeldt D, Herlitz J, et al. Effect on mortality of metoprolol in acute myocardial infarction. A double-blind randomized trial. *Lancet* 1981 (Oct. 17); 823-827.
18. Hjalmarson A, Herlitz J, Holmberg S, Ryden L, Swedberg K, Vedin A, Waagstein F, Waldenstrom A, Waldenstrom J, Wedel H, Wilhelmsen L, and Wilhelmsson C: The Göteborg Metoprolol Trial; Effects on mortality and morbidity in acute myocardial infarction. *Circulation* 1983; 67 (suppl I): I26- I32.
19. Imhof P: Les bêta-bloquants et le sport. *Médecine et Hygiène* 1979; 1348.
20. Johnsson G. Influence of metoprolol and propranolol on hemodynamic effects induced by adrenaline and physical work. *Acta Pharmacol Toxicol* 1975; 36 (suppl V): 59-68.
21. Johnsson G, Regardh C-G, and Sölvell L. Combined pharmacokinetic and pharmacodynamic studies in man of the adrenergic β_1 -receptor antagonist metoprolol. *Acta Pharmacol Toxicol* 1975; 36 (suppl V): 31-44.
22. Johnsson G, Svedmyr N, and Thiringer G. Effects of intravenous propranolol and metoprolol and their interaction with isoprenaline on pulmonary function, heart rate and blood pressure in asthmatics. *Eur J Clin Pharmacol* 1975; 8: 175-180.
23. Kendall MJ, John VA, Quarterman CP, and Welling PG. A single and multiple dose pharmacokinetic and pharmacodynamic comparison of conventional and slow-release metoprolol. *Eur J Clin Pharmacol* 1980; 17: 87-92.
24. Keyrilainen O, and Uusitalo A. Effects of metoprolol in angina pectoris. A subacute study with exercise tests and a long-term tolerability study. *Acta Med Scand* 1976; 199: 491-497.
25. Lennard MS. The polymorphic oxidation of beta-adrenoceptor antagonists. *Pharmacol Ther* 1989;41:461-77.
26. Logan RL, Goldstraw PW, Kiddle GB, Flacks LM, and Watt M. Sustained release metoprolol: a comparison with conventional formulation in the treatment of hypertension. *NZ Med J* 1981; 687: 1- 2.

27. Lorimer AR, Barbour MB, and Lawrie TDV. An evaluation of the effect on resting and exercise blood pressure of some first line treatments in hypertension. *Ann Clin Res* 1983; 15: 30-34.
28. Malek I, Waagstein F, Hjalmarson A, Holmberg S, and Swedberg K. Hemodynamic effects of the cardioselective β -blocking agent metoprolol in acute myocardial infarction. *Acta Med Scand* 1978; 204: 195-201.
29. The Miami Trial Research Group. Metoprolol in acute myocardial infarction (MIAMI). A randomized placebo-controlled international trial. *Eur Heart J* 1985; 6: 199-226.
30. Olsson G, Rehnqvist N, Sjogren A, Eerhart L, and Lundman T. Long-term treatment with metoprolol after myocardial infarction: Effect on 3 year mortality and morbidity. *J Am Coll Cardiol* 1985; 5 (6): 1428-1437.
31. Quarterman CP, Kendall MJ, and Welling PG. Plasma levels and negative chronotropic effect of metoprolol following single doses of a conventional and sustained-release formulation. *Europ J Clin Pharmacol* 1979; 15: 97-103.
32. Regardh CG, Borg KO, Johansson R, Johnsson G, Palmer L. Pharmacokinetic studies on the selective β_1 -receptor antagonist metoprolol in man. *J Pharmacokinet Biopharm* 1974;2(4):347-64.
33. Regardh CG, Johnsson G. Clinical pharmacokinetics of metoprolol. *Clin Pharmacokinet* 1980;5:557-69.
34. Richards S. Cutaneous side-effects of beta-adrenergic blockers. *Aust J Derm* 1985;26:25-8.
35. Ryden L, Ariniego R, Arnman K, Herlitz J, Hjalmarson A, Holmberg S, Reyes C, Smedgard P, Svedberg K, Vedin A, Waagstein F, Waldenstrom A, Wilhelmsson C, Wedel H, and Yamamoto M. A double-blind trial of metoprolol in acute myocardial infarction. Effects on ventricular tachyarrhythmias. *N Eng J Med* 1983; 308: 614-618.
36. Salathia KS, Barber JM, McIlmoyle EL, Nicholas J, Evans AE, Elwood JH, Cran G, Shanks RG, and Boyle D McC. Very early intervention with metoprolol in suspected acute myocardial infarction. *Eur Heart J* 1985; 6: 190-198.
37. Singh BN, et al. Effects of cardioselective β adrenergic receptor blockade on specific airways resistance in normal subjects and in patients with bronchial asthma. *Clin Pharmacol Ther* 1976; 19 (5, Pt 1): 493-501.
38. Stenberg J, Wasir H, Amery A, Sannerstedt R, and Werko L. Comparative hemodynamic studies in man of adrenergic β_1 -receptor agents without (H 93/26 = metoprolol) or with (H 87/07) intrinsic sympathicomimetic activity. *Acta Pharmacol Toxicol* 1975; 36 (suppl V): 76-84.

39. Stockley IH, editor. Drug Interactions. A source book of adverse interaction their mechanisms, clinical importance and management. Amiodarone-beta blockers 96-7. Disopyramide + beta-blockers 99-100. Lignocaine (Lidocaine) + beta-blockers) 108-9. Quinidine + beta-blockers 121-2. Clonidine + beta-blockers 362-4. Prazozin + beta-blockers 383-4. Beta-blockers + cimetidine 413-4. Beta-blockers + rifampicin 427-8. Beta-blockers + sympathonimetics 429-31. Digitalis glycosides + beta-blockers 506. Hypoglycamic agents-beta blockers 548-550. Anaesthetics, general+ beta-blockers 697. Oxford: Blackwell Science, 1992.
40. Trafford JAP, McGonicle R, Bowles J, O'Neal H, Bennet-Jones DN, Halford L, and Evans R. A two year clinical evaluation of atenolol and metoprolol in the treatment of hypertension. Br J Clin Pract 1982; 36: 350-352.
41. Van Joost T, Smitt JHS. Skin reactions to propranolol and cross sensitivity to β -adrenoreceptor blocking agents. Arch Dermatol 1981; 117:600-1.
42. Waagstein F, and Hjalmarson AC. Double-blind study of the effect of cardioselective β -blockade on chest pain in acute myocardial infarction. Acta Med Scand 1975; (Suppl 587): 201-208.
43. Waal-Manning, HJ. Hypertension: Which β -blocker? Drugs 1976; 12: 412-441.
44. Wagner F, Kalusche D, Trenk D, Jähnchen E, Roskamm H. Drug interaction between propafenone and metoprolol. Br J Clin Pharmacol 1987; 24:213-20.
45. LOPRESOR[®] (metoprolol tartrate), Control # 235180, Product Monograph, Novartis Pharmaceuticals Canada Inc., Date of Revision April 14, 2020.

PART III: CONSUMER INFORMATION**Pr JAMP-METOPROLOL-L****Metoprolol Tartrate Tablets USP**

This leaflet is part III of a three-part "Product Monograph" published when JAMP-METOPROLOL-L was approved for sale in Canada and is designed specifically for Consumers. This leaflet is a summary and will not tell you everything about JAMP-METOPROLOL-L. Contact your doctor or pharmacist if you have any questions about the drug.

This information applies JAMP-METOPROLOL-L tablets.

ABOUT TO MEDICATION**What the medication is used for:**

JAMP-METOPROLOL-L is used alone or in combination with another medicine for the following conditions:

- to treat high blood pressure
- to treat angina (chest pain triggered by exercise)
- to help to protect the heart after a heart attack (myocardial infarction)

What it does:

JAMP-METOPROLOL-L is a beta-blocker. It helps to control high blood pressure or other heart-related problems.

When it should not be used:

You should not be treated with JAMP-METOPROLOL-L if you:

- are allergic to metoprolol, to any of the other ingredients in JAMP-METOPROLOL-L or to another beta-blocker
- have breathlessness, difficulty breathing when lying down, swelling of the feet or legs (signs of a heart disorder)
- have a slow or irregular heartbeat
- have sudden and oppressive chest pain (sign of heart attack)
- have very poor blood circulation in your limbs (for example, very cold, pale hands or feet, or pain in your leg muscles when you walk)
- have low blood pressure
- have non-treated tumor of the medulla of the adrenal glands (pheochromocytoma)
- have asthma or had history of difficulty breathing with wheezing or coughing
- have severe skin problems
- suffer from severe drop in blood pressure, dizziness, fast heartbeat, rapid and shallow breathing, cold clammy skin (signs of a heart disorder named cardiogenic shock).

What the medicinal ingredient is:

metoprolol tartrate.

What the nonmedicinal ingredients are:

JAMP-METOPROLOL-L tablets contain: microcrystalline cellulose lactose, sodium starch glycolate magnesium stearate, povidone, colloidal silicon dioxide. In addition to these ingredients the 25mg tablet contains Opadry 03F580000 White

(HPMC 2910/Hypromellose, titanium dioxide, macrogol/Polyethylene glycol, talc), the 50mg tablet contains Opadry 03F540232 Pink (HPMC 2910/Hypromellose, titanium dioxide, macrogol/Polyethylene glycol, talc, FD&C Red #40, D&C Yellow #10, D&C Red #30) and the 100mg tablet contains Opadry 03F505098 Blue (HPMC 2910/Hypromellose, titanium dioxide, macrogol/Polyethylene glycol, talc, FD&C Blue #2, FD&C Blue #1).

What dosage forms it comes in:

JAMP-METOPROLOL-L Tablets 25mg, 50 mg and 100 mg

WARNINGS AND PRECAUTIONS**BEFORE you use JAMP-METOPROLOL-L talk to your doctor or pharmacist if you:**

- have heart, liver or kidney disease
- have high or low levels of sugar in the blood (diabetes)
- are at risk for serious allergic reactions
- have chest pain when you are at rest
- have poor blood circulation in your limbs (for example, very cold, pale hands or feet, or pain in your leg muscles when you walk)
- have a tumor of the medulla or of the adrenal glands (pheochromocytoma), you would require an additional treatment to JAMP-METOPROLOL-L
- have allergy to or an intolerance to lactose. This applies only to JAMP-METOPROLOL-L
- have an overactive thyroid gland
- have respiratory disease such as asthma, or sometimes get breathlessness and wheezing
- have a severe syndrome named oculomucocutaneous syndrome whose signs include severe conjunctivitis (red, irritated and watery eye), skin rash and ear infection
- are pregnant, or intend to become pregnant. JAMP-METOPROLOL-L should not be used during pregnancy. Your doctor will discuss with you the potential risks of taking Jamp- Metoprolol-L during pregnancy
- are breast feeding. If your doctor decides that you must continue to take JAMP-METOPROLOL-L you should stop breast-feeding as JAMP-METOPROLOL-L passes into breast milk
- are under 18 years old.

If you need to undergo an operation where an anesthetic is used, tell your anesthetic professional that you are taking JAMP-METOPROLOL-L.

Driving and using machines: If you experience dizziness, tiredness or blurred vision during your treatment with JAMP-METOPROLOL-L, do not drive, use machinery, or perform other tasks that need full attention until you know how you respond to JAMP-METOPROLOL-L. Drinking alcohol may increase tiredness.

INTERACTIONS WITH THIS MEDICATION

Tell your doctor or pharmacist if you are taking or have recently taken any other medicines, including herbal and non-

prescription medicines. Some other medicines may interact with JAMP-METOPROLOL-L. These include:

- medicines used to treat high blood pressure,
- medicines used to treat chest pain (angina) (e.g. nitroglycerin),
- medicines used to treat irregular heartbeat (e.g. amiodarone, propafenone, quinidine, disopyramidetocainide, procainamide, ajmalineamiodarone, flecainide, digitalis glycosides such as digoxin, lidocaine),
- anaesthetics, medicines used during surgical operations,
- adrenaline or similar substances (sympathomimetics), which are found in some eye and nose drops, and in some cough medicines or remedies for the common cold (e.g. noradrenaline, isoprenaline, ephedrine, phenylephrine, phenylpropanolamine, and xanthine derivatives),
- insulin, or medicines used to treat high levels of sugar in the blood (diabetes),
- medicines used to relieve pain or inflammation (non-steroidal anti-inflammatory drugs such as COX-2 inhibitors),
- antibiotics (e.g. rifampicin),
- antivirals (e.g. ritonavir),
- antihistamines used to treat hay fever (e.g. diphenhydramine),
- medicines used to treat malaria (e.g. hydroxychloroquine or quinine),
- antipsychotic medicines (e.g. thioridazine, chlorpromazine, fluphenazine, haloperidol),
- antidepressants (e.g. fluoxetine, paroxetine, fluvoxamine, sertraline, clomipramine, desipramine or bupropion),
- MOA Inhibitors and adrenergic neuron blockers (e.g. reserpine or guanethidine),
- antifungals (e.g. terbinafine),
- ergot alkaloids, used in the prevention and treatment of migraine headaches,
- dipyridamole, used to reduce the risk of blood clots,
- alcohol.
- fingolimod, a medicine used to treat multiple sclerosis
- Aldesleukin, a medicine used to treat kidney cancer, or other medicines that may cause a decrease in blood pressure

PROPER USE OF THIS MEDICATION

Usual dose:

Follow your doctor’s instructions carefully. Do not exceed the recommended dosage.

Your doctor will tell you exactly how many tablets of JAMP-METOPROLOL-L to take based on your individual requirements.

High Blood Pressure: the usual maintenance dose is 100-200 mg daily.

Angina: the usual maintenance dose is 200 mg daily.

To Protect the Heart After a Heart Attack: the usual maintenance dose is 100 mg twice daily.

In general, the daily dosage is in the following range: 100 to 200 mg daily, either once daily (in the morning), or divided into two separate doses (one in the morning and one in the evening).

Depending on how you respond to the treatment, your doctor may suggest a higher or lower dose.

JAMP-METOPROLOL-L tablets should be swallowed whole without being chewed, preferably with or following a meal.

Do not change the dose or stop the treatment without talking to your doctor. If you stop taking JAMP-METOPROLOL-L suddenly, your condition may become worse. Your doctor may want you to reduce the dose gradually before stopping treatment altogether.

Overdose:

In case of drug overdose, contact a health care practitioner, hospital emergency department or regional Poison Control Centre immediately, even if there are no symptoms.

Some of the effects of an overdose of JAMP-METOPROLOL-L are: an abnormally slow heartbeat or an irregular heartbeat, very low blood pressure, breathlessness, difficulty breathing when lying down, swelling of the feet, loss of consciousness, blue discoloration of the lips, tongue, skin, seizures, nausea and vomiting, sudden and oppressive chest pain and death.

Missed Dose:

If you forget to take a dose of JAMP-METOPROLOL-L, take the missed dose as soon as you remember it. However, if it is almost time for the next dose, skip the missed dose and go back to your regular dosing schedule. Do not take a double dose.

SIDE EFFECTS AND WHAT TO DO ABOUT THEM

Like other medications, JAMP-METOPROLOL-L may cause some side effects. These side effects may be minor and temporary. However, some may be serious and need medical attention.

Side effects may include: sleep disturbance, fatigue and tiredness especially with activity, headache, dry mouth, nausea and vomiting, diarrhea or constipation and abdominal pain, numbness, a tingling sensation in the extremities signs of paresthesia, increased sensitivity of the skin to sun, sweating, hair loss, increased weight, and lower back pain.

Based on your health history, your doctor should take blood tests for as long as you are being treated with JAMP-METOPROLOL-L. They may monitor:

- blood sugar
- heart, liver, kidney and thyroid function
- how JAMP-METOPROLOL-L is affecting other medications that you are taking.

| SERIOUS SIDE EFFECTS, HOW OFTEN THEY HAPPEN AND WHAT TO DO ABOUT THEM | | | | |
|---|--|-------------------------------------|--------------|--|
| Symptom / effect | | Talk with your doctor or pharmacist | | Stop taking drug and immediately seek assistance |
| | | Only if severe | In all cases | |
| Common | Low Blood Pressure: feeling of lightheadedness or fainting especially when getting up from a lying or sitting position | | √ | |
| | Chest Pain | | √ | |
| | Bronchospasm: difficulty breathing with wheezing or coughing | | √ | |
| | Heart Failure: shortness of breath, leg swelling and tiredness especially with activity | | | √ |
| Uncommon | Edema: ankle swelling | √ | | |
| | Slow or irregular heartbeat (palpitations) | | √ | |
| | New or Worsening Psoriasis: skin rash (in the form of itchy rash, thickened patches of red/silver skin) | √ | | |

| SERIOUS SIDE EFFECTS, HOW OFTEN THEY HAPPEN AND WHAT TO DO ABOUT THEM | | | | |
|---|---|-------------------------------------|--------------|--|
| Symptom / effect | | Talk with your doctor or pharmacist | | Stop taking drug and immediately seek assistance |
| | | Only if severe | In all cases | |
| | Allergic Reaction: rash, hives, swelling of the face, throat, lips, difficulty swallowing or breathing | | | √ |
| | Liver Disorder: yellowing of the skin or eyes, dark urine, abdominal pain, nausea, vomiting, loss of appetite | | √ | |
| | Sexuality: Abnormal curvature of the penis | √ | | |
| | Change in sex drive | √ | | |
| | Change in ability to achieve or maintain an erection | √ | | |
| | Hallucinations: see or hear things that are not there | | √ | |
| | Depression: feels sad, loss of interest in usual activities changes in sleep and eating patterns | | √ | |

SERIOUS SIDE EFFECTS, HOW OFTEN THEY HAPPEN AND WHAT TO DO ABOUT THEM

| Symptom / effect | | Talk with your doctor or pharmacist | | Stop taking drug and immediately seek assistance |
|------------------|--|-------------------------------------|--------------|--|
| | | Only if severe | In all cases | |
| | Change in personality and confusion | | √ | |
| | Visual Disturbance: blurred vision, dry or irritated eyes | √ | | |
| | Hearing Disturbance: noises, reduced or loss of hearing | √ | | |
| | Parasesthesia: Colder than usual hands or feet | √ | | |
| | Gangrene; toes or fingers cold to the touch, discoloured and painful | | | √ |
| | Kidney Disorder: change in frequency of urination, swelling of extremities, fatigue, skin rash, itching, nausea, vomiting | √ | | |
| | High Blood Pressure: headaches, vision disorders, nausea and vomiting | | √ | |

SERIOUS SIDE EFFECTS, HOW OFTEN THEY HAPPEN AND WHAT TO DO ABOUT THEM

| Symptom / effect | | Talk with your doctor or pharmacist | | Stop taking drug and immediately seek assistance |
|------------------|---|-------------------------------------|--------------|--|
| | | Only if severe | In all cases | |
| | Blood clots: swelling, pain and redness in an arm or leg that can be warm to touch. You may develop sudden chest pain, difficulty breathing and heart palpitations | | | √ |
| | Decreased Platelets: bruising, bleeding, fatigue and weakness | | √ | |
| | Retroperitoneal fibrosis: lower back pain, kidney failure (low or no urine produced), high blood pressure, blood clot in the legs (See Blood Clots above) | | | √ |
| | Arthritis: stiff sore joints | | √ | |
| | Oculomucocutaneous Syndrome: red, irritated and watery eyes, skin rash and ear infection | | | √ |

This is not a complete list of side effects. For any unexpected effects while taking JAMP-METOPROLOL-L contact your doctor or pharmacist.

HOW TO STORE IT

Store JAMP-METOPROLOL-L tablets between 15°-30°C.
Protected from moisture and light.

Keep all medicines out of the reach and sight of children.

This medicine is prescribed for your specific medical problem and for your own use only. Do not give to other people.

Do not use outdated medicines. Discard them safely out of the reach of children or take them to your pharmacist who will dispose of them for you.

Reporting Side Effects

You can report any suspected side effects associated with the use of health products to Health Canada by:

- Visiting the Web page on Adverse Reaction Reporting (<https://www.canada.ca/en/healthcanada/services/drugs-health-products/medeffect-canada.html>) for information on how to report online, by mail or by fax; or
- Calling toll-free at 1-866-234-2345.

NOTE: Contact your health professional if you need information about how to manage your side effects. The Canada Vigilance Program does not provide medical advice.

MORE INFORMATION

If you want more information about JAMP-METOPROLOL-L:

- Talk to your healthcare professional
- Find the full product monograph that is prepared for healthcare professionals and includes this Patient Medication Information by visiting the <https://health-products.canada.ca/dpd-bdpp/index-eng.jsp>; the manufacturer's website (www.jamppharma.com), or by calling 1-866-399-9091.

This leaflet was prepared by
JAMP Pharma Corporation
1310 rue Nobel
Boucherville Québec
J4B 5H3, Canada

Revision date: January 25, 2021