

**PRODUCT MONOGRAPH**

**<sup>Pr</sup>NAT-CETIRIZINE**

**Cetirizine hydrochloride Tablets Natco Standard, 20 mg**

**Histamine H<sub>1</sub> Receptor Antagonist**

**Natco Pharma (Canada) Inc.  
2000 Argentia Road, Plaza 1, Suite 200,  
Mississauga  
Canada L5N 1P7**

**Date of Revision:  
March 03, 2021**

Submission Control No: 242780

## Table of Contents

<b>PART I: HEALTH PROFESSIONAL INFORMATION</b> .....	<b>3</b>
SUMMARY PRODUCT INFORMATION.....	3
INDICATIONS AND CLINICAL USE .....	3
CONTRAINDICATIONS .....	3
WARNINGS AND PRECAUTIONS.....	4
ADVERSE REACTIONS.....	5
DRUG INTERACTIONS.....	7
DOSAGE AND ADMINISTRATION .....	8
OVERDOSAGE .....	8
ACTION AND CLINICAL PHARMACOLOGY.....	9
STORAGE AND STABILITY .....	11
DOSAGE FORMS, COMPOSITION AND PACKAGING.....	<a href="#">11</a>
<b>PART II: SCIENTIFIC INFORMATION</b> .....	<b>12</b>
PHARMACEUTICAL INFORMATION.....	12
CLINICAL TRIALS .....	13
DETAILED PHARMACOLOGY.....	16
TOXICOLOGY .....	19
REFERENCES .....	28
<b>PART III: CONSUMER INFORMATION</b> .....	<b>30</b>

**NAT-CETIRIZINE**  
**(cetirizine hydrochloride)**

**PART I: HEALTH PROFESSIONAL INFORMATION**

**SUMMARY PRODUCT INFORMATION**

<b>Route of Administration</b>	<b>Dosage Form / Strength</b>	<b>Nonmedicinal Ingredients</b>
Oral	<b>Prescription :</b> 20 mg tablet	Microcrystalline cellulose, crospovidone, colloidal anhydrous silica, magnesium stearate, hypromellose, titanium dioxide and polyethylene glycol.

**INDICATIONS AND CLINICAL USE**

**Adults and children 12 years of age and over:**

NAT-CETIRIZINE (cetirizine hydrochloride) 20 mg tablets is indicated for the fast relief of nasal and non-nasal symptoms associated with seasonal and perennial allergic rhinitis (i.e. sneezing, rhinorrhea, post-nasal discharge, nasal congestion/stuffiness, tearing and redness of the eyes, itchy nose/throat) and chronic idiopathic urticaria (e.g. pruritus and hives).

**Pediatrics (< 12 years of age):**

NAT-CETIRIZINE (20 mg tablets) is not recommended in children less than 12 years of age, as safety and efficacy have not been established in this age group.

**CONTRAINDICATIONS**

NAT-CETIRIZINE (cetirizine hydrochloride) is contraindicated in those patients with a known hypersensitivity to it, to its parent compound, hydroxyzine, or to piperazine derivatives, in patients who are hypersensitive to any other ingredient in the formulation, or in patients with severe renal impairment (less than 10 mL/min creatinine clearance).

**WARNINGS AND PRECAUTIONS**

**General**

Severe skin reactions such as acute generalized exanthematous pustulosis (AGEP) have been reported very rarely with cetirizine-containing products. This acute pustular eruption may exhibit an early or delayed onset with numerous small, mostly non-follicular pustules arising on a widespread edematous erythema mainly localized on the skin folds, trunk, and upper extremities, which may be accompanied by fever. Patients should be carefully monitored.

If symptoms persist or worsen, or if new symptoms occur, the patient should discontinue use and consult a physician.

### **Activities Requiring Mental Alertness**

Studies using objective measurements have shown no effect of cetirizine hydrochloride at a dose of 10 mg on cognitive function, motor performance or sleep latency in healthy volunteers. However, in clinical trials the appearance of some CNS effects, particularly somnolence, have been observed. If drowsiness occurs, patients should be advised not to drive or operate machinery. Patients should be advised to avoid concurrent use of cetirizine hydrochloride with sedating substances, including alcohol, because additional reductions in alertness and additional impairment of CNS performance may occur. (See Drug Interactions).

### **Special Populations**

#### **Pregnant Women**

There are no adequate and well-controlled studies in pregnant women. Until such data become available, NAT-CETIRIZINE should not be used during pregnancy, unless advised otherwise by a physician.

No teratogenic effects were caused by oral doses as high as 60, 188 and 133 times the maximum clinically studied human dose in mice, rats and rabbits, respectively. No effects on reproduction and fertility were observed at doses as high as 40 and 10 times the maximum recommended human dose in male and female mice, respectively. An oral dose 60 times the maximum clinically studied human dose in female mice did not affect parturition or lactation. Although the animal studies are not indicative of any adverse effects during pregnancy at clinically relevant doses, such studies are not always predictive of a human response.

#### **Nursing Women**

Use of NAT-CETIRIZINE in nursing mothers is not recommended, unless directed otherwise by a physician. Studies in beagle dogs indicate that approximately 3% of the dose is excreted in milk. The extent of excretion in human milk is unknown.

#### **Pediatrics**

NAT-CETIRIZINE should not be administered to children below 12 years of age (See DOSAGE AND ADMINISTRATION). It is recommended to use an appropriate pediatric formulation of cetirizine.

#### **Geriatrics**

Cetirizine hydrochloride was well tolerated by patients aged 65 and over. Clearance of cetirizine hydrochloride is reduced in proportion to creatinine clearance. In patients whose creatinine clearance is reduced (i.e., those with moderate renal impairment), a starting dose of 5 mg/day is recommended (see HUMAN PHARMACOKINETICS).

#### **Hepatic Impairment**

Occasional instances of liver function test (transaminase) elevations have occurred during cetirizine hydrochloride therapy. This incidence was 1.6% in the short-term trials and 4.4% in the 6 month trials. These liver enzyme elevations, mainly ALT, were generally reversible. There was

no evidence of jaundice or hepatitis, and the clinical significance is presently unknown. Consequently, NAT-CETIRIZINE should be used with caution in patients with pre-existing liver disease. In patients with moderate or severe hepatic impairment, a starting dose of 5 mg is recommended.

## ADVERSE REACTIONS

### **Clinical Trial Adverse Drug Reactions**

*Because clinical trials are conducted under very specific conditions the adverse reaction rates observed in the clinical trials may not reflect the rates observed in practice and should not be compared to the rates in the clinical trials of another drug. Adverse drug reaction information from clinical trials is useful for identifying drug-related adverse events and for approximating rates.*

In clinical development programs (domestic and international), cetirizine hydrochloride has been evaluated in more than 6000 treated patients at daily doses ranging from 5 to 20 mg. The most common adverse reactions were headache and somnolence (see paragraph below). The incidence of headache associated with cetirizine hydrochloride was not different from placebo. The incidence of somnolence associated with cetirizine hydrochloride was dose related and predominantly mild to moderate.

Most adverse reactions reported during cetirizine hydrochloride therapy in clinical trials were mild to moderate. The incidence of discontinuation due to adverse reactions in patients receiving cetirizine hydrochloride was not significantly different from placebo (1.0% vs 0.6%, respectively, in placebo-controlled trials). There was no difference by gender or by body weight with regard to the incidence of adverse reactions.

Occasional instances of transient, reversible hepatic transaminase elevations have occurred during cetirizine hydrochloride therapy, without evidence of jaundice, hepatitis or other clinical findings.

Adverse events which were reported at an incidence of greater than 2% and greater than placebo in clinical trials with the 20 mg tablet (one to four weeks duration) are listed in Table 1.

**TABLE 1**  
**ADVERSE REACTIONS REPORTED IN PLACEBO-CONTROLLED UNITED STATES**  
**CETIRIZINE HYDROCHLORIDE TRIALS (TOTAL DAILY DOSE 20 mg) AT RATES**  
**OF ≥2% AND GREATER THAN PLACEBO**  
**(Percent Incidence)**

<b>Adverse Experience</b>	<b>Cetirizine Hydrochloride 20 mg Tablet (N=272)</b>	<b>Placebo (N=671)</b>	<b>Difference of Percentage</b>
Somnolence	23.9%	7.7%	16.2
Dry Mouth	7.7	1.5	6.2
Fatigue	7.0	2.4	4.6

The following events were observed infrequently (equal to or less than 2%), in 3982 patients who received cetirizine hydrochloride in worldwide trials, including an open study of 6 months duration; a causal relationship with cetirizine hydrochloride administration has not been established.

**Application Site:** application site reaction, injection site inflammation

**Autonomic Nervous System:** anorexia, urinary retention, flushing, saliva increased

**Cardiovascular:** palpitation, tachycardia, hypertension, arrhythmia, cardiac failure

**Central and Peripheral Nervous Systems:** fatigue, dizziness, insomnia, nervousness  
paresthesia, confusion, hyperkinesia, hypertonia, migraine, tremor, vertigo, leg cramps,  
ataxia, dysphonia, coordination abnormal, hyperesthesia, hypoesthesia, myelitis, paralysis,  
ptosis, speech disorder, twitching, visual field defect

**Endocrine:** thyroid disorder

**Gastrointestinal:** nausea, pharyngitis, appetite increased, dyspepsia, abdominal pain, diarrhea,  
flatulence, constipation, vomiting, stomatitis ulcerative, tongue disorder, tooth caries  
aggravated, stomatitis, tongue discoloration, tongue edema, gastritis, hemorrhage rectum,  
hemorrhoids, melena, hepatic function abnormal

**Genitourinary:** polyuria, urinary tract infection, cystitis, dysuria, hematuria, urine abnormal

**Hearing and vestibular:** earache, tinnitus, deafness, ototoxicity

**Metabolic/Nutritional:** thirst, edema, dehydration, diabetes mellitus

**Musculoskeletal:** myalgia, arthralgia, bone disorder, arthrosis, tendon disorder, arthritis, muscle  
weakness, Psychiatric: depression, emotional lability, concentration impaired, anxiety,  
depersonalization, paroniria, thinking abnormal, agitation, amnesia, libido decreased, euphoria

**Resistance Mechanism:** healing impaired, herpes simplex, infection, infection fungal, infection  
viral

**Respiratory System:** epistaxis, rhinitis, coughing, respiratory disorder, bronchospasm,  
dyspnea, upper respiratory tract infection, hyperventilation, sinusitis, sputum increased,  
bronchitis, pneumonia

**Reproductive:** dysmenorrhea, menstrual disorder, breast pain female, intermenstrual bleeding,  
leukorrhea, menorrhagia, pregnancy unintended, vaginitis, testes disorder

**Reticuloendothelial:** lymphadenopathy

**Skin:** pruritus, rash, skin disorder, skin dry, urticaria, acne, dermatitis, rash erythematous,  
sweating increased, alopecia, angioedema, furunculosis, bullous eruption, eczema,  
hyperkeratosis, hypertrichosis, photosensitivity reaction, photosensitivity toxic reaction, rash  
maculopapular, seborrhea, purpura

**Special Senses:** taste perversion, taste loss, parosmia

**Vision:** eye abnormality, vision abnormal, eye pain, conjunctivitis, xerophthalmia, glaucoma,  
ocular hemorrhage

**Body as a Whole:** weight increase, back pain, malaise, pain, chest pain, fever, asthenia, edema  
generalized, edema periorbital, edema peripheral, rigors, edema legs, face edema, hot flushes,  
abdomen enlarged, allergic reaction, nasal polyp

### **Less Common Clinical Trial Adverse Drug Reactions (<1%)**

Weight gain was reported as an adverse event in 0.4% of cetirizine patients in placebo controlled trials. In an open study of 6 months duration, the mean weight gain was 2.8% after 20 weeks, with no further increase at 26 weeks.

### **Abnormal Hematologic and Clinical Chemistry Findings**

Occasional instances of transient, reversible hepatic transaminase elevations have occurred during cetirizine therapy.

### **Post-Market Adverse Drug Reactions**

Additional adverse drug reactions identified during post-marketing experience with cetirizine include: blurred vision, eye swelling, feeling abnormal, enuresis, hepatic function abnormal (increased transaminases, alkaline phosphatase, alanine aminotransferase, aspartate aminotransferase and  $\gamma$ -glutamyl transferase), erectile dysfunction, hallucination, dysgeusia, dyskinesia, dystonia, memory impairment, tremor, arthralgia, urinary retention, pruritus upon withdrawal and weight increased.

In post-marketing experience the following additional rare, but potential severe adverse events have been reported: hemolytic anemia, thrombocytopenia, orofacial dyskinesia, severe hypotension, anaphylaxis, hepatitis, glomerulonephritis, stillbirth, cholestasis and acute generalized exanthematous pustulosis. In addition, isolated cases of the following adverse drug reactions have been reported: convulsions, syncope, aggression, and hypersensitivity.

## **DRUG INTERACTIONS**

### **Overview**

Interaction studies with cetirizine hydrochloride and alcohol or diazepam indicate that at therapeutic doses, cetirizine hydrochloride does not increase alcohol-induced or diazepam-induced impairment of motor and mental performance.

The drug should not be used with sedating substances such as alcohol and some other medications, including anti-anxiety medications, sleep aids, antihistamines, antidepressants, muscle relaxants or prescription analgesics due to possible interactions.

### **Drug-Drug Interactions**

No clinically significant drug interactions have been found with theophylline, pseudoephedrine, cimetidine, erythromycin and ketoconazole. Epidemiologic data suggests that there also would not be interaction with other macrolide antibiotics or imidazole antifungals. In clinical trials, cetirizine hydrochloride has been safely administered with beta-agonists, non-steroidal anti-inflammatory drugs, oral contraceptives, narcotic analgesics, corticosteroids, H<sub>2</sub>-antagonists, cephalosporins, penicillins, thyroid hormones and thiazide diuretics. If drowsiness occurs, concurrent use of cetirizine hydrochloride with sedating substances should be avoided because additional reductions in alertness and additional impairment of CNS performance may occur. (See Activities Requiring Mental Alertness).

Based on (1) its relatively low level of metabolic elimination, (2) no effect on corrected QT intervals at plasma concentrations three times the maximal therapeutic levels, and (3) no apparent interactions with ketoconazole or erythromycin, cetirizine is unlikely to have clinically significant interactions with other macrolides such as clarithromycin or other imidazole antifungals such as itraconazole in patients with normal renal and hepatic function. Although no data with these other drugs are available at the present time, there is no

epidemiological evidence (the safety database comprised 6,490 patients evaluated in U.S. and Canadian studies) of interactions between macrolide antibiotics and/or imidazole antifungals taken orally, and cetirizine/hydroxyzine. The epidemiologic data do not suggest an increase of adverse events, cardiac or non-cardiac, in patients treated with cetirizine and concomitant macrolide or imidazole antifungal medication.

## **DOSAGE AND ADMINISTRATION**

### **Dosing Considerations**

The recommended initial dose of cetirizine hydrochloride is 5 or 10 mg per day, depending on symptom severity. NAT-CETIRIZINE 20 mg tablets are recommended only if lower doses (non-prescription) of cetirizine hydrochloride do not provide sufficient response for patients 12 years of age and older.

### **Recommended Dosage and Dosage Adjustment**

#### **Adults and Children 12 Years of Age and Over**

The recommended dose is one 20 mg tablet once daily, with or without food.

The maximum recommended daily dose is 20 mg.

#### **Geriatrics ( $\geq 65$ Years of age)**

NAT-CETIRIZINE 20 mg tablets are not recommended in patients 65 years and older. The recommended starting dose is 5 mg once daily.

#### **Patients with hepatic impairment**

NAT-CETIRIZINE 20 mg tablets are not recommended in patients with moderate or severe hepatic impairment. The recommended starting dose is 5 mg once daily.

#### **Patients with renal impairment**

NAT-CETIRIZINE 20 mg tablets are not recommended for patients with moderate renal impairment. In patients with moderate renal impairment, the recommended starting dose is 5 mg once daily.

NAT-CETIRIZINE is contraindicated in patients with severe renal impairment (less than 10 mL/min creatinine clearance).

#### **Pediatrics (< 12 years of age)**

NAT-CETIRIZINE 20 mg tablets should not be administered to children under 12 years of age. It is recommended to use an appropriate pediatric formulation of cetirizine.

## **OVERDOSAGE**

Overdose has been reported with cetirizine hydrochloride. Symptoms observed after an overdose of cetirizine are mainly associated with CNS effects or with symptoms that could suggest an



anticholinergic effect. Adverse events reported after an intake of at least 5 times the recommended daily dose are: confusion, diarrhea, dizziness, fatigue, headache, malaise, mydriasis, pruritus, restlessness, sedation, somnolence, stupor, tachycardia, hypertension, tremor, and urinary retention. Hyperactivity and severe lethargy have been seen in children. If an acute overdose occurs, evacuation of the stomach should be considered during the first few hours after this overdose. Treatment should be symptomatic and supportive taking into account any concomitantly ingested medications. There is no known specific antidote to cetirizine hydrochloride. Cetirizine hydrochloride is not effectively removed by dialysis, and dialysis will be ineffective unless a dialyzable agent has been concomitantly ingested. The minimal lethal oral dose in rodents is at least 590 times the maximum clinically studied dose.

For management of a suspected drug overdose, contact your regional Poison Control Centre.
---

## **ACTION AND CLINICAL PHARMACOLOGY**

### **Mechanism of Action**

Cetirizine hydrochloride, an active human metabolite of hydroxyzine, is a histamine H<sub>1</sub> receptor antagonist anti-allergic compound; its principal effects are mediated via selective inhibition of peripheral H<sub>1</sub> receptors. Cetirizine hydrochloride is distinguished from other histamine H<sub>1</sub> receptor antagonists by the presence of a carboxylic acid function. This difference may be partly responsible for the selectivity of cetirizine hydrochloride seen in pharmacologic models and its distinctive pharmacokinetic properties in humans.

### **Pharmacodynamics**

The antihistaminic activity of cetirizine hydrochloride has been well documented in a variety of animal and human models. *In vivo* animal models have shown negligible anticholinergic or antiserotonergic activity. *In vitro* receptor binding studies have detected no measurable affinity for other than H<sub>1</sub> receptors. Autoradiographic studies have shown negligible penetration into the brain. Systemically administered cetirizine does not significantly occupy cerebral H<sub>1</sub> receptors. Several studies involving objective and subjective tests in healthy volunteers have demonstrated that cetirizine hydrochloride at doses up to 10 mg did not significantly differ from placebo with respect to CNS impairment, daytime drowsiness, reaction times, mental alertness, task performance, objective CNS depression and various other tests of cognitive function.

Cetirizine hydrochloride does not exacerbate asthma and is effective in a variety of histamine mediated disorders. In adults, oral doses of 5-20 mg in humans strongly inhibit the skin wheal and flare response caused by the intradermal injection of histamine. The onset of activity occurs within 20 (50% of subjects) to 60 (95% of subjects) minutes and persists for at least 24 hours following a single dose. The effects of intradermal injection of various other mediators or histamine releasers as well as components of the allergy cascade, including allergic inflammatory response to cutaneous antigen challenge are also inhibited.

### **Pharmacokinetics**

**Absorption:** In adults, cetirizine hydrochloride is rapidly absorbed after oral administration. Peak plasma levels after a 10 mg dose are approximately 300 ng/mL and occur at about 1 hour. Co-

administration of cetirizine hydrochloride with food does not affect bioavailability as measured by AUC but absorption is delayed by about 1 hour, with 23% lower  $C_{max}$ .

While a high-fat meal does not impact the extent of absorption of cetirizine from the orally disintegrating tablet (ODT) as measured by  $AUC_T$ , absorption is delayed by approximately 3 hours and  $C_{max}$  is reduced by approximately 37% when the ODT is administered with a high-fat meal relative to the ODT administered under fasted conditions.

**Distribution:** Plasma protein binding is 93% in the concentration range observed in clinical studies. The apparent volume of distribution is 0.45 L/kg, suggestive of significant extravascular distribution.

**Metabolism:** In adults, cetirizine hydrochloride is less extensively metabolized than other antihistamines and approximately 60% of an administered dose is excreted unchanged in 24 hours. High bioavailability is associated with generally low inter-subject variation in blood levels. It is attributable primarily to low first-pass metabolism. Only one metabolite has been identified in humans - the product of oxidative dealkylation of the terminal carboxymethyl group. The antihistaminic activity of this metabolite is negligible.

**Excretion:** The plasma elimination half-life is approximately 8 to 9 hours and does not change with multiple dosing. Pharmacokinetics are dose independent and plasma levels are proportional to the dose administered over the clinically studied range of 5 to 20 mg.

**TABLE 2**  
**TYPICAL PHARMACOKINETIC PARAMETERS OF CETIRIZINE IN ADULTS**

Parameter	Adults 10 mg Single dose
$C_{max}$ (ng/mL)	300
$T_{max}$ (hr)	1.1
$T_{1/2}$ (hr)	8.0
AUC (ng.hr/mL)	2871
Urinary recovery (%)	60

### **Special Populations and Conditions**

**Geriatrics:** The clearance of cetirizine hydrochloride is reduced in elderly patients, but only in proportion to the decrease in creatinine clearance. Thus, in 16 patients with a mean age of 77 years, half-life increased to 12 hours. Cetirizine hydrochloride blood levels were monitored in a clinical trial of 59 patients aged 60 to 82, who received 10 mg daily for 3 weeks and no undue

accumulation of cetirizine hydrochloride was found.

**Hepatic Insufficiency/Renal Insufficiency:** In patients with mild to moderate hepatic and renal impairment, total body clearance of cetirizine hydrochloride is reduced and AUC and half-life increased by about 2 to 3 fold. Clearance is reduced in proportion to the decline in creatinine clearance. Plasma levels are unaffected by hemodialysis. The plasma elimination half-life in dialysis patients is approximately 20 hours and the plasma AUC is increased by about threefold.

## **STORAGE AND STABILITY**

Recommended storage: Store between 15°-30°C. Protect from moisture.

## **DOSAGE FORMS, COMPOSITION AND PACKAGING**

NAT-CETIRIZINE (cetirizine hydrochloride) 20 mg Tablet: 20 mg of cetirizine hydrochloride. The 20 mg tablets are only available by prescription.

NAT-CETIRIZINE 20 mg tablets are white to off-white, oval rectangular film coated tablets with 'C break line 2' on one side and plain on the other side. The non-medicinal ingredients are: microcrystalline cellulose, crospovidone, colloidal anhydrous silica, magnesium stearate; hypromellose, titanium dioxide and polyethylene glycol. Available in plastic bottles of 100's and 500's tablets.

## PART II: SCIENTIFIC INFORMATION

### PHARMACEUTICAL INFORMATION


#### Drug Substance

- Common Name: Cetirizine hydrochloride (USP)  
Cetirizine dihydrochloride (Ph.Eur.)
- Proper Name: Cetirizine dihydrochloride
- Chemical Name: (RS)-2-(2-(4-[4-chlorophenyl]phenylmethyl)-1-piperazinyl)ethoxy) acetic acid, dihydrochloride
- Molecular Formula:  $C_{21}H_{27}Cl_3N_2O_3$
- Molecular Weight: 461.8 g/mol

#### Physicochemical properties:

Cetirizine hydrochloride is a white or almost white powder. It is freely soluble in water and practically insoluble in methylene chloride and acetone.

#### Structural Formula:



## CLINICAL TRIALS

### Comparative Bioavailability Studies

A randomized, two-treatment, two-sequence, two-period, crossover, single 1 x 20 mg dose bioequivalence study of NAT-CETIRIZINE (Cetirizine hydrochloride tablets) 20 mg (Natco Pharma (Canada) Inc.) and REACTINE® (Cetirizine hydrochloride) Tablets 20 mg (McNeil Consumer Healthcare, division of Johnson & Johnson Inc.) was conducted in 18 healthy Asian male subjects from 24-39 years of age under fasting conditions. A summary of the bioavailability data from 15 subjects who completed the study is presented in the following table.

**SUMMARY TABLE OF THE COMPARATIVE BIOAVAILABILITY DATA**

CETIRIZINE (1 x 20 mg cetirizine hydrochloride) From measured data				
Geometric Mean Arithmetic Mean (CV%)				
Parameter	Test*	Reference†	% Ratio of Geometric Means	90% Confidence Interval
AUC <sub>T</sub> (ng.hr/mL)	6954.3 7042.09 (15.5)	6618.2993 6726.72 (17.5)	105.1	101.4 - 108.9
AUC <sub>I</sub> (ng.hr/mL)	7225.0 7329.7 (16.7)	6900.7 7035.1 (19.3)	104.7	101.0 - 108.6
C <sub>max</sub> (ng/mL)	792.5 800.6 (14.3)	710.3 726.3 (18.3)	111.6	105.2 - 118.4
T <sub>max</sub> § (hr)	0.67 (0.50-1.25)	1.25 (0.50-2.50)		
T <sub>1/2</sub> € (hr)	9.30 (22.7)	9.61 (26.0)		

\*NAT-CETIRIZINE (Cetirizine hydrochloride) Tablets 20 mg (Natco Pharma (Canada) Inc.)

†REACTINE® (Cetirizine hydrochloride) Tablets 20 mg (McNeil Consumer Healthcare, division of Johnson & Johnson Inc.) was purchased from Canada.

§ Expressed as the median (range) only.

€ Expressed as arithmetic mean (CV%) only.

Randomized, multi-centre, double-blind, placebo-controlled clinical trials have demonstrated the effectiveness of cetirizine hydrochloride in relieving the symptoms associated with seasonal allergic rhinitis, perennial allergic rhinitis and chronic idiopathic urticaria. The clinical trials have shown only weak anticholinergic effects. There is no evidence that tolerance to the antihistaminic or anti-allergy effects of cetirizine hydrochloride develops or that it has any abuse potential or dependency liability.

In adults, objective and subjective tests in healthy volunteers, have demonstrated that cetirizine hydrochloride at doses up to 10 mg did not significantly differ from placebo with respect to CNS impairment, daytime drowsiness, reaction times, mental alertness, task performance, objective

CNS depression and various other tests of cognitive function.

Specific electrocardiographic (ECG) studies in healthy adult volunteers at doses up to 60 mg per day (three times the maximum clinically studied dose) for 1 week did not prolong QTc intervals nor was there any evidence of QTc prolongation in clinical trials which included ECG evaluations.

Cetirizine given at the maximum clinically studied dose of 20 mg daily did not prolong the QTc when given in combination with either ketoconazole 400 mg o.d. or erythromycin 500 mg q8h for 10 days. Moreover, cetirizine did not significantly alter the pharmacokinetics of either ketoconazole or erythromycin nor were the pharmacokinetics of cetirizine altered by either ketoconazole or erythromycin.

Three well-controlled clinical trials assessed the effects of cetirizine (10 mg) on nasal congestion as an individual symptom score when it was part of the constellation of symptoms associated with allergic rhinitis (as assessed in adults with SAR). Cetirizine proved to be significantly more effective than placebo in improving nasal congestion.

Improvement in quality of life (QOL) with cetirizine hydrochloride (10 mg) in patients with allergic rhinitis has been demonstrated in a number of published studies (2 to 6 weeks duration) using a variety of validated QOL measurement tools (see Table 3). Improvement in the following QOL domains were observed: physical, social and work activities, vitality and social functioning, practical problems, symptom distress (nasal, eye), sleep problems, and emotional difficulties.

**TABLE 3**

**RANDOMIZED, PARALLEL, DOUBLE-BLIND, PLACEBO-CONTROLLED CLINICAL STUDIES DEMONSTRATING IMPROVEMENT IN QUALITY OF LIFE DOMAINS FOLLOWING REGULAR CETIRIZINE HYDROCHLORIDE 10 mg (p.o.) USE (2-6 WEEKS)**

<b>Study</b>	<b>Study Description</b>	<b>Result</b>
Bousquet J. <i>et al</i> (1996)	Objective to determine effect of cetirizine HCl 10 mg on QOL in patients with allergic rhinitis. Duration of study: 6 weeks. Total of 122 subjects completed cetirizine arm versus 126 subjects in placebo arm. Validated QOL measurement tool used: SF - 36*	Quality of life and nasal symptoms were measured after 1 and 6 weeks of treatment using SF-36 questionnaire. After 6 weeks, percentage of days without rhinitis or only mild rhinitis was significantly greater in the cetirizine group compared with the placebo group. Cetirizine improved all nine QOL dimensions (from p = 0.01 to p<0.0001) after 1 and 6 weeks of cetirizine treatment.
Burtin B. <i>et al</i> (2000)	Investigate extent to which cetirizine HCl 10 mg continues to improve QOL after long-term treatment (6 weeks) versus shorter-term treatment (1 week). Validated QOL measurement tool used: SF-36*. Note: This is an additional analysis of the study published by Bousquet <i>et al</i> (1996).	Further 5 week course of therapy maintains QOL improvements seen after 1 week.
Murray JJ. <i>et al</i> (2002)	Evaluate the health-related quality of life effects, safety and efficacy of cetirizine HCl 10 mg in treatment of seasonal allergic rhinitis. Patients completing the 2-week treatment period included 413 in the cetirizine group and 396 in the placebo group. Validated QOL measurement tool used: RQLQ <sup>†</sup>	QOL scores were measured following 2 weeks of treatment. The cetirizine-treated patient group experienced greater (p<0.001) improvement in overall RQLQ and individual domain scores, compared to the placebo patient group.
Noonan MJ. <i>et al</i> (2003)	Test the effect of cetirizine HCl 10 mg once daily on the health-related QOL (HRQL) of adult patients 18-65 years of age with allergic rhinitis. Study duration: 2 weeks. 196 subjects completed the cetirizine study arm; 183 completed the placebo study arm. Validated QOL measurement tool used: RQLQ <sup>†</sup>	The cetirizine-treated patient group reported greater improvement in overall HRQL (p<0.001) and in each of the seven domains of RQLQ after two weeks (p<0.05 to p<0.001) versus the placebo patient group.

\* SF-36: Medical Outcome Short-Form Health Survey

† RQLQ: Rhinoconjunctivitis Quality of Life Questionnaire

## DETAILED PHARMACOLOGY

### ANIMAL PHARMACOLOGY

Cetirizine hydrochloride has been shown to be a potent H<sub>1</sub> antagonist in 14 animal studies carried out to evaluate the antihistaminic activity of the drug in vivo. The selectivity of cetirizine hydrochloride for H<sub>1</sub> receptors has also been demonstrated in isolated organ studies and receptor binding studies in vitro. Cetirizine hydrochloride has been shown to inhibit endogenous and exogenous histamine-induced bronchial and cutaneous reactions.

Autoradiographic studies with radiolabeled cetirizine hydrochloride in the rat have shown negligible penetration of the brain. *Ex vivo* experiments in the mouse have shown that systemically administered cetirizine hydrochloride minimally occupies cerebral H<sub>1</sub> receptors. In various animal behaviour models and neuropharmacologic studies in mice, rats and dogs, cetirizine hydrochloride exhibits a lack of significant central nervous system effects up to doses of 15 mg/kg to 46 mg/kg given p.o. or i.p. These doses are 30 to 1000 times higher than the dose required to exert an antihistaminic effect on cutaneous reactions.

### HUMAN PHARMACOLOGY

#### Pharmacodynamics

Studies in normal volunteers show that cetirizine hydrochloride at doses of 5 to 20 mg strongly inhibits the skin wheal and flare caused by the intradermal injection of histamine. The onset of activity corresponds with the occurrence of maximal plasma levels, and significant blockade persists for at least 24 hours after a single dose. The effects of intradermal injection of various other mediators or histamine releasers are also inhibited by cetirizine hydrochloride, as is cold-induced urticaria.

In mildly asthmatic subjects, cetirizine hydrochloride at 5 to 20 mg is highly effective in blocking bronchoconstriction due to nebulized histamine, with virtually total blockade after a 20 mg dose; a modest reduction of resting bronchial tone is also seen.

Studies in normal subjects using objective assessments of psychomotor performances showed that cetirizine hydrochloride at doses up to 20 mg did not produce significant changes in the Multiple Sleep Latency test, a measure of daytime drowsiness, in comparison with placebo. However, hydroxyzine 25 mg caused a statistically significant decrease in time to sleep onset. When the Flicker Fusion Threshold was used to measure mental alertness, cetirizine hydrochloride did not produce significant change but hydroxyzine significantly reduced the mental alertness. In this study, cetirizine hydrochloride 10 and 20 mg and hydroxyzine 25 mg had equipotent antihistaminic activity as determined by the suppression of skin wheal response to histamine.

Several combined placebo and positive control studies in normal subjects using a multiple crossover design with objective and subjective assessments of CNS and performance impairment showed that cetirizine hydrochloride 10 mg did not differ from placebo. Positive controls i.e. sedating antihistamines, e.g. diphenhydramine, hydroxyzine, triprolidine, were included in these trials to verify that the tests were able to detect impairment. Objective tests



included: Multiple Sleep Latency Test (EEG monitoring), Critical Flicker-Fusion (CFF), Choice Reaction Time (CRT), Continuous Tracking Test (CTT), word testing, simulated driving tests and assembly line tests (SALT), actual road-driving tests. Subjective tests included: Visual Analog Scale (VAS) reporting, Stanford Sleepiness Scale (SSS) by the subject as well subjective assessments by driving instructors.

Due to the association of torsades and QT prolongation with newer antihistamines, and the metabolic/pharmacokinetic interaction of antihistamines with erythromycin and ketoconazole, three studies were performed to evaluate the pharmacokinetic effects and ECG effects of cetirizine, and the possible interactions of cetirizine with ketoconazole and erythromycin. These studies show that cetirizine, alone or in combination with erythromycin or ketoconazole, does not cause clinically significant QTc prolongation. Furthermore, no effects on the pharmacokinetics of erythromycin or ketoconazole and no effects of these two compounds on the pharmacokinetics of cetirizine were seen.

**Protocol 90CK16-0497:** There were no statistically significant differences among the treatments in mean QTc prior to daily dosing, indicating that multiple dosing with cetirizine at both the maximal clinically studied daily dose (20 mg) or three times the maximal clinically studied dose (60 mg once daily) has no effect on QTc relative to a placebo effect. Furthermore, cetirizine did not have any statistically significant effect on QT (uncorrected) or on heart rate as measured by RR interval. This finding was consistent over all dose days as indicated by no statistically significant treatment-by-day interaction for each of the three parameters. This suggests that, within the first 7 days of treatment, cetirizine produces neither an early transient effect nor a late-appearing cumulative effect. Also, there were no significant differences with respect to the mean changes in QTc, QT, and RR from pre-dosing to 1, 2, 4, and 6 hours after dosing indicating that a dose of cetirizine has no acute effect on QT interval or heart rate relative to placebo at any of the post-dose hours for up to 7 days of treatment. The plasma cetirizine concentration-time profiles were dose proportional.

Four subjects (19.1%) during cetirizine 20 mg treatment and 6 subjects (28.6%) during cetirizine 60 mg experienced at least one 10% prolongation of QTc as compared to 6 subjects (28.6%) on placebo. These incidence rates were not significantly different. The largest prolongations observed were 15.6%, 19.0%, and 15.4% over baseline for placebo, cetirizine 20 mg, and cetirizine 60 mg, respectively.

Protocol 92KC16-0604: The objectives of this study were to determine whether cetirizine, in the presence of erythromycin, induces a prolongation of the QT interval and to determine whether there are pharmacokinetic interactions between cetirizine and erythromycin in young, healthy males. This was a randomized, multiple dose, open (the cardiologist was blinded), two-way crossover study with a washout period. The two treatment regimens administered in the study were the following:

**Regimen 1**

Day 1:	placebo o.d.
Days 2-6:	20 mg cetirizine o.d.
Days 7-16:	500 mg erythromycin q8h and 20 mg cetirizine o.d.

**Regimen 2**

Day 1: placebo o.d.  
Days 2-6: 500 mg erythromycin q8h  
Days 7-16: 500 mg erythromycin q8h and  
20 mg cetirizine o.d.

The mean change from baseline Hodges QTc after 5 days of dosing with cetirizine alone and erythromycin alone was -5, 10 msec and 3.01 msec, respectively. After an additional 10 days of dosing with combination treatment, the mean change from baseline was -3.71 msec for combination treatment following cetirizine alone and -0.39 msec for combination treatment following erythromycin alone. Using these mean changes, the drug interaction effect was estimated to be 0.03 msec, which is not statistically significantly different from zero. This result indicates that any possible effect on changes in Hodges QTc attributable to either drug alone is not altered by the presence of the other, and that the effect on Hodges QTc of combination dosing is the sum of the individual effects. The estimated effect of cetirizine is -5.08 msec which is a statistically significant reduction from baseline. The estimated erythromycin effect of 3.03 msec was not statistically significant. These results indicate that cetirizine did not induce a mean prolongation of Hodges QTc, and since the effect of combination dosing was just the sum of each component (estimated to be -2.05), there was no significant mean prolongation associated with combination treatment.

No subject experienced a 10% prolongation of Hodges QTc over baseline during cetirizine alone treatment. Eight subjects experienced at least 1 prolongation of 10% or greater. Two subjects (13.3%) had a 10% or greater increase during treatment with erythromycin alone, 2 subjects (14.3%) during combination treatment following cetirizine and 4 subjects (26.7%) had an occurrence during combination treatment following erythromycin. The maximum prolongation in any subject in the study was 17.8% which occurred during erythromycin treatment alone. There was no significant pharmacokinetic interaction between cetirizine and erythromycin when administered concomitantly in therapeutic dosages and regimens.

**Protocol 92CK16-0603:** The objectives of this study were to determine whether cetirizine, in the presence of ketoconazole, induces a prolongation of the QT interval and to determine whether there are pharmacokinetic interactions between cetirizine and ketoconazole in young, healthy males. This was a randomized, multiple dose, open (the cardiologist was blinded), two-way crossover study. The two treatment regimens administered in the study were the following:

**Regimen 1**

Day 1: placebo o.d.  
Days 2-6: 400 mg ketoconazole o.d.  
Days 7-16: 400 mg ketoconazole o.d. and  
20 mg cetirizine o.d.

**Regimen 2**

Day 1: placebo o.d.  
Days 2-6: placebo o.d.  
Days 7-16: 20 mg cetirizine o.d.

There was no statistically significant drug interaction effect on the change in Hodges QTc from baseline. This indicates that the effect of the combination on changes in Hodges QTc is equal to

the sum of the individual component effects. The effects of each drug alone on change in Hodges QTc from baseline were statistically significant, with a mean increase from baseline of 8.16 msec and 8.32 msec for cetirizine and ketoconazole, respectively. Based on these findings, the effect of combination treatment on changes in Hodges QTc is estimated to be 16.48 msec.

No subject experienced a 10% or greater QTc prolongation during the 5 days placebo treatment. Two subjects (13.3%) experienced an increase in QTc of 10% or greater during the 10 day cetirizine treatment, 1 subject (6.3%) during the 5 day ketoconazole treatment and 5 subjects (31.3%) had an occurrence during combination treatment (2 in study phase I and 3 in study phase II). The maximum prolongation in any subject in the study was 14.3%, which occurred during combination treatment. Cetirizine did not significantly affect ketoconazole plasma pharmacokinetics.

Using Bazett's formula for QTc, 3 subjects had a total of 12 occurrences of a QTc >440 msec. There was 1 occurrence on placebo, 4 on cetirizine treatment and 7 on combined treatment. These occurrences of QTc >440 msec were episodic and not sustained.

The results of the study of protocol 90CK16-0497 demonstrate that cetirizine alone in multiple doses up to 60 mg (three times the maximum recommended dose of 20 mg) does not cause a prolongation of the QTc. Cetirizine did not increase mean QTc nor increase the percentage of patients who had 10% increases or greater in post-dose QTc. The pharmacokinetics of cetirizine were linear over the dose range and no dose related increase in QTc was seen. The results of study protocols 92CK16-0603 and 0604 demonstrate there was no significant interaction of cetirizine with either ketoconazole or erythromycin on QTc. Cetirizine given at the maximum recommended dose of 20 mg daily did not prolong the QTc when given in combination with either ketoconazole 400 mg o.d. or erythromycin 500 mg q8h for 10 days. Moreover, cetirizine did not significantly alter the pharmacokinetics of either ketoconazole or erythromycin nor were the pharmacokinetics of cetirizine altered by either ketoconazole or erythromycin.

With regard to QTc effect of cetirizine alone in the interaction studies, a small clinically insignificant decrease was seen in the erythromycin-cetirizine interaction study, and a small clinically insignificant increase in QTc was seen in the cetirizine-ketoconazole study. However, this small increase may be the result of other factors. For example, in the study of protocol 0497, a small increase in QTc was seen with placebo. In order to facilitate a comparison of the data in the 20-60 mg cetirizine study (protocol 90CK16-0497) with that in the two interaction studies, an analysis was done using the Hodges QTc formula and statistical models similar to the interaction study analyses. Based on this analysis, QTc increases of 5.4 msec, 3.0 msec and 7.3 msec for placebo, 20 mg and 60 mg cetirizine, respectively, were observed at the end of the 7 day treatment period. A shortened RR interval was found in all treatment groups, including placebo. The increase associated with placebo treatment indicates that other factors may affect QTc such as deconditioning during confinement, which is essentially a time effect.

## **TOXICOLOGY**

### **Acute Toxicity Studies**

#### **1. Rodents**

Cetirizine hydrochloride was administered orally or intravenously to 10 fasted animals/sex/dose level. Clinical signs, food consumption, and mortality were observed for 14 days; body weights were recorded at 1 and 2 weeks, and all animals were necropsied. Results revealed no differences in clinical signs nor lethality for either sex. The oral non-lethal dose for the most sensitive sex was 250 times the expected maximal clinical dose (EMCD) of 0.4 mg/kg (20 mg/50 kg). The severity of symptoms was dose related. The main symptoms were dose related. The main symptoms were cyanosis and dyspnea. Following oral administration in rats, mortalities occurred within 24 hours; after IV administration, all deaths occurred within 10 minutes and survivors recovered within 1-3 hours. In mice, mortalities were seen in the first 3 days after oral and within 24 hours in most cases after IV administration. The results of the rodent acute toxicity studies are summarized in Table 4.

**TABLE 4**

Species	Sex	Maximum Non-Lethal Dose mg/kg		LD <sub>50</sub> (95% Confidence Limits) mg/kg		LD <sub>50</sub> Ratio
		PO	IV	PO	IV	PO/IV
Rats (Wistar)	M	237	ND*	703 (305-1175)	66 (58-96)	10.65
	F	237	42	865 (553-1353)	70 (61-82)	12.36
Mice (NMRI)	M	237	240	600 (375-1391)	336 (301-476)	1.79
	F	100	240	752 (432-5114)	301 (264-366)	2.50

\* ND = Not determined.

## 2. Dogs

Cetirizine hydrochloride was administered orally to groups of 2 fasted beagle dogs (1M; 1F) at doses of 40, 80, 160, or 320 mg/kg and IV in the cephalic vein at a dose of 70 mg/kg to 2 fasted beagle dogs (1M; 1F) and 125 mg/kg to 1 fasted female beagle dog. Clinical signs, food consumption, and mortality were observed daily for 14 days.

**Oral:** No symptoms were observed at 40 mg/kg; at 80 mg/kg an increase of heart rate was seen; at higher doses vomiting was observed; in this study the non-lethal oral dose was approximately 320 mg/kg, 800 times the EMCD.

**IV:** At 70 mg/kg salivation and hematuria were observed; at 125 mg/kg the treated male died, thus the maximum non-lethal IV dose in these conditions, was 70 mg/kg, 175 times the EMCD. Administration of cetirizine pediatric solution and the corresponding vehicle to Beagle dogs at a single dose of 20 mg/kg produced no significant signs of toxicity.

## **Chronic Toxicity Studies**

### **1. Fifteen-day Study in Mice**

Fifteen-day oral studies comparing gavage dosing with diet dosing in mice (6/sex/level) at dose levels of 5, 10, 20, 40, 80, or 160 mg/kg/day revealed that similar hepatic findings, consisting of increased liver weights and periacinar hepatocytic hypertrophy, were induced by both routes of administration. The findings were similar in character, incidence and severity. Periacinar hepatocytic steatosis occurred in only 3 mice (2 male, 1 female) at 160 mg/kg by gavage dosing.

### **2. Four-Week Study in Mice**

Dietary administration of cetirizine hydrochloride to mice (16/sex/level) at dose levels of 1, 3, 9, 27 or 81 mg/kg/day for 4 weeks resulted in hepatic changes which were more pronounced in males than in female mice. In males, treatment resulted in minimal to moderate centrilobular hepatic hypertrophy at dose levels of 3 mg/kg or greater.

In females, treatment resulted in microsomal enzyme induction at dose levels of 9 mg/kg or greater, and elevated serum triglyceride levels at 27 and 81 mg/kg. Increased liver weights, hepatic hypertrophy and/or steatosis did not occur in female mice.

Electron microscopical examination of the livers of male mice revealed a moderate or minimal proliferation of smooth endoplasmic reticulum and apparent relative decrease in the amount of rough endoplasmic reticulum in centrilobular hepatocytes in all male mice receiving 27mg/kg/day. Smooth endoplasmic reticulum proliferation was also present in 5/10 male mice receiving 9 mg/kg/day.

### **3. Four-Week Reversibility Study in Mice**

In order to determine the reversibility of the hepatic changes in mice, cetirizine hydrochloride was administered to male and female mice (36/sex/level) by dietary admixture at dose levels of 40, or 160 mg/kg/day for 4 weeks followed by recovery periods of 4 and 13 weeks. After 4 weeks of cetirizine hydrochloride treatment, hepatic and/or hepatic-related changes similar to those observed in previously conducted mouse toxicity studies were induced.

Following the 4-week recovery period, the serum biochemical parameters, hepatic lipid levels, microsomal drug metabolizing enzyme activities, and liver electron microscopic results were essentially similar to controls. Although still evident at this time period, the increased liver weights, as well as the macroscopic and microscopic liver findings indicated a trend towards reversibility. By 13 weeks of recovery, the hepatic changes were no longer apparent.

### **4. Fifteen-day Study in Rats**

Fifteen-day oral studies comparing gavage dosing with diet dosing in rats (6/sex/level) at dose levels of 5, 10, 20, 40, 80, or 160 mg/kg/day revealed that hepatic changes, consisting of liver weight increases, periacinar hepatocytic hypertrophy and large droplet hepatic steatosis, were induced by both routes of administration. However, the findings occurred more frequently and tended to be more severe with dietary dosing.

### **5. One-Month Gavage Study in Rats**

In a preliminary study to investigate target organ toxicity, cetirizine hydrochloride was

administered to rats (10/sex/level) by oral gavage for 1 month at dose levels of 25, 75, and 225 mg/kg/day. At 225 mg/kg, treatment was associated with increased serum alpha-2-globulin values and increased liver weights in male and female, increased serum cholesterol values in female rats, and hepatic steatosis and necrosis in all 10 male rats. Hepatic steatosis and necrosis were also reported in 3 male rats at 75 mg/kg. There were no significant treatment-related elevations in serum enzyme levels in male rats, including alkaline phosphatase, LDH, ASAT, ALAT, and sorbitol dehydrogenase (SDH). All H&E and Oil Red O stained liver sections from male and female animals were re-evaluated. Results indicate midzonal and/or centrilobular hepatic hypertrophy in male and female rats at 75 and 225 mg/kg; hepatic steatosis in 2 males at 75 mg/kg and 10 males and 1 female rat at 225 mg/kg, and an area of centrilobular necrosis in one lobe from 1 male rat at 225 mg/kg. Centrilobular necrosis was not evident in male animals at the mid-dose or in 9/10 male rats at the high-dose.

#### **6. Four-Week Diet Study in Rats**

Dietary administration of cetirizine hydrochloride to rats (16/sex/level) at dose levels of 2, 6, 18, 54, or 160 mg/kg/day for 4 weeks resulted in hepatic changes which, as in oral gavage studies, were more pronounced in males than in females. At 54 and 160 mg/kg, increased liver weights were reported in male rats, and centrilobular hepatic hypertrophy, microsomal enzyme induction, and mid-zonal/centrilobular hepatic steatosis were observed in male and female rats. Other findings included lower food consumption for females at 18 (94.3%), 54 (94.7%) and 160 (92.1%) mg/kg, and lower body weight gains in male and female rats at 160 mg/kg. A slight increase in erythropoietic activity in the spleen, characterized by increases in erythrocytic parameters and white blood cells, increased spleen weights, and a minimally increased cellularity of the splenic red pulp, was observed in male treated rats, particularly at 54 or 160 mg/kg. However, there were no associated microscopic bone marrow changes.

#### **7. Four-Week Reversibility Study in Rats**

In order to determine the reversibility of the hepatic changes in rats, cetirizine hydrochloride was administered to male and female rats (30/sex/level) by dietary admixture at dose levels of 40, or 160 mg/kg/day for 4 weeks followed by recovery periods of 4 and 13 weeks. After 4 weeks of cetirizine hydrochloride treatment, hepatic and/or hepatic-related changes similar to those observed in previously conducted rat toxicity studies were induced. Following the 4-week recovery period, serum biochemical parameters, hepatic lipid levels, microsomal drug metabolizing enzyme activities, liver weights, liver electron microscopic findings, and hepatic macroscopic and microscopic findings were similar to controls, indicating a reversibility of all observed liver and liver-related effects.

#### **8. Six-Month Gavage Study in Rats**

Oral administration of cetirizine hydrochloride to rats (25/sex/level) by gavage at dose levels of 8, 25, or 75 mg/kg/day for 6 months produced hepatic changes consisting of hypertrophy, increased liver weights, altered serum biochemical values (after 3 months treatment) and steatosis. The modifications were less pronounced after 6 months than at 3 months and males were affected to a greater extent than females.

At the 3-month interim sacrifice, hepatic hypertrophy was observed in both sexes at all treatment levels. The degree was dose-related, ranging from minimal to moderate. Increased liver weights

were reported in males at 75 mg/kg and in females at 25 and 75 mg/kg. Minimal to slight hepatic steatosis was observed in 1 male at 25 mg/kg and in 3 males and 1 female at 75 mg/kg. The steatosis was associated with slightly reduced serum triglyceride levels at 75 mg/kg. Serum SDH levels were elevated in males at all treatment levels and in females at 75 mg/kg.

After 6 months treatment, the incidence of hepatic hypertrophy was lower than reported at 3 months, and the incidence of hepatic steatosis was similar to 3-month results. Results of electron microscopic examination of livers from several control and high-dose treated animals revealed definite smooth endoplasmic reticulum (SER) proliferation in male treated rats and a slight SER proliferation in 1 female treated rat.

### **9. One-Month Study in Dogs**

Cetirizine hydrochloride was administered orally (capsules) to beagle dogs (3/sex/level) at dosage levels of 15, 45 or 135 mg/kg/day for 1 month. At 15 and 45 mg/kg, cetirizine hydrochloride was well tolerated with only an increased incidence of vomiting, 2.5% and 7.1%, respectively, above control incidence (0.6%) reported. At 135 mg/kg, treatment resulted in an increased incidence of vomiting (17.2%); body tremor; salivation; ataxia; body weight loss and reduced food intake in 2 female dogs at the end of the treatment period; decreased mean urine specific gravity associated with an increased urine volume; a slight increase in mean serum alkaline phosphatase and a slight increase in alpha-2-globulins for females at 4 weeks. Histopathological examination of tissues from all dogs on study revealed no treatment-related alterations.

### **10. Six-Month Study in Dogs**

In a 6-month study (with a 3-month interim sacrifice), cetirizine hydrochloride was administered orally (capsules) to beagle dogs (5/sex/level) at dosage levels of 8, 25, or 75 mg/kg/day. At 8 mg/kg for 6 months, 20 times the EMCD, cetirizine hydrochloride administration was well tolerated with only a very slight increased incidence of vomiting (0.8%) over control incidence (0.3%) reported. At 25 and 75 mg/kg, 62.5 and 187.5 times the EMCD, cetirizine hydrochloride treatment resulted in an increased incidence of vomiting, 1.6% and 4.0% respectively, and decreased body weight gains in female dogs, 27% and 47% respectively, after 27 weeks of treatment. In addition, at 75 mg/kg, 1 male dog died and a 2nd male dog became moribund and was sacrificed during the study. Histopathological examination of tissues did not reveal any treatment-related lesions.

### **11. Two-Week Study in Monkeys**

In a 2-week duration study, cetirizine hydrochloride was administered by oral gavage to cynomolgus monkeys (1/sex/level) at dose levels of 50, 100, or 200 mg/kg/day. At 200 mg/kg, vomiting, salivation, and other signs indicative of a debilitating condition were reported. Both monkeys at this level exhibited a progressive loss in body weight and a marked reduction in food consumption during the treatment period. One monkey each at 50 (124 g) and 100 mg/kg (183 g) also exhibited a body weight loss during the treatment period. Histopathological examinations revealed minimal to moderate fatty infiltration in centrilobular hepatocytes from both monkeys at 200 mg/kg.

### **12. Four-Week Study in Monkeys**

In a 4-week study, cetirizine hydrochloride was administered by oral gavage to cynomolgus

monkeys (3/sex/level) at doses of 17, 50 or 150 mg/kg/day. Vomiting, huddled posture, poor coat condition, limb tremors, abnormal scratching motions and a reduced body temperature were observed in animals treated at 150 mg/kg. One female monkey at 150 mg/kg became moribund and was sacrificed on day 22 of dosing. This animal exhibited a debilitated body condition; a marked reduction in food consumption, an overall body weight loss of 622 g; elevated serum urea, GPT and GOT levels with decreased serum calcium and triglycerides, and a slow heart rate with sinus arrhythmia noted from an ECG recording made immediately prior to sacrifice. No treatment-related microscopic findings were reported. Marked body weight losses were recorded for the majority of monkeys at 150 mg/kg. Results of laboratory investigations performed during week 4 revealed decreased Ornithine Carbomoyltransferase (OCT) and SDH levels for the groups receiving 50 or 150 mg/kg, and increased triglyceride levels for the group receiving 150 mg/kg. No changes in the activities of measured hepatic microsomal enzymes were detected and no treatment-related microscopic abnormalities were observed.

### **13. One-Year Study in Dogs**

The oral (capsule) administration of cetirizine hydrochloride to beagle dogs (5/sex/level) at dose levels of 4, 15, or 60 mg/kg/day for 52 weeks was well tolerated and did not produce any significant toxicological findings. A dose-related increased incidence of vomiting, up to 6.3% at 60 mg/kg, within 1-hour of dose administration occurred in all treatment groups. There were no other treatment-related clinical signs. At necropsy, hepatic drug metabolizing enzyme activities were evaluated for each dog. Cetirizine hydrochloride, at dose levels up to 60 mg/kg/day, did not cause any induction of hepatic microsomal drug metabolizing enzymes, microsomal protein levels, or cytochrome P-450.

### **14. One-Year Study in Monkeys**

The oral (gavage) administration of cetirizine hydrochloride to cynomolgus monkeys (5/sex/level) at dose levels of 5, 15 or 45 mg/kg/day for 52 weeks was well tolerated and did not produce any significant toxicological findings. A dose-related increased incidence of salivation at or just after dose administration was reported in all dose groups. At necropsy, hepatic drug metabolizing enzyme activities were evaluated for each monkey. Cetirizine hydrochloride, at dose levels up to 45 mg/kg/day, did not cause any induction of hepatic microsomal drug metabolizing enzymes, microsomal protein levels, or cytochrome P-450.

## **Carcinogenicity**

### **1. Two-Year Study in Mice**

Dietary administration of cetirizine hydrochloride to mice (52/sex/level) at dose levels of 1, 4, or 16 mg/kg/day for 104 weeks, produced no evidence of a carcinogenic potential at doses 40 times the maximum clinically studied human daily dose (20 mg).

### **2. Two-Year Study in Rats**

Dietary administration of cetirizine hydrochloride to rats (50/sex/level) at dose levels of 3, 8, or 20 mg/kg/day for 104 weeks produced no evidence of a carcinogenic potential at doses 50 times higher than the maximum clinically studied human daily dose.

Non-neoplastic treatment-related microscopic findings consisted of a tendency towards an increased incidence of centrilobular vacuolation and fat deposition in the liver in male rats at 8 and



20 mg/kg, and of a slight, not dose-related, increased incidence of ulceration of the non-glandular stomach in female rats.

## **Teratology and Reproduction**

### **1. Reproduction and Fertility Study in Mice**

Cetirizine hydrochloride was administered by oral gavage at dose levels of 0, 4, 16 and 64 mg/kg/day to groups of 20 male and 40 female COBS CD-1 mice, in a reproduction and fertility study. There were no effects on male and female fertility or reproductive performance, or on pup development through 2 generations at oral doses up to 16 mg/kg, 40 times the expected maximum clinical dose (EMCD) of 20 mg.

### **2. Teratology**

**a. Teratology Study in Mice:** Cetirizine hydrochloride was administered by oral gavage at dose levels of 6, 24, and 96 mg/kg/day to groups of 30 time-mated COBS CD-1 female mice from day 6 to day 15 of gestation. Cetirizine hydrochloride at dose levels up to 96 mg/kg/day from gestation days 6 through 15 was not embryo-feto-toxic nor teratogenic.

**b. Teratology Study in Rats:** Cetirizine hydrochloride, administered by oral gavage at dose levels of 8, 25, 75 and 225 mg/kg/day to mated Sprague Dawley female rats (25 - 26 animals/level) from day 6 to day 15 of gestation, was not teratogenic. The incidence of the major malformations was not dose-related and the 2 fetuses exhibiting these malformations were both runts, 1 at 8 mg/kg (agnathia with displacement of the eyes and ears and left microphthalmia) and 1 at 225 mg/kg (left microphthalmia). Although the limited in-house historical data on this species of rat (1225-2800 fetuses) did not report agnathia or microphthalmia, these types of malformations have been reported at a low incidence in control data for CD rats. The no-effect level for maternal toxicity was 25 mg/kg, and the no-effect level for embryo-feto-toxicity, although not clearly established, was approximately 8 mg/kg. At 8 mg/kg, the incidence of reduced ossification of parietal, interparietal, and hyoid cranial bones was slightly higher than control incidence, but, considered to be within normal variability.

**c. Teratology Study in Rabbits:** Cetirizine hydrochloride, administered by oral gavage at dose levels of 15, 45, and 135 mg/kg/day to mated New Zealand White female rabbits 16/level at 15 and 45 mg/kg; 18/level at 135 mg/kg; 17 in the control group) from day 6 to day 18 of gestation was not teratogenic. The no-effect level for maternal toxicity and embryo-feto-toxicity was 15 mg/kg, 37.5 times the EMCD. At 15 mg/kg, maternal body weight gain was slightly decreased during the post-treatment period.

**d. Conclusion:** The above described anomalies, irregularly found in all cetirizine hydrochloride treated groups, did not occur in a dose-related fashion; moreover, these sorts of anomalies are known to occur spontaneously in untreated animal populations. In addition, many of the anomalies observed occurred in small fetuses, and at doses associated with maternal toxicity. Consequently a definitive causal relationship with cetirizine hydrochloride cannot be ruled out.

### **3. Peri- and Post-Natal Development Study in Mice**

Cetirizine hydrochloride was administered by oral gavage to groups of 32 time-mated COBS CD-1

female mice at dose levels of 0, 6, 24 or 96 mg/kg/day from day 15 of gestation and continued up to sacrifice of the dams on, or shortly after, day 21 post partum (weaning). Cetirizine hydrochloride, at dose levels of 6 and 24 mg/kg/day, up to 60 times the EMCD, from day 15 of gestation to weaning of pups, did not produce any adverse effect on perinatal conditions or progeny development. At 96 mg/kg, cetirizine hydrochloride treatment was associated with slight maternal effects and lower mean pup weights after birth, at 4 to 21 days of lactation.

### **Mutagenicity**

The mutagenic potential of cetirizine hydrochloride was assessed in in vitro non-mammalian cell systems as well as in in vitro and in vivo mammalian cell systems. Cetirizine hydrochloride was not mutagenic.

### **Toxicology Summary**

The principle findings in rodent subchronic oral toxicity studies were related to the liver and consisted of hypertrophy of hepatocytes, proliferation of smooth endoplasmic reticulum (SER), microsomal enzyme induction, increased liver weights, hepatic steatosis, hepatic necrosis, elevated or reduced serum triglyceride levels, and increased serum GPT, OCT and SDH values. Of these findings, the SER proliferation associated with microsomal enzyme induction and hepatic hypertrophy followed by increased liver weights are probably pharmacological responses to cetirizine hydrochloride treatment rather than toxicological. The hepatotoxic findings consisting of hepatic steatosis and necrosis, and altered biochemical parameters appear to be related to the marked hepatic metabolism of cetirizine hydrochloride in rodents. Significant safety margins, calculated for rodent hepatotoxicity, ranged from 20 to greater than 370 times the EMCD of 20 mg depending on species, route of administration, and duration of treatment.

Similar liver-related findings were not evident in dogs receiving cetirizine hydrochloride orally for 1 month at doses up to 338 times the EMCD or for 6 months and 1 year respectively at doses up to 188 and 150 times the EMCD, nor were liver-related changes observed in cynomolgus monkeys receiving cetirizine hydrochloride for 1 month and 1 year respectively at doses up to 375 and 112.5 times the EMCD.

The dietary administration of cetirizine hydrochloride to mice at doses up to 16 mg/kg/day, 40 times the EMCD, and to rats at doses up to 20 mg/kg/day, 50 times the EMCD, for 104 weeks showed no indications of carcinogenic potential.

Re-analysis of the data demonstrated that no adverse effects on embryo-fetal viability, body weight or morphology were produced by maternally toxic dosages in development toxicity (Segment II) studies in the rat (225 mg/kg/day, 563 times the maximum clinically studied human dose), rabbit (135 mg/kg/day, 338 times the maximum clinically studied human dose) and mouse (96 mg/kg/day, 240 times the maximum clinically study human dose.)

Cetirizine is a major human metabolite of hydroxyzine (50 mg hydroxyzine = 20 mg cetirizine). Thus, the long-term experience with hydroxyzine also provides an indication of the safety of cetirizine in pregnancy. During 30 years of clinical use, hydroxyzine has not been associated with an increase of any human congenital malformation above the expected background incidence. Thus, human exposure to cetirizine has occurred for more than 30 years without any indication

that it or its parent compound, hydroxyzine, is a human teratogen. The effect of hydroxyzine on human pregnancies have been studied in a large epidemiology study [the Collaborative Perinatal Project]. The study did not report any increase in human congenital malformation as a consequence of the use of hydroxyzine.

The only other reported effect of hydroxyzine on pregnancy in a laboratory species was abortion in rhesus monkeys at dosages of 5 to 12 mg/kg. Three abortions and 2 normal offspring were produced after administration of 5 to 12 mg/kg dosages during organogenesis. The rhesus monkey is known to have a high incidence of abortion. The absence of expected control procedures in this old study, and the use of only 5 animals precludes drawing a causal relationship of this observation with hydroxyzine.

## REFERENCES

1. Bousquet J, Duchateau J, et al. Improvement of quality of life by treatment with cetirizine in patients with perennial allergic rhinitis as determined by a French version of the SF-36 questionnaire. *J Allergy Clin Immunol* 1996; 98(2):309-316.
2. Burtin B, Duchateau J, Pignat JC, Donnelly F, Bousquet J. Further improvement of quality of life by cetirizine in perennial allergic rhinitis as a function of treatment duration. *J Invest Allergy Clin Immunol* 2000; 10(2): 66-70.
3. Day JH, Briscoe M, Rafeiro E, Chapman D, Kramer B. Comparative onset of action and symptom relief with cetirizine, loratadine, or placebo in an environmental exposure unit in subjects with seasonal allergic rhinitis: confirmation of a test system. *Annals of Allergy, Asthma and Immunology* 2001; 87: 474-481.
4. Go MJT, Wuite J, Arendt C, Bernheim J. Double-blind, placebo controlled comparison of cetirizine and terfenadine in chronic idiopathic urticaria. *Acta Ther* 1989; 15(1):77-86.
5. Howarth PH, Stern M, Roi L, Reynolds R, Bousquet J. Double-blind, placebo-controlled study comparing the efficacy and safety of fexofenadine hydrochloride (120 and 180 mg once daily) and cetirizine in seasonal allergic rhinitis. *J Allergy Clin Immunol* 1999; 104: 927-33.
6. Hyo S, Fujieda S, Kawada R, Kitazawa S, Takenaka H. The efficacy of short-term administration of 3 antihistamines vs placebo under natural exposure to Japanese cedar pollen. *Ann Allergy Asthma Immunol.* 2005; 94:457-464.
7. Juhlin L, Arendt C. Treatment of chronic urticaria with cetirizine dihydrochloride non-sedating antihistamine. *Br J Dermatol* 1988; 119(1):67-72.
8. Juhlin L, De Vos C, Rihoux JP. Inhibiting effect of cetirizine on histamine-induced and 48/80-induced wheals and flares, experimental dermographism, and cold-induced urticaria. *J Allergy Clin Immunol* 1987; 80(4):599-602.
9. Kontou-Fili K, Paleologos G, Herakleous M. Suppression of histamine-induced skin reactions by loratadine and cetirizine diHCl. *Eur J Clin Pharmacol* 1989; 36:617-9.
10. Matzke GR, Yeh J, Awni WM, Halstenson CE, Chung M. Pharmacokinetics of cetirizine in the elderly and patients with renal insufficiency. *Ann Allergy* 1987; 59(6 Pt 2):25-30.
11. Murray JJ, Nathan RA, Bronsky EA, Olufade, AO, Chapman, D. Kramer B. Comparative Evaluation of Cetirizine in the Management of Seasonal Allergic Rhinitis: Impact on Symptoms, Quality of Life, Productivity, and Activity Impairment. *Allergy and Asthma Proc.* 2002; 23(6): 391-398.

12. Noonan MJ, Raphael GD, et al: The health-related quality of life effects of once-daily cetirizine HCl in patients with seasonal allergic rhinitis: A randomized double-blind, placebo-controlled trial. *Clinical & Experimental Allergy* 2003; 33(3):351-358.
13. Panayotopoulos SM, Panayotopoulou ES. Efficacy of cetirizine in the treatment of seasonal allergic rhinoconjunctivitis. *Ann Allergy* 1990; 65(2):146-8.
14. Seidel WF, Cohen S, Bliwise NG, Dement WC. Cetirizine effects on objective measures of daytime sleepiness and performance. *Ann Allergy* 1987; 59(6 Pt 2):58-62.
15. Snyder SH, Snowman AM. Receptor effects of cetirizine. *Ann Allergy* 1987; 59(6 Pt 2):4-8.
16. Wang DY, Hanotte F, De Vos C, Clement P. Effect of cetirizine, levocetirizine, and dextrocetirizine on histamine-induced nasal response in healthy adult volunteers. *Allergy* 2001; 56: 339-343.
17. Wood SG, John BA, Chasseaud LF, Yeh J, Chung M. The metabolism and pharmacokinetics of <sup>14</sup>C-cetirizine in humans. *Ann Allergy* 1987; 59(6 Pt 2):31-4.
18. Product Monograph for REACTINE® 20 mg Tablet, Date of revision: April 6, 2020, Control No: 235677, McNeil Consumer Healthcare, division of Johnson & Johnson Inc., Markham, Canada.

## PART III: CONSUMER INFORMATION

### PrNAT-CETIRIZINE

#### (Cetirizine Hydrochloride Tablets Natco Standard)

This leaflet is part III of a three-part "Product Monograph" published when NAT-CETIRIZINE was approved for sale in Canada and is designed specifically for Consumers. This leaflet is a summary and will not tell you everything about NAT-CETIRIZINE. Contact your doctor or pharmacist if you have any questions about the drug.

#### ABOUT THIS MEDICATION

##### What the medication is used for:

NAT-CETIRIZINE 20 mg tablets are used in adults and children 12 years of age and older for the fast, 24-hour relief of nasal and non-nasal seasonal (e.g. trees, grass, pollen, ragweed/hay fever) and year-round (e.g. dust, animal dander, mold) allergy symptoms, including: sneezing, runny nose, post-nasal discharge, nasal congestion/stuffiness, itchy, watery eyes, itchy nose/throat, and itching and hives due to allergic skin reactions.

##### What it does:

Your allergy symptoms are simply your body's over reaction in trying to protect you from allergens such as dust, ragweed, grass and tree pollen, animal dander or mold. When allergens are detected, the cells in your body release a chemical called histamine which binds to specific histamine receptors in your skin and tissues. The resulting reaction causes itchy, watery eyes, sneezing and runny nose. NAT-CETIRIZINE helps relieve your allergy symptoms by blocking these receptor sites before histamine binds there, preventing or reducing many of the symptoms of an allergic reaction. The anti-inflammatory properties of NAT-CETIRIZINE also help by reducing swelling and related symptoms, including redness and hives (red, itchy bumps or welts on your skin).

##### When it should not be used:

Do not take NAT-CETIRIZINE if you:

- are allergic (hypersensitive) to cetirizine hydrochloride, hydroxyzine, piperazine derivatives or any of the other ingredients in NAT-CETIRIZINE
- have serious kidney problems

##### What the medicinal ingredient is:

Cetirizine hydrochloride

##### What the nonmedicinal ingredients are:

Microcrystalline cellulose, croscovidone, colloidal anhydrous silica, magnesium stearate, hypromellose, titanium dioxide and polyethylene glycol.

##### What dosage forms it comes in:

Tablets: 20 mg

#### WARNINGS AND PRECAUTIONS

BEFORE you use NAT-CETIRIZINE 20 mg tablets, talk to your doctor or pharmacist if you:

- have kidney or liver problems
- are pregnant or planning to become pregnant
- are breastfeeding or planning to breastfeed
- are 65 years of age or older

**Driving and using machines :** NAT-CETIRIZINE can cause drowsiness. If drowsiness occurs, do not drive or operate machinery.

#### INTERACTIONS WITH THIS MEDICATION

The following may interact with NAT-CETIRIZINE:

- alcohol
- other substances that cause sedation such as anti-anxiety medications, sleep aids, antihistamines, antidepressants, muscle relaxants and pain medication.

#### PROPER USE OF THIS MEDICATION

NAT-CETIRIZINE 20 mg tablets can be taken with or without food.

##### Usual dose:

**Children 12 years of age and over and Adults:** One tablet once per day.

NAT-CETIRIZINE 20 mg tablets should **not** be taken by:

- **Adults 65 years of age and over**
- **Patients with liver or kidney problems**
- **Children less than 12 years of age**

##### Overdose:

If you think you have taken too much NAT-CETIRIZINE contact your healthcare professional, hospital emergency department or regional Poison Control Centre immediately, even if there are no symptoms.

Symptoms of an overdose include: confusion, diarrhea, dizziness, sleepiness, headache, fatigue, feeling of bodily discomfort, large pupils, itching, restlessness, sedation, lowered level of consciousness, rapid heartbeat, high blood pressure, tremor and inability to urinate. Extreme activity and extreme lack of energy have been seen in children.

##### Missed Dose:

If you miss taking your dose on time, do not worry; take your dose when you remember. Do not take more than one dose in 24 hours.

#### SIDE EFFECTS AND WHAT TO DO ABOUT THEM

Side effects may include:

- headache
- sleepiness, fatigue
- dry mouth
- loss of taste
- nausea
- dizziness
- trouble sleeping (insomnia)
- eye pain or swelling
- blurred vision
- erectile dysfunction

<b>SERIOUS SIDE EFFECTS, HOW OFTEN THEY HAPPEN AND WHAT TO DO ABOUT THEM</b>			
Symptom/effect	Talk with your doctor or pharmacist		Stop taking drug and seek immediate medical help
	Only if severe	In all cases	
<b>VERY RARE:</b>			
Agitation, aggressiveness		✓	
<b>Allergic Reaction:</b> rash, hives, swelling of the face, lips, tongue or throat, difficulty swallowing or breathing			✓
Difficulty urinating		✓	
<b>Hallucinations:</b> seeing or hearing things that are not really there			✓
<b>Liver problems:</b> yellowing of the skin or eyes		✓	
Memory problems, partial or total memory loss		✓	
New rash or itching after stopping drug		✓	
Restlessness with increased body movement		✓	
<b>Severe Skin Reaction:</b> rash with small pimples and redness on the skin folds, trunk and arms, with or without fever			✓

*This is not a complete list of side effects. For any unexpected effects while taking NAT-CETIRIZINE, contact your doctor or pharmacist.*

### **HOW TO STORE IT**

Store at temperatures between 15°C and 30°C. Protect from moisture.

Keep out of reach and sight of children.

### **REPORTING SIDE EFFECTS**

You can report any suspected side effects associated with the use of health products to Health Canada by:

- Visiting the Web page on Adverse Reaction Reporting (<https://www.canada.ca/en/health-canada/services/drugs-health-products/medeffect-canada/adverse-reaction-reporting.html>) for information on how to report online, by mail or by fax; or
- Calling toll-free at 1-866-234-2345.

*NOTE: Contact your health professional if you need information about how to manage your side effects. The Canada Vigilance Program does not provide medical advice.*

### **MORE INFORMATION**

**If you want more information about NAT-CETIRIZINE:**

- Talk to your healthcare professional
- Find the full product monograph that is prepared for healthcare professionals and includes this Consumer Information by visiting the Health Canada website (<https://health-products.canada.ca/dpd-bdpp/index-eng.jsp>); the manufacturer's website [www.natcopharma.ca](http://www.natcopharma.ca), or by calling 1-800-296-9329.

This leaflet is prepared by:  
 Natco Pharma (Canada) Inc.  
 2000 Argentia Road, Plaza 1, Suite 200  
 Mississauga, Ontario  
 L5N 1P7, Canada

Last revised: March 03, 2021