

PRODUCT MONOGRAPH

Pr SANDOZ SAXAGLIPTIN

Saxagliptin Tablets

Saxagliptin (as saxagliptin hydrochloride dihydrate)

2.5 mg and 5 mg

Oral Antihyperglycemic Agent

DPP-4 inhibitor

Incretin Enhancer

Sandoz Canada Inc.
110 Rue de Lauzon
Boucherville, Québec
J4B 1E6

Date of Revision:
February 25, 2019

Submission Control No: 224039

Table of Contents

PART I: HEALTH PROFESSIONAL INFORMATION.....3
SUMMARY PRODUCT INFORMATION3
INDICATIONS AND CLINICAL USE.....3
CONTRAINDICATIONS4
WARNINGS AND PRECAUTIONS.....4
ADVERSE REACTIONS.....7
DRUG INTERACTIONS15
DOSAGE AND ADMINISTRATION19
OVERDOSAGE20
ACTION AND CLINICAL PHARMACOLOGY20
STORAGE AND STABILITY23
DOSAGE FORMS, COMPOSITION AND PACKAGING24

PART II: SCIENTIFIC INFORMATION.....25
PHARMACEUTICAL INFORMATION.....25
CLINICAL TRIALS26
DETAILED PHARMACOLOGY38
TOXICOLOGY38
REFERENCES40

PART III: CONSUMER INFORMATION.....41

Pr SANDOZ SAXAGLIPTIN
Saxagliptin Tablets
Saxagliptin (as saxagliptin hydrochloride dihydrate)
2.5 mg and 5 mg

PART I: HEALTH PROFESSIONAL INFORMATION

SUMMARY PRODUCT INFORMATION

Route of Administration	Dosage Form / Strength	Clinically Relevant Nonmedicinal Ingredients
oral	Tablet / 2.5 mg, 5 mg	Lactose monohydrate, lactose anhydrous. <i>For a complete listing see Dosage Forms, Composition and Packaging section.</i>

INDICATIONS AND CLINICAL USE

Combination with metformin:

Sandoz Saxagliptin is indicated in patients with type 2 diabetes mellitus to improve glycemic control in combination with metformin when metformin used alone, with diet and exercise, does not provide adequate glycemic control (see CLINICAL TRIALS).

Combination with sulfonylurea:

Sandoz Saxagliptin is indicated in patients with type 2 diabetes mellitus to improve glycemic control in combination with sulfonylurea when sulfonylurea used alone, with diet and exercise, does not provide adequate glycemic control (see CLINICAL TRIALS).

Combination with insulin (with or without metformin):

Sandoz Saxagliptin is indicated in patients with type 2 diabetes mellitus to improve glycemic control in combination with premixed or long/intermediate acting insulin (with or without metformin) when premixed or long/intermediate acting insulin (with or without metformin) used alone, with diet and exercise, do not provide adequate glycemic control (see CLINICAL TRIALS).

Combination with metformin and a sulfonylurea:

Sandoz Saxagliptin is indicated in patients with type 2 diabetes mellitus to improve glycemic control in combination with metformin and a sulfonylurea when dual therapy with these two agents, with diet and exercise, does not provide adequate glycemic control (see CLINICAL TRIALS).

Geriatrics (≥ 65 years of age):

No dosage adjustment is required based on age, however greater sensitivity of some older individuals cannot be ruled out (see WARNINGS AND PRECAUTIONS, DOSAGE AND ADMINISTRATION and ACTION AND CLINICAL PHARMACOLOGY).

Pediatrics (< 18 years of age): Safety and effectiveness of saxagliptin in pediatric patients have not been established. Therefore, Sandoz Saxagliptin should not be used in this patient population.

CONTRAINDICATIONS

Patients who have had a history of any hypersensitivity reaction, including anaphylaxis or angioedema, to saxagliptin or to another DPP-4 inhibitor, or to any ingredient in the formulation (see WARNINGS AND PRECAUTIONS, Hypersensitivity Reactions and ADVERSE REACTIONS, Post-Market Adverse Drug Reactions). For a complete listing, see DOSAGE FORMS, COMPOSITION AND PACKAGING.

Sandoz Saxagliptin (saxagliptin) is contraindicated in patients with diabetic ketoacidosis, diabetic coma/precoma or Type I diabetes mellitus. These conditions should be treated with insulin.

WARNINGS AND PRECAUTIONS

Cardiovascular

Patients with Congestive Heart Failure: In a post-market placebo-controlled cardiovascular outcomes trial (SAVOR), hospitalization for heart failure occurred at a greater rate in the saxagliptin group (3.5%) compared to the placebo group (2.8%) [HR =1.27; 95% confidence interval 1.07, 1.51]. In the SAVOR trial, 2105 (12.8%) patients had a history of congestive heart failure, of whom 1056 were randomized to saxagliptin treatment. Caution is warranted if Sandoz Saxagliptin is used in patients with history of congestive heart failure (especially in those patients who also have renal impairment and/or history of myocardial infarction [MI]). During therapy with Sandoz Saxagliptin, patients should be observed for signs and symptoms of heart failure. Patients should be advised of characteristic symptoms of heart failure, and to immediately report such symptoms. If heart failure develops, discontinue Sandoz Saxagliptin and manage according to current standards of care (see ADVERSE REACTIONS, Post-Marketing, Cardiovascular Safety).

Endocrine and Metabolism

A lower dose of sulfonylurea or insulin may be required to reduce the risk of hypoglycemia when used in combination with Sandoz Saxagliptin (see ADVERSE REACTIONS, Clinical Trial Adverse Drug Reactions and DOSAGE AND ADMINISTRATION).

Loss of control of blood glucose: When a patient stabilized on Sandoz Saxagliptin is exposed to stress such as fever, trauma, infection, or surgery, a loss of control of blood glucose may occur. At such times, it may be necessary to temporarily discontinue Sandoz Saxagliptin and administer insulin.

Musculoskeletal and connective tissue disorders

Severe and Disabling Arthralgia: Severe and disabling arthralgias have been reported post-marketing in patients taking saxagliptin or other DPP-4 inhibitors. Onset of symptoms following initiation of drug therapy varied from one day to years. Saxagliptin is considered a possible cause for severe joint pain. Patients experienced relief of symptoms upon discontinuation of the

medication and some experienced recurrence of symptoms with reintroduction of saxagliptin or another DPP-4 inhibitor. If a patient treated with Sandoz Saxagliptin, presents with severe joint pain, discontinuation of Sandoz Saxagliptin and replacement with other antidiabetic medications should be considered (see ADVERSE REACTIONS, Post-Market Adverse Drug Reactions).

Use with potent CYP 3A4 inducers

Using CYP3A4 inducers like carbamazepine, dexamethasone, phenobarbital, phenytoin, and rifampin may reduce the glycemic lowering effect of saxagliptin (see DRUG INTERACTIONS).

Lactose

Sandoz Saxagliptin tablets contain lactose monohydrate. Patients with hereditary problems of galactose intolerance, the Lapp lactase deficiency or glucose-galactose malabsorption should not take this product.

Hypersensitivity Reactions

There have been post-marketing reports of serious hypersensitivity reactions, including anaphylaxis and angioedema, in patients treated with saxagliptin and other members of this class. Exfoliative skin conditions including Stevens-Johnson syndrome have also been reported in patients treated with saxagliptin and other members of this class, although causality with saxagliptin has not been established. Onset of these reactions occurred within the first 3 months after initiation of the treatment, with some reports occurring after the first dose. If a hypersensitivity reaction to saxagliptin is suspected, discontinue Sandoz Saxagliptin, assess for other potential causes for the event, and institute alternative treatment for diabetes (see CONTRAINDICATIONS and ADVERSE REACTIONS, Post-Market Adverse Drug Reactions).

Pancreatitis

There have been post-marketing reports of acute and chronic pancreatitis in patients taking saxagliptin. Reports of fatal and non-fatal hemorrhagic or necrotizing pancreatitis were noted in patients taking other members of this class. After initiation of Sandoz Saxagliptin patients should be observed carefully for signs and symptoms of pancreatitis. If pancreatitis is suspected, Sandoz Saxagliptin should promptly be discontinued and appropriate management should be initiated. It is unknown whether patients with a history of pancreatitis are at increased risk for the development of pancreatitis while using Sandoz Saxagliptin. Risk factors for pancreatitis include a history of pancreatitis, gallstones, alcoholism, or hypertriglyceridemia.

Immune

Immunocompromised patients: A dose-related mean decrease in absolute lymphocyte count was observed with saxagliptin. When clinically indicated, such as in settings of unusual or prolonged infection, lymphocyte count should be measured. The effect of saxagliptin on lymphocyte counts in patients with lymphocyte abnormalities (e.g. human immunodeficiency virus) is unknown. See Clinical trial adverse drug reactions, Abnormal Hematologic and Clinical Chemistry Findings.

Immunocompromised patients, such as patients who have undergone organ transplantation or patients diagnosed with human immunodeficiency syndrome, have not been studied in the saxagliptin clinical program. Therefore, the efficacy and safety profile of saxagliptin in these patients has not been established.

Peri-operative considerations

See Endocrine and Metabolism section – Loss of control of blood glucose.

Skin

Ulcerative and necrotic skin lesions have been reported in monkeys in non-clinical toxicology studies (see Part II: TOXICOLOGY, Chronic Toxicity). Although skin lesions were not observed at an increased incidence in clinical trials, there is limited experience in patients with diabetic skin complications.

Rash is noted as an adverse event for saxagliptin (see ADVERSE REACTIONS, Clinical trial adverse drug reactions). In keeping with routine care of the diabetic patient, monitoring for skin disorders is recommended.

Bullous pemphigoid: Post-marketing cases of bullous pemphigoid requiring hospitalization have been reported with the use of Saxagliptin and other DPP-4 inhibitors. In reported cases, patients typically recovered with topical or systemic immunosuppressive treatment and discontinuation of the DPP-4 inhibitor.

Tell patients to immediately report development of blisters or erosions while receiving Sandoz Saxagliptin. If bullous pemphigoid is suspected, Sandoz Saxagliptin should be discontinued and referral to a dermatologist should be considered for diagnosis and appropriate treatment.

Special Populations

Pregnant Women: There are no adequate and well-controlled studies in pregnant women. As animal reproduction studies are not always predictive of human response, Sandoz Saxagliptin is not recommended for use in pregnancy (see TOXICOLOGY).

Nursing Women: Saxagliptin is secreted in the milk of lactating rats. It is not known whether saxagliptin is excreted in human milk. Therefore, Sandoz Saxagliptin should not be used by a woman who is nursing.

Pediatrics (< 18 years of age): Safety and effectiveness of saxagliptin in pediatric patients have not been established. Therefore, Sandoz Saxagliptin should not be used in this patient population.

Geriatrics (≥ 65 years of age): Of the total number of subjects (N=4148) studied in controlled clinical safety and efficacy studies of saxagliptin, 634 (15.3%) patients were 65 years and over, of which 59 (1.4%) patients were 75 years and over. No overall differences in safety or effectiveness were observed between subjects 65 years and over and younger subjects. While this clinical experience has not identified differences in responses between the elderly and younger patients, greater sensitivity of some older individuals cannot be ruled out.

Saxagliptin and its major metabolite are known to be eliminated in part by the kidney. Renal function should be assessed prior to initiating Sandoz Saxagliptin and periodically thereafter in geriatric patients because they are more likely to have decreased renal function. Care should be

taken in prescribing saxagliptin in this population based on renal function (see WARNINGS AND PRECAUTIONS, DOSAGE AND ADMINISTRATION and ACTION AND CLINICAL PHARMACOLOGY).

Hepatic Insufficiency: There are limited clinical data in patients with hepatic impairment taking multiple doses of saxagliptin. The use of Sandoz Saxagliptin in patients with moderate to severe hepatic impairment is not recommended (see ACTION AND CLINICAL PHARMACOLOGY).

Renal Impairment: Saxagliptin should be used with caution in patients with severe renal impairment, and is not recommended for use in patients with end-stage renal disease (ESRD) requiring hemodialysis. In patients with eGFR < 45 mL/min/1.73m², the dose is 2.5 mg once daily. (See DOSAGE AND ADMINISTRATION, Renal Impairment).

Assessment of renal function is recommended prior to initiation of Sandoz Saxagliptin, and periodically thereafter (see DOSAGE AND ADMINISTRATION, ACTION AND CLINICAL PHARMACOLOGY and CLINICAL TRIALS – Patients with Renal Impairment).

Monitoring and Laboratory Tests

Response should be monitored by periodic measurements of blood glucose and HbA_{1c} (A1C) levels. Assessment of renal function is recommended prior to initiation of Sandoz Saxagliptin and periodically thereafter. Patients with a history of heart failure or other risk factors for heart failure, including renal impairment, should be closely monitored for signs and symptoms of heart failure.

ADVERSE REACTIONS

Adverse Drug Reaction Overview

Saxagliptin was generally well tolerated in controlled clinical studies as an add-on to metformin, as an add-on to sulfonylurea, as an add-on to insulin (with or without metformin) and as an add-on to metformin and a sulfonylurea with the overall incidence of adverse events similar to that reported with placebo.

In a placebo-controlled clinical study of patients receiving saxagliptin 5 mg or placebo as an add-on to metformin, the incidence of serious adverse events was 9.9% and 5.6% respectively. The most commonly reported adverse events, reported regardless of causality and more common with saxagliptin than placebo, were nasopharyngitis and bronchitis. Discontinuation of therapy due to adverse events occurred in 7.3% and 4.5% of patients, respectively.

In a placebo-controlled clinical study of patients receiving saxagliptin 5 mg or placebo as an add-on to sulfonylurea (glyburide), the incidence of serious adverse events was 3.6% and 5.6% respectively. The most commonly reported adverse events, reported regardless of causality and more common with saxagliptin than placebo, were hypoglycemia and urinary tract infection. Discontinuation of therapy due to adverse events occurred in 4.7% and 3.4% of patients, respectively.

In a placebo-controlled clinical study of patients receiving saxagliptin 5 mg or placebo as an add-on to insulin (with or without metformin), the incidence of serious adverse events was 8.2% and 8.6%

respectively. The most commonly reported adverse events, reported regardless of causality and more common with saxagliptin than placebo, were headache and bronchitis. Discontinuation of therapy due to adverse events occurred in 3.0% and 2.0% of patients, respectively.

In a placebo-controlled clinical study of patients receiving saxagliptin 5 mg or placebo as an add-on to metformin and a sulfonylurea, the incidence of serious adverse events was 2.3% and 5.5% respectively. The most commonly reported adverse events, reported regardless of causality and more common with saxagliptin than placebo, were hypoglycemia, hypertension and diarrhea. Discontinuation of therapy due to adverse events occurred in 0.8 % and 2.3% of patients, respectively.

Clinical Trial Adverse Drug Reactions

Because clinical trials are conducted under very specific conditions the adverse reaction rates observed in the clinical trials may not reflect the rates observed in practice and should not be compared to the rates in the clinical trials of another drug. Adverse drug reaction information from clinical trials is useful for identifying drug-related adverse events and for approximating rates.

Adverse events, reported regardless of causality assessment, in $\geq 2\%$ of patients treated with either saxagliptin 5 mg or placebo as an add-on to metformin or add-on to sulfonylurea (glyburide) are shown in Table 1.

Table 1 Adverse Events (Regardless of Investigator Assessment of Causality) in the Add-on to Metformin^a Study and in Add-on to Sulfonylurea^b Study (24-week Short Term Study and the Long Term Extension) Reported in $\geq 2\%$ of Patients Treated with Either Saxagliptin 5 mg or Placebo in at Least One Study

Body System/Organ Class Adverse Event	Number of Patients (%) Add-on to Metformin		Number of Patients (%) Add-on to Sulfonylurea	
	Saxagliptin 5 mg N=191	Placebo N=179	Saxagliptin 5 mg N=253	Placebo N= 267
Blood and lymphatic system disorders				
Anemia	11 (5.8)	3 (1.7)	2 (0.8)	3 (1.1)
Eosinophilia	6 (3.1)	0	1 (0.4)	1 (0.4)
Cardiac disorders				
Coronary artery disease	4 (2.1)	0	1 (0.4)	1 (0.4)
Gastrointestinal disorders				
Diarrhea	14 (7.3)	23 (12.8)	13 (5.1)	23 (8.6)
Dyspepsia	11 (5.8)	8 (4.5)	9 (3.6)	7 (2.6)
Toothache	8 (4.2)	11 (6.1)	8 (3.2)	8 (3.0)
Abdominal pain	7 (3.7)	2 (1.1)	6 (2.4)	4 (1.5)
Abdominal pain upper	7 (3.7)	5 (2.8)	10 (4.0)	8 (3.0)
Nausea	7 (3.7)	8 (4.5)	4 (1.6)	4 (1.5)
Vomiting	7 (3.7)	7 (3.9)	4 (1.6)	4 (1.5)
Constipation	5 (2.6)	3 (1.7)	2 (0.8)	3 (1.1)
Gastroesophageal reflux disease	4 (2.1)	1 (0.6)	2 (0.8)	6 (2.2)

Body System/Organ Class Adverse Event	Number of Patients (%) Add-on to Metformin		Number of Patients (%) Add-on to Sulfonylurea	
	Saxagliptin 5 mg N=191	Placebo N=179	Saxagliptin 5 mg N=253	Placebo N= 267
Gastritis	2 (1.0)	2 (1.1)	5 (2.0)	8 (3.0)
General disorders and administration site conditions				
Edema peripheral	11 (5.8)	9 (5.0)	5 (2.0)	7 (2.6)
Chest pain	5 (2.6)	2 (1.1)	4 (1.6)	3 (1.1)
Fatigue	5 (2.6)	7 (3.9)	8 (3.2)	3 (1.1)
Asthenia	0	2 (1.1)	5 (2.0)	8 (3.0)
Infections and infestations				
Influenza	22 (11.5)	23 (12.8)	16 (6.3)	26 (9.7)
Nasopharyngitis	21 (11.0)	19 (10.6)	24 (9.5)	27 (10.1)
Bronchitis	18 (9.4)	11 (6.1)	8 (3.2)	7 (2.6)
Upper respiratory tract infection	17 (8.9)	14 (7.8)	22 (8.7)	22 (8.2)
Urinary tract infection	15 (7.9)	12 (6.7)	35 (13.8)	29 (10.9)
Sinusitis	10 (5.2)	9 (5.0)	4 (1.6)	3 (1.1)
Gastroenteritis	5 (2.6)	3 (1.7)	7 (2.8)	7 (2.6)
Tooth infection	5 (2.6)	3 (1.7)	2 (0.8)	2 (0.7)
Gastroenteritis viral	4 (2.1)	2 (1.1)	1 (0.4)	1 (0.4)
Pharyngitis	2 (1.0)	4 (2.2)	19 (7.5)	14 (5.2)
Viral infection	1 (0.5)	4 (2.2)	4 (1.6)	6 (2.2)
Pharyngotonsillitis	1 (0.5)	1 (0.6)	5 (2.0)	10 (3.7)
Injury, poisoning, and procedural complications				
Limb Injury	3 (1.6)	1 (0.6)	0	7 (2.6)
Investigations				
Blood creatine phosphokinase increased	4 (2.1)	2 (1.1)	8 (3.2)	4 (1.5)
Alanine aminotransferase increased	1 (0.5)	4 (2.2)	4 (1.6)	3 (1.1)
Metabolism and nutrition disorders				
Hypoglycemia ^c	17 (8.9)	18 (10.1)	50 (19.8)	49 (18.4)
Hypertriglyceridemia	6 (3.1)	2 (1.1)	9 (3.6)	5 (1.9)
Dyslipidemia	3 (1.6)	4 (2.2)	11 (4.3)	10 (3.7)
Musculoskeletal and connective tissue disorders				
Arthralgia	16 (8.4)	9 (5.0)	17 (6.7)	20 (7.5)
Back pain	15 (7.9)	16 (8.9)	16 (6.3)	17 (6.4)
Osteoarthritis	8 (4.2)	4 (2.2)	2 (0.8)	7 (2.6)
Myalgia	6 (3.1)	4 (2.2)	6 (2.4)	5 (1.9)
Pain in extremity	6 (3.1)	13 (7.3)	12 (4.7)	18 (6.7)
Exostosis	4 (2.1)	2 (1.1)	1 (0.4)	1 (0.4)
Musculoskeletal pain	4 (2.1)	9 (5.0)	4 (1.6)	9 (3.4)
Muscle spasms	3 (1.6)	4 (2.2)	3 (1.2)	4 (1.5)
Nervous system disorders				
Headache	17 (8.9)	20 (11.2)	25 (9.9)	19 (7.1)
Dizziness	8 (4.2)	9 (5.0)	3 (1.2)	11 (4.1)
Parasthesia	0	2 (1.1)	1 (0.4)	6 (2.2)
Psychiatric disorders				
Anxiety	8 (4.2)	5 (2.8)	5 (2.0)	4 (1.5)
Depression	6 (3.1)	4 (2.2)	6 (2.4)	2 (0.7)
Renal and urinary disorders				
Microalbuminuria	5 (2.6)	4 (2.2)	3 (1.2)	2 (0.7)
Nephrolithiasis	4 (2.1)	3 (1.7)	0	4 (1.5)
Dysuria	0	4 (2.2)	5 (2.0)	7 (2.6)
Respiratory, thoracic, and mediastinal disorders				

Body System/Organ Class Adverse Event	Number of Patients (%) Add-on to Metformin		Number of Patients (%) Add-on to Sulfonylurea	
	Saxagliptin 5 mg N=191	Placebo N=179	Saxagliptin 5 mg N=253	Placebo N= 267
Cough	7 (3.7)	9 (5.0)	14 (5.5)	16 (6.0)
Pharyngolaryngeal pain	5 (2.6)	3 (1.7)	3 (1.2)	4 (1.5)
Skin and subcutaneous tissue disorders				
Rash	6 (3.1)	5 (2.8)	1 (0.4)	2 (0.7)
Alopecia	4 (2.1)	0	0	1 (0.4)
Pruritus	3 (1.6)	1 (0.6)	2 (0.8)	6 (2.2)
Vascular disorders				
Hypertension	9 (4.7)	12 (6.7)	21 (8.3)	13 (4.9)

^a The mean duration of exposure to double-blind study medication, including exposure after the initiation of rescue medication, was 75 weeks (Standard Deviation = 34) for saxagliptin 5 mg plus metformin and 68 weeks (Standard Deviation= 35) for placebo plus metformin groups.

^b The mean duration of exposure to double-blind study medication, including exposure after the initiation of rescue medication, was 50 weeks (Standard Deviation = 17) for saxagliptin 5 mg plus glyburide and 48 weeks (Standard Deviation= 17) for placebo plus up-titrated glyburide groups.

^c “Hypoglycemia” includes events of Hypoglycemia and Blood Glucose Decreased.

Rash-related adverse events in the add-on to metformin study (24-week short-term and long-term extension) were reported in 4.2% and 2.8% of patients who received saxagliptin 5 mg and placebo, respectively. In the add-on to sulfonylurea study (24-week short-term and long-term extension) rash-related events were reported in 1.6 % and 1.1% of patients who received saxagliptin 5 mg and placebo, respectively.

In a pooled analysis of the 24-week placebo-controlled clinical trials, hypersensitivity-related events, such as urticaria and facial edema were reported in 1.5% and 0.4% of patients who received saxagliptin 5 mg and placebo, respectively. None of these events in patients who received saxagliptin required hospitalization or were reported as life-threatening by the investigators. One saxagliptin-treated patient in this pooled analysis discontinued due to generalized urticaria and facial edema.

In the short-term 24-week add-on to sulfonylurea study (glyburide 7.5 mg), the overall incidence of hypoglycemia was higher for saxagliptin 5 mg versus placebo (14.6% versus 10.1%). The incidence of confirmed hypoglycemic events, defined as symptoms of hypoglycemia accompanied by a fingerstick glucose value of ≤ 2.8 mmol/L, was similar for saxagliptin 5 mg treated group (0.8%) and placebo group (0.7%). In the long-term extension of the add-on to sulfonylurea study, the overall incidence of hypoglycemia was similar for saxagliptin 5 mg (19.8%) versus uptitrated sulfonylurea plus placebo (19.8% versus 18.4%).

The adverse event of hypertension was reported in more patients on saxagliptin 5 mg plus glyburide (8.3%) versus placebo plus glyburide (4.9%) in the add-on to sulfonylurea trial. Analysis of the mean systolic and diastolic blood pressure values did not reveal clinically meaningful changes.

Adverse reactions, reported regardless of causality assessment, in ≥ 2 % of patients treated with either saxagliptin 5 mg or placebo as an add-on to insulin (with or without metformin) are shown in Table 2.

Table 2 Adverse Reactions (Regardless of Investigator Assessment of Causality) in the Add-on to Insulin Study ^a (24-week Short Term Study and the Long Term Extension) Reported in $\geq 2\%$ of Patients Treated with Either Saxagliptin 5 mg or Placebo

Body System/Organ Class Adverse Event	Number of Patients (%) Add-on to Insulin (with or without Metformin)	
	Saxagliptin 5 mg + Insulin N=304	Placebo + Insulin N=151
Blood and lymphatic system disorders		
Anemia	6 (2.0)	4 (2.6)
Gastrointestinal disorders		
Diarrhea	14 (4.6)	7 (4.6)
Constipation	12 (3.9)	5 (3.3)
Abdominal pain	8 (2.6)	2 (1.3)
Gastritis	8 (2.6)	2 (1.3)
Nausea	5 (1.6)	5 (3.3)
General disorders and administration site conditions		
Edema peripheral	9 (3.0)	5 (3.3)
Infections and infestations		
Urinary tract infection	24 (7.9)	12 (7.9)
Nasopharyngitis	19 (6.3)	10 (6.6)
Upper respiratory tract infection	19 (6.3)	11 (7.3)
Bronchitis	16 (5.3)	5 (3.3)
Pharyngitis	11 (3.6)	8 (5.3)
Influenza	10 (3.3)	14 (9.3)
Cystitis	8 (2.6)	3 (2.0)
Gastroenteritis	7 (2.3)	2 (1.3)
Investigations		
Blood creatine phosphokinase increased	7 (2.3)	1 (0.7)
Metabolism and nutrition disorders		
Hypoglycemia ^b	69 (22.7)	40 (26.5)
Musculoskeletal and connective tissue disorders		
Arthralgia	13 (4.3)	5 (3.3)
Back pain	10 (3.3)	6 (4.0)
Osteoarthritis	7 (2.3)	0
Pain in extremity	7 (2.3)	10 (6.6)
Musculoskeletal pain	3 (1.0)	6 (4.0)
Nervous system disorders		
Headache	18 (5.9)	6 (4.0)
Dizziness	8 (2.6)	3 (2.0)
Respiratory, thoracic, and mediastinal disorders		
Cough	7 (2.3)	6 (4.0)
Vascular disorders		
Hypertension	9 (3.0)	8 (5.3)
Hypertensive crisis ^c	6 (2.0)	1 (0.7)

^a The mean duration of exposure to double-blind study medication, including exposure after changes in insulin medication, was 47 week (Standard Deviation = 13) for Saxagliptin 5 mg plus insulin and 47 weeks (Standard Deviation = 13) for placebo plus insulin groups.

^b "Hypoglycemia" includes events of Hypoglycemia and Blood Glucose Decreased.

^c Term as reported; cases do not meet medically accepted definition of hypertensive crisis.

In the short-term 24-week add-on to insulin study, the overall incidence of reported hypoglycemia was 18.4% for saxagliptin 5 mg and 19.9% for placebo. The incidence of confirmed hypoglycemic events, defined as symptoms of hypoglycemia accompanied by a fingerstick glucose value of ≤ 2.8 mmol/L, was 5.3% for the Saxagliptin 5 mg treated group versus 3.3% for the placebo group. In the long-term extension of the add-on to insulin study, the overall incidence of hypoglycemia was lower for Saxagliptin 5 mg (22.7%) versus placebo (26.5%) plus insulin with or without metformin.

Adverse reactions, reported regardless of causality assessment, in ≥ 2 % of patients treated with either saxagliptin 5 mg or placebo as an add-on to metformin and a sulfonylurea are shown in Table 3.

Table 3 Adverse Reactions (Regardless of Investigator Assessment of Causality) in the Add-on to Metformin and a Sulfonylurea (SU) Study^a (24-week) Reported in ≥ 2 % of Patients Treated with Either Saxagliptin 5 mg or Placebo

Body System/Organ Class Adverse Event	Number of Patients (%) Add-on to Metformin and SU	
	Saxagliptin 5 mg + Metformin + SU N=129	Placebo + Metformin + SU N=128
Blood and lymphatic system disorders		
Anemia	1 (0.8)	5 (3.9)
Gastrointestinal disorders		
Diarrhea	7 (5.4)	5 (3.9)
Flatulence	4 (3.1)	0
Gastritis	3 (2.3)	3 (2.3)
Nausea	2 (1.6)	4 (3.1)
Constipation	1 (0.8)	3 (2.3)
Infections and infestations		
Nasopharyngitis	8 (6.2)	12 (9.4)
Upper respiratory tract infection	6 (4.7)	6 (4.7)
Urinary tract infection	4 (3.1)	8 (6.3)
Pharyngitis	0	3 (2.3)
Oral candidiasis	0	3 (2.3)
Metabolism and nutrition disorders		
Hypoglycemia ^b	13 (10.1)	8 (6.3)
Dyslipidemia	5 (3.9)	7 (5.5)
Hyperglycemia	4 (3.1)	4 (3.1)
Musculoskeletal and connective tissue disorders		
Pain in extremity	2 (1.6)	4 (3.1)
Arthralgia	2 (1.6)	3 (2.3)
Back pain	1 (0.8)	4 (3.1)
Nervous system disorders		
Headache	4 (3.1)	3 (2.3)
Dizziness	3 (2.3)	2 (1.6)
Neuropathy peripheral	3 (2.3)	0
Psychiatric Disorders		
Insomnia	0	3 (2.3)
Respiratory, thoracic, and mediastinal disorders		
Cough	4 (3.1)	1 (0.8)

Skin and subcutaneous tissue disorders		
Rash	2 (1.6)	3 (2.3)
Vascular disorders		
Hypertension	7 (5.4)	2 (1.6)

^aThe mean duration of exposure to double-blind study medication was 159 days (Standard Deviation = 31) in the saxagliptin 5 mg group and 160 days (Standard Deviation = 30) for the placebo group.

^b“Hypoglycemia” includes events of Hypoglycemia and Blood Glucose Decrease.

Serious Adverse Reactions (reported in < 2% of patients) and Adverse Reactions of Interest* (reported in < 2% of patients and in at least 2 patients), Regardless of Investigator Assessment of Causality and Frequency > Placebo, in the Add-on to Metformin, Add-on to Sulfonylurea, Add-on to Insulin (with or without Metformin) Studies (24-week short-term and the long-term extensions) and in the Add-on to Metformin and a Sulfonylurea Study (24-week)

* System Organ Classes were considered to be of interest based on the adverse event profile of the DPP-4 inhibitor class of drugs, non-clinical data for saxagliptin, as well as the patient population.

Blood and lymphatic system disorders*: eosinophilia, lymphopenia, iron deficiency anemia, normochromic normocytic anemia,

Cardiac disorders*: coronary artery disease, left ventricular hypertrophy, atrioventricular block first degree, bundle branch block left, mitral valve incompetence, myocardial ischemia, palpitations, ventricular extrasystoles, acute myocardial infarction, atrioventricular block complete, cardiac failure congestive, cardiogenic shock

Gastrointestinal disorders: abdominal pain, diarrhea, salivary gland mass, vomiting

Hepatobiliary disorders: cholecystitis acute, cholecystitis, hepatitis

Immune system disorders*: hypersensitivity, sarcoidosis

Infections and infestations: cellulitis orbital, clostridium difficile colitis, urosepsis, diverticulitis, lower respiratory tract infection

Injury, poisoning and procedural complications: road traffic accident, ankle fracture, fall, gastrointestinal injury, incisional hernia, limb injury, skin laceration

Investigations*: aspartate aminotransferase increased, c-reactive protein increased, electrocardiogram repolarisation abnormality, blood cholesterol increased, blood pressure increased, electrocardiogram abnormal, lymphocyte count decreased

Metabolism and nutrition disorders: dehydration

Musculoskeletal and connective tissue disorders: arthralgia, osteoarthritis

Neoplasms benign, malignant and unspecified (including cysts and polyps): myelodysplastic syndrome, pancreatic cancer, laryngeal cancer

Nervous system disorders: altered state of consciousness, dizziness

Renal and urinary disorders: calculus ureteric, calculus urinary, renal impairment

Respiratory, thoracic and mediastinal disorders: hemoptysis, pulmonary embolism

Skin and subcutaneous tissue disorders*: rash, alopecia, dermatitis atopic, hyperhidrosis, rash papular, skin lesion, dermatitis contact, dermatitis, dry skin, seborrhoeic dermatitis, urticaria

Surgical and medical procedures: sterilisation

Abnormal Hematologic and Clinical Chemistry Findings

Absolute Lymphocyte Counts: A dose-related mean decrease in absolute lymphocyte count was observed with saxagliptin. From a baseline absolute lymphocyte count of approximately 2200 cells/mcL, a mean decrease of approximately 100 cells/mcL relative to placebo was observed in a pooled analysis of the placebo-controlled clinical studies. The proportion of patients who were reported to have a lymphocyte count \leq 750 cells/ mcL was 1.5% in the saxagliptin 5 mg group and 0.4% in the placebo group. The decreases in lymphocyte count were not associated with clinically relevant adverse reactions. When clinically indicated, such as in settings of unusual or prolonged infection, lymphocyte count should be measured. The effect of saxagliptin on lymphocyte counts in patients with lymphocyte abnormalities (e.g. human immunodeficiency virus) is unknown.

Platelets: Saxagliptin did not demonstrate a clinically meaningful or consistent effect on platelet count in the double-blind, controlled clinical safety and efficacy trials. In the add-on to insulin trial, there was a -2.6% decrease from baseline in platelet count in the saxagliptin group compared with a -0.1% decrease in the placebo group. An event of thrombocytopenia, consistent with a diagnosis of idiopathic thrombocytopenic purpura, was observed in the clinical program. The relationship of this event to saxagliptin is not known.

Urinary white and red blood cell counts: In the add-on to insulin trial, there was a higher percentage of saxagliptin patients, compared to placebo patients who presented with marked urinary red blood cell counts (15.1% saxagliptin versus 3.2% placebo) and urinary white blood cell counts (30.4% versus 18.9%). No consistent findings of urine laboratory abnormalities have been observed in the overall saxagliptin clinical program. No imbalances were observed for either URBC or UWBC in the pooled analysis of Phase 2/3 studies.

Post-Market Adverse Drug Reactions

Additional adverse reactions have been identified during post-marketing use of saxagliptin. Because these reactions are reported voluntarily from a population of uncertain size, it is generally not possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

Blood and lymphatic system disorders: idiopathic thrombocytopenic purpura

Gastrointestinal disorders: acute and chronic pancreatitis (see WARNINGS AND PRECAUTIONS)

Immune system disorders: Hypersensitivity reactions, including anaphylaxis, angioedema, rash, urticaria and exfoliative skin conditions, including Stevens-Johnson syndrome (see CONTRAINDICATIONS and WARNINGS AND PRECAUTIONS, Hypersensitivity Reactions)

Musculoskeletal and connective tissue disorders: severe and disabling arthralgia (see WARNINGS AND PRECAUTIONS, Musculoskeletal and connective tissue disorders)

Skin and subcutaneous tissue disorders: bullous pemphigoid.

Post-Marketing, Cardiovascular Safety

The Saxagliptin Assessment of Vascular Outcomes Recorded in Patients with Diabetes Mellitus-Thrombolysis in Myocardial Infarction (SAVOR) Trial was a CV outcome trial in 16,492 type 2 diabetic patients (median HbA1c = 7.6%) (12959 with established CV disease; 3533 with multiple risk factors only) who were randomized to saxagliptin (n=8280) or placebo (n=8212). The study population also included those ≥ 65 years (n=8561) and ≥ 75 years (n=2330), with normal or mild renal impairment (n=13,916) as well as moderate (n=2240) or severe (n=336) renal impairment. Subjects were followed for a mean duration of 2 years.

The primary endpoint was a composite endpoint consisting of the time-to-first occurrence of any of the following major adverse CV events (MACE): CV death, nonfatal myocardial infarction, or nonfatal ischemic stroke.

The trial established that the upper bound of the 2-sided 95% CI for the estimated risk ratio comparing the incidence of the primary composite endpoint observed with saxagliptin to that observed in the placebo group was < 1.3 . The study did not demonstrate the superiority of saxagliptin compared with placebo when added to current background therapy, in reducing the primary MACE endpoint (HR 1.00; 95% CI: 0.89, 1.12; $p = 0.986$).

Hospitalization for heart failure, occurred at a greater rate in the saxagliptin group (3.5%) compared with the placebo group (2.8%) [HR = 1.27; (95% CI 1.07, 1.51)]. Subjects on saxagliptin with a baseline history of congestive heart failure, especially those who also had renal impairment and/or MI, were at higher absolute risk for hospitalization for heart failure.

DRUG INTERACTIONS

The metabolism of saxagliptin is primarily mediated by P450 3A4/5 (CYP3A4/5).

In *in vitro* studies, saxagliptin and its major pharmacologically active metabolite neither inhibited nor induced CYP3A4. In addition, in *in vitro* studies, saxagliptin and its major pharmacologically active metabolite neither inhibited CYP1A2, 2A6, 2B6, 2C9, 2C19, 2D6, 2E1, nor induced CYP1A2, 2B6, 2C9. Therefore, saxagliptin is unlikely to alter the metabolic clearance of coadministered drugs that are metabolized by these enzymes. Saxagliptin is neither a significant inhibitor of P-glycoprotein (P-gp) nor an inducer of P-gp, and is unlikely to cause interactions with drugs that utilize these pathways.

The *in vitro* protein binding of saxagliptin and its major metabolite in human serum is below measurable levels. Thus, protein binding would not have a meaningful influence on the pharmacokinetics of saxagliptin or other drugs.

Drug-Drug Interactions

Effect of other drugs on saxagliptin

In studies conducted in healthy subjects as described below, the pharmacokinetics of saxagliptin and its major metabolite were not meaningfully altered by metformin, glyburide, pioglitazone, digoxin, simvastatin, diltiazem, ketoconazole, omeprazole, aluminum hydroxide + magnesium

hydroxide + simethicone combination, or famotidine. These drugs are considered unlikely to cause a clinically meaningful interaction with saxagliptin.

CYP3A4/5 Inducers: The coadministration of saxagliptin and CYP3A4/5 inducers, other than rifampin (such as carbamazepine, dexamethasone, phenobarbital and phenytoin) have not been studied and may result in decreased plasma concentration of saxagliptin and increased concentration of its major metabolite. Glycemic control should be carefully assessed when saxagliptin is used concomitantly with a potent CYP3A4 inducer.

Metformin: Coadministration of a single dose of saxagliptin (100 mg) and metformin (1000 mg), an OCT-1 and OCT-2 substrate, decreased the C_{max} of saxagliptin by 21%; however, the AUC was unchanged. Therefore, metformin is considered unlikely to cause a clinically meaningful interaction with saxagliptin. Meaningful interactions of saxagliptin with other OCT-1 and OCT-2 substrates would not be expected.

Glyburide: Coadministration of a single dose of saxagliptin (10 mg) and glyburide (5 mg), a CYP2C9 substrate, did not affect the pharmacokinetics of saxagliptin. Therefore, glyburide is considered unlikely to cause a clinically meaningful interaction with saxagliptin. Meaningful interactions of saxagliptin with other CYP2C9 substrates would not be expected.

Pioglitazone: Coadministration of multiple once-daily doses of saxagliptin (10 mg) and pioglitazone (45 mg), a CYP2C8 (major) and CYP3A4 (minor) substrate, did not alter the pharmacokinetics of saxagliptin. Therefore, pioglitazone is considered unlikely to cause a clinically meaningful interaction with saxagliptin. Meaningful interactions of saxagliptin with other CYP2C8 substrates would not be expected.

Digoxin: Coadministration of multiple once-daily doses of saxagliptin (10 mg) and digoxin (0.25 mg), a P-gp substrate, did not alter the pharmacokinetics of saxagliptin. Therefore, digoxin is considered unlikely to cause a clinically meaningful interaction with saxagliptin. Meaningful interactions of saxagliptin with other P-gp substrates would not be expected.

Simvastatin: Coadministration of multiple once-daily doses of saxagliptin (10 mg) and simvastatin (40 mg), a CYP3A4/5 substrate, increased the C_{max} of saxagliptin by 21%; however, the AUC of saxagliptin was unchanged. Therefore, simvastatin is considered unlikely to cause a clinically meaningful interaction with saxagliptin. Meaningful interactions of saxagliptin would not be expected with other substrates of CYP3A4/5.

Diltiazem: Coadministration of a single dose of saxagliptin (10 mg) and diltiazem (360 mg long-acting formulation at steady state), a moderate inhibitor of CYP3A4/5, increased the C_{max} and AUC for saxagliptin by 63% and 109%, respectively. This coadministration was also associated with 44% and 34% decreases in C_{max} and AUC(INF) values, respectively of its major metabolite. Therefore, diltiazem is considered unlikely to cause a clinically meaningful interaction with saxagliptin. Meaningful interactions of saxagliptin with other moderate CYP3A4/5 inhibitors would not be expected.

Ketoconazole: Coadministration of a single dose of saxagliptin (100 mg) and ketoconazole (200 mg every 12 hours at steady state), a potent inhibitor of CYP3A4/5 and P-gp, increased the C_{max} and AUC for saxagliptin by 62% and 145 % respectively. This coadministration was also associated with 95% and 88% decreases in C_{max} and AUC(INF) values, respectively of its major metabolite. Following coadministration of a single dose of saxagliptin at 20 times the recommended dose (100 mg) with ketoconazole, transient flu-like symptoms and a transient decrease in absolute lymphocyte count were observed. Additionally, transient decreases in absolute lymphocyte count were observed without any flu-like symptoms following coadministration of a single dose of saxagliptin at 4 times the recommended dose (20 mg) with ketoconazole.

Rifampin (Rifampicin): Coadministration of a single dose of saxagliptin (5 mg) with the potent CYP3A4/5 and P-gp inducer rifampin (600 mg once daily at steady state), decreased the C_{max} and AUC of saxagliptin by 53% and 76%, respectively. There was a corresponding increase in C_{max} (39%) but no significant change in plasma AUC of the active metabolite. There was no change in the maximum DPP4 inhibition (% I_{max}) and only a 6% decrease in the mean area under the effect time curve for DPP4 inhibition (AUEC) over a 24-hour period (the dosing interval for saxagliptin) when saxagliptin was coadministered with rifampin; however, a shorter DPP4 inhibition T-HALF was observed during the rifampin coadministration period (25.9 hours for saxagliptin-alone versus 14.5 hours for saxagliptin plus rifampin). See also WARNINGS AND PRECAUTIONS, Use with potent CYP3A4 inducers.

Omeprazole: Coadministration of multiple once-daily doses of saxagliptin (10 mg) and omeprazole (40 mg), a CYP2C19 (major) and CYP3A4 substrate, an inhibitor of CYP2C19, and an inducer of MRP-3, did not alter the pharmacokinetics of saxagliptin. Therefore, omeprazole is considered unlikely to cause a clinically meaningful interaction with saxagliptin. Meaningful interactions of saxagliptin with other CYP2C19 inhibitors or MRP-3 inducers would not be expected.

Aluminum hydroxide + magnesium hydroxide + simethicone: Coadministration of a single dose of saxagliptin (10 mg) and a liquid containing aluminum hydroxide (2400 mg), magnesium hydroxide (2400 mg), and simethicone (240 mg) decreased the C_{max} of saxagliptin by 26%; however, the AUC of saxagliptin was unchanged. Therefore, meaningful interactions of saxagliptin with antacid and antigas formulations of this type would not be expected.

Famotidine: Administration of a single dose of saxagliptin (10 mg) three hours after a single dose of famotidine (40 mg), an inhibitor of hOCT-1, hOCT-2, and hOCT-3, increased the C_{max} of saxagliptin by 14%; however, the AUC of saxagliptin was unchanged. Therefore, famotidine is considered unlikely to cause a clinically meaningful interaction with saxagliptin. Meaningful interactions of saxagliptin would not be expected with other inhibitors of hOCT-1, hOCT-2, and hOCT-3.

Effect of saxagliptin on other drugs

In studies conducted in healthy subjects, as described below, saxagliptin did not meaningfully alter the pharmacokinetics of metformin, glyburide, pioglitazone, digoxin, simvastatin, diltiazem, ketoconazole, or an estrogen/progestin combined oral contraceptive. Saxagliptin is considered unlikely to cause a clinically meaningful interaction with these drugs.

Metformin: Coadministration of a single dose of saxagliptin (100 mg) and metformin (1000 mg), an OCT-1 and OCT-2 substrate, did not alter the pharmacokinetics of metformin in healthy subjects. Therefore, saxagliptin is considered unlikely to cause a clinically meaningful interaction with metformin. Saxagliptin is not an inhibitor of OCT-1 and OCT-2-mediated transport.

Glyburide: Coadministration of a single dose of saxagliptin (10 mg) and glyburide (5 mg), a CYP2C9 substrate, increased the plasma C_{max} of glyburide by 16%; however, the AUC of glyburide was unchanged. Therefore, saxagliptin is considered unlikely to cause a clinically meaningful interaction with glyburide. Saxagliptin does not meaningfully inhibit CYP2C9-mediated metabolism.

Pioglitazone: Coadministration of multiple once-daily doses of saxagliptin (10 mg) and pioglitazone (45 mg), a CYP2C8 substrate, increased the plasma C_{max} of pioglitazone by 14%; however, the AUC of pioglitazone was unchanged. Therefore, saxagliptin is considered unlikely to cause a clinically meaningful interaction with pioglitazone. Saxagliptin does not meaningfully inhibit or induce CYP2C8-mediated metabolism.

Digoxin: Coadministration of multiple once-daily doses of saxagliptin (10 mg) and digoxin (0.25 mg), a P-gp substrate, did not alter the pharmacokinetics of digoxin. Therefore, saxagliptin is considered unlikely to cause a clinically meaningful interaction with digoxin. Saxagliptin is not an inhibitor or inducer of P-gp-mediated transport.

Simvastatin: Coadministration of multiple once-daily doses of saxagliptin (10 mg) and simvastatin (40 mg), a CYP3A4/5 substrate, did not alter the pharmacokinetics of simvastatin. Therefore, saxagliptin is considered unlikely to cause a clinically meaningful interaction with simvastatin. Saxagliptin is not an inhibitor or inducer of CYP3A4/5-mediated metabolism.

Diltiazem: Coadministration of multiple once-daily doses of saxagliptin (10 mg) and diltiazem (360 mg long-acting formulation at steady state), a moderate inhibitor of CYP3A4/5, increased the plasma C_{max} of diltiazem by 16%; however, the AUC of diltiazem was unchanged. Therefore, saxagliptin is considered unlikely to cause a clinically meaningful interaction with diltiazem.

Ketoconazole: Coadministration of a single dose of saxagliptin (100 mg) and multiple doses of ketoconazole (200 mg every 12 hours at steady state), a potent inhibitor of CYP3A4/5 and P-gp, decreased the geometric means for C_{max} and AUC(INF) of ketoconazole by 16% and by 13% respectively, relative to those observed following administration of 200 mg ketoconazole q 12 h alone.

Oral Contraceptives: Coadministration of multiple once-daily doses of saxagliptin (5 mg) and a monophasic combined oral contraceptive containing 0.035 mg ethinyl estradiol/0.250 mg norgestimate for 21 days, did not alter the steady state pharmacokinetics of the primary active estrogen component, ethinyl estradiol, or the primary active progestin component, norelgestromin. The plasma AUC of norgestrel, an active metabolite of norelgestromin, was increased by 13% and the plasma C_{max} of norgestrel was increased by 17%. This small magnitude change in AUC and C_{max} of norgestrel is not considered to be clinically meaningful. Based on these findings, saxagliptin would not be expected to meaningfully alter the pharmacokinetics of an estrogen/progestin

combined oral contraceptive.

Drug-Food Interactions

There are no known interactions with food. Grapefruit juice being a weak inhibitor of CYP3A4 gut wall metabolism may give rise to modest increases in plasma levels of saxagliptin.

Drug-Herb Interactions

Interactions with herbal products have not been established.

Drug-Laboratory Interactions

Interactions with laboratory tests have not been established.

Drug-Lifestyle Interactions

No studies of the effects of saxagliptin on the ability to drive and use machines have been performed. However, saxagliptin is not expected to affect the ability to drive and use machines.

DOSAGE AND ADMINISTRATION

Dosing Considerations

Sandoz Saxagliptin may be taken with or without food.

Recommended Dose and Dosage Adjustment

The recommended dose of Sandoz Saxagliptin is 5 mg once daily.

Renal Impairment: Assessment of renal function is recommended prior to initiation of Sandoz Saxagliptin and periodically thereafter (see WARNINGS AND PRECAUTIONS, ACTION AND CLINICAL PHARMACOLOGY and CLINICAL TRIALS – Patients with Renal Impairment).

Mild renal impairment

- No dosage adjustment for Sandoz Saxagliptin is recommended for patients with mild renal impairment (eGFR 60-89 mL/min/1.73m² (by Modified Diet in Renal Disease [MDRD] eGFR equation)).

Moderate renal impairment

- No dosage adjustment is required for patients with eGFR \geq 45 mL/min/1.73 m².
- For patients with moderate renal impairment with eGFR < 45 mL/min/1.73 m², the dose is 2.5 mg once daily.

Severe renal impairment

Saxagliptin should be used with caution in patients with severe renal impairment. Sandoz Saxagliptin is not recommended for patients with end-stage renal disease (ESRD) requiring hemodialysis. For patients with severe renal impairment (eGFR < 30 mL/min/1.73m²) the recommended dose is 2.5 mg once daily.

Hepatic Impairment: Use of Sandoz Saxagliptin in patients with moderate to severe hepatic impairment is not recommended due to lack of clinical experience with this patient population.

Pediatrics (<18 years of age): Safety and effectiveness of Sandoz Saxagliptin in pediatric patients have not been established. Therefore, Sandoz Saxagliptin should not be used in this patient population.

Geriatrics (≥65 years of age): No dosage adjustment for Sandoz Saxagliptin is required based solely on age (see WARNINGS AND PRECAUTIONS, CLINICAL TRIALS and ACTION AND CLINICAL PHARMACOLOGY).

Missed Dose

If a dose of Sandoz Saxagliptin is missed, it should be taken as soon as the patient remembers. A double dose of Sandoz Saxagliptin should not be taken on the same day.

Administration

Sandoz Saxagliptin tablets must not be split or cut.

OVERDOSAGE

For management of a suspected drug overdose, contact your regional Poison Control Centre.

In the event of an overdose, appropriate supportive treatment should be initiated as dictated by the patient's clinical status. Saxagliptin and its major metabolite are removed by hemodialysis (23% of dose over 4 hours).

ACTION AND CLINICAL PHARMACOLOGY

Mechanism of Action

Saxagliptin is a potent, selective, reversible, competitive, DPP-4 inhibitor. Saxagliptin demonstrates selectivity for DPP-4 versus other DPP enzymes, including DPP-8 and DPP-9. Saxagliptin has extended binding to the DPP-4 active site, prolonging its inhibition of DPP-4. Saxagliptin exerts its actions in patients with type 2 diabetes by slowing the inactivation of incretin hormones, including glucagon-like peptide-1 (GLP-1). The concentration of active (intact) GLP-1 incretin hormone is increased.

Incretin hormones are released by the intestine throughout the day and concentrations are increased in response to a meal. These hormones are rapidly inactivated by the enzyme DPP-4. The incretins are part of an endogenous system involved in the physiologic regulation of glucose homeostasis. When blood glucose concentrations are elevated, GLP-1 and GIP increase insulin synthesis and release from pancreatic beta cells. GLP-1 also lowers glucagon secretion from pancreatic alpha cells, leading to reduced hepatic glucose production.

The concentration of GLP-1 is reduced in patients with type 2 diabetes⁹, but saxagliptin increases active GLP-1 concentration. By increasing active GLP-1 concentration, saxagliptin increases postprandial insulin release and decreases postprandial glucagon concentrations in the circulation in a glucose-dependent manner. In patients with type 2 diabetes with hyperglycemia, these changes in insulin and glucagon levels may lead to lower hemoglobin A1C (HbA1c) and lower fasting and

postprandial glucose concentrations.⁵

Pharmacodynamics

In patients with type 2 diabetes, administration of saxagliptin led to dose-dependent inhibition of DPP-4 enzyme activity for a 24-hour period. After an oral glucose load or a meal, this DPP-4 inhibition resulted in a 2- to 3-fold increase in circulating levels of active GLP-1, decreased postprandial glucagon concentrations, and increased glucose-dependent beta cell responsiveness with higher postprandial insulin and C-peptide concentrations. The rise in insulin and decrease in glucagon were associated with lower fasting glucose concentrations and reduced glucose excursion following an oral glucose load or a meal.

Cardiac electrophysiology: In a randomized, double-blind, placebo-controlled, 4-way crossover, active comparator study, 40 healthy subjects were administered saxagliptin 40 mg (8 times the RHD), saxagliptin 10 mg (2 times the RHD), or placebo once daily for 4 days, or a single dose of moxifloxacin 400 mg as a positive control. The saxagliptin 10 mg and 40 mg treatments were not associated with any prolongation of the QTc, QRS, or PR intervals. In the saxagliptin 10 mg treatment a significant increase in heart rate was observed at 0.5, 1, 1.5, 4, and 12 h post-dosing, with a maximum placebo- and baseline-corrected mean increase of 3.75 (90% 1.55, 5.95) beats per minute at 0.5 post-dosing when the baseline-corrected change in the placebo treatment at this time was -1.4 (90% CI -3.0, 0.1) beats per minute. Significant increases in heart rate were also observed in the saxagliptin 40 mg treatment at 0.5, 4, and 12 hours post-dosing, with a maximum placebo- and baseline-corrected mean increase of 4.5 (90% CI 2.23, 6.82) beats per minute at 4 hours post-dose dose when the baseline-corrected change in the placebo treatment at this time was -3.3 (90% CI -5.0, -1.6) beats per minute. The effect of the recommended 5 mg dose was not investigated in this study.

Pharmacokinetics

The pharmacokinetics of saxagliptin has been extensively characterized in healthy subjects and patients with type 2 diabetes (Table 4).

Table 4 Summary of saxagliptin's pharmacokinetic parameters in healthy subjects

	C_{max} (ng/mL)	$t_{1/2}$ (h)	AUC ng.h/mL	Renal Clearance (mL/min)
Single oral dose (5 mg) mean	24	2.5	78	230

Saxagliptin was rapidly absorbed after oral administration, with maximum saxagliptin plasma concentrations (C_{max}) usually attained within two hours after administration in the fasted state. The C_{max} and AUC values increased proportionally to the increment in the saxagliptin dose. Following a 5 mg single oral dose of saxagliptin to healthy subjects, the mean plasma AUC(INF) values for saxagliptin and its major metabolite were 78 ng·h/mL and 214 ng·h/mL, respectively. The corresponding plasma C_{max} values were 24 ng/mL and 47 ng/mL, respectively. The intra-subject coefficients of variation for saxagliptin C_{max} and AUC were less than 12%.

Following a single oral dose of 5 mg saxagliptin to healthy subjects, the mean plasma terminal half-life ($t_{1/2}$) for saxagliptin was 2.5 hours, and the mean $t_{1/2}$ value for plasma DPP-4 inhibition was

26.9 hours. The inhibition of plasma DPP-4 activity by saxagliptin occurs for at least 24-hours after oral administration of saxagliptin. No appreciable accumulation was observed with repeated once-daily dosing at any dose level. No dose- and time-dependence was observed in the clearance of saxagliptin and its major metabolite over 14 days of once-daily dosing with saxagliptin at doses ranging from 2.5 mg to 400 mg. Results from population-based exposure modeling suggest that the pharmacokinetics of saxagliptin and its major metabolite were similar in healthy subjects and in patients with type 2 diabetes.

Absorption: Saxagliptin may be administered with or without food. The amount of saxagliptin absorbed following an oral dose is at least 75%. Food had relatively modest effects on the pharmacokinetics of saxagliptin in healthy subjects. Administration with a high-fat meal resulted in no change in saxagliptin C_{max} and a 27% increase in AUC compared with the fasted state. The time for saxagliptin to reach C_{max} (T_{max}) was increased by approximately 0.5 hours with food compared with the fasted state. These changes were not considered to be clinically meaningful.

Distribution: The *in vitro* protein binding of saxagliptin and its major metabolite in human serum is below measurable levels. Thus, changes in blood protein levels in various disease states (e.g., renal or hepatic impairment) are not expected to alter the disposition of saxagliptin.

Metabolism: The metabolism of saxagliptin is primarily mediated by cytochrome P450 3A4/5 (CYP3A4/5). The major metabolite of saxagliptin is also a selective, reversible, competitive DPP-4 inhibitor, half as potent as saxagliptin.

Excretion: Saxagliptin is eliminated by both renal and hepatic pathways. Following a single 50 mg dose of ¹⁴C-saxagliptin, 24%, 36%, and 75% of the dose was excreted in the urine as saxagliptin, its major metabolite, and total radioactivity, respectively. The average renal clearance of saxagliptin (~230 mL/min) was greater than the average estimated glomerular filtration rate (~120 mL/min), suggesting some active renal excretion. For the major metabolite, renal clearance values were comparable to estimated glomerular filtration rate. A total of 22% of the administered radioactivity was recovered in feces representing the fraction of the saxagliptin dose excreted in bile and/or unabsorbed drug from the gastrointestinal tract.

Pharmacokinetics of the Major Metabolite: The C_{max} and AUC values for the major metabolite of saxagliptin increased proportionally to the increment in the saxagliptin dose. Following single oral doses of 2.5 mg to 400 mg saxagliptin in the fed or fasted states, the mean AUC values for the major metabolite ranged from 2- and 7 times higher than the parent saxagliptin exposures on a molar basis. Following a single oral dose of 5 mg saxagliptin in the fasted state, the mean terminal half-life ($t_{1/2}$) value for the major metabolite was 3.1 hours and no appreciable accumulation was observed upon repeated once-daily dosing at any dose.

Special Populations and Conditions

Pediatrics (< 18 years of age): Pharmacokinetics in the pediatric population have not been studied. Therefore, Sandoz Saxagliptin should not be used in this patient population.

Geriatrics (≥ 65 years of age): No dosage adjustment is necessary based on age alone. Elderly subjects (65-80 years) had 23% and 59% higher geometric mean C_{max} and geometric mean AUC values, respectively, for parent saxagliptin than young subjects (18-40 years). Differences in major metabolite pharmacokinetics between elderly and young subjects generally reflected the differences observed in parent saxagliptin pharmacokinetics. The difference between the pharmacokinetics of saxagliptin and the major metabolite in young and elderly subjects is likely to be due to multiple factors including declining renal function and metabolic capacity with increasing age.

Gender: No dosage adjustment is necessary based on gender. There were no differences observed in saxagliptin pharmacokinetics between males and females. Compared to males, females had approximately 25% higher exposure values for the major metabolite than males, but the clinical relevance of this difference is unknown.

Race: No dosage adjustment is necessary based on race. An exposure modeling analysis compared the pharmacokinetics of saxagliptin and its major metabolite in 309 white subjects with 105 non-white subjects (consisting of 6 race groups). No significant difference in the pharmacokinetics of saxagliptin and its major metabolite were detected between these two populations.

Body Mass Index: No dosage adjustment is recommended based on body mass index (BMI).

Renal Impairment: A single-dose, open-label study was conducted to evaluate the pharmacokinetics of saxagliptin (10 mg dose) in subjects with varying degrees of chronic renal impairment compared to subjects with normal renal function.

In subjects with approximately $eGFR \geq 45$ mL/min/1.73 m² by MDRD $eGFR$ equation, the increases of the AUC values of saxagliptin and its major metabolite were not clinically relevant. Dosage adjustment in these patients is not recommended.

In subjects with renal impairment with approximately $eGFR < 45$ mL/min/1.73 m², the AUC values of saxagliptin and its major metabolite were more than 2.1- and 4.5-fold higher, respectively, than AUC values in subjects with normal renal function. The dose of Sandoz Saxagliptin should be reduced to 2.5 mg once daily in patients with renal impairment with approximately $eGFR < 45$ mL/min/1.73 m² (see WARNINGS AND PRECAUTIONS, DOSAGE AND ADMINISTRATION and CLINICAL TRIALS – Patients with Renal Impairment).

Hepatic Impairment: In subjects with hepatic impairment (Child-Pugh classes A, B, and C), mean C_{max} and AUC of saxagliptin were up to 8% and 77% higher, respectively, compared to healthy matched controls following administration of a single 10 mg dose of saxagliptin. The corresponding C_{max} and AUC of the major metabolite were up to 59% and 33% lower, respectively, compared to healthy matched controls. Use in moderate to severe hepatic impairment is not recommended.

STORAGE AND STABILITY

Store at room temperature (15-30°C). Store in the original packaging. Protect from moisture.

Keep in a safe place out of reach of children and pets. Unused medication should not be disposed of down the drain or in household garbage.

DOSAGE FORMS, COMPOSITION AND PACKAGING

Sandoz Saxagliptin 2.5 mg tablets are light yellow, biconvex, round, film-coated tablets with “SG” embossed on one side and “2.5” on the other side.

Sandoz Saxagliptin 5 mg tablets are light pink, biconvex, round, film-coated tablets with “SG” embossed on one side and “5.0” on the other side.

The 2.5 mg tablets are provided in blisters in cartons of 30 (3 blister strips of 10 tablets each) and bottles of 30.

The 5 mg tablets are provided in blisters in cartons of 30 (3 blister strips of 10 tablets each) and bottles of 30.

Composition

Each 2.5 mg tablet contains 2.5 mg of saxagliptin (as saxagliptin hydrochloride dihydrate).

Each 5 mg tablet contains 5 mg of saxagliptin (as saxagliptin hydrochloride dihydrate).

Sandoz Saxagliptin tablets contain the following non-medicinal ingredients: lactose monohydrate, microcrystalline cellulose, croscarmellose sodium, magnesium stearate and (dilute) hydrochloric acid.

In addition, the film coating contains the following inactive ingredients: hypromellose, macrogol, lactose anhydrous, titanium dioxide and yellow iron oxide (2.5 mg) or red iron oxide (5 mg).

PART II: SCIENTIFIC INFORMATION

PHARMACEUTICAL INFORMATION

Drug Substance

Common name: Saxagliptin Hydrochloride Dihydrate


Chemical name: (1S, 3S, 5S)-2-[(2S)-amino(3-hydroxytricyclo [3.3.1.1^{3,7}] dec-1-yl) acetyl]-2-azabicyclo [3.1.0] hexane-3-carbonitrile, hydrochloride, dihydrate.

Or

(1S, 3S, 5S)-2-[(2S)-2-amino-2-(3-hydroxyadamantan-1-yl) acetyl]-2-azabicyclo [3.1.0] hexane-3-carbonitrile hydrochloride dihydrate.

Molecular formula and molecular mass: $C_{18}H_{25}N_3O_2 \cdot HCl \cdot 2H_2O$; 387.90 g/mol

Structural formula:



Physicochemical properties: Saxagliptin Hydrochloride is a white to off white crystalline powder. The substance is freely soluble in water, soluble in methanol and ethanol; and insoluble in ethyl acetate.

CLINICAL TRIALS

Study demographics and trial design

Table 5 - Summary of patient demographics for clinical trials in specific indication

Trial design	Dosage, route of administration and duration	Study subjects per treatment arm Subjects ≥ 65 years of age Subjects ≥ 75 years of age (n = number)	Mean age (Range)	Gender (% M/F)
Add-on Combination Therapy with Metformin⁴				
Multicentre, randomized, double-blind, placebo-controlled	open-label metformin (1500 – 2500 mg) plus saxagliptin 5 mg or placebo Oral, 24 weeks	Saxagliptin 5 mg N=191 ≥ 65 years N=32 ≥ 75 years N=2 Placebo N= 179 ≥ 65 years n=26 ≥ 75 years n=3	55 years (26 - 76)	54/46
Add-on Combination Therapy with a Sulfonylurea³				
Multicentre, randomized, double-blind, placebo-controlled	open-label glyburide 7.5 mg plus saxagliptin 5 mg or open-label glyburide 7.5 mg plus double-blind glyburide 2.5 mg (total daily dose of 10 mg titratable to 15 mg) plus placebo Oral, 24 weeks	Saxagliptin 5 mg N=253 ≥ 65 years N=42 ≥ 75 years N=3 Placebo N=267 ≥ 65 years n=52 ≥ 75 years n=5	55 years (18 - 77)	45/55
Add-on Combination Therapy with Insulin (with or without Metformin)				
Multicentre, randomized, double-blind, placebo-controlled	open-label insulin (≥ 30 units/day, ≤ 150 units/day) alone or with metformin plus saxagliptin 5 mg or open label insulin (≥ 30 units/day, ≤ 150 units/day) alone or with metformin plus placebo Oral, 24 weeks	Saxagliptin 5 mg N=304 ≥ 65 years N=71 ≥ 75 years N=6 Placebo N=151 ≥ 65 years n=33 ≥ 75 years n=3	57 years (18 – 77)	41/59
Add-on Combination Therapy with Metformin and a Sulfonylurea				

Trial design	Dosage, route of administration and duration	Study subjects per treatment arm Subjects ≥ 65 years of age Subjects ≥ 75 years of age (n = number)	Mean age (Range)	Gender (% M/F)
Multicentre, randomized, double-blind, placebo-controlled	open-label metformin (≥ 1500 mg) and a sulfonylurea (≥ 50% of the maximum dose) plus saxagliptin 5 mg or open-label metformin (≥ 1500 mg) and a sulfonylurea (≥ 50% of the maximum dose) plus placebo Oral, 24 weeks	Saxagliptin 5 mg N=129 ≥ 65 years N=28 ≥ 75 years N=2 Placebo N=128 ≥ 65 years n=33 ≥ 75 years n=7	57 years (25 – 83)	60/40

Study results

In patients with type 2 diabetes, the treatment with saxagliptin 5 mg produced clinically relevant and statistically significant improvements in hemoglobin A1c (A1C), fasting plasma glucose (FPG), and postprandial glucose (PPG), including 2-hour PPG following standard oral glucose tolerance test (OGTT), compared to control.

Add-On Combination Therapy with Metformin

A total of 743 patients with type 2 diabetes participated in this randomized, double-blind, placebo-controlled study of 24-week duration, to evaluate the efficacy and safety of saxagliptin in combination with metformin in patients with inadequate glycemic control (A1C ≥ 7% and ≤ 10%) on metformin alone. Patients were required to be on a stable dose of metformin (1500 mg to 2550 mg daily) for at least 8 weeks to be enrolled in this study.

Patients who met eligibility criteria were enrolled in a single-blind, two-week, dietary and exercise placebo lead-in period during which patients received metformin at their pre-study dose, up to 2500 mg daily for the duration of the study. Following the lead-in period, eligible patients were randomized to 2.5 mg, 5 mg, or 10 mg of saxagliptin or placebo in addition to their current dose of open-label metformin. Patients who failed to meet specific glycemic goals during the study were treated with pioglitazone rescue therapy, added on to placebo or saxagliptin plus metformin. Dose titrations of saxagliptin and metformin were not allowed in this study.

In combination with metformin, saxagliptin 5 mg provided significant improvements in A1C, FPG, and PPG compared with the placebo plus metformin group (Table 6).

Table 6 Glycemic Parameters at Week 24 in a Placebo-Controlled Study of Saxagliptin in Combination with Metformin[§]

Efficacy Parameter	Saxagliptin 5 mg + Metformin	Placebo + Metformin
A1C (%)	N=186	N=175
Baseline (mean)	8.1	8.1
Change from baseline (adjusted mean ±)	-0.7	0.1
Difference from placebo (adjusted mean ±)	-0.8 ^a	-
95% Confidence Interval	(-1.0, -0.6)	
Percent of patients achieving A1C <7%	44% ^a (81/186)	17% (29/175)
FPG (mmol/L)	N=187	N=176
Baseline (mean)	9.9	9.7
Change from baseline (adjusted mean ±)	-1.2	0.07
Difference from placebo (adjusted mean ±)	-1.3 ^a	
95% Confidence Interval	(-1.7, -0.9)	
2-hour PPG (mmol/L)	N=155	N=135
Baseline (mean)	16.4	16.4
Change from baseline (adjusted mean ±)	-3.2	-1.0
Difference from placebo (adjusted mean ±)	-2.2 ^a	
95% Confidence Interval	(-3.1, -1.3)	
3-hour PPG AUC (mmol*min/L)	N=146	N=131
Baseline (mean)	2721	2631
Change from baseline (adjusted mean ±)	-532	-183
Difference from placebo (adjusted mean ±)	-349 ^a	
95% Confidence Interval	(-478, -221)	


§ Intent-to-treat population using last observation on study prior to pioglitazone rescue therapy.

± Least squares mean adjusted for baseline value.

a p-value <0.0001 compared to placebo

The mean percent change from baseline in A1C over the 24-week period is shown in Figure 1. The proportion of patients achieving A1C <7% (regardless of baseline value) was significantly greater in the saxagliptin 5 mg plus metformin treatment (43.5%) groups compared with the placebo plus metformin group (16.6%). Significant reductions in 2-hour PPG level following standard oral glucose tolerance test were observed in the saxagliptin 5 mg plus metformin treatment group (-3.2 mmol/L) compared with -1.0 mmol/L in the placebo plus metformin group. The proportion of patients who discontinued for lack of glycemic control or who were rescued for meeting prespecified glycemic criteria was higher in the placebo plus metformin group (27%) than in the saxagliptin 5 mg plus metformin group (13%). Higher baseline A1C was associated with a greater adjusted mean change from baseline in A1C with saxagliptin 5 mg. The effect of saxagliptin on lipid endpoints in this study was similar to placebo. Similar changes in body weight were observed in patients who received saxagliptin and placebo therapy (-0.9 kg and -0.9 kg, respectively).

Figure 1 Mean Change from Baseline in A1C in a Placebo-Controlled Study on Saxagliptin in Combination with Metformin *



*Intent-to-treat population using last observation on study prior to pioglitazone rescue therapy. Mean change from baseline (LOCF).

Controlled Long-Term Study Extension

Patients who completed all visits during the initial 24-week study period without need for hyperglycemia rescue therapy were eligible to enter a controlled double blind long-term study extension. Of the patients that started the 24-week treatment, 162 (84.8%) and 149 (83.2%) patients were taking saxagliptin 5 mg plus metformin and placebo plus metformin respectively. Patients who received saxagliptin in the initial 24-week study period maintained the same dose of saxagliptin in the long-term extension. Treatment with saxagliptin 5 mg plus metformin was associated with a greater reduction in A1C than in the placebo plus metformin group, and the effect relative to placebo was sustained at Week 50 and Week 102 compared to placebo. The A1C change for saxagliptin 5 mg plus metformin (n=100 observed, n=187 LOCF) compared with placebo plus metformin (n=59 observed, n=175 LOCF) was -0.7% at Week 50. The A1C change for saxagliptin 5 mg plus metformin (n=31 observed, n=184 LOCF) compared with placebo plus metformin (n=15 observed, n=172 LOCF) was -0.7% at Week 102.

Add-On Combination Therapy with a Sulfonylurea

A total of 768 patients with type 2 diabetes participated in this randomized, double-blind, placebo-controlled study of 24-weeks duration, to evaluate the efficacy and safety of saxagliptin in combination with sulfonylurea (SU) in patients with inadequate glycemic control at enrollment (A1C \geq 7.5% to \leq 10%) on a submaximal dose of SU alone. Patients were required to be on a submaximal dose of SU for 2 months or greater to be enrolled in this study. In this study,

saxagliptin in combination with a fixed, intermediate dose of SU was compared to titration to a higher dose of SU.

Patients who met eligibility criteria were enrolled in a single-blind, 4-week, dietary and exercise lead-in period and placed on glyburide 7.5 mg once daily. Following the lead-in period, eligible patients with A1C $\geq 7\%$ to $\leq 10\%$ were randomized to either 2.5 mg or 5 mg of saxagliptin plus 7.5 mg glyburide, or placebo plus a 10 mg total daily dose of glyburide. Patients who received placebo were eligible to have glyburide up-titrated to a total daily dose of 15 mg. Up titration of glyburide was not allowed in patients who received saxagliptin 2.5 or 5 mg. Glyburide could be down-titrated in any treatment group once during the 24-week study period due to hypoglycemia as deemed necessary by the investigator. Approximately 92% of patients in the placebo plus glyburide group were up-titrated to a final total daily dose of 15 mg during the study period. Patients who failed to meet specific glycemic goals during the study were treated with metformin rescue, added on to the saxagliptin plus glyburide or the placebo plus up-titrated glyburide group. Dose titration of saxagliptin was not permitted during the study.

In combination with glyburide, saxagliptin 5 mg provided significant improvements in A1C, FPG, and PPG compared with the placebo plus up-titrated glyburide group (Table 7). The mean percent change from baseline in A1C over the 24-week period is shown in Figure 2. The proportion of patients achieving A1C $<7.0\%$ (regardless of baseline value) was significantly greater in the saxagliptin 5 mg plus glyburide treatment group (22.8%) compared with the placebo plus up-titrated glyburide group (9.1%). Significant reductions in 2 hour PPG level following standard oral glucose tolerance test were observed in the saxagliptin 5 mg plus glyburide treatment group (-1.9 mmol/L) compared with 0.4 mmol/L in the placebo plus up-titrated glyburide group. The proportion of patients who discontinued for lack of glycemic control or who were rescued for meeting prespecified glycemic criteria was higher in the placebo plus up-titrated glyburide group (30%) than in the saxagliptin 5 mg plus glyburide group (17%). Higher baseline A1C was associated with a greater adjusted mean change from baseline in A1C with saxagliptin 5 mg. The effect of saxagliptin on lipid endpoints in this study was similar to placebo. Small increases in body weight were seen in patients treated with saxagliptin 5 mg plus glyburide and with placebo plus up-titrated glyburide (0.8 kg versus 0.3 kg, $p=0.012$).

Table 7 Glycemic Parameters at Week 24 in a Placebo-Controlled Study of Saxagliptin in Combination with Glyburide[§]

Efficacy Parameter	Saxagliptin 5 mg + Glyburide 7.5 mg	Placebo + Up-Titrated Glyburide
A1C (%)	N=250	N=264
Baseline (mean)	8.5	8.4
Change from baseline (adjusted mean ±)	-0.6	0.1
Difference from up-titrated glyburide (adjusted mean ±)	-0.7 ^a	-
95% Confidence Interval	(-0.9, -0.6)	
Percent of patients achieving A1C <7%	23% ^a (57/250)	9% (24/264)
FPG (mmol/L)	N=252	N=265
Baseline (mean)	9.7	9.7
Change from baseline (adjusted mean ±)	-0.5	0.04
Difference from up-titrated glyburide (adjusted mean ±)	-0.6 ^b	
95% Confidence Interval	(-0.9, -0.2)	
2-hour PPG (mmol/L)	N=202	N=206
Baseline (mean)	17.5	17.9
Change from baseline (adjusted mean ±)	-1.9	0.4
Difference from placebo (adjusted mean ±)	-2.3 ^a	
95% Confidence Interval	(-2.9, -1.7)	
3-hour PPG AUC (mmol*min/L)	N=195	N=204
Baseline (mean)	2794	2875
Change from baseline (adjusted mean ±)	-278	66
Difference from up-titrated glyburide (adjusted mean ±)	-344 ^a	
95% Confidence Interval	(-433, -254)	


§ Intent-to-treat population using last observation on study prior to metformin rescue therapy.

± Least squares mean adjusted for baseline value.

a p-value <0.0001 compared to placebo + up-titrated glyburide

b p-value=0.0020 compared to placebo + up-titrated glyburide

Figure 2 Mean Change from Baseline in A1C in a Placebo-Controlled Study on Saxagliptin in Combination with Glyburide *



* Intent-to-treat population using last observation on study prior to metformin rescue therapy. Mean change from baseline (LOCF).

Controlled Long-Term Study Extension

Patients who were rescued (based on predefined glucose levels) during the initial 24-week study period as well as those who completed all visits during the initial 24-week study period without the need for rescue therapy were eligible to enter a controlled double blind long-term study extension. Of the patients that started the 24-week treatment, 227 (89.7%) and 235 (88%) patients were taking saxagliptin 5 mg plus glyburide and placebo plus up-titrated glyburide respectively. Patients who received saxagliptin in the initial 24-week study period maintained the same dose of saxagliptin in the long-term extension. The A1C change for saxagliptin 5 mg plus glyburide (n=99 observed, n=243 LOCF) compared with placebo plus up-titrated glyburide (n=61 observed, n=253 LOCF) was -0.6% at Week 50.

Add-On Combination Therapy with Insulin (with or without Metformin)

A total of 455 patients with type 2 diabetes participated in this randomized, double-blind, placebo-controlled trial of 24-week duration to evaluate the efficacy and safety of saxagliptin in combination with insulin in patients with inadequate glycemic control (A1C $\geq 7.5\%$ and $\leq 11\%$) on insulin alone (N=141) or on insulin in combination with a stable dose of metformin (N=314). Patients were required to be on a stable dose of insulin (≥ 30 units to ≤ 150 units daily) with $\leq 20\%$ variation in total daily dose for ≥ 8 weeks prior to screening with or without metformin. Patients were on intermediate- or long-acting (basal) insulin or premixed insulin.

Patients using short-acting insulins were excluded unless the short-acting insulin was administered as part of a premixed insulin.

Patients who met eligibility criteria were enrolled in a single-blind, four-week, dietary and exercise placebo lead-in period during which patients received insulin (and metformin, if applicable) at their prestudy dose(s). Following the lead-in period, eligible patients were randomized to saxagliptin 5 mg or placebo in addition to continuing their current dose of insulin (and metformin, if applicable). Patients maintained a stable dose of insulin when possible. Patients who failed to meet specific glycemic goals or who increased their insulin dose by >20% were rescued and subsequently switched to a flexible insulin dose regimen. Dose titrations of saxagliptin and metformin (if applicable) were not allowed in this study.

Saxagliptin 5 mg add-on to insulin with or without metformin provided significant improvements in A1C and PPG compared with placebo add-on to insulin with or without metformin (Table 8). Similar A1C reductions versus placebo were achieved for patients using saxagliptin 5 mg add-on to insulin alone and saxagliptin 5 mg add-on to insulin in combination with metformin (−0.4% and −0.4%, respectively). The proportion of patients who discontinued for lack of glycemic control or who were rescued was 23% in the saxagliptin 5 mg add-on to insulin group and 32% in the placebo add-on to insulin group. The mean daily insulin dose at baseline was 53 units in patients treated with saxagliptin 5 mg and 55 units in patients treated with placebo. The mean change from baseline in daily dose of insulin was an increase of 2 units for the saxagliptin 5 mg group and 5 units for the placebo group.

Table 8 Glycemic Parameters at Week 24 in a Placebo-Controlled Trial of Saxagliptin as Add-On Combination Therapy with Insulin*

Efficacy Parameter	Saxagliptin 5 mg + Insulin (+/- Metformin) N=304	Placebo + Insulin (+/- Metformin) N=151
A1C (%)	N=300	N=149
Baseline (mean)	8.7	8.7
Change from baseline (adjusted mean)	-0.7	-0.3
Difference from placebo (adjusted mean)	-0.4 [‡]	-
95% Confidence Interval	(-0.6, -0.2)	
Percent of patients achieving A1C <7%	17% [§] (52/300)	7% (10/149)
2-hour PPG (mmol/L)	N=262	N=129
Baseline (mean)	13.9	14.2
Change from baseline (adjusted mean ±)	-1.5	-0.2
Difference from placebo (adjusted mean ±)	-1.3 [¶]	
95% Confidence Interval	(-2.1, -0.5)	
FPG (mmol/L)	N=300	N=149
Baseline (mean)	9.6	9.6
Change from baseline (adjusted mean ±)	-0.6	-0.3
Difference from placebo (adjusted mean ±)	-0.2 [#]	
95% Confidence Interval	(-0.7, 0.3)	
Mean Total Daily Dose of Insulin (unit)	N=299	N=151
Baseline (mean)	53	55
Change from baseline (adjusted mean ±)	2	5
Difference from placebo (adjusted mean ±)	-3 [§]	
95% Confidence Interval	(-6, -1)	

* Intent-to-treat population using last observation on study or last observation prior to insulin rescue therapy for patients needing rescue. Mean Total Daily Dose of Insulin: Intent-to-treat population using last observation on study.

† Least squares mean adjusted for baseline value and metformin use at baseline.

‡ p-value <0.0001 compared to placebo + insulin

§ Significance not tested

¶ p-value = 0.0016 compared to placebo + insulin

Not statistically significant

Controlled Long-Term Study Extension

Following completion of the 24-week short-term treatment period, patients were eligible to enter a controlled double blind long-term treatment period. Patients continued to take the same blinded study medication that they were assigned during the short-term treatment period (saxagliptin 5 mg or placebo added on to insulin with or without metformin). During the long-term treatment extension, changes in both the dose and type of insulin were allowed. Of the patients that continued into the long-term treatment period, 268 (88.2% of randomized) patients and 134 (88.7% of randomized) patients were taking saxagliptin 5 mg and placebo plus insulin with or without metformin, respectively. Results from the extension period demonstrated that reductions from baseline A1C seen in the saxagliptin 5 mg add-on to insulin group compared with the placebo add-

on to insulin group were sustained to Week 52; the A1C change for saxagliptin 5 mg (n=244 observed) compared with placebo (n=124 observed) was -0.4% at Week 52. Results were similar for subjects using metformin and not using metformin at baseline. Increases from baseline in mean total daily dose of insulin (MTDDI) were seen in both treatment groups through Week 52, with a numerically smaller increase in the saxagliptin 5 mg group (5 units saxagliptin versus 6 units Placebo).

Add-On Combination Therapy with Metformin and a Sulfonylurea

A total of 257 patients with type 2 diabetes participated in this 24-week, randomized, double-blind, placebo-controlled trial to evaluate the efficacy and safety of saxagliptin in combination with metformin and a sulfonylurea in patients with inadequate glycemic control (A1C \geq 7% and \leq 10%) on a stable combined dose of metformin (\geq 1500 mg) and sulfonylurea (\geq 50% of the maximum recommended dose) for at least eight weeks prior to enrollment.

Patients who met eligibility criteria were entered in a 2-week enrollment period to allow assessment of inclusion/exclusion criteria. Following the 2-week enrollment period, eligible patients were randomized to either double-blind saxagliptin (5 mg once daily) or double-blind matching placebo for 24 weeks. During the 24-week double-blind treatment period, patients continued metformin and sulfonylurea at the same constant dose ascertained during enrollment. Sulfonylurea could be down titrated once in the case of a major hypoglycemic event or recurring minor hypoglycemic events. In the absence of hypoglycemia, titration (up or down) of study medication during the treatment period was prohibited.

Saxagliptin, in combination with metformin and a sulfonylurea, provided significant improvements in A1C and PPG compared with placebo in combination with metformin and a sulfonylurea (Table 9).

Table 9 Glycemic Parameters at Week 24 in a Placebo-Controlled Trial of Saxagliptin as Add-On Combination Therapy with Metformin and a Sulfonylurea*

Efficacy Parameter	Saxagliptin 5 mg + Metformin and a Sulfonylurea N=129	Placebo + Metformin and a Sulfonylurea N=128
A1C (%)	N=127	N=127
Baseline (mean)	8.4	8.2
Change from baseline (adjusted mean)	-0.7	-0.1
Difference from placebo (adjusted mean)	-0.7 [‡]	
95% Confidence Interval	(-0.9, -0.5)	
Percent of patients achieving A1C <7%	31% [§] (39/127)	9% (12/127)
2-hour PPG (mmol/L)	N=115	N=113
Baseline (mean)	14.85	14.54
Change from baseline (adjusted mean ±)	-0.65	0.28
Difference from placebo (adjusted mean ±)	-0.93 [¶]	
95% Confidence Interval	(-1.77, -0.09)	
FPG (mmol/L)	N=121	N=123
Baseline (mean)	8.99	8.63
Change from baseline (adjusted mean)	-0.29	0.15
Difference from placebo (adjusted mean)	-0.44 [#]	
95% Confidence Interval	(-0.94, 0.06)	

* Intent-to-treat population using last observation prior to discontinuation.

† Least squares mean adjusted for baseline value.

‡ p-value <0.0001 compared to placebo + metformin and a sulfonylurea

§ Significance not tested.

¶ p-value = 0.0301 compared to placebo + metformin and a sulfonylurea

Not statistically significant.

Patients with Renal Impairment

A 12-week, randomized, double-blind, placebo-controlled study was conducted to evaluate the treatment effect of saxagliptin 2.5 mg once daily compared with placebo in 170 patients with type 2 diabetes and renal impairment (85 patients on saxagliptin: moderate [n=48], severe [n=18]; or ESRD [n=19] and 85 patients on placebo). In this study, 98.2% of the patients entered the study on and continued antihyperglycemic medications (insulin and/or oral antihyperglycemic drug) other than the study drug (75.3% on insulin and 31.2% on oral antihyperglycemic drug; some received both).

Treatment with saxagliptin 2.5 mg provided significant improvement in A1C versus placebo (mean reduction from baseline at Week 12 of -0.9% for the saxagliptin group and -0.4% for the placebo group, p=0.007). Improvements in A1C following treatment with saxagliptin 2.5 mg were sustained up to Week 52, however the number of patients who completed 52 weeks without modification of other antihyperglycemic medications was low (26 subjects in the saxagliptin group versus 34 subjects in the placebo group).

There was no adverse effect on renal function at Week 12 or Week 52, consistent with experience

in patients with normal renal function in clinical trials. The safety results and profile of saxagliptin throughout the study were consistent with those previously observed in clinical trials.

Comparative Bioavailability Studies

A single dose, single center, randomized, two-way crossover comparative bioavailability study of Sandoz Saxagliptin (saxagliptin hydrochloride) 5 mg tablets (by Sandoz Canada Inc.) and Onglyza® (saxagliptin hydrochloride) 5 mg tablets (by AstraZeneca Canada Inc.), administered as a single 1 x 5 mg dose, was conducted in forty (40) healthy volunteers (14 males and 26 females) under fasting conditions. Please see the table below for the results obtained from the study:

SUMMARY TABLE OF THE COMPARATIVE BIOAVAILABILITY DATA

Saxagliptin (1 x 5 mg) From measured data uncorrected for potency Geometric LS Mean Arithmetic Mean (CV %)				
Parameter	Test*	Reference†	% Ratio of Geometric LS Means	90% Confidence Interval (%)
AUC _T (ng·h/mL)	105.037	104.741	100.3	97.2 - 103.4
	108.637 (26.9)	107.765 (25.2)		
AUC _∞ (ng·h/mL)	106.883	106.678	100.2	97.2 - 103.3
	110.470 (26.7)	109.698 (25.2)		
C _{max} (ng/mL)	31.710	30.053	105.5	99.6 - 111.8
	33.277 (29.0)	31.088 (25.0)		
T _{max} [§] (h)	0.55 (0.33 - 2.00)	0.75 (0.33 - 2.00)		
T _{1/2} ^ε (h)	3.51 (40.9)	3.41 (26.9)		

* Sandoz Saxagliptin 5 mg saxagliptin tablet manufactured by Lek Pharmaceuticals.

† Onglyza® 5 mg saxagliptin tablet, AstraZeneca Canada Inc, Canada (country of purchase).

§ Expressed as median (range)

ε Expressed as the arithmetic mean only (CV%)

DETAILED PHARMACOLOGY

Saxagliptin and its major metabolite are potent reversible inhibitors of DPP-4 *in vitro* with selectivity for DPP-4 versus other enzymes, including other DPP family members such as DPP-8 and DPP-9. Saxagliptin and its major metabolite have extended binding to the DPP-4 active site, prolonging their activity, but do not have extended duration of binding to other enzymes, including DPP-8 and DPP-9. Saxagliptin was a potent inhibitor of T-cell cell surface DPP activity in cell based assays, but did not inhibit T-cell activation either *in vitro* or *in vivo*.

Saxagliptin, when dosed orally, demonstrated dose-related inhibition of DPP-4 in *ex vivo* assays in rats, dogs and cynomolgus monkeys. In acute *in vivo* studies, saxagliptin increased concentrations of intact GLP-1 in response to a meal in lean rats (maximum effect at 1 mg/kg). Saxagliptin also increased plasma insulin and lowered plasma glucose following an oral glucose tolerance test in obese insulin resistant and diabetic animal rodent models (maximum effect range 0.4 to 1.3 mg/kg). In chronic dosing studies using the progressively diabetic ZDF rat model, saxagliptin (4 mg/kg/day) delayed development of fasting hyperglycemia and the results of oral glucose tolerance tests showed significantly improved glucose homeostasis. These results are consistent with the mechanism of action of saxagliptin and its effects as an anti-hyperglycemic agent.

TOXICOLOGY

Acute Toxicity

Saxagliptin was observed to be well tolerated at single doses up to 2000 mg/kg in mice and rats and 25 mg/kg in cynomolgus monkeys. In rodents, 4000 mg/kg resulted in transient decreases in body-weight gain and activity and/or lethality. In monkeys, overt toxicity and lethality were observed at 50 mg/kg.

Chronic Toxicity

The potential toxicity of saxagliptin was evaluated in a number of repeat-dose studies in mice, rats, dogs and monkeys. Saxagliptin administered to rats for 6 months at doses of 2, 20 and 100 mg/kg/day was well tolerated, causing only at the high dose, minimal splenic lymphoid hyperplasia and pulmonary histiocytosis. The no-observed-adverse-effect-level (20 mg/kg/day) was 36 times (males) and 78 times (females) the human exposure based on the recommended human dose of 5 mg/day (RHD). In dogs, saxagliptin administered orally at 5 and 10 mg/kg/day for 12 months caused toxicity in the intestinal tract, as evidenced by bloody and mucoid feces. The no-observed-adverse-effect-level was 1 mg/kg/day, 4 times the RHD. In monkeys, major target organ changes included skin lesions (scabs, erosions, and ulceration), lymphoid hyperplasia (primarily spleen and bone marrow) and multi-tissue mononuclear-cell infiltrates. Skin healing during the dosing period was observed with recovery of both skin and microscopic changes following a drug-free recovery period. The AUCs at the no effect level for these changes were 1 to 3 times the RHD.

Carcinogenicity

Two-year carcinogenicity studies were conducted in mice and rats at oral doses of 50, 250, and 600 mg/kg/day and 25, 75, 150, and 300 mg/kg/day, respectively. Saxagliptin did not induce tumors in either mice or rats at the highest doses evaluated. The highest doses evaluated in mice were

equivalent to approximately 900 (males) and 1210 (females) times the human exposure at the recommended human dose of 5 mg/day (RHD). In rats, AUC exposures were approximately 370 (males) and 2300 (females) times the RHD.

Mutagenesis

The mutagenic and clastogenic potential of saxagliptin was tested at high concentrations and exposures in a battery of genetic toxicity studies including an *in vitro* Ames bacterial assay, an *in vitro* cytogenetics assay in primary human lymphocytes, an *in vivo* oral micronucleus assay in rats, an *in vivo* oral DNA repair study in rats, and an oral *in vivo/in vitro* cytogenetics study in rat peripheral blood lymphocytes. Saxagliptin was not mutagenic or clastogenic based on the combined outcomes of these studies. The major metabolite was not mutagenic in an *in vitro* Ames bacterial assay.

Reproduction

In a rat fertility study, males were treated with oral gavage doses of 100, 200, and 400 mg/kg/day for two weeks prior to mating, during mating, and up to scheduled termination (approximately four weeks total) and females were treated with oral gavage doses of 125, 300, and 750 mg/kg/day for two weeks prior to mating through gestation day 7. No adverse effects on fertility were observed at 200 mg/kg/day (males) or 125 mg/kg/day (females) resulting in respective exposures (AUC) of approximately 630 (males) and 805 (females) times human exposure at the RHD. At higher, maternally toxic doses (300 and 750 mg/kg/day), increased fetal resorptions were observed (approximately 2150 and 6375 times the RHD). Additional effects on estrous cycling, fertility, ovulation, and implantation were observed at 750 mg/kg (approximately 6375 times the RHD).

Development

Saxagliptin was not teratogenic at any dose evaluated in rats or rabbits. At high doses in rats, saxagliptin caused a minor and reversible developmental delay in ossification of the fetal pelvis at ≥ 240 mg/kg/day (≥ 1560 times the human exposure [AUC] at the RHD). Maternal toxicity and reduced fetal body weights were observed at 900 mg/kg/day (8290 times the RHD). In rabbits, the effects of saxagliptin were limited to minor skeletal variations observed only at maternally toxic doses (200 mg/kg/day, exposures 1420 times the RHD).

Saxagliptin administered to female rats from gestation day 6 to lactation day 20 resulted in decreased body weights in male and female offspring only at maternally toxic doses (≥ 250 mg/kg/day, exposures ≥ 1690 times the RHD). No functional or behavioral toxicity was observed in offspring of rats administered saxagliptin at any dose.

REFERENCES

1. Augeri DJ, Robl JA, Betebenner DA et al. Discovery and preclinical profile of saxagliptin (BMS-477118): A highly potent, long-acting, orally active dipeptidyl peptidase IV inhibitor for the treatment of type 2 diabetes. *J Med. Chem.* 2005;48:5025-37
2. Canadian Diabetes Association Clinical Practice Guidelines Expert Committee. 2008 Clinical Practice Guidelines for the Prevention and Management of Diabetes in Canada. *Canadian J Diabetes* 2008;32(Suppl 1):S1-S201.
3. Chacra AR, Tan GH, Apanovitch A et al. Saxagliptin added to a submaximal dose of sulphonylurea improves glycaemic control compared with uptitration of sulphonylurea in patients with type 2 diabetes: a randomised controlled trial. *Int J Clin Pract* 2009;63(9):1395-1406.
4. DeFronzo RA, Hissa MN, Garber AJ et al. The efficacy and safety of saxagliptin when added to metformin therapy in patients with inadequately controlled type 2 diabetes with metformin alone. *Diabetes Care* 2009;32:1649-55
5. Drucker DJ. Enhancing incretin action for the treatment of type 2 diabetes. *Diabetes Care* 2003;26:2929-40
6. Kim YB, Kopcho LM, Kirby MS et al. Mechanism of Gly-Pro-pNA cleavage catalyzed by dipeptidyl peptidase-IV and its inhibition by saxagliptin (BMS-477118). *Arch Biochem Biophys* 2006;445:9-18.
7. Metzler WJ, Yanchunas J, Weigelt C et al. Involvement of DPP-IV catalytic residues in enzyme saxagliptin complex formation. *Protein Sci.* 2008;17:240-50.
8. Rosenstock J, Sankoh S and List JF. Glucose-lowering activity of the dipeptidyl peptidase-4 inhibitor saxagliptin in drug-naïve patients with type 2 diabetes. *Diabetes, Obesity and Met* 2008;10:376–86.
9. Vilsboll T, Krarup T, Deacon C, Madsbad S, Holst J. Reduced postprandial concentrations of intact biologically active glucagons-like peptide 1 in type 2 diabetic patients. *Diabetes* 2001;50(3):609-613.
10. Scirica BM, Bhatt DL, Braunwald E, Steg PG, Davidson J et al. Saxagliptin and cardiovascular outcomes in patients with type 2 diabetes mellitus. *N Engl J Med* 2013; 369:1317-1326.
11. Product Monograph, ONGLYZA[®], saxagliptin tablets (as saxagliptin hydrochloride) 2.5 mg and 5 mg, Control #:212634, Date of Revision: December 12, 2018.

PART III: CONSUMER INFORMATION

Pr SANDOZ SAXAGLIPTIN
 Saxagliptin Tablets
 Saxagliptin (as saxagliptin hydrochloride dihydrate)
 2.5 mg and 5 mg

This leaflet is part III of a three-part "Product Monograph" published when Sandoz Saxagliptin was approved for sale in Canada and is designed specifically for consumers. This leaflet is a summary and will not tell you everything about Sandoz Saxagliptin. Contact your doctor or pharmacist if you have any questions about the drug.

ABOUT THIS MEDICATION

What the medication is used for:

Sandoz Saxagliptin is used to improve blood sugar levels in adult patients with type 2 diabetes, in combination with;

- metformin,
- sulfonylurea,
- metformin and a sulfonylurea or
- insulin (with or without metformin)

when diet and exercise and metformin, sulfonylurea, metformin and a sulfonylurea or insulin (with or without metformin) alone have failed to adequately control blood sugar levels.

What it does:

Sandoz Saxagliptin belongs to a class of medicines called DPP-4 inhibitors (dipeptidyl peptidase-4 inhibitors). Sandoz Saxagliptin helps to improve blood sugar levels in response to a meal. Sandoz Saxagliptin also lowers blood sugar levels between meals, and helps to decrease the amount of sugar made by your body.

What is Type 2 Diabetes?

Insulin is a hormone that helps control the level of sugar (glucose) in your blood. Type 2 diabetes is a condition in which your body does not make enough insulin and/or the insulin that your body produces does not work as well as it should. When this happens, glucose can build up in the blood. This can lead to serious problems.

When it should not be used:

- Do not take Sandoz Saxagliptin if you are allergic to any of the ingredients in Sandoz Saxagliptin or if you are allergic to the other drugs belonging to DPP-4 class.
- Do not take Sandoz Saxagliptin if you have diabetic ketoacidosis (accumulation of ketones in the blood and urine), diabetic pre-coma or diabetic coma.
- Do not take Sandoz Saxagliptin if you have Type 1 diabetes.

What the medicinal ingredient is:

Saxagliptin hydrochloride

What the nonmedicinal ingredients are:

Non-medicinal ingredients: lactose monohydrate, microcrystalline cellulose, croscarmellose sodium, magnesium stearate and (dilute) hydrochloric acid. In addition, the film coating contains the following inactive ingredients: hypromellose, macrogol, lactose anhydrous, titanium dioxide and yellow iron oxide (2.5 mg) or red iron oxide (5 mg).

What dosage forms it comes in:

Sandoz Saxagliptin is supplied in 2.5 mg and 5 mg (as saxagliptin hydrochloride) film-coated tablets.

WARNINGS AND PRECAUTIONS

BEFORE you use Sandoz Saxagliptin talk to your doctor or pharmacist if:

- you have had an allergic reaction to other DPP-4 inhibitors
- you have or have had any kidney problems
- you have or have had liver problems
- you have or have had heart failure
- you have been told you have a reduced immune system [e.g. you have had organ transplantation or have been diagnosed with human immunodeficiency syndrome (HIV/AIDS)]
- you are pregnant or planning to become pregnant
- you are breast-feeding or plan to breast-feed
- you have or have had pancreas problems such as inflammation of the pancreas (pancreatitis)

Heart Failure has been seen in patients treated with Sandoz Saxagliptin. **Heart Failure** is when your heart is unable to pump enough blood to meet the needs of the body. You are at greater risk of **Heart Failure** if you have or have had:

- heart or blood vessel disease including heart failure and heart attack
- kidney disease
- several risk factors of getting heart disease

Symptoms of heart failure include one or more of the following: tiredness, swollen ankles, a fast increase in weight and increased shortness of breath especially when lying down. This is serious. You must talk to your physician immediately or go to the hospital if this happens to you.

Sandoz Saxagliptin should not be used in patients with end-stage renal disease (ESRD) who require dialysis.

Lactose monohydrate is a non-medicinal ingredient in Sandoz Saxagliptin. Do not take Sandoz Saxagliptin if a doctor has told you that you have one of the following hereditary diseases: galactose intolerance, Lapp lactase deficiency, or glucose-galactose malabsorption.

Your blood sugar may get too high (hyperglycemia) if you have fever, infection, surgery, or trauma (stress conditions). In such

cases contact your doctor as your medication may need to be adjusted.

INTERACTIONS WITH THIS MEDICATION

Talk to your doctor or pharmacist about all the drugs you take. This includes prescription drugs, as well as those you buy yourself, and herbal supplements.

PROPER USE OF THIS MEDICATION

Follow the directions given to you by your doctor.

Usual Dose:

The usual recommended adult dose is 5 mg, once daily, taken with or without food.

You may need a lower dose of Sandoz Saxagliptin if your kidneys are not working well.

Sandoz Saxagliptin tablets must not be split or cut.

Overdose:

If you think you have taken too much Sandoz Saxagliptin, call your doctor or pharmacist right away, or contact your local poison control centre immediately.

Missed Dose:

If you miss a dose of Sandoz Saxagliptin, take it as soon as you remember. If you do not remember until it is almost time for your next dose, skip the missed dose and go back to your regular schedule. Do not take a double dose.

SIDE EFFECTS AND WHAT TO DO ABOUT THEM

The common side effects of Sandoz Saxagliptin include: upper respiratory tract infection, urinary tract infection and headache. Hypoglycemia may occur more frequently in people who already take a sulfonylurea or insulin. If you have symptoms of low blood sugar, you should check your blood sugar and treat if low, then call your healthcare provider. Symptoms of low blood sugar include shaking, sweating, rapid heartbeat, change in vision, hunger, headache, and change in mood. Tell your physician or pharmacist if you develop any unusual side effects, or if any of the above side effects do not go away or get worse.

SERIOUS SIDE EFFECTS, HOW OFTEN THEY HAPPEN AND WHAT TO DO ABOUT THEM				
Symptom / Effect		Talk with your doctor or pharmacist		Stop taking drug and call your doctor or pharmacist
		Only if severe	In all cases	
Uncommon	Pancreatitis (inflammation of the pancreas): prolonged severe abdominal pain which may be accompanied by vomiting		✓	✓
	Severe disabling joint pain		✓	
Very rare	Allergic (hypersensitivity) reactions: rash, hives, swelling of the face, lips and throat that may cause difficulty in breathing or swallowing		✓	✓
	Bullous pemphigoid (serious skin reaction): blistering of the skin, redness, peeling skin		✓	
Unknown	Heart failure (a weakness of the heart): tiredness, swollen ankles, increasing shortness of breath especially when lying down and a fast increase in weight.			✓

This is not a complete list of side effects. For any unexpected effects while taking Sandoz Saxagliptin, contact your doctor or pharmacist.

HOW TO STORE IT

Store at room temperature (15-30°C). Store in the original packaging. Protect from moisture. Keep well out of reach and sight of children.

SERIOUS SIDE EFFECTS, HOW OFTEN THEY HAPPEN AND WHAT TO DO ABOUT THEM

Symptom / Effect	Talk with your doctor or pharmacist		Stop taking drug and call your doctor or pharmacist
	Only if severe	In all cases	

REPORTING SIDE EFFECTS

You can report any suspected side effects associated with the use of health products to Health Canada by:

- Visiting the Web page on Adverse Reaction Reporting (<https://www.canada.ca/en/health-canada/services/drugs-health-products/medeffect-canada.html>) for information on how to report online, by mail or by fax; or
- Calling toll-free at 1-866-234-2345.

NOTE: Contact your health professional if you need information about how to manage your side effects. The Canada Vigilance Program does not provide medical advice.

MORE INFORMATION

If you want more information about Sandoz Saxagliptin:

- Talk to your healthcare professional
- Find the full product monograph that is prepared for healthcare professionals and includes this Patient Medication Information by visiting the Health Canada website (<https://health-products.canada.ca/dpd-bdpp/index-eng.jsp>); the manufacturer's website www.sandoz.com, or by calling 1-800-361-3062.

Or
by written request at:

110 Rue de Lauzon Boucherville, Québec, Canada, J4B 1E6

Or by e-mail at: medinfo@sandoz.com

This leaflet was prepared by Sandoz Canada Inc.

Last revised: February 25, 2019