

PRODUCT MONOGRAPH
INCLUDING PATIENT MEDICATION INFORMATION

Pr ATHENA-ONDANSETRON ODT

Ondansetron orally disintegrating tablets

Orally disintegrating tablets, 4 mg and 8 mg, oral

Manufacturer Standard

Antiemetic

5-HT₃ receptor antagonist

ATC code A04AA01

Athena Pharmaceutiques SAS
Espace Arnold de Ville,
12 rue Georges Blandon,
78430 Louveciennes,
France

Date of Initial Authorization:
JUN 21, 2021

Date of Revision:
JAN 17, 2022

Imported by:
Athena Canada Inc.
Vaughan, Ontario
L4K 3P3

Submission Control Number: 255400

RECENT MAJOR LABEL CHANGES

1 INDICATIONS	01/2022
4 DOSAGE AND ADMINISTRATION, 4.1 Dosing Considerations	01/2022
7 WARNINGS AND PRECAUTIONS, Cardiovascular, Myocardial Ischemia and Coronary Artery Spasm	01/2022
7 WARNING AND PRECAUTIONS, Special Populations, 7.1.1 Pregnant Women	01/2022

TABLE OF CONTENTS

Sections or subsections that are not applicable at the time of authorization are not listed.

RECENT MAJOR LABEL CHANGES	2
TABLE OF CONTENTS	2
PART I: HEALTH PROFESSIONAL INFORMATION	4
1 INDICATIONS	4
1.1 Pediatrics (< 18 years of age)	4
1.2 Geriatrics (≥ 65 years of age)	4
2 CONTRAINDICATIONS	4
4 DOSAGE AND ADMINISTRATION	5
4.1 Dosing Considerations.....	5
4.2 Recommended Dose and Dosage Adjustment.....	6
4.4 Administration.....	8
5 OVERDOSAGE	8
6 DOSAGE FORMS, STRENGTHS, COMPOSITION AND PACKAGING	8
7 WARNINGS AND PRECAUTIONS	9
7.1 Special Populations.....	12
7.1.1 Pregnant Women.....	12
7.1.2 Breast-feeding	12
7.1.3 Pediatrics.....	12
7.1.4 Geriatrics.....	12
8 ADVERSE REACTIONS	13
8.2 Clinical Trial Adverse Reactions	13
8.5 Post-Market Adverse Reactions.....	14

9	DRUG INTERACTIONS	15
	9.1 Serious Drug Interactions	15
	9.2 Drug Interactions Overview	15
	9.4 Drug-Drug Interactions.....	16
	9.5 Drug-Food Interactions	18
	9.6 Drug-Herb Interactions.....	18
	9.7 Drug-Laboratory Test Interactions	18
10	CLINICAL PHARMACOLOGY.....	18
	10.1 Mechanism of Action	18
	10.2 Pharmacodynamics.....	18
	10.3 Pharmacokinetics.....	20
11	STORAGE, STABILITY AND DISPOSAL.....	23
	PART II: SCIENTIFIC INFORMATION	24
13	PHARMACEUTICAL INFORMATION	24
14	CLINICAL TRIALS	24
	14.1 Trial Design and Study Demographics	24
	14.2 Study Results	24
	14.3 Comparative Bioavailability Studies	26
16	NON-CLINICAL TOXICOLOGY	27
17	SUPPORTING PRODUCT MONOGRAPHS	31
	PATIENT MEDICATION INFORMATION	32

PART I: HEALTH PROFESSIONAL INFORMATION

1 INDICATIONS

Adults (18 - 64 years of age)

ATHENA-ONDANSETRON ODT (ondansetron) is indicated for:

- the prevention of nausea and vomiting associated with mildly and moderately emetogenic chemotherapy and radiotherapy.
- the maintenance of antiemesis following intravenous doses of ondansetron used for the prevention of nausea and vomiting associated with highly emetogenic chemotherapy, including cisplatin.
- the prevention and treatment of post-operative nausea and vomiting.

1.1 Pediatrics (< 18 years of age)

Post-Chemotherapy Induced Nausea and Vomiting

- **Pediatrics (4 - 12 years of age):** ATHENA-ONDANSETRON ODT was effective and well tolerated when given to children 4-12 years of age.
- **Pediatrics (< 4 years of age):** ATHENA-ONDANSETRON ODT is not indicated for the treatment of children less than 4 years old.

Post-Radiotherapy Induced Nausea and Vomiting

ATHENA-ONDANSETRON ODT is not indicated for this use in any pediatric population.

Post-Operative Nausea and Vomiting

ATHENA-ONDANSETRON ODT is not indicated for this use in any pediatric population.

1.2 Geriatrics (≥ 65 years of age)

Post-Chemotherapy and Radiotherapy Induced Nausea and Vomiting

Efficacy and tolerance of ATHENA-ONDANSETRON ODT were similar to that observed in younger adults. See [4.2 Recommended Dose and Dosage Adjustment, Use in Elderly \(≥ 65 years of age\); 7 WARNINGS AND PRECAUTIONS Geriatrics](#).

Post-Operative Nausea and Vomiting

Clinical experience in the use of ATHENA-ONDANSETRON ODT in the prevention and treatment of post-operative nausea and vomiting is limited; ATHENA-ONDANSETRON ODT is not indicated for this use in the geriatric population.

2 CONTRAINDICATIONS

- ATHENA-ONDANSETRON ODT (ondansetron) is contraindicated in patients with a history of hypersensitivity to the drug or any components of its formulations. For a complete listing, see [6 DOSAGE FORMS, STRENGTHS, COMPOSITION AND PACKAGING](#).
- The concomitant use of apomorphine with ondansetron is contraindicated based on reports of profound hypotension and loss of consciousness when apomorphine was administered with ondansetron.

4 DOSAGE AND ADMINISTRATION

4.1 Dosing Considerations

Note: Athena Pharmaceutiques SAS only markets ATHENA-ONDANSETRON ODT (ondansetron orally disintegrating tablets). When injectable ondansetron is used, the product monograph for ondansetron hydrochloride dihydrate injection should be consulted.

Ondansetron clearance is reduced in patients with moderate or severe hepatic impairment. Their total daily dose should not exceed 8 mg, which may be given as a single intravenous or oral dose. See [7 WARNINGS AND PRECAUTIONS, Hepatic](#).

Ondansetron has important cardiac side-effects (dose-dependent QTc prolongation, coronary artery spasm, myocardial ischemia, and sequelae). These effects are reported more often with intravenous administration, and are expected to be greater with a faster rate of infusion. See [7 WARNINGS AND PRECAUTIONS, Cardiovascular, QTc Interval Prolongation; and Myocardial Ischemia and Coronary Artery Spasm](#); [9.4 Drug-Drug Interactions, QTc-Prolonging Drugs](#); [10.2 Pharmacodynamics, Electrocardiography](#).

Though ondansetron efficacy and tolerance were similar for elderly compared to younger adults in chemotherapy clinical trials, exposure-response modelling predicted a greater effect on QTcF in patients ≥ 75 years of age compared to young adults. See [10.3 Pharmacokinetics, Geriatrics](#).

Dosing considerations that reduce cardiac risks:

- Use the minimum effective dose.
- Use oral formulations if possible (lower C_{max}).

The efficacy of ondansetron in highly emetogenic chemotherapy requires the addition of a single intravenous dose of dexamethasone sodium phosphate 20 mg administered prior to chemotherapy.

The efficacy of twice daily dosage regimens for the treatment of post-chemotherapy emesis has been established only in adult patients receiving **less** emetogenic chemotherapy. The appropriateness of twice versus three times daily dosage regimens for other patient groups should be based on an assessment of the needs and responsiveness of the individual patient.

4.2 Recommended Dose and Dosage Adjustment

Highly Emetogenic Chemotherapy Induced Nausea and Vomiting

Caution: To reduce cardiac risks, carefully follow the dosing guidelines under [4.1 Dosing Considerations](#).

Note: Athena Pharmaceutiques SAS only markets ATHENA-ONDANSETRON ODT (ondansetron orally disintegrating tablets). When injectable ondansetron is used, the product monograph for ondansetron hydrochloride dihydrate injection should be consulted.

Initial Dose for Prevention of Emesis during the First 24 h Following Chemotherapy:

ATHENA-ONDANSETRON ODT is not appropriate. An intravenous formulation of ondansetron should be used as an initial dose prior to and in the first 24 hours following chemotherapy.

Post-chemotherapy:

After the first 24 hours, ondansetron 8 mg may be taken orally every 8 hours for up to 5 days.

Less Emetogenic Chemotherapy Induced Nausea and Vomiting

Caution: To reduce cardiac risks, carefully follow the dosing guidelines under [4.1 Dosing Considerations](#).

Note: Athena Pharmaceutiques SAS only markets ATHENA-ONDANSETRON ODT (ondansetron orally disintegrating tablets). When injectable ondansetron is used, the product monograph for ondansetron hydrochloride dihydrate injection should be consulted.

- **Use in Adults**
 - Initial dose:
 - 8 mg orally, given 1-2 hours before chemotherapy.
 - After chemotherapy: doses of 8 mg orally, twice daily, for up to 5 days.
- **Use in Children 4 - 12 years of age**
 - After chemotherapy: 4 mg orally, every 8 hours.
- **Use in Children < 4 years of age**

ATHENA-ONDANSETRON ODT is not indicated for this use in this pediatric population.
- **Use in Elderly (≥ 65 years of age)**

See [10.3 Pharmacokinetics, Geriatrics](#).

Radiotherapy Induced Nausea and Vomiting

Product Monograph

Athena-Ondansetron ODT (ondansetron)

Page 6 of 40

Note: Athena Pharmaceutiques SAS only markets ATHENA-ONDANSETRON ODT (ondansetron orally disintegrating tablets). When injectable ondansetron is used, the product monograph for ondansetron hydrochloride dihydrate injection should be consulted.

Caution: To reduce cardiac risks, carefully follow the dosing guidelines under [4.1 Dosing Considerations](#).

- **Use in Adults**
 - Initial dose: 8 mg orally, given 1 to 2 hours before radiotherapy.
 - After radiotherapy: 8 mg orally, given every 8 hours, for up to 5 days after a course of treatment.

- **Use in Children (< 18 years of age)**

ATHENA-ONDANSETRON ODT is not indicated for this use in the pediatric population.

- **Use in Elderly (≥ 65 years of age)**

See [Use in Adults](#), and [4.1 Dosing Considerations](#); [10.3 Pharmacokinetics, Geriatrics](#).

Post-Operative Nausea and Vomiting

Note: Athena Pharmaceutiques SAS only markets ATHENA-ONDANSETRON ODT (ondansetron orally disintegrating tablets). When injectable ondansetron is used, the product monograph for ondansetron hydrochloride dihydrate injection should be consulted.

- **Use in Adults**
 - For prevention:
 - One hour prior to anaesthesia: 16 mg orally.
 - OR**
 - At induction of anaesthesia: 4 mg IV, undiluted, infused - preferably over 2-5 minutes, and not less than 30 seconds.
 - For established post-operative nausea and vomiting:
 - A single dose of 4 mg IV, infused - preferably over 2-5 minutes, and not less than 30 seconds.

- **Use in Children (< 18 years of age)**

ATHENA-ONDANSETRON ODT is not indicated for this use in the pediatric population.

- **Use in Elderly (≥ 65 years of age)**

ATHENA-ONDANSETRON ODT is not indicated for this use in the elderly.

4.4 Administration

To take ATHENA-ONDANSETRON ODT (ondansetron orally disintegrating tablets):

1. Tear along the perforations of the foil to separate off one tablet within its blister unit.
2. Gently push the ATHENA-ONDANSETRON ODT out of the blister aluminium foil and remove it with dry fingers.
3. Immediately place ATHENA-ONDANSETRON ODT on top of the tongue. Allow it to melt. Apply gentle pressure if required. Swallow the pasty mass formed.

5 OVERDOSAGE

At present there is little information concerning overdose with ondansetron. Individual doses of 84 mg and 145 mg and total daily doses as large as 252 mg have been administered with only mild side effects. There is no specific antidote for ondansetron, therefore, in cases of suspected overdose, symptomatic and supportive therapy should be given as appropriate.

The use of Ipecac to treat overdose with ondansetron is not recommended as patients are unlikely to respond due to the antiemetic action of ondansetron itself.

“Sudden blindness” (amaurosis) of 2 to 3 minutes duration plus severe constipation occurred in one patient that was administered 72 mg of ondansetron intravenously as a single dose. Hypotension (and faintness) occurred in another patient that took 48 mg of oral ondansetron. Following infusion of 32 mg over only a 4-minute period, a vasovagal episode with transient second-degree heart block was observed. Neuromuscular abnormalities, autonomic instability, somnolence, and a brief generalized tonic-clonic seizure (which resolved after a dose of benzodiazepine) were observed in a 12 months’ old infant who ingested seven or eight 8-mg ondansetron tablets (approximately forty times the recommended 0.1 - 0.15 mg/kg dose for a paediatric patient). In all instances, the events resolved completely.

Ondansetron prolongs QT interval in a dose-dependent fashion (see [10.2 Pharmacodynamics](#)). ECG monitoring is recommended in cases of overdose.

Cases consistent with serotonin syndrome have been reported in young children following oral overdose.

For management of a suspected drug overdose, contact your regional poison control centre.

6 DOSAGE FORMS, STRENGTHS, COMPOSITION AND PACKAGING

Table 1: Dosage Forms, Strengths, Composition and Packaging

Route of Administration	Dosage Form / Strength / Composition	Non-medicinal Ingredients
-------------------------	--------------------------------------	---------------------------

Oral	Orally disintegrating tablets / 4 mg and 8 mg ondansetron (base)	Amino methacrylate copolymer, aspartame (see 7 WARNINGS AND PRECAUTIONS General), colloidal silicon dioxide, magnesium stearate, mannitol, polyvinyl pyrrolidone and strawberry flavour.
------	---	---

Description

ATHENA-ONDANSETRON ODT 4 mg and 8 mg orally disintegrating tablets:

White to off-white, round, flat, bevelled edged tablets with no markings on either side.

Each 4 mg tablet contains 4 mg ondansetron (base) and each 8 mg tablet contains 8 mg ondansetron (base). ATHENA-ONDANSETRON ODT orally disintegrating tablets also contain amino methacrylate copolymer, aspartame, colloidal silicon dioxide, magnesium stearate, mannitol, polyvinyl pyrrolidone and strawberry flavour.

ATHENA-ONDANSETRON ODT orally disintegrating tablets (ondansetron) are packaged in foil blister packs of 10 orally disintegrating tablets (2 rows of 5 orally disintegrating tablets) per blister strip per box.

7 WARNINGS AND PRECAUTIONS

General

ATHENA-ONDANSETRON ODT (ondansetron) contains aspartame and therefore should be taken with caution in patients with phenylketonuria.

ATHENA-ONDANSETRON ODT is not effective in preventing motion-induced nausea and vomiting.

Cardiovascular

QTc Interval Prolongation

Ondansetron prolongs the QT interval (see [10.2 Pharmacodynamics, Electrocardiography](#)). The magnitude of QTc prolongation will depend on the peak serum ondansetron concentration (C_{max}), which is substantially determined by the route of administration, the dose and the infusion rate of intravenous ondansetron. In addition, post-marketing cases of torsade de pointes have been reported in patients using ondansetron. Torsade de pointes is a polymorphic ventricular tachyarrhythmia. Generally, the risk of torsade de pointes increases with the magnitude of QTc prolongation produced by the drug. Torsade de pointes may be asymptomatic or experienced by the patient as dizziness, palpitations, syncope, or seizures. If sustained, torsade de pointes can progress to ventricular fibrillation and sudden cardiac death.

Avoid ondansetron in patients with congenital long QT syndrome. Ondansetron should be administered with caution to patients who have or may develop prolongation of QTc, including congestive heart failure, bradyarrhythmias or patients taking other medicinal products that lead to either QT prolongation or electrolyte abnormalities. See [9.4 Drug-Drug Interactions](#).

Hypokalaemia, hypocalcaemia and hypomagnesaemia should be corrected prior to ondansetron administration.

Additional risk factors for torsade de pointes in the general population include, but are not limited to, the following:

- female gender;
- age 65 years or older;
- baseline prolongation of the QT/QTc interval;
- presence of genetic variants affecting cardiac ion channels or regulatory proteins;
- family history of sudden cardiac death at < 50 years;
- cardiac disease (e.g., myocardial ischemia or infarction, left ventricular hypertrophy, cardiomyopathy, conduction system disease);
- history of arrhythmias (especially ventricular arrhythmias, atrial fibrillation, or recent conversion from atrial fibrillation);
- bradycardia (< 50 beats per minute);
- acute neurological events (e.g., intracranial or subarachnoid haemorrhage, stroke, intracranial trauma);
- nutritional deficits (e.g., eating disorders, extreme diets);
- diabetes mellitus;
- autonomic neuropathy.

Myocardial Ischemia and Coronary Artery Spasm

Ondansetron can cause coronary artery vasospasm and myocardial ischemia which may lead to myocardial infarction. In some cases, the symptoms appeared immediately after IV infusion, or shortly after oral administration, including after low doses in patients without significant known pre-existing cardiovascular disease or other risk factors. Caution is advised during and after ondansetron administration, and close monitoring is recommended in patients with known or suspected ischemic or vasospastic coronary artery disease or other significant underlying cardiovascular disease.

Driving and Operating Machinery

In psychomotor testing ondansetron does not impair performance nor cause sedation.

Gastrointestinal

As ondansetron is known to increase large bowel transit time, patients with signs of subacute intestinal obstruction should be monitored following administration.

Hepatic/Biliary/Pancreatic

Abnormal liver function test results have been reported, as well as liver failure in clinical trial cancer patients. See [8.2 Clinical Trial Adverse Reactions, Hepatic](#); and [8.5 Post-Market Adverse Reactions](#).

Immune

Cross-reactive hypersensitivity has been reported between different 5-HT₃ antagonists. Patients who have experienced hypersensitivity reactions to one 5-HT₃ antagonist have experienced more severe reactions upon being challenged with another drug of the same class. The use of a different 5-HT₃ receptor antagonist is not recommended as a replacement in cases in which a patient has experienced even a mild hypersensitivity type reaction to another 5-HT₃ antagonist.

Neurologic

Serotonin syndrome/Neuroleptic Malignant Syndrome: Cases of life-threatening serotonin syndrome or neuroleptic malignant syndrome-like events have been reported with 5-HT₃ receptor antagonist antiemetics, including ondansetron, when given in combination with other serotonergic and/or neuroleptic drugs. Serotonin syndrome symptoms may include mental status changes (e.g., agitation, hallucinations, coma), autonomic instability (e.g., tachycardia, labile blood pressure, hyperthermia), neuromuscular aberrations (e.g., hyperreflexia, incoordination) and/or gastrointestinal symptoms (e.g., nausea, vomiting, diarrhoea). As these syndromes may result in potentially life-threatening conditions, treatment should be discontinued if such events occur and supportive symptomatic treatment should be initiated. If concomitant treatment of ATHENA-ONDANSETRON ODT with a drug affecting the serotonergic neurotransmitter system is clinically warranted, careful observation of the patient is advised, particularly during treatment initiation and dose increases (see [9 DRUG INTERACTIONS](#)).

Reproductive Health: Female and Male Potential

Pregnancy status should be verified for females of reproductive potential prior to starting the treatment with ATHENA-ONDANSETRON ODT.

Females of reproductive potential should be advised that it is possible that ATHENA-ONDANSETRON ODT can cause harm to the developing foetus. Sexually active females of reproductive potential should use effective contraception (methods that result in less than 1% pregnancy rates) when using ATHENA-ONDANSETRON ODT and for two days after stopping treatment with ATHENA-ONDANSETRON ODT.

7.1 Special Populations

7.1.1 Pregnant Women

The use of ondansetron in pregnancy is not recommended. Ondansetron use during early pregnancy has been associated with a small increase in orofacial malformations. Despite some limitations in methodology, several human epidemiological studies noted an increase in orofacial clefts in infants of women administered ondansetron during the first trimester of pregnancy. Regarding cardiac malformations, the epidemiological studies showed conflicting results.

Ondansetron is not teratogenic in animals (see 16 [NON-CLINICAL TOXICOLOGY, Reproductive and Developmental Toxicology](#)).

7.1.2 Breast-feeding

Ondansetron is excreted in the milk of lactating rats. It is not known if it is excreted in human milk, however, nursing is not recommended during treatment with ondansetron.

7.1.3 Pediatrics

Insufficient information is available to provide dosage recommendations for children 3 years of age or younger.

7.1.4 Geriatrics

Early Phase I studies in healthy elderly volunteers showed a slight age-related decrease in clearance, and an increase in half-life of ondansetron. A greater effect on QTcF is predicted in patients ≥ 75 years of age compared to young adults, based on exposure-response modelling. See [10.3 Pharmacokinetics, Geriatrics](#).

8 ADVERSE REACTIONS

8.2 Clinical Trial Adverse Reactions

Because clinical trials are conducted under very specific conditions the adverse reaction rates observed in the clinical trials may not reflect the rates observed in practice and should not be compared to the rates in the clinical trials of another drug. Adverse drug reaction information from clinical trials is useful for identifying drug-related adverse events and for approximating rates.

Ondansetron has been administered to over 2500 patients worldwide in controlled clinical trials and has been well tolerated.

The most frequent adverse events reported in controlled clinical trials were headache (11%) and constipation (4%). Other adverse events include sensations of flushing or warmth (< 1%).

Cardiovascular

There have been rare reports of tachycardia, angina (chest pain), bradycardia, hypotension, syncope and electrocardiographical alterations.

Central Nervous System

There have been rare reports of seizures. Movement disorders and dyskinesia have been reported in two large clinical trials of ondansetron at a rate of 0.1 – 0.3%.

Dermatological

Rash has occurred in approximately 1% of patients receiving ondansetron.

Eye Disorder

Rare cases of transient visual disturbances (e.g. blurred vision) have been reported during or shortly after intravenous administration of ondansetron, particularly at rates equal to or greater than 30 mg in 15 minutes.

Hepatic/ Biliary / Pancreatic

There were transient increases of SGOT and SGPT of over twice the upper limit of normal in approximately 5% of patients. These increases did not appear to be related to dose or duration of therapy. There have been reports of liver failure and death in patients with cancer receiving concurrent medications including potentially hepatotoxic cytotoxic chemotherapy and antibiotics. The etiology of the liver failure is unclear.

Hypersensitivity

Rare cases of immediate hypersensitivity reactions sometimes severe, including anaphylaxis, bronchospasm, urticaria and angioedema have been reported.

Local Reactions

Pain, redness and burning at the site of injection have been reported.

Metabolic

There have been rare reports of hypokalaemia.

Other

There have been reports of abdominal pain, weakness and xerostomia

8.5 Post-Market Adverse Reactions

Over 250 million patient treatment days of ondansetron have been supplied since the launch of the product worldwide. The following events have been spontaneously reported during post-approval use of ondansetron, although the link to ondansetron cannot always be clearly established.

The adverse event profiles in children and adolescents were comparable to that seen in adults.

Immune Disorder

Rare cases of hypersensitivity reactions, sometimes severe (e.g., laryngeal oedema, stridor, laryngospasm and cardiopulmonary arrest) have also been reported.

Cardiovascular Disorders

There have been rare reports (< 0.01%) of myocardial infarction, myocardial ischemia, angina, chest pain with or without ST segment depression, arrhythmias (including ventricular or supraventricular tachycardia, premature ventricular contractions, and atrial fibrillation), electrocardiographic alterations (including second degree heart block), palpitations and syncope.

Rarely and predominantly with intravenous ondansetron, transient ECG changes including QTc interval prolongation, Torsade de Pointes, ventricular fibrillation, coronary artery spasm, myocardial ischemia, cardiac arrest, and sudden death have been reported. (see [7 WARNINGS AND PRECAUTIONS, Cardiovascular](#)).

Eye Disorder

There have been very rare cases of transient blindness following ondansetron treatment, generally within the recommended dosing range and predominantly during intravenous administration.

The majority of blindness cases reported resolved within 20 minutes. Although most patients had received chemotherapeutic agents, including cisplatin a few cases of transient blindness occurred following ondansetron administration for the treatment of post-operative nausea or vomiting and in the absence of cisplatin treatment. Some cases of transient blindness were reported as cortical in origin.

Hepatic/ Biliary / Pancreatic

Occasional asymptomatic increases in liver function tests have been reported.

Nervous System Disorders

Transient episodes of dizziness (<0.1%) have been reported predominantly during or upon completion of IV infusion of ondansetron.

Uncommon reports (< 1%) suggestive of extrapyramidal reactions including oculogyric crisis/dystonic reactions (e.g., oro-facial dyskinesia, opisthotonos, tremor, etc.), movement disorders and dyskinesia have been reported without definitive evidence of persistent clinical sequelae.

Serotonin syndrome and neuroleptic malignant syndrome-like events have been reported with 5-HT₃ receptor antagonist antiemetics, including ondansetron, when given in combination with other serotonergic and/or neuroleptic drugs (see [7 WARNINGS AND PRECAUTIONS, Neurologic](#)).

Respiratory, Thoracic and Mediastinal Disorders

There have also been rare reports of hiccups.

Skin and Subcutaneous Tissue Disorders

Very rare reports have been received for bullous skin and mucosal reactions, including fatal cases. These reports include toxic skin eruptions, such as Stevens-Johnson syndrome and toxic epidermal necrolysis, and have occurred in patients taking other medications that can be associated with bullous skin and mucosal reactions.

9 DRUG INTERACTIONS

9.1 Serious Drug Interactions

Serious Drug Interactions (see [9.4 Drug-Drug Interactions](#))

- Apomorphine (see [2 CONTRAINDICATIONS](#))
- QTc-Prolonging drugs
- Serotonergic agents

9.2 Drug Interactions Overview

Ondansetron is extensively metabolised by multiple hepatic cytochrome P450 enzymes (predominantly CYP3A4, also CYP2D6 and CYP1A2), and clearance is reduced in hepatic insufficiency (see [10.3 Pharmacokinetics, Hepatic Insufficiency](#)). CYP 3A4 inducers can increase ondansetron clearance (see [9.4. Drug-Drug Interactions, CYP 3A4 Inducers](#)).

Ondansetron does not itself appear to induce or inhibit the cytochrome P450 drug-

metabolizing enzyme system of the liver.

9.4 Drug-Drug Interactions

CYP 3A4 inducers

Despite the multiplicity of metabolic enzymes capable of metabolising ondansetron, which might be expected to compensate for an increase or decrease in enzyme activity, patients treated with inducers of CYP3A4 (i.e. phenytoin, carbamazepine, and rifampicin) demonstrated an increase in oral clearance of ondansetron and a decrease in ondansetron blood concentrations.

In a pharmacokinetic study of 16 epileptic patients maintained chronically on carbamazepine or phenytoin (CYP 3A4 inducers), reduction in AUC, C_{max} and $T_{1/2}$ of ondansetron was observed. This resulted in a significant increase in clearance. However, on the basis of the inter-subject variability in the available data, no dosage adjustment can be recommended.

CYP 2D6 inhibitors

No effect in ondansetron clearance secondary to enzyme inhibition has been identified to date.

QTc-Prolonging Drugs

The concomitant use of ATHENA-ONDANSETRON ODT with another QTc-prolonging drug should be carefully considered to determine that the therapeutic benefit outweighs the potential risk. Drugs that have been associated with QTc interval prolongation and/or torsade de pointes include, but are not limited to, the examples in the following list.

Chemical/pharmacological classes are listed if some, although not necessarily all, class members have been implicated in QTc prolongation and/or torsade de pointes:

- Class IA antiarrhythmics (e.g., quinidine, procainamide, disopyramide);
- Class III antiarrhythmics (e.g., amiodarone, sotalol, ibutilide, dronedarone);
- Class 1C antiarrhythmics (e.g., flecainide, propafenone);
- antiemetics (e.g., dolasetron, palonosetron, granisetron, droperidol, chlorpromazine, prochlorperazine);
- tyrosine kinase inhibitors (e.g., vandetanib, sunitinib, nilotinib, lapatinib);
- antipsychotics (e.g., chlorpromazine, pimozide, haloperidol, ziprasidone);
- antidepressants (e.g., citalopram, fluoxetine, venlafaxine, tricyclic/tetracyclic antidepressants e.g., amitriptyline, imipramine, maprotiline);
- opioids (e.g., methadone);
- domperidone;

- macrolide antibiotics and analogues (e.g., erythromycin, clarithromycin, telithromycin, tacrolimus);
- quinolone antibiotics (e.g., moxifloxacin, levofloxacin, ciprofloxacin);
- antimalarials (e.g., quinine, chloroquine);
- azole antifungals (e.g., ketoconazole, fluconazole, voriconazole);
- histone deacetylase inhibitors (e.g., vorinostat);
- beta-2 adrenoceptor agonists (e.g., salmeterol, formoterol).

Drugs that Cause Electrolyte Abnormalities

The use of ondansetron with drugs that can disrupt electrolyte levels should be avoided. Such drugs include, but are not limited to, the following:

- loop, thiazide, and related diuretics;
- laxatives and enemas;
- amphotericin B;
- high dose corticosteroids.

The above lists of potentially interacting drugs are not comprehensive. Current information sources should be consulted for newly approved drugs that prolong the QTc interval or cause electrolyte disturbances, as well as for older drugs for which these effects have recently been established.

Tramadol

Data from small studies indicate that ondansetron may reduce the analgesic effect of tramadol.

Apomorphine

Based on reports of profound hypotension and loss of consciousness when ondansetron was administered with apomorphine hydrochloride, concomitant use with apomorphine is contraindicated (see [2 CONTRAINDICATIONS](#)).

Serotonergic Drugs

As with other serotonergic agents, serotonin syndrome, a potentially life-threatening condition, may occur with 5-HT₃ receptor antagonist antiemetic treatment when given in combination with other agents that may affect the serotonergic neurotransmitter system [including triptans, SSRIs, SNRIs, lithium, sibutramine, fentanyl and its analogues, dextromethorphan, tramadol, tapentadol, meperidine, methadone and pertazocine or St. John's Wort (*Hypericum perforatum*)] and with drugs which impair metabolism of serotonin (such as MAOIs, including linezolid (an antibiotic which is a reversible non-selective MAOI) and methylene blue) (see [7 WARNINGS AND PRECAUTIONS, Neurologic](#)).

9.5 Drug-Food Interactions

Interactions with food have not been established.

9.6 Drug-Herb Interactions

Interactions with food have not been established.

9.7 Drug-Laboratory Test Interactions

Interactions with food have not been established.

10 CLINICAL PHARMACOLOGY

10.1 Mechanism of Action

Ondansetron is a selective antagonist of the serotonin receptor subtype, 5-HT₃. Its precise mode of action in the control of chemotherapy induced nausea and vomiting is not known.

Cytotoxic chemotherapy and radiotherapy are associated with the release of serotonin (5-HT) from enterochromaffin cells of the small intestine, presumably initiating a vomiting reflex through stimulation of 5-HT₃ receptors located on vagal afferents. Ondansetron may block the initiation of this reflex. Activation of vagal afferents may also cause a central release of serotonin from the chemoreceptor trigger zone of the area postrema, located on the floor of the fourth ventricle. Thus, the antiemetic effect of ondansetron is probably due to the selective antagonism of 5-HT₃ receptors on neurons located in either the peripheral or central nervous systems, or both.

The mechanisms of ondansetron's antiemetic action in post-operative nausea and vomiting are not known.

10.2 Pharmacodynamics

Serotonin receptors of the 5-HT₃ type are present both peripherally and on vagal nerve terminals. Ondansetron probably acts by preventing activation of these receptors or receptors located in other regions of the central nervous system. Both the peripheral and central nervous systems appear to be involved since both abdominal vagotomy and microinjection of ondansetron and other 5-HT₃ antagonists directly into the area postrema eliminate cisplatin-induced emesis, while 5-HT₁-like (methiothepin maleate) and 5-HT₂ (ketanserin) antagonists have no effect.

Ondansetron is highly selective for 5-HT₃ receptors and shows negligible binding to other receptors such as 5-HT₁-like, 5-HT₂, α ₁ and α ₂ adrenoceptors, β ₁ and β ₂ adrenoceptors, D₁ and D₂ muscarinic, nicotinic, GABA_A, H₁ and H₂ receptors.

The pharmacological specificity of ondansetron may explain the observed lack of

extrapyramidal side effects often seen following similar therapy with metoclopramide, which preferentially binds to dopamine receptors of the D2 subtype.

In vivo pharmacodynamic studies have investigated the effects of ondansetron on gastric emptying, small bowel transit time and oesophageal motility.

Both oral (16 mg tid) and intravenous (5-10 mg) doses of ondansetron failed to produce a significant effect on gastric emptying in both healthy volunteers and in patients suffering from delayed gastric emptying. However, in one study intravenous doses of 8 mg did increase gastric emptying in over half the volunteers tested.


Intravenous infusion of either 1 mg or 5 mg ondansetron tended to increase small bowel transit times and single intravenous doses of 10 mg ondansetron have been reported to decrease sphincter pressure in the lower oesophagus in some subjects.

Electrocardiography

The effect of ondansetron on the QTc interval was evaluated in a double blind, randomized, placebo and positive (moxifloxacin) controlled, crossover study in 58 healthy adult men and women. Ondansetron was tested at single doses of 8 mg and 32 mg infused intravenously over 15 minutes. At the highest tested dose of 32 mg, prolongation of the Fridericia-corrected QTc interval ($QT/RR^{0.33}=QTcF$) was observed from 15 min to 4 h after the start of the 15 min infusion, with a maximum mean (upper limit of 90% CI) difference in QTcF from placebo after baseline-correction of 19.6 (21.5) msec at 20 min. At the lower tested dose of 8 mg, QTc prolongation was observed from 15 min to 1 h after the start of the 15 minute infusion, with a maximum mean (upper limit of 90% CI) difference in QTcF from placebo after baseline - correction of 5.8 (7.8) msec at 15 min. The magnitude of QTc prolongation with ondansetron is expected to be greater if the infusion rate is faster than 15 minutes. The 32 mg intravenous dose of ondansetron must not be administered.

No treatment-related effects on the QRS duration or the PR interval were observed at either the 8 or 32 mg dose.

LS Mean Difference (90% CI) in QTcF Interval Between Treatment and Placebo Over Time



An ECG assessment study has not been performed for orally administered ondansetron. On the basis of pharmacokinetic-pharmacodynamic modelling, an 8 mg oral dose of ondansetron is predicted to cause a mean QTcF increase of 0.7 ms (90% CI -2.1, 3.3) at steady-state, assuming a mean maximal plasma concentration of 24.7 ng/mL (95% CI 21.1, 29.0).

The magnitude of QTc prolongation at the recommended 5 mg/m² dose in pediatrics has not been studied, but pharmacokinetic-pharmacodynamic modelling predicts a mean increase of 6.6 ms (90% CI 2.8, 10.7) at maximal plasma concentrations.

10.3 Pharmacokinetics

Absorption

- Oral administration


Pharmacokinetic studies in human volunteers showed peak plasma levels of 20-30 ng/mL at around 1½ hours after an 8 mg oral dose of ondansetron. Repeat dosing of an 8 mg tablet every 8 hours for 6 days increased the peak plasma value to 40 ng/mL.

- Intravenous administration


Pharmacokinetic studies in human volunteers: An 8 mg infusion of ondansetron resulted in peak plasma levels of 80-100 ng/mL. A continuous intravenous infusion of 1 mg/hour after the initial 8 mg loading dose of ondansetron maintained plasma levels over 30 ng/mL during the following 24-hour period.

Mean plasma concentration-time curves for ondansetron following 8 mg and 32 mg dose:

Linear Scale



Semi-logarithmic Scale



Distribution

The absolute bioavailability of ondansetron in humans was approximately 60% and the plasma protein binding was approximately 73%.

See also [16 NON-CLINICAL TOXICOLOGY, Non-clinical pharmacodynamics](#).

Metabolism

In vitro metabolism studies have shown that ondansetron is a substrate for human hepatic cytochrome P450 enzymes, including CYP1A2, CYP2D6 and CYP3A4. In terms of overall ondansetron turnover, CYP3A4 played the predominant role. Because of the multiplicity of metabolic enzymes capable of metabolising ondansetron, it is likely that inhibition or loss of one enzyme (e.g. CYP2D6 enzyme deficiency) will be compensated by others and may result in little change in overall rates of ondansetron clearance. CYP 3A4 inducers can increase

clearance (see [9.4 Drug-Drug Interactions, CYP 3A4 inducers](#)).

Elimination

Following extensive metabolism of an orally or intravenously administered dose, ondansetron is excreted in the urine and faeces. In humans, less than 10% of the dose is excreted unchanged in the urine. The major urinary metabolites are glucuronide conjugates (45%), sulphate conjugates (20%) and hydroxylation products (10%).

The half-life of ondansetron after either an 8 mg oral dose or intravenous dose was approximately 3-4 hours and may be extended to 6-8 hours in the elderly.

Special Populations and Conditions

Geriatrics

Early Phase I studies in healthy elderly volunteers showed a slight age-related decrease in clearance, and an increase in half-life of ondansetron. However, wide inter-subject variability resulted in considerable overlap in pharmacokinetic parameters between young (< 65 years of age) and elderly subjects (\geq 65 years of age) and there were no overall differences in safety or efficacy observed between young and elderly cancer patients enrolled in CINV clinical trials. (See [4.2 Recommended Dose and Dosage Adjustment, Use in Elderly](#))

Based on more recent ondansetron plasma concentrations and exposure-response modelling, a greater effect on QTcF is predicted in patients \geq 75 years of age compared to young adults. Specific dosing information is provided for patients over 65 years of age and over 75 years of age for intravenous dosing. (See [4.2 Recommended Dose and Dosage Adjustment, Use in Elderly](#))

Genetic Polymorphism

CYP 2D6

The elimination half-life and plasma levels of a single 8 mg intravenous dose of ondansetron did not differ between subjects classified as poor and extensive metabolisers of sparteine and debrisoquine (CYP 2D6 substrates). No alteration of daily dosage or frequency of ondansetron dosing is recommended for patients known to be CYP 2D6 poor metabolisers.

Hepatic Insufficiency

The clearance of an 8 mg intravenous dose of ondansetron was significantly reduced and the serum half-life significantly prolonged in subjects with severe impairment of hepatic function. In patients with moderate or severe impairment of hepatic function, reductions in dosage are therefore recommended and a total daily dose of 8 mg should not be exceeded. This may be given as a single intravenous or oral dose.

There is no experience in patients who are clinically jaundiced.

Renal Insufficiency

No alteration of daily dosage, frequency of dosing, or route of administration is required in patients with impaired renal function.

11 STORAGE, STABILITY AND DISPOSAL


ATHENA-ONDANSETRON ODT (ondansetron) orally disintegrating tablets should be stored at room temperature between 15 - 30°C.

PART II: SCIENTIFIC INFORMATION

13 PHARMACEUTICAL INFORMATION

Drug Substance

Proper name:	ondansetron
Chemical name:	1,2,3,9-tetrahydro-9-methyl-3-[(2-methyl-1H-imidazol-1-yl)methyl]-4H-carbazol-4-one.
Molecular Formula:	C ₁₈ H ₁₉ N ₃ O
Molecular mass:	293.4 g/mol
Structural formula:	



Description and Solubility:

Ondansetron is a white to off-white powder. It is soluble at pH 1.2. Practically insoluble in water. Solubility decreases with increasing pH from very slightly soluble at pH 3.5 and pH 5.4 to practically insoluble at pH 8. Soluble in chloroform and slightly soluble in acetonitrile and methanol.

14 CLINICAL TRIALS

14.1 Trial Design and Study Demographics

The clinical trial data on which the original indication was authorized is not available.

14.2 Study Results

Clinical trial results showing the number and percentage of patients exhibiting a complete response to ondansetron (0 emetic episodes) are shown in the tables below for both post-operative and chemotherapy induced emesis.

Dose	Ondansetron* 3 doses of 0.15 mg/kg	Placebo* 3 doses of placebo	Ondansetron 8 mg IV + 1 mg/hr, 24 hours	Ondansetron 8 mg IV	Ondansetron 32 mg IV
# of patients	14	14	168	152	173
Treatment Response:					
0 emetic episodes	2 (14%)	0 (0%)	92 (55%)	82 (54%)	97 (56%)
1 – 2 emetic episodes	8 (57%)	0 (0%)	-	-	-

*Results are from an initial study using a different dosing regimen.

Table 3: Prevention of Post-Operative Emesis – Response Over 24 Hours*

Dose	Oral Prevention			Intravenous Prevention		
	Ondansetron 16 mg od	Placebo	<i>p value</i>	Ondansetron 4 mg IV	Placebo	<i>p value</i>
# of patients	253	250		136	139	
Treatment Response:						
0 emetic episodes	126 (50%)	79 (32%)	< 0.001	103 (76%)	62 (46%)	< 0.001

* The majority of patients included in the prevention and treatment of post-operative nausea and vomiting studies using Ondansetron have been adult women receiving balanced anaesthesia for gynaecological surgery.

Table 4: Treatment of Post-Operative Emesis – Response Over 24 Hours*

Intravenous Treatment			
Dose	Ondansetron 4 mg IV	Placebo	<i>p value</i>
# of patients	104	117	
Treatment Response:			
0 emetic episodes	49 (47%)	19 (16%)	< 0.001

*The majority of patients included in the prevention and treatment of post-operative nausea and vomiting studies using Ondansetron have been adult women receiving balanced anaesthesia for gynaecological surgery.

Table 5: Prevention of Radiotherapy Induced Emesis – Response Over 24 Hours*

Oral Treatment			
Dose	Ondansetron 8 mg PO tid*	Metoclopramide 10 mg PO tid*	<i>p value</i>
# of patients	38	44	
Treatment Response: 0 emetic episodes	37 (97%)	20 (45%)	< 0.001

*results from a study of adult male and female patients receiving single high dose radiotherapy (800 to 1,000 cGy) over an anterior or posterior field size of ≥ 80 cm² to the abdomen.

*Patients received the first dose of Ondansetron 8 mg tablets or metoclopramide (10 mg) 1-2 hours before radiotherapy. If radiotherapy was given in the morning, 2 additional doses of study treatment were given (1 tablet late afternoon and 1 tablet before bedtime). If radiotherapy was given in the afternoon, patients took only 1 further tablet that day before bedtime. Patients continued oral medication on a 3 times a day basis for 3-5 days.

14.3 Comparative Bioavailability Studies

A single dose, single center, randomized, two-way crossover comparative bioavailability study of ATHENA-ONDANSETRON ODT 8 mg orally disintegrating tablets (manufactured by Athena Drug Delivery Solutions Pvt. Ltd., India) and Zofran® ODT 8 mg orally disintegrating tablets, USP (manufacturer GlaxoSmithKline Inc., Canada), administered as a single 1 x 8 mg dose, was conducted in healthy male and female volunteers (n=24) under fasting conditions. Please see the table below for the results obtained from the study:

SUMMARY TABLE OF THE COMPARATIVE BIOAVAILABILITY DATA

Ondansetron (1 x 8 mg) From measured data Geometric LS Mean Arithmetic Mean (CV %)				
Parameter	Ondansetron* 8 mg Orally Disintegrating Tablet	Zofran® ODT† 8 mg Orally Disintegrating Tablet, USP, GlaxoSmithKline Inc., Canada	% Ratio of Geometric LS Means	90% Confidence Interval
AUC _{0-T} (ng·h/mL)	251.700 272.615 (41.7%)	274.714 299.283 (41.4%)	91.6	86.6 - 96.9
AUC _{0-∞} (ng·h/mL)	267.378 292.464 (45.2%)	293.295 322.736 (44.5%)	91.2	86.0 - 96.7
C _{max} (ng/mL)	36.783 38.877 (34.2%)	39.853 42.657 (36.7%)	92.3	87.6 - 97.2
T _{max} ‡ (h)	2.00 (1.50 - 3.03)	2.00 (1.25 - 5.00)		
T _{half} § (h)	6.09 (19.2%)	6.22 (21.7)		

* Athena-Ondansetron ODT 8 mg orally disintegrating tablets.

† Zofran® ODT 8 mg orally disintegrating tablets manufactured by GlaxoSmithKline Canada Inc.

‡ Expressed as the median (range) only.

§ Expressed as the arithmetic mean (CV%) only.

16 NON-CLINICAL TOXICOLOGY

Non-clinical pharmacodynamics

Ferret model

The ferret provides an excellent model for demonstrating the antiemetic action of drugs. Emesis can be induced by antineoplastic drugs or whole body irradiation. Behavioural changes associated with these treatments are noted in these animals and may also provide a parallel for the human experience of nausea.

CINV

The antiemetic action of ondansetron has been evaluated in both male and female ferrets given cisplatin (9-10 mg/kg), cyclophosphamide (200 mg/kg) or irradiation (2 and 8 Gy, 250 kV). Intravenous doses of ondansetron (0.1-1 mg/kg) abolished cisplatin-induced emesis for up to 2 hours. In cyclophosphamide-induced emesis, subcutaneous doses of 0.5 mg/kg ondansetron completely eliminated vomiting, significantly reduced retching and delayed the onset of these responses.

RINV

The radiation-induced emesis, 0.5 mg/kg ondansetron alone completely and rapidly eliminated retching and vomiting.

Dexamethasone

The antiemetic effects of ondansetron (0.1 mg/kg) in combination with dexamethasone (2-5 mg/kg) were potentiated in ferrets with cyclophosphamide-induced emesis, compared with ondansetron alone. Ondansetron with dexamethasone produced a significant reduction in retching (65%) and vomiting (72%).

Gastric emptying

Among its secondary effects, ondansetron has also been shown to cause a dose-dependent increase in the rate of gastric emptying in the guinea pig, which is significant at doses of 0.01-0.1 mg/kg. As gastric stasis is frequently associated with nausea, stimulation of gastric motility may be a beneficial action of ondansetron. In the cat, dog and monkey, ondansetron has little effect on heart rate, blood pressure or ECG at intravenous doses up to 3 mg/kg.

QT-prolongation

A study in cloned human cardiac ion channels has shown ondansetron has the potential to affect cardiac repolarisation via blockade of hERG potassium channels at clinically relevant concentrations. Dose-dependent QT prolongation has been observed in a thorough QT study in human volunteers (see [10.2 Pharmacodynamics – Electrocardiography](#)).

Non-clinical pharmacokinetics

In mice, rats, rabbits and dogs dosed at 1 mg/kg orally and/or intravenously, the plasma half-life of ondansetron was less than 1 hour, but the half-lives of its metabolites were significantly longer. Peak plasma concentrations of ondansetron in rats and dogs ranged from 351 to 419 ng/mL for the IV dose and 8 to 15 ng/mL for the oral dose.

Plasma levels were linear over a 30-fold dose range. In repeat-dose studies there was no apparent accumulation of ondansetron.

Ondansetron is almost completely absorbed in animals, and is rapidly metabolized by N-demethylation and hydroxylation of the indole ring, followed by conjugation with glucuronic acid and sulphate. There is significant first-pass metabolism after oral doses.

Ondansetron and its metabolites are rapidly and widely distributed in tissues, reaching higher levels than the corresponding plasma levels. In the rat and dog, ondansetron binds reversibly to tissues containing melanin and elastin. In rats and man, plasma protein binding is about 73%, while it is slightly lower in the dog (60%). Ondansetron and its metabolites cross the blood-brain barrier to only a slight extent.

General Toxicology:

Acute Toxicity

Single doses of ondansetron up to the LD₅₀ in mice and in rats were generally well tolerated. Reactions, including tremor and convulsive behaviour, occurred only at near lethal levels.

Table 6: Acute Toxicity

Species	LD ₅₀ (mg/kg)	
	Oral	IV
Mice	10-30	1.0-2.5
Rats	100-150	15-20

All deaths resulted from the acute effects of treatment, the observed clinical signs being consistent with the central nervous system effects associated with behavioural depression. These effects were not associated with any apparent histopathological changes in the brain. No target organ toxicity was identified.

Long term Toxicity

Table 7: Subacute Toxicity Studies

Species	Route	Dose (mg/kg/day)	Duration of Study	Results
Rats	Oral	160	7 weeks	Well tolerated
	IV	12	5 weeks	Well tolerated
Dogs	Oral	7.5 – 25	5 weeks	Transient post-dosing clinical reactions associated with behavioural depression (at highest dose levels)
	IV	2 - 8	5 weeks	

Maximum daily dose levels in rats were found to be higher when doses were gradually increased. Identical doses were rapidly lethal to rats not previously exposed to ondansetron. Post-dosing reactions, in both rats and dogs, included ataxia, exophthalmia, mydriasis, tremor and respiratory changes. Increases in liver enzymes (SGPT and SGOT) were noted at high dose levels. Dogs dosed at 6.75 mg/kg/day intravenously exhibited vein irritancy in the form of

constriction and thickening, creating resistance to needle penetration. The changes were noted after seven days treatment but were reversed by decreasing the dose concentration.

Table 8: Chronic Toxicity

Species	Duration	Max. no-effect Dose (mg/kg/day)	Effects
Rat	18 months	1	Usually transient and restricted to highest dose
Dog	12 months	12	

Carcinogenicity

Table 9: Carcinogenicity Studies

Species	Route	Dose (mg/kg/day)	Duration of Study	Results
Mice	Oral	1-40 (max. oral dose 30)	2 years	No treatment related increases in tumour incidence.
Rats	Oral	1-25 (max. oral dose 10)	2 years	Proportion of benign/malignant tumours also remained Consistent with the pathological background of the Animals studied.

There was no evidence of a tumourigenic effect of ondansetron in any tissue.

Mutagenicity

No evidence of mutagenicity was observed in microbial mutagen tests using mutant strains of *Salmonella typhimurium*, *Escherichia coli* or *Saccharomyces cerevisiae*, with or without a rat-liver post-mitochondrial metabolizing system.

There was also no evidence of damage to genetic material noted in in vitro V-79 mammalian cell mutation studies, in vitro chromosome aberration tests using human peripheral lymphocytes, or in vivo chromosome aberration assays in mouse bone marrow.

Reproductive and Developmental Toxicology

Ondansetron was not teratogenic in rats and rabbits at dosages up to the maximum non-convulsive level, (rat: 15 mg/kg/day, rabbit: 30 mg/kg/day; the maternal dose was approximately 6 and 24 times the maximum recommended human oral dose of 24 mg/day, respectively, based on body surface area). No adverse effects on pregnancy or foetal and post-natal development were detected in rats and no foetal abnormalities were observed in rabbits after oral administration of ondansetron.

A slight maternal toxicity was observed at the highest dose level in intravenous organogenesis (4.0 mg/kg/day) studies in the rabbit. Effects included maternal body weight loss and increased incidence of early foetal death. In a rat fertility study, there was a dose-related decrease in the proportion of surviving pups of the F2 generation; however, the significance of this is unclear.

Administration of ondansetron to pregnant rats and rabbits indicated there was foetal exposure to low levels of ondansetron and its metabolites. Ondansetron is retained in the foetal eye presumably bound to melanin. In rats, the transfer of ondansetron and its metabolites into breast milk was extensive. The concentration of unchanged ondansetron in breast milk was higher than in corresponding plasma samples.

Daily administration of ondansetron at dosages up to 15 mg/kg/day to pregnant rats (a maternal dose of approximately 6 times the maximum recommended human oral dose of 24 mg/day, based on body surface area) from day 17 of pregnancy to litter day 22 had no effects on pregnancy of the parental generation or on post-natal development and mating of the F1 generation. Foetal development of the F2 generation was comparable to controls; however, the number of implantations and viable foetuses was reduced in the highest dosage group when compared with controls.

17 SUPPORTING PRODUCT MONOGRAPHS

1. Zofran ODT (Oral Disintegrating Tablets), Control No.: 252778, Product Monograph, Novartis Pharmaceuticals Canada Inc., (November 9, 2021).

PATIENT MEDICATION INFORMATION

READ THIS FOR SAFE AND EFFECTIVE USE OF YOUR MEDICINE

PrATHENA-ONDANSETRON ODT

(Ondansetron Orally Disintegrating Tablets)

Read this carefully before you start taking **ATHENA-ONDANSETRON ODT** and each time you get a refill. This leaflet is a summary and will not tell you everything about this drug. Talk to your healthcare professional about your medical condition and treatment and ask if there is any new information about **ATHENA-ONDANSETRON ODT**.

What is ATHENA-ONDANSETRON ODT used for?

Children (4 to 17 years of age):

ATHENA-ONDANSETRON ODT (ondansetron) is used to prevent nausea and vomiting during chemotherapy.

Adults (18 to 64 years of age):

ATHENA-ONDANSETRON ODT is used:

- to prevent nausea and vomiting during chemotherapy and radiotherapy, and
- to prevent or treat nausea and vomiting after surgery.

Geriatrics (65 years of age and older):

ATHENA-ONDANSETRON ODT is used to prevent nausea and vomiting during chemotherapy and radiotherapy.

How does ATHENA-ONDANSETRON ODT work?

ATHENA-ONDANSETRON ODT is a medication known as an antiemetic. Treatments such as cancer chemotherapy and radiotherapy are associated with the release of a natural substance (serotonin). The release of serotonin, can make you feel sick and vomit. The way that ATHENA-ONDANSETRON ODT works is not known, but it is thought to help stop the effects of serotonin to reduce the effects of nausea and vomiting.

What are the ingredients in ATHENA-ONDANSETRON ODT?

- Medicinal ingredient: ondansetron.
- Non-medicinal ingredients: amino methacrylate copolymer, aspartame, colloidal silicon dioxide, magnesium stearate, mannitol, polyvinyl pyrrolidone and

strawberry flavour.

ATHENA-ONDANSETRON ODT is available in the following dosage form:

- Orally disintegrating tablets: 4 mg and 8 mg of ondansetron.

Do not use ATHENA-ONDANSETRON ODT if:

- you are allergic to ondansetron or to any other ingredients in ATHENA-ONDANSETRON ODT.
- you are taking a medicine called apomorphine (used to treat Parkinson's disease).

To help avoid side effects and ensure proper use, talk to your healthcare professional before you take ATHENA-ONDANSETRON ODT. Talk about any health conditions or problems you may have, including if you:

- have had an allergic reaction to medicines that are similar to ATHENA-ONDANSETRON ODT such as medicines containing granisetron or palonosetron.
- are pregnant or planning to become pregnant. ATHENA-ONDANSETRON ODT is not recommended for use during pregnancy.
- are breast feeding or planning to breastfeed. ATHENA-ONDANSETRON ODT can pass into your breast milk and affect your baby.
- have liver problems.
- have signs of intestinal obstruction or blockage.
- have or have had heart or blood vessel problems, including if you are at a higher risk for these problems. Risk factors include, but are not limited to, if you:
 - have family members who have or have had heart or blood vessel problems,
 - smoke,
 - have high blood pressure,
 - have high cholesterol levels,
 - have diabetes, or
 - are overweight.
- have a condition called phenylketonuria, as ATHENA-ONDANSETRON ODT contains aspartame.
- are taking medications that affect the serotonin in your body (e.g., serotonergic and neuroleptic medications). If you are unsure, ask your healthcare professional.
- have QT/QTc prolongation (a heart rhythm condition) or a family history of QT/QTc prolongation.
- are taking medications that may lead to QT/QTc prolongation or electrolyte imbalances. If you are unsure, ask your healthcare professional.
- have low blood levels of potassium, magnesium, or calcium.

Other warnings you should know about:

Serotonin syndrome: ATHENA-ONDANSETRON ODT can cause serotonin syndrome, a rare but potentially life-threatening condition. It can cause serious changes in how your brain, muscles and digestive system work. You may develop serotonin syndrome if you take ATHENA-ONDANSETRON ODT with certain anti-depressants or migraine medications.

Serotonin syndrome symptoms include:

- fever, sweating, shivering, diarrhea, nausea, vomiting;
- muscle shakes, jerks, twitches or stiffness, overactive reflexes, loss of coordination;
- fast heartbeat, changes in blood pressure;
- confusion, agitation, restlessness, hallucinations, mood changes, unconsciousness, and coma.

Myocardial ischemia (lack of blood flow to the heart): Treatment with ATHENA-ONDANSETRON ODT can cause myocardial ischemia which can lead to a heart attack. This may happen shortly after ATHENA-ONDANSETRON ODT administration. Some symptoms of myocardial ischemia can include sudden chest pain, pressure or discomfort, feeling faint, feeling anxious, shortness of breath, irregular heartbeat, nausea, and sudden heavy sweating. Your healthcare professional will monitor your health during and after administration of ATHENA-ONDANSETRON ODT. However, if you notice any symptoms of myocardial ischemia, tell your healthcare professional right away. They may reduce or stop your treatment, and may recommend another therapy.

QT/QTc prolongation: ATHENA-ONDANSETRON ODT can affect the electrical activity of your heart known as QT/QTc prolongation. This effect can be measured with an electrocardiogram (ECG). In rare cases, QT/QTc prolongation can cause changes to the rhythm of your heart (e.g., fast, slow or irregular heartbeats). This can lead to dizziness, palpitations (sensation of rapid, pounding, or irregular heart beat), fainting, or death. You are at a higher risk if you have a heart disease, are taking certain interacting medicines, are a female, or are over the age of 65 years. It is important to follow the instructions of your healthcare professional with regard to dosing or any special tests. If you experience any symptoms of a possible heart rhythm problem, you should seek immediate medical attention.

Severe allergic reactions: ATHENA-ONDANSETRON ODT can cause allergic reactions in certain individuals. Symptoms of a severe allergic reaction can include wheezing, sudden chest pain, tightness of the chest, heart throbbing, swelling of eyelids, face or lips, or develop a skin rash, skin lumps or hives. If you notice any signs of a severe allergic reaction, **contact your healthcare professional immediately. Do not take any more medicine unless your healthcare professional tells you to do so.**

Pregnancy:

- If you are pregnant, there are specific risks for your unborn baby that you must discuss with your healthcare professional.
- If you are able to get pregnant, you may be asked to take a pregnancy test before starting your treatment with ATHENA-ONDANSETRON ODT.
- You should use effective birth control while you are taking ATHENA-ONDANSETRON ODT, and for at least 2 days after stopping ATHENA-ONDANSETRON ODT. Ask your healthcare professional about options of effective birth control.
- If you become pregnant while taking ATHENA-ONDANSETRON ODT, tell your healthcare professional right away.

Tell your healthcare professional about all the medicines you take, including any drugs, vitamins, minerals, natural supplements or alternative medicines.

The following may interact with ATHENA-ONDANSETRON ODT:

- medicines called CYP3A4 inducers (e.g., phenytoin, carbamazepine, and rifampicin);
- medicines used to treat heart rhythm disorders (e.g., quinidine, procainamide, disopyramide, amiodarone, sotalol, ibutilide, dronedarone, flecainide, and propafenone);
- medicines used to treat vomiting and nausea called antiemetics (e.g., dolasetron, palonosetron, granisetron, droperidol, chlorpromazine, prochlorperazine, and domperidone);
- medicines called tyrosine kinase inhibitors (e.g., vandetanib, sunitinib, nilotinib, and lapatinib);
- medicines used to manage psychosis or schizophrenia called antipsychotics (e.g., chlorpromazine, pimozide, haloperidol, and ziprasidone);
- medicines used to treat depression called antidepressants (e.g., citalopram, fluoxetine, venlafaxine, tricyclic/tetracyclic antidepressants, amitriptyline, imipramine, and maprotiline);
- medicines used to treat pain called opioids (e.g., methadone and tramadol);
- medicines used to treat bacterial infections called antibiotics (e.g., erythromycin, clarithromycin, telithromycin, tacrolimus, moxifloxacin, levofloxacin, and ciprofloxacin);
- medicines used to treat malaria called antimalarials (e.g., quinine and chloroquine);
- medicines used to treat fungal infections called azole antifungals (e.g., ketoconazole, fluconazole, and voriconazole);
- medicines used to treat cancer (e.g., vorinostat);
- medicines called beta-2 adrenoceptor agonists (e.g., salmeterol and formoterol);
- medicines that can affect electrolyte levels (e.g., diuretics, laxatives, enemas,

- amphotericin B, and high doses of corticosteroids);
- a medicine used to treat Parkinson’s Disease called apomorphine;
 - medicines called serotonergic drugs that can affect the serotonin in the body (e.g., triptans, Selective Serotonin-Reuptake Inhibitors (SSRIs), Serotonin Noradrenalin Reuptake Inhibitors (SNRIs), lithium, sibutramine, fentanyl and its analogues, dextromethorphan, tramadol, tapentadol, meperidine, methadone, pertazocine, St. John’s Wort (*Hypericum perforatum*), monoamine oxidase inhibitors (MAOIs), linezolid, and methylene blue).

If you are unsure about any medications you are taking, ask your healthcare professional.

How to take ATHENA-ONDANSETRON ODT:

- The label on the container of your medicine should tell you how often to take your medicine and how many doses you should take each time. If not, or if you are not sure, consult your healthcare professional.
- **Do not** take more doses, or take them more often than your healthcare professional prescribes. If, however, you vomit within one hour of taking your medicine, you should take the same amount of medicine again. If vomiting persists, consult your healthcare professional.

To take ATHENA-ONDANSETRON ODT orally disintegrating tablets:

1. Tear along the perforations of the foil to separate off one tablet within its blister unit.
2. Gently push the ATHENA-ONDANSETRON ODT out of the blister aluminium foil and remove it with dry fingers.
3. Immediately place ATHENA-ONDANSETRON ODT on top of the tongue. Allow it to melt. Apply gentle pressure if required. Swallow the pasty mass formed.

Usual dose:

Take ATHENA-ONDANSETRON ODT exactly as your healthcare professional has told you. Your healthcare professional will determine the right dose and length of ATHENA-ONDANSETRON ODT for you. Your dose will depend on your medical condition, age, current health, and if you take certain other medications. Your healthcare professional may monitor your health throughout your treatment and may interrupt, reduce or stop your dose.

Overdose:

If you think you, or a person you are caring for, have taken too much ATHENA-ONDANSETRON ODT, contact a healthcare professional, hospital emergency department, or regional poison control centre immediately, even if there are no symptoms.

Missed Dose:

If you miss a dose and do not feel sick, take the next dose when it is due.

If you forget to take your medicine and feel sick or vomit, take a dose as soon as possible.

If your healthcare professional decides to stop the treatment, do not keep any leftover medicine unless your healthcare professional tells you to.

What are possible side effects from using ATHENA-ONDANSETRON ODT?

These are not all the possible side effects you may have when taking ATHENA-ONDANSETRON ODT. If you experience any side effects not listed here, tell your healthcare professional.

Some side effects may include:

- feeling of flushing or warmth;
- hiccups.

There is no need to stop taking your medicine, but you should tell your healthcare professional about these symptoms at your next visit.

If you feel unwell or have any symptoms that you do not understand, you should contact your healthcare professional immediately.

Serious side effects and what to do about them			
Symptom / effect	Talk to your healthcare professional		Stop taking drug and get immediate medical help
	Only if severe	In all cases	
UNCOMMON			
Heart problems (disorders affecting your heart muscle, valves or rhythm): chest pain, chest discomfort, high blood pressure, irregular heart rhythm, shortness of breath, or fainting.			✓
Seizures: loss of consciousness with uncontrollable shaking, visual disturbances (e.g., blurred vision).			✓

Serious side effects and what to do about them			
Symptom / effect	Talk to your healthcare professional		Stop taking drug and get immediate medical help
	Only if severe	In all cases	
Movement disorders (including dyskinesia): loss of coordination or balance, speech or limb movements, muscle spasms, difficulty walking, tremor, upward rolling of the eyes, or abnormal muscular stiffness.			✓
RARE			
Eye problems such as blurred vision		✓	
Immediate severe allergic reaction: swelling of the mouth, throat, difficulty in breathing, rash, hives, or increased heart rate.			✓
Serotonin syndrome: a reaction which may cause feelings of agitation or restlessness, flushing, muscle twitching, involuntary eye movements, heavy sweating, high body temperature (> 38°C), or rigid muscles.			✓
Liver problems: yellowing of your skin and eyes (jaundice), unusual dark urine and pale stools, pain or swelling in the right upper abdomen, unusual tiredness, nausea, or vomiting.			✓
Hypokalemia (low level of potassium in the blood): muscle weakness, muscle spasms, cramping, constipation, feeling of skipped heart beats or palpitations, fatigue, tingling, or numbness			✓

Serious side effects and what to do about them			
Symptom / effect	Talk to your healthcare professional		Stop taking drug and get immediate medical help
	Only if severe	In all cases	
Prolongation of QT interval (a heart rhythm condition): irregular heartbeat, palpitations, dizziness, fainting, loss of consciousness, or seizures.			✓
Hypotension (low blood pressure): dizziness, fainting, light-headedness, blurred vision, nausea, or vomiting.			✓
Myocardial ischemia (lack of blood flow to the heart which can lead to heart attack): sudden chest pain, pressure or discomfort, feeling faint, feeling anxious, shortness of breath, irregular heartbeat, nausea, or sudden heavy sweating.			✓
VERY RARE			
Eye problems such as temporary blindness		✓	
Stevens-Johnson Syndrome (SJS) and Toxic Epidermal Necrolysis (TEN) (severe skin reactions): redness, blistering or peeling of the skin and/or inside of the lips, eyes, mouth, nasal passages or genitals, fever, chills, headache, cough, body aches, or swollen glands.			✓

If you have a troublesome symptom or side effect that is not listed here or becomes bad enough to interfere with your daily activities, tell your healthcare professional.

Reporting Side Effects

You can report any suspected side effects associated with the use of health products to Health Canada by:

- Visiting the Web page on Adverse Reaction Reporting (<https://www.canada.ca/en/health-canada/services/drugs-health-products/medeffect-canada/adverse-reaction-reporting.html>) for information on how to report online, by mail or by fax; or
- Calling toll-free at 1-866-234-2345.

NOTE: Contact your healthcare professional if you need information about how to manage your side effects. The Canada Vigilance Program does not provide medical advice.

Storage:

ATHENA-ONDANSETRON ODT orally disintegrating tablets should be stored at room temperature between 15 - 30°C.

If you want more information about ATHENA-ONDANSETRON ODT:

- Talk to your healthcare professional.
- Find the full Product Monograph that is prepared for healthcare professionals and includes this Patient Medication Information by visiting the Health Canada website (<https://www.canada.ca/en/health-canada/services/drugs-health-products/drug-products/drug-product-database.html>); the manufacturer's website www.athena.ca or by calling toll-free 1.888.620.5236.

This leaflet was prepared by Athena Pharmaceutiques SAS.

Last revised: JAN 17, 2022