

PRODUCT MONOGRAPH  
INCLUDING PATIENT MEDICATION INFORMATION

Pr **SOHONOS**<sup>™</sup>

palovarotene capsules

1 mg, 1.5 mg, 2.5 mg, 5 mg, 10 mg

Retinoid/Retinoic acid receptor gamma (RAR $\gamma$ ) selective agonist

ATC code: M09AX11

Ipsen Biopharmaceuticals Canada Inc.  
5050 Satellite Drive, Suite 500  
Mississauga ON  
L4W 0G1  
[www.ipsen.ca](http://www.ipsen.ca)

Date of Initial Authorization:

Date of Revision:  
JAN 20, 2022

Submission Control Number: 252065

SOHONOS is a trademark of Clementia Pharmaceuticals Inc. ©2021. All rights reserved.

## TABLE OF CONTENTS

Sections or subsections that are not applicable at the time of authorization are not listed.

<b>TABLE OF CONTENTS</b> .....	<b>2</b>
<b>PART I: HEALTH PROFESSIONAL INFORMATION</b> .....	<b>4</b>
<b>1 INDICATIONS</b> .....	<b>4</b>
1.1 Pediatrics .....	4
1.2 Geriatrics.....	4
<b>2 CONTRAINDICATIONS</b> .....	<b>4</b>
<b>3 SERIOUS WARNINGS AND PRECAUTIONS BOX</b> .....	<b>4</b>
<b>4 DOSAGE AND ADMINISTRATION</b> .....	<b>5</b>
4.1 Dosing Considerations.....	5
4.2 Recommended Dose and Dosage Adjustment .....	5
4.4 Administration.....	7
4.5 Missed Dose.....	7
<b>5 OVERDOSAGE</b> .....	<b>7</b>
<b>6 DOSAGE FORMS, STRENGTHS, COMPOSITION AND PACKAGING</b> .....	<b>7</b>
<b>7 WARNINGS AND PRECAUTIONS</b> .....	<b>8</b>
7.1 Special Populations.....	12
7.1.1 Pregnant Women .....	12
7.1.2 Breast-feeding.....	13
7.1.3 Pediatrics.....	13
7.1.4 Geriatrics.....	13
<b>8 ADVERSE REACTIONS</b> .....	<b>13</b>
8.1 Adverse Reaction Overview.....	13
8.2 Clinical Trial Adverse Reactions.....	14
8.2.1 Clinical Trial Adverse Reactions – Pediatrics .....	16
8.3 Less Common Clinical Trial Adverse Reactions with SOHONOS (<5%).....	16
8.4 Abnormal Laboratory Findings: Hematologic, Clinical Chemistry and Other Quantitative Data .....	16
<b>9 DRUG INTERACTIONS</b> .....	<b>18</b>
9.2 Drug Interactions Overview.....	18
9.3 Drug-Behavioural Interactions.....	19
9.4 Drug-Drug Interactions.....	19

9.5	Drug-Food Interactions .....	20
9.6	Drug-Herb Interactions.....	20
9.7	Drug-Laboratory Test Interactions.....	20
<b>10</b>	<b>CLINICAL PHARMACOLOGY.....</b>	<b>20</b>
10.1	Mechanism of Action.....	20
10.2	Pharmacodynamics.....	21
10.3	Pharmacokinetics.....	21
<b>11</b>	<b>STORAGE, STABILITY AND DISPOSAL .....</b>	<b>23</b>
<b>12</b>	<b>SPECIAL HANDLING INSTRUCTIONS .....</b>	<b>23</b>
<b>PART II: SCIENTIFIC INFORMATION .....</b>		<b>24</b>
<b>13</b>	<b>PHARMACEUTICAL INFORMATION .....</b>	<b>24</b>
<b>14</b>	<b>CLINICAL TRIALS .....</b>	<b>25</b>
14.1	Clinical Trials by Indication.....	25
<b>15</b>	<b>MICROBIOLOGY.....</b>	<b>30</b>
<b>16</b>	<b>NON-CLINICAL TOXICOLOGY .....</b>	<b>31</b>
<b>PATIENT MEDICATION INFORMATION .....</b>		<b>33</b>

## PART I: HEALTH PROFESSIONAL INFORMATION

### 1 INDICATIONS

SOHONOS (palovarotene capsules) is indicated to reduce the formation of heterotopic ossification in adults and children aged 8 years and above for females and 10 years and above for males with Fibrodysplasia (myositis) Ossificans Progressiva.

#### 1.1 Pediatrics

**Pediatrics (<8 years (females)/<10 years (males)):** Do not use SOHONOS in females below 8 years and males below 10 years. Based on the data submitted and reviewed by Health Canada, the safety and efficacy of SOHONOS in these pediatric patients has not been established. Therefore, Health Canada has not authorized an indication for pediatric use in females below 8 years and males below 10 years (see [7.1.3 Pediatrics](#)).

**Pediatrics (≥8 years (females)/10 years (males)):** Based on the data submitted and reviewed by Health Canada, the safety and efficacy of SOHONOS in pediatric patients has been established. Therefore, Health Canada has authorized an indication for pediatric use (see [7.1.3 Pediatrics](#)).

#### 1.2 Geriatrics

**Geriatrics (≥ 65 years):** Clinical studies of SOHONOS did not include sufficient numbers of subjects aged 65 and over to determine whether they respond differently from younger subjects. There were no clinically relevant differences in SOHONOS pharmacokinetics between young males and elderly males (see [7.1.4 Geriatrics](#)).

### 2 CONTRAINDICATIONS

- Patients who are pregnant or breastfeeding.
- Patients of childbearing potential unless all the conditions for pregnancy prevention are met, or they are not at risk of pregnancy due to physical limitations as assessed by the physician.
- Patients with a history of allergy or hypersensitivity to retinoids, or to any component of this product. For a complete listing, see [6 DOSAGE FORMS, STRENGTHS, COMPOSITION AND PACKAGING](#).

### 3 SERIOUS WARNINGS AND PRECAUTIONS BOX

#### Serious Warnings and Precautions

- SOHONOS is a member of the retinoid class of drugs that is associated with birth defects in humans. SOHONOS must not be used by patients who are, or intend to become, pregnant due to the risk of teratogenicity. To minimize fetal exposure, SOHONOS is to be administered only if all conditions for pregnancy prevention are met (see [7 WARNINGS AND PRECAUTIONS, Reproductive Health: Female and Male Potential, Teratogenic Risk](#) and [Special Populations: Pregnant Women](#)).
- SOHONOS has been shown to cause premature physal closure in growing children with FOP; periodic monitoring (every 3 months) is recommended (see [7 WARNINGS AND PRECAUTIONS, Musculoskeletal, Monitoring Recommendations](#)).

## 4 DOSAGE AND ADMINISTRATION

### 4.1 Dosing Considerations

Pregnancy testing and contraceptive measures must be followed prior to dosing SOHONOS in patients of childbearing potential.

Patients who are pregnant, or who intend to become pregnant, should avoid contact with SOHONOS. If pregnancy does occur during treatment with SOHONOS or for one month after its discontinuation, SOHONOS treatment must be immediately stopped if not already discontinued and patient should discuss with their physician.

To avoid unintended exposure, if emptying the capsule contents onto soft food, caregivers administering SOHONOS should wear disposable gloves when handling, and use disposable paper towels and a container to collect waste (e.g. a resealable bag).

### 4.2 Recommended Dose and Dosage Adjustment

#### Chronic Regimen

Recommended dosing: 5 mg once daily (chronic treatment)

Chronic treatment should stop at the time of initiation of flare-up treatment, and re-initiated after completion of the flare-up treatment.

Weight-adjusted dosage is required in children who are under 14 years of age (see [Table 1](#)).

#### Flare-up Regimen

Recommended Dosing: 20 mg once daily for 4 weeks, followed by 10 mg once daily for 8 weeks for a total of 12 weeks (20/10 mg flare-up treatment) even if symptoms resolve earlier.

In the presence of persistent flare-up symptoms, treatment may be extended in 4-week intervals with 10 mg SOHONOS.

Should the patient experience another flare-up (new flare-up location or marked worsening of the original flare-up) at any time during flare-up treatment, the aforementioned flare-up 12-week treatment protocol should be restarted.

Treatment should begin at the onset of the first symptom indicative of an FOP flare-up, or substantial high-risk traumatic event likely to lead to a flare-up, under the guidance of a health care professional.

Weight-adjusted dosage is required in children who are under 14 years of age (see [Table 1](#)).

#### Dose Adjustment in Children Under 14 Years of Age

SOHONOS dosage is weight-adjusted in patients under 14 years of age (see [Table 1](#)). The health care professional should prescribe the most appropriate dosage based on weight for children aged from 8 years (females) and 10 years (males) to less than 14 years (see [Musculoskeletal](#) and [14 CLINICAL TRIALS](#)).

**Table 1. Weight-Adjusted Dosage for Children ≥ 8 Years (Females)/10 Years (Males) but <14 Years**

<b>Weight</b>	<b>Chronic Dosage</b>	<b>Flare-up Dosage (Weeks 1 to 4)</b>	<b>Flare-up Dosage (Weeks 5 to 12)</b>
≥60 kg*	5 mg	20 mg	10 mg
40 to <60 kg	4 mg	15 mg	7.5 mg
20 to <40 kg	3 mg	12.5 mg	6 mg
10 to <20 kg	2.5 mg	10 mg	5 mg

\*All children ≥14 years of age and adults should receive the dose in the ≥60 kg weight category

### **Dose Modification for Adverse Reactions**

If the patient experiences intolerable adverse reactions during either the chronic or the flare-up (Weeks 1-12) SOHONOS treatment, the daily dose should be reduced in a stepwise way at the discretion of the physician (see [Table 2](#) for the chronic dosing and [Table 3](#) and [Table 4](#) for the flare-up dosing). If the chronic dose reduction is not tolerated, consider discontinuation of the chronic dosing and only treating flare-ups. Additional dose reduction should occur if adverse reactions continue to be intolerable. If the patient is already receiving the lowest dose, then consideration should be given to discontinue therapy, temporarily or permanently, or switch to flare-up treatment only (see [Flare-up only regimen](#)). Subsequent flare-up treatment should be initiated at the same reduced treatment dose that was tolerated previously.

**Table 2. Dose Reduction: Chronic Dosing**

<b>Weight</b>	<b>Chronic Dosage</b>
≥60 kg*	2.5 mg
40 to <60 kg	2 mg
20 to <40 kg	1.5 mg
10 to <20 kg	1 mg

\*All children ≥14 years of age and adults should receive the dose in the ≥60 kg weight category

**Table 3. Dose Reduction: Flare-up Dosing Step 1**

<b>Weight</b>	<b>Flare-up Dosage (Weeks 1 to 4)</b>	<b>Flare-up Dosage (Weeks 5 to 12)</b>
≥60 kg*	15 mg	7.5 mg
40 to <60 kg	12.5 mg	5 mg
20 to <40 kg	10 mg	4 mg
10 to <20 kg	7.5 mg	3 mg

\*All children ≥14 years of age and adults should receive the dose in the ≥60 kg weight category

**Table 4. Dose Reduction: Flare-up Dosing Step 2**

Weight	Flare-up Dosage (Weeks 1 to 4)	Flare-up Dosage (Weeks 5 to 12)
≥60 kg*	10 mg	5 mg
40 to <60 kg	7.5 mg	4 mg
20 to <40 kg	6 mg	3 mg
10 to <20 kg	5 mg	2.5 mg

\*All children ≥14 years of age and adults should receive the dose in the ≥60 kg weight category

#### Flare-up only regimen:

If the patient experiences intolerable adverse reactions while taking chronic daily treatment and dose reduction does not alleviate the adverse reactions, then the patient may take SOHONOS only at the time of flare-up (or substantial high-risk traumatic event).

#### 4.4 Administration

SOHONOS should be taken with food, preferably at the same time each day. SOHONOS may be swallowed whole, or capsules may be opened and the contents emptied onto a teaspoon of soft food such as applesauce, yogurt, pudding, milk, oatmeal, rice cereal or liquid nutritional supplement, and taken immediately. If not taken immediately, it can be taken after a maximum of 1 hour after the sprinkling, provided it was maintained at room temperature and not exposed to direct sunlight.

#### 4.5 Missed Dose

If a dose of medication is missed by less than 6 hours, patients should take the missed dose as soon as possible. If the dose has been missed by more than 6 hours, the patient should skip the missed dose and continue with the next scheduled dose. The patient should not take two doses at the same time or in the same day.

### 5 OVERDOSAGE

No clinical experience with an overdose of SOHONOS has been reported. A single supratherapeutic dose of 50 mg SOHONOS had no apparent effects on vital signs, ECGs, or clinical laboratory parameters. Any overdose should be treated with supportive care according to the signs and symptoms exhibited by the patient.

For management of a suspected drug overdose, contact your regional poison control centre.

### 6 DOSAGE FORMS, STRENGTHS, COMPOSITION AND PACKAGING

**Table 5. Dosage Forms, Strengths, Composition and Packaging**

Route of Administration	Dosage Form / Strength/Composition	Non-medicinal Ingredients
Oral	Capsule 1 mg, 1.5 mg, 2.5 mg, 5 mg, 10 mg palovarotene	Croscarmellose sodium, gelatin, lactose monohydrate, magnesium stearate, microcrystalline cellulose, pharmaceutical grade printing ink, povidone, sodium lauryl sulfate, and titanium dioxide.

Capsules are packaged into a blister composed of PVC/PCTFE (polyvinylchloride/poly-chloro-tri-fluoro-ethylene) backed with push-through aluminium foil. The blisters are then packaged into a carton.

1 mg hard-gelatin capsules are white, opaque, size “0”, elongated, printed with black ink “PVO 1” on the body, containing white to off-white powder; available in cartons of 28 capsules (2 x 14 blister strips).

1.5 mg hard-gelatin capsules are white, opaque, size “0”, elongated, printed with black ink “PVO 1.5” on the body, containing white to off-white powder; available in cartons of 28 capsules (2 x 14 blister strips).

2.5 mg hard-gelatin capsules are white, opaque, size “0”, elongated, printed with black ink “PVO 2.5” on the body, containing white to off-white powder; available in cartons of 28 capsules (2 x 14 blister strips).

5 mg hard-gelatin capsules are white, opaque, size “0”, elongated, printed with black ink “PVO 5” on the body, containing white to off-white powder; available in cartons of 28 capsules (2 x 14 blister strips).

10 mg hard-gelatin capsules are white, opaque, size “0”, elongated, printed with black ink “PVO 10” on the body, containing white to off-white powder; available in cartons of 28 capsules (2 x 14 blister strips).

## 7 WARNINGS AND PRECAUTIONS

Please see **3 SERIOUS WARNINGS AND PRECAUTIONS BOX**.

### General

#### Conditions of Use

SOHONOS is contraindicated in patients of childbearing potential unless all of the following conditions for pregnancy prevention are met, or they are not at risk for pregnancy (e.g. patient has undergone a hysterectomy, bilateral oophorectomy, bilateral tubal ligation or has been medically confirmed to be postmenopausal).

#### Patients of Childbearing Potential

- The potential for pregnancy must be assessed for all patients.
- The patient must understand the teratogenic risk and the need to rapidly consult their physician if there is a risk of pregnancy or if they might be pregnant.
- Patients of child-bearing potential must use at least one highly effective method of contraception (i.e. IUD) or two effective methods (i.e. combined hormonal contraception in combination with another method of contraception such as a barrier method) during treatment with SOHONOS.
- The patient must understand and accept to undergo regular pregnancy testing before treatment with SOHONOS, during treatment and 1 month after stopping treatment.
- The patient must understand and accept the need for effective contraception, without interruption, 1 month before starting treatment, throughout the entire duration of treatment and for 1 month after the end of treatment.

- For those patients taking the flare-up regimen only, patients must continue to use effective contraception even during periods when SOHONOS is not being taken as the timing of flare-ups may not be predictable.
- The patient acknowledges that they understand the hazards and necessary precautions associated with the use of SOHONOS.
- The patient is informed and understands the potential consequences of pregnancy and the need to rapidly consult their physician if there is a risk of pregnancy or if they might be pregnant.

These conditions also concern patients of childbearing potential who are not currently sexually active unless the prescriber attests that there are compelling reasons to indicate that there is no risk of pregnancy (see [2 CONTRAINDICATIONS](#)).

### **Driving and Operating Machinery**

SOHONOS may influence the ability to drive and use machines (see [Ophthalmic, Nyctalopia](#)).

### **Monitoring and Laboratory Tests**

#### **Pregnancy Testing**

Medically documented blood or urine pregnancy tests are recommended to be performed, as follows:

#### *Prior to Starting Therapy*

At least one month after the patient has started using contraception, and shortly (preferably a few days) prior to the first prescription, the patient should undergo a pregnancy test. This test should ensure the patient is not pregnant when starting treatment with SOHONOS.

#### *Follow-up Pregnancy Testing During Treatment*

Monthly pregnancy testing is recommended.

#### *End of Treatment*

One month after stopping treatment, patients should undergo a final pregnancy test.

#### **Growth**

Periodic monitoring every 3 months of physcal growth plates is recommended in growing children (see [Musculoskeletal](#)).

### **Mucocutaneous**

Mucocutaneous effects are the most commonly reported adverse reactions across all doses during clinical studies with SOHONOS (98%) and were generally mild to moderate in severity; 4.3% of the mucocutaneous adverse reactions were severe. Mucocutaneous adverse reactions observed in over 10% of subjects were: dry skin (78%), dry lips (55%), alopecia (41%), pruritis (40%), erythema (32%), rash (30%), skin exfoliation [skin peeling] (31%), rash (30%), drug eruption (17%), eczema (14%) and skin irritation (12%). SOHONOS may contribute to an increased risk of skin and soft tissue infections, particularly paronychia (14%) and decubitus ulcer (6%), due to a decreased skin barrier from mucocutaneous effects such as dry and

peeling skin. One subject in a pivotal study developed perianal bleeding from dry perianal skin. The event was not felt to be due to a gastrointestinal bleeding source.

Some of these mucocutaneous adverse reactions led to dose reductions which occurred more frequently during flare-up dosing (35%) than during chronic treatment (3%), suggesting a dose response relationship. Prophylactic measures to minimize risk and/or treat the mucocutaneous effects are recommended (e.g. skin emollients, sunscreen, lip moisturizers, artificial tears, or other helpful treatments). Dose reductions may be required (see [Recommended Dose and Dosage Adjustment](#)).

### Photosensitivity

Photosensitivity reactions, such as exaggerated sunburn reactions (e.g. burning, erythema, blistering) involving areas exposed to the sun have been associated with the use of retinoids. Although SOHONOS was negative for phototoxicity when tested in vitro, precautionary measures for phototoxicity are still recommended. Excessive exposure to sun or artificial ultraviolet light should be avoided, and protection from sunlight should be used when exposure cannot be avoided (ie: use of sunscreens, wearing protective clothing and use of sunglasses).

## **Musculoskeletal**

### Bone Safety

Premature physeal closure (PPC) has been demonstrated to be an important risk associated with SOHONOS treatment in growing children. In clinical studies, PPC can be irreversible. Twenty-four (24%) SOHONOS treated pediatric subjects <18 years of age reported PPCs, comprising 10 of 39 subjects from ≥8/10 to <14 years (26%) and 14 of 25 subjects <8/10 years (56%). Overall, 8 subjects had severe PPC; of these, 5 subjects were younger children ≤7 years of age. SOHONOS is not indicated in children aged below 8 years for females and below 10 years for males.

Potential long-term consequences of PPC include growth arrest, leg length discrepancy, disproportionate growth (epiphyseal growth plate closure preferentially affecting the lower extremities), angular deformity in affected joints, and gait disturbance. Consistent with the retinoid literature, all but one of the PPC events was observed first in the knee. Additionally, femur and tibia length growth also demonstrated symmetric growth although differences leading to leg length discrepancy would likely take a longer amount of time to manifest. Should asymmetric physeal fusion occur, angular deformity would likely take a longer amount of time to manifest. Given the relatively short follow-up times to date of SOHONOS, long-term consequences on angular deformity have not been established in subjects treated with SOHONOS. Upon initiation of treatment with SOHONOS, all growing children should be considered to be at risk of developing PPC and the potential long-term consequences (see [8.2.1 Clinical Trial Adverse Reactions – Pediatrics](#)).

### *Monitoring Recommendations:*

Periodic monitoring every 3 months of physeal growth plates is recommended in growing children. Should evidence of adverse effects on growth and/or PPC be observed, further evaluation and increased monitoring may be required. The decision to temporarily interrupt SOHONOS during the evaluation period or permanently discontinue should be made based on individual benefit-risk determination.

There are no clear characteristics that define or predict who will develop PPC, over what period, or after what duration of SOHONOS exposure. Therefore, prior to starting treatment with SOHONOS, it is recommended that all growing children undergo baseline clinical and

radiological assessments including, but not limited to, an assessment of skeletal maturity via hand/wrist and knee x-rays, standard growth curves, and pubertal staging. Continued monitoring of linear growth and skeletal maturity via x-ray is recommended every 3 months until patients reach skeletal maturity or final adult height (frequency of monitoring may depend on patient characteristics such as age, pubertal status and skeletal maturity). Once the patients have reached skeletal maturity or the patient has reached final adult height, no further monitoring for PPC is necessary.

Should a patient exhibit signs of PPC based on clinical and radiologic evaluations, an assessment of the benefits and risks should be made in order to determine the appropriateness of continued treatment versus temporary or permanent discontinuation of SOHONOS, until the patient reaches skeletal maturity.

## **Ophthalmic**

### *Xerophthalmia/Conjunctivitis*

25% of subjects experienced xerophthalmia. Subjects were treated with saline or lubricating eye drops and symptoms improved.

### *Nyctalopia*

Night blindness (nyctalopia) has been identified as a potentially dangerous effect associated with systemic retinoids. This may be dose-dependent, making driving a vehicle at night potentially hazardous during treatment. Night blindness is generally reversible after cessation of treatment but can also persist in some cases. A single mild adverse event of night blindness in an FOP subject receiving SOHONOS reported in the clinical trial resolved after 134 days without treatment discontinuation or dose de-escalation. (see [Driving and Operating Machinery](#)).

## **Psychiatric**

Depression, depression aggravated, anxiety, mood alterations and suicidal thoughts and behaviours have been reported in patients treated with systemic retinoids and individuals with a personal history of psychiatric illness appear to be more susceptible. There is a relatively high background prevalence (24%) of depression in untreated patients with FOP with 9% of patients having a medical history of depression when enrolled in SOHONOS clinical trials.

In FOP clinical trials, there was no treatment-related increase in suicide ideation, suicidal behaviour or psychiatric disorders overall relative to untreated FOP subjects. Particular care should be taken in patients with a history of psychiatric illness. Patients should be monitored for signs of depression and referred for appropriate treatment if necessary.

## **Reproductive Health: Female and Male Potential (see [2 CONTRAINDICATIONS](#) and [7.1.1 Pregnant Women](#))**

- **Fertility**

### *Females*

A fertility study conducted on female rats revealed no effects of SOHONOS at up to 1mg/kg/day on reproductive function, or fertility and early embryonic development. At 3mg/kg/day lower numbers of implantation sites and live embryos were observed. These effects were considered secondary to decreased food intake and not a primary effect of SOHONOS. There were no findings in female reproductive organs in rats or dogs in chronic toxicity studies (see [16 NON-CLINICAL TOXICOLOGY](#)).

## Males

A fertility study conducted on male rats revealed no effects of SOHONOS at up to 1 mg/kg/day on reproductive function, fertility, or early embryonic development. However, some testicular toxicity was observed in rats at 5 mg/kg/day for 4 weeks, a dose that produced systemic toxicity and deaths. No evidence of testicular toxicity was observed in rats or dogs in chronic toxicity studies (see [16 NON-CLINICAL TOXICOLOGY](#)).

- **Teratogenic Risk**

The teratogenic potential of systemic retinoids is well established. Studies in pregnant rats have shown that administration of SOHONOS resulted in fetal malformations typical of retinoids (i.e. cleft palate, misshapen skull bones, short long bones). There have been no reports of pregnancy or *in utero* exposure reported in clinical studies with SOHONOS (see [Reproductive and Developmental Toxicology](#)).

### Retinoid Class Effects: Pregnancy and Teratogenicity

- The class of retinoids are known to cause severe birth defects in a very high percentage of infants born to patients who became pregnant during treatment with retinoids in any amount, even for a short period of time.
- Birth defects which have been documented following exposure to other retinoids, including: CNS (hydrocephalus, hydranencephaly, microcephaly, posterior fossa abnormalities, cranial nerve dysfunction, cerebellar malformation); craniofacial (anotia, microtia, low set ears, small or absent external auditory canals, microphthalmia, facial dysmorphism, cleft palate); cardiac (septal defects, aortic arch abnormalities, tetralogy of Fallot); thymus gland abnormalities; and parathyroid hormone deficiency. Cases of IQ scores less than 85 with or without other abnormalities have been reported in patients taking retinoids.
- Potentially any exposed fetus can be affected. There are no accurate means of determining whether an exposed fetus has been affected.
- SOHONOS should only be prescribed by physicians knowledgeable in the [SOHONOS Pregnancy Prevention Plan](#).

## 7.1 Special Populations

### 7.1.1 Pregnant Women

SOHONOS is contraindicated during pregnancy and in patients of childbearing potential who are sexually active and not using contraception because SOHONOS can cause fetal harm when administered to a pregnant patient. In animal reproduction studies, SOHONOS induced fetal malformations typical of retinoids when orally administered to pregnant rats during the period of organogenesis at a dose of  $\geq 0.25$  mg/kg/day. There are no available clinical data on SOHONOS use in pregnant patients to inform drug-associated risks. If pregnancy occurs in a patient treated with SOHONOS, treatment must be stopped, and the patient should be counselled to refer to their physician for evaluation and advice. If pregnancy occurs within 1 month of treatment discontinuation, there remains a risk of severe and serious malformation of the fetus. The patient should be counselled to refer to her physician (see [2 CONTRAINDICATIONS](#) and [Reproductive and Developmental Toxicology](#)).

A registry has been established to collect information about the effect of SOHONOS exposure, including exposure during pregnancy. The registry is a prospective observational study actively

collecting information on SOHONOS exposure, including during pregnancy and associated pregnancy outcomes. Physicians are encouraged to enroll patients in the SOHONOS registry, or for more information regarding the registry, by calling 1-855-215-2288. If the patient does not wish to participate in the registry, physicians and patients are still encouraged to report pregnancies by calling 1-855-215-2288. Educational materials can be found at [www.ipsen.com/canadaen/SOEDUMAT](http://www.ipsen.com/canadaen/SOEDUMAT).

### 7.1.2 Breast-feeding

No data are available on the presence of SOHONOS or its main metabolites in human breast milk, or the effects of SOHONOS on the breastfed child, or on milk production. Due to the potential for serious adverse reactions from SOHONOS in a breastfed child, patients who are breastfeeding should not take SOHONOS and should not breastfeed for at least 1 month following cessation of SOHONOS.

### 7.1.3 Pediatrics

The safety and effectiveness of SOHONOS for the treatment of FOP has been established in females  $\geq 8$  and males  $\geq 10$  years of age. SOHONOS should not be used in children aged below 8 years for females and below 10 years for males. Clinical studies in subjects aged 4 years and above have shown that growing patients with open epiphyses are at risk of developing PPC of growth plates when treated with SOHONOS (see [Musculoskeletal](#)).

Bone X-rays should be conducted every 3 months for pediatric patients before physal closure (frequency of monitoring may depend on patient characteristics such as age, pubertal status and skeletal maturity). Should a patient exhibit signs of PPC based on clinical and radiologic evaluations, an assessment of the benefits and risks should be made in order to determine the appropriateness of continued treatment versus temporary or permanent discontinuation of SOHONOS, until the patient reaches skeletal maturity.

### 7.1.4 Geriatrics

Clinical studies of SOHONOS did not include sufficient numbers of subjects aged 65 and over to determine whether they respond differently from younger subjects. The oldest subject in the clinical trials was 61 years of age. The median age at the time of death in FOP is 40 years of age. There were no clinically relevant differences in SOHONOS pharmacokinetics between young males and elderly males. In general, dose selection for an elderly patient ( $\geq 65$  years) should be done with caution.

## 8 ADVERSE REACTIONS

### 8.1 Adverse Reaction Overview

The most commonly reported adverse reactions (more than 10%) reported in FOP subjects aged  $\geq 8$  years (females)/10 years (males) years old were: cutaneous, including dry skin (78%), pruritus (55%), alopecia (41%), rash (39%), erythema (32%), skin exfoliation (31%), skin reaction (24%) drug eruption (17%), and skin irritation (12%); gastrointestinal, including lip dry (55%), chapped lips (17%), dry mouth (13%), cheilitis (11%) and nausea (11%); infectious, including paronychia (14%); musculoskeletal, including arthralgia (14%); ocular, including dry eye (26%); injury, poisoning and procedural complications, including skin abrasion (21%); respiratory, including epistaxis (12%) and neurological, including headache (17%). The

majority of adverse reactions were mild or moderate in severity across all SOHONOS FOP trials.

Serious adverse reactions of PPC reported in pediatric FOP subjects aged  $\geq 8$  years (females)/10 years (males) to  $< 14$  years occurred in 10/39 subjects (26%) and  $< 8$  years (females) and 10 years (males) occurred in 14/25 subjects (56%). Close monitoring of PPC is recommended (see **Musculoskeletal**). The serious adverse reaction of cellulitis occurred in 2 subjects (1.4%) with FOP aged  $\geq 8$  (females)/10 (males) years old. All others were reported in single subjects and included the following: anaemia, ankle fracture, seizure and epiphyseal disorder (frayed metaphyseal edge), each in 0.7% of subjects.

Adverse events (AEs) leading to permanent discontinuation occurred in 7% of SOHONOS treated subjects. Among them, the most common AEs leading to treatment discontinuation were dry skin (1.4%) and cellulitis, furuncle, localised infection, parainfluenza virus infection, epiphyses premature fusion (PPC), mobility decreased, erythema, malnutrition and intentional self-injury, each in 0.7% of subjects. No study drug discontinuations were reported in placebo/untreated subjects due to AEs.

Mucocutaneous AEs most commonly led to dose reductions during SOHONOS 20/10 mg flare-up treatment (35%) and during chronic treatment (3%).

Overall, the most common AEs leading to dose reduction were mucocutaneous including pruritus (9%), dry skin (8%), drug eruption (7%), and skin exfoliation (4%). In FOP clinical trials, there was no treatment-related increase in suicide ideation, suicidal behaviour or psychiatric disorders overall relative to untreated FOP subjects.

There have been no reports of pregnancy or in utero exposure reported in clinical studies with SOHONOS.

## 8.2 Clinical Trial Adverse Reactions

Clinical trials are conducted under very specific conditions. The adverse reaction rates observed in the clinical trials may not reflect the rates observed in practice and should not be compared to the rates in the clinical trials of another drug. Adverse reaction information from clinical trials may be useful in identifying and approximating rates of adverse drug reactions in real-world use.

The data from the FOP clinical trials described below reflect exposure to SOHONOS in a total of 164 subjects (including 139 subjects aged  $\geq 8$  (females) and  $\geq 10$  (males) years of age) for a mean duration of 94.1 weeks, to a maximum of 3.8 years. Subjects received either:

- Chronic/flare-up regimen: 5 mg daily dose with a 20/10 mg dose for 12 weeks at the time of flare-up (4 weeks of 20 mg once daily followed by 10 mg once daily for 8 weeks)
- Flare-up regimen only: either the 20/10 mg dose for 12 weeks, a 10/5 mg dose for 6 weeks (10 mg once daily for 2 weeks followed by 5 mg once daily for 4 weeks) or a 5/2.5 mg dose for 6 weeks (5 mg once daily for 2 weeks followed by 2.5 mg once daily for 4 weeks).

**Table 6. Adverse Reactions Reported in FOP Subjects Aged ≥ 8 (Females)/ 10 (Males) Years in All Clinical Trials (Incidence More Than ≥5% in the SOHONOS Exposed Patients)**

System Organ Class Preferred Term	SOHONOS Treatment Period		SOHONOS Total** (n=139) n (%)
	Chronic* 5 mg (n=130) n (%)	Flare-up* 20/10 mg (n=100) n (%)	
<b>Eye disorders</b>	21 (16)	31 (31)	46 (33)
Dry eye	13 (10)	21 (21)	36 (26)
Ocular hyperemia	1 (1)	7 (7)	9 (7)
<b>Gastrointestinal disorders</b>	71 (55)	52 (52)	105 (76)
Lip dry	48 (37)	29 (29)	77 (55)
Chapped lips	10 (8)	12 (12)	24 (17)
Dry mouth	9 (7)	6 (6)	18 (13)
Nausea	7 (5)	6 (6)	15 (11)
Vomiting	6 (5)	5 (5)	12 (9)
Cheilitis	5 (4)	11 (11)	15 (11)
Diarrhea	4 (3)	2 (2)	8 (6)
<b>Infections and infestations</b>	42 (32)	36 (36)	68 (49)
Paronychia	11 (9)	11 (11)	20 (14)
<b>Injury, poisoning and procedural complications</b>	24 (19)	25 (25)	46 (33)
Skin abrasion	10 (8)	20 (20)	29 (21)
Sunburn	4 (3)	3 (3)	8 (6)
<b>Metabolism and nutrition disorders</b>	8 (6)	13 (13)	25 (18)
Decreased appetite	3 (2)	7 (7)	11 (8)
<b>Musculoskeletal and connective tissue disorders</b>	27 (21)	30 (30)	49 (35)
Arthralgia	8 (6)	13 (13)	19 (14)
Epiphysis premature fusion <sup>1</sup>	5 (4)	2 (2)	7 (5)
Back Pain	3 (2)	4 (4)	8 (6)
Joint swelling	2 (2)	5 (5)	7 (5)
<b>Nervous system disorders</b>	13 (10)	21 (21)	32 (23)
Headache	9 (7)	10 (10)	23 (17)
<b>Psychiatric disorders</b>	15 (12)	16 (16)	31 (22)
Depressed mood	4 (3)	3 (3)	7 (5)
Irritability	3 (2)	2 (2)	8 (6)
<b>Respiratory, thoracic and mediastinal disorders</b>	23 (18)	14 (14)	38 (27)
Epistaxis	11 (9)	7 (7)	17 (12)
<b>Skin and subcutaneous tissue disorders</b>	113 (87)	90 (90)	136 (98)
Dry skin	75 (58)	58 (58)	109 (78)
Alopecia	31 (24)	29 (29)	57 (41)
Pruritus <sup>2</sup>	43 (33)	45 (45)	76 (55)
Rash <sup>3</sup>	33 (25)	29 (29)	54 (39)
Skin exfoliation	20 (15)	28 (28)	43 (31)
Erythema	19 (15)	28 (28)	44 (32)
Drug eruption	12 (9)	15 (15)	23 (17)
Skin irritation	8 (6)	8 (8)	16 (12)
Onychoclasia	5 (4)	5 (5)	11 (8)
Skin fissures	4 (3)	9 (9)	11 (8)

System Organ Class Preferred Term	SOHONOS Treatment Period		
	Chronic*	Flare-up*	SOHONOS Total**
	5 mg (n=130) n (%)	20/10 mg (n=100) n (%)	(n=139) n (%)
Blister	4 (3)	5 (5)	9 (7)
Skin reaction <sup>4</sup>	17 (13)	17 (17)	33 (24)
Decubitus ulcer	3 (2)	5 (5)	8 (6)
Madarosis	3 (2)	4 (4)	7 (5)
Ingrowing nail	2 (2)	5 (5)	7 (5)
<b>Vascular disorders</b>	2 (2)	8 (8)	9 (7)
Flushing	1 (1)	6 (6)	7 (5)

\*ADRs were captured as subjects were receiving chronic or flare-up treatment in Phase 2 and 3 clinical studies regardless of prior chronic treatment

\*\*inclusive of all dosing regimens across all Phase 2 and 3 FOP studies

<sup>1</sup> Epiphysis premature fusion is PT used to capture premature physal closure or PPC

<sup>2</sup> Pruritus includes pruritus and pruritus generalized

<sup>3</sup> Rash includes rash, rash generalized and rash maculo-papular

<sup>4</sup> Skin reaction includes skin reaction, dermatitis and eczema

### 8.2.1 Clinical Trial Adverse Reactions – Pediatrics

Subjects <18 years with open epiphyses were assessed for growth during the clinical study. PPC was observed in 24 of 102 subjects (24%) <18 years of age and was more common in younger (<8/10 years: 14 of 25 subjects, 56%) compared with older subjects (≥8/10 to <14 years: 10 of 39 subjects, 26%).

### 8.3 Less Common Clinical Trial Adverse Reactions with SOHONOS (<5%)

Blood and lymphatic system disorders: anemia

Gastrointestinal disorders: abdominal pain; gastroesophageal reflux disease

General disorders and administrative site conditions: fatigue; ankle fracture

Infections and infestations: cellulitis; conjunctivitis; skin infection

Neoplasms benign, malignant and unspecified (including cysts and polyps): pyogenic granuloma

Nervous system disorders: seizure

Psychiatric disorders: suicidal ideation

Renal and urinary disorders: proteinuria

Skin and subcutaneous tissue disorders: urticaria; skin fragility; swelling face

### 8.4 Abnormal Laboratory Findings: Hematologic, Clinical Chemistry and Other Quantitative Data

Incidence of baseline and new-onset PCS hematology, chemistry and ECG values during chronic and flare-up treatment in subjects ≥8 years (females)/10 years (males) of age are summarized in [Table 7](#), [Table 8](#), and [Table 9](#). Overall, for both chronic and flare-up treatment there were no new-onset PCS abnormalities for AST/ALT or AST/ALT >3 × ULN together with total bilirubin >2 × ULN.

**Table 7. Incidence of New-onset Potentially Clinically Significant (PCS) Hematology Values in Subjects ≥8 (Females)/10 (Males) Years**

Parameter**	Chronic SOHONOS Treatment Period 5 mg (N=130)	Flare-up SOHONOS Treatment Period			Untreated* (NHS) (N=91)
		Flare-up 10/5 mg (N=25)	Flare-up 20/10 mg (N=99)	Flare-up Total (N=106)	
Hemoglobin					
N'	127	25	96	103	68
New-onset PCS high	0	0	0	0	0
New-onset PCS low	2 (1.6)	0	1 (1.0)	1 (1.0)	0
Hematocrit					
N'	127	25	96	103	68
New-onset PCS high	0	0	1 (1.0)	1 (1.0)	0
New-onset PCS low	1 (0.8)	0	5 (5.2)	5 (4.9)	0
Platelets					
N'	127	25	96	103	68
New-onset PCS high	1 (0.8)	0	1 (1.0)	1 (1.0)	0
New-onset PCS low	1 (0.8)	0	0	0	0
WBC					
N'	127	25	96	103	68
New-onset PCS high	3 (2.4)	0	8 (8.3)	9 (8.7)	0
New-onset PCS low	0	0	0	0	0

N'=number of subjects at the relevant visit; PCS=potentially clinically significant; WBC=white blood cell; NHS=Natural History Study

\*Assessments taken at Study Months 12, 24, and 36

\*\*Parameter new-onset PCS, n(%)

PCS defined as hemoglobin <9.5 g/dL; >19 g/dL; hematocrit <34% or >54%; platelets <75×10<sup>9</sup> L or >700×10<sup>9</sup>/L; WBC <2.8×10<sup>9</sup>/L or >16×10<sup>9</sup>/L

New-onset PCS values were defined as values that were not PCS at screen/baseline and were PCS after screen/baseline.

**Table 8. Incidence of New-onset Potentially Clinically Significant (PCS) Chemistry Values in Subjects ≥8 (Females)/10 (Males) Years**

Parameter**	Chronic SOHONOS Treatment Period 5 mg (N=130)	Flare-up SOHONOS Treatment Period			Untreated* (NHS) (N=91)
		Flare-up 10/5 mg (N=25)	Flare-up 20/10 mg (N=99)	Flare-up Total (N=106)	
Bilirubin					
N'	127	25	97	104	68
New-onset PCS high	0	1 (4.0)	2 (2.1)	2 (1.9)	0
Lipase					
N'	127	25	97	104	68
New-onset PCS high	3 (2.4)	2 (8.0)	5 (5.2)	7 (6.7)	1 (1.5)

N'=number of subjects at the relevant visit; PCS=potentially clinically significant; NHS=Natural History Study

\*Assessments taken at Study Months 12, 24, and 36

\*\*Parameter new-onset PCS, n(%)

PCS defined as: total bilirubin >2 mg/dL; lipase >3× ULN

New-onset PCS values were defined as values that were not PCS at screen/baseline and were PCS after screen/baseline.

**Table 9. Incidence of New-onset Potentially Clinically Significant (PCS) ECG Values in Subjects ≥8 (Females)/10 (Males) Years**

Parameter**	Chronic SOHONOS Treatment Period 5 mg (N=130)	Flare-up SOHONOS Treatment Period			Untreated* (NHS) (N=91)
		Flare-up 10/5 mg (N=25)	Flare-up 20/10 mg (N=99)	Flare-up Total (N=106)	
QTcF/QTcB					
N'	123	25	32	39	72
New-onset PCS high	0	0	0	0	0
Median PR Duration					
N'	123	25	32	40	76
New-onset PCS high	12 (9.8)	3 (12.0)	3 (9.4)	6 (15.0)	3 (3.9)
Median QT duration					
N'	123	25	32	39	72
New-onset PCS high	3 (2.4)	0	0	0	1 (1.4)
Median QRS duration					
N'	123	25	32	40	76
New-onset PCS high	35 (28.5)	6 (24.0)	9 (28.1)	13 (32.5)	17 (22.4)

N'=number of subjects at the relevant visit; PCS=potentially clinically significant; NHS=Natural History Study

\*Assessments taken at Study Months 12, 24, and 36

\*\*Parameter new-onset PCS, n(%)

Following are the PCS criteria for ECG parameters. PR interval high: >200 ms only OR increase from baseline ≥20 ms only OR 3) >200 ms and increase from baseline ≥20 ms; QRS interval high: >100 ms only OR increase from baseline ≥10 ms only OR >100 ms and increase from baseline ≥10 ms; QT interval high: >500 ms only OR increase from baseline ≥60 ms only OR >500 ms and increase from baseline ≥60 ms; QTcF or QTcB intervals high: >500 ms only OR increase from baseline ≥60 ms only OR >500 ms and increase from baseline ≥60 ms. New-onset PCS values were defined as values that were not PCS at screen/baseline and were PCS after screen/baseline.

## 9 DRUG INTERACTIONS

### 9.2 Drug Interactions Overview

#### CYP3A4 Inhibitors and Inducers

Co-administration of SOHONOS with ketoconazole, a strong CYP3A4 inhibitor, increased the steady-state exposures of SOHONOS approximately 121% and 212%, based on C<sub>max</sub> and AUC, respectively. Avoid concomitant use of a strong CYP3A4 inhibitor such as azole antifungals (e.g. ketoconazole, itraconazole), and macrolide antibiotics (e.g. clarithromycin) with SOHONOS. Advise patients to avoid grapefruit or grapefruit juice that are known to inhibit cytochrome P450 during SOHONOS treatment.

Coadministration of SOHONOS with rifampicin/rifampin, a strong CYP 3A4 inducer, decreased the exposure of SOHONOS approximately 81% and 89%, based on C<sub>max</sub> and AUC, respectively. Avoid concomitant use of strong CYP3A4 inducers (e.g. carbamazepine, phenytoin, rifabutin, St John's wort extract) with SOHONOS.

## Vitamin A

SOHONOS belongs to the same pharmacological class as vitamin A. Therefore, the use of both vitamin A and SOHONOS at the same time may lead to additive effects. Concomitant administration of vitamin A and/or other oral retinoids with SOHONOS must be avoided because of the risk of hypervitaminosis A.

## Tetracyclines

Systemic retinoid use has been associated with cases of benign intracranial hypertension (also called pseudotumor cerebri), some of which involved the concomitant use of tetracyclines. Avoid coadministration of SOHONOS with tetracyclines derivatives.

## Systemic corticosteroids

Corticosteroids were administered per standard of care in the FOP clinical trials (e.g. prednisone at 2 mg/kg to a maximum dose of 100 mg daily for 4 days). In the population PK analysis, there was no evidence that administration of prednisone affected SOHONOS pharmacokinetics. SOHONOS had no effect on the pharmacokinetics of prednisone or its metabolite prednisolone.

### **9.3 Drug-Behavioural Interactions**

Interactions with behavioural risks have not been established.

### **9.4 Drug-Drug Interactions**

The drugs listed in this table are based on either drug interaction case reports or studies, or potential interactions due to the expected magnitude and seriousness of the interaction (i.e. those identified as contraindicated).

**Table 10. Established or Potential Drug-Drug Interactions**

<b>Proper/Common Name</b>	<b>Source of Evidence</b>	<b>Effect</b>	<b>Clinical Comment</b>
Ketoconazole	CT	Co-administration increased the exposure of SOHONOS at steady-state approximately 212% based on AUC.	Avoid concomitant use of a strong CYP3A4 inhibitor such asazole antifungals (e.g. ketoconazole, itraconazole) and macrolide antibiotics (e.g. clarithromycin) with SOHONOS. Advise patients to avoid grapefruit or grapefruit juice that are known to inhibit cytochrome P450 during SOHONOS treatment.
Rifampicin/rifampin	CT	Co-administration decreased the exposure of SOHONOS approximately 89% based on AUC.	Avoid concomitant use of strong CYP3A4 inducers (e.g. carbamazepine, phenytoin, rifabutin, St John's wort extract) with SOHONOS.

Proper/Common Name	Source of Evidence	Effect	Clinical Comment
Tetracyclines	T	Systemic retinoid use has been associated with cases of benign intracranial hypertension (also called pseudotumor cerebri), some of which involved the concomitant use of tetracyclines.	Avoid co-administration of SOHONOS with tetracyclines derivatives.
Vitamin A	T	Use of both vitamin A and SOHONOS at the same time may lead to additive effects.	Avoid concomitant administration of vitamin A and/or other oral retinoids with SOHONOS.

Legend: CT = Clinical Trial; T = Theoretical

## 9.5 Drug-Food Interactions

Oral absorption of SOHONOS is increased when given with food. For this reason, SOHONOS should be taken with food. Advise patients to avoid grapefruit or grapefruit juice that are known to inhibit cytochrome P450 during SOHONOS treatment.

## 9.6 Drug-Herb Interactions

Concomitant use of a strong CYP3A4 inducer such as St John's wort extract should be avoided with SOHONOS (see [Table 10](#)).

## 9.7 Drug-Laboratory Test Interactions

Interactions with laboratory tests have not been established.

# 10 CLINICAL PHARMACOLOGY

## 10.1 Mechanism of Action

Palovarotene is an orally bioavailable retinoic acid receptor gamma (RAR $\gamma$ ) selective agonist. FOP is a genetic condition caused by a gain-of-function mutation in the ACVR1/ALK2 gene, which encodes activin receptor type 1A/activin receptor-like kinase 2, a bone morphogenetic protein (BMP) type I receptor. This ALK2 gain-of-function mutation aberrantly activates the BMP-mothers against decapentaplegic homolog (Smad)1/5/8 signaling pathway, diverting normal soft tissue (muscles, tendons and ligaments) injury repair mechanisms away from tissue regeneration by promoting chondrogenesis and heterotopic bone formation.

RAR $\gamma$  is expressed in chondrogenic cells and chondrocytes operating as an unliganded transcriptional repressor. In a human FOP fibroblast cell line carrying the gain-of-function R206H ALK2 receptor mutation, palovarotene inhibited BMP-mediated Smad1/5 signaling. In animal models of FOP and injury-induced heterotopic ossification (HO), palovarotene reduced new HO and maintained joint mobility by decreasing mast cell infiltration and fibroproliferative response at the site of injury. Through binding to RAR $\gamma$ , palovarotene decreases BMP signaling and inhibits SMAD1/5/8 signaling, which are deeply involved in the pathogenesis of myositis ossificans, hence in FOP. By interfering with these pathways, palovarotene prevents chondrogenesis and ultimately HO by enabling normal muscle tissue repair or regeneration to take place, which reduces damage to muscle tissue.

## 10.2 Pharmacodynamics

### Cardiac Electrophysiology

In a randomized, double-blind, placebo- and positive-controlled, 4-way crossover ECG assessment study in healthy adult subjects (N=31) receiving single 20 mg (therapeutic) and 50 mg (supratherapeutic) doses of palovarotene, no pharmacodynamic effects on QTc interval, QRS duration, PR interval, or heart rate were observed.

## 10.3 Pharmacokinetics

The pharmacokinetics of palovarotene after oral administration have been well characterized from single and multiple dose studies in healthy subjects, and in patients with FOP. Oral absorption of palovarotene is increased when given with food. For this reason, palovarotene was given with food in all the FOP trials.

In a population PK analysis, the pharmacokinetics of palovarotene were linear and dose proportional from 0.02 to 50 mg. Steady-state is achieved by Day 3.

**Table 11. Summary of Steady-State Palovarotene Pharmacokinetic Parameters in Patients with FOP**

Mean (SD) Steady State PK Parameters	C <sub>max</sub> (ng/mL)	T <sub>max</sub> (h)	t <sub>1/2</sub> (h)	AUC <sub>0-τ</sub> (ng.h/mL)	CL/F (L/h)
5 mg	40.6 (16.2)	3.00 (2.67, 6.07)	4.9 (1.4)	264 (98.4)	17.8 (8.64)
10 mg	78.4 (33.3)	3.00 (2.75, 10.0)	4.3 (0.7)	540 (226)	17.0 (7.80)
20 mg	165 (72.7)	3.00 (2.83, 10.0)	4.4 (1.2)	1060 (449)	16.5 (7.43)

### Absorption

After administration of palovarotene 20 mg once daily for 14 days after a standard breakfast to healthy adult subjects, the median T<sub>max</sub> was 4.6 hours, the average C<sub>max</sub> was 140 ng/mL, and the average AUC<sub>(0-τ)</sub> was 942 ng\*hr/mL. Little or no accumulation was observed following once-daily dosing. The mean steady-state trough plasma concentration was 3.5 ng/mL after once-daily 20 mg palovarotene.

Administration of a single dose of palovarotene (2 x 10 mg) in 23 subjects after a high-fat (approximately 50% of total caloric content of the meal), high-calorie (approximately 800 to 1200 calories) meal resulted in an approximate 40% increase in AUC<sub>τ</sub>, an approximate 16% increase in C<sub>max</sub> and an approximate delay in T<sub>max</sub> by 2 hours when compared with administration of palovarotene (2 x 10 mg) under fasting conditions.

Plasma palovarotene AUC<sub>τ</sub> and C<sub>max</sub> were comparable when either swallowed whole following a high-fat, high-calorie breakfast or the contents sprinkled onto applesauce following a high-fat, high-calorie breakfast.

### Distribution

Mean protein binding of palovarotene is 99.0% *in vitro*. The mean apparent volume of distribution at steady-state (V<sub>ss</sub>/F) is 319 L following 20 mg once daily doses.

## Metabolism

Palovarotene is extensively metabolized by CYP3A4 and to a minor extent by CYP2C8 and CYP2C19. Five metabolites, M1 (6,7-dihydroxy), M2 (6-hydroxy), M3 (7-hydroxy), M4a (6-oxo), and M4b (7-oxo), were observed for palovarotene and reached steady-state by Day 4 with high variability in plasma levels. Following administration of [<sup>14</sup>C]-radiolabelled palovarotene, the contribution of palovarotene and its four known major metabolites (M2, M3, M4a, and M4b) represented collectively 40% of the total exposure in plasma. The pharmacological activity of M3 and M4b is approximately 1.7% and 4.2% of the activity of the parent drug based on an in vitro RAR $\gamma$  transactivation assay.

## Elimination

After administration of palovarotene 20 mg once daily for 14 days following a standard breakfast, the mean elimination half-life is 8.7 hours. From a population PK analysis, the apparent total body clearance (CL/F) is estimated at 19.9 L/h across dose levels and across studies, supporting dose linearity.

Following administration of a 1 mg dose of [<sup>14</sup>C]-radiolabeled palovarotene in healthy subjects, 97.1% of the dose was recovered in the feces and 3.2% in the urine. More than 92% of the dose was recovered in the first 6 days postdose and mass balance was achieved with 100% of the dose recovered by Day 14.

## Special Populations and Conditions

Based on a population PK analysis, there was no evidence that age, sex, race, smoking status or health status affected palovarotene PK. Body weight was found to have a significant impact on palovarotene PK resulting in increasing exposure with decreasing weight at the same dose.

### Semen Exposure in Males:

Administration of SOHONOS to a male patient is considered unlikely to affect development of an embryo or fetus carried by a pregnant female sexual partner exposed to SOHONOS via the patient's semen. Based on the results of a clinical study in 24 healthy male subjects, and the maximal palovarotene amount that has been quantified in a single ejaculate (33 ng or ~0.00017% of the daily dose administered), the maximum potential fetal exposure to SOHONOS through semen is estimated to be 0.0066 ng/mL which is less than 1/100th of the exposure at the NOAEL for effects on embryofetal development.

**Pediatrics:** Weight-adjusted doses for skeletally immature children in the FOP studies were selected for four defined weight categories (see [4.2 Recommended Dose and Dosage Adjustment](#)) to provide similar exposure to adolescents and adults receiving doses from 5 to 20 mg SOHONOS. The appropriateness of the selected weight-based dosing was assessed in the population PK analysis. Based on simulations in a pediatric population, the derived steady-state exposure metrics (AUC<sub>0- $\tau$</sub> , and C<sub>max,ss</sub>) following weight-based dosing for 5, 10 and 20 mg (or dose equivalent) in skeletally immature pediatric patients is presented by weight category in [Table 12](#) below.

**Table 12. Summary of Steady-State Exposure (50%Median) Following 5, 10 and 20mg SOHONOS in Pediatric Population by Weight Category**

Dose	Weight Category / Dose	C <sub>max, ss</sub> (ng/mL)*	AUC <sub>24, ss</sub> (ng·hr/mL)*
5 mg QD	<20 kg / 2.5 mg	50.6 (47.1-54.0)	263 (243-282)
	20-40 kg / 3 mg	40.0 (38.2-42.2)	237 (225-248)
	40-60 kg / 4 mg	36.7 (34.3-39.0)	242 (228-254)
	≥60 kg / 5 mg	35.9 (32.8-39.6)	252 (232-277)
10 mg QD	<20 kg / 5 mg	101 (94.1-108)	527 (485-563)
	20-40 kg / 6 mg	80.0 (76.3-84.4)	474 (450-495)
	40-60 kg / 7.5 mg	68.8 (64.4-73.2)	454 (428-476)
	≥60 kg / 10 mg	71.9 (65.7-79.1)	504 (464-554)
20 mg QD	<20 kg / 10 mg	202 (188-216)	1054 (971-1126)
	20-40 kg / 12.5 mg	167 (159-176)	988 (938-1032)
	40-60 kg / 15 mg	138 (129-146)	909 (856-952)
	≥60 kg / 20 mg	144 (131-158)	1008 (928-1107)

QD=once daily  
\* 50<sup>th</sup> Median (90% PI)

The simulated overall exposure (AUC<sub>24, ss</sub>) was comparable for the equivalent doses across the different weight groups, indicating that the weight-based dose scheme provides similar exposure between the pediatric weight groups.

**Hepatic Insufficiency** Based on population pharmacokinetic analysis (n=701), no clinically significant differences in PK were observed in subjects with mild hepatic impairment (n=47; 7%), therefore no dose adjustment is required in these patients. The effect of moderate to severe hepatic impairment on palovarotene pharmacokinetics has not been evaluated. Therefore, use caution in patients with moderate hepatic impairment. Palovarotene is not recommended in patients with severe hepatic impairment.

**Renal Insufficiency** Based on a population pharmacokinetic analysis (n=701), no clinically significant differences in the PK of palovarotene were observed in subjects with mild (n=158, 22.5%) and moderate (n=24, 3.4%) renal impairment. The effect of severe renal impairment on palovarotene pharmacokinetics has not been evaluated. Therefore, use in patients with severe renal impairment is not recommended.

## 11 STORAGE, STABILITY AND DISPOSAL

Store at room temperature (15° to 30°C). Keep blister strip in the carton in order to protect from light.

## 12 SPECIAL HANDLING INSTRUCTIONS

Return any unused SOHONOS (palovarotene) capsules to the pharmacist.

## PART II: SCIENTIFIC INFORMATION

### 13 PHARMACEUTICAL INFORMATION

#### Drug Substance

Proper/Common name: Palovarotene


Chemical name: 4-[(E)-2(5,5,8,8-Tetramethyl-3-pyrazol-1-ylmethyl-5,6,7,8-tetrahydro-naphthalen-2-yl)-vinyl]-benzoic acid

or

4-[(1E)-2-[5,6,7,8-Tetrahydro-5,5,8,8-tetramethyl-3-(1 H-pyrazol-1-ylmethyl)-2-naphthalenyl]-ethenyl]-benzoic acid

Molecular formula and molecular mass: C<sub>27</sub>H<sub>30</sub>N<sub>2</sub>O<sub>2</sub> 414.54 g/mol

Structural formula:



**Table 13. Physicochemical Properties**

Description:	White to off-white solid. Crystalline.
Solubility:	
Water	Practically insoluble or insoluble (0.04 µg/mL)
0.01M Hydrochloric Acid, pH 2.0	Practically insoluble or insoluble (not detected)
0.01M Potassium Acetate, pH 4.0	Practically insoluble or insoluble (not detected)
0.01M Potassium Acetate, pH 5.0	Practically insoluble or insoluble (0.006 µg/mL)
0.01M Potassium Acetate, pH 6.0	Practically insoluble or insoluble (0.04 µg/mL)
0.01M Potassium Phosphate Monobasic, pH 7.0	Practically insoluble or insoluble (0.2 µg/mL)
0.01M Potassium Phosphate Monobasic, pH 8.0	Practically insoluble or insoluble (1.8 µg/mL)
Ethanol (absolute)	Slightly soluble (1 mg/mL)
Isopropanol	Very slightly soluble (0.7 mg/mL)
Methanol	Very slightly soluble (0.3 mg/mL)
Acetonitrile	Very slightly soluble (0.3 mg/mL)

## 14 CLINICAL TRIALS

### 14.1 Clinical Trials by Indication

#### Fibrodysplasia Ossificans Progressiva (FOP)

Reduce the formation of HO in adults & children  $\geq 8$  (females)/10 (males) years of age with FOP

**Table 14. Summary of Patient Demographics for Clinical Trials in FOP**

Study	Study Design	Dosage, Route of Administration and Duration	Study Subjects (n)	Median age (Range)	Sex
PVO-1A-301 (MOVE)	Phase 3, multi-center, open-label study	5 mg QD for up to 48 months, with dose escalation for flare-up treatment to 20 mg QD for 4 weeks, then 10 mg QD for 8 weeks (total of 12 weeks; or until flare-up resolution if treatment extended beyond 12 weeks). Weight-based dosing according to 4 categories (10 to <20 kg, 20 to <40 kg, 40 to <60 kg, and $\geq 60$ kg).	Total enrolled: 107  Principal safety set: 99 <ul style="list-style-type: none"> <li>• &lt;8/10 yrs: 20</li> <li>• <math>\geq 8/10</math> to 14 yrs: 42</li> <li>• &gt;14 yrs: 37</li> </ul>	13 (4, 61)	53.5% M
PVO-1A-202	Phase 2, multi-center, open-label study conducted in 3 parts	Part A: Oral QD 10 mg for 2 weeks, then 5 mg for 4 weeks for the next 2 subsequent flare-ups. Weight-based dosing when children 6+ years enrolled from Study PVO-1A-201.	Total enrolled: 40 <ul style="list-style-type: none"> <li>• &lt;8/10 yrs: 4</li> <li>• <math>\geq 8/10</math> to 14 yrs: 8</li> <li>• &gt;14 yrs: 28</li> </ul>	21 (7, 53)	45.0% M
		Part B: Adult Cohort (chronic/flare-up regimen): 5 mg QD for up to 24 months, with dose escalation for flare-up treatment to 20 mg QD for 4 weeks, then 10 mg QD for 8 weeks (total of 12 weeks or until flare-up resolution if treatment extended beyond 12 weeks).  Pediatric Cohort (flare-up regimen only): same as flare-up dosing in the Adult Cohort except dosing is weight-adjusted	Total enrolled: 54 <ul style="list-style-type: none"> <li>• &lt;8/10 yrs: 3</li> <li>• <math>\geq 8/10</math> to 14 yrs: 9</li> <li>• &gt;14 yrs: 42</li> </ul>	19 (7, 54)	42.6% M

Study	Study Design	Dosage, Route of Administration and Duration	Study Subjects (n)	Median age (Range)	Sex
		Part C: All subjects receive chronic/flare-up regimen from Part B (weight-adjusted for skeletally immature subjects). Part C is ongoing.	Total enrolled: 48 <ul style="list-style-type: none"> <li>• &lt;8/10 yrs: 0</li> <li>• ≥8/10 to 14 yrs: 8</li> <li>• &gt;14 yrs: 40</li> </ul>	20 (9, 48)	47.9 % M
PVO-1A-201	Multicenter, randomized, placebo controlled adaptive dose finding study	Cohort 1: oral QD 10 mg for 2 weeks, then 5.0 mg for 4 weeks, or placebo  Cohort 2: oral QD 10 mg for 2 weeks, then 5.0 mg for 4 weeks; oral QD 5 mg for 2 weeks, then 2.5 mg for 4 weeks; or placebo	Total enrolled: 40 <ul style="list-style-type: none"> <li>• &lt;8/10 yrs: 4</li> <li>• ≥8/10 to 14 yrs: 9</li> <li>• &gt;14 yrs: 27</li> </ul>	21 (7, 53)	45.0% M
PVO-1A-001	Natural history study (NHS)	Dosage not applicable (non-interventional study); 3-year follow-up.	Total enrolled: 114  Principal safety set :111 <ul style="list-style-type: none"> <li>• &lt;8/10 yrs: 23</li> <li>• ≥8/10 to 14 yrs: 30</li> <li>• &gt;14 yrs: 58</li> </ul>	15 (4, 56)	54.1% M

QD=once daily

The MOVE study (PVO-1A-301) was a Phase 3, single arm study in subjects with FOP aged 4 years and older. The study evaluated the efficacy and safety of the chronic/flare-up SOHONOS regimen in preventing new HO as assessed by low-dose, whole body CT (WBCT) imaging (excluding head) as compared to data from the Natural History Study (NHS, PVO-1A-001). All WBCT images from treated subjects in the MOVE study and untreated subjects in the NHS were read in a manner blinded to study origination. Of the 107 subjects enrolled in the MOVE study, 99 had the R206H mutation and 8 had other FOP mutations. Of the 99 with the R206H mutation, 97 had at least one postbaseline HO volume measurement and were included in the Full Analysis Set.

In the MOVE study, subjects received SOHONOS 5 mg daily with increased dosing at the time of a flare-up (defined as at least one symptom (e.g. pain, swelling, redness) consistent with a previous flare-up or a substantial high-risk traumatic event likely to lead to a flare-up). Flare-up treatment was extended in 4-week increments for persistent symptoms. Anytime during flare-up treatment, the 12-week treatment restarted if subject had another flare-up or a substantial high-risk traumatic event. The dosing was adjusted according to body weight in skeletally immature children (children who had not reached at least 90% skeletal maturity defined as a bone age of ≥12 years 0 months for girls and ≥14 years 0 months for boys). The treatment groups assessed in the chronic/flare-up regimen were well matched for baseline demographics compared to subjects in the NHS study.

The median age (range) of subjects who contributed to the efficacy analysis in the target population ≥8 (females)/10 (males) in the SOHONOS group (N=77) was 14 (8, 61) years; and 18 (9, 56) years in the untreated group (N=76) from the NHS. There were more male than female subjects in both the SOHONOS (54.4% and 45.6%, respectively) and untreated (51.1% and 48.9%, respectively) groups.

**Table 15. PVO-1A-301 (MOVE) Study: Baseline Demographics and FOP Characteristics in Subjects  $\geq 8$  (Females)/10 (Males) Years**

	SOHONOS (N = 79)	Untreated* (N = 88)
Age at FOP diagnosis (years)		
Mean (SD)	6.6 (5.01)	7.5 (5.30)
Time from FOP diagnosis to enrollment (years)		
Mean (SD)	11.4 (9.75)	13.4 (9.69)
Associated Clinical Findings		
Cervical Spine Malformations	35 (44.3%)	45 (51.1%)
Hearing Loss	36 (45.6%)	31 (35.2%)
Thumb Malformations	37 (46.8%)	49 (55.7%)
Shortened Femoral Necks	10 (12.7%)	13 (14.8%)
Osteochondromas		
Tibia	31 (39.2%)	33 (37.5%)
Femur	5 (6.3%)	6 (6.8%)
Humerus	2 (2.5%)	2 (2.3%)
Number of flare-ups within past 12 months		
Mean (SD)	1.2 (1.5)	1.5 (2.0)
Median (min, max)	1.0 (0, 6)	1.0 (0, 10)
Time since last flare-up, months		
Mean (SD)	27.0 (40.4)	21.3 (33.8)
Median (min, max)	11 (1, 199)	7 (0, 181)

FOP= fibrodysplasia ossificans progressiva; SD=standard deviation

\*The untreated group includes subjects from the Natural History Study

The flare-up only regimen was assessed in the Phase 2 program, including the double-blind, placebo-controlled Study PVO-1A-201 and open-label extension Study PVO-1A-202. Subjects participating in Study PVO-1A-201 were randomized in a 3:3:2 fashion to SOHONOS 10 mg for 2 weeks, then 5 mg for 4 weeks (10/5 mg treatment), SOHONOS 5 mg for 2 weeks, then 2.5 mg for 4 weeks (5/2.5 mg treatment) or placebo for 6 weeks, followed by a 6-week observation period for all groups.

In Study PVO-1A-202 Part A subjects experiencing another flare-up received SOHONOS 10/5 mg treatment in an open label manner. In Study PVO-1A-202 Part B, subjects who were at least 90% skeletally mature received chronic 5 mg daily treatment with increased dosing at the time of a flare-up to 20 mg for 4 weeks followed by 10 mg for 8 weeks (chronic/flare-up regimen), with continuation of treatment in 4-week increments for persistent symptoms. Skeletally immature subjects received the 20/10 mg flare-up treatment (weight-adjusted) in Part B.

The median (range) age of subjects in the target population ( $\geq 8$  (females)/10 (males)) was 11 (7, 34) years (N=12); and 15 (7, 53) years in the untreated group (N=38). The percentage of male subjects in the target population was 33.3% and 47.0% in the untreated group.

### Results of the MOVE Study (PVO-1A-301) in FOP

The mean annualized new HO volume from post hoc analyses in the MOVE study (PVO-1A-301) in the overall population for treated and untreated subjects is shown in [Table 16](#) and in the indicated population of subjects  $\geq 8$  (females)/10 (males) for treated and untreated subjects is shown in [Table 17](#) and [Figure 1](#). Post hoc analyses were performed after the prespecified analysis using a Bayesian compound Poisson model with square-root transformation demonstrated that the futility boundary had been crossed. These results revealed that using the square-root transformation of the data in the Bayesian model moved the statistical

conclusion from significant therapeutic benefit of SOHONOS to showing futility. Additional analyses using the Bayesian and weighted linear mixed effect (wLME) models of annualized new HO volume without square-root transformation (including all raw data) demonstrated the efficacy of SOHONOS.

**Table 16. Mean Annualized New HO Volume in the Overall Population in the MOVE Study (PVO-1A-301)**

		<b>SOHONOS (N=97)</b>	<b>Untreated (N=98)</b>
New HO (mm <sup>3</sup> )	Mean (SEM)	8821 (2972)	23318 (4938)
	% reduction (SOHONOS vs untreated)	62%	
	LS mean (SEM)	8969 (4085)	20579 (3365)
	% reduction (SOHONOS vs untreated)	56%	
		<b>wLME estimate (95% CI)</b>	<b>Nominal p-value</b>
	Treatment difference	-11611 (-21977, 1244)	0.0292

CI=confidence interval; HO=heterotopic ossification; LS mean=least square mean; SEM=standard error of the mean; wLME=weighted linear mixed effect.

Note: The new HO weighted linear mixed effect (wLME) LS mean estimate and SEM are from a mixed model with dependent variable annualized new HO and independent variables including fixed effects of treatment and baseline total HO/baseline age and a random subject effect.


**Table 17. Mean Annualized New HO Volume in Subjects ≥8 Years (Females)/10 Years (Males) of Age in the MOVE Study (PVO-1A-301)**

		<b>SOHONOS (N=77)</b>	<b>Untreated (N=76)</b>
New HO (mm <sup>3</sup> )	Mean (SEM)	10655 (3641)	24777 (6192)
	% reduction (SOHONOS vs untreated)	57%	
	LS mean (SEM)	10572 (4943)	21617 (4165)
	% reduction (SOHONOS vs untreated)	51%	
		<b>wLME estimate (95% CI)</b>	<b>Nominal p-value</b>
	Treatment difference	-11045 (-24196, 2105)	0.0953

CI=confidence interval; FAS=Full Analysis Set; HO=heterotopic ossification; LS mean=least square mean; SEM=standard error of mean; wLME=weighted linear mixed effect model.

Note: The annualized new HO wLME LS mean estimate and SEM are from a mixed model with dependent variable annualized new HO and independent variables including fixed effects of treatment and baseline total HO/baseline age and a random subject effect.

**Figure 1. Mean Annualized New HO Volume in Subjects ≥8 (Females)/10 (Males) Years in the MOVE Study (PVO-1A-301)**



HO=heterotopic ossification; SE=standard error.

**Results of Studies PVO-1A-201 and PVO-1A-202 in FOP**

In the overall population, the Phase 2 studies demonstrated a reduction of 72% (p-value of 0.02) in new HO volume in the 15 flare-ups treated with the 20/10 mg flare-up only dose (3045 mm<sup>3</sup>) compared to 47 placebo/untreated flare-ups (10,780 mm<sup>3</sup>) (Table 18). In the target population, the Phase 2 studies demonstrated a reduction of 72% (p-value of 0.04) in new HO volume in the 14 flare-ups treated with the 20/10 mg flare-up only dose (3262 mm<sup>3</sup>) compared with 43 placebo/untreated flare-ups (11,712 mm<sup>3</sup>) (Table 19 and Figure 2). These results are supported by the 10/5 mg flare-up dose in both the target population (2807 mm<sup>3</sup>; 76% reduction; p-value of 0.10) and total population (3010 mm<sup>3</sup>; 72% reduction; p-value of 0.11).

**Table 18. Flare-up New HO at Week 12 for the Placebo/Untreated and SOHONOS Treated Flare-ups in the Overall Population**

		<b>SOHONOS 20/10 mg (M=15)</b>	<b>PBO/Untreated (M=47)</b>
New HO volume (mm <sup>3</sup> ) (including new HO volumes = 0 <sup>a</sup> )	Mean (Std Err)	3045 (1408)	10780 (4841)
	Nominal p-Value <sup>b</sup>	0.02	

HO=heterotopic ossification; M=number of flare-ups analysed; PBO=placebo; Std Err=standard error.

<sup>a</sup> When a flare-up had no baseline HO, 0 was assigned for analysis.

<sup>b</sup> Based on hierarchical bootstrap estimates, which included a 4-level treatment factor (SOHONOS 20/10 mg Episodic; SOHONOS 20/10 mg Chronic; SOHONOS 10/5 mg; Placebo/Untreated) and covariates Flare-up Location at hip (Yes vs. No), Steroid Use flare-up (Yes vs. No), Sex (Male vs. Female), Age (Years) at flare-up Day 1.

**Table 19. Flare-up New HO at Week 12 for the Placebo/Untreated and SOHONOS Treated Flare-ups in Subjects ≥8 Years (Females)/10 Years (Males) of Age**


		SOHONOS 20/10 mg (M=17)	PBO/Untreated (M=46)
Number of flare-ups analyzed		14	43
New HO volume (mm <sup>3</sup> ) (including new HO volumes = 0 <sup>a</sup> )	Mean (Std Err)	3262 (1494)	11712 (5274)
	Nominal p-Value <sup>b</sup>	0.04	

HO=heterotopic ossification; M=number of flare-ups; PBO=placebo; Std Err=standard error.

<sup>a</sup> When a flare-up had no baseline HO, 0 was assigned for analysis.

<sup>b</sup> Based on hierarchical bootstrap estimates, which included a 4-level treatment factor (SOHONOS 20/10 mg Episodic; SOHONOS 20/10 mg Chronic; SOHONOS 10/5 mg; Placebo/Untreated) and covariates Flare-up Location at hip/shoulder/knee (Yes vs. No), Steroid Use flare-up (Yes vs. No), Sex (Male vs. Female), Age (Years) at flare-up Day 1.

**Figure 2. Mean Volume of New HO at Week 12 in SOHONOS vs Placebo/Untreated Flare-ups in Subjects ≥8 (Females)/10 (Males)Years of Age**



Numbers in parentheses are the number of flare-ups; HO=heterotopic ossification; SE=standard error.

## 15 MICROBIOLOGY

No microbiological information is required for this drug product.

## 16 NON-CLINICAL TOXICOLOGY

### General Toxicology:

The toxicity of palovarotene was evaluated in repeat-dose daily oral toxicity studies lasting up to 6 weeks in juvenile rats (0.1, 0.5, and 1.2 mg/kg/day), up to 26 weeks in adult rats (0.3, 0.6 and 1.0 mg/kg/day), up to 4 weeks in adult rabbits (0.5, 1 and 2.5 mg/kg/day), and up to 39 weeks in adult dogs (0.012, 0.025, 0.040 and 0.120 mg/kg/day). The toxicity of repeated daily doses of palovarotene was similar to that of vitamin A and other retinoids. The dose-limiting toxicities of palovarotene were a subset of typical retinoid toxicities, primarily mucocutaneous toxicity in all nonclinical species and skeletal effects in rats. In this species, the effects of palovarotene were similar regardless of age; effects targeted skeleton and skin in both juvenile rats as young as 1 week of age and in adult rats. As would be expected, skin effects were dose-limiting in adult rats while skeletal effects were dose-limiting in younger rats. The toxicity profile of palovarotene metabolites was similar to that of palovarotene; no unique or unexpected toxicities were identified.

### *Mucocutaneous effects*

Mucocutaneous toxicity occurred in dose-related manner in adult animals, involved mainly the skin and was characterized microscopically by erythema, edema, epithelial hyperplasia, hyperkeratosis and/or hypergranulosis in the epidermis. Their intensity increased in a time-dependent manner up to 4 weeks of treatment, and they were reversible with discontinuation of dosing. In juvenile rats, skin lesions were limited to dry skin, thin fur cover, and/or scab and generally were no longer noted during the recovery period. These effects are generally observed at all dose levels, starting below the clinically relevant exposures.

Squamous epithelium of other tissues, such as nonglandular mucosa of the stomach (forestomach, rodent-specific), esophagus (rabbits only, non-adverse), and conjunctivae of the eye and surface of inner ear (dogs only), was also affected. The changes identified in the esophagus of rabbits after 4 weeks of oral administration at 1 mg/kg/day and above (i.e. below the highest clinical exposure) did not compromise the integrity of the esophageal mucosa or the health of the animal and were not interpreted to be toxicologically significant. No effects were observed in the non-squamous mucosa of glandular stomach or intestinal tissues in rats, dogs, or rabbits, which suggests that the potential for gastrointestinal toxicity is low.

### *Skeletal effects*

The potential effects of chronic administration of palovarotene in pediatric patients were investigated in juvenile rats given daily oral doses of palovarotene at 0.1, 0.5, or 1.2 mg/kg/day throughout the period of skeletal growth (from weaning through puberty). Palovarotene produced dose-dependent effects on bone size, shape, and mass and/or geometry, all of which appeared to result from impaired physal cartilage maturation/differentiation. In particular, narrowing/closure of the growth plate in the proximal femur resulted in changes in femoral head shape at  $\geq 0.5$  mg/kg/day and avascular necrosis (AVN) of the femoral head at 1.2 mg/kg/day. In addition, reduced bone mass and/or geometry were noted at  $\geq 0.5$  mg/kg/day. Ongoing recovery was noted at 0.5 mg/kg/day only. The NOAEL for skeletal toxicity (i.e. 0.1 mg/kg/day) was below the range of clinically relevant exposures.

In adult rats, repeated daily oral doses of palovarotene at 0.3, 0.6 and 1.0 mg/kg/day for 26 weeks caused, at all doses, chondrodystrophy in the growth plate, which remains open into adulthood in this species. Chondrodystrophy was associated with bone brittleness/fracture in rats when treated at 5 mg/kg/day (i.e. at exposure similar to the highest clinical exposure) for 4 weeks. The effects on bone growth are not considered relevant to humans once the growth

plates have closed (i.e. near the end of puberty: approximately 14-15 years for girls and 15-17 years for boys).

## **Carcinogenicity**

Carcinogenicity studies have not been conducted with palovarotene. Neither palovarotene nor its metabolites were mutagenic in Ames assays. Palovarotene did not have any clastogenic effect in the in vivo mouse micronucleus study at up to two doses of 25 mg/kg given on two consecutive days. Clastogenic activity was noted for palovarotene and its metabolites in the in vitro chromosomal aberration assay with human peripheral blood lymphocytes but only at cytotoxic concentrations that reduced the mitotic index by more than 50%. Specifically, for palovarotene, the threshold for clastogenic activity was between 30,000 and 50,000 ng/mL with the frequency of aberrations being increased at test concentrations  $\geq 50,000$  ng/mL but not at concentrations  $\leq 30,000$  ng/mL. For palovarotene metabolites, the frequency of cells with structural aberrations was significantly increased at a metabolite mixture concentration of 250,000 ng/mL (3-hour exposure) whereas following 24-hour exposure, no statistically significant increases of the frequency of chromosomal aberrations were observed. The concentrations at which these effects occurred with palovarotene or its human metabolites were orders of magnitude higher than the anticipated human maximal  $C_{max}$  at the highest proposed clinical dose (138-202 ng/mL at 20 mg/day). Based on these results, palovarotene and its metabolites are not considered to pose a genotoxic hazard to human subjects.

## **Reproductive and Developmental Toxicology:**

### *Impairment of Fertility*

Palovarotene given orally to female rats for 2 weeks prior to mating and up to Day 7 postcoitum at doses of 0.3 and 1 mg/kg/day revealed no effects on reproductive function, fertility or early embryonic development. Prolonged periods of diestrus and a slightly lower ovulation rate were noted at 3 mg/kg/day (high dose). Consequently, the 1 mg/kg/day dose (below the range of clinically relevant exposures) was considered as the NOAEL. Oral administration of palovarotene to male rats for 9 weeks (i.e. before cohabitation through two weeks after mating) at 0.3 and 1 mg/kg/day did not affect reproductive function, fertility, or early embryonic development. The dose of 1 mg/kg/day, which is below the range of clinically relevant exposures, was therefore considered as the NOAEL.

Evidence of testicular toxicity (including seminiferous tubule degeneration) was seen in male rats given daily oral doses at 5 mg/kg/day for 4 weeks, which exceeded the 4-week MTD. No testicular findings were observed at lower doses (0.04, 0.2 and 1 mg/kg/day) in this study. Moreover, no findings in female reproductive organs or evidence of testicular toxicity were observed in rat or dog chronic toxicity studies (i.e. up to 26 or 39 weeks, respectively).

### *Animal Reproduction*

Oral administration of palovarotene to pregnant rats from GD (Gestation Day) 6 to GD17 at doses of 0.25 and 1.25 mg/kg/day induced slight maternal toxicity (reduced maternal body weight gain and/or decreased food consumption). It also induced foetal malformations typical of retinoids (e.g. cleft palate, misshapen skull bones, short/long bones). The NOAEL for both maternal and foetal toxicity was therefore set at 0.01 mg/kg/day, which is below the range of clinical relevant exposures.

## PATIENT MEDICATION INFORMATION

### READ THIS FOR SAFE AND EFFECTIVE USE OF YOUR MEDICINE

Pr **SOHONOS**<sup>TM</sup>

#### palovarotene capsules

Read this carefully before you start taking **SOHONOS** and each time you get a refill. This leaflet is a summary and will not tell you everything about this drug. Talk to your healthcare professional about your medical condition and treatment and ask if there is any new information about **SOHONOS**.

#### **Serious Warnings and Precautions**

- **SOHONOS** can cause bone growth to stop early in growing children. This may lead to possible short height and differences in arm or leg length. Your doctor will explain the risks to you (i.e. the patient) or caregiver and child. Children who are actively growing should be closely monitored (every 3 months or as determined by your doctor) during treatment with **SOHONOS**.
- **SOHONOS** can seriously harm an unborn baby. Do **NOT** use **SOHONOS** if you are pregnant, think you may be pregnant or are planning to become pregnant. If you are of childbearing potential, you should **only** take **SOHONOS** if **ALL** the conditions described below under “**SOHONOS Pregnancy Prevention Plan**” are met.

#### **SOHONOS Pregnancy Prevention Plan**

Patients who could get pregnant must meet **ALL** of the conditions below to use **SOHONOS**.

##### **1. You must NOT take SOHONOS if you are pregnant, or breastfeeding**

**SOHONOS** can seriously harm an unborn baby. It can cause serious defects to the unborn baby's brain, face, ears, eyes, heart and certain glands (thymus gland and parathyroid gland). It also makes a miscarriage more likely. This may happen even if **SOHONOS** is taken only for a short time during pregnancy.

##### **You must NOT**

- take **SOHONOS** if you are pregnant or if you think you might be pregnant.
- take **SOHONOS** if you are breastfeeding. The medicine is likely to pass into your milk and may harm your baby.
- take **SOHONOS** if you could get pregnant during treatment.
- get pregnant for one month after stopping this treatment because some medicine may still be left in your body.

##### **2. If you could get pregnant, you and your doctor must discuss the strict rules to follow before, during and after taking SOHONOS.**

- Your doctor must explain the risk of harm to the unborn baby. You must understand why you must not get pregnant and what you need to do to prevent pregnancy.
- You must have talked about contraception (birth control) with your doctor. The doctor will give you information on how to avoid getting pregnant. Your doctor may send you to a specialist for birth control advice.

- You must have talked about pregnancy testing with your doctor. You must understand and agree to have regular pregnancy testing.
- Your doctor must explain to you what to do if you become pregnant, or think you might be pregnant.

**3. You must avoid getting pregnant by using effective birth control before, during and after taking SOHONOS.**

- You must be able and willing to comply with the mandatory birth control measures. Discuss with your doctor which methods would be suitable for you.
- You must use birth control even if you are not sexually active (unless your doctor decides this is not necessary).
- You must use birth control for a month before taking SOHONOS, during treatment and for a month afterwards.
- You must use at least one highly effective method of birth control (for example an intrauterine device) or, two effective methods that work in different ways (for example a hormonal birth control pill and a condom).

If you are taking SOHONOS **only** to treat flare-ups, you must continue to use effective birth control even during times when you are not taking SOHONOS. This is because the timing of your flare-ups may not be predictable.


**4. You must have pregnancy testing before, during and after taking SOHONOS**

- Before you start treatment, you must take a pregnancy test. The test must show that you are not pregnant when starting treatment with SOHONOS.
- You must have regular pregnancy tests, every month during treatment with SOHONOS. Your doctor may also ask you to take a pregnancy test 1 month after stopping SOHONOS because some medicine may still be left in your body.

**5. If you become pregnant while taking SOHONOS or within one month after stopping treatment, you must contact your doctor right away**

- If you are still taking SOHONOS, you must stop treatment and tell your doctor right away.
- You and your doctor must discuss:
  - If you want to continue with your pregnancy
  - The serious risk of your baby having severe birth defects
- You or your doctor should report the pregnancy by calling 1-855-215-2288. If you agree, your doctor can also enroll you in the SOHONOS registry by calling the same number (1-855-215-2288). Educational materials can be found at [www.ipsen.com/canadaen/SOEDUMAT](http://www.ipsen.com/canadaen/SOEDUMAT).

**If your pre-treatment counselling did not include an in-depth conversation about the SOHONOS Pregnancy Prevention Plan, please contact your doctor.**



### What is SOHONOS used for?

- SOHONOS is used to reduce the formation of heterotopic ossification. This is a condition where bone forms in soft tissues outside the skeleton. It is used in adults and children (females 8 years and older, males 10 years and older) who have the genetic disorder fibrodysplasia (myositis) ossificans progressiva, also called FOP.

### How does SOHONOS work?

SOHONOS contains palovarotene which belongs to a group of medicines called retinoids. It works by preventing bone formation in muscles, tendons or soft tissue.

### What are the ingredients in SOHONOS?

Medicinal ingredients: palovarotene

Non-medicinal ingredients: Croscarmellose sodium, gelatin, lactose monohydrate, magnesium stearate, microcrystalline cellulose, pharmaceutical grade printing ink, povidone, sodium lauryl sulfate, and titanium dioxide.

### SOHONOS comes in the following dosage forms:

Capsules: 1 mg, 1.5 mg, 2.5 mg, 5 mg and 10 mg

### Do not use SOHONOS if:

- You are pregnant or breastfeeding. Physicians and patients can report pregnancies by calling 1-855-215-2288.

- There is a chance you could become pregnant. You must follow the precautions under “**SOHONOS Pregnancy Prevention Plan**”.
- You are allergic to palovarotene, or any of the other ingredients of SOHONOS.
- You are allergic to other retinoids.

**To help avoid side effects and ensure proper use, talk to your healthcare professional before you take SOHONOS. Talk about any health conditions or problems you may have, including if you:**

- are pregnant, think you may be pregnant or are planning to have a baby.
- are breastfeeding.
- have liver or kidney problems.
- have been told by your doctor that you have an intolerance to some sugars. This is because SOHONOS contains lactose.

#### **Other warnings you should know about:**

SOHONOS can cause:

- **Sensitivity to sunlight and ultraviolet light and skin effects** (e.g. dry skin, loss of hair, itching, rashes). To reduce these side effects, your healthcare professional may recommend the use of:
  - Skin moisturizers, sunscreen with a broad-spectrum SPF value of 15 or higher, lip moisturisers, or other helpful treatments
  - Protective clothing and sunglasses to help reduce exposure to sunlight.
- **Dry eyes.** Your healthcare professional may recommend use of artificial tears (drops to lubricate the eye).

#### **Check-ups and testing**

You will have regular visits with your healthcare professional during treatment with SOHONOS. They will:

- Check for problems with normal bone growth (in growing children), every 3 months (or as determined by your healthcare professional).
- Check for signs and symptoms of mental health problems, such as depression, anxiety, mood alterations, suicidal thoughts and behaviours.
- Follow-up on monthly pregnancy testing in patients of childbearing potential.

#### **Driving and using machines**

SOHONOS may affect your ability to see in the dark. Before you drive or do tasks that require special attention, wait until you know how you respond to SOHONOS.

#### **Fertility**

It is not known if SOHONOS affects your ability to have children. Talk to your healthcare professional if this is a concern for you.

**Tell your healthcare professional about all the medicines you take, including any drugs, vitamins, minerals, natural supplements or alternative medicines.**

**The following may interact with SOHONOS:**

- Medicines used for skin conditions, known as ‘retinoids’ such as acitretin, alitretinoin, isotretinoin, tretinoin, adapalene and tazarotene.
- Medicines or vitamin supplements containing Vitamin A. Vitamin A in high doses has many of the same side effects as SOHONOS. Taking both together may increase your chance of getting side effects.
- Medicines used to treat fungal infections (antifungals), such as itraconazole and ketoconazole.
- Medicines used to treat bacterial infections (antibiotics) such as clarithromycin, rifampicin/rifampin, rifabutin, and tetracyclines.
- Medicines used to treat depression.
- Medicines used to treat epilepsy or seizures such as phenytoin and carbamazepine.
- Herbal preparations containing St. John’s Wort (*Hypericum perforatum*), sometimes used for treating depression or depression-related conditions such as anxiety.

### How to take SOHONOS:

- Take SOHONOS exactly as your healthcare professional tells you. Check with your healthcare professional if you are not sure.
- Swallow whole with food (after a meal), preferably at the same time each day.
- Avoid taking grapefruit-containing products for as long as you are using this medicine. It may increase the amount of SOHONOS in your blood.
- If you are unable to swallow capsules:
  - Open the capsules and sprinkle contents onto a teaspoon of soft food (such as applesauce, yogurt or pudding).
  - Swallow the mixture. If you want to take it later, the mixture can be stored for up to one hour at room temperature, provided that it is not exposed to direct sunlight.

### Instructions for handling SOHONOS:

- If you are pregnant, or are planning on becoming pregnant, you should avoid contact with SOHONOS.
- For caregivers emptying the capsule contents onto soft food:
  - Wear disposable gloves to avoid unintended exposure.
  - Use disposable paper towels and a container (e.g. a re-sealable bag) to collect waste.

### Usual dose:

- Your doctor will decide the dose of SOHONOS you need to take daily. This may depend on your age, weight and symptoms. **Do NOT change your dose of SOHONOS before discussing with your doctor.**
- If you have side effects during treatment, tell your doctor. They may need to adjust your dose, temporarily stop or completely stop your treatment with SOHONOS.
- The recommended dosage regimen of SOHONOS is a combination of:
  - Chronic treatment: daily treatment; and
  - Flare-up treatment: higher-dose treatment taken at the time of a flare-up.**Do NOT take both chronic treatment and flare-up treatment at the same time.** Your doctor will decide which treatment you should follow.
- **Contact your doctor right away when:**

- **You notice any symptom of a flare-up.** Your doctor will explain to you the signs and symptoms that you should look for. Symptoms may include but are not limited to: localised pain, swelling/inflammation in soft tissues such as muscles and tendons, redness, warmth, decreased joint range of motion, and stiffness.
- **You have a soft tissue injury.** This can happen during surgery, vaccinations, dental procedures, muscle weakness, bumps/bruises/falls or flu-like illnesses. Your doctor may start you on treatment to prevent a flare-up in these cases.
- **Your flare-up symptoms do not go away.** Your doctor may extend your flare-up treatment until the flare-up resolves.
- **You have another flare-up at any time during treatment.** This may be a new flare-up location or worsening of the original flare-up. Your doctor may need to consider restarting the 12-week flare-up treatment.
- **Your flare-up has resolved.** Your doctor will determine if you should return to the chronic daily treatment.

### Overdose:

If you think you, or a person you are caring for, have taken too much SOHONOS, contact a healthcare professional, hospital emergency department, or regional poison control centre immediately, even if there are no symptoms.

### Missed Dose:

- If you miss a dose, take the dose as soon as you remember. Take your next dose at the usual time.
- If you miss a dose by more than 6 hours, skip the missed dose. Continue with the next scheduled dose at the usual time.
- Do NOT take two doses at the same time or in the same day to make up for a dose you missed.

### What are possible side effects from using SOHONOS?

These are not all the possible side effects you may have when taking SOHONOS. If you experience side effects not listed here, tell your healthcare professional.

Side effects you may experience include:

- dry skin
- hair loss
- itching
- skin shedding
- rash
- hives
- eczema
- skin growth
- nose bleed
- headache
- dry and scaly lips
- dry mouth

- lip inflammation (cheilitis)
- nausea
- diarrhea
- acid reflux disease
- infection of the nail
- joint pain
- back pain
- dry eye
- sensitivity to sunlight

<b>Serious side effects and what to do about them</b>			
<b>Symptom / effect</b>	<b>Talk to your healthcare professional</b>		<b>Stop taking drug and get immediate medical help</b>
	<b>Only if severe</b>	<b>In all cases</b>	
<b>COMMON</b>			
<b>Mental health problems such as depression or psychosis (a severe mental disturbance)</b> <ul style="list-style-type: none"> <li>• changes in your mood such as becoming depressed, feeling sad, or having crying spells</li> <li>• losing interest in your usual activities</li> <li>• changes in your normal sleep patterns</li> <li>• becoming more irritable or aggressive than usual (for example, temper outbursts, thoughts of violence)</li> <li>• losing your appetite, becoming unusually tired</li> <li>• having trouble concentrating</li> <li>• withdrawing from family and friends</li> <li>• having thoughts about taking your own life (suicidal thoughts)</li> </ul>		X	
<b>Bone/growth changes such as premature closure of the growth plate (in growing children):</b> <ul style="list-style-type: none"> <li>• slowing or stopping of growth which may affect height</li> <li>• differences in leg length which may affect walking</li> </ul>		X	
<b>Anemia</b> (decreased number of red blood cells): fatigue, loss of energy or weakness, looking pale, shortness of breath		X	
<b>Cellulitis</b> (skin infection): pain, tenderness, swelling, redness of the skin		X	
<b>Conjunctivitis</b> (eye infection): itchy, red eyes with discharge, swelling		X	

If you have a troublesome symptom or side effect that is not listed here or becomes bad enough to interfere with your daily activities, tell your healthcare professional.

### **Reporting Side Effects**

You can report any suspected side effects associated with the use of health products to Health Canada by:

- Visiting the Web page on Adverse Reaction Reporting (<https://www.canada.ca/en/health-canada/services/drugs-health-products/medeffect-canada.html>) for information on how to report online, by mail or by fax; or
- Calling toll-free at 1-866-234-2345.

*NOTE: Contact your health professional if you need information about how to manage your side effects. The Canada Vigilance Program does not provide medical advice.*

### **Storage:**

Store at room temperature (15° to 30°C). Keep blister strip in the carton in order to protect from light.

Keep out of reach and sight of children.

### **If you want more information about SOHONOS:**

- Talk to your healthcare professional
- Find the full product monograph that is prepared for healthcare professionals and includes this Patient Medication Information by visiting the Health Canada website: <https://www.canada.ca/en/health-canada/services/drugs-health-products/drug-products/drug-product-database.html>; the manufacturer's website [www.ipсен.ca](http://www.ipсен.ca), or by calling 1-855-215-2288.

This leaflet was prepared by Ipsen Biopharmaceuticals Canada Inc.

Last Revised JAN 20, 2022