

PRODUCT MONOGRAPH

INCLUDING PATIENT MEDICATION INFORMATION

PrSULFAMETHOXAZOLE AND TRIMETHOPRIM INJECTION, BP

sulfamethoxazole (80 mg/mL) + trimethoprim (16 mg/mL)
sterile solution for the preparation of intravenous infusions

Antibacterial Agent

Juno Pharmaceuticals Corp
402-2233 Argentia Road,
Mississauga, ON, L5N 2X7

Date of Preparation: August 18, 2022

Control Number: 258006

TABLE OF CONTENTS

RECENT MAJOR LABEL CHANGES	Error! Bookmark not defined.
TABLE OF CONTENTS	2
PART I: HEALTH PROFESSIONAL INFORMATION	4
1 INDICATIONS	4
1.1 Pediatrics (2 months – 12 years of age).....	5
1.2 Geriatrics	5
2 CONTRAINDICATIONS	5
3 SERIOUS WARNINGS AND PRECAUTIONS BOX	6
4 DOSAGE AND ADMINISTRATION	6
4.1 Dosing Considerations	6
4.2 Recommended Dose and Dosage Adjustment	6
4.3 Reconstitution	8
4.4 Administration	9
5 OVERDOSAGE	9
6 DOSAGE FORMS, STRENGTHS, COMPOSITION AND PACKAGING	9
7 WARNINGS AND PRECAUTIONS	10
7.1 Special Populations.....	15
7.1.1 Pregnant Women	15
7.1.2 Breast-feeding	15
7.1.3 Pediatrics (2 months – 12 years of age).....	15
7.1.4 Geriatrics	16
8 ADVERSE REACTIONS	16
8.1 Adverse Reaction Overview	16
9 DRUG INTERACTIONS	18
9.1 Drug-Behavioural Interactions.....	18
9.2 Drug-Drug Interactions	18
9.3 Drug-Herb Interactions.....	20
9.4 Drug-Food Interactions.....	20
9.5 Drug-Laboratory Test Interactions	20
10 ACTION AND CLINICAL PHARMACOLOGY	20
10.1 Mechanism of Action.....	20
10.2 Pharmacokinetics.....	21
11 STORAGE, STABILITY AND DISPOSAL	23
PART II: SCIENTIFIC INFORMATION	24
13 PHARMACEUTICAL INFORMATION	24
14 CLINICAL TRIALS	25
15 MICROBIOLOGY	25

16	NON-CLINICAL TOXICOLOGY	30
17	SUPPORTING PRODUCT MONOGRAPHS	33
	PATIENT MEDICATION INFORMATION	34

PART I: HEALTH PROFESSIONAL INFORMATION

1 INDICATIONS

Sulfamethoxazole and Trimethoprim Injection, BP has been effective in the treatment of infections associated with the following gram-positive and gram-negative organisms:

Gram-Negative Organisms

Haemophilus influenzae
Neisseria gonorrhoeae
Escherichia coli
Klebsiella species
Enterobacter (Aerobacter) aerogenes
Proteus mirabilis
Proteus vulgaris
Salmonella species
Shigella species
Vibrio cholerae

Gram-Positive Organisms

Streptococcus pyogenes
Streptococcus viridans
Staphylococcus albus
Staphylococcus aureus
Diplococcus pneumoniae

Other Organisms

Brucella melitensis
Nocardia asteroides
Nocardia brasiliensis
Paracoccidioides brasiliensis
Pneumocystis jiroveci
Streptomyces somaliensis

Sensitivity tests should be performed wherever possible to determine choice of therapy. These tests should be repeated if there is a failure to respond, relapse or early recurrence.

Sulfamethoxazole and Trimethoprim Injection, BP may be indicated for the following infections when caused by susceptible strains of the above organisms.

Urinary Tract Infections:

Treatment of acute uncomplicated urinary tract infections*. It is recommended that initial episodes of uncomplicated urinary tract infections be treated with a single effective antibacterial agent rather than the combination.

Upper and Lower Respiratory Tract Infections:

Treatment of acute exacerbations of chronic bronchitis.

Treatment of *Pneumocystis jiroveci* pneumonia*. Sulfamethoxazole and Trimethoprim Injection, BP is also indicated in the treatment of infants and children with a diagnosis of *Pneumocystis jiroveci* pneumonitis, especially if they are immunosuppressed.

Gastrointestinal Tract Infections:

Treatment of cholera, as an adjunct to fluid and electrolyte replacement, when the organism has been shown to be sensitive *in vitro*.

Treatment of bacillary dysentery*.

Other Infections:

Treatment of nocardiosis*. Brucellosis (second line therapy), when used in combination with gentamicin or rifampicin.

Sulfamethoxazole and Trimethoprim Injection, BP is not indicated in infections associated with *Pseudomonas*, *Mycoplasma*, nor when the infection is caused by a virus.

This drug has not yet been fully evaluated in streptococcal infections.

*Sulfamethoxazole and trimethoprim have been investigated clinically in these indications.

To reduce the development of drug-resistant bacteria and maintain the effectiveness of Sulfamethoxazole and Trimethoprim Injection, BP and other antibacterial drugs, Sulfamethoxazole and Trimethoprim Injection, BP should be used only to treat infections that are proven or strongly suspected to be caused by susceptible bacteria. When culture and susceptibility information are available, they should be considered in selecting or modifying antibacterial therapy. In the absence of such data, local epidemiology and susceptibility patterns may contribute to the empiric selection of therapy.

1.1 Pediatrics (2 months – 12 years of age)

Pediatrics: Based on the data submitted and reviewed by Health Canada, safety and efficacy of trimethoprim-sulfamethoxazole in pediatric patients has been established; therefore, Health Canada has authorized an indication for pediatric use.

However, Sulfamethoxazole and Trimethoprim Injection, BP is not authorized for pediatric patients younger than 2 months of age (see [2 CONTRAINDICATIONS](#)).

1.2 Geriatrics

Geriatrics: Based on the data submitted and reviewed by Health Canada, the safety and efficacy of trimethoprim-sulfamethoxazole injection in geriatric patients has been established; therefore, Health Canada has authorized an indication for geriatric use. However, there may be an increased risk of severe adverse reactions in elderly patients, particularly when complicating conditions exist (see [7 Warnings and Precautions](#) and [8 Adverse Reactions](#)). Appropriate dosage adjustments should be made (see [4 Dosage and Administration](#)).

2 CONTRAINDICATIONS

Sulfamethoxazole and Trimethoprim Injection, BP is contraindicated in patients who are hypersensitive to this drug or to any ingredient in the formulation, including any non-medicinal ingredient, or component of the container. For a complete listing, see Dosage Forms, Strengths, Composition and Packaging.

Sulfamethoxazole and Trimethoprim Injection, BP is contraindicated in patients with a history of drug-induced immune thrombocytopenia, with use of trimethoprim and/or sulfonamides, and in

patients with documented megaloblastic anemia due to folate deficiency, evidence of marked parenchymal damage, or blood dyscrasias.

Sulfamethoxazole and Trimethoprim Injection, BP is contraindicated in patients with marked renal impairment where repeated measurements monitoring plasma drug concentrations cannot be performed (see also [7 Warnings and Precautions](#)).

Sulfamethoxazole and Trimethoprim Injection, BP is contraindicated in pregnant patients and in nursing mothers, because sulfonamides pass the placenta and are excreted in the milk and may cause kernicterus.

Sulfamethoxazole and Trimethoprim Injection, BP is contraindicated in premature babies and full term infants less than two months of age.

Sulfamethoxazole and Trimethoprim Injection, BP should not be given to patients with acute porphyria.

Sulfamethoxazole and Trimethoprim Injection, BP must not be given in combination with dofetilide (See [9 DRUG INTERACTIONS](#)).

3 **SERIOUS WARNINGS AND PRECAUTIONS BOX**

Serious Warnings and Precautions

- Sulfamethoxazole and Trimethoprim Injection, BP can cause severe skin reactions that may be life threatening including Steven-Johnson syndrome (SJS), toxic epidermal necrolysis (TEN), drug reaction with eosinophilia and systemic symptoms (DRESS), acute febrile neutrophilic dermatosis (Sweet's syndrome) and acute generalised exanthematous pustulosis (AGEP) (see [7 WARNINGS AND PRECAUTIONS, Skin](#)).
- Sulfamethoxazole and Trimethoprim Injection, BP can cause fulminant hepatic necrosis that may be life-threatening (see [7 WARNINGS AND PRECAUTIONS, Hepatic/Biliary/Pancreatic](#)).
- Sulfamethoxazole and Trimethoprim Injection, BP can cause agranulocytosis (see [7 WARNINGS AND PRECAUTIONS, Hematologic](#)).
- Sulfamethoxazole and Trimethoprim Injection, BP can cause aplastic anemia (see [7 WARNINGS AND PRECAUTIONS, Hematologic](#)).
- Sulfamethoxazole and Trimethoprim Injection, BP can cause immune thrombocytopenia that may be life-threatening (see [7 WARNINGS AND PRECAUTIONS, Immune](#)).
- Sulfamethoxazole and Trimethoprim Injection, BP can cause an allergic reaction in the lungs and in the airways that may be life-threatening (see [7 WARNINGS AND PRECAUTIONS, Respiratory](#)).

4 **DOSAGE AND ADMINISTRATION**

4.1 **Dosing Considerations**

Sulfamethoxazole and Trimethoprim Injection, BP may be used only in patients who are unable to take oral medication or where there is a need for rapid attainment of high serum concentrations. Oral treatment should be substituted as soon as possible.

4.2 **Recommended Dose and Dosage Adjustment**

Serious Systemic Infections

Adults

The intravenous dosage of Sulfamethoxazole and Trimethoprim Injection, BP depends on the severity of the infection. A dose of 160 to 240 mg trimethoprim + 800 to 1200 mg sulfamethoxazole may be given every 6, 8 or 12 hours. This dose must be properly diluted (see Pharmaceutical Information) and infused over a period of one-half to one hour.

Children (2 months–12 years of age):

The recommended daily dosage for children is 5 to 10 mg trimethoprim/kg body weight/day and 25 to 50 mg sulfamethoxazole/kg body weight/day. This daily dosage must be properly diluted and administered in equally divided doses by infusion over a period of one-half to one hour. Sulfamethoxazole and Trimethoprim Injection, BP is not recommended for pediatric patients younger than 2 months of age ([see 1.1 Pediatrics](#)).

Volume of Undiluted Sulfamethoxazole and Trimethoprim Injection, BP per Body Weight* (conversion factor 0.31 to 0.63 mL/kg)				
Body Weight (kg)	Volume of Undiluted Sulfamethoxazole and Trimethoprim Injection, BP for Infusion (mL)			
	Total Daily Dose	Dose Every		
		12 Hours (b.i.d.)	8 Hours (t.i.d.)	6 Hours (q.i.d.)
5	1.6 - 3.2	0.8 - 1.6	0.5 - 1.1	0.4 - 0.8
10	3.1 - 6.3	1.6 - 3.2	1.0 - 2.1	0.8 - 1.6
20	6.2 - 12.6	3.1 - 6.3	2.1 - 4.2	1.6 - 3.2
40	12.4 - 25.2	6.2 - 12.6	4.1 - 8.4	3.1 - 6.3
60	18.6 - 37.8	9.3 - 18.9	6.2 - 12.6	4.7 - 9.5

* Sulfamethoxazole and Trimethoprim Injection, BP must be properly diluted (see Pharmaceutical Information) and administered in equally divided doses.

The daily dosage of Sulfamethoxazole and Trimethoprim Injection, BP must be given in two equally divided doses.

Treatment should be continued until the patient has been symptom free for two days; the majority will require treatment for at least five days.

Other diseases, including certain tropical diseases rarely seen in Canada have also been successfully treated with Sulfamethoxazole and Trimethoprim Injection, BP. The duration of treatment is as follows:

Disease	Duration
Cholera	7 days
Nocardiosis	12 weeks
Brucellosis	2 weeks to 3 months

Pneumocystis jiroveci Pneumonitis

Children and Adults

The recommended daily intravenous dosage is 20 mg trimethoprim/kg body weight + 100 mg sulfamethoxazole/kg body weight. This daily dosage is to be divided into four equal doses infused over a period of one-half to one hour, at six-hour intervals, until oral therapy can be instituted.

Volume of Undiluted Sulfamethoxazole and Trimethoprim Injection, BP per Body Weight* (conversion factor 1.25 mL/kg)		
Body Weight (kg)	Volume of Undiluted Sulfamethoxazole and Trimethoprim Injection, BP (mL)	
	Total Daily Dose	Dose Every 6 Hours (q.i.d.)
5	6.3	1.6
10	12.5	3.1
20	25.0	6.3
40	50.0	12.5
60	75.0	18.8
80	100.0	25.0

* Sulfamethoxazole and Trimethoprim Injection, BP must be properly diluted (see Pharmaceutical Information) and administered at six-hour intervals.

Therapy should be continued for a total treatment period of at least two weeks. The aim is to obtain peak plasma or serum levels of trimethoprim of greater than or equal to 5 micrograms/mL (see [8 Adverse Reactions](#)).

Patients with Impaired Renal Function

When renal function is impaired, a reduced dosage should be employed using the following table:

Creatinine Clearance (mL/min)	Recommended Dose Regimen
Above 25	Usual standard regimen
15-25	Half the usual regimen
Below 15	Use not recommended

4.3 Reconstitution

CAUTION: Direct intravenous injection is not recommended. Sulfamethoxazole and Trimethoprim Injection, BP must be diluted in one of the following diluents:

- Ringer's Solution
- Sodium chloride 0.9% Solution
- Dextrose 5% Solution
- Dextrose 10% Solution
- 10% Dextran 40 in Sodium chloride 0.9% Solution
- 10% Dextran 40 in Dextrose 5% Solution

	Volume of Diluent (mL)	Approx. Available Volume(mL)	Nominal Concentration (mg/mL)	
			Trimethoprim	Sulfamethoxazole
5 mL ampoule	125	130	0.62	3.1

The prepared solution must be kept at room temperature and administration started within five hours. Do not mix the prepared infusion solution with other drugs or solutions. If, upon visual inspection, there is cloudiness or evidence of precipitation after mixing, the solution should be discarded and a fresh solution prepared.

4.4 Administration

Sulfamethoxazole and Trimethoprim Injection, BP for infusion is for administration ONLY by the I.V. route and must be diluted before administration (see Reconstitution).

The duration of the infusion should be approximately one-half to one hour, but this needs to be balanced against the fluid requirements of the patient

5 OVERDOSAGE

Symptoms and Treatment

Acute

The amount of a single dose of sulfamethoxazole and trimethoprim that is either associated with symptoms of overdose or is likely to be life-threatening has not been reported. Signs and symptoms of overdose reported with sulfonamides include anorexia, colic, nausea, vomiting, dizziness, headache, drowsiness, and unconsciousness. Pyrexia, hematuria, and crystalluria may be noted. Blood dyscrasias and jaundice are potential late manifestations of overdose. Signs of acute overdose with trimethoprim include nausea, vomiting, dizziness, headache, mental depression, confusion, and bone marrow depression.

General principles of treatment include the forcing oral fluids; and the administration of intravenous fluids if urine output is low and renal function is normal. Acidification of the urine will increase renal elimination of trimethoprim. Inducing diuresis plus alkalinisation of urine will enhance the elimination of sulfamethoxazole. Alkalinisation will reduce the rate of elimination of trimethoprim. The patient should be monitored with blood counts and appropriate blood chemistries, including electrolytes. If a significant blood dyscrasia or jaundice occurs, specific therapy should be instituted for these complications. Peritoneal dialysis is not effective and hemodialysis is only moderately effective in eliminating trimethoprim and sulfamethoxazole.

There is no known antidote for sulfonamide poisoning; however, calcium folinate (leucovorin), 3 to 6 mg I.M. for 5 to 7 days, is an effective antidote for adverse effects in the hemopoietic system caused by trimethoprim.

Chronic

Use of sulfamethoxazole-trimethoprim at high doses and/or for extended periods of time may cause confusion and bone marrow depression manifested as thrombocytopenia, leukopenia, and/or megaloblastic anemia. If signs of bone marrow depression occur, the patient should be given leucovorin; 5 to 15 mg leucovorin daily has been recommended by some investigators.

For management of a suspected drug overdose, contact your regional poison control centre.

6 DOSAGE FORMS, STRENGTHS, COMPOSITION AND PACKAGING

Table 1 – Dosage Forms, Strengths, Composition and Packaging.

Route of Administration	Dosage Form Strength/Composition	Non-medicinal Ingredients
Intravenous	Strong Sterile Co-Trimoxazole Solution: Each mL of Sulfamethoxazole and Trimethoprim Injection, BP contains 16 mg trimethoprim, 80 mg sulfamethoxazole.	Propylene glycol (450 mg), tromethamine (14 mg), sodium hydroxide (12.65 mg and for pH adjustment), ethanol 96% (0.132 mL), and water for injection (q.s).

Sulfamethoxazole and Trimethoprim Injection, BP contains trimethoprim (16 mg/mL) and sulfamethoxazole (80 mg/mL). Ampoules of 5 mL packaged 10 ampoules per carton. Sulfamethoxazole and Trimethoprim Injection, BP is preservative free.

7 WARNINGS AND PRECAUTIONS

General

Sulfamethoxazole and Trimethoprim Injection, BP should only be used where, in the judgement of the physician, the benefit of treatment outweighs any possible risks; consideration should be given to the use of a single effective antibacterial agent.

The possibility of superinfection with a non-sensitive organism should be borne in mind.

Local irritation and inflammation due to extravascular infiltration of the infusion has been observed with sulfamethoxazole-trimethoprim. If these occur, the infusion should be discontinued and restarted at another site.

Life threatening adverse reaction

Fatalities associated with the administration of sulfonamides and sulfamethoxazole-trimethoprim, although rare, have occurred due to severe reactions, including Stevens-Johnson syndrome, toxic epidermal necrolysis, drug reaction with eosinophilia and systemic symptoms (DRESS), acute febrile neutrophilic dermatosis (Sweet's syndrome), acute generalised exanthematous pustulosis (AGEP), fulminant hepatic necrosis, agranulocytosis, immune thrombocytopenia, aplastic anemia, other blood dyscrasias, and hypersensitivity of the respiratory tract. (see [3 SERIOUS WARNINGS AND PRECAUTIONS BOX](#)).

Circulatory shock with fever, severe hypotension, and confusion requiring intravenous fluid resuscitation and vasopressors has occurred within minutes to hours of re-challenge with trimethoprim-sulfamethoxazole in patients with history of recent (days to weeks) exposure to sulfamethoxazole-trimethoprim.

Driving and Operating Machinery

Due caution should be exercised when driving or operating a vehicle or potentially dangerous machinery.

Endocrine and Metabolism

Metabolic acidosis

Trimethoprim-sulfamethoxazole has been associated with metabolic acidosis when other possible underlying causes have been excluded. Close monitoring is always advisable when metabolic acidosis is suspected.

Fluid overload

Fluid Overload is possible, especially when very high doses are being administered to patients with underlying cardio-pulmonary disease.

Urinary output

An adequate urinary output should be maintained at all times. Evidence of crystalluria *in vivo* is rare, although sulfonamide crystals have been noted in cooled urine from treated patients. In patients suffering from hypoalbuminaemia the risk may be increased.

Patients with hyperkalaemia and hyponatraemia

Caution should be exercised in administering trimethoprim to patients at risk of hyperkalemia and hyponatremia. Serum potassium and sodium and renal function should be closely monitored, and dosage should be adjusted for renal function (see Warnings and Precautions – Renal and Dosage and Administration).

The risk factors for hyperkalemia are high trimethoprim dosage (20 mg/kg/day), renal insufficiency (serum creatinine \geq 1.2 mg/dl), hypoaldosteronism, older age, dietary potassium and other drugs that impair potassium excretion. The likely mechanism is via trimethoprim inhibition of sodium channels in the distal nephron, similar to that of the potassium-sparing diuretic amiloride.

Hyperkalemia is generally reversible on discontinuation of trimethoprim. In patients presented with hyperkalemia due to sulfamethoxazole-trimethoprim injection, Sulfamethoxazole and Trimethoprim Injection, BP can be discontinued and appropriate standard potassium-lowering therapy instituted.

Phenylketonuric Patients

Trimethoprim has been noted to impair phenylalanine metabolism but this is of no significance in phenylketonuric patients on appropriate dietary restriction.

Gastrointestinal

***Clostridium difficile* – Associated Disease (CDAD)**

Clostridium difficile-associated disease (CDAD) has been reported with use of many antibacterial agents, including sulfamethoxazole-trimethoprim. CDAD may range in severity from mild diarrhea to fatal colitis. It is important to consider this diagnosis in patients who present with diarrhea, or symptoms of colitis, pseudomembranous colitis, toxic megacolon, or perforation of colon subsequent to the administration of any antibacterial agent. CDAD has been reported to occur over 2 months after the administration of antibacterial agents.

Treatment with antibacterial agents may alter the normal flora of the colon and may permit overgrowth of *C. difficile*. *Clostridium difficile* produces toxins A and B, which contribute to the development of CDAD. CDAD may cause significant morbidity and mortality. CDAD can be refractory to antimicrobial therapy. If the diagnosis of CDAD is suspected or confirmed, appropriate

therapeutic measures should be initiated. Mild cases of CDAD usually respond to discontinuation of antibacterial agents not directed against *C. difficile*. In moderate to severe cases, consideration should be given to management with fluids and electrolytes, protein supplementation, and treatment with an antibacterial agent clinically effective against *C. difficile*. Surgical evaluation should be instituted as clinically indicated, as surgical intervention may be required in certain severe cases (see [8 Adverse Reactions](#)).

Patients with gastrointestinal tract infection

Clinicians should be aware that first line therapy in the management of all patients with diarrheal disease is the maintenance of adequate hydration.

Hematology

Sulfamethoxazole and Trimethoprim Injection, BP can cause agranulocytosis, where the number of white cells in the blood becomes dangerously low. Symptoms of this can include pyrexia, chills, oropharyngeal pain, asthenia. Patients can also experience fast heart rate or fast breathing. If patients experience any of these symptoms, the treatment with Sulfamethoxazole and Trimethoprim Injection, BP should be immediately stopped.

Sulfamethoxazole and Trimethoprim Injection, BP can cause aplastic anemia, where the bone marrow is unable to make enough blood cells from being damaged. Symptoms of this can include fatigue, dyspnoea, pallor, contusion, pyrexia, chills, oropharyngeal pain, and malaise. If patients experience any of these symptoms, the treatment with Sulfamethoxazole and Trimethoprim Injection, BP should be immediately stopped

Folate

Regular monthly blood counts are advisable when trimethoprim-sulfamethoxazole is given for long periods, or to folate deficient patients or to the elderly, since there exists a possibility of asymptomatic changes in haematological laboratory indices due to lack of available folate. Supplementation with folic acid may be considered during treatment but this should be initiated with caution due to possible interference with antimicrobial efficacy (see *Interactions*). Changes indicative of folic acid impairment have, in certain specific situations, been reversed by folic acid therapy.

Special care should be exercised when treating suspected folate-deficient patients; folate supplementation should be considered.

Patients with serious haematological disorders

Except under careful supervision trimethoprim-sulfamethoxazole should not be given to patients with serious hematological disorders (see [8 Adverse Reactions](#)).

Hepatic/ Biliary/ Pancreatic:

Sulfamethoxazole and Trimethoprim Injection, BP can cause fulminant hepatic necrosis that may be life-threatening. Symptoms include jaundice, upper right abdominal pain, abdominal distension, nausea and vomiting. If patients experience any of these symptoms, the treatment with Sulfamethoxazole and Trimethoprim Injection, BP should be immediately stopped.

Sulfamethoxazole and Trimethoprim Injection, BP should be given with caution to patients with impaired renal or hepatic function, to those with possible folate deficiency (e.g., the elderly, chronic alcoholics, rheumatoid arthritics, patients receiving anticonvulsant therapy, patients with malabsorption syndrome, and patients in malnutrition states), and to those with severe allergy or bronchial asthma.

Immune

Use in the Treatment of and Prophylaxis for Pneumocystis jiroveci Pneumonia in Patients with Acquired Immunodeficiency Syndrome (AIDS):

The incidence of side effects, particularly rash, severe hypersensitivity reactions, fever, leukopenia, neutropenia, thrombocytopenia and elevated aminotransferase (transaminase) values in AIDS patients who are being treated with sulfamethoxazole-trimethoprim for Pneumocystis jiroveci pneumonia (PJP) has been reported to be greatly increased compared with the incidence normally associated with the use of sulfamethoxazole-trimethoprim injection in non-AIDS patients. If signs of bone marrow depression occur, the patient should be given calcium folinate supplementation (5-10 mg/day). The incidence of hyperkalemia and hyponatremia appears to be increased in AIDS patients receiving sulfamethoxazole-trimethoprim. Adverse effects are generally less severe in patients receiving for prophylaxis. A history of mild intolerance to sulfamethoxazole-trimethoprim in AIDS patients does not appear to predict intolerance of subsequent secondary prophylaxis. However, if a patient develops skin rash or any sign of adverse reaction, therapy with sulfamethoxazole-trimethoprim should be re-evaluated (see Warnings and Precautions). Rhabdomyolysis has been reported in HIV positive patients receiving trimethoprim-sulfamethoxazole for prophylaxis or treatment of PJP. In some cases, rhabdomyolysis led to acute renal failure requiring emergency dialysis.

Severe hypersensitivity reactions have also been reported in HIV-infected patients on re-exposure to sulfamethoxazole-trimethoprim, sometimes after a dosage interval of a few days. Concomitant administration of intravenous diphenhydramine may permit continued infusion.

The concomitant use of leucovorin with sulfamethoxazole-trimethoprim for the acute treatment of Pneumocystis jiroveci pneumonia in patients with HIV infection was associated with increased rates of treatment failure and morbidity in a placebo-controlled study (see 9.4 Drug-Drug Interactions).

Sulfamethoxazole and Trimethoprim Injection, BP can cause immune thrombocytopenia that may be life-threatening. Symptoms of this include being easily bruised, a rash on the skin that appears tiny pinpoint-sized reddish or purple spots, usually on the lower legs, gingival bleeding or epistaxis, and blood in urine or haematochezia. If patients experience any of these symptoms, the treatment with Sulfamethoxazole and Trimethoprim Injection, BP should be immediately stopped.

Monitoring and Laboratory Tests

Sulfamethoxazole and Trimethoprim Injection, BP may affect the results of thyroid function tests.

Renal

In patients with renal impairment, a reduced or less frequent dosage is recommended in order to avoid accumulation of trimethoprim in the blood (see Dosage and Administration). Non-ionic diffusion is the main factor in the renal handling of trimethoprim, and as renal failure advances, trimethoprim excretion decreases. For such patients, serum assays are necessary.

Sulfamethoxazole and Trimethoprim Injection, BP should not be used when the serum creatinine level is above 2 mg per 100 mL, in order to avoid possible permanent impairment of renal function.

Patients with glucose-6-phosphate dehydrogenase deficiency

In glucose-6-phosphate dehydrogenase-deficient individuals, hemolysis may occur. This reaction is frequently dose-related.

Patients with or at risk of acute porphyria

The administration of Sulfamethoxazole and Trimethoprim Injection, BP to patients known or suspected to be at risk of acute porphyria should be avoided. Both trimethoprim and sulfonamides (although not specifically sulfamethoxazole) have been associated with clinical exacerbation of acute porphyria.

Functional inhibition of the renal tubular secretion of creatinine may produce a spurious fall in the estimated rate of creatinine clearance.

Respiratory:

Sulfamethoxazole and Trimethoprim Injection, BP can cause an allergic reaction in the lungs and in the airways, where the airways can close up and make breathing difficult and can be life-threatening if the person does not get medical help. Symptoms include dyspnoea, cough, wheezing, and chest discomfort. If patients experience any of these symptoms, the treatment with Sulfamethoxazole and Trimethoprim Injection, BP should be immediately stopped.

Acute and delayed lung injury, cough, shortness of breath, and pulmonary infiltrates potentially representing hypersensitivity reactions of the respiratory tract have been reported in association with trimethoprim-sulfamethoxazole treatment.

Other severe pulmonary adverse reactions occurring within days to week of trimethoprim-sulfamethoxazole initiation and resulting in prolonged respiratory failure requiring mechanical ventilation or extracorporeal membrane oxygenation (ECMO), lung transplantation or death have also been reported in patients and otherwise healthy individuals treated with trimethoprim-sulfamethoxazole products.

Skin

Sulfamethoxazole and Trimethoprim Injection, BP can cause severe skin reactions that may be life-threatening, including: Steven-Johnson syndrome (SJS), toxic epidermal necrolysis (TEN), drug reaction with eosinophilia and systemic symptoms (DRESS), acute febrile neutrophilic dermatosis (Sweet's syndrome) and acute generalised exanthematous pustulosis (AGEP). Symptoms include: rash erythematous, blister, skin exfoliation, pyrexia, body aches, eosinophilia, blisters and sores or ulcers on mucous membranes (mouth, nose and genitals), ocular hyperaemia and eye swelling. If patients experience any of these symptoms, the treatment with Sulfamethoxazole and Trimethoprim Injection, BP should be immediately stopped.

Patients should be advised of the signs and symptoms and advised to monitor closely for skin reactions. The highest risk for occurrence of SJS or TEN is within the first weeks of treatment and of DRESS is within the first two to eight weeks after drug administration. The best results in managing SJS, TEN, DRESS, AGEP and Sweet's syndrome come from early diagnosis and immediate discontinuation of any suspect drug. Early withdrawal is associated with a better prognosis. If the patient has developed SJS, TEN, DRESS, AGEP and Sweet's syndrome with the

use of trimethoprim-sulfamethoxazole, trimethoprim-sulfamethoxazole must not be re-started in this patient at any time.

Susceptibility /Resistance

Development of Drug Resistant Bacteria

Prescribing Sulfamethoxazole and Trimethoprim Injection, BP in the absence of a proven or strongly suspected bacterial infection is unlikely to provide benefit to the patient and risks the development of drug-resistant bacteria.

Treatment of streptococcal pharyngitis due to Group A beta-haemolytic streptococci

Sulfamethoxazole-trimethoprim **should not be used in the treatment of streptococcal pharyngitis**. Clinical studies have documented that patients with group A β -hemolytic streptococcal tonsillopharyngitis have a greater incidence of bacteriologic failure when treated with trimethoprim-sulfamethoxazole than to those patients treated with penicillin, as evidenced by failure to eradicate this organism from the tonsillopharyngeal area.

7.1 Special Populations

7.1.1 Pregnant Women

Trimethoprim and sulfamethoxazole cross the placenta and their safety in human pregnancy has not been established. Trimethoprim is a folate antagonist and, in animal studies, both agents have been shown to cause fetal abnormalities. At doses in excess of the recommended human therapeutic dose, trimethoprim and sulfamethoxazole have been reported to cause cleft palate and other fetal abnormalities in rats, findings typical of a folate antagonist. Effects with trimethoprim were preventable by administration of dietary folate. In rabbits, fetal loss was seen at doses of trimethoprim in excess of human therapeutic doses. Case-control studies have shown that there may be an association between exposure to folate antagonists and birth defects in humans. Therefore, Sulfamethoxazole and Trimethoprim Injection, BP should be avoided in pregnancy, particularly in the first trimester, unless the potential benefit to the mother outweighs the potential risk to the fetus; folate supplementation should be considered if Sulfamethoxazole and Trimethoprim Injection, BP is used in pregnancy.

Sulfamethoxazole competes with bilirubin for binding to plasma albumin. As significant maternally derived drug levels persist for several days in the newborn, there may be a risk of precipitating or exacerbating neonatal hyperbilirubinaemia, with an associated theoretical risk of kernicterus, when sulfamethoxazole-trimethoprim is administered to the mother near the time of delivery. This theoretical risk is particularly relevant in infants at increased risk of hyperbilirubinaemia, such as those who are preterm and those with glucose-6-phosphatedehydrogenase deficiency.

7.1.2 Breast-feeding

Trimethoprim and sulfamethoxazole are excreted in breast milk. Administration of Sulfamethoxazole and Trimethoprim Injection, BP should be avoided in late pregnancy and in lactating mothers where the mother or infant has, or is at particular risk of developing hyperbilirubinaemia. Additionally, administration of Sulfamethoxazole and Trimethoprim Injection, BP should be avoided in infants younger than eight weeks in view of predisposition of young infants to hyperbilirubinaemia.

7.1.3 Pediatrics (2 months – 12 years of age)

Based on the data submitted and reviewed by Health Canada, the safety and efficacy of trimethoprim-sulfamethoxazole in pediatric patients have been established; therefore, Health Canada has authorized an indication for pediatric use.

However, Sulfamethoxazole and Trimethoprim Injection, BP is not recommended for pediatric patients younger than 2 months of age (see [2 Contraindications](#)).

7.1.4 Geriatrics

Based on the data submitted and reviewed by Health Canada, the safety and efficacy of trimethoprim-sulfamethoxazole Injection in geriatric patients have been established; therefore, Health Canada has authorized an indication for geriatric use.

However, there may be an increased risk of severe adverse reactions in elderly patients, particularly when complicating conditions exist, e.g., impaired kidney and/or liver function, or concomitant use of other drugs. Severe skin reactions, or generalized bone marrow suppression (see Warnings and Precautions and Adverse Reactions), or a specific decrease in platelets (with or without purpura) are the most frequently reported severe adverse reactions in elderly patients. In those concurrently receiving certain diuretics, primarily thiazides, an increased incidence of thrombocytopenia with or without purpura has been reported. Appropriate dosage adjustments should be made for patients with impaired kidney function (see [4 Dosage and Administration](#)).

Close supervision is recommended when Sulfamethoxazole and Trimethoprim Injection, BP is used in elderly patients or in patients taking high doses of Sulfamethoxazole and Trimethoprim Injection, BP as these patients may be more susceptible to hyperkalemia and hyponatremia.

Special care should be exercised when treating the elderly or suspected folate-deficient patients; folate supplementation should be considered.

8 ADVERSE REACTIONS

8.1 Adverse Reaction Overview

The most common adverse effects are hyperkalemia, anorexia, monilial overgrowth, headache, local thrombophlebitis at the site of injection, gastrointestinal disturbances (nausea, vomiting, diarrhea) and allergic skin reactions (such as rash and urticaria). FATALITIES ASSOCIATED WITH THE ADMINISTRATION OF SULFONAMIDES AND SULFAMETHOXAZOLE-TRIMETHOPRIM ALTHOUGH RARE, HAVE OCCURRED DUE TO SEVERE REACTIONS, INCLUDING STEVENS-JOHNSON SYNDROME, DRUG REACTION WITH EOSINOPHILIA AND SYSTEMIC SYMPTOMS (DRESS), TOXIC EPIDERMAL NECROLYSIS, FULMINANT HEPATIC NECROSIS, AGRANULOCYTOSIS, APLASTIC ANEMIA, OTHER BLOOD DYSCRASIAS, AND HYPERSENSITIVITY OF THE RESPIRATORY TRACT (See [3 Serious Warnings and Precautions Box](#) and [7 Warnings And Precautions](#)).

General

Weakness, fatigue, insomnia, vision troubles, alopecia, epistaxis, local thrombophlebitis at the site of injection, edema. Monilial overgrowth is common.

Allergic

Anaphylactic reaction, allergic myocarditis, erythema multiforme, toxicoderma, exfoliative dermatitis, angioedema, pyrexia, chills, hypersensitivity vasculitis resembling Henoch-Schönlein

purpura, serum sickness, serum sickness-like syndrome, generalized allergic reactions, generalized skin eruptions, fixed drug eruption, photosensitivity, conjunctival and scleral injection, pruritus, urticaria, and rash. In addition, periarteritis nodosa and systemic lupus erythematosus and anaphylactoid reactions (sweating and collapse) have been reported.

Severe hypersensitivity reactions associated with PJP including rash, pyrexia, neutropenia, thrombocytopenia, hepatic enzyme increased, rhabdomyolysis, hyperkalemia, hyponatremia have been reported at the high dosages used for PJP management, necessitating cessation of therapy.

Rhabdomyolysis has been reported in HIV positive patients receiving trimethoprim-sulfamethoxazole for prophylaxis or treatment of PJP.

Cardiovascular

QT interval prolongation (ventricular tachycardia).

Endocrine and Metabolism

The sulfonamides bear certain chemical similarities to some goitrogens, diuretics (acetazolamide and the thiazides), and oral hypoglycemic agents. Cross-sensitivity may exist with these agents. Diuresis and hypoglycemia have occurred rarely in patients receiving sulfonamides.

Decreased appetite, metabolic acidosis, renal tubular acidosis, hyperkalemia, hyponatremia, hypoglycemia (See Also Warnings and Precautions).

Gastrointestinal

Pseudomembranous enterocolitis, pancreatitis, stomatitis, glossitis, dry mouth, nausea, vomiting, pyrosis, gastric intolerance, gastritis or gastroenteritis, dyspepsia, emesis, abdominal pain, constipation, flatulence, diarrhea.

Genitourinary

Impaired renal function (sometimes reported as renal failure), interstitial nephritis, kidney changes (as indicated by abnormal elevations in blood urea nitrogen, blood non-protein nitrogen, serum creatinine and urine protein levels), toxic nephrosis with oliguria and anuria, crystalluria, hematuria, urgency, and dysuria.

Hematologic

Leukopenia, neutropenia, thrombocytopenia, megaloblastic anemia, aplastic and hemolytic anemia, methemoglobinemia, purpura, agranulocytosis, hypoprothrombinemia, eosinophilia, haemolysis in certain susceptible G-6-PD deficient patients and bone marrow depression.

Hepatic/Biliary/Pancreatic

Hepatitis, including cholestatic jaundice and hepatic necrosis, jaundice, elevation of serum transaminase, alkaline phosphatase and bilirubin.

Hepatic changes including fatalities have been recorded in at-risk patients. Cholestatic jaundice and hepatic necrosis may be fatal.

Musculoskeletal

Arthralgia, rhabdomyolysis and myalgia.

Neurologic

Aseptic meningitis, convulsions, peripheral neuritis, ataxia, tremor, vertigo, tinnitus, headache.

Aseptic meningitis was rapidly reversible on withdrawal of the drug, but recurred in a number of cases on re-exposure to either sulfamethoxazole-trimethoprim or to trimethoprim alone.

Ophthalmologic

Uveitis

Psychiatric

Hallucinations, depression, apathy, nervousness, dizziness, psychotic disorder.

Respiratory

Lung infiltration, cough, dyspnea, acute eosinophilic pneumonia, acute and delayed lung injury, interstitial lung disease, and acute respiratory failure.

Skin and subcutaneous tissue disorders

Severe cutaneous adverse reactions (SCARs). Acute febrile neutrophilic dermatosis (Sweet's syndrome) (Frequency Not Known). Acute generalized exanthematous pustulosis (AGEP) (Frequency Not Known).

Stevens-Johnson syndrome (SJS), toxic epidermal necrolysis (TEN), and drug reaction with eosinophilia and systemic symptoms (DRESS) (Frequency Not Known) have been reported to be life-threatening (see [7 Warnings and Precautions](#)).

9 DRUG INTERACTIONS

9.1 Drug-Behavioural Interactions

Patients should be instructed to maintain an adequate fluid intake in order to prevent crystalluria and stoneformation.

Patients should tell their physician of all dietary regimens and supplements.

9.2 Drug-Drug Interactions

Diuretics (thiazides): in elderly patients concurrently receiving certain diuretics, primarily thiazides, an increased incidence of thrombocytopenia with or without purpura has been reported.

Pyrimethamine: occasional reports suggest that patients receiving pyrimethamine at doses in excess of 25 mg weekly may develop megaloblastic anemia should sulfamethoxazole-trimethoprim be prescribed concurrently.

Zidovudine: in some situations, concomitant treatment with zidovudine may increase risk of hematological adverse reactions to sulfamethoxazole-trimethoprim. If concomitant treatment is necessary, consideration should be given to monitoring of hematological parameters.

Lamivudine: administration of sulfamethoxazole-trimethoprim 800 mg/160 mg causes a 40% increase in lamivudine exposure because of the trimethoprim component. Lamivudine has no effect on the pharmacokinetics of trimethoprim or sulfamethoxazole.

Warfarin: it has been reported that sulfamethoxazole-trimethoprim may prolong the prothrombin time in patients who are receiving the anticoagulant warfarin. This interaction should be kept in mind when trimethoprim-sulfamethoxazole is given to patients already on anticoagulant therapy, and the coagulation time should be reassessed.

Phenytoin: sulfamethoxazole-trimethoprim may inhibit the hepatic metabolism of phenytoin. Sulfamethoxazole-trimethoprim given at a common clinical dosage, increased the phenytoin half-life by 39% and decreased the phenytoin metabolic clearance rate by 27%. When administering these drugs concurrently, one should be alert for possible excessive phenytoin effect. Close monitoring of the patient's condition and serum phenytoin levels is advisable.

Methotrexate: sulfonamides can also displace methotrexate from plasma protein binding sites, thus increasing free methotrexate concentrations. Folate supplementation should be considered. If sulfamethoxazole-trimethoprim is considered appropriate therapy in patients receiving other anti-folate drugs, a folate supplementation should be considered.

Digoxin: concomitant use of trimethoprim with digoxin has been shown to increase plasma digoxin levels in a proportion of elderly patients.

Hyperkalaemia: caution should be exercised in patients taking any other drugs that can cause hyperkalemia, for example ACE inhibitors, angiotensin receptor blockers and potassium sparing diuretics such as spironolactone. Concomitant use of trimethoprim-sulfamethoxazole (cotrimoxazole) may result in clinically relevant hyperkalaemia.

Cyclosporin: reversible deterioration in renal function has been observed in patients treated with sulfamethoxazole-trimethoprim and cyclosporin following renal transplantation.

When trimethoprim is administered simultaneously with drugs that form cations at physiological pH, and are also partly excreted by active renal secretion (e.g., procainamide, amantadine), there is the possibility of competitive inhibition of this process which may lead to an increase in plasma concentration of one or both of the drugs.

Interaction with sulphonylurea hypoglycemic agents is uncommon but potentiation has been reported.

Rifampicin: concurrent use of rifampicin and sulfamethoxazole-trimethoprim results in a shortening of the plasma half life of trimethoprim after a period of about one week. This is not thought to be of clinical significance.

Trimethoprim is an inhibitor of cytochrome P450 2C8 enzyme and may interact with other drugs that are primarily metabolized by the 2C8 isoform. Sulfamethoxazole is an inhibitor of cytochrome P450 2C9 and may interact with other drugs that are primarily metabolized by the 2C9 isoform.

Repaglinide: trimethoprim may increase the exposure of repaglinide which may result in hypoglycaemia.

Folinic acid: folinic acid supplementation has been shown to interfere with the antimicrobial efficacy of trimethoprim-sulfamethoxazole. This has been observed in *Pneumocystis jiroveci* pneumonia prophylaxis and treatment.

Azathioprine: There are conflicting clinical reports of interactions between azathioprine and trimethoprim-sulfamethoxazole, resulting in serious haematological abnormalities.

Dofetilide

Increased plasma levels of dofetilide have been reported after co-administration of trimethoprim and dofetilide. Dofetilide can cause serious ventricular arrhythmias associated with QT interval prolongation, including torsades de pointes, which are directly related to the dofetilide plasma concentration. Concomitant administration of dofetilide and trimethoprim is contraindicated (see [2 Contraindications](#)).

Leucovorin: The concomitant use of leucovorin with sulfamethoxazole-trimethoprim for the acute treatment of *Pneumocystis jiroveci* pneumonia in patients with HIV infection was associated with increased rates of treatment failure and morbidity in a placebo-controlled study.

9.3 Drug-Herb Interactions

Interactions with herbal products have not been established.

9.4 Drug-Food Interactions

Caution should be exercised in patients following potassium enriched dietary regimens ([See 4.4 Administration](#)).

9.5 Drug-Laboratory Test Interactions

Sulfamethoxazole-trimethoprim, specifically the trimethoprim component, can interfere with a serum methotrexate assay as determined by the competitive binding protein technique (CBPA) when a bacterial dihydrofolate reductase is used as the binding protein. No interference occurs, however, if methotrexate is measured by a radioimmunoassay (RIA).


The presence of trimethoprim and sulfamethoxazole may also interfere with the Jaffé alkaline picrate reaction assay for creatinine, resulting in overestimations of about 10% in the range of normal values.

10 ACTION AND CLINICAL PHARMACOLOGY

10.1 Mechanism of Action

Sulfamethoxazole and Trimethoprim Injection, BP is an antibacterial agent with a wide spectrum of activity. It contains two active antibacterial components, sulfamethoxazole and trimethoprim, which act synergistically on many species of bacteria.

Figure 1



Sulfamethoxazole and trimethoprim act sequentially in two successive steps in the biosynthesis of nucleic acids. Trimethoprim is an inhibitor of dihydrofolate reductase, the enzyme which reduces dihydrofolic acid to its tetrahydro form. This biochemical step is essential in the production of the folate coenzymes which are involved in the biosynthesis of thymine, purine, serine and methionine. Sulfamethoxazole exerts its antibacterial activity by competing with para-aminobenzoic acid.

Most pathogenic bacteria meet their need for dihydrofolic acid by synthesizing it from para-aminobenzoic acid, pteridine and glutamic acid. Animals, in contrast, depend on exogenous sources for their needs of folic acid and do not rely upon intracellular synthesis.

Under usual circumstances, sulfamethoxazole or trimethoprim acting alone do not produce complete block in this biosynthesis of nucleic acids. Instead, they cause sufficient reduction in the synthesis of folate coenzymes to produce bacteriostasis. When the two agents act together, the superimposition of their effects produces a complete block in the synthesis, leading to death of the organism. Thus the effect of the dual action is to reduce the minimum inhibitory concentrations (MIC) of each agent (synergism) and to convert a bacteriostatic action to a bactericidal action.

The activity of sulfamethoxazole-trimethoprim therefore depends upon the ability of both sulfamethoxazole and trimethoprim to affect the folate metabolism of the bacterium; however, for sulfamethoxazole-trimethoprim to be therapeutic it must not affect the folate metabolism of the host. Since sulfamethoxazole affects only the *de novo* synthesis of dihydrofolic acid by bacteria, it does not affect folate metabolism of animals. Since in animals, as in bacteria, the folates have to be recycled to the active form by dihydrofolate reductase, trimethoprim could be expected to affect mammalian folate metabolism. Trimethoprim, however, was especially selected from similar folate inhibitors because of its low toxicity for animals and high toxicity for bacteria. This difference has since been shown to be due to the fact that the affinity of trimethoprim for the dihydrofolate reductase of bacteria is some 40,000 times greater than for the corresponding mammalian enzyme.

10.2 Pharmacokinetics

Peak plasma levels of trimethoprim and sulfamethoxazole are higher and achieved more rapidly after one hour of intravenous infusion of sulfamethoxazole-trimethoprim than after oral administration of an equivalent dose of a trimethoprim-sulfamethoxazole oral presentation. Plasma concentration, elimination half-life and urinary excretion rates show no significant differences following either the oral or intravenous route of administration.

Absorption: Both trimethoprim and sulfamethoxazole are rapidly absorbed following oral administration. Detectable levels of both drugs appear in the blood in about five minutes with significant levels being reached within an hour. Peak blood levels for both compounds are attained usually in two to four hours, are maintained for about seven hours, and detectable amounts are still present after 24 hours. When the two drugs are administered together, the individual blood levels are similar to those achieved when the drugs are administered separately, thus indicating no effect in absorption of one drug by the other.

Distribution: The ratio of one part trimethoprim to five parts sulfamethoxazole achieves drug concentrations in the blood in the ratio of approximately 1:20, a ratio considered to be optimal against a wide range of bacteria. Unlike sulfamethoxazole, trimethoprim concentrates in tissues; biopsy material from a small number of patients taking trimethoprim preoperatively indicated that the concentration of trimethoprim in the tissues exceeded that of the plasma sampled at the same time - most significant in the lung (by 10 times). A similar pattern occurs in animals. Levels of trimethoprim in the sputum were also found to be higher than in the plasma following oral administration of trimethoprim-sulfamethoxazole.

The concentrations of both drugs have also been found to be well-maintained in lymph and tissue fluids. In serum, the degree of protein-binding by trimethoprim varies with the concentration, but it normally is about 44% bound to plasma protein. Sulfamethoxazole was found to be about 70% bound to plasma protein. Addition of sulfamethoxazole reduced the binding of trimethoprim by 3 to 4%, but there was no change in the protein-binding of sulfamethoxazole (about 66%) at therapeutically attainable concentrations of the two drugs.

Metabolism and Elimination: Studies conducted on the individual components administered separately, indicate that in the presence of a high fluid intake, approximately 50%, and in the presence of a low fluid intake, approximately 40% of the orally ingested trimethoprim is excreted unchanged in the urine within 24 hours. Sulfamethoxazole is more extensively metabolised than TMP, via acetylation, oxidation or glucuronidation. Over a 72 hour period, approximately 85% of the dose can be accounted for in the urine as unchanged drug plus the major (N4- acetylated) metabolite. Approximately 10% of the excreted drug is in the form of metabolites with little or no antibacterial activity. Some trimethoprim is excreted in the bile, where concentrations twice those of plasma are obtained, but as it is almost completely reabsorbed; very little appears in the feces. Studies with radio-labelled trimethoprim indicated that it is almost completely absorbed following oral administration in man; less than 4% of the radioactivity appeared in the feces over a period of six days. Radioactivity was eliminated from the plasma and urine at almost identical rates; almost all of an oral dose being excreted in the urine within 48 hours. The biological half-life of trimethoprim was calculated to be 10 hours (range of 6.2 to 12 hours in four patients), which corresponds well to the half-life of 9 to 11 hours determined in man for sulfamethoxazole.

From 25% to 50% of the orally ingested sulfamethoxazole is excreted in the urine within 24 hours. Of the excreted drug, approximately half is the N4 acetylated derivative, a fifth is the N4 conjugate, a sixth is the unchanged parent compound, and about a tenth is another N4 free compound.

Although the amount of each drug excreted is similar when given separately or in combination, the method of excretion by the kidney is quite different. Sharpstone demonstrated that there is net tubular reabsorption of filtered sulfamethoxazole, at least in patients with normal renal function,

whereas with trimethoprim there is a tubular secretory mechanism of excretion in patients with normal or impaired renal function.

Renal clearance of sulfamethoxazole increased with rising urine flow-rate, was independent of urine pH when this was less than 7, but increased with alkalinization of the urine above a pH of 7. The clearance of trimethoprim was unaffected by alteration in urine flow-rate but increased sharply with falling urine pH.

Special Patient Populations

Renal impairment

In patients with impaired renal function, sulfamethoxazole excretion was only slightly decreased, whereas trimethoprim excretion decreased markedly in severe renal impairment.

Hepatic impairment

Caution should be exercised when treating patients with severe hepatic impairment as there may decrease the absorption and biotransformation of trimethoprim and sulfamethoxazole.

Older patients

In older patients, a slight reduction in renal clearance of sulfamethoxazole but not trimethoprim has been observed.

Paediatric population

See Recommended Dose and Dosage Adjustment.

11 STORAGE, STABILITY AND DISPOSAL

Sulfamethoxazole and Trimethoprim Injection, BP should be stored at room temperature between 15° and 30°C and protected from light. *Please refer to Dosage and Administration for in-use stability instruction.*

PART II: SCIENTIFIC INFORMATION

13 PHARMACEUTICAL INFORMATION

Drug Substance


The active ingredients of Sulfamethoxazole and Trimethoprim Injection, BP are a combination of trimethoprim and sulfamethoxazole which has been established in a ratio of 1:5.

Proper name: Sulfamethoxazole

Chemical name: N¹-(5 - methyl - 3 - isoxazoly) sulfanilamide

Molecular formula and molecular mass: C₁₀H₁₁N₃O₃S 253.31g/mol

Structural formula:



Physicochemical properties: Sulfamethoxazole is a white or almost white crystalline powder


Melting point: 169.0 °C-172.0 °C

Proper name: Trimethoprim

Chemical name: 5-(3,4,5 Trimethoxybenzyl) Pyrimidine-2,4-diamine

Molecular formula and molecular mass: C₁₄H₁₈N₄O₃ 290.32 g/mol

Structural formula:



Physicochemical properties: Trimethoprim is a white to yellowish-white powder.

Melting point: 199-203°C

Solubility: Very slightly soluble in water. Slightly soluble in alcohol.

14 CLINICAL TRIALS

The clinical trial data on which the original indication was authorized is not available.

15 MICROBIOLOGY

Trimethoprim-sulfamethoxazole is bactericidal *in vitro* against the gram-negative and gram-positive organisms listed in Table 4.

In vitro Activity: Trimethoprim is, in general, more active than sulfamethoxazole against most bacterial species (see Table 2). Notable exceptions to this include *Neisseria gonorrhoeae* and *Pseudomonas aeruginosa* (which is, in general, insensitive to these drugs).

Table 2
COMPARISON OF ACTIVITY OF TRIMETHOPRIM AND SULFAMETHOXAZOLE *IN VITRO*

	MIC (mcg/mL)	
	Trimethoprim	Sulfamethoxazole
<i>Streptococcus pyogenes</i>	0.4	100 (± 25)
<i>Diplococcus pneumoniae Type II</i>	1	32 (± 16)
<i>Viridans streptococci</i>	0.25	8
<i>Streptococcus faecalis</i>	0.5	100
<i>Streptococcus agalactiae</i>	4	50
<i>Staphylococcus aureus</i>	0.2	4
<i>Erysipelothrix rhusiopathiae</i>	8	>100
<i>Corynebacterium pyogenes</i>	0.4	>100
<i>Corynebacterium diphtheriae</i>	0.4	>100
<i>Clostridium perfringens</i>	50	16 (± 8)
<i>Mycobacterium tuberculosis</i>	250	>1000
<i>Nocardia asteroides</i>	10	5
<i>Escherichia coli</i>	0.2	8
<i>Citrobacter freundii</i>	0.1	3
<i>Klebsiella pneumoniae</i>	0.5	16
<i>Klebsiella rhinoscleromatis</i>	0.5	10
<i>Enterobacter aerogenes</i>	3	>100
<i>Salmonella typhi</i>	0.4	4
<i>Salmonella typhimurium</i>	0.3	10
<i>Shigella spp.</i>	0.4	4
<i>Vibrio comma</i>	0.8	32
<i>Pasteurella septica</i>	0.1	8
<i>Haemophilus influenzae</i>	0.12	>50
<i>Bordetella pertussis</i>	3	100

	MIC (mcg/mL)	
	Trimethoprim	Sulfamethoxazole
<i>Moraxella lacunata</i>	4	8 (± 2)
<i>Proteus spp.</i>	1	8
<i>Providencia B</i>	1	30
<i>Pseudomonas aeruginosa</i>	>100	25
<i>Pseudomonas pseudomallei</i>	4	10
<i>Neisseria gonorrhoeae</i>	12	1.6
<i>Neisseria meningitidis</i>	8	1.5

The activities were compared in the Wellcome Nutrient Agar containing 5% lysed horse blood. For *Neisseria* and *Haemophilus* spp., the medium was heated at 80°C for 5 minutes and in the case of *Mycobacterium tuberculosis*, Peizer and Schacter medium was used.

Demonstration of Synergy


By testing trimethoprim and sulfamethoxazole, both separately and in combination, synergy can be demonstrated *in vitro*. Synergy is indicated by one or all of the following:

1. by a reduction in the MIC of each drug when the drugs are used in combination
2. by an increase in the size of the zone of inhibition around the combination disc; and
3. by an increase in bactericidal activity when the drugs are used in combination

The reduction in the MIC varies with the ratio of the drugs present and it has been demonstrated that the optimum ratio, as measured by maximum reduction in the MIC's of both drugs, is that in which the drugs are present in proportions corresponding to their respective MIC when acting singly. It should be emphasized, however, that potentiation occurs over a wide range of ratios. With an excess of one of the drugs, the proportion of the other drug may be markedly reduced below that of the optimum ratio, yet still produce a synergistic effect.

Figure 2

CONCENTRATION OF SULFAMETHOXAZOLE



CONCENTRATION OF SULFAMETHOXAZOLE: Isobologram showing the synergy existing between trimethoprim and sulfamethoxazole. Concentrations required to produce 50% inhibition of bacterial growth.

Because of the wide variation in sensitivities of organisms to trimethoprim and sulfamethoxazole, the optimum ratio is also variable and could be different for each organism. Since, in general, trimethoprim is about 20 to 100 times more active than sulfamethoxazole, when examining strains for enhanced susceptibility to the combination investigators have generally preferred to use a fixed ratio, choosing one near the modal optimum. The 1:20 ratio is used most frequently, and examples of the increase in activity are shown in Table 3.

Table 3
EFFECT ON MIC OF COMBINING 1 PART OF TRIMETHOPRIM WITH 20 PARTS OF SULFAMETHOXAZOLE

Organism	MIC mcg/mL			
	Sulfamethoxazole		Trimethoprim	
	Alone	Mixture	Alone	Mixture
<i>Streptococcus pyogenes</i>	>100	1.0	1.0	0.050
<i>Diplococcus pneumoniae</i>	30	2.0	2.0	0.100
<i>Staphylococcus aureus</i>	3	0.3	1.0	0.015
<i>Haemophilus influenzae</i>	10	0.3	1.0	0.015
<i>Bordetella pertussis</i>	50	4.0	3.0	0.200
<i>Klebsiella pneumoniae</i>	>100	4.0	1.0	0.200
<i>Klebsiella aerogenes</i>	>100	4.0	1.0	0.200
<i>Escherichia coli</i>	3	1.0	0.3	0.050
<i>Salmonella typhimurium</i>	10	1.0	0.3	0.050
<i>Shigella sonnei</i>	10	1.0	0.3	0.050
<i>Proteus vulgaris</i>	30	3.0	3.0	0.150
<i>Neisseria gonorrhoeae</i>	27	1.0	14.4	0.540

Table 4 shows the consolidated reported incidence of sensitivity of 49 165 strains of 40 species to trimethoprim plus sulfamethoxazole from 28 studies. A standard sensitivity disc containing 1.25

mcg trimethoprim and 23.75 mcg sulfamethoxazole was used in each study, but the medium varied.

Table 4
INCIDENCE OF SENSITIVITY TO TRIMETHOPRIM + SULFAMETHOXAZOLE

Species	Total No.of Strains	Sensitive Strains	
		Total	%
<i>S. aureus</i>	4 929	4 280	86.8
<i>S. epidermidis</i>	99	83	83.8
<i>D. pneumoniae</i>	140	140	100.0
<i>St. Pyogenes</i>	757	699	92.3
<i>St. viridans</i>	873	803	91.9
<i>Streptococci</i>	191	102	53.4
<i>St. agalactiae</i>	20	20	100.0
<i>Enterococci</i>	7 394	3 798	51.4
<i>Escherichia coli</i>	18 903	16 851	89.1
<i>Klebsiella</i>	1 365	1 109	81.2
<i>K. pneumoniae</i>	12	12	100.0
<i>Proteus spp.</i>	3 142	2 436	77.5
<i>Pr. vulgaris</i>	610	402	65.9
<i>Pr. mirabilis</i>	2 730	2 337	85.6
<i>Pr. morgani</i>	183	160	87.4
<i>Pr. rettgeri</i>	498	431	86.5
<i>Providencia A</i>	133	104	78.2
<i>Klebs.-Enterobacter</i>	670	458	68.4
<i>Kl. edwardsii</i>	2	2	100.0
<i>Enterobacter</i>	1 344	1 169	86.9
<i>Ent. cloacae</i>	193	187	96.9
<i>Salmonella</i>	594	586	98.6
<i>Hafnia</i>	92	82	89.1
<i>Shigella</i>	226	222	98.2
<i>Sh. dysenteriae</i>	12	8	66.7
<i>Ps. aeruginosa</i>	3 081	600	19.5
<i>Ps. pseudomallei</i>	12	6	50.0
<i>Citrobacter</i>	202	184	91.1
<i>Serratia</i>	28	26	92.9
<i>Paracolobactrum</i>	84	59	70.2
<i>Haemophilus influenzae</i>	284	218	76.8
<i>Flavobacterium</i>	2	2	100.0

Species	Total No. of Strains	Sensitive Strains	
		Total	%
<i>Achromobacter</i>	160	124	77.5
<i>Arizona sp.</i>	18	10	55.6
<i>Alcaligenes sp.</i>	150	127	84.7
<i>AD group</i>	4	4	100.0
<i>Cory. diphtheriae</i>	2	2	100.0
<i>Acinetobacter</i>	16	15	93.7
<i>Aeromonas</i>	4	4	100.0
<i>Neisseria Meningitidis</i>	6	6	100.0
<i>Neisseria gonorrhoeae</i>	32	31	97.0

The resistance of *Bacteroides* spp. and *Lactobacilli* is of special interest, for they comprise the major portion of the flora of the gut. Trimethoprim plus sulfamethoxazole given daily for 10 days to 12 adult volunteers, eliminated all members of the *Enterobacteriaceae* family from the feces but did not affect either of the former bacterial groups. This lack of effect of these major groups probably accounts for the infrequent occurrence of intestinal upsets during therapy with sulfamethoxazole-trimethoprim.

Trimethoprim and Sulfonamide-Resistant Strains

The theoretical basis for the synergistic effect of sulfamethoxazole-trimethoprim is that sulfamethoxazole reduces the amount of dihydrofolate synthesized by the infecting organism (usually causing bacteriostasis), and an additional small amount of trimethoprim produces a complete block in the conversion of the folate to its active form (usually causing bacterial death).

When examined by conventional susceptibility methods, an organism is regarded as resistant to sulfonamides when its macroscopic growth is not affected. "Resistance" by this definition does not necessarily mean that the sulfonamide has not reduced the folate biosynthesis of the organism. There is indirect enzymatic evidence that the dihydrofolate content of such sulfonamide-resistant strains is, in fact, reduced in the presence of sulfonamides, although not to the same extent as that of sulfonamide-sensitive strains. Therefore, in the presence of sulfamethoxazole, the effect of trimethoprim on these sulfonamide-resistant strains should be increased because the amount of substrate against which the trimethoprim competes is reduced. *Streptococcus faecalis* is often regarded as being indifferent to the presence of sulfonamides, yet the susceptibility of this organism (and of sulfonamide-resistant strains of *Escherichia coli*) can be shown to be enhanced markedly with the addition of trimethoprim. Perhaps even more convincing evidence can be obtained by the diffusion method.

Although sulfonamide sensitivity discs produce no zones of inhibition with *Streptococcus faecalis*, discs containing 23.75 mcg sulfamethoxazole plus 1.25 mcg trimethoprim produce larger zones of inhibited growth of *Streptococcus faecalis* than do discs containing 1.25 mcg trimethoprim. The difference in size is abolished when para-aminobenzoic acid is present.

Reversal of Trimethoprim Activity

Trimethoprim acts by interfering with the conversion of dihydrofolic acid to tetrahydrofolic acid. Therefore, the presence of an exogenous source of the latter should, theoretically, diminish or even abolish the antibacterial activity of sulfamethoxazole-trimethoprim in the host.

In vitro, 1 mcg/mL folic acid affects only the sensitivities of *Streptococcus faecalis*, an organism known to utilize exogenous folates. *In vivo*, when administered subcutaneously to mice infected with *Diplococcus pneumoniae*, *Streptococcus pyogenes*, *Escherichia coli*, *Proteus vulgaris*, *Salmonella schottmuelleri* and *Salmonella typhimurium*, folic acid does not affect the ability of trimethoprim to potentiate the antibacterial activity of sulfamethoxazole.

The *in vitro* interference with the action of trimethoprim and the sulfonamides by thymidine also raises the question of whether thymidine could affect *in vivo* activity. Experimental studies in the hamster indicate that thymidine is degraded rapidly *in vivo*. In experiments in mice where large doses of the nucleoside were given intraperitoneally, it did not interfere with the protection afforded by trimethoprim and sulfamethoxazole against *Proteus vulgaris*.

Resistance Development

During the serial passage in the presence of trimethoprim, little change in sensitivity occurs with light inocula; however, resistance develops rapidly with heavy inocula. With sulfonamide-sensitive strains, the emergence of these mutants is markedly delayed by the presence of sulfamethoxazole. The delaying effect of the sulfonamide depends, however, on the degree of sulfonamide resistance and is minimal with highly resistant strains.

Recently, R factors conferring high degrees of trimethoprim resistance have been identified in members of the Enterobacteriaceae family isolated from man and animals. A factor conferring high trimethoprim and sulfonamide resistance was detected in a strain of *Escherichia coli* and in a strain of *Klebsiella aerogenes*. Both strains were isolated from infected urine of human patients.

16 NON-CLINICAL TOXICOLOGY

Acute Studies

Acute toxicity studies in rats of the separate components and of trimethoprim and sulfamethoxazole combined in a ratio of 1:5 demonstrated the following LD50 values:

	Sulfamethoxazole	Trimethoprim	1:5
Rats (Adult) oral	2000 mg/kg	1500 mg/kg	6500 mg/kg
Rats (Neonates) oral	1360 mg/kg	195 mg/kg	1160 mg/kg

Subacute Studies

Daily dosages of 33, 100, and 300 mg/kg of trimethoprim and 133, 400, and 1200 mg/kg of sulfamethoxazole were given to young, sexually immature rhesus monkeys for one month. The compounds were also given in combination; the lower doses of each being combined, and similarly the higher ones. Effects on weight gain were seen. Loss in weight was noted with high and medium dosage groups with the combined drugs. Changes were induced in hemopoiesis which were consistent with trimethoprim action in interfering with dihydrofolate reductase activity. Also high doses of the sulfonamide produce hypoplastic hemopoietic changes. Half the animals on high dose levels showed increased blood urea concentrations.

A similar study in rats produced similar results. In addition, some changes associated with the sulfonamide were noted in the thyroid and in the pituitary, such as increased weight and epithelial changes. Fatty changes were also seen in the liver of monkeys and rats on the medium and high dose levels.

Chronic Studies

Six Months

Toxicity studies of six months duration were conducted in rats and monkeys with a combination of trimethoprim and sulfamethoxazole (1:2 ratio) with total daily oral doses ranging between 99 and 900 mg/kg.

Doses of 99 mg/kg daily for six months were well-tolerated in both species with minimal signs of toxicity; 300 mg/kg was well-tolerated by monkeys, but in rats impaired growth was seen and 2 of the 10 animals in this dosage group died. With 900 mg/kg, marked effects on growth and on survival occurred in both species.

Histopathological examinations were made on more than 20 different tissues from each species; these showed depression of hematopoiesis in both species in the 300 and 900 mg/kg dosage groups and minor changes with 99 mg/kg. These bone marrow changes were related to trimethoprim's interference with dihydrofolate reductase activity. Other tissue changes attributed to drug action seen in the rat, but not in the monkey, were thyroid hyperplasia and pituitary cytological effects, both associated with the sulfonamide moiety.

12-14 Months

Oral toxicity studies with trimethoprim and sulfamethoxazole, singly or in a 1:5 combination, were conducted in the monkey and in the rat. For the monkey, dose levels ranging from 10 + 50 to 60 + 300 mg/kg six days per week were employed for a period of 52 weeks; for the rat, dose levels ranging from 5 + 25 to 120 + 600 mg/kg per day were employed for a duration of 60 weeks.

In the monkey, the 1:5 combination did not produce any significant compound-related effects, except for a slight reduction in weight gain in the 60 + 300 mg/kg dose group.

In the rat study, thyroid hyperplasia of a dose-related severity was seen after 13 weeks in all animals receiving sulfamethoxazole. This hyperplasia progressed to nodularity or adenoma formation in some rats after 52 weeks at doses as low as 50 mg/kg per day, and to local vascular invasion and lung metastases after 60 weeks at doses as low as 150 mg/kg per day. Pituitary changes (large pale cells, often vacuolated), considered to be secondary to the thyroid change, were found in a few rats in all the sulfamethoxazole treated groups.

The phenomenon of thyroid hyperplasia in rats has been produced in this species by a number of sulfonamides and antithyroid drugs. The thyroid hyperplasia which occurs under the influence of these drugs is considered to be compensatory to the failure of thyroid hormone synthesis; it has been stated that this hyperplasia can be prevented or reversed by thyroid hormone. The progression of thyroid hyperplasia to nodule or adenoma formation is an observation in rats which has been reported previously in the literature on the antithyroid drugs, thiouracil and thiourea. It is considered that in these studies the production of thyroid tumours was due, not to any direct carcinogenic action of the drugs, but rather to the excessive and prolonged stimulation of the thyroid epithelium by the thyrotropic secretion of the pituitary.

Other changes associated with sulfamethoxazole treatment in our animal studies were: a dose-related increased alkaline phosphatase, a dose-related reduction in mean body weight gain, slight depression of hematopoiesis, testicular atrophy, focal renal calcification, and slightly increased fat vacuolation of the liver and kidney.

Human Tolerance Studies

Chronic Tolerance and Toxicity Study

A double-blind, placebo-controlled trial designed to study human tolerance and possible toxic effects of an orally administered 1:5 trimethoprim/sulfamethoxazole combination, was completed in 36 normal healthy men for 13 weeks. At the two dose levels investigated (80 + 400 and 160 + 800 mg/kg three times daily) the drug appeared to be well-tolerated, with only a few minor, easily

reversible side-effects occurring. The trial had to be stopped in two subjects due to recurrent black tongue.

Thyroid Function Study

Thyroid function tests (protein-bound iodine and serum cholesterol determinations) were conducted in 25 patients who received two to four sulfamethoxazole-trimethoprim tablets daily for a duration of 35 to 760 days. In none of these patients was there evidence of depression of thyroid function. One patient showed a diffused stroma of the thyroid and an increased¹³¹I uptake, and another patient had a small diffused goiter after 120 days of treatment.

Reproduction and Teratology

A three-phase investigation, comprised of a fertility and general reproductive performance study and a perinatal and postnatal study in the rat, and teratology studies in the rat and rabbit, were conducted with an orally administered 1:5 combination of trimethoprim to sulfamethoxazole. The dose levels investigated, singly or combined, were 70 + 350, 30 + 150, 15 + 75, 0 + 350, 0 + 150, 0 + 75, and 14 + 0 or 70 + 0 mg/kg.

Some drug-related effects noted in the investigation were: a reduced body weight gain by eight weeks in males in the Fertility and General Reproductive Performance Study at dose levels of 150 or 350 mg/kg of sulfamethoxazole, alone or in combination, and an increased incidence of maternal mortality in the rabbit teratology study at the same dose levels. In one of the 18 litters of the high combination group, four of the eight pups were abnormal. Two had bone malformations and two had curled tails, missing or small kidneys, absence of eyelid and one also had misshapen lateral ventricles of the brain. The instances of small, underdeveloped kidneys were such as to raise a question of dose relationships. In the teratogenicity study in rats, instances of small, underdeveloped kidneys were seen: in control group - 0; in combination groups (420 mg/kg dose) - 6 (8.5%), (180 mg/kg dose) - 3 (4.7%), (90 mg/kg dose) - 2 (3.2%). Other malformations noted in a group receiving 420 mg/kg, were one instance of incomplete nasal septum and two fetuses with abnormally large openings in the lateral ventricles. Fertility and general reproductive performance, and early and late fetal development were not affected by the dose regimen employed.

Fertility

In these studies, the animals were dosed per os with a 1:5 mixture of trimethoprim to sulfamethoxazole daily from 60 days before mating until the end of weaning.

In the rat, at 600 mg/kg there was a slight, non-significant lowering of the pregnancy rate when compared with controls. The number of live progeny per litter at birth and at weaning was less than in controls. A slight treatment-related disturbance of estrus and of sperm count was also noted.

With 200 mg/kg the pregnancy rate was slightly lower than in controls, but the other effects seen with the higher dose were not noted.

In the rabbit, daily oral doses of 600 mg/kg produced vomiting, even with divided doses, and was therefore abandoned. Two hundred mg/kg did not have a significant effect on the pregnancy rate, on the number of live births per litter, or on the mean weight of progeny at birth or at weaning.

Teratogenicity

In these studies, rats and rabbits were dosed by stomach tube daily from days 8 to 16 of pregnancy, or on a single day of pregnancy (rat only). Trimethoprim and sulfamethoxazole were used alone, in a 1:4 combination, and in a 1:2 combination.

For the rat, dosing with 500 mg/kg of trimethoprim on any single day of gestation between days 8 to 16 had no effect on the dams or their fetuses. A single dose of 2000 mg/kg of trimethoprim was lethal to most fetuses when given on the eighth or ninth day, and it produced a very high incidence of malformations when given on days 10, 11 or 12. However, the incidence of these malformations dropped off precipitately when dosing was on the 13th day or later.

The most common abnormality seen with either compound in the rat, when dosing was daily on days 8 to 16 of pregnancy, was cleft palate which occurred with 200 mg/kg of trimethoprim alone and with 640 mg/kg of sulfamethoxazole alone. Higher doses of trimethoprim produced bony defects and exencephaly, related to its action in interfering with dihydrofolate reductase activity. The abnormalities could be prevented by the administration of folic acid subcutaneously. No fetal abnormalities were found at daily doses of 160 mg/kg or less of trimethoprim, or 512 mg/kg or less of sulfamethoxazole.

Using compounds in a 1:4 trimethoprim/sulfamethoxazole combination, fetal malformations appeared at between 128 mg/kg and 160 mg/kg of trimethoprim and 512 mg/kg and 640 mg/kg of sulfamethoxazole. There appeared to be a distinct synergism with the 1 to 2 mixture.

In rabbits given the drug daily during organogenesis (days 8 to 16), no teratogenic effect was revealed with the 1:4 mixture or its components. While no important effect on the incidence of dead fetuses was noted with daily doses of 125 mg/kg of trimethoprim, 500 mg/kg or less of sulfamethoxazole, or with 312.5 mg/kg of the combination, the incidence of fetuses dying before full term was higher than for controls in the groups given trimethoprim except at the 62.5 mg/kg dose. Pregnant does tolerated the combination better than sulfamethoxazole alone.

17 SUPPORTING PRODUCT MONOGRAPHS

1. SEPTRA® Injection, solution, sulfamethoxazole (80 mg/mL) + trimethoprim (16 mg/mL), submission control 259323, Product Monograph, Aspen Pharmacare Canada Inc. MAR 28, 2022.

**READ THIS FOR SAFE AND EFFECTIVE USE OF YOUR MEDICINE
PATIENT MEDICATION INFORMATION**

**^PSulfamethoxazole and Trimethoprim Injection, BP
sterile solution for the preparation of intravenous infusions**

Read this carefully before you start taking Sulfamethoxazole and Trimethoprim Injection, BP and each time you get a refill. This leaflet is a summary and will not tell you everything about this drug. Talk to your healthcare professional about your medical condition and treatment and ask if there is any new information about Sulfamethoxazole and Trimethoprim Injection, BP.

Serious Warnings and Precautions

Sulfamethoxazole and Trimethoprim Injection, BP can cause serious side effects. If you get any of these side effects, stop taking Sulfamethoxazole and Trimethoprim Injection, BP and get immediate medical help. **See “Serious side effects and what to do about them” table for symptoms of these side effects.**

- Severe skin reactions that may be life threatening. These include Steven-Johnson syndrome, toxic epidermal necrolysis and drug reaction with eosinophilia and systemic symptoms (DRESS), acute febrile neutrophilic dermatosis (Sweet’s syndrome) and acute generalised exanthematous pustulosis.
- Blood problems including:
 - Agranulocytosis, where the number of white cells in the blood becomes dangerously low.
 - A blood disease called aplastic anemia, where the bone marrow is unable to make enough blood cells from being damaged.
 - A blood disease called immune thrombocytopenia, which can be life-threatening.
- Allergic reaction in the lungs and in the airways, where the airways can close up and make breathing difficult, and can be life-threatening if the person does not get medical help.

For further information and symptoms see:

- the **“To help avoid side effects and ensure proper use,...”** section
- the **“What are possible side effects from using Sulfamethoxazole and Trimethoprim Injection, BP?”** section

What is Sulfamethoxazole and Trimethoprim Injection, BP used for?

Sulfamethoxazole and Trimethoprim Injection, BP is used to treat:

- urinary tract infections;
- lung infections such as bronchitis or pneumonia (a lung infection caused by fungi);
- gastrointestinal (stomach and bowel) infections such as cholera or dysentery;
- nocardiosis, an infection of the lungs or other parts of the body;
- brucellosis which is a disease spread from animals to humans. When used to treat brucellosis it is used along with another medicine, gentamicin or rifampicin.

Antibacterial drugs like Sulfamethoxazole and Trimethoprim Injection, BP treat only bacterial infections. They do not treat viral infections. Although you may feel better early in treatment, Sulfamethoxazole and Trimethoprim Injection, BP should be used exactly as directed. Misuse or

overuse of Sulfamethoxazole and Trimethoprim Injection, BP could lead to the growth of bacteria that will not be killed by Sulfamethoxazole and Trimethoprim Injection, BP (resistance). This means that Sulfamethoxazole and Trimethoprim Injection, BP may not work for you in the future.

How does Sulfamethoxazole and Trimethoprim Injection, BP work?

Sulfamethoxazole and Trimethoprim Injection, BP contains two different antibiotics called sulfamethoxazole and trimethoprim. They work together to kill or to slow or stop the growth of bacteria or fungi that cause disease. This means that Sulfamethoxazole and Trimethoprim Injection, BP can be given to prevent or to treat certain kinds of infectious diseases.

What are the ingredients in Sulfamethoxazole and Trimethoprim Injection, BP?

Medicinal ingredients: trimethoprim and sulfamethoxazole

Non-medicinal ingredients: propylene glycol, tromethamine, sodium hydroxide, ethanol 96% and water for injection.

Sulfamethoxazole and Trimethoprim Injection, BP comes in the following dosage forms:

Solution for the preparation of intravenous infusions: sulfamethoxazole (80 mg / mL) + trimethoprim (16 mg / mL).

Do not use Sulfamethoxazole and Trimethoprim Injection, BP if:

- you are allergic to sulfamethoxazole, trimethoprim or any of the other ingredients in Sulfamethoxazole and Trimethoprim Injection;
- you are allergic to sulphonamide medicines. Examples include diabetes medicines (such as gliclazide and glibenclamide). Talk to your healthcare professional if you are allergic to a medicine and you are not sure if it is a sulphonamide medicine causing bruises or bleeding (thrombocytopenia);
- you have liver problems;
- you have kidney problems;
- you have blood problems;
- you are pregnant;
- you are breastfeeding;
- you have been told that you have a rare blood problem called porphyria, which can affect your skin or nervous system.

Children

If it is for your child, Sulfamethoxazole and Trimethoprim Injection, BP should not be given if they are less than 2 months old.

If you are not sure if any of the above apply to you, talk to your healthcare professional before being given Sulfamethoxazole and Trimethoprim Injection, BP.

To help avoid side effects and ensure proper use, talk to your healthcare professional before you take Sulfamethoxazole and Trimethoprim Injection, BP. Talk about any health conditions or problems you may have, including if you:

- have severe allergies or asthma;
- you don't have enough folic acid (a vitamin) in your body called folate deficiency;
- you are underweight or malnourished;
- you have a disease called glucose-6-phosphate dehydrogenase deficiency;
- you are at risk for a rare blood disorder called porphyria, which can affect your skin or nervous system;

- you have been told by your healthcare professional that you have a high level of potassium in your blood;
- you have been told by your healthcare professional that you have a low level of sodium or albumin in your blood;
- you have been told by your healthcare professional that you have any serious disorder of the blood or blood forming tissues such as low blood cell counts;
- you have existing heart or lung disease, which may lead to a build-up of fluid in your body;
- you have hereditary disorder called phenylketonuria and are not on a special diet to help your condition;
- you are human immunodeficiency virus (HIV) positive or have a condition called Acquired Immuno deficiency Syndrome (AIDS);
- If you have kidney problems.

Other warnings you should know about:

Use in Elderly Patients

Elderly patients are more likely to get serious side effects when receiving Sulfamethoxazole and Trimethoprim Injection, BP. This is increased if you have kidney or liver disease or are taking some types of other medicines, such as diuretics.

Pregnancy and Breastfeeding

Talk to your healthcare professional before taking this medicine if you are planning to get pregnant or planning to breastfeed. You should not receive Sulfamethoxazole and Trimethoprim Injection, BP if you are pregnant or are breastfeeding.

Use in Patients with Acquired Immunodeficiency Syndrome (AIDS)

If you have AIDS, you may be more likely to get side effects when receiving Sulfamethoxazole and Trimethoprim Injection, BP. These may include rash, severe allergic reactions, fever or low blood cell counts.

Gastrointestinal - C. difficile colitis

Sulfamethoxazole and Trimethoprim Injection, BP may increase your risk of being infected with a bacteria called *C. difficile*. Symptoms include watery diarrhea that happens three or more times per day or diarrhea associated with abdominal cramping.

Driving and using machines

After you are given Sulfamethoxazole and Trimethoprim Injection, BP you may feel weak, tired, dizzy, or confused. Before driving a vehicle or using machinery wait to see how you feel after being given Sulfamethoxazole and Trimethoprim Injection, BP.

Use in patients who have taken sulfamethoxazole-trimethoprim recently such as in the past days to weeks

If you have recently taken the medicines in Sulfamethoxazole and Trimethoprim Injection, BP (sulfamethoxazole and trimethoprim), you could get serious reactions after taking Sulfamethoxazole and Trimethoprim Injection, BP. These include circulatory shock with fever, very low blood pressure and confusion. See the “Serious side effects and what to do about them” table for symptoms.

Tell your healthcare professional about all the medicines you take, including any drugs, vitamins, minerals, natural supplements or alternative medicines.

The following may interact with Sulfamethoxazole and Trimethoprim Injection, BP:

- medicines to remove excess water from the body known as diuretics (water pills) such as spironolactone;
- medicines to treat malaria such as pyrimethamine;
- medicines to treat Human Immunodeficiency Virus (HIV) such as zidovudine or lamivudine;
- medicines to thin the blood such as warfarin;
- medicines to treat epilepsy (fits) and seizures such as phenytoin;
- medicines to treat cancer or arthritis such as methotrexate;
- medicines to treat heart conditions such as digoxin or procainamide;
- medicines that can increase the amount of potassium in your blood, such as steroids (like prednisolone) and heart and high blood pressure medicines;
- medicines used after organ transplantation such as cyclosporine;
- medicines to treat Parkinson's disease, multiple sclerosis, the flu or shingles such as amantadine;
- medicines to treat diabetes, such as glibenclamide, glipizide or tolbutamide (sulphonylureas) and repaglinide;
- medicines to treat bacterial infections such as rifampicin;
- medicines used after cancer treatment or to help with low levels of folate such as folic acid;
- medicines to help prevent pregnancy such as contraceptives.
- medicine used to treat irregular heartbeats, such as dofetilide.

Drug-Food Interactions

- Tell your healthcare professional if you have special dietary needs, especially if you are following a potassium rich diet. Potassium rich foods include beans, dark leafy greens, potatoes, squash, yogurt, fish, avocados, mushrooms and bananas.

How to take Sulfamethoxazole and Trimethoprim Injection, BP:

- Sulfamethoxazole and Trimethoprim Injection, BP will be given to you by a healthcare professional.
- It will be given to you as a continuous infusion into your vein. This is where a medicine is slowly given to you over a period of time.
- You should drink plenty of fluids while receiving Sulfamethoxazole and Trimethoprim Injection, BP.

Usual dose:

- Your healthcare professional will decide how much Sulfamethoxazole and Trimethoprim Injection, BP you will receive.
- The dose you are given and how often you are given it will depend on:
 - your infection;
 - your weight;
 - your age.

Overdose:

If you have been given too much Sulfamethoxazole and Trimethoprim Injection, BP you may have the following signs or symptoms:

- anorexia (extreme fear of gaining weight);
- colic (severe pain in the abdomen caused by gas);
- nausea and vomiting;
- dizziness, drowsiness or confusion;
- fainting;
- headache;
- Pyrexia (fever);
- hematuria (blood in urine);
- crystalluria (cloudy urine);
- jaundice (yellowing of the skin or whites of the eye)
- Feeling depressed

If you think you have taken too much Sulfamethoxazole and Trimethoprim Injection, BP, contact your healthcare professional, hospital emergency department or regional poison control centre immediately, even if there are no symptoms.

Missed Dose:

Your healthcare professional will inject this medicine into you. If you miss a scheduled injection talk to your healthcare professional as soon as possible.

What are possible side effects from using Sulfamethoxazole and Trimethoprim Injection, BP?

These are not all the possible side effects you may feel when taking Sulfamethoxazole and Trimethoprim Injection, BP. If you experience any side effects not listed here, contact your healthcare professional.

Side effects may include:

- Feeling weak or tired;
- Trouble sleeping;
- Vision problems;
- Hair loss;
- Nose bleed;
- Inflammation at the site of injection;
- Swelling of body tissues with fluid;
- Chills;
- Sensitivity to sunlight;
- Palpitations (heart beat that feels too fast, strong or irregular);
- Cold sores, ulcers, or soreness of your tongue or inside of your mouth;
- Dry mouth;
- Heartburn;
- Abdominal pain or gas;
- Constipation;
- Nausea, vomiting and diarrhea;
- Loss of appetite;

- Passing more or less urine than usual; difficulty reaching bathroom in time;
 - Muscle and joint pain or muscle weakness;
 - Tingling or numbness in your hands and feet;
 - Problems controlling your movements;
 - Uncontrollable shaking;
 - Vertigo (sensation of movement or feeling off balance);
 - Ringing or other unusual sounds in your ears;
 - Headache;
 - Inflammation of your eye that causes pain and redness;
 - Depression;
 - Apathy (indifference and a lack of motivation);
 - Feeling unsteady or dizzy.
 - anorexia (extreme fear of gaining weight)
- Sulfamethoxazole and Trimethoprim Injection, BP can cause abnormal blood test results.

Serious side effects and what to do about them			
Symptom / effect	Talk to your healthcare professional		Stop taking drug and get immediate medical help
	Only if severe	In all cases	
Allergic reactions: swelling of face, mouth, tongue or throat which may be red and painful and/or cause difficulty in swallowing; red patches on the skin; rash; hives; fever (high temperature); joint pain; feeling sick (nausea); being sick (vomiting); chest pain.			√
Stevens-Johnson syndrome (severe skin rash): redness, blistering and/or peeling of the skin and/or inside of the lips, eyes, mouth, nasal passages or genitals, accompanied by fever, chills, headache, cough, body aches or swollen glands			√
Toxic Epidermal Necrolysis (severe skin reaction): redness, blistering and/or peeling of large areas of the skin			√
Drug reaction with eosinophilia and systemic symptoms (DRESS) (serious skin reaction that may affect more than one organ): fever, severe skin rash, peeling skin, and abnormal blood and liver function tests			√
Acute febrile neutrophilic dermatosis (Sweet's syndrome) (a serious skin condition): Purple or red, raised, painful sores on the limbs, face or neck, fever			√
Acute generalised exanthematous pustulosis (serious skin reaction): widespread skin rash and redness, skin blisters, swelling of the face, fever.			√
Lung problems: Difficulty breathing, blue colour to skin, cough, confusion, drowsiness, chest pain, wheezing, shortness of breath, rapid breathing, tightness of the chest, fever, weight loss, loss of appetite, tiredness and weakness.			√
Aseptic meningitis (inflammation of the protective lining of the brain that is not caused by infection): including sudden headache or stiffness of your neck, accompanied by fever, nausea, vomiting, sensitivity to light.			√
Acute inflammation of the small and large intestine <i>Pseudomembranous colitis</i> : including watery or bloody			√

Serious side effects and what to do about them			
Symptom / effect	Talk to your healthcare professional		Stop taking drug and get immediate medical help
	Only if severe	In all cases	
diarrhea, abdominal cramps, pain or tenderness, fever, nausea, dehydration.			
Fits (convulsions or seizures)			√
Heart problems: Increased heart rate, chest pain, shortness of breath).			√
Pancreatitis (Acute inflammation of the pancreas): including upper abdominal pain that spreads to the back, swollen and tender abdomen, nausea, vomiting, fever.			√
Hypoglycaemia (an abnormally low level of sugar in the blood): including dizziness or light-headedness, shakiness, nervousness or anxiety, feeling confused, sweating, chills.			√
Problems with your urine; pain or difficulty passing urine, blood or cloudiness in your urine.			√
Hepatitis (inflammation of the liver): fatigue, fever, body ache, abdominal pain, dark urine or pale stools, difficulty to urinate.			√
Jaundice (yellowing of the skin and whites of the eyes).			√
An infection called thrush or candidiasis which can affect your mouth or vagina.			√
Hallucinations : Seeing, hearing, smelling, tasting or feeling things that don't exist outside your mind			√
Immune thrombocytopenia : including being easily bruised, a rash on the skin that appears tiny pinpoint-sized reddish or purple spots, usually on the lower legs, bleeding from the gums or nose, and blood in the urine or stool.			√
Aplastic anemia : including feeling tired, feeling short of breath, pale skin, unexplained or easy bruising, fever, chills, sore throat, and a general feeling of being unwell.			√
Severe Cutaneous Adverse Reactions (SCAR) (severe skin reactions that may also affect other organs): <ul style="list-style-type: none"> • Skin peeling, scaling, or blistering (with or without pus) which may also affect your eyes, mouth, nose or genitals, itching, severe rash, bumps under the skin, skin pain, skin color changes (redness, yellowing, purplish) • Swelling and redness of eyes or face • Flu-like feeling, fever, chills, body aches, swollen glands, cough Shortness of breath, chest pain or discomfort 			√
Angioedema : swelling of the face, hands, feet, genitals, tongue or throat, difficulty swallowing or breathing; swelling of the digestive tract which may cause diarrhea, nausea or vomiting			√
Erythema multiforme (an allergic skin reaction): raised red or purple skin patches, possibly with blister or crust in the center; possibly swollen lips, mild itching or burning			√

If you have a troublesome symptom or side effect that is not listed here or becomes bad enough to interfere with your daily activities, talk to your healthcare professional.

Reporting Side Effects

You can report any suspected side effects associated with the use of health products to Health Canada by:

- Visiting the Web page on Adverse Reaction Reporting (<https://www.canada.ca/en/health-canada/services/drugs-health-products/medeffect-canada.html>) for information on how to report online, by mail or by fax; or
- Calling toll-free at 1-866-234-2345.

NOTE: Contact your health professional if you need information about how to manage your side effects. The Canada Vigilance Program does not provide medical advice.

Storage:

Store Sulfamethoxazole and Trimethoprim Injection, BP between 15° and 30°C. Protect from light. Keep out of reach and sight of children.

If you want more information about Sulfamethoxazole and Trimethoprim Injection, BP:

- Talk to your healthcare professional
- Find the full product monograph that is prepared for healthcare professionals and includes this Patient Medication Information by visiting the Health Canada website (<https://health-products.canada.ca/dpd-bdpp/index-eng.jsp>) or by calling 1-855-819-0505.

This leaflet was prepared by
Juno Pharmaceuticals Corp.
402-2233 Argentia Road,
Mississauga, ON, L5N 2X7

Revised: August 18, 2022