

PRODUCT MONOGRAPH
INCLUDING PATIENT MEDICATION INFORMATION

PrCAMZYOS™

mavacamten capsules

Capsules, 2.5 mg, 5 mg, 10mg and 15 mg, Oral

Cardiac myosin inhibitor

Bristol-Myers Squibb Canada
Montreal, Canada

Date of Initial Authorization:
November 8, 2022

Submission Control Number: 258772

™ of Myocardia, Inc., used under license by Bristol-Myers Squibb Canada

RECENT MAJOR LABEL CHANGES

Not applicable.

TABLE OF CONTENTS

Sections or subsections that are not applicable at the time of authorization are not listed .

RECENT MAJOR LABEL CHANGES 2

TABLE OF CONTENTS 2

PART I: HEALTH PROFESSIONAL INFORMATION 4

1 INDICATIONS 4

 1.1 Pediatrics 4

 1.2 Geriatrics..... 4

2 CONTRAINDICATIONS 4

3 SERIOUS WARNINGS AND PRECAUTIONS BOX 4

4 DOSAGE AND ADMINISTRATION 5

 4.1 Dosing Considerations 5

 4.2 Recommended Dose and Dosage Adjustment..... 5

 4.5 Missed Dose 8

5 OVERDOSAGE 8

6 DOSAGE FORMS, STRENGTHS, COMPOSITION AND PACKAGING 9

7 WARNINGS AND PRECAUTIONS 10

 7.1 Special Populations..... 11

 7.1.1 Pregnant Women..... 11

 7.1.2 Breast-feeding..... 11

 7.1.3 Pediatrics..... 11

 7.1.4 Geriatrics 11

8 ADVERSE REACTIONS 12

 8.1 Adverse Reaction Overview..... 12

 8.2 Clinical Trial Adverse Reactions 12

9 DRUG INTERACTIONS 13

 9.2 Drug Interactions Overview..... 13

 9.4 Drug-Drug Interactions 14

9.5	Drug-Food Interactions.....	18
9.6	Drug-Herb Interactions.....	18
9.7	Drug-Laboratory Test Interactions.....	18
10	CLINICAL PHARMACOLOGY.....	18
10.1	Mechanism of Action.....	18
10.2	Pharmacodynamics.....	18
10.3	Pharmacokinetics.....	19
11	STORAGE, STABILITY AND DISPOSAL.....	21
	PART II: SCIENTIFIC INFORMATION.....	22
13	PHARMACEUTICAL INFORMATION.....	22
14	CLINICAL TRIALS.....	22
14.1	Clinical Trials by Indication.....	22
15	MICROBIOLOGY.....	28
16	NON-CLINICAL TOXICOLOGY.....	28
	PATIENT MEDICATION INFORMATION.....	30

PART I: HEALTH PROFESSIONAL INFORMATION

1 INDICATIONS

CAMZYOS (mavacamten capsules) is indicated for the treatment of symptomatic obstructive hypertrophic cardiomyopathy (oHCM) of New York Heart Association (NYHA) Class II-III in adult patients.

1.1 Pediatrics

Pediatrics (< 18 years of age): The safety and efficacy have not been established; therefore, Health Canada has not authorized an indication for pediatric use.

1.2 Geriatrics

Geriatrics: Evidence from clinical studies suggests that safety, effectiveness, and pharmacokinetics were consistent between elderly patients (≥ 65 years) and younger patients (18 to < 65 years).

2 CONTRAINDICATIONS

CAMZYOS is contraindicated:

- with concomitant use of moderate or strong cytochrome P450 (CYP) enzyme CYP 2C19 inhibitors, or strong CYP 3A4 inhibitors (see [7 WARNINGS AND PRECAUTIONS](#), Risk of heart failure or loss of response to mavacamten due to drug-drug interactions, and [9 DRUG INTERACTIONS](#)), due to risk of developing left ventricular dysfunction
- with concomitant use of moderate or strong inducers to both CYP 2C19 and CYP 3A4 (see [7 WARNINGS AND PRECAUTIONS](#), Risk of heart failure or loss of response to mavacamten due to drug-drug interactions, and [9 DRUG INTERACTIONS](#)), due to risk of loss of therapeutic effect
- during pregnancy (see [7 WARNINGS AND PRECAUTIONS](#), Reproductive Health, and Pregnant Women)
- in patients who are hypersensitive to this drug or to any ingredient in the formulation, including any non-medicinal ingredient, or component of the container. For a complete listing, see [6 DOSAGE FORMS, STRENGTHS, COMPOSITION AND PACKAGING](#).

3 SERIOUS WARNINGS AND PRECAUTIONS BOX

Serious Warnings and Precautions	
Risk of Heart Failure	
	<ul style="list-style-type: none">• CAMZYOS reduces left ventricular ejection fraction (LVEF) and can cause heart failure due to systolic dysfunction (see 7 WARNINGS AND PRECAUTIONS, Heart Failure due to Systolic Dysfunction).• Echocardiogram assessments of LVEF and left ventricular outflow tract (LVOT) gradient are required prior to, and regularly during, treatment with CAMZYOS. Initiation of CAMZYOS in patients with LVEF <55% is not recommended. Interrupt CAMZYOS treatment if LVEF is <50% at any visit or if the patient experiences heart failure symptoms or worsening clinical status

(see [4 DOSAGE AND ADMINISTRATION](#), and [7 WARNINGS AND PRECAUTIONS](#), Cardiovascular).

- Concomitant use of CAMZYOS with certain cytochrome P450 inhibitors may increase the risk of heart failure due to systolic dysfunction (see [7 WARNINGS AND PRECAUTIONS](#), Cardiovascular).

4 DOSAGE AND ADMINISTRATION

4.1 Dosing Considerations

CAMZYOS should be administered under the supervision of a physician experienced in the treatment of HCM.

Dosage must be individualized based on clinical status and echocardiographic assessment of patient response.

CAMZYOS has generally been used together with a beta-blocker or a non-dihydropyridine calcium channel blocker (CCB). Concomitant use with other negative inotropic agents, or with combined use of such agents, is not recommended (see [7 WARNINGS AND PRECAUTIONS](#), Concomitant Use with Drugs That Reduce Cardiac Contractility).

Testing prior to initiation

Prior to initiating treatment with CAMZYOS, assess LVEF by echocardiography (see [7 WARNINGS AND PRECAUTIONS](#), Cardiovascular). Initiation of treatment with CAMZYOS in patients with LVEF <55% is not recommended.

Treatment monitoring

It is important to regularly monitor the patient's symptoms of oHCM, LVOT gradient with Valsalva maneuver and LVEF using echocardiogram assessments.

Once an individualized maintenance dose is achieved (see [4.2 Recommended Dose and Dosage Adjustment](#), *Treatment Initiation*), monitor patients every 12 weeks (see [4.2 Recommended Dose and Dosage Adjustment](#), *Treatment Maintenance*).

If at any visit LVEF is <50%, treatment with CAMZYOS should be interrupted for at least four (4) weeks (see [4.2 Recommended Dose and Dosage Adjustment](#)).

Assessment of LVEF is also recommended if clinical status changes or in patients with an intercurrent illness such as serious infections or with clinically relevant arrhythmias, including atrial fibrillation and other tachyarrhythmias. In these circumstances, the dose of CAMZYOS may need to be reduced or temporarily discontinued (see [7 WARNINGS AND PRECAUTIONS](#), Cardiovascular). Dose increases are not recommended if a patient is experiencing an intercurrent illness which may impair systolic function.

4.2 Recommended Dose and Dosage Adjustment


Treatment initiation, over the first 12 weeks

The recommended starting dose of CAMZYOS is 5 mg taken orally once daily. The capsule should be swallowed whole with water and can be taken with or without food.

Patients should be assessed for early clinical response four (4) weeks after initiation of treatment with

CAMZYOS. If LVOT gradient with Valsalva maneuver is < 20 mmHg, the dose of CAMZYOS should be decreased to 2.5 mg once daily. Otherwise maintain at 5 mg once daily. Thereafter, follow-up visits should occur at eight (8) weeks and twelve (12) weeks after treatment initiation, with dose adjustments, as appropriate, see Figure 1, below.

Figure 1: Treatment Initiation Phase




Treatment Maintenance, from 12 weeks after initiation

The patient’s individualized dose of CAMZYOS will be either 2.5, 5, 10 or 15 mg once daily.


After the 12-week treatment initiation phase, patient response to CAMZYOS should continue to be monitored every 12 weeks, including echocardiogram assessment of LVEF and LVOT gradient. If symptoms of oHCM persist, while LVOT gradient with Valsalva maneuver is ≥ 30 mmHg, a step-wise dose increase may be considered in patients with an LVEF ≥ 55%, see Figure 2, below.

Figure 2: Treatment Maintenance Phase



Following treatment interruption of CAMZYOS due to LVEF < 50%, patients should be monitored carefully to determine if treatment can be resumed, see Figure 3, below.

Figure 3: Following Treatment Interruption due to LVEF < 50%



Concomitant therapy

CAMZYOS is contraindicated with concomitant use of moderate or strong CYP2C19 inhibitors, or strong CYP3A4 inhibitors (see [2 CONTRAINDICATIONS](#)). Concomitant use of CAMZYOS with certain cytochrome P450 inhibitors may increase the risk of heart failure due to systolic dysfunction.

In patients who are on stable therapy with a weak CYP 2C19 inhibitor or a moderate CYP 3A4 inhibitor, initiate CAMZYOS at the recommended starting dosage of 5 mg orally once daily.

In patients treated with CAMZYOS, who then intend to initiate a dose of a weak CYP 2C19 inhibitor or a moderate CYP 3A4 inhibitor, reduce dosage of CAMZYOS by one level, i.e., 15 → 10 mg; 10 → 5 mg; or 5 → 2.5 mg. Avoid initiation of a weak CYP 2C19 inhibitor or a moderate CYP 3A4 inhibitor in patients

who are on stable treatment with 2.5 mg of CAMZYOS because a lower once-daily dose of CAMZYOS is not available.

Special populations

Geriatric patients (65 years and above)

No specific dose adjustments are recommended for patients aged 65 years and older (see [1.2 Geriatrics](#)).

Pediatric patients (below 18 years)

Health Canada has not authorized an indication for pediatric use (see [1.1 Pediatrics](#)).

Renal Impairment

No dosage adjustment is required in patients with mild (eGFR 60 - <90 mL/min/1.73m²) and moderate (eGFR 30 - <60 mL/min/1.73m²) renal impairment. CAMZYOS should be used with caution in patients with severe (eGFR < 30 mL/min/1.73m²) renal impairment, as CAMZYOS has not been studied in this population (see [10.3 Pharmacokinetics](#)).

Hepatic impairment

No dosage adjustment is required in patients with mild (Child-Pugh Class A) and moderate (Child-Pugh Class B) hepatic impairment. CAMZYOS is not recommended for patients with severe (Child-Pugh Class C) hepatic impairment, as CAMZYOS has not been studied in this population (see [10.3 Pharmacokinetics](#)).

4.5 Missed Dose

If a dose is missed, it should be taken as soon as possible, and the next scheduled dose should be taken at the usual time the following day. Two separate doses should not be taken in a single day.

5 OVERDOSAGE

Human experience of overdose with CAMZYOS is limited. Systolic dysfunction is the most likely adverse effect of overdosage of CAMZYOS.

CAMZYOS has been given as a single dose of up to 144 mg in patients with HCM. There was one reported serious adverse reaction consisting of vasovagal reaction, hypotension, and asystole lasting 38 seconds at that dose. In healthy subjects, doses of up to 25 mg have been administered for up to 25 days. Three out of 8 participants treated at the 25 mg dose level experienced 20% or greater reductions in LVEF. A death of an infant was reported after accidental ingestion of three 15 mg capsules of CAMZYOS. Systolic dysfunction is the most likely result of overdosage of CAMZYOS.

Management of overdose

Treatment of overdose with CAMZYOS consists of discontinuation of CAMZYOS treatment, and instituting supportive measures to maintain hemodynamic status, e.g., initiation of inotropic support with adrenergic agents, if appropriate, along with close monitoring of vital signs and LVEF. Early administration of activated charcoal may be considered in case of mavacamten overdose to reduce absorption, although this measure has not been specifically studied.

For management of a suspected drug overdose, contact your regional poison control centre.

6 DOSAGE FORMS, STRENGTHS, COMPOSITION AND PACKAGING

Table 1 – Dosage Forms, Strengths, Composition and Packaging

Route of Administration	Dosage Form / Strength/Composition	Non-medicinal Ingredients
Oral	Capsules 2.5 mg, 5 mg, 10 mg and 15 mg	<p>Croscarmellose sodium, hypromellose, magnesium stearate (non-bovine), mannitol, silicon dioxide.</p> <p>Shell 2.5mg: gelatin (bovine and/ or porcine), black iron oxide, red iron oxide, titanium dioxide.</p> <p>Shell 5 mg: gelatin (bovine and/ or porcine), titanium dioxide, yellow iron oxide.</p> <p>Shell 10 mg: gelatin (bovine and/ or porcine), titanium dioxide, red iron oxide.</p> <p>Shell 10 mg: gelatin (bovine and/ or porcine), titanium dioxide, red iron oxide.</p> <p>Shell 15 mg: gelatin (bovine and/ or porcine), titanium dioxide, black iron oxide.</p> <p>Imprinting ink: black iron oxide, potassium hydroxide, propylene glycol, shellac, strong ammonia solution</p>

Description

2.5 mg: Light purple opaque cap, hard capsule, size 2, imprinted with “2.5 mg” in black, and white opaque body imprinted with “Mava” in black, both in radial direction. The capsule contains white to off-white powder.

5 mg: Yellow opaque cap, hard capsule, size 2, imprinted with “5 mg” in black, and white opaque body imprinted with “Mava” in black, both in radial direction. The capsule contains white to off-white powder.

10 mg: Pink opaque cap, hard capsule, size 2, imprinted with “10 mg” in black, and white opaque body imprinted with “Mava” in black, both in radial direction. The capsule contains white to off-white powder.

15 mg: Gray opaque cap, hard capsule, size 2, imprinted with “15 mg” in black, and white opaque body imprinted with “Mava” in black, both in radial direction. The capsule contains white to off-white powder.

Supplied in bottles of 30 capsules.

7 WARNINGS AND PRECAUTIONS

Cardiovascular

Heart Failure due to Systolic Dysfunction

CAMZYOS reduces LVEF and can cause heart failure due to systolic dysfunction. Patients with intercurrent illness, such as serious infections, or clinically relevant arrhythmias, including atrial fibrillation and other tachyarrhythmias may be at greater risk of systolic dysfunction and progressing to heart failure. The development of signs or symptoms of heart failure in an individual patient or elevation in N-terminal (NT)-pro b-type natriuretic peptide (NT-proBNP) at any time should prompt an evaluation of cardiac function. LVEF should be measured prior to initiating treatment and closely monitored thereafter. Treatment interruption may be necessary to ensure that LVEF remains $\geq 50\%$ (see [3 SERIOUS WARNINGS AND PRECAUTIONS BOX](#), and [4.1 Dosing Considerations](#)).

Risk of heart failure or loss of response to mavacamten due to drug-drug interactions

CAMZYOS is primarily metabolized by CYP 2C19, and to a lesser extent by CYP 3A4 (see [10.3 Pharmacokinetics, Metabolism](#)). Concomitant use of CAMZYOS with drugs that substantially inhibit these enzymes may lead to life-threatening adverse events, such as heart failure (see [2 CONTRAINDICATIONS](#), and [9 DRUG INTERACTIONS](#)). Conversely, concomitant use with drugs that substantially induce these enzymes may lead to loss of therapeutic effectiveness of mavacamten (see [2 CONTRAINDICATIONS](#), and [9 DRUG INTERACTIONS](#)).

Advise patients of the potential for drug interactions and with over-the-counter medications, such as omeprazole and esomeprazole

Concomitant Use with Drugs That Reduce Cardiac Contractility

Expect additive negative inotropic effects of CAMZYOS with other drugs that reduce cardiac contractility. The concomitant use of CAMZYOS with disopyramide or ranolazine, or its use in patients taking beta blockers in combination with verapamil or diltiazem has not been studied. Therefore, such use is not recommended. If concomitant therapy with any negative inotrope is initiated with CAMZYOS, or if the dose of a negative inotrope is increased, monitor LVEF closely until stable doses and stable clinical response have been achieved (see [9.2 DRUG INTERACTIONS](#), Drugs That Reduce Cardiac Contractility).

Driving and Operating Machinery

Dizziness may occur following administration of CAMZYOS. Patients should be advised not to drive or use machines if they experience dizziness. Exercise caution when driving or operating a vehicle or potentially dangerous machinery.

Reproductive Health: Female and Male Potential

- **Teratogenic Risk**

Based on animal studies, mavacamten may decrease embryonic viability, impair fetal and post-natal growth, and cause embryo-fetal malformations when administered during pregnancy (see [16 NON-CLINICAL TOXICOLOGY](#), Reproductive and Developmental Toxicology). Therefore, use of CAMZYOS during pregnancy is contraindicated (see [2 CONTRAINDICATIONS](#)).

- **Contraception**

Mavacamten was shown to transfer to semen in a dose-dependent manner in human participants and to cross the placenta in rabbits. Advise females of reproductive potential to avoid becoming pregnant and to use highly effective contraception during treatment with CAMZYOS and for at least 4 months after discontinuing treatment.

Confirm a negative pregnancy test in women of reproductive potential prior to initiation of treatment (see [16 NON-CLINICAL TOXICOLOGY](#), Reproductive and Developmental Toxicology).

7.1 Special Populations

7.1.1 Pregnant Women

There are no data on the developmental risk associated with the use of CAMZYOS in pregnant women. CAMZYOS was well-tolerated by pregnant animals. However, based on animal data, CAMZYOS may cause fetal malformations when administered during pregnancy (see [16 NON-CLINICAL TOXICOLOGY](#), Reproductive and Developmental Toxicology).

CAMZYOS treatment is contraindicated during pregnancy (see [2 CONTRAINDICATIONS](#)). Females of reproductive potential who undergo treatment with CAMZYOS should be informed of the potential hazard to the fetus and should be advised to avoid becoming pregnant prior to or during treatment and for at least 4 months after discontinuation.

If the patient becomes pregnant while receiving the drug, CAMZYOS treatment must be immediately discontinued.

7.1.2 Breast-feeding

Caution should be exercised because many drugs can be excreted in human milk. Since it is unknown whether mavacamten or its metabolites are excreted in human milk, a decision must be made whether to discontinue breast-feeding during treatment and for 4 months after the last dose, or to discontinue treatment, taking into account benefit-risk of treatment with mavacamten (see [16 NON-CLINICAL TOXICOLOGY](#), Reproductive and Developmental Toxicology).

7.1.3 Pediatrics

Pediatrics (< 18 years of age): The safety and efficacy have not been established; therefore, Health Canada has not authorized an indication for pediatric use.

7.1.4 Geriatrics

Geriatrics: Clinical trials of CAMZYOS included 95 patients of 263 (36.1%) who were aged 65 years and over, of whom, 17 of 263 patients (6.5%) were aged 75 years or older. Safety, effectiveness, and pharmacokinetics appeared to be consistent between older patients (≥ 65 years) and younger patients (18 to < 65 years).

8 ADVERSE REACTIONS

8.1 Adverse Reaction Overview

CAMZYOS reduces LVEF and can cause heart failure due to systolic dysfunction (see [3 SERIOUS WARNINGS AND PRECAUTIONS BOX](#), and [7 WARNINGS AND PRECAUTIONS](#), Cardiovascular).

Effects on systolic function

In the pivotal trial, EXPLORER-HCM, routine echocardiographic screening measures determined that 7 patients (5.7%) in the CAMZYOS group and 2 patients (1.6%) in the placebo group experienced reversible reductions in LVEF < 50%, to a median LVEF of 48% (range: 35-49%) while on treatment for a period of 30 weeks. None of the 7 patients receiving mavacamten had systolic dysfunction leading to heart failure. In 3 of the 7 CAMZYOS patients and in 1 of the 2 placebo patients, these reductions were observed in asymptomatic patients. In all 7 patients treated with CAMZYOS, LVEF recovered following interruption of CAMZYOS.

The safety of CAMZYOS was evaluated in EXPLORER-HCM (see [14 CLINICAL TRIALS](#)). The most commonly reported on-treatment adverse events in mavacamten-treated patients were dizziness (17.1%) and headache (11.4%).

Serious adverse events were observed in 8.1% of mavacamten-treated patients and 8.6% of placebo-treated patients to Week 30. Two cases of stress cardiomyopathy were observed in mavacamten patients. Syncope was reported in two mavacamten-treated patients, compared to one treated with placebo. There were no deaths in the mavacamten group.

Study drug was permanently discontinued in 1.6% of mavacamten-treated patients due to adverse events, consisting of one patient with syncope and one with atrial fibrillation.

8.2 Clinical Trial Adverse Reactions

Clinical trials are conducted under very specific conditions. The adverse reaction rates observed in the clinical trials; therefore, may not reflect the rates observed in practice and should not be compared to the rates in the clinical trials of another drug. Adverse reaction information from clinical trials may be useful in identifying and approximating rates of adverse drug reactions in real-world use.

The safety of CAMZYOS was evaluated in EXPLORER-HCM, a Phase 3, double-blind, randomized, placebo-controlled trial. Of the 251 oHCM adult patients in this trial, 123 patients were treated with a daily dose of 2.5 mg, 5 mg, 10 mg or 15 mg of CAMZYOS, while 128 were treated with placebo. CAMZYOS-treated patients received a median duration of exposure to mavacamten of 30.4 weeks.

Median age was 59 years (range: 26 to 82), with 36.1% of patients ≥ 65 years, and 6.5% of patients ≥ 75 years. 94% were White, and 54% male.

Adverse events that occurred $\geq 5\%$ of mavacamten-treated patients and that were more common than in placebo-treated patients at Week 30 are summarised in Table 2, below.

Table 2: Adverse Events in ≥ 5% of Mavacamten-treated Patients in EXPLORER-HCM

	CAMZYOS N=123 n (%)	placebo N=128 n (%)
Cardiac disorders		
Atrial fibrillation	8 (6.5%)	9 (7.0%)
Palpitations	7 (5.7%)	9 (7.0%)
Infections		
Nasopharyngitis	10 (8.1)	6 (4.7)
Musculoskeletal		
Back pain	9 (7.3)	7 (5.5)
Nervous system		
Dizziness	21 (17.1%)	15 (11.7%)
Headache	14 (11.4%)	10 (7.8%)
Respiratory		
Cough	8 (6.5%)	4 (3.1%)
Dyspnea	8 (6.5%)	10 (7.8%)

A long-term open-label extension trial of 224 oHCM patients who completed the EXPLORER-HCM trial was conducted. After a mean duration of 32 weeks in this trial, the most common reported adverse events included, fatigue (6.7%), atrial fibrillation (4.9%), headache (4.9%), dyspnea (4.5%), and dizziness (4.0%). SAE were reported in 8.5% of patients, in which, 3 cases of cardiac failure and 2 of atrial fibrillation were noted. All three cases of cardiac failure resolved.

9 DRUG INTERACTIONS

9.2 Drug Interactions Overview

Mavacamten is primarily metabolized by CYP 2C19 and to a lesser extent by CYP 3A4. Moderate and strong CYP 3A4 inhibitors/inducers or CYP 2C19 inhibitors/inducers may significantly affect the exposure of mavacamten (see [7 WARNINGS AND PRECAUTIONS](#), Cardiovascular, and 9.4 Drug-Drug Interactions).

Drugs That Reduce Cardiac Contractility

Additive negative inotropic effects are expected when mavacamten used concomitantly with other drugs that reduce cardiac contractility

In the pivotal clinical trial, EXPLORER-HCM, 119 of 123 patients who were treated with CAMZYOS received concomitant therapy with one of either beta blockers (n=94), verapamil (n=19), or diltiazem (n=6) (see [14 CLINICAL TRIALS](#)). There is limited information available on the potential for a pharmacodynamic interaction between CAMZYOS and other drugs that also reduce cardiac

contractility. If treatment with a new negative inotrope is initiated, or if the dose of a negative inotrope is increased, in a patient receiving CAMZYOS, close medical supervision with echocardiographic monitoring of LVEF should be provided until stable doses and clinical response have been achieved (see [7 WARNINGS AND PRECAUTIONS](#), Cardiovascular).

Effect on Transporters

In vitro data indicate that mavacamten is not an inhibitor of major efflux transporters (P-gp, BCRP, BSEP, MATE1, or MATE2-K) at therapeutic concentrations, nor is it an inhibitor of major uptake transporters (organic anion transporting polypeptides [OATPs], organic cation transporters [OCTs], or organic anion transporters [OATs]) at therapeutic concentrations.

Effect on CYP Enzyme Substrates

Concomitant use of mavacamten may reduce plasma concentrations of drugs which are CYP 2B6, CYP 2C8, CYP 2C9, or CYP 2C19 substrates.

9.4 Drug-Drug Interactions

The drugs listed in this table are based on either drug interaction case reports or studies, or potential interactions due to the expected magnitude and seriousness of the interaction (i.e., those identified as contraindicated).

Table 3 Established or Potential Drug-Drug Interactions

[Proper/Common name]	Source of Evidence	Effect	Clinical comment
Omeprazole	CT	Co-administration of mavacamten (15 mg) with the weak CYP 2C19 inhibitor omeprazole (20 mg QD) resulted in a 48% increase in mavacamten AUC _{inf} with no effect on C _{max} in combined healthy normal metabolizers and rapid metabolizers for CYP 2C19.	Consider additional monitoring of LVEF for initiation of weak CYP 2C19 inhibitor and adjust the dose based on clinical assessment. Intermittent co-administration of weak CYP 2C19 inhibitors (such as omeprazole or esomeprazole) with mavacamten is not recommended. Avoid initiation of concomitant weak CYP 2C19 inhibitors in patients who are on stable treatment with 2.5 mg of mavacamten.

Ticlopidine	PBPK modeling	Co-administration of mavacamten (15 mg QD) with the strong CYP 2C19 inhibitor ticlopidine (219.57 mg BID) resulted in a 98% and 59% increase in mavacamten AUC _{TAU} and C _{max} , respectively, in virtual healthy CYP 2C19 normal metabolizers.	No clinical drug interaction study was conducted with moderate or strong CYP 2C19 inhibitors. Concomitant use of mavacamten with moderate or strong CYP 2C19 inhibitors is contraindicated due to risk of heart failure.
Verapamil	CT	Co-administration of mavacamten (25 mg) with the moderate CYP 3A4 inhibitor verapamil sustained release (240 mg) resulted in a 16% and 52% increase in mavacamten AUC _{inf} and C _{max} , respectively, in combined healthy intermediate metabolizers and normal metabolizers for CYP 2C19.	Consider additional monitoring of LVEF when initiating a moderate CYP 3A4 inhibitor and adjust the dose based on clinical assessment. Avoid concomitant use of mavacamten in patients on combination therapy of a calcium channel blocker (e.g., verapamil, diltiazem) and a beta blocker.
Diltiazem	PBPK modeling	Co-administration of mavacamten (15 mg QD) with the moderate CYP 3A4 inhibitor diltiazem (60 mg TID) was predicted to result in a 19% and 12% increase in mavacamten AUC _{TAU} and C _{max} , respectively, in healthy intermediate or normal metabolizers of CYP 2C19. In healthy CYP 2C19 poor metabolizers, a 55% and 42% increase in mavacamten AUC _{TAU} and C _{max} , respectively, was predicted.	Consider additional monitoring of LVEF when initiating a moderate CYP 3A4 inhibitor and adjust the dose based on clinical assessment. Avoid initiation of concomitant moderate CYP 3A4 inhibitors in patients who are on stable treatment with 2.5 mg of mavacamten.

Itraconazole	PBPK modeling	Co-administration of mavacamten (15 mg QD) with the strong CYP 3A4 inhibitor itraconazole (200 mg QD) was predicted to result in an increase of up to 76% and 54% in AUC _{TAU} and C _{max} , respectively in healthy CYP 2C19 poor metabolizers. In CYP 2C19 normal metabolizers an increase of 29% and 17% in AUC _{TAU} and C _{max} , respectively was predicted.	<p>Concomitant use of mavacamten with a strong CYP 3A4 inhibitor is contraindicated.</p> <p>Concomitant use with a strong CYP 3A4 inhibitor may increase the risk of heart failure due to systolic dysfunction.</p>
Rifampin	PBPK modeling	Co-administration of mavacamten (a single 15-mg dose) with a strong CYP 3A4 and CYP 2C19 inducer (rifampin 600 mg daily dose) following a 7-day lead-in induction period, was predicted to result in a decrease in mavacamten of up to 69% and 7% in AUC _{0-T} and C _{max} , respectively in CYP 2C19 normal metabolizers and poor metabolizers.	<p>Mavacamten is contraindicated with concomitant use of moderate or strong inducers to both CYP 2C19 and CYP 3A4.</p> <p>Concomitant use with a moderate to strong inducer of CYP 2C19 and/or CYP 3A4 may reduce the efficacy of mavacamten.</p>

Carbamazepine	PBPK modeling	Co-administration of mavacamten (a single 15-mg dose) with the strong CYP 3A4 inducer carbamazepine (400 mg BID) following a 14-day lead-in induction period, was predicted to result in a decrease in mavacamten of 13% (range 9% to 28%) and 1% (range 0% to 2%) in AUC_{0-T} and C_{max} , respectively in CYP 2C19 normal metabolizers. In CYP 2C19 poor metabolizers a decrease of 30% (range 30% to 44%) and 1% (range 0% to 2%) in AUC_{0-T} and C_{max} was predicted.	Mavacamten is contraindicated with concomitant use of moderate or strong inducers to both CYP 2C19 and CYP 3A4. Concomitant use with a moderate to strong inducer of CYP 2C19 and/or CYP 3A4 may reduce the efficacy of mavacamten.
Midazolam	CT	Co-administration of a 16-day course of mavacamten (25 mg on days 1 and 2, followed by 15 mg daily for 14 days) resulted in a 13% and 7% in AUC_{inf} and C_{max} of midazolam (substrate of CYP 3A4), respectively, in healthy CYP 2C19 normal metabolizers.	Dosage adjustment may be required in the targeted patient population.
Ethinyl estradiol and norethindrone	CT	In healthy women, co-administration of a 17-day course of mavacamten (25 mg on days 1 and 2, followed by 15 mg daily for 15 days) did not decrease the exposure of a single dose of ethinyl estradiol (35 mcg) and norethindrone (1 mg), which are the components of typical oral contraceptives and substrates for CYP 3A4.	No dosage adjustment is required.

Legend: CT = Clinical Trial; PBPK = Physiologically Based Pharmacokinetic.

The PBPK model was developed based on *in vitro* and *in vivo* data from single and multiple dose studies

and drug-drug interaction studies with omeprazole and verapamil. No clinical drug-drug interaction studies with strong CYP 2C19 or CYP 3A4 inhibitors or inducers were available.

9.5 Drug-Food Interactions

Refer to Section [10.3 Pharmacokinetics](#) for details on the effect of food on the absorption of mavacamten.

9.6 Drug-Herb Interactions

Interactions with herbal products have not been established.

9.7 Drug-Laboratory Test Interactions

Interactions with laboratory tests have not been established.

10 CLINICAL PHARMACOLOGY

10.1 Mechanism of Action

Mavacamten is a selective, allosteric, and reversible cardiac myosin inhibitor. Mavacamten modulates the number of myosin heads that can enter power generating states, reducing force-producing systolic and residual diastolic cross-bridge formation. Mavacamten also shifts the overall myosin population towards an energy-sparing, but recruitable, super-relaxed state. Excess cross-bridge formation and dysregulation of the super-relaxed state of myosin are mechanistic hallmarks of HCM, which can result in hypercontractility, impaired relaxation, excess energy consumption, and myocardial wall stress. In HCM patients, myosin inhibition with mavacamten may attenuate myocardial hypercontractility, and can improve myocardial relaxation, reduce dynamic LVOT obstruction, and improve cardiac filling pressures and cardiac structure, the cardiac biomarkers, including NT-proBNP, and exercise capacity, as measured by peak oxygen uptake (pVO_2).

10.2 Pharmacodynamics

LVEF

A reduction in left ejection fraction may be expected with mavacamten treatment. In the EXPLORER-HCM trial, mean resting LVEF was 74% at baseline in both treatment groups. Consistent with the mechanism of action of CAMZYOS, the mean absolute change from baseline in LVEF was -4% in the CAMZYOS group and 0% in the placebo group over the 30-week treatment period. At Week 38, following an 8-week interruption of trial drug, mean LVEF was similar to baseline for both treatment groups.

LVOT obstruction

In the EXPLORER-HCM trial, patients achieved reductions in mean post-exercise resting and Valsalva-provoked LVOT (vLVOT) gradient, which were sustained throughout the 30-week trial. At Week 30, mean change from baseline in post-exercise LVOT resting and Valsalva LVOT gradients were -47 mmHg and -49 mmHg, respectively, for the CAMZYOS group, and -10 mmHg and -12 mmHg, respectively, for the placebo group. These differences were highly significant in favour of mavacamten. Significant vLVOT gradient differences were observed by Week 4 and then maintained throughout the study treatment period. At Week 38, following 8 weeks of trial drug washout, mean LVOT gradients were similar to baseline for both treatment groups.

Cardiac Structure

In EXPLORER-HCM study, echocardiographic measurements of cardiac structure showed a mean reduction from baseline at Week 30 in left ventricular mass index (LVMI) in the mavacamten group of -7.4 g/m^2 , versus an increase in LVMI in the placebo group of $+9.0 \text{ g/m}^2$. There was also a mean reduction from baseline in left atrial volume index (LAVI) in the mavacamten group of -7.5 mL/m^2 , versus no change in the placebo group at -0.1 mL/m^2 .

Cardiac Biomarkers

In the EXPLORER-HCM trial, profound reductions in N-terminal pro-B-type natriuretic peptide (NT-proBNP), a biomarker of cardiac wall stress, were observed by Week 4, from 783 ng/L at baseline to 242 ng/L, and sustained through to the end of study treatment. At Week 30, mavacamten treatment was associated with a median decrease of NT-proBNP from baseline of 556 ng/L, compared to a median decrease of only 5 ng/L for placebo.

Cardiac Electrophysiology

In a Phase 1 multiple ascending dose study in healthy subjects, sustained exposure to mavacamten at suprathreshold levels (18.5 mg and 25 mg once daily for up to 28 days) leading to marked depression of systolic function was associated with QTc prolongation.

In HCM patients, the QT interval may be intrinsically prolonged due to the underlying disease, in association with ventricular pacing, or in association with drugs with potential for QT prolongation commonly used in HCM population. An exposure-response analysis across clinical studies in HCM patients treated with mavacamten did not demonstrate a concentration dependent QTc prolongation in the therapeutic exposure range. The effects of co-administration of CAMZYOS with QT prolonging drugs or in patients with potassium channel variants resulting in a long QT interval have not been characterized.

10.3 Pharmacokinetics

Mavacamten has a variable terminal $t_{1/2}$ that depends on CYP 2C19 metabolic status (6-9 days in normal metabolizers (NMs) and 23 days in poor metabolizers (PMs)). Exposure to mavacamten increased approximately dose proportionally between 2 mg and 48 mg. The geometric mean accumulation was approximately 7-fold for AUC_{0-24h} and 2 to 2.6-fold for C_{max} after multiple once-daily dosing of mavacamten.

Table 4 Summary of mavacamten Pharmacokinetic Parameters in Healthy Volunteers after Multiple Doses at Steady State

PK Parameter on Day 28	C_{max} (ng/mL)	T_{max} (h)	$t_{1/2}$ (h)	AUC_{0-24} (ng.hr/mL)	CL/F (mL/hr)	Vd/F (L)
12.5 mg dose	412	2.7	45.4	6185	2021	129

Legend: C_{max} = Maximum observed measured plasma concentration over time spans specified; T_{max} = Time of the maximum observed plasma concentration; $t_{1/2}$ = Apparent first-order terminal elimination half-life; AUC_{0-24} = Area under the plasma concentration-time curve over the last 24-h dosing interval CL/F= Apparent clearance, calculated as $Dose/AUC_{0-\infty}$; Vd/F= Apparent volume of distribution

Absorption: Mavacamten is rapidly absorbed (T_{max} of 1 hour) after oral administration with an

estimated oral bioavailability of approximately 85% within the clinical dose range. The increase in mavacamten exposure is generally dose proportional after once daily doses of mavacamten (2 mg to 48 mg).

Effect of food

Administration of the 15 mg mavacamten capsule with a high-fat calorie meal delayed absorption by 3 hours (median T_{max} of 4 hours) and decreased the C_{max} and AUC_T by 55% and 12% respectively. CAMZYOS may be administered without regard to food.

Distribution: Specific studies to assess distribution of mavacamten have not been conducted in humans. Plasma protein binding of mavacamten is 97% to 98% as observed in a clinical study.

Based upon measurements of mavacamten in semen of 10 male subjects who received either an 18.5 mg (n=4) or 25 mg (n=6) once daily dosing for up to 28 days, the mean (SD) mavacamten semen-to-plasma ratio was 0.039 (0.0047) and 0.044 (0.016), respectively.

Metabolism: Mavacamten is extensively metabolized, primarily through CYP 2C19 (74%), CYP 3A4 (18%), and CYP 2C9 (8%). Three metabolites have been detected in human plasma. The exposure of the most abundant metabolite MYK 1078 in human plasma was less than 4% of the exposure of mavacamten, and the other two metabolites had exposures less than 3% of the exposure of mavacamten indicating the metabolites would have minimal to no impact on the overall activity of mavacamten.

Elimination: Mavacamten is cleared from plasma primarily by metabolism through cytochrome P450 enzymes. Terminal half-life is 6-9 days in CYP 2C19 NMs or 23 days in PMs, depending on CYP 2C19 metabolic status. Drug accumulation occurs with an accumulation ratio about 2-fold for C_{max} and about 7-fold for AUC in CYP 2C19 NMs. The accumulation depends on the metabolism status for CYP 2C19 with the largest accumulation observed in CYP 2C19 PMs. At steady-state, the peak-to-trough plasma concentration ratio with once daily dosing is approximately 1.5. Intersubject PK variability is moderate, with a coefficient of variation for exposure in of approximately 30 to 50% for C_{max} and AUC.

Following a single 25 mg dose of ¹⁴C labeled mavacamten, 7% and 85% of the total radioactivity was recovered in the feces and urine, respectively. Unchanged drug accounted for approximately 1% and 3% of the administered dose in the feces and urine, respectively.

Special Populations and Conditions

- **Pediatrics:** The safety and efficacy of CAMZYOS in pediatric patients (< 18 years of age) have not been established, therefore, Health Canada has not authorized an indication for pediatric use (see [1 INDICATIONS](#)).
- **Geriatrics:** Safety, effectiveness, and pharmacokinetics were appeared to be consistent between elderly patients (≥ 65 years) and younger patients (18 to < 65 years) (see [7 WARNINGS AND PRECAUTIONS](#)).

- **Genetic Polymorphism**

CYP 2C19 Poor Metabolizers (PMs): After a single dose of 15 mg mavacamten, C_{max} and AUC_{inf} increased by 47% and 241%, respectively, in CYP 2C19 PMs compared to normal metabolizers (NMs). Mean half-life is prolonged in CYP 2C19 PMs compared to NMs (23 days vs 6-9 days, respectively). The prevalence of CYP 2C19 PMs is approximately 2% in Caucasians, and 18% in the Asian population.

- **Hepatic Insufficiency:** A single dose PK study was conducted in patients with mild (Child-Pugh Class A) or moderate (Child-Pugh Class B) hepatic impairment, as well as a control group with normal hepatic function. Mavacamten exposures (AUC) increased 3.24-fold and 1.87-fold in patients with mild and moderate hepatic impairment, respectively, compared to patients with normal hepatic function. There was no effect of hepatic function on C_{max} , consistent with no change in the rate of absorption and/or volume of distribution. A dedicated PK study has not been conducted in patients with severe (Child-Pugh Class C) hepatic impairment (see [4.2 Recommended Dose and Dosage Adjustment](#), Special Populations).
- **Renal Insufficiency:** Approximately 3% of a mavacamten dose is excreted in the urine as parent drug. A dedicated PK study has not been conducted in patients with renal impairment. A population PK analysis estimated a 1.17-fold increase in the median steady-state exposure in subjects with eGFR of 45 mL/min/1.73 m², relative to the reference subject who had a median eGFR of 95 mL/min/1.73 m² (see [4.2 Recommended Dose and Dosage Adjustment](#), Special Populations).

11 STORAGE, STABILITY AND DISPOSAL

Store at room temperature (15 to 30°C).

Keep out of reach and sight of children.

PART II: SCIENTIFIC INFORMATION

13 PHARMACEUTICAL INFORMATION


Drug Substance

Proper name: mavacamten

Chemical name: 3-(1-methylethyl)-6-[[[(1S)-1-phenylethyl]amino]-2,4(1H,3H)-pyrimidinedione.

Molecular formula and molecular mass: C₁₅H₁₉N₃O₂ and 273.33 g/mol

Structural formula:



Physicochemical properties: mavacamten is a white to off-white powder that is practically insoluble in water and aqueous buffers, sparingly soluble in methanol and ethanol, and freely soluble in DMSO and NMP.

14 CLINICAL TRIALS

14.1 Clinical Trials by Indication

Obstructive Hypertrophic Cardiomyopathy

Table 5 - Summary of pivotal trial in oHCM

Study Name	Study design	Dosage, route of administration and duration	Study subjects (n)	Mean age (Range)	Sex
EXPLORER-HCM	Phase 3, double-blind, randomized, placebo-controlled, multicenter, international, parallel-group trial to evaluate the safety and efficacy in patients with oHCM over 30 weeks.	Oral starting dose of 5 mg once daily of CAMZYOS or matching placebo. The dose could then be either down-titrated to 2.5 mg or up-titrated to a maximum dose of 15 mg once daily based on PD responses, specifically LVEF and Valsalva LVOT gradient, as well as mavacamten plasma concentration.	123 patients received CAMZYOS	CAMZYOS: 59 years (26, 82)	CAMZYOS: 66 (54%) males 57 (46%) females
			128 patients received placebo	Placebo: 59 years (18, 81)	Placebo: 83 (65%) males 45 (35%) females

The efficacy of CAMZYOS was evaluated in EXPLORER-HCM, a Phase 3, double-blind, randomized, placebo-controlled, multicenter, international, parallel-group trial in 251 adult patients with symptomatic NYHA Class II and III obstructive hypertrophic cardiomyopathy (oHCM), with left

ventricular ejection fraction (LVEF) \geq 55%, and LVOT peak gradient \geq 50 mmHg at rest or with provocation. The majority of patients received conventional background HCM treatment, for 96% of patients in the mavacamten arm (beta-blockers 76%, calcium channel blockers 20%), and 87% in the placebo arm (beta-blockers 74%, calcium channel blockers 13%).

Patients were randomized in a 1:1 ratio to receive either a starting dose of 5 mg of CAMZYOS (n=123) or placebo (n=128) once daily for 30 weeks, with a starting dose of 5 mg and the dose periodically adjusted to optimize the patient's therapeutic response, as measured by a decrease in LVOT gradient with Valsalva manoeuvre, and to maintain LVEF \geq 50%, while being further informed by plasma concentrations of CAMZYOS.

Treatment assignment was stratified by severity of disease at baseline, i.e., NYHA Class (II or III), current treatment with beta-blockers (yes or no), type of ergometer (treadmill or exercise bicycle) used for assessment of peak oxygen consumption (pVO₂), and consent for the cardiac magnetic resonance substudy (yes or no). Patients on background dual treatment with beta-blockers and calcium channel blockers or taking disopyramide or ranolazine were excluded. Patients with known infiltrative or storage disorders causing cardiac hypertrophy that may mimic oHCM, such as Fabry disease, amyloidosis, or Noonan syndrome with LV hypertrophy, were also excluded.

Patient demographics and baseline characteristics are presented in Table 6 and Table 7, below.

Table 6- Patient Demographics in EXPLORER-HCM

Characteristics	CAMZYOS N=123	Placebo N=128
Age (years)		
Mean (minimum, maximum)	59 (26, 82)	59 (18, 81)
Sex, n (%)		
Male	66 (54)	83 (65)
Female	57 (46)	45 (35)
Mean body mass index, kg/m ² (SD)	30 (5)	29 (6)
Mean heart rate, beats/min (SD)	63 (10)	62 (11)
Mean blood pressure, mm Hg (SD)		
Systolic	128 (16)	128 (15)
Diastolic	75 (11)	76 (10)
Race, n (%)		
Alaska Native	0	1 (1)
Asian	4 (3)	2 (1)
Black or African American	1 (1)	5 (4)
Unknown	3 (2)	6 (5)
White	115 (94)	114 (89)
Ethnicity		
Hispanic or Latino	8 (7)	4 (3)
Not Hispanic or Latino	114 (93)	119 (93)
Family history of HCM	33 (27)	36 (28)
Medical history, n (%)		
Hypertension	57 (46)	53 (41)
Hyperlipidemia	27 (22)	39 (30)
Coronary artery disease	12 (10)	6 (5)
Obesity	15 (12)	14 (11)
Type 2 diabetes	6 (5)	7 (6)
Asthma	17 (14)	11 (9)

Chronic obstructive pulmonary disease	2 (2)	3 (2)
---------------------------------------	-------	-------

Table 7 - Baseline Disease Characteristics in EXPLORER-HCM

Characteristics	CAMZYOS N=123	Placebo N=128
NYHA functional class, n (%)		
NYHA Class II	88 (72)	95 (74)
NYHA Class III	35 (28)	33 (26)
pVO ₂ , mL/kg per min	18.9 (4.9)	19.9 (4.9)
Background HCM treatment, n (%)		
Beta-blockers	94 (76)	95 (74)
Non-hydropyridine calcium channel blockers	25 (20)	17 (13)
Echocardiography parameters		
Mean interventricular septum thickness, mm (SD)	17 (3)	17 (3)
Systolic anterior motion of mitral valve, n (%)	97 (82)	102 (81)
Mean LAVI (SD), mL/m ²	40 (12)	41 (14)
Mean LVEF (%)	74 (6)	74 (6)
Mean LVOT gradient with Valsalva maneuver, mmHg (SD)	72 (32)	74 (32)
Mean LVOT gradient post-exercise mmHg (SD)	86 (34)	84 (36)
Critical cardiac history		
Atrial fibrillation, n (%)	12 (10)	23 (18)
Implantable cardioverter defibrillator (ICD), n (%)	27 (22)	29 (23)
Prior invasive septal reduction therapies, n (%)	11 (9)	8 (6)

SD = Standard deviation

In the EXPLORER-HCM trial, 81% (100/123) of patients were receiving either the 5 mg or 10 mg dose at the end of the treatment period, with 49% (60/123) receiving the 5 mg dose.

Primary endpoint

The primary endpoint was comprised of a composite of change at Week 30 from baseline in exercise capacity measured by pVO₂ and symptoms measured by NYHA functional classification, defined as an improvement of pVO₂ by ≥ 1.5 mL/kg/min AND an improvement in NYHA class by at least 1 OR an improvement of pVO₂ by ≥ 3.0 mL/kg/min AND no worsening in NYHA class.

Peak VO₂ is a direct, objective, and reproducible measure of exercise capacity, and it has been endorsed for the clinical evaluation of the efficacy of therapeutic intervention in patients with HCM based on its high reproducibility, low variance, and a narrow dynamic range at maximal effort.

At Week 30, patients receiving CAMZYOS had significantly greater improvement compared to placebo group across the primary composite endpoint, see Table 8, below.

Table 8 - Results of EXPLORER-HCM in oHCM

Primary Endpoints	CAMZYOS N = 123 n (%)	Placebo N = 128 n (%)	Treatment difference (95% CI)	p-value
Patients Achieving Primary Endpoint at Week 30	45 (37%)	22 (17%)	19 (8.67, 30.13)	0.0005
Condition 1: Patients with Change from Baseline in $pVO_2 \geq 1.5$ mL/kg/min AND Improvement in NYHA Class ≥ 1 at Week 30, n (%)	41 (33%)	18 (14%)	19 (8.99 – 29.55)	
Condition 2: Patients with Change from Baseline in $pVO_2 \geq 3.0$ mL/kg/min AND No Worsening in NYHA Class at Week 30, n (%)	29 (24%)	14 (11%)	13 (3.39 – 21.89)	
Subgroup of Condition 2: Patients with Change from Baseline in $pVO_2 \geq 3.0$ mL/kg/min AND Improvement in NYHA Class ≥ 1 at Week 30, n (%)	25 (20%)	10 (8%)	12.5 (4.0 - 21.0)	

Although the benefit of mavacamten was smaller in patients on background beta blocker therapy compared to those who were not (attenuated improvement in pVO_2), analyses of other secondary endpoints (symptoms, LVOT gradient) suggest that patients might benefit from mavacamten treatment regardless of beta blocker use.

Secondary endpoints

The treatment effects of CAMZYOS on LVOT obstruction, functional capacity, and health status were also assessed by change from baseline through Week 30 in post-exercise LVOT peak gradient, change in pVO_2 , proportion of patients with improvement in NYHA class, Kansas City Cardiomyopathy Questionnaire 23 (KCCQ 23) Clinical Summary Score (CSS), and Hypertrophic Cardiomyopathy Symptom Questionnaire (HCMSQ) Shortness of Breath (SoB) domain score. At Week 30, patients receiving CAMZYOS had significantly greater improvement compared to placebo group across all secondary endpoints, see Table 9 and Table 10, below.

Table 9 - Results of Secondary Endpoints in EXPLORER-HCM in oHCM

Secondary Endpoints	Mavacamten N=123	Placebo N=128	Treatment Difference (95% CI)	p-value
Change from baseline post-exercise LVOT peak gradient at Week 30, mmHg, Mean (SD)	-47 (40)	-10 (30)	-35 (-43, -28)	<0.0001
Change from baseline to Week 30 in pVO ₂ , mL/kg/min, Mean (SD)	1.4 (3)	-0.05 (3)	1.4 (0.6, 2)	<0.0006
Patients with improvement of NYHA class \geq 1 at Week 30, N, (%)	80 (65%)	40(31%)	34 (22, 45)	<0.0001

Table 10: Results of Patients Reported Outcomes Secondary Endpoints in EXPLORER-HCM in oHCM

Secondary Endpoints	Baseline, Mean (SD)		Week 30, Mean (SD)		Change from Baseline to Week 30, Mean (SD)		Difference, (95% CI) and p-value
	Mavacamten	Placebo	Mavacamten	placebo	Mavacamten	Placebo	
KCCQ-23 CSS [†]	N=99 71 (16)	N=97 71 (19)	N=108 82 (16)	N= 113 73 (20)	N=92 14 (14)	N=88 4 (14)	9 (5, 13) p<0.0001
HCMSQ SoB [‡]	N=108 4.9 (2.5)	N=109 4.5 (3.2)	N=92 2.0 (2.6)	N=97 3.7 (3.0)	N=85 -2.8 (2.7)	N=88 -0.9 (2.4)	-1.8 (-2.4, -1.2) p< 0.0001

[†] The KCCQ-23 CSS is a validated patient reported outcome in oHCM and is composed of the physical limitations and the total symptom burden scores of the KCCQ-23.

The Clinical Summary Score (CSS) ranges from 0 to 100, with higher scores representing better health status.

[‡] The HCMSQ SoB domain score is a validated patient reported outcome in oHCM and measures frequency and severity of shortness of breath. The HCMSQ Shortness of Breath (SoB) domain score ranges from 0 to 18, with lower scores representing less shortness of breath.

At Week 30, 50% (61 of 123) of patients treated with CAMZYOS attained NYHA Class I status, compared to a 21% (27 of 128) of patients in the placebo group. NYHA Class changes from baseline to Week 30 are shown in Table 11, below.


Table 11: NYHA Class Status at Baseline, and achieved at Week 30

NYHA* Class	Mavacamten (N = 123)			Placebo (N = 128)		
	Class II n (%)	Class III n (%)	Total n (%)	Class II n (%)	Class III n (%)	Total n (%)
Baseline	88 (71.5)	35 (28.5)	123 (100.0)	95 (74.2)	33 (25.8)	128 (100.0)
Week 30, n (%)						
Class I	52 (42.3)	9 (7.3)	61 (49.6)	24 (18.8)	3 (2.3)	27 (21.1)
Class II	33 (26.8)	19 (15.4)	52 (42.3)	61 (47.7)	13 (10.2)	74 (57.8)
Class III	1 (0.8)	7 (5.7)	8 (6.5)	9 (7.0)	16 (12.5)	25 (19.5)
Missing	2 (1.6)	0	2 (1.6)	1 (0.8)	1 (0.8)	2 (1.6)

*NYHA = New York Heart Association


The KCCQ-23 CSS favored CAMZYOS compared to placebo at Week 30. The mean improvement from baseline on the KCCQ-23 CSS was significantly greater in the CAMZYOS group compared to placebo at Week 30 ($p < 0.0001$), with effects observed as early as 6 weeks and sustained over the full treatment period to Week 30, see Figure 4, below.

Figure 4: KCCQ-23 CSS: Mean Change from Baseline Over Time



The HCMSQ SoB domain favored CAMZYOS compared to placebo at Week 30. The mean improvement from baseline on the HCMSQ SoB domain was significantly greater in the CAMZYOS group compared to placebo at Week 30 ($p < 0.0001$), with effects observed as early as 4 weeks and sustained over the full treatment period to Week 30, see Figure 5, below.

Figure 5: HCMSQ Shortness of Breath Domain: Mean Change from Baseline Over Time



15 MICROBIOLOGY

Not Applicable

16 NON-CLINICAL TOXICOLOGY

General Toxicology

The safety of mavacamten has been evaluated in rats and dogs dosed for up to 6 and 9 months respectively. Noted toxicities, including echocardiographic findings of reduced systolic performance and cardiac dilation, death, due to heart failure, and, in rats, increased heart weights likely secondary to cardiac hypertrophy in response to decreased contractility, were consistent with the mavacamten mechanism of action and primary pharmacological activity. Other findings included cardiac osseous metaplasia in rats and QTc prolongation in dogs. The NOAELs in the 6-month rat and 9-month dog studies were 0.3 and 0.18 mg/kg/day, respectively. Plasma exposures (AUC) at the NOAELs in rats and dogs were respectively 0.1× and 0.3× of the average exposures in humans at the MRHD.

Carcinogenicity

There was no evidence of carcinogenicity at the highest oral mavacamten doses tested in a 6-month rasH2 transgenic mouse study (3 mg/kg/day) or a 2-year rat study (0.6 mg/kg/day). Exposures (AUC) in mice were up to 3-fold higher than the MRHD, while exposures (AUC) in rats were only 0.2-fold compared to the MRHD.

Genotoxicity

Neither mavacamten nor its isomer were found to be genotoxic in a reverse mutation bacterial test (Ames test), a human *in vitro* lymphocyte clastogenicity assay, or a rat *in vivo* micronucleus assay (mavacamten only).

Reproductive and Developmental Toxicology

Fertility

There was no evidence of effects of mavacamten on mating and fertility in male or female rats at oral doses up to 1.2 mg/kg/day. Parental plasma exposures (AUC) of mavacamten at the highest doses tested were only 0.8× the exposures in humans at the MRHD.

Embryo-fetal Development

Mavacamten administered orally to pregnant rats during the period of organogenesis decreased mean fetal body weight and increased post-implantation loss and fetal malformations (heart malformations, total situs inversus, fused sternbrae and other skeletal malformations) in the high dose group (1.5 mg/kg/day). No post-natal adverse effects were noted in surviving pups nor in dams. Maternal plasma exposures (AUC) at the no effect dose for embryo-fetal development in rats (0.75 mg/kg/day) were only 0.3× the average exposures in humans at the MRHD.

Mavacamten administered orally to pregnant rabbits during the period of organogenesis caused fetal malformations (dilatation of pulmonary trunk and/or aortic arch, higher incidence of fused sternbrae) at doses of 1.2 mg/kg/day and higher. A higher incidence of pre- and post-implantation loss was noted at the highest dose (2.0 mg/kg/day). Mavacamten was also associated with cardiac deaths (mild bilateral ventricular dilatation noted in both cases, without microscopic evidence of heart failure) in 9% of pregnant rabbits receiving the highest dose after 10 and 12 days of treatment. Plasma exposure (AUC) at the no effect dose for embryo-fetal development in rabbits (0.6 mg/kg/day) was only 0.4× relative to average exposures in humans at the MRHD.

Together, these results suggest that mavacamten has the potential for significant toxicity to human embryos and fetuses, albeit with remaining uncertainties related to timing and duration of exposure. Mavacamten was shown to cross the placenta and was measured in embryonic and extra-embryonic tissues. Rabbit fetuses were estimated to be exposed to 0.15-fold of the maternal exposure. It is unknown if continuous exposure before conception and throughout gestation would lead to accumulation of mavacamten in embryonic tissues. It is unknown if mavacamten transfers to maternal milk.

PATIENT MEDICATION INFORMATION

READ THIS FOR SAFE AND EFFECTIVE USE OF YOUR MEDICINE

Pr **CAMZYOS™**

mavacamten capsules

Read this carefully before you start taking **CAMZYOS** and each time you get a refill. This leaflet is a summary and will not tell you everything about this drug. Talk to your healthcare professional about your medical condition and treatment and ask if there is any new information about **CAMZYOS**.

Serious Warnings and Precautions

CAMZYOS may cause serious side effects, including:

- **Heart failure** (a condition where the heart cannot pump blood with enough force): It is a serious condition that can lead to death. You must have echocardiograms before you take your first dose and regularly during your treatment with CAMZYOS. This is to help your healthcare professional understand how your heart is responding to CAMZYOS. People who develop a serious infection or irregular heartbeat have a greater risk of heart failure during treatment with CAMZYOS. Tell your healthcare professional **right away** if you develop new or worsening:
 - shortness of breath;
 - swelling in your legs;
 - chest pain;
 - a racing sensation in your heart (palpitations);
 - fatigue;
 - rapid weight gain.
- **The risk of heart failure is also increased when CAMZYOS is taken with certain other medicines.** Tell your healthcare professional about all of the medicines you take before and during your treatment with CAMZYOS. These include prescribed medications and those obtained over-the-counter. Your healthcare professional will tell you if you can take them during your treatment with CAMZYOS.
- **Because of the serious risk of heart failure,** your healthcare professional will make sure you understand how to take CAMZYOS safely before you start your treatment. This includes returning for echocardiograms when advised by your healthcare professional.

See the **Serious side effects and what to do about them** table for more information on this and other serious side effects.

What is CAMZYOS used for?

CAMZYOS is used to treat obstructive hypertrophic cardiomyopathy (oHCM) in adults with symptoms of the disorder.

oHCM is a disorder in which the heart muscle contracts too much and becomes abnormally thick. It usually affects the septum, which is the wall between the two lower chambers of the heart (ventricles).

The thickened heart muscle can become stiff and prevent the heart from adequately pumping blood to the rest of the body.

How does CAMZYOS work?

CAMZYOS belongs to a group of medicines called cardiac myosin inhibitors. It works by relaxing the muscles in your heart, which improves your heart's ability to pump blood to the rest of your body. This may improve your symptoms and your ability to be active.

What are the ingredients in CAMZYOS?

Medicinal ingredient: Mavacamten

Non-medicinal ingredients: Croscarmellose sodium, hypromellose, magnesium stearate (non-bovine), mannitol, and silicon dioxide.

Capsule shell 2.5 mg: black iron oxide, gelatin (bovine and/ or porcine), red iron oxide, titanium dioxide.

Capsule shell 5 mg: gelatin (bovine and/ or porcine), titanium dioxide, yellow iron oxide.

Capsule shell 10 mg: gelatin (bovine and/ or porcine), red iron oxide, titanium dioxide.

Capsule shell 15 mg: black iron oxide, gelatin (bovine and/ or porcine), titanium dioxide.

Imprinting Ink: black iron oxide, potassium hydroxide, propylene glycol, shellac, strong ammonia solution.

CAMZYOS comes in the following dosage forms:

Capsules: 2.5 mg, 5 mg, 10 mg and 15 mg.

Do not use CAMZYOS if:

- you are allergic to mavacamten or any of the other ingredients in CAMZYOS.
- you are taking medicines that are:
 - moderate and strong CYP2C19 inhibitors, or strong CYP3A4 inhibitors;
 - moderate and strong CYP2C19 or CYP3A4 inducers.Ask your healthcare professional if you are unsure.
- you are pregnant, planning to become pregnant, or think you might be pregnant.

To help avoid side effects and ensure proper use, talk to your healthcare professional before you take CAMZYOS. Talk about any health conditions or problems you may have, including if you:

- are taking any prescription and over-the-counter medicines, vitamins, herbal supplements and grapefruit juice. Especially tell your healthcare professional if you take:
 - medicines used to treat excess stomach acid (e.g., omeprazole or esomeprazole)
 - medicines used to treat high blood pressure (e.g., beta blockers, verapamil, diltiazem).
 - disopyramide (used to treat an irregular heart beat).
 - ranolazine (used to treat chest pain)
- have an irregular heart beat (arrhythmia).
- currently suffer from another serious illness (e.g., serious infection).
- have severe liver or kidney problems.

- are able to become pregnant.
- are breastfeeding or planning to breastfeed.

Other warnings you should know about:

Contraception (for female patients): If you are female and are able to become pregnant:

- your healthcare professional will ask you to take a pregnancy test to confirm that you are not pregnant before you start your treatment with CAMZYOS;
- you should use an effective birth control method (contraception) during your treatment with CAMZYOS and for up to 4 months after your last dose of CAMZYOS.

Pregnancy: Do not take CAMZYOS during pregnancy. CAMZYOS may harm your unborn baby. Your healthcare professional will discuss the potential risks with you. If you discover that you are pregnant while taking CAMZYOS, stop taking this medicine and tell your healthcare professional **right away**.

Breast-feeding: It is not known if CAMZYOS can pass into breast milk and harm your baby. You and your healthcare professional will decide whether you should take CAMZYOS or breastfeed. You should not do both at the same time. If the decision is to take CAMZYOS, you should not breastfeed during treatment with CAMZYOS and for up to 4 months after the last dose of CAMZYOS.

Driving or using machines: CAMZYOS may make you feel dizzy. Avoid driving, using machinery, or doing activities that require you to be alert if you feel dizzy.

Tell your healthcare professional about all the medicines you take, including any drugs, vitamins, minerals, natural supplements or alternative medicines.

Serious Drug Interactions

Do not take CAMZYOS with medicines that are:

- moderate and strong CYP2C19 inhibitors, or strong CYP3A4 inhibitors (e.g., ticlopidine, itraconazole, ketoconazole);
- moderate and strong CYP2C19 or CYP3A4 inducers (e.g., rifampicin, carbamazepine).

Ask your healthcare professional if you are unsure you are taking any of these types of medicines.

The following may interact with CAMZYOS:

- medicines used to treat excess stomach acid (e.g., omeprazole or esomeprazole)
- disopyramide (used to treat an irregular heart beat)
- ranolazine (used to treat chest pain)
- midazolam (used to produce sleepiness and relieve anxiety before surgery)

Do not stop or change the dose of a medicine or start a new medicine without telling your healthcare professional. This is because some medicines can affect the way CAMZYOS works. Some medicines can increase the levels of CAMZYOS in your body and make it more likely for you to get side effects that are possibly severe. Some medicines can reduce the levels of CAMZYOS in your body and may reduce its beneficial effects.

Know the medicines you take. Keep an updated list of them to show or tell your healthcare professional and pharmacist when you get a new medicine.

How to take CAMZYOS:

- Take CAMZYOS exactly as your healthcare professional has told you to take it.
- Take CAMZYOS once a day, with or without food.
- Swallow the capsule whole with water. Do not break, open, or chew the capsule.
- Do not change your dose of CAMZYOS without talking to your healthcare professional first.
- Your healthcare professional may change your dose, temporarily stop, or permanently stop your treatment with CAMZYOS if you have certain side effects. Tell your healthcare professional **right away** if you are experiencing any side effects during your treatment.

Usual dose:

- Your healthcare professional will decide on the dose that is right for you. The dose of CAMZYOS prescribed to you will depend on your heart function. Your healthcare professional will assess your heart function before you take CAMZYOS and regularly during treatment and adjust your dose during treatment if needed.
- The recommended starting dose is 5 mg once a day.

Overdose:

Symptoms of an overdose with CAMZYOS may include:

- fainting;
- low blood pressure, which may make you feel dizzy or light-headed;
- heart unable to pump enough blood to the rest of the body (heart failure);
- heart suddenly stops beating (cardiac arrest).

If you think you, or a person you are caring for, have taken too much CAMZYOS, contact a healthcare professional, hospital emergency department, or regional poison control centre immediately, even if there are no symptoms.

Missed Dose:

If you miss a dose, take it as soon as possible and take your next scheduled dose at your usual time the next day. **Do not take two doses on the same day to make up for a missed dose.**

What are possible side effects from using CAMZYOS?

These are not all the possible side effects you may have when taking CAMZYOS. If you experience any side effects not listed here, tell your healthcare professional.

Side effects with CAMZYOS may include:

- dizziness
- headache
- fainting
- fatigue

- cough

Serious side effects and what to do about them			
Symptom / effect	Talk to your healthcare professional		Stop taking drug and get immediate medical help
	Only if severe	In all cases	
COMMON			
Heart Failure (a condition where the heart cannot pump blood with enough force): shortness of breath, fatigue and weakness, swelling in ankles, legs and feet, chest pain, cough, fluid retention, rapid weight gain, lack of appetite, nausea, rapid or irregular heartbeat, reduced ability to exercise.		✓	

If you have a troublesome symptom or side effect that is not listed here or becomes bad enough to interfere with your daily activities, tell your healthcare professional.

Reporting Side Effects

You can report any suspected side effects associated with the use of health products to Health Canada by:

- Visiting the Web page on Adverse Reaction Reporting (<https://www.canada.ca/en/health-canada/services/drugs-health-products/medeffect-canada/adverse-reaction-reporting.html>) for information on how to report online, by mail or by fax; or
- Calling toll-free at 1-866-234-2345.

NOTE: Contact your health professional if you need information about how to manage your side effects. The Canada Vigilance Program does not provide medical advice.

Storage:

- Store at room temperature (15°C to 30°C).
- Keep out of reach and sight of children.

If you want more information about CAMZYOS:

- Talk to your healthcare professional
- Find the full product monograph that is prepared for healthcare professionals and includes this Patient Medication Information by visiting the Health Canada website:

(<https://www.canada.ca/en/health-canada/services/drugs-health-products/drug-products/drug-product-database.html>); the manufacturer's website <https://www.bms.com/ca/en>, or by calling 1-866-463-6267.

This leaflet was prepared by Bristol-Myers Squibb Canada.

Last Revised November 8, 2022.