

PRODUCT MONOGRAPH
INCLUDING PATIENT MEDICATION INFORMATION

Pr **MYLAN-ZIPRASIDONE**

Ziprasidone Capsules

Capsules, 20 mg, 40 mg, 60 mg and 80 mg of Ziprasidone (as Ziprasidone Hydrochloride Monohydrate),
Oral

Antipsychotic Agent

Mylan Pharmaceuticals ULC
85 Advance Road
Etobicoke, ON
M8Z 2S6

Date of Initial Authorization
March 15, 2018

Date of Revision:
April 18, 2023

Submission Control No: 270286

RECENT MAJOR LABEL CHANGES

<u>4 Dosage and Administration, 4.1 Dosing Considerations</u>	04/2023
<u>7 Warnings and Precautions, Serotonin toxicity / Serotonin syndrome</u>	04/2023
<u>7 Warnings and Precautions, Skin</u>	04/2023
<u>7 Warnings and Precautions, 7.1.3 Pediatrics</u>	04/2023
<u>7 Warnings and Precautions, 7.1.4 Geriatrics</u>	04/2023

TABLE OF CONTENTS

Sections or subsections that are not applicable at the time of authorization are not listed.

RECENT MAJOR LABEL CHANGES 2

PART I: HEALTH PROFESSIONAL INFORMATION 4

1. INDICATIONS..... 4

 1.1 PEDIATRICS..... 4

 1.2 GERIATRICS 4

2. CONTRAINDICATIONS 4

3. SERIOUS WARNINGS AND PRECAUTIONS 5

4. DOSAGE AND ADMINISTRATION 5

 4.1 DOSING CONSIDERATIONS 5

 4.2 RECOMMENDED DOSE AND DOSAGE ADJUSTMENT 5

 4.4 ADMINISTRATION 6

 4.5 MISSED DOSE 7

5. OVERDOSAGE..... 7

6. DOSAGE FORMS, STRENGTHS, COMPOSITION AND PACKAGING 8

7. WARNINGS AND PRECAUTIONS 9

 7.1 SPECIAL POPULATIONS 19

 7.1.1 Pregnant Women 20

 7.1.2 Breast-feeding 21

 7.1.3 Pediatrics 21

 7.1.4 Geriatrics 21

8. ADVERSE REACTIONS 22

 8.1 ADVERSE REACTION OVERVIEW 22

8.2	CLINICAL TRIAL ADVERSE REACTIONS	23
8.2.1	Clinical Trial Adverse Reactions – Pediatrics	26
8.3	LESS COMMON CLINICAL TRIAL ADVERSE REACTIONS	29
8.5	POST-MARKET ADVERSE REACTIONS	30
9.	DRUG INTERACTIONS	31
9.2	DRUG INTERACTIONS OVERVIEW	31
9.3	DRUG BEHAVIOURAL INTERACTIONS.....	31
9.4	DRUG-DRUG INTERACTIONS	32
9.5	DRUG-FOOD INTERACTIONS.....	33
9.6	DRUG-HERB INTERACTIONS	33
9.7	DRUG-LABORATORY TEST INTERACTIONS	33
10.	CLINICAL PHARMACOLOGY	34
10.1	MECHANISM OF ACTION.....	34
10.2	PHARMACODYNAMICS.....	34
10.3	PHARMACOKINETICS	34
11.	STORAGE, STABILITY AND DISPOSAL	36
12.	SPECIAL HANDLING INSTRUCTIONS	37
PART II: SCIENTIFIC INFORMATION		38
13.	PHARMACEUTICAL INFORMATION	38
14.	CLINICAL TRIALS	39
14.1	TRIAL DESIGN AND STUDY DEMOGRAPHICS.....	39
14.2	STUDY RESULTS.....	39
14.3	COMPARATIVE BIOAVAILABILITY STUDIES.....	40
15.	MICROBIOLOGY	41
16.	NON-CLINICAL TOXICOLOGY.....	41
17.	SUPPORTING PRODUCT MONOGRAPHS	44
PATIENT MEDICATION INFORMATION		45

PART I: HEALTH PROFESSIONAL INFORMATION

1. INDICATIONS

- **Schizophrenia**

Mylan-Ziprasidone (ziprasidone hydrochloride) is indicated for the treatment of schizophrenia. The prescriber should consider the finding of ziprasidone's greater capacity to prolong the QT/QTc interval compared to other antipsychotic drugs (see **2 CONTRAINDICATIONS, 7 WARNINGS AND PRECAUTIONS**).

The efficacy of ziprasidone was established in short-term (4- and 6-week) controlled trials of schizophrenic inpatients (see **14 CLINICAL TRIALS**).

Mylan-Ziprasidone has been shown to be effective in maintaining clinical improvement during long-term therapy (1-year). The health professional who elects to use Mylan-Ziprasidone for extended periods should periodically re-evaluate the long-term usefulness of the drug for the individual patient.

1.1 Pediatrics

Pediatrics (<18 years of age): Based on the data submitted and reviewed by Health Canada, the safety and efficacy of ziprasidone hydrochloride in pediatric patients has not been established; therefore, Health Canada has not authorized an indication for pediatric use. (see **7.1.3 Pediatrics**).

1.2 Geriatrics

Mylan-Ziprasidone is not indicated for use in elderly patients with dementia (see **3 SERIOUS WARNINGS AND PRECAUTIONS** and **10.3 Pharmacokinetics, Special Populations and Conditions**).

The greater frequency of decreased hepatic, renal, or cardiac function, characteristic of the elderly, as well as concomitant disease and use of other drugs may impact the pharmacokinetics of ziprasidone hydrochloride in this population. Caution should be used when treating geriatric patients with Mylan-Ziprasidone (See **10.3 Pharmacokinetics, Special Populations and Conditions**, and **4 DOSAGE AND ADMINISTRATION**).

2. CONTRAINDICATIONS

- **QT Prolongation:** Because of ziprasidone's dose-related prolongation of the QT interval and the known association of fatal arrhythmias with QT prolongation by some other drugs, Mylan-Ziprasidone is contraindicated in patients with:
 - known history of QT prolongation (including congenital long QT syndrome);
 - recent acute myocardial infarction; or

- uncompensated heart failure (see 7 WARNINGS AND PRECAUTIONS, Cardiovascular).

Pharmacokinetic/pharmacodynamic studies between ziprasidone and other drugs that prolong the QT interval have not been performed. An additive effect of ziprasidone and other drugs that prolong the QT interval cannot be excluded. Therefore, ziprasidone should not be given with dofetilide, sotalol, quinidine, other Class Ia and III anti-arrhythmias, mesoridazine, thioridazine, chlorpromazine, droperidol, pimozide, sparfloxacin, gatifloxacin, moxifloxacin, halofantrine, mefloquine, pentamidine, arsenic trioxide, levomethadyl acetate, dolasetron mesylate, probucol or tacrolimus. Ziprasidone is also contraindicated with drugs that have demonstrated QT prolongation as one of their pharmacodynamic effects and have this effect described in their respective Product Monograph as a contraindication or a warning (see 7 WARNINGS AND PRECAUTIONS, Cardiovascular).

- **Hypersensitivity:** Mylan-Ziprasidone is contraindicated in patients who are hypersensitive to ziprasidone or to any ingredient in the formulation or component of the container. For a complete listing see 6 DOSAGE FORMS, STRENGTHS, COMPOSITION AND PACKAGING.

3. SERIOUS WARNINGS AND PRECAUTIONS

Serious Warnings and Precautions

Increased Mortality in Elderly Patients with Dementia

Elderly patients with dementia treated with atypical antipsychotic drugs are at an increased risk of death compared to placebo. Analyses of 13 placebo-controlled trials with various antipsychotics (modal duration of 10 weeks) in these patients showed a mean 1.6-fold increase in the death rate in the drug-treated patients. Although the causes of death were varied, most of the deaths appeared to be either cardiovascular (e.g., heart failure, sudden death) or infectious (e.g., pneumonia) in nature (see 7.1.4 Geriatrics, Use in Geriatric Patients with Dementia).

4. DOSAGE AND ADMINISTRATION

4.1 Dosing Considerations

- Dosage adjustments are generally not required on the basis of age, gender, race, or renal impairment.
- The absorption of ziprasidone is increased up to two-fold in the presence of a meal.
- Concomitant treatment with other drugs that have been consistently observed to prolong the QT/QTc interval should be avoided. See also: 7 WARNINGS and PRECAUTIONS, Recommendations Regarding Risk Factors for QT Prolongation.

4.2 Recommended Dose and Dosage Adjustment

- **Pediatrics:** Health Canada has not authorized an indication for pediatric use (see [7.1.3 Pediatrics](#))
- **Adults:**
 - **Schizophrenia**

Initial Treatment:

Mylan-Ziprasidone may be administered at an initial daily dose of 40 mg BID with a meal. However, individual patients may benefit from an initial dose of 20 mg BID. Daily dosage may subsequently be adjusted on the basis of clinical status up to 80 mg BID. Dosage adjustments, if indicated, should generally occur at intervals of not less than 2 days, since steady-state is achieved within 1 to 3 days.

Efficacy in schizophrenia was studied in a dose range of 20 to 100 mg BID in short-term, placebo- controlled clinical trials. There were trends toward dose response within the range of 20 to 80 mg BID, but results were not consistent. An increase to a dose greater than 80 mg BID is not generally recommended. The safety of doses above 100 mg BID has not been systematically evaluated in clinical trials.

Maintenance Treatment:

It is recommended that responding patients with schizophrenia be continued on Mylan-Ziprasidone at the lowest dose needed to maintain remission. The efficacy of ziprasidone hydrochloride 20, 40, or 80 mg BID in maintenance treatment has been established over a 12-month treatment period.

Patients should be periodically reassessed to determine the need for maintenance treatment. While there is no body of evidence available to answer the question of how long the patient should be treated with ziprasidone hydrochloride, the effectiveness of maintenance treatment is well established for many other antipsychotic drugs.

- **Geriatrics:** Given the greater sensitivity of this population, a lower starting dose, slower titration, and careful monitoring during the initial dosing period may be considered for elderly patients when clinical factors warrant (see [7.1.4 Geriatrics](#)).

Mylan-Ziprasidone is not indicated for treatment of elderly patients with dementia (see [3 **SERIOUS WARNINGS AND PRECAUTIONS**](#)).

- **Hepatic Impairment:** Lower doses should be considered for hepatic insufficiency, considering that <1% of ziprasidone is cleared renally, and there is a lack of experience with ziprasidone in patients with severe hepatic impairment.

4.4 Administration

Mylan-Ziprasidone should be administered BID, with a meal. Swallow capsules whole, do not open,

crush, or chew the capsules.

4.5 Missed Dose

The missed dose should be taken at the next scheduled dose. Doses should not be doubled.

5. OVERDOSAGE

Symptoms

In premarketing trials, accidental or intentional overdose of ziprasidone was documented in 10 patients. All of these patients survived without sequelae. In the patient taking the largest confirmed amount, 3240 mg, the only symptoms reported were minimal sedation, slurring of speech, and transitory hypertension (200/95).

In post-marketing use, the most common adverse events reported in association with ziprasidone overdose generally included extrapyramidal symptoms, somnolence, tremor, and anxiety. Hypertension, hypotension, diarrhea, tachycardia, and prolongation of the QTc and QRS intervals have also been reported.

Treatment

There is no specific antidote to ziprasidone, and it is not dialyzable. The possibility of multiple drug involvement should be considered.

In case of acute overdose, establish and maintain an airway and ensure adequate oxygenation and ventilation. Intravenous access should be established and gastric lavage (after intubation, if patient is unconscious) and administration of activated charcoal together with a laxative should be considered. The possibility of obtundation, seizure, or dystonic reaction of the head and neck following overdose may create a risk of aspiration with induced emesis.

Cardiovascular monitoring should commence immediately and should include continuous electrocardiographic monitoring to detect possible arrhythmias. If antiarrhythmic therapy is administered, disopyramide, procainamide, and quinidine carry a theoretical hazard of additive QT-prolonging effects that might be additive to those of ziprasidone.

Hypotension and circulatory collapse should be treated with appropriate measures such as intravenous fluids. If sympathomimetic agents are used for vascular support, epinephrine and dopamine should not be used, since beta stimulation combined with α_1 -antagonism associated with ziprasidone may worsen hypotension. Similarly, it is reasonable to expect that the alpha-adrenergic-blocking properties of bretylium might be additive to those of ziprasidone, resulting in problematic hypotension.

In cases of severe extrapyramidal symptoms, anticholinergic medication should be administered.

Close medical supervision and monitoring should continue until the patient recovers.

For management of a suspected drug overdose, contact your regional poison control centre.

6. DOSAGE FORMS, STRENGTHS, COMPOSITION AND PACKAGING

Table – Dosage Forms, Strengths, Composition and Packaging

Route of Administration	Dosage Form / Strength /Composition	Non-medicinal Ingredients
Oral	Capsules / 20 mg, 40 mg, 60 mg and 80 mg Ziprasidone (as Ziprasidone Hydrochloride Monohydrate)	Lactose monohydrate, pollacrillin potassium, povidone, magnesium stearate. Capsule shell: black iron oxide, butyl alcohol, dehydrated alcohol, FD&C blue #1 (20 mg, 40 mg, and 80 mg strengths only), gelatin, isopropyl alcohol, potassium hydroxide, propylene glycol, shellac, strong ammonia solution, titanium dioxide.

Availability

Mylan-Ziprasidone 20 mg, 40 mg, 60 mg, 80 mg capsules are supplied in HDPE bottles of 60s and 100s.

Composition

Mylan-Ziprasidone 20 mg Capsules

Mylan-Ziprasidone 20 mg capsules are hard-shell gelatin capsules filled with white to pink colored powder, and have a No. 4, blue opaque cap and white opaque body. The capsule is axially printed with "Mylan" over "ZH 20" in black ink on the cap and body.

Each capsule contains 20 mg of ziprasidone and the following non-medicinal ingredients: lactose monohydrate, pollacrillin potassium, povidone, magnesium stearate.

Capsule shell: black iron oxide, butyl alcohol, dehydrated alcohol, FD&C blue #1, gelatin, isopropyl alcohol, potassium hydroxide, propylene glycol, shellac, strong ammonia solution, titanium dioxide.

Mylan-Ziprasidone 40 mg Capsules

Mylan-Ziprasidone 40 mg capsules are hard-shell gelatin capsules filled with white to pink colored powder, and have a No. 3, blue opaque cap and blue opaque body. The capsule is axially printed with "Mylan" over "ZH 40" in black ink on the cap and body.

Each capsule contains 40 mg of ziprasidone and the following non-medicinal ingredients: lactose monohydrate, pollacrillin potassium, povidone, magnesium stearate.

Capsule shell: black iron oxide, butyl alcohol, dehydrated alcohol, FD&C blue #1, gelatin, isopropyl alcohol, potassium hydroxide, propylene glycol, shellac, strong ammonia solution, titanium dioxide.

Mylan-Ziprasidone 60 mg Capsules

Mylan-Ziprasidone 60 mg capsules are hard-shell gelatin capsules filled with white to pink colored powder, and have a No. 2, white opaque cap and white opaque body. The capsule is axially printed with "Mylan" over "ZH 60" in black ink on the cap and body.

Each capsule contains 60 mg of ziprasidone and the following non-medicinal ingredients: lactose monohydrate, pollacrillin potassium, povidone, magnesium stearate.

Capsule shell: black iron oxide, butyl alcohol, dehydrated alcohol, gelatin, isopropyl alcohol, potassium hydroxide, propylene glycol, shellac, strong ammonia solution, titanium dioxide.

Mylan-Ziprasidone 80 mg Capsules

Mylan-Ziprasidone 80 mg capsules are hard-shell gelatin capsules filled with white to pink coloured powder, and have a No. 1, blue opaque cap and white opaque body. The capsule is axially printed with "Mylan" over "ZH 80" in black ink on the cap and body.

Each capsule contains 80 mg of ziprasidone and the following non-medicinal ingredients: lactose monohydrate, pollacrillin potassium, povidone, magnesium stearate.

Capsule shell: black iron oxide, butyl alcohol, dehydrated alcohol, FD&C blue #1, gelatin, isopropyl alcohol, potassium hydroxide, propylene glycol, shellac, strong ammonia solution, titanium dioxide.

7. WARNINGS AND PRECAUTIONS

Please see 3 SERIOUS WARNINGS AND PRECAUTIONS.

General

Disruption of the body's ability to reduce core body temperature has been attributed to antipsychotic agents. Appropriate care is advised when prescribing ziprasidone for patients who will be experiencing conditions which may contribute to an elevation in core body temperature, e.g., exercising strenuously, exposure to extreme heat, receiving concomitant medication with anticholinergic activity, or being subject to dehydration.

Ziprasidone has not been evaluated or used to any appreciable extent in patients with a recent history of myocardial infarction or unstable heart disease. Patients with these diagnoses were excluded from pre-marketing clinical studies. Because of the risks of QT/QTc prolongation and orthostatic hypotension with ziprasidone, caution should be observed in cardiac patients (see 2 CONTRAINDICATIONS, as well as 7 WARNINGS AND PRECAUTIONS, QT Prolongation and Orthostatic Hypotension).

Mylan-Ziprasidone capsules contain lactose. This should be considered when prescribing to patients

with rare hereditary problems of galactose intolerance, lactase deficiency or glucose-galactose malabsorption.

Carcinogenesis and Mutagenesis

For animal data, see 16 NON-CLINICAL TOXICOLOGY.

Cardiovascular

- **QT Prolongation** (see also 2 CONTRAINDICATIONS):

Mylan-Ziprasidone is associated with moderate QT/QTc interval prolongation, as described in the subsections below.

Recommendations Regarding Risk Factors for QT Prolongation

Many drugs that cause QT/QTc prolongation are suspected to increase the risk of a rare, potentially fatal polymorphic ventricular tachyarrhythmia known as torsades de pointes. Generally, the risk of torsades de pointes increases with magnitude of the QT/QTc prolongation produced by the drug.

Torsades may be asymptomatic or experienced by the patient as dizziness, palpitations, syncope or seizures. If sustained, torsades de pointes can progress to ventricular fibrillation and sudden cardiac death.

As per Health Canada's QT/QTc Guidelines, in the general population, certain circumstances may increase the risk of the occurrence of torsades de pointes in association with the use of drugs that prolong the QT/QTc interval, including (1) bradycardia; (2) electrolyte disturbances (e.g., hypokalemia, hypomagnesemia, or hypocalcemia); (3) concomitant use of other drugs that prolong the QT/QTc interval; (4) presence of congenital prolongation of the QT interval; (5) family history of sudden cardiac death at <50 years; (6) personal history of cardiac disease or arrhythmias; (7) acute neurological events, e.g., stroke; (8) being female or 65 years of age or older; (9) nutritional deficits e.g., eating disorders; (10) diabetes mellitus. Therefore:

- Ziprasidone should not be used in combination with other drugs that are known to prolong the QT/QTc interval (see 2 CONTRAINDICATIONS and 9.4 Drug-Drug Interactions). Additionally, clinicians should be alert to the identification of other drugs that have been consistently observed to prolong the QT/QTc interval. Such drugs should not be prescribed with ziprasidone.
- Ziprasidone should also not be used in patients with congenital long QT syndrome, or in patients with a history of cardiac arrhythmias, with recent acute myocardial infarction, or with uncompensated heart failure (see 2 CONTRAINDICATIONS).
- If patients with stable cardiac disease are treated, an ECG review should be considered before treatment is started.

- Persistently prolonged QT/QTc intervals may also increase the risk of further prolongation and arrhythmia, but it is not clear that routine screening ECG measures are effective in detecting such patients. Rather, if cardiac symptoms, such as palpitations, vertigo, syncope or seizures occur then the possibility of a malignant cardiac arrhythmia should be considered and a cardiac evaluation including an ECG should be performed. If the QTc interval for a patient is >500 msec, then it is recommended that the treatment be stopped.
- It is recommended that patients being considered for ziprasidone treatment who are at risk for significant electrolyte disturbances (e.g., diuretic therapy, protracted diarrhea or vomiting, water intoxication, eating disorder, and alcoholism), have baseline serum potassium and magnesium measurements performed and levels corrected if necessary. Hypokalemia (and/or hypomagnesemia) may increase the risk of QT prolongation and arrhythmia. It is essential to periodically monitor serum electrolytes in patients for whom diuretic therapy is introduced during ziprasidone treatment.
- Patients receiving treatment with drugs that prolong the QT/QTc interval should be counselled appropriately, regarding risk factors, symptoms suggestive of arrhythmia and risk management strategies ([see 9.4 Drug-Drug Interactions](#)).

Studies Specifically Designed to Assess QT Prolongation

Comparative study (128-054): Six antipsychotics:

A study directly comparing the QT/QTc prolonging effect of ziprasidone with several other drugs effective in the treatment of schizophrenia was conducted in patient volunteers (n=28-35 per drug). In the first phase of the trial, ECGs were obtained at the time of maximum plasma concentration when the drug was administered alone. In the second phase of the trial, ECGs were obtained at the time of maximum plasma concentration while the drug was co-administered with an inhibitor of the CYP450 metabolism of the drug.

In the first phase of the study, the mean change in QTc from baseline was calculated for each drug, using a sample-based correction that makes adjustments for the effect of heart rate on the QT interval. The mean increase in QTc from baseline for ziprasidone (baseline correction) at 160 mg/day was 15.9 msec, which was approximately 9 to 14 msec greater than for four of the comparator drugs (haloperidol at 15 mg/day [7.1 msec], quetiapine at 750 mg/day [5.7 msec], risperidone at 16 mg/day [3.6 msec], and olanzapine at 20 mg /day [1.7 msec], but was approximately 14 msec less than the prolongation observed for thioridazine at 300 mg/day [30.1 msec].

In the second phase of the study, the effect of ziprasidone on QTc length (16.6 msec) was not augmented by the presence of a metabolic inhibitor (ketoconazole 200 mg BID). The mean increase for the other comparator drugs was haloperidol [13.3 msec], quetiapine [8.0 msec], olanzapine [3.0 msec], and risperidone [2.6 msec], compared to thioridazine [29.6 msec].

QT Effects at 2x maximum recommended ziprasidone dose:

A study examining the effect of 3 doses of orally administered ziprasidone (including twice the recommended clinical dose, n=29) and haloperidol (the highest dose level was comparably high, n=30) on the QTc interval was conducted in clinically stable patients with schizophrenia and schizoaffective

disorder. The study comprised 4 consecutive periods, including drug tapering (phase 1), wash out (phase 2), drug therapy (phase 3) followed by the study drug wash out and initiation of outpatient drug therapy (phase 4). Serial baseline electrocardiograms (ECGs) were collected under controlled conditions on the last day (day 0) of period 2 at times matched to those collected during study drug administration (phase 3) at the time of estimated peak drug exposure. At each steady-state dose level, three ECGs and a pharmacokinetic sample were collected at the predicted time of peak exposure to administered drug (T_{max}). One of the three ECGs was collected at T_{max} and the other two were collected one hour on either side of T_{max} .

The mean increase in QTc from baseline for ziprasidone at 40 mg/day was 4.5 msec, and at 160 mg/day was 19.5 msec (comparable to the study described above). A further increase in dose to 320 mg/day (twice the maximum recommended clinical dose) led to an increase in QTc of 22.5 msec, which was only 3 msec more than after 160 mg/day in this study, suggesting a plateau. In comparison, there was no mean QTc increase apparent at the lowest haloperidol dose (2.5 mg/day). At the 2 higher doses of haloperidol, (15 and 30 mg/day), mean QTc increases ranged from 6.6 to 7.2 msec. No subject in either treatment group experienced a QTc interval ≥ 450 msec or an increase from baseline of ≥ 75 msec.

Data from Non-QT specific ziprasidone studies

In placebo-controlled trials, ziprasidone increased the QTc interval compared to placebo by approximately 10 msec at the highest recommended daily dose of 160 mg, which was the basis for subsequent QT-specific studies. The clinical trial data for ziprasidone did not reveal an excess risk of mortality for ziprasidone compared to other antipsychotic drugs or placebo.

Electrocardiogram readings revealing QTc intervals exceeding the potentially clinically relevant threshold of 500 msec in clinical trials with ziprasidone occurred in: 2/3266 (0.06%) patients receiving ziprasidone hydrochloride and 1/538 (0.19%) patients receiving placebo. One patient had a history of prolonged QTc and a screening measurement of 489 msec; QTc was 503 msec during ziprasidone treatment. The other patient, who was receiving ziprasidone for more than 6.5 years without interruption, had a QTc of 503 msec at week 189, and 435 msec 19 weeks later, while maintained on the same oral dose of ziprasidone. There were confounding factors that contributed to the occurrence of these cases.

Post-Marketing Data (see also 8.5 Post-Market Adverse Reactions)

There have been rare post-marketing reports of torsades de pointes (in the presence of multiple confounding factors) (see 8.5 Post-Market Adverse Reactions). Torsades de pointes have not been observed in association with the use of ziprasidone at recommended doses in clinical trials, but experience is too limited to rule out increased risk.

In view of the clinical trial data demonstrating a moderate QT prolongation effect of ziprasidone hydrochloride, a review of 5-year, post-marketing spontaneous data from the FDA AERS database was conducted using a set of heart-related search terms.

Small elevations in spontaneous reporting rates were observed for ziprasidone compared with two other atypical antipsychotics, for both fatal cases, and "all" cases (i.e., fatal plus non-fatal).

Accumulated case reports should not be used as a basis for determining the incidence of a reaction or estimating risk for a particular product, as neither the total number of reactions occurring, nor the number of patients exposed to the health product is known. Because of the multiple factors that influence reporting, quantitative comparisons of health product safety cannot be made from the data. Comparison of reporting rates cannot be employed to confirm or refute a hypothesis, due to well-known, inherent limitations with spontaneous reporting of adverse events.

- **Orthostatic Hypotension:**

Ziprasidone may induce orthostatic hypotension associated with dizziness, tachycardia and, in some patients, syncope, especially during the initial dose-titration period, probably reflecting its α_1 -adrenergic antagonist properties. Syncope was reported in 0.6% (22/3834) of the patients treated with ziprasidone.

Ziprasidone should be used with particular caution in patients with known cardiovascular disease (history of myocardial infarction or ischemic heart disease, heart failure or conduction abnormalities), cerebrovascular disease or conditions which would predispose patients to hypotension (dehydration, hypovolemia and treatment with antihypertensive medications). Patients with a history of clinically significant cardiac disorders were excluded from the trials.

Dependence/Tolerance

Ziprasidone hydrochloride has not been systematically studied, in animals or humans, for its potential for abuse, tolerance or physical dependence. While clinical trials did not reveal any tendency for drug-seeking behaviour, these observations were not systematic and it is not possible to predict on the basis of this limited experience the extent to which ziprasidone hydrochloride will be misused, diverted and/or abused once marketed. Consequently, patients should be evaluated carefully for a history of drug abuse, and such patients should be observed closely for signs of Mylan-Ziprasidone misuse or abuse (e.g., development of tolerance, increases in dose, drug-seeking behaviour).

Driving and Operating Machinery

Since ziprasidone has the potential to impair judgment, thinking, or motor skills, patients should be cautioned about performing activities requiring mental alertness, such as operating a motor vehicle (including automobiles) or operating hazardous machinery, until they are reasonably certain that ziprasidone therapy does not affect them adversely.

Endocrine and Metabolism

- **Hyperglycemia:**

As with some other antipsychotics, hyperglycemia, exacerbation of pre-existing diabetes, and diabetic coma have been reported very rarely during the use of ziprasidone hydrochloride.

However, no causal relationship with ziprasidone hydrochloride has been established (see **8.5 Post-Market Adverse Reactions**). Diabetic ketoacidosis (DKA) has occurred in patients with no reported history of hyperglycemia. Patients should have baseline and periodic monitoring of blood glucose and body weight.

Assessment of the relationship between atypical antipsychotic use and glucose abnormalities is complicated by the possibility of an increased background risk of diabetes mellitus in patients with schizophrenia and the increasing incidence of diabetes mellitus in the general population. Given these confounders, and that there is no data in drug-naïve patients, the relationship between atypical antipsychotic use and hyperglycemia-related adverse events is not completely understood. However, epidemiological studies, which did not include ziprasidone hydrochloride, suggest an increased risk of treatment-emergent hyperglycemia-related adverse events in patients treated with the atypical antipsychotics. Because ziprasidone hydrochloride was not marketed at the time these studies were performed, it is not known if ziprasidone hydrochloride is associated with this increased risk. Precise risk estimates for hyperglycemia-related adverse events in patients treated with atypical antipsychotics are not available.

Any patient treated with atypical antipsychotics should be monitored for symptoms of hyperglycemia, including polydipsia, polyuria, polyphagia, and weakness. Patients who develop symptoms of hyperglycemia during treatment with atypical antipsychotics should undergo fasting blood glucose testing. In some cases, hyperglycemia has resolved when the atypical antipsychotic was discontinued; however, some patients required continuation of anti-diabetic treatment despite discontinuation of the suspect drug. Patients with risk factors for diabetes mellitus (e.g., obesity, family history of diabetes) who are starting treatment with atypical antipsychotics should undergo fasting blood glucose testing at the beginning of treatment and periodically during treatment. Patients with an established diagnosis of diabetes mellitus who are started on atypical antipsychotics should be monitored regularly for worsening of glucose control.

- **Hyperprolactinemia:**

As with other drugs that antagonize dopamine D₂ receptors and/or serotonin 5-HT₂ receptors, Mylan-Ziprasidone may elevate prolactin levels in humans. Elevations associated with Mylan-Ziprasidone treatment are generally mild and may decline during administration.

Increased prolactin levels were also observed in animal studies with this compound, and were associated with an increase in mammary gland neoplasia in mice; a similar effect was not observed in rats (see **16 NON-CLINICAL TOXICOLOGY, Carcinogenicity**).

Tissue culture experiments indicate that approximately one-third of human breast cancers are prolactin-dependent in vitro, Mylan-Ziprasidone should only be administered to patients with previously detected breast cancer if the benefits outweigh the potential risks.

Although disturbances such as galactorrhea, amenorrhea, gynecomastia, and impotence have been reported with prolactin-elevating compounds, the clinical significance of elevated serum prolactin levels is unknown for most patients. Long-standing hyperprolactinemia when associated with

hypogonadism may lead to decreased bone mineral density in both female and male subjects. Caution should be exercised when considering ziprasidone treatment in patients with pituitary tumors. Neither clinical studies nor epidemiological studies conducted to date have shown an association between chronic administration of this class of drugs and tumorigenesis in humans. The available evidence is considered too limited to be conclusive at this time.

Gastrointestinal

Patients should be advised of the risk of severe constipation during Mylan-Ziprasidone treatment, and that they should tell their doctor if constipation occurs or worsens, as they may need laxatives (see [8.5 Post Market Adverse Reactions](#)).

Genitourinary

- **Priapism:**

Rare cases of priapism have been reported with antipsychotic use, such as ziprasidone hydrochloride. This adverse reaction, as with other psychotropic drugs, did not appear to be dose-dependent and did not correlate with duration of treatment. The likely mechanism of action of priapism is a relative decrease in sympathetic tone. Severe priapism may require surgical intervention.

Hematologic

- **Venous Thromboembolism:**

Venous thromboembolism (VTE), including fatal pulmonary embolism, has been reported with antipsychotic drugs, including ziprasidone hydrochloride, in case reports and/or observational studies. When prescribing Mylan-Ziprasidone, all possible risk factors for VTE should be identified before and during treatment with Mylan-Ziprasidone and preventive measures undertaken.

- **Neutropenia, granulocytopenia, and agranulocytosis:**

Neutropenia, granulocytopenia, and agranulocytosis have been reported during antipsychotic use (see [8.5 Post Market Adverse Reactions](#)). Therefore, it is recommended that patients have their complete blood count (CBC) tested prior to starting Mylan-Ziprasidone and then periodically throughout treatment.

Possible risk factors for leukopenia/neutropenia include pre-existing low white blood cell count (WBC) and history of drug induced leukopenia/neutropenia. Patients with a pre-existing low WBC or a history of drug induced leukopenia/neutropenia should have their complete blood count (CBC) monitored frequently during the first few months of therapy and Mylan-Ziprasidone should be discontinued at the first sign of decline in WBC, in the absence of other causative factors.

Patients with neutropenia should be carefully monitored for fever or other symptoms or signs of infection and treated promptly if such symptoms or signs occur. Patients with severe neutropenia (absolute neutrophil count $<1 \times 10^9/L$) should discontinue Mylan-Ziprasidone and have their WBC

followed until recovery.

Hepatic/Biliary/Pancreatic

In patients with hepatic insufficiency, lower doses should be considered (see [4.2 Recommended Dose and Dosage Adjustment, Hepatic Impairment](#) and [10.3 Pharmacokinetics, Hepatic Insufficiency](#)).

Monitoring and Laboratory Tests

Patients being considered for ziprasidone treatment that are at risk of significant electrolyte disturbances should have baseline serum potassium and magnesium measurements. Low serum potassium and magnesium should be repleted before proceeding with treatment. Patients who are started on diuretics during ziprasidone therapy need periodic monitoring of serum potassium and magnesium. Ziprasidone should be discontinued in patients who are found to have persistent QTc measurements >500 msec (see [7 WARNINGS AND PRECAUTIONS, QT Prolongation](#)).

Neurologic

- **Neuroleptic Malignant Syndrome (NMS) :**

A potentially fatal symptom complex sometimes referred to as NMS has been reported in association with the administration of antipsychotic drugs, including ziprasidone hydrochloride.

Clinical manifestations of NMS are hyperpyrexia, muscle rigidity, altered mental status, and evidence of autonomic instability (irregular pulse or blood pressure, tachycardia, diaphoresis, and cardiac dysrhythmia). Additional signs may include elevated creatinine phosphokinase, myoglobinuria (rhabdomyolysis), and acute renal failure.

In arriving at a diagnosis, it is important to identify cases where the clinical presentation includes both serious medical illness (e.g., pneumonia, systemic infection, etc.), and untreated or inadequately treated extrapyramidal signs and symptoms (EPS). Other important considerations in the differential diagnosis include central anticholinergic toxicity, heat stroke, drug fever, and primary central nervous system pathology.

The management of NMS should include: 1) immediate discontinuation of antipsychotic drugs including Mylan-Ziprasidone and other drugs not essential to concurrent therapy; 2) intensive symptomatic treatment and medical monitoring; and 3) treatment of any concomitant serious medical problems for which specific treatments are available. There is no general agreement about specific pharmacological treatment regimens for uncomplicated NMS.

If a patient requires antipsychotic drug treatment after recovery from NMS, the potential reintroduction of drug therapy should be carefully considered. The patient should be carefully monitored, since recurrences of NMS have been reported.

- **Tardive Dyskinesia (TD):**

A syndrome consisting of potentially irreversible, involuntary and disabling dyskinesic movements may develop in patients treated with antipsychotic drugs. Although the prevalence of the tardive dyskinesia syndrome appears to be highest among the elderly, especially elderly women, it is impossible to rely upon prevalence estimates to predict, at the inception of antipsychotic treatment, which patients are likely to develop the tardive dyskinesia syndrome. Whether antipsychotic drug products differ in their potential to cause tardive dyskinesia is unknown.

The risk of developing tardive dyskinesia, and the likelihood that it will become irreversible are believed to increase as the duration of treatment and the total cumulative dose of antipsychotic drugs administered to the patient increase. However, the syndrome can develop, although much less commonly, after relatively brief treatment periods at low doses.

There is no known treatment for established cases of tardive dyskinesia, although the syndrome may remit, partially or completely, if antipsychotic treatment is withdrawn. Antipsychotic treatment itself however may suppress (or partially suppress) the signs and symptoms of the tardive dyskinesia syndrome and thereby may possibly mask the underlying process. The effect that symptomatic suppression has upon the long-term course of the tardive dyskinesia syndrome is unknown.

Given these considerations, Mylan-Ziprasidone should be prescribed in a manner that is most likely to minimize the occurrence of tardive dyskinesia. Chronic antipsychotic treatment should generally be reserved for patients who 1) suffer from a chronic illness that is known to respond to antipsychotic drugs, and 2) for whom alternative, equally effective but potentially less harmful treatments are not available or appropriate. In patients who do require chronic treatment, the smallest dose and the shortest duration of treatment producing a satisfactory clinical response should be sought. The need for continued treatment should be reassessed periodically.

If signs and symptoms of tardive dyskinesia appear in a patient on Mylan-Ziprasidone, drug discontinuation should be considered. However, some patients may require treatment with Mylan-Ziprasidone despite the presence of the tardive dyskinesia syndrome.

- **Falls:**

Antipsychotic drugs (which includes ziprasidone) may cause somnolence, postural hypotension, and motor and sensory instability, which could lead to falls and, consequently, fractures or other injuries. For patients with diseases, conditions, or medications that could exacerbate these effects, a fall risk assessment should be completed when initiating antipsychotic treatment and recurrently for patients on long-term antipsychotic therapy.

- **Somnolence:**

Somnolence was a commonly reported adverse event in patients treated with ziprasidone hydrochloride. In the 4- and 6- week placebo-controlled trials in patients with schizophrenia, somnolence was reported in 14% of patients on ziprasidone compared to 7% of patients on placebo (see **8 ADVERSE REACTIONS**).

- **Seizures:**

During clinical trials, seizures occurred in 0.4% of patients treated with ziprasidone. There were confounding factors that may have contributed to the occurrence of seizures in many of these cases. Nevertheless, as with other antipsychotic drugs, ziprasidone should be used cautiously in patients with a history of seizures or with conditions that potentially lower the seizure threshold, e.g., Alzheimer's dementia. Conditions that lower the seizure threshold may be more prevalent in a population of 65 years or older.

Serotonin toxicity / Serotonin syndrome:

Serotonin toxicity, also known as serotonin syndrome, is a potentially life-threatening condition in patients taking multiple serotonergic agents or who have had considerable exposure to a single serotonin-augmenting drug. In isolated cases, there have been reports of serotonin syndrome temporally associated with the therapeutic use of ziprasidone in combination with other serotonergic medicinal products such as SSRIs (see **8.5 Post-Market Adverse Reactions**).

Serotonin toxicity is characterized by neuromuscular excitation, autonomic stimulation (e.g. tachycardia, flushing) and altered mental state (e.g. anxiety, agitation, hypomania). In accordance with the Hunter Criteria, serotonin toxicity diagnosis is likely when, in the presence of at least one serotonergic agent, one of the following is observed:

- spontaneous clonus
- inducible clonus or ocular clonus with agitation or diaphoresis
- tremor and hyperreflexia
- hypertonia and body temperature >38°C and ocular clonus or inducible clonus.

If concomitant treatment with Mylan-Ziprasidone and serotonergic agents is clinically warranted, careful observation of the patient is advised. If serotonin toxicity is suspected, discontinuation of the serotonergic agents should be considered.

Psychiatric

- **Suicide:**

The possibility of a suicide attempt is inherent in psychotic illness; thus, close supervision and appropriate clinical management of high-risk patients should accompany drug therapy.

Prescriptions for Mylan-Ziprasidone should be written for the smallest quantity of capsules consistent with good patient management, in order to reduce the risk of overdose.

Renal

Dose adjustments are not required for patients with renal impairment (see **10.3 Pharmacokinetics, Renal Insufficiency**).

Reproductive Health: Female and Male Potential

There are no adequate and well-controlled studies in women and men exposed to ziprasidone.

Skin

- **Severe Cutaneous Adverse Reactions:**

Severe cutaneous adverse reactions (SCARs), including Stevens-Johnson syndrome (SJS), toxic epidermal necrolysis (TEN) and drug reaction with eosinophilia and systemic symptoms (DRESS) are potentially life threatening adverse drug reactions that have been reported with atypical antipsychotic exposure. SCARs commonly present as a combination of the following symptoms: malaise, mucosal ulceration, extensive cutaneous rash or exfoliative dermatitis, fever, lymphadenopathy and possible eosinophilia. Discontinue Mylan-Ziprasidone if severe cutaneous adverse reactions occur (see [8.5 Post-Market Adverse Reactions](#)).

Drug reaction with eosinophilia and systemic symptoms (DRESS) has been reported with ziprasidone exposure. DRESS consists of a combination of three or more of the following: cutaneous reaction (such as rash or exfoliative dermatitis), eosinophilia, fever, lymphadenopathy and one or more systemic complications such as hepatitis, nephritis, pneumonitis, myocarditis, and pericarditis.

Other severe cutaneous adverse reactions, such as Stevens-Johnson syndrome, have been reported with ziprasidone exposure (see [8.5 Post Market Adverse Reactions](#)).

Severe cutaneous adverse reactions are sometimes fatal. Discontinue ziprasidone if severe cutaneous adverse reactions occur.

- **Rash:**

In pre-marketing trials with ziprasidone, about 5% of patients developed rash (173/3834) and/or urticaria (12/3834), with discontinuation in about one-sixth of these cases. The occurrence of rash was related to dose of ziprasidone, although the finding might also be explained by the longer exposure time in the higher dose patients. Several patients with rash had signs and symptoms of associated systemic illness, e.g., elevated white blood cells (WBC). Most patients improved promptly with adjunctive treatment with antihistamines or steroids and/or upon discontinuation of ziprasidone, and all patients experiencing these events were reported to recover completely. Upon appearance of rash for which an alternative etiology cannot be identified, ziprasidone should be discontinued.

7.1 Special Populations

7.1.1 Pregnant Women

There are no adequate and well-controlled studies in pregnant women. Women of childbearing potential receiving ziprasidone should therefore be counselled on the need to use an effective method of contraception during treatment with Mylan-Ziprasidone.

Patients should be advised to notify their health professional if they become pregnant or intend to become pregnant. Mylan-Ziprasidone should not be used during pregnancy unless the expected benefits to the mother markedly outweigh the potential risks to the fetus.

Teratogenic effects:

In animal studies, ziprasidone demonstrated developmental toxicity, including possible teratogenic effects at doses similar to human therapeutic doses. When ziprasidone was administered to pregnant rabbits during the period of organogenesis, an increased incidence of fetal structural abnormalities (ventricular septal defects and other cardiovascular malformations and kidney alterations) was observed at a dose of 30 mg/kg/day (3 times the maximum recommended human dose (MRHD) of 200 mg/day on a mg/m² basis). There was no evidence to suggest that these developmental effects were secondary to maternal toxicity. The developmental no-effect dose was 10 mg/kg/day (equivalent to the MRHD on an mg/m² basis).

In rats, embryofetal toxicity (decreased fetal weights, delayed skeletal ossification) was observed following administration of 10 to 160 mg/kg/day (0.5 to 8 times the MRHD on a mg/m² basis) during organogenesis or throughout gestation, but there was no evidence of teratogenicity. Doses of 40 and 160 mg/kg/day (2 and 8 times the MRHD on a mg/m² basis) were associated with maternal toxicity. The developmental no-effect dose was 5 mg/kg/day (0.2 times the MRHD on a mg/m² basis).

There was an increase in the number of pups born dead and a decrease in postnatal survival through the first 4 days of lactation among the offspring of female rats treated during gestation and lactation with doses of 10 mg/kg/day (0.5 times the MRHD on a mg/m² basis) or greater. Offspring developmental delays and neurobehavioral functional impairment were observed at doses of 5 mg/kg/day (0.2 times the MRHD on a mg/m² basis) or greater. A no-effect level was not established for these effects.

Non teratogenic effects:

Neonates exposed to antipsychotic drugs (including Mylan-Ziprasidone) during the third trimester of pregnancy are at risk for extrapyramidal and/or withdrawal symptoms following delivery. There have been reports of agitation, hypertonia, hypotonia, tremor, somnolence, respiratory distress and feeding disorder in these neonates. These complications have varied in severity; while in some cases symptoms have been self-limited, in other cases neonates have required intensive care unit support and prolonged hospitalization.

Mylan-Ziprasidone should not be used during pregnancy unless the expected benefits to the mother markedly outweigh the potential risks to the fetus.

Labor and Delivery

The effect of ziprasidone hydrochloride on labor and delivery in humans is not known.

7.1.2 Breast-feeding

There are no adequate and well-controlled studies in lactating women. Limited data indicate that ziprasidone is excreted into breast milk at very low levels. It is recommended that women taking Mylan-Ziprasidone should not breast-feed.

7.1.3 Pediatrics

Pediatrics (< 18 years of age): Based on the data submitted and reviewed by Health Canada, the safety and efficacy of ziprasidone hydrochloride in pediatric patients under the age of 18 years has not been established; therefore, Health Canada has not authorized an indication for pediatric use and its use is not recommended.

Weight gain has been observed with atypical antipsychotic use in pediatric and adolescent patient populations. Independent of any drug-specific effects, weight gain can be associated with adverse changes in other metabolic parameters (e.g., glucose and lipid metabolism). Abnormal childhood weight and metabolic status can have adverse effects on cardiovascular outcomes in adulthood. Weight gain and adverse effects on other metabolic parameters associated with atypical antipsychotics can be more frequent or more severe in pediatric and adolescent patients than in the adult patients.

The following adverse events in the two studies in pediatrics are of note because they are not typical of the adult population treated with ziprasidone: abnormally decreased bicarbonate values; elevations in testosterone, elevations in insulin, total neutrophils, monocytes and ALT; fatigue; abdominal pain, insomnia; restlessness. There are also adverse events of note due to a greater incidence rate compared to adults, or a greater differential over placebo: blurred vision; extrapyramidal symptoms (aggregated); sedation/somnolence; nausea; vomiting; and elevations in serum prolactin (See [8.2.1 Clinical Trial Adverse Reactions - Pediatrics](#)).

The long-term safety, including cardiometabolic effects and effects on growth, maturation and behavioural development in patients under 18 years of age has not been systematically evaluated.

The ziprasidone hydrochloride pediatric study in schizophrenia was terminated due to lack of efficacy (Placebo-controlled study A1281134, and open-label extension 1135)

7.1.4 Geriatrics

The number of patients 65 years or older with schizophrenia or related disorders, exposed to ziprasidone hydrochloride during clinical trials was limited (n=109). In general, there was no indication of any different tolerability of ziprasidone or for reduced clearance of ziprasidone in the elderly compared to younger adults. Nevertheless, geriatric patients generally have decreased cardiac, hepatic

and renal function, and more frequent use of concomitant medication. The presence of multiple factors that might increase the pharmacodynamic response to ziprasidone, or cause poorer tolerance or orthostasis, should lead to consideration of a lower starting dose, slower titration, and careful monitoring during the initial dosing period for elderly patients.

Use in Geriatric Patients with Dementia

- **Overall Mortality**

Elderly patients with dementia treated with atypical antipsychotic drugs have an increased mortality compared to placebo in a meta-analysis of 13 controlled trials of various atypical antipsychotic drugs. Mylan-Ziprasidone is not indicated for the treatment of elderly patients with dementia (e.g., dementia-related psychosis) (see 3 SERIOUS WARNINGS AND PRECAUTIONS).

- **Dysphagia**

Esophageal dysmotility and aspiration have been associated with antipsychotic drug use. Aspiration pneumonia is a common cause of morbidity and mortality in elderly patients, in particular those with advanced Alzheimer's dementia. Ziprasidone and other antipsychotic drugs should be used cautiously in patients at risk for aspiration pneumonia.

- **Cerebrovascular Adverse Events (CVAEs), including Stroke in Elderly Patients with Dementia**

In placebo-controlled trials with some atypical antipsychotics, there was a higher incidence of cerebrovascular adverse events (cerebrovascular accidents and transient ischemic attacks) including fatalities compared to placebo-treated subjects. Mylan-Ziprasidone is not indicated for the treatment of patients with dementia (e.g., dementia-related psychosis). (see 3 SERIOUS WARNINGS AND PRECAUTIONS).

- **Lower Threshold for Seizure**

Conditions that lower the seizure threshold, such as Alzheimer's dementia, may be more prevalent in a population of 65 years or older. Ziprasidone and other antipsychotic drugs should be used cautiously in patients with a history of seizures or with conditions that potentially lower the seizure threshold.

8. ADVERSE REACTIONS

8.1 Adverse Reaction Overview

In the schizophrenia studies, the most commonly observed adverse events associated with the use of ziprasidone (incidence of 5% or greater) and observed at a rate on ziprasidone at least twice that of placebo were somnolence (14%), extrapyramidal symptoms (14%), and respiratory tract infection (8%).

An analysis for dose response in the schizophrenia 4-study pool revealed an apparent relation of adverse event to dose for the following events: asthenia, postural hypotension, anorexia, dry mouth, increased salivation, arthralgia, anxiety, dizziness, dystonia, hypertonia, somnolence, tremor, rhinitis, rash, and abnormal vision.

Explorations for interactions on the basis of gender did not reveal any clinically meaningful differences in the adverse event occurrence on the basis of this demographic factor.

8.2 Clinical Trial Adverse Reactions

Clinical trials are conducted under very specific conditions. The adverse reaction rates observed in the clinical trials; therefore, may not reflect the rates observed in practice and should not be compared to the rates in the clinical trials of another drug. Adverse reaction information from clinical trials may be useful in identifying and approximating rates of adverse drug reactions in real-world use.

The following findings are based on a pool of two 6-week, and two 4-week placebo-controlled trials for schizophrenia in which ziprasidone was administered in doses ranging from 10 to 200 mg/day.

Schizophrenia

A total of 4.1% (29/702) of patients treated with ziprasidone hydrochloride in short-term, placebo-controlled studies discontinued treatment due to an adverse event, compared with 2.2% (6/273) on placebo and 8.2% (7/85) on the active control drug. The most common event associated with dropout was rash, including 7 dropouts for rash among ziprasidone patients (1%) compared to no placebo patients (see **7 WARNINGS AND PRECAUTIONS, Rash**).

Table 1 enumerates the incidence, rounded to the nearest percent, of treatment-emergent adverse events that occurred during acute therapy (up to 6 weeks) in predominantly schizophrenic patients, including only those events that occurred in 1% or more of patients treated with ziprasidone and for which the incidence in patients treated with ziprasidone was greater than the incidence in placebo-treated patients.

Table 1. Treatment-Emergent Adverse Event Incidence in Short-Term Placebo-Controlled Trials – Schizophrenia

Body System	Percentage of patients reporting	
	Ziprasidone Hydrochloride (n = 702)	Placebo (n = 273)
Body as a Whole		
Asthenia	5	3
Accidental Injury	4	2
Chest Pain	3	2
Cardiovascular		
Tachycardia	2	1

Body System	Percentage of patients reporting	
	Ziprasidone Hydrochloride (n = 702)	Placebo (n = 273)
Postural Hypotension	1	0
Digestive		
Nausea	10	7
Constipation	9	8
Dyspepsia	8	7
Diarrhea	5	4
Dry Mouth	4	2
Anorexia	2	1
Musculoskeletal		
Myalgia	1	0
Nervous		
Extrapyramidal Symptoms*	14	8
Somnolence	14	7
Akathisia	8	7
Dizziness**	8	6
Respiratory		
Respiratory Tract Infection	8	3
Rhinitis	4	2
Cough Increased	3	1
Skin and Appendages		
Rash	4	3
Fungal Dermatitis	2	1
Special Senses		
Abnormal Vision	3	2

* Extrapyramidal Symptoms includes the following adverse event terms: extrapyramidal syndrome, hypertonia, dystonia, dyskinesia, hypokinesia, tremor, paralysis and twitching. None of these adverse events occurred individually at an incidence greater than 5% in schizophrenia trials.

** Dizziness includes the adverse event terms dizziness and lightheadedness

Explorations for interactions on the basis of gender did not reveal any clinically meaningful differences in the adverse event occurrence on the basis of this demographic factor.

In the schizophrenia studies, the most commonly observed adverse events associated with the use of ziprasidone (incidence of 5% or greater) and observed at a rate on ziprasidone at least twice that of placebo were somnolence (14%), extrapyramidal symptoms (14%), and respiratory tract infection (8%).

An analysis for dose response in the schizophrenia 4-study pool revealed an apparent relation of adverse event to dose for the following events: asthenia, postural hypotension, anorexia, dry mouth, increased salivation, arthralgia, anxiety, dizziness, dystonia, hypertonia, somnolence, tremor, rhinitis, rash, and abnormal vision.

Extrapyramidal Symptoms (EPS): The incidence of reported EPS for ziprasidone-treated patients in the short-term, placebo-controlled trials was 14% vs. 8% for placebo. Medications to treat EPS and

movement disorders were allowed; but if clinically meaningful movement disorder side effects were observed by the clinician or volunteered by the subject, or if a clinically meaningful movement disorder, present at screening, increased in severity or required the administration of anticholinergics or propranolol, these symptoms and their severity were recorded as adverse event. Objectively collected data from those trials on the Simpson Angus Rating Scale (for EPS) and the Barnes Akathisia Scale (for akathisia) did not generally show a difference between ziprasidone and placebo.

Table 2. Treatment-Emergent Extrapyramidal Symptoms Assessed by Adverse Events Incidence in Short-Term, Schizophrenia Placebo-Controlled Trials

Extrapyramidal Symptoms	Percentage of Subjects Reporting Event	
	Ziprasidone Hydrochloride N=702	Placebo N=273
Dystonic events ¹	4.0%	2.2%
Parkinsonism events ²	10.7%	5.1%
Akathisia events ³	8.4%	7.0%
Dyskinetic events ⁴	1.9%	2.9%
Residual events ⁵	0.3%	0.4%
Any extrapyramidal event	21.7%	15%

¹ Patients with the following COSTART terms were counted in this category: dystonia, oculogyric crisis

² Patients with the following COSTART terms were counted in this category: abnormal gait, akinesia, cogwheel rigidity, extrapyramidal syndrome, hypokinesia, muscular hypertonia, tremor

³ Patients with the following COSTART terms were counted in this category: akathisia

⁴ Patients with the following COSTART terms were counted in this category: dyskinesia, paralysis, tardive dyskinesia

⁵ Patients with the following COSTART terms were counted in this category: twitching

In short-term, placebo-controlled trials for patients with a psychotic disorder other than schizophrenia-related, the incidence of treatment-emergent EPS assessed by adverse events was greater in ziprasidone-treated patients than placebo (40% vs 16%, respectively). As with schizophrenic patient population, objectively collected data from those trials on the Simpson Angus Rating Scale (for EPS) and the Barnes Akathisia Scale (for akathisia) did not generally show a difference between ziprasidone and placebo.

Dystonia: Symptoms of dystonia, prolonged abnormal contractions of muscle groups, may occur in susceptible individuals during the first few days of treatment. Dystonic symptoms include: spasm of the neck muscles, sometimes progressing to tightness of the throat, swallowing difficulty, difficulty breathing, and/or protrusion of the tongue. While these symptoms can occur at low doses, they occur more frequently and with greater severity with high potency and at higher doses of first generation antipsychotic drugs. An elevated risk of acute dystonia is observed in males and younger age groups.

Vital Sign Changes: Mylan-Ziprasidone is associated with orthostatic hypotension (see **7 WARNINGS AND PRECAUTIONS, Orthostatic Hypotension**).

ECG Changes: Ziprasidone is associated with an increase in the QTc interval (see **7 WARNINGS AND**

PRECAUTIONS, QT Prolongation). In schizophrenia trials, ziprasidone was associated with a mean increase in heart rate of 1.4 beats per minute compared to a 0.2 beats per minute decrease among placebo patients.

Weight Gain: The proportions of patients meeting a weight gain criterion of $\geq 7\%$ of body weight were compared in a pool of four 4- and 6-week placebo-controlled schizophrenia clinical trials, revealing a statistically significantly greater incidence of weight gain for ziprasidone (10%) compared to placebo (4%). A median weight gain of 0.5 kg was observed in ziprasidone patients compared to no median weight change in placebo patients. In this set of clinical trials, weight gain was reported as an adverse event in 0.4% of ziprasidone and 0.4% of placebo patients.

During long-term therapy with ziprasidone, a categorization of patients at baseline on the basis of body mass index (BMI) revealed the greatest mean weight gain and highest incidence of clinically significant weight gain ($>7\%$ of body weight) in patients with low BMI (<23) compared to normal (23-27) or overweight patients (>27). There was a mean weight gain of 1.4 kg for those patients with a “low” baseline BMI, no mean change for patients with a “normal” BMI, and a 1.3 kg mean weight loss for patients who entered the program with a “high” BMI.

8.2.1 Clinical Trial Adverse Reactions – Pediatrics

The safety and efficacy of ziprasidone hydrochloride in children under the age of 18 years have not been established and its use is not recommended (see [7.1.3 Pediatrics](#)).

All of the adverse events described above for adults should be considered in the case of children and adolescents taking Mylan-Ziprasidone. In placebo-controlled clinical trials, the most frequent adverse reactions (reported with a frequency $> 10\%$) were sedation, somnolence, headache, fatigue, dizziness, nausea, vomiting, and decreased appetite and extrapyramidal disorders. The pediatric safety profile of ziprasidone was similar to the adult profile, except for an increased incidence of sedation and somnolence in pediatric patients.

Additional adverse events of note from the study in the pediatric population, are summarized in the following table. The listed events are those that are either i) worse in children than in adults (greater frequency rates compared to studies in adults of the same disorder, or greater difference from placebo rates, or greater severity), or ii) identified only in pediatric populations, and for which drug rates are greater than placebo.

Table 3 Adverse Events during short-term (6 weeks) treatment of children and adolescents with Schizophrenia (ages 13 -17 years), that were identified as worse in children compared to adults or unique to children, and greater than placebo.

Body System and MedDRA term	Percentage of Subjects With Adverse events	
	Ziprasidone Hydrochloride (n=193)	Placebo (n=90)

Gastrointestinal Disorder		
Nausea	10%	2%
Vomiting	6%	3%
General Disorders		
Fatigue	9%	4%
Investigations		
Prolactin increased ^a	20% 24% (male) 14% (female)	5% 8% (male) 0% (female)
Bicarbonate decreased ^b	17%	6%
Monocytes increased ^c	7%	2%
Nervous system disorders		
Extrapyramidal Symptoms ^d	25%	7%
Sedation, Somnolence	24%	7%
Dizziness	9%	1%

Includes data up to 6 days after last dose of study drug.
Subjects were counted only once per treatment in each row.
Medical Dictionary for Regulatory Activities (v14.0) coding applied.

- ^a Prolactin levels: >19.47 µg/L males (>1.1 ULN=17.7 µg/L); >32.12 µg/L females (>1.1 ULN=29.2 µg/L) at any time. One subject had an increased prolactin level greater than 60 µg/L (female, on ziprasidone, 98.3 µg/L). For the term prolactin increased, “n” for ziprasidone hydrochloride = 50 and for placebo = 20
- ^b Bicarbonate levels: < 19.8 mEq/L (<0.9 LLN= 22 per µL) at any time. Lowest observed ziprasidone hydrochloride value was 15 mEq/L. For the term bicarbonate decreased, “n” for ziprasidone hydrochloride = 95 and for placebo = 49
- ^c Monocyte levels: > 11% (>1.2 ULN=9-10%) at any time. Highest observed ziprasidone hydrochloride value was 17% . For the term monocytes increased, “n” for ziprasidone hydrochloride = 172 and for placebo = 84
- ^d “Extrapyramidal symptoms” includes the following adverse reaction terms: gait disturbance, muscle rigidity, muscle spasms, muscle twitching, torticollis, musculoskeletal stiffness, akathisia, drooling, dyskinesia, dystonia, extrapyramidal disorder, hypokinesia, parkinsonian rest tremor, tremor, masked facies.

Weight Gain in Children and Adolescents: In one 6-week, placebo-controlled trial in adolescent patients (13-17 years of age) with schizophrenia, the mean change in body weight was -0.1 kg in the ziprasidone group and 0.0 kg in the placebo group. Of ziprasidone hydrochloride treated patients 4 % gained > 7% of their bodyweight, compared to 0% of placebo-treated patients.

In the open-label study that enrolled patients from the above schizophrenia trial, 34% of patients (76/221) completed 26 weeks of therapy with ziprasidone, and 13 % of the patients gained ≥ 7% of their body weight, not adjusted for normal growth. After 26 weeks of treatment, the mean increase in body weight was 1.7 kg. In order to adjust for normal growth over 26 weeks an increase of at least 0.5 standard deviation from baseline in BMI Z-score was used as a measure of a clinically significant change; 8% of patients on ziprasidone hydrochloride met this criterion after 26 weeks of treatment.

In child and adolescent patients (10-17 years of age) with a psychotic disorder other than schizophrenia-related, who completed 26 weeks of therapy with ziprasidone (67/169), 16% of patients met the criterion of a clinically significant weight change (adjusted for normal growth), as above.

Extrapyramidal Symptoms (EPS) in Children and Adolescent Population: The pediatric EPS data are from one short-term placebo controlled, monotherapy study in each of a schizophrenic patient population (6 week duration; 13-17 years of age), and a patient population with other psychotic disorder (4 week duration; 10-17 years of age). Across the two placebo-controlled studies, the incidences of adverse events potentially related to EPS showed a greater differential between ziprasidone and placebo compared to trials of adults with these indications (Incidence of EPS on ziprasidone was 25% and 30%, for schizophrenia and other, respectively, compared to 7% for placebo in both studies). The identified events included akathisia, drooling, dyskinesia, dystonia, gait disturbance, extrapyramidal disorder, muscle spasms/twitching, musculoskeletal stiffness, tremor, and torticollis.

Schizophrenia Open-label Data: In the 26-week, open label extension study that enrolled subjects from the placebo-controlled schizophrenia trial, a total of 15% (34/221) experienced EPS during that period. Of these, 8% (3/34) experienced an EPS event as severe. Of the subjects experiencing EPS 30% (10/34) had dose reductions, and none were discontinued from drug. In the schizophrenia patient population, EPS events of note in addition to the aggregated events listed above include: tardive dyskinesia, restless legs syndrome, masked facies and oculogyric crisis.

While the majority of extrapyramidal adverse events resolved within the duration of each study period, there were cases in which the events remained unresolved at the end of both the controlled and open-label trials.

QTc Prolongation Effects: Ziprasidone was associated with a mild to moderate dose-related prolongation of the QT interval in the pediatric clinical trials. There are insufficient data to determine whether this population is more vulnerable than adults to QT prolongation effects from ziprasidone hydrochloride.

Laboratory Abnormalities: The laboratory abnormalities observed during the double-blind phase of the two pediatric studies are presented in Table 3 (schizophrenia). Of note in the open-label phases, an ALT value of 763 U/L was observed in a patient, along with akathisia and fatigue, and an elevated AST value. No follow-up information is available. Decreased bicarbonate was reported at 23% (47/ 201) in the open-label schizophrenia study, and increased prolactin was reported at 20% (17 /86). One subject in the open-label schizophrenia study had an adverse event of increased prolactin of moderate severity.

Suicide-Related Events: In the ziprasidone pediatric studies, periodic searches were conducted of the adverse event database of each study to identify all possibly suicide-related adverse events (PSRAEs). These were reviewed by a blinded independent panel of experts and classified according to the C-CASA suicidality classification system (Columbia Classification Algorithm for Suicide Assessment). The incidence rates below exclude events classified as overdose due to dosing error.

In one 6-week, placebo-controlled trial in adolescent patients (13-17 years of age) with schizophrenia, the incidence of PSRAEs was 2.3% (5/193) for ziprasidone and 2.2% (2/90) for placebo. In the 26-week, open-label study that enrolled patients from the above trial (N=221). The incidence of PSRAEs was

4.1% (9/221). There was one completed suicide (a 17 year old female with a diagnosis of schizophrenia, disorganized type, receiving 160 mg ziprasidone).

In 26-week open label follow-up study in child and adolescent patients (10-17 years of age) with psychotic disorder other than schizophrenia-related, the incidence of PSRAEs was 9.3% (15/162).

The safety and efficacy of ziprasidone hydrochloride in children under the age of 18 years have not been established and its use is not recommended.

8.3 Less Common Clinical Trial Adverse Reactions

Less Common Clinical Trial Adverse Events (<1%) – Schizophrenia

All reported treatment-emergent events are included except those already listed in Table 1 or in other sections. It is important to emphasize that, although the events reported occurred during treatment with ziprasidone hydrochloride capsules, they were not necessarily caused by the therapy.

The adverse events are categorized by body system and listed in order of decreasing frequency according to the following definitions: frequent adverse events are those occurring in at least 1/100 patients (only those not already listed in the tabulated results from placebo-controlled trials appear in this listing); infrequent adverse events are those occurring in 1/100 to 1/1000 patients; rare events are those occurring in fewer than 1/1000 patients.

Body As a Whole – Frequent: abdominal pain, flu syndrome, fever, accidental fall, face edema, chills, photosensitivity reaction, flank pain, hypothermia, motor vehicle accident; Rare: feeling hot.

Cardiovascular System – Frequent: hypertension; Infrequent: bradycardia, angina pectoris, atrial fibrillation; Rare: first degree AV block, bundle branch block, phlebitis, pulmonary embolus, cardiomegaly, cerebral infarct, cerebrovascular accident, deep thrombophlebitis, myocarditis, thrombophlebitis.

Digestive System – Frequent: vomiting; Infrequent: rectal hemorrhage, dysphagia, tongue edema; Rare: gum hemorrhage, jaundice, fecal impaction, gamma glutamyl transpeptidase increased, hematemesis, cholestatic jaundice, hepatitis, hepatomegaly, leukoplakia of mouth, fatty liver deposit, melena.

Endocrine – Rare: hypothyroidism, hyperthyroidism, thyroiditis.

Hemic and Lymphatic System – Infrequent: anemia, ecchymosis, leukocytosis, leukopenia, eosinophilia, lymphadenopathy; Rare: thrombocytopenia, hypochromic anemia, lymphocytosis, monocytosis, basophilia, lymphedema, polycythemia, thrombocythemia.

Metabolic and Nutritional Disorders – Infrequent: thirst, transaminase increased, peripheral edema, hyperglycemia, creatine phosphokinase increased, alkaline phosphatase increased, hypercholesterolemia, dehydration, lactic dehydrogenase increased, albuminuria, hypokalemia; Rare:

BUN increased, creatinine increased, hyperlipemia, hypocholesteremia, hyperkalemia, hypochloremia, hypoglycemia, hyponatremia, hypoproteinemia, glucose tolerance decreased, gout, hyperchloremia, hyperuricemia, hypocalcemia, hypoglycemic reaction, hypomagnesemia, ketosis, respiratory alkalosis.

Musculoskeletal System – Infrequent: tenosynovitis; Rare: myopathy.

Nervous System – Frequent: agitation, tremor, dyskinesia, hostility, paresthesia, confusion, vertigo, hypokinesia, hyperkinesia, abnormal gait, oculogyric crisis, hypesthesia, ataxia, amnesia, cogwheel rigidity, delirium, hypotonia, akinesia, dysarthria, withdrawal syndrome, buccoglossal syndrome, choreoathetosis, diplopia, incoordination, neuropathy; Rare: myoclonus, nystagmus, torticollis, circumoral paresthesia, opisthotonos, reflexes increased, trismus.

Respiratory System – Frequent: dyspnea; Infrequent: pneumonia, epistaxis; Rare: hemoptysis, laryngismus.

Skin and Appendages – Infrequent: maculopapular rash, urticaria, alopecia, eczema, exfoliative dermatitis, contact dermatitis, vesicubullous rash. Drug reaction with eosinophilia and systemic symptoms (DRESS).

Special Senses – Infrequent: conjunctivitis, dry eyes, tinnitus, blepharitis, cataract, photophobia; Rare: eye hemorrhage, visual field defect, keratitis, keratoconjunctivitis.

Urogenital System – Infrequent: impotence, abnormal ejaculation, amenorrhea, hematuria, menorrhagia, female lactation, polyuria, urinary retention, metrorrhagia, male sexual dysfunction, anorgasmia, glycosuria; Rare: gynecomastia, vaginal hemorrhage, nocturia, oliguria, female sexual dysfunction, uterine hemorrhage.

8.5 Post-Market Adverse Reactions

Adverse event reports not listed above that have been received from spontaneous post-marketing reports for ziprasidone hydrochloride since market introduction are shown below (no causal relationship with ziprasidone has been established).

Cardiac Disorders: Tachycardia, torsades de pointes (in the presence of multiple confounding factors – see 7 WARNINGS AND PRECAUTIONS, QT Prolongation);

Gastrointestinal Disorders: Dysphagia, swollen tongue, severe constipation (see 7 WARNINGS AND PRECAUTIONS, Gastrointestinal);

Hemic and Lymphatic System - neutropenia, granulocytopenia and agranulocytosis;

Immune System Disorders: Allergic;

Metabolic and Nutritional Disorders: diabetic coma, lipids abnormal;

Nervous System Disorders: Facial droop, neuroleptic malignant syndrome, serotonin syndrome (alone or in combination with serotonergic medicinal products), tardive dyskinesia;

Psychiatric Disorders: Insomnia, mania/hypomania;

Renal and Urinary Disorders: Enuresis, urinary incontinence;

Reproductive System and Breast Disorders: Galactorrhea, priapism;

Skin and subcutaneous Tissue Disorders: Angioedema, rash; Stevens Johnson Syndrome, Drug reaction with eosinophilia and systemic symptoms (DRESS);

Vascular Disorders: Postural hypotension, syncope;

Sleep-related Disorders: Atypical antipsychotic drugs, including ziprasidone, have been reported with cases of sleep apnoea, with or without concomitant weight gain. In patients who have a history of or are at risk for sleep apnoea, Mylan-Ziprasidone should be prescribed with caution.

Risks of somnambulism (sleep walking) and sleep-related eating disorder have been reported with the use of atypical antipsychotics including ziprasidone.

9. DRUG INTERACTIONS

9.2 Drug Interactions Overview

Based upon the pharmacodynamic and pharmacokinetic profile of ziprasidone, possible interactions could be anticipated:

- Ziprasidone should not be used with any drug that prolongs the QT interval (see [2 CONTRAINDICATIONS](#) and [7 WARNINGS AND PRECAUTIONS](#)).
- Given the primary CNS effects of ziprasidone, caution should be used when it is taken in combination with alcohol and other centrally acting drugs.
- Because of its potential for inducing hypotension, ziprasidone may enhance the effects of certain antihypertensive agents.
- Concomitant use of CYP3A4 inhibitors/inducers could cause changes in the concentration of ziprasidone.
- Ziprasidone may antagonize the effects of levodopa and dopamine agonists.

9.3 Drug Behavioural Interactions

Smoking: Based on in vitro studies utilizing human liver enzymes, ziprasidone is a substrate for CYP1A2, however, the contribution of this pathway is minor. Consistent with these in vitro results, population pharmacokinetic evaluation has not revealed any significant differences in ziprasidone

pharmacokinetics between smokers and nonsmokers.

CNS Drugs/Alcohol: Given the primary CNS effects of ziprasidone, caution should be used when it is taken in combination with other centrally acting drugs, including alcohol and cannabis.

9.4 Drug-Drug Interactions

The drugs listed in this table are based on either drug interaction case reports or studies, or potential interactions due to the expected magnitude and seriousness of the interaction (i.e., those identified as contraindicated).

Table 4 – Established or Potential Drug-Drug Interactions

Proper/Common name	Source of Evidence	Effect	Clinical comment
CNS Drugs (e.g. antiparkinson agents, SSRIs, antipsychotics)	T		Given the primary CNS effects of ziprasidone, caution should be used when it is taken in combination with other centrally acting drugs.
Mood Stabilisers (e.g. lithium)	CT	Ziprasidone, at a dose of 40 mg BID, administered concomitantly with lithium at a dose of 450 mg BID for 7 days did not affect the steady-state level or renal clearance of lithium.	As ziprasidone and lithium are associated with cardiac conduction changes, the combination may pose a risk for pharmacodynamic interaction, including arrhythmias.
CYP3A4 Inducers / Inhibitors Ketoconazole	CT	Potent inhibitor of CYP3A4, at a dose of 400 mg per day for 5 days, increased the AUC and C _{max} of ziprasidone (80 mg BID) by approximately 35-40%. The serum concentration of S-methyl-dihydroziprasidone, at the expected T _{max} of ziprasidone, was increased by 55% during ketoconazole treatment. No additional QTc prolongation was observed. Other potent inhibitors of CYP3A4 would be expected to have similar effects.	Coadministration of potent CYP3A4 inhibitors has the potential of increasing ziprasidone serum concentrations. The clinical importance of this potential has not been clearly defined.
Carbamazepine	CT	An inducer of CYP3A4; administration of 200 mg BID for 25 days resulted in a decrease of	This effect may be greater when higher doses of carbamazepine are administered.

Proper/Common name	Source of Evidence	Effect	Clinical comment
Rifampin	T	approximately 36% in the AUC of ziprasidone (20 mg BID). An inducer of CYP3A4; co-administration could cause decreased concentrations of ziprasidone.	
Drugs that prolong QT interval (e.g. dofetilide, sotalol, quinidine, other Class Ia and III anti-arrhythmias, mesoridazine, thioridazine, chlorpromazine, droperidol, pimozide, sparfloxacin, gatifloxacin, moxifloxacin, halofantrine, mefloquine, pentamidine, arsenic trioxide, levomethadyl acetate, dolasetron mesylate, probucol or tacrolimus)	T	Ziprasidone causes a mild to moderate prolongation of the QT interval.	An additive effect of ziprasidone and other drugs that prolong the QT interval cannot be excluded.

Legend: C = Case Study; CT = Clinical Trial; T = Theoretical

9.5 Drug-Food Interactions

The absorption of ziprasidone is increased up to two-fold in the presence of food, so Mylan-Ziprasidone should always be taken with food (see [4.4 Administration](#)). Do not take with grapefruit or grapefruit juice as they are potent CYP3A4 inhibitors and could increase the concentration of ziprasidone.

9.6 Drug-Herb Interactions

Interactions between Mylan-Ziprasidone and the herbal remedy St. John's Wort may occur and may result in decreased concentrations of ziprasidone.

9.7 Drug-Laboratory Test Interactions

Interactions with laboratory tests have not been established.

10. CLINICAL PHARMACOLOGY

10.1 Mechanism of Action

The mechanism of action of ziprasidone, as with other drugs having efficacy in schizophrenia, is unknown. However, it has been proposed that the efficacy of this drug in schizophrenia is mediated through a combination of dopamine type 2 (D₂) and serotonin type 2 (5HT₂) antagonism.

Antagonism at receptors other than dopamine and 5HT₂ with similar receptor affinities may explain some of the other therapeutic and side effects of ziprasidone. Antagonism of histamine H₁ receptors may explain the somnolence observed with ziprasidone. Antagonism of α_1 -adrenergic receptors may explain the orthostatic hypotension observed with ziprasidone.

10.2 Pharmacodynamics

Ziprasidone exhibited high in vitro binding affinity for the dopamine D₂ and D₃, the serotonin 5-HT_{2A}, 5-HT_{2C}, 5-HT_{1A}, 5-HT_{1D}, and α_1 -adrenergic receptors (K_i = 4.8, 7.2, 0.4, 1.3, 3.4, 2, and 10 nM, respectively), and moderate affinity for the histamine H₁ receptor (K_i=47 nM). Ziprasidone functioned as an antagonist at the D₂, 5-HT_{2A}, and 5-HT_{1D} receptors, and as an agonist at the 5-HT_{1A} receptor.

Ziprasidone inhibited synaptic reuptake of serotonin and norepinephrine. No appreciable affinity was exhibited for other receptor/binding sites tested, including the cholinergic muscarinic receptor (IC₅₀ >1 mcM).

Cardiac Electrophysiology

Ziprasidone causes a mild to moderate dose-related prolongation of the QT interval.

10.3 Pharmacokinetics

Overview

The multiple-dose pharmacokinetics of ziprasidone are dose-proportional within the proposed clinical dose range, and ziprasidone accumulation is predictable with multiple dosing. Steady-state is attained within 1-3 days when dosing as recommended. The mean terminal phase half-life after multiple dosing in normal volunteers and schizophrenic patients is between 6 and 10 hours, with a range of individual values from 3 to 18 hours.

Ziprasidone's activity is primarily due to the parent drug. Ziprasidone is extensively metabolized after oral administration with only a small amount excreted in the urine (<1%) or feces (<4%) as unchanged drug. Ziprasidone is unlikely to interfere with the metabolism of drugs metabolized by cytochrome P450 enzymes.

Absorption

Following oral administration of multiple doses of ziprasidone with food, peak serum concentrations typically occur 6 to 8 hours post-dose. The absolute bioavailability of a 20 mg dose under fed conditions is approximately 60%. The absorption of ziprasidone is increased up to two-fold in the presence of food.

Distribution

Ziprasidone has a mean apparent volume of distribution of 1.5 L/kg. Twice daily dosing generally leads to attainment of steady state within 1–3 days.

Ziprasidone is greater than 99% bound to plasma proteins, binding primarily to albumin and α_1 -acid glycoprotein. The in vitro plasma protein binding of ziprasidone was not altered by warfarin or propranolol, two highly protein-bound drugs, nor did ziprasidone alter the binding of these drugs in human plasma. Thus, the potential for drug interactions with ziprasidone due to displacement is minimal.

Metabolism

Ziprasidone is extensively metabolized after oral administration with only a small amount excreted as unchanged drug in the urine (<1%) or feces (<4%). Unchanged ziprasidone represents about 44% of total drug-related material in serum.

Ziprasidone is primarily cleared via three metabolic routes (one route begins with reduction, the other two with oxidation) to yield four major circulating metabolites: S-methyldihydroziprasidone, via reduction then methylation, and benisothiazolepiperazine (BITP) sulphoxide, BITP-sulphone, and ziprasidone sulphoxide via oxidation routes.

Based on in vivo abundance of excretory metabolites, approximately two-thirds of ziprasidone metabolic clearance is mediated via reduction and methylation to generate S-methyldihydroziprasidone, while cytochrome P450-catalyzed oxidation mediates less than one third of ziprasidone clearance.

In vitro studies using human liver subcellular fractions indicate that the metabolite S-methyldihydroziprasidone is generated in two steps: the reduction reaction is mediated by aldehyde oxidase and potentially also by glutathione, while the subsequent methylation is mediated by thiol methyltransferase. There are no known clinically relevant inhibitors or inducers of aldehyde oxidase. In vitro studies using human liver microsomes and recombinant enzymes indicate that CYP3A4 is the major CYP contributing to the two routes of oxidative metabolism of ziprasidone. CYP1A2 may contribute to a much lesser extent.

Elimination

The mean terminal phase half-life after multiple dosing in normal volunteers and schizophrenic

patients is between 6 and 10 hours, with a range of individual values from 3 to 18 hours. Approximately 20% of ziprasidone dose is excreted in the urine, with approximately 66% being eliminated in the feces. S-methyldihydroziprasidone is mainly eliminated by biliary excretion and CYP3A4 metabolism. The sulphoxide is eliminated through renal excretion and by secondary metabolism catalyzed by CYP3A4.

Special Populations and Conditions

- **Pediatrics (< 18 years):** Safety and efficacy of ziprasidone hydrochloride in children have not been established; therefore, Health Canada has not authorized an indication for pediatric use.
- **Geriatrics and Sex:** In a multiple-dose (8 days of treatment) study involving n=32 subjects, there was no difference in the pharmacokinetics of ziprasidone between men and women or between elderly (>65 years) and young (18 to 45 years) subjects. Additionally, population pharmacokinetic evaluation of patients in controlled trials has revealed no evidence of clinically significant age or gender-related differences in the pharmacokinetics of ziprasidone. Dosage modifications for age or gender are, therefore, not recommended.
- **Ethnic Origin:** No specific pharmacokinetic study was conducted to investigate the effects of race. Population pharmacokinetic evaluation has revealed no evidence of clinically significant race-related differences in the pharmacokinetics of ziprasidone. Dosage modifications for race are, therefore, not recommended.
- **Hepatic Insufficiency:** As ziprasidone is cleared substantially by the liver, the presence of hepatic impairment would be expected to increase the AUC of ziprasidone; a multiple-dose study at the lowest therapeutic dose of 20 mg BID for 5 days in subjects with clinically significant (Childs-Pugh Class A and B) cirrhosis (n=13) revealed an increase in AUC₀₋₁₂ of 19% and 34% respectively, compared to a matched control group (n=13). A half-life of 7.1 hours was observed in subjects with cirrhosis compared to 4.8 hours in the control group. The effect of liver impairment on the serum concentrations of the metabolites is unknown.
- **Renal Insufficiency:** Because ziprasidone is highly metabolized with less than 1% of the drug excreted unchanged, renal impairment alone is unlikely to have a major impact on the pharmacokinetics of ziprasidone. The pharmacokinetic characteristics of ziprasidone following 8 days of treatment with 20 mg BID were similar among subjects with varying degrees of renal impairment (n=27), and subjects with normal renal function (n=9), indicating that dosage adjustment based upon the degree of renal impairment is not required. Ziprasidone is not removed by hemodialysis.

11. STORAGE, STABILITY AND DISPOSAL

Mylan-Ziprasidone should be stored at controlled room temperature between 15°C and 30°C.


12. SPECIAL HANDLING INSTRUCTIONS

No special handling is necessary for this drug product.

PART II: SCIENTIFIC INFORMATION

13. PHARMACEUTICAL INFORMATION

Drug Substance

Proper name:	Ziprasidone Hydrochloride Monohydrate
Chemical name:	5-[2-[4-(1,2-Benzisothiazol-3-yl)-1-piperazinyl]ethyl]-6-chloro-1,3-dihydro-2H-indol-2-one hydrochloride monohydrate or 5-(2-(4-(1,2-Benzisothiazol-3-yl)piperazinyl)ethyl)-6-chlorooxindole hydrochloride monohydrate
Molecular formula and molecular mass:	Molecular formula : $C_{21}H_{21}ClN_4OS.HCl.H_2O$ Molecular weight: 467.42 g/mol
Structural formula:	
Physicochemical properties:	Description: White to pink colour powder. Solubility: Soluble in Formic acid (anhydrous). Insoluble in n-Hexane. pH: 4.06 (1% water suspension) pKa : 6.3 Partition coefficient (n-octanol/water) at room temperature: 0.93 Hygroscopicity: Not hygroscopic in nature

14. CLINICAL TRIALS

14.1 Trial Design and Study Demographics

Schizophrenia Trials

The efficacy of oral ziprasidone hydrochloride in the treatment of schizophrenia was established in 4 short-term (4- to 6-week) and 1 long-term (52-week) placebo-controlled trials of psychotic inpatients who met DSM-III-R criteria for schizophrenia. Each study included 2-3 fixed doses of ziprasidone as well as placebo. Four (4) of the 5 trials were able to distinguish ziprasidone from placebo; 1 short-term study did not. Although a single fixed-dose haloperidol arm was included as a comparative treatment in 1 of the 3 short-term trials, this single study was inadequate to provide a reliable and valid comparison of ziprasidone and haloperidol.

Several instruments were used for assessing psychiatric signs and symptoms in these studies, among them the Brief Psychiatric Rating Scale (BPRS) and the Positive and Negative Syndrome Scale (PANSS), both multi-item inventories of psychopathology traditionally used to evaluate the effects of drug treatment in psychosis. The BPRS psychosis cluster (conceptual disorganization, hallucinatory behaviour, suspiciousness, and unusual thought content) is considered a particularly useful subset for assessing actively psychotic schizophrenic patients. A second traditional assessment, the Clinical Global Impression (CGI), reflects the impression of a skilled observer, fully familiar with the manifestations of schizophrenia, about the overall clinical state of the patient. In addition, the Scale for the Assessment of Negative Symptoms (SANS) was employed in some clinical trials.

14.2 Study Results

Schizophrenia Trials

- In a 4-week, placebo-controlled trial (n=139) comparing 2 fixed doses of ziprasidone hydrochloride (20 and 60 mg BID) with placebo, only the 60 mg BID dose was superior to placebo, on the BPRS total score and the CGI-S score. This higher dose group was not superior to placebo on the BPRS psychosis cluster or on the SANS.
- In a 6-week, placebo-controlled trial (n=302) comparing 2 fixed dose of ziprasidone hydrochloride (40 and 80 mg BID) with placebo, both dose groups were superior to placebo on the BPRS total score, the BPRS psychosis cluster score, the CGI severity score, and the PANSS total and negative subscale scores. Although the 80 mg BID dose group has a numerically greater effect than 40 mg BID dose group, the difference was not statistically significant.
- In a 6-week, placebo-controlled trial (n=419) comparing 3 fixed doses of ziprasidone hydrochloride (20, 60 and 100 mg BID) with placebo, all three dose groups were superior to placebo on the PANSS total score, the BPRS total score, the BPRS psychosis cluster, and the CGI-S. Only the 100 mg BID dose group was superior to placebo on the PANSS negative subscale score. There was no clear evidence for a dose- response relationship within the 20 mg BID and

100 mg BID dose range.

- In a 4-week, placebo-controlled trial (n=200) comparing 3 fixed doses of ziprasidone (5, 20 and 40 mg BID), none of the dose groups was statistically superior to placebo on any outcome of interest.
- A double-blind, randomized, parallel-group study was conducted in n=294 symptomatically stable inpatients with DSM-III-R diagnosis of chronic schizophrenia, who had been hospitalized for a period of not less than 2 months at study entry. Patients were randomized to 1 of 3 fixed doses of ziprasidone (20, 40, or 80 mg BID) or placebo and followed for 52 weeks. Patients were observed for “impending psychotic relapse”, defined as 2 consecutive study visit assessments showing a score of ≥ 6 (much worse or very much worse) on the CGI-improvement scale, and/or a score of ≥ 6 (moderately severe) on the hostility or uncooperativeness items of the PANSS. Ziprasidone was significantly superior to placebo in both time to relapse and rate of relapse, with no significant difference between the different dose groups.

There were insufficient data to examine population subsets based on age and race. Examination of population subsets based on gender did not reveal any differential responsiveness.

An analysis of the effect of ziprasidone on patients with clinically significant depressive symptoms (Montgomery-Asberg Depression Rating Scale, MADRS) ≥ 14 was conducted in 2 multicentre, placebo-controlled studies in acute schizophrenia. A statistically significant improvement versus placebo ($p < 0.05$) in the MADRS was observed in patients receiving ziprasidone 60 mg twice daily (n=32) in one study and 80 mg twice daily (n=56) in another study. The validity of this scale in patients with schizophrenia however is not established.

14.3 Comparative Bioavailability Studies

A blinded, randomized, two-treatment, two-period, single oral dose (1 x 20 mg), crossover comparative bioavailability study of Mylan-Ziprasidone capsules, 20 mg (Mylan Pharmaceuticals ULC) and ZELDOX[®] capsules, 20 mg (Pfizer Canada Inc.), was conducted in healthy, adult male subjects under fed conditions. Comparative bioavailability data from 32 subjects that were included in the statistical analysis are presented in the following table:

SUMMARY TABLE OF THE COMPARATIVE BIOAVAILABILITY DATA

Ziprasidone (1 x 20 mg) Geometric Mean Arithmetic Mean (CV %)				
Parameter	Test ¹	Reference ²	% Ratio of Geometric Means	90% Confidence Interval
AUC _T (ng·h/mL)	588.6 628.3 (36.4)	676.5 714.6 (35.1)	87.0	83.2 – 91.0

Ziprasidone (1 x 20 mg) Geometric Mean Arithmetic Mean (CV %)				
Parameter	Test ¹	Reference ²	% Ratio of Geometric Means	90% Confidence Interval
AUC _t (ng·h/mL)	602.4 642.9 (36.5)	693.0 731.9 (35.1)	86.9	83.2 – 90.8
C _{max} (ng/mL)	64.3 68.5 (35.1)	74.5 80.6 (40.8)	86.3	76.2 – 97.8
T _{max} ³ (h)	4.5 (1.0-10.0)	4.8 (2.0-10.0)		
T _½ ⁴ (h)	6.1 (19.0)	6.2 (18.9)		

¹ Mylan-Ziprasidone (ziprasidone as ziprasidone hydrochloride monohydrate) capsules, 20 mg (Mylan Pharmaceuticals ULC)

² ZELDOX® (ziprasidone as ziprasidone hydrochloride monohydrate) capsules, 20 mg (Pfizer Canada Inc.)

³ Expressed as the median (range) only

⁴ Expressed as the arithmetic mean (CV %) only

15. MICROBIOLOGY

No microbiological information is required for this drug product.

16. NON-CLINICAL TOXICOLOGY

General Toxicology:

Decreased motor activity occurred at all dose levels (10, 40 and 200 mg daily) in S-D rats fed ziprasidone orally over a 6 month period compared to controls. This was seen in both sexes but was more pronounced in the intermediate and high dose groups. Dose-related decreases in body weight gain were observed mostly in males. Several high dose males exhibited aggressive behaviour upon handling. Stress related or secondary changes such as adrenal hypertrophy were observed at the intermediate and high dose groups. The No Observed Adverse Effect Level (NOAEL) was determined to be 10 mg/kg/day.

In a 12 month toxicity study performed on Beagle dogs fed 5, 10 or 20 mg ziprasidone per day orally, all treatment-related effects were consistent with the pharmacologic properties of the compound, and included sedation, tremors, face pressing, pawing, cage biting, limb lifting/unusual postures, increased activity, aggressive behaviour, prolapse of the nictitating membrane and mammary gland development (intermediate dose females). Significant weight loss occurred in high dose males. There were no treatment-related effects noted in clinical pathology or histopathology parameters. Plasma drug concentrations exhibited a wide animal-to-animal variation within a given treatment group. Also, a difference in exposure was noted between male and female animals at the high and intermediate dose levels. The NOAEL was determined to be 10 mg/kg/day.

Carcinogenicity:

Mice

Ziprasidone was administered in the diet of CD-1 mice (50/sex/dose) at an initial dose of 50 mg/kg for the three treated groups. Two groups of 50/sex control mice received unsupplemented feed. On Day 15, the mid and high doses were increased to 100 mg/kg, and on Day 29 the high dose was increased to 200 mg/kg. The final dose levels were therefore 50, 100 and 200 mg/kg. It was concluded, at the end of this 24-month study, that treatment at the mid and high doses produced a statistically significant reduction both in body weight gain during the growth phase of the animals and in body weight in mice at the end of the study compared to controls. This was associated with a reduction in food and water consumption. Histopathological findings were limited to females and consisted of a dose-related increase in the incidence of hyperplasia and neoplasia in the pituitary gland (shown immunohistochemically to be prolactin-producing) and secondary changes in the mammary gland, ovaries and uterus. These findings were seen at 50 to 200 mg/kg/day, corresponding to systemic exposure about 1-4 times greater than that in humans; a no-effect dose level for these effects was not established.

Proliferative changes in the pituitary and mammary glands are not unexpected findings in rodents following treatment with this class of compounds, and are associated with increased prolactin concentrations.

Rats

Ziprasidone was administered in the diet of Long-Evans rats (50/sex/level) for 2 years at dose levels of 2, 6 and 12 mg/kg/day. All groups (low, intermediate and high) began at 2 mg/kg/day, and after 2 weeks the intermediate and high dose groups were raised to 6 mg/kg/day. After another two weeks, the high dose level group was increased to 12 mg/kg/day. Two identical control groups (50/sex/group) received non-medicated diet.

At dose levels up to 12 mg/kg/day, causing body weight decrements of approximately 10 to 20% relative to controls, ziprasidone showed no oncogenic potential in the rat.

Genotoxicity:

The results of a comprehensive battery of *in vivo* and *in vitro* genetic toxicology studies were generally negative, the exception being a slight increase in mutation frequency in *Salmonella typhimurium* TA 1537, but only at or near insoluble levels. Such results were not considered to represent a genotoxic hazard by ziprasidone due to the small response which lacked a true dose-relationship (positive only at the highest level tested), a reduction to non-significant levels by microsomal enzymes, a lack of mutagenic activity in urine from drug-treated mice, findings that gene mutation assays in mammalian cells *in vitro* were negative, and there was no indication of chromosomal mutation induction in mammalian cells either *in vivo* or *in vitro*.

Reproductive and Developmental Teratology:

There are no adequate and well-controlled studies in pregnant women.

Fertility and reproduction studies were conducted in rats. In a study where rats were fed ziprasidone 5, 10 or 40 mg/kg/day starting 10 days prior to mating for males and 2 weeks prior to mating and through gestation and lactation for females, sedation occurred at all dose levels but fertility was unaffected. Post-natal functional development testing indicated a slight delay in development that would be predicted based on the deficits in the body weights of the pups.

The NOAEL for fertility, defined as successful copulation and pregnancy, is 40 mg/kg, the highest dose tested. The NOAEL for reproduction and fetal/neonatal outcome is 5 mg/kg based on decreased gestational body weight gain at the 10 and 40 mg/kg dose levels, altered estrous cycles, decreased number of implantation sites, and number of viable pups at birth in litters from F0 dams at 40 mg/kg and decreased fetal body weights in the F1 offspring at the 10 and 40 mg/kg dose level. For all adult animals who were directly treated, the NOAEL is 5 mg/kg based on non-reproductive parameters.

In another studies where rats were fed ziprasidone 10, 40 or 160 mg/kg/day from 4 weeks prior to mating for males and 2 weeks prior to mating through gestation and post partum 10 days for females, sedation noted for all treatment groups. Food intake and body weights were decreased in a dose-related manner in male rats in all treated groups. Other signs included rough hair coat in males at 160 mg/kg and chromodacryorrhea in animals from both the 40 and 160 mg/kg dose groups.

Fertility was decreased in mating groups containing female rats treated with 160 mg/kg. The number of pups per litter was decreased at 160 mg/kg/day, while the proportion of pups born alive were decreased in litters from animals treated with 160 mg/kg. Survival of pups to postnatal day 4 decreased in all treated litters compared to control litters, particularly in the high dose group. The sedation observed in the dams after dosing was likely responsible for the decreased pup survival in the 160 mg/kg dose group.

Teratology studies were performed on female SD rats fed 5, 10 or 40 mg/kg/day of ziprasidone for 13 gestation days (days 6 – 18) and through lactation day 21. None of the dams died as a consequence of treatment. Mild to moderate sedation occurred at all dose levels, but did not interfere with food consumption, parturition, lactation or adequate maternal care of offspring. Mean body weight was significantly lower for the 40 mg/kg dams throughout gestation and lactation. Food consumption was not affected for any treated group.

The NOAEL for maternal effects is 10 mg/kg based on body weight inhibitions seen at 40 mg/kg. The NOAEL for postnatal development and behaviour of offspring is 5 mg/kg based on body weight inhibitions at 10 and 40 mg/kg, increased number of pups born dead and reduced number of pups alive on post natal day 4, delays in eye opening and air righting, and increased motor activity in females at 40 mg/kg.

Teratology studies were also performed on female rabbits fed 30 mg/kg/day of ziprasidone for 13 gestation days (days 6 – 18). All animals survived the dosing period, except one high dose animal which was found moribund and sacrificed on gestational day 27. The only clinical sign noted was occasional

loose or soft stool in 6/29 animals of the 30 mg/kg group. Mean maternal body weight gain and food consumption, indications of maternal toxicity, were significantly decreased during part or all of the treatment period. Reproductive parameters, mean fetal body weights and placental weights were unaffected by treatment. Visceral examination of the fetuses showed one fetus in each of the control and 30 mg/kg dose groups with a ventricular septal defect.

The findings support the conclusion that ziprasidone is not teratogenic in rabbits and confirm the previously noted NOAEL for fetuses to be 30 mg/kg.

Special Toxicology:

Ziprasidone is not a Class B Poison or a harmful substance upon either oral or dermal exposure. It is not considered a corrosive material, and it is not an ocular irritant. It did not produce a phototoxic reaction in BALB/c mice or induce either a systemic anaphylaxis reaction or passive cutaneous anaphylaxis reaction in guinea pigs.

17. SUPPORTING PRODUCT MONOGRAPHS

ZELDOX® (Capsules, 20 mg, 40 mg, 60 mg and 80 mg), submission control: 255625. Product Monograph, Upjohn Canada ULC. (DEC 22, 2021)

PATIENT MEDICATION INFORMATION

READ THIS FOR SAFE AND EFFECTIVE USE OF YOUR MEDICINE

Pr **MYLAN-ZIPRASIDONE**

Ziprasidone Capsules

Read this carefully before you start taking **Mylan-Ziprasidone** and each time you get a refill. This leaflet is a summary and will not tell you everything about this drug. Talk to your healthcare professional about your medical condition and treatment and ask if there is any new information about **Mylan-Ziprasidone**.

Serious Warnings and Precautions

Mylan-Ziprasidone can increase the risk of death when used in elderly people with dementia. The reasons for death varied but most were related to heart or circulatory problems or infections.

What is Mylan-Ziprasidone used for?

Mylan-Ziprasidone is used in adults to treat:

- schizophrenia

How does Mylan-Ziprasidone work?

Mylan-Ziprasidone belongs to a group of medicines called atypical antipsychotics. Exactly how Mylan-Ziprasidone works is unknown, however it seems to readjust the balance of the chemicals called dopamine and serotonin which are out of balance in people with schizophrenia. You should see an improvement in your symptoms within a few days of starting treatment with Mylan-Ziprasidone.

What are the ingredients in Mylan-Ziprasidone?

Medicinal ingredient: Ziprasidone hydrochloride monohydrate.

Non-medicinal ingredients: Lactose monohydrate, pollacrillin potassium, povidone, magnesium stearate.

Capsule shell (20 mg, 40 mg, 80 mg): black iron oxide, butyl alcohol, dehydrated alcohol, FD&C blue #1, gelatin, isopropyl alcohol, potassium hydroxide, propylene glycol, shellac, strong ammonia solution, titanium dioxide.

Capsule shell (60 mg): black iron oxide, butyl alcohol, dehydrated alcohol, gelatin, isopropyl alcohol, potassium hydroxide, propylene glycol, shellac, strong ammonia solution, titanium dioxide.

Mylan-Ziprasidone comes in the following dosage forms:

Capsules containing 20, 40, 60 and 80 mg of ziprasidone (as ziprasidone hydrochloride monohydrate).

Do not use Mylan-Ziprasidone if:

- You are allergic to ziprasidone hydrochloride.
- You are allergic to any of the other ingredients in Mylan-Ziprasidone or to a component of the container.
- You have the following heart conditions:
 - have QT prolongation (abnormal electrical activity of the heart) including a condition called long QT syndrome (a specific heart rhythm problem)
 - have had a recent heart attack
 - have heart failure
 - have certain irregularities of heart rhythm (discuss the specifics with your doctor)
- You are taking any of the following medicines:
 - heart medicines, such as dofetilide, sotalol, quinidine, other Class Ia and III anti-arrhythmics,
 - medicines which have been shown to have an effect on heart rhythm as indicated in their respective Patient Medication Information leaflet,
 - other anti-psychotic medicines such as mesoridazine, thioridazine, chlorpromazine, droperidol, pimozide,
 - other medications such as sparfloxacin (antibiotic medicine), levomethadyl acetate (opiate dependence medicine), dolasetron mesylate (nausea medicine), probucol (cholesterol lowering medicine) or tacrolimus (immunosuppressant medicine).
- Mylan-Ziprasidone is not recommended for use in children and adolescents under the age of 18 years.

To help avoid side effects and ensure proper use, talk to your healthcare professional before you take Mylan-Ziprasidone. Talk about any health conditions or problems you may have, including if you:

- are taking or have recently taken any prescription medicines
- are taking any over-the-counter medicines you can buy without a prescription, including natural/herbal remedies
- have had any problems with your heart
- have a family history of heart disease
- are taking any medications for heart disease or blood pressure that makes you prone to low blood pressure
- have a history of stroke or “mini-stroke”
- have liver problems
- have had any problem with fainting or dizziness
- have ever had blackouts or seizures
- have diabetes or a family history of diabetes
- are allergic to any medicines
- drink alcohol or use recreational drugs
- abuse drugs or have abused drugs in the past
- exercise vigorously or work in hot or sunny places
- suffer from lactose intolerance because Mylan-Ziprasidone capsules contain lactose

- have low white blood cell counts
- have low levels of potassium or magnesium in your blood
- are dehydrated or overhydrated
- suffer from electrolyte imbalances (which can occur during diuretic therapy, severe episodes of diarrhea or vomiting, if you suffer from an eating disorder or alcoholism)
- have or have had breast cancer
- have risk factors for developing blood clots such as: a family history of blood clots, age over 65, smoking, obesity, recent major surgery (such as hip or knee replacement), not being able to move due to air travel or other reason, or take oral contraceptives ("The Pill"). All possible risk factors for developing blood clots should be identified before and while you are taking Mylan-Ziprasidone and eliminated when possible.
- have a condition that increases your risk of having a seizure (such as Alzheimer's dementia)

Other warnings you should know about:

Heart Rhythm Problems:

You should not take Mylan-Ziprasidone if you have QT prolongation (abnormal electrical activity of the heart) including a condition called long QT syndrome (a specific heart rhythm problem) or if you are taking certain medicines. See the "Do not use Mylan-Ziprasidone if" section above for more information. Tell your doctor if you already have heart conditions, or if you are taking certain other medicines that may also change the way the electrical current in the heart works. Mylan-Ziprasidone may change the way the electrical current in your heart works and in rare cases cause dangerous heart rhythm abnormalities.

It is important to tell your physician, pharmacist or other healthcare professional that you are taking Mylan-Ziprasidone before you start taking any other drugs, including over-the-counter medications and natural/herbal remedies.

If you experience any symptoms of possible heart rhythm problems, such as dizziness, palpitations (sensation of rapid, pounding, or irregular heart beat), fainting, or seizures, you should get immediate medical help.

Pregnancy:

Tell your doctor if you are pregnant, think you might be pregnant or are planning to become pregnant. You should not take Mylan-Ziprasidone if you are pregnant. You must use a reliable method of birth control while you are taking Mylan-Ziprasidone. Talk to your doctor about reliable methods of birth control.

Breastfeeding:

Tell your doctor if you are breastfeeding or planning to breastfeed. Mylan-Ziprasidone is released into breast milk. It is not known if this is safe for your baby. You should not breastfeed a baby if you are taking Mylan-Ziprasidone.

Effects on Newborns:

In some cases, babies born to a mother taking ziprasidone capsules during pregnancy have experienced symptoms that are severe and require the newborn to be hospitalized. Be prepared to seek immediate emergency medical attention for your newborn if they have difficulty breathing, are overly sleepy, have muscle stiffness, or floppy muscles (like a rag doll), are shaking, or are having

difficulty feeding.

Alcohol:

The effects of alcohol could be made worse while taking Mylan-Ziprasidone. You should not drink alcohol while taking Mylan-Ziprasidone.

Driving and using machines:

Mylan-Ziprasidone can impair your judgment, thinking and motor skills and make you sleepy or dizzy. This may impair your ability to drive or to use machines. Wait until you know how Mylan-Ziprasidone affects you before driving or using machines. Do not drive or use machines if Mylan-Ziprasidone impairs your ability to do so safely.

Electrolyte imbalance:

Tell your doctor or pharmacist if you have diarrhea, vomiting, if you suffer from an eating disorder or alcoholism or any other illness that can cause you to lose fluids. Your doctor may want to check your blood to make sure that you have the right amount of important salts (“electrolytes”). This is because an imbalance in electrolytes is a risk factor for heart problems, which may occur more frequently with Mylan-Ziprasidone than with other anti-psychotics. Disordered eating, alcoholism and water intoxication are also risk factors for imbalance in electrolytes.

Since medications of the same drug class as Mylan-Ziprasidone may interfere with the ability of the body to adjust to heat, it is best to avoid becoming overheated or dehydrated while taking Mylan-Ziprasidone. Avoid vigorous exercise or being exposed to extreme heat.

Laboratory tests:

Your doctor should check your body weight before starting Mylan-Ziprasidone and during treatment. Your doctor should take blood tests before starting Mylan-Ziprasidone. They will monitor blood sugar, and the number of infection fighting white blood cells. Your doctor should continue to monitor your blood for as long as you are being treated.

If you have high levels of prolactin (measured with a blood test) and a condition called hypogonadism you may be at increased risk of breaking a bone due to osteoporosis. This occurs in both men and women.

Falls:

Feeling sleepy, a fall in blood pressure when you stand up from sitting or lying down, movement problems have been reported with the use of antipsychotic drugs. This can lead to falls that may cause fractures or other fall-related injuries.

Tell your healthcare professional about all the medicines you take, including any drugs, vitamins, minerals, natural supplements or alternative medicines.

The following may interact with Mylan-Ziprasidone:

How to take Mylan-Ziprasidone:

- Take Mylan-Ziprasidone exactly as your doctor has told you to.
- Swallow Mylan-Ziprasidone capsules whole with a glass of water.
- Do not open, crush or chew the capsules.

- Always take Mylan-Ziprasidone with a meal.
- Take Mylan-Ziprasidone at the same time each day.
- Do not change your dose or stop taking your medicine without your doctor’s approval.
- Dosage directions should be followed carefully. Never take more than the prescribed dose.
- Remember to keep taking Mylan-Ziprasidone, even when you feel better, to avoid relapse of symptoms. Mylan-Ziprasidone should be taken for as long as you and your doctor believe it is helping you.
- Never give Mylan-Ziprasidone to anyone else as this medicine has been prescribed only for you.

Overdose:

If you think you, or a person you are caring for, have taken too much Mylan-Ziprasidone, contact a healthcare professional, hospital emergency department, or regional poison control centre immediately, even if there are no symptoms. Take the medication package with you.

Missed Dose:

If you miss a dose of Mylan-Ziprasidone, wait until your next scheduled dose. **Do not take 2 doses at once.**

What are possible side effects from using Mylan-Ziprasidone?

These are not all the possible side effects you may have when taking Mylan-Ziprasidone. If you experience any side effects not listed here, tell your healthcare professional.

Side effects may include:

- feeling unusually tired or sleepy
- nausea or upset stomach
- restlessness
- diarrhea
- increased cough/runny nose

Serious side effects and what to do about them			
Symptom / effect	Talk to your healthcare professional		Stop taking drug and get immediate medical help
	Only if severe	In all cases	
UNKNOWN			
Blood disorders: fever, chills, weakness, sore throat, sores in the mouth or throat, bleeding gums, bone pain, low blood pressure, fast heartbeat, and trouble breathing, muscle weakness, fatigue, difficulty swallowing, difficulty breathing, protrusion of the tongue		✓	
Falls due to feeling dizzy, or faint		✓	
COMMON			

Serious side effects and what to do about them			
Symptom / effect	Talk to your healthcare professional		Stop taking drug and get immediate medical help
	Only if severe	In all cases	
Skin rash on its own	✓		
Dystonia (movement disorder): muscle spasms that you cannot control, neck spasms, tightness of the throat, difficulty swallowing, breathing problems, tongue sticking out			✓
Extrapyramidal symptoms (movement disorder): feeling restless, tense, involuntary muscle contractions, continuous spasms, rigidity, slowness of movement, tremor, jerky movements		✓	
Tardive dyskinesia (movement disorder): muscle twitching or abnormal movement of your face or tongue		✓	
Stroke (bleeding or blood clot in the brain): sudden weakness or numbness of the face, arms, or legs and speech or vision problems			✓
Weight gain	✓		
UNCOMMON			
Heart rhythm problems: dizziness, palpitations (sensation of rapid, pounding or irregular heart beat), fainting, or seizures			✓
Blood clots: swelling, pain and redness in an arm or leg that can be warm to touch. You may develop sudden chest pain, difficulty breathing and heart palpitations.		✓	
Seizure: loss of consciousness with uncontrollable shaking, “fit”			✓
Allergic reaction: skin rash, hives, swelling of throat and tongue, difficulty breathing			✓
Hyperprolactinemia (elevated prolactin levels): irregular menstrual cycles, production and discharge of breast milk, abnormal		✓	

Serious side effects and what to do about them			
Symptom / effect	Talk to your healthcare professional		Stop taking drug and get immediate medical help
	Only if severe	In all cases	
hair growth, infertility			
Serotonin toxicity (too much serotonin in the body): agitation, restlessness, confusion, rapid heart rate, dilated pupils, loss of muscle coordination, twitching muscles, rigidity, heavy sweating, diarrhea			✓
RARE			
Neuroleptic malignant syndrome (nervous system disorder): High fever with pronounced muscle stiffness, state of confusion, rapid or irregular heartbeat, profuse sweating			✓
Priapism: Long lasting (greater than 4 hours in duration) and painful erection of the penis.			✓
Feeling very hot and unable to cool down (generally as a result of several factor together, such as vigorous exercise, dehydration, warm conditions)		✓	
New or worsening constipation		✓	
Severe allergic reaction: skin rash, fever, chest pain and abdominal pain			✓
Skin rash and fever with swollen glands			✓
Severe skin reactions: Fever, severe rash, swollen lymph glands, flu-like feeling, blisters and peeling skin that may start in and around the mouth, nose, eyes and genitals and spread to other areas of the body, yellow skin or eyes, shortness of breath, dry cough, chest pain or discomfort, feeling thirsty, urinating less often, less urine			✓
VERY RARE			
Hyperglycemia (high blood sugar): extreme thirst, frequent urination,		✓	

Serious side effects and what to do about them			
Symptom / effect	Talk to your healthcare professional		Stop taking drug and get immediate medical help
	Only if severe	In all cases	
excessive hunger, weakness, nausea, vomiting, fruity-smelling breath			

If you have a troublesome symptom or side effect that is not listed here or becomes bad enough to interfere with your daily activities, tell your healthcare professional.

Reporting Side Effects

You can report any suspected side effects associated with the use of health products to Health Canada by:

- Visiting the Web page on Adverse Reaction Reporting (<https://www.canada.ca/en/health-canada/services/drugs-health-products/medeffect-canada/adverse-reaction-reporting.html>) for information on how to report online, by mail or by fax; or
- Calling toll-free at 1-866-234-2345.

NOTE: Contact your health professional if you need information about how to manage your side effects. The Canada Vigilance Program does not provide medical advice.

Storage:

Keep Mylan-Ziprasidone and all medicines out of the reach and sight of children.

Store Mylan-Ziprasidone at controlled room temperature (15°C to 30°C).

If your doctor tells you to stop taking Mylan-Ziprasidone or if your medicine has expired, return any leftover medicine to your pharmacist for proper discarding.

If you want more information about Mylan-Ziprasidone:

- Talk to your healthcare professional
- Find the full product monograph that is prepared for healthcare professionals and includes this Patient Medication Information by visiting the Health Canada website: (<https://www.canada.ca/en/health-canada/services/drugs-health-products/drug-products/drug-product-database.html>); the manufacturer's website www.mylan.ca, or by calling 1-844-596-9526.

This leaflet was prepared by Mylan Pharmaceuticals ULC
Etobicoke, Ontario M8Z 2S6

Last revised: APR 18, 2023



Mylan Pharmaceuticals ULC
Etobicoke, ON M8Z 2S6
1-844-596-9526
www.mylan.ca