

PRODUCT MONOGRAPH
INCLUDING PATIENT MEDICATION INFORMATION

PrANAFRANIL[®]

clomipramine hydrochloride tablets

Tablets, 10 mg, 25 mg and 50 mg, Oral

ATC code: N06AA04

Antidepressant / Antiobsessional

Apotex Inc.
150 Signet Drive
Toronto, Ontario
M9L 1T9

Date of Initial Authorization:
SEP 28, 2020

Date of Revision:
MAY 02, 2023

Submission Control Number: 268209

RECENT MAJOR LABEL CHANGES

1 Indications, 1.1 Pediatrics	05/2023
2 Contraindications	05/2023
3 Serious Warnings and Precautions Box	05/2023
4 Dosage and Administration, 4.1 Dosing Considerations	05/2023
4 Dosage and Administration, 4.2 Recommended Dose and Dosage Adjustment	05/2023
7 Warnings and Precautions, Endocrine and Metabolism	05/2023
7 Warnings and Precautions, Hematologic	05/2023
7 Warnings and Precautions, Musculoskeletal	05/2023
7 Warnings and Precautions, Neurologic	05/2023
7 Warnings and Precautions, Ophthalmologic	05/2023
7 Warnings and Precautions, Peri-Operative Considerations	05/2023
7 Warnings and Precautions, Psychiatric	05/2023
7 Warnings and Precautions, Skin	05/2023
7 Warnings and Precautions, 7.1.3 Pediatrics	05/2023
7 Warnings and Precautions, 7.1.4 Geriatrics	05/2023

TABLE OF CONTENTS

Sections or subsections that are not applicable at the time of authorization are not listed.

RECENT MAJOR LABEL CHANGES	2
TABLE OF CONTENTS	2
PART I: HEALTH PROFESSIONAL INFORMATION	5
1 INDICATIONS.....	5
1.1 Pediatrics.....	5
1.2 Geriatrics	5
2 CONTRAINDICATIONS	6
3 SERIOUS WARNINGS AND PRECAUTIONS BOX	6
4 DOSAGE AND ADMINISTRATION	7
4.1 Dosing Considerations	7

4.2	Recommended Dose and Dosage Adjustment	9
4.4	Administration.....	11
4.5	Missed Dose	11
5	OVERDOSAGE.....	11
6	DOSAGE FORMS, STRENGTHS, COMPOSITION AND PACKAGING	13
7	WARNINGS AND PRECAUTIONS	13
7.1	Special Populations.....	21
7.1.1	Pregnant Women	21
7.1.2	Breast-feeding	21
7.1.3	Pediatrics.....	21
7.1.4	Geriatrics	22
8	ADVERSE REACTIONS	22
8.1	Adverse Reactions Overview	22
8.2	Clinical Trial Adverse Reactions	22
8.5	Post-Market Adverse Reactions	25
9	DRUG INTERACTIONS	25
9.1	Serious Drug Interactions	25
9.2	Drug Interactions Overview.....	26
9.3	Drug-Behavioural Interactions.....	27
9.4	Drug-Drug Interactions.....	27
9.5	Drug-Food Interactions.....	33
9.6	Drug-Herb Interactions.....	34
9.7	Drug-Laboratory Test Interactions.....	34
10	CLINICAL PHARMACOLOGY	34
10.1	Mechanism of Action.....	34
10.2	Pharmacodynamics	34
10.3	Pharmacokinetics	36
11	STORAGE, STABILITY AND DISPOSAL	37
12	SPECIAL HANDLING INSTRUCTIONS	38
PART II: SCIENTIFIC INFORMATION		39
13	PHARMACEUTICAL INFORMATION	39

14	CLINICAL TRIALS	39
15	MICROBIOLOGY	39
16	NON-CLINICAL TOXICOLOGY	39
	PATIENT MEDICATION INFORMATION.....	42

PART I: HEALTH PROFESSIONAL INFORMATION

1 INDICATIONS

ANAFRANIL (clomipramine hydrochloride tablets) is indicated for:

- the treatment of depression. ANAFRANIL also appears to have a mild sedative effect which may be helpful in alleviating the anxiety component often accompanying depression.
- the treatment of obsessions and compulsions in patients with obsessive compulsive disorder (OCD). The obsessions and compulsions must cause marked distress, be time-consuming, or significantly interfere with social or occupational functioning.

The effectiveness of ANAFRANIL for long-term use (e.g., for more than 10 weeks) has not been systematically evaluated in placebo-controlled trials. The healthcare professional who elects to prescribe ANAFRANIL for extended periods should periodically re-evaluate the long-term usefulness of the drug for the individual patient (see [4 DOSAGE AND ADMINISTRATION](#)).

1.1 Pediatrics

Depression

Pediatrics (<18 years):

No data are available to Health Canada; therefore, Health Canada has not authorized an indication for pediatric use in children under 18 years of age (see [3 SERIOUS WARNINGS AND PRECAUTIONS BOX](#); [7 WARNINGS AND PRECAUTIONS Clinical Worsening and Suicide Risk](#)).

Obsessive Compulsive Disorder

Pediatrics (<10 years):

No data are available to Health Canada; therefore, Health Canada has not authorized an indication for pediatric use in children under 10 years of age.

Pediatrics (10 to 17 years): Limited information is available for the use of ANAFRANIL in children aged 10 to 17 years, therefore, ANAFRANIL is not recommended for use in this population (see [4.2 Recommended Dose and Dosage Adjustment, Obsessive Compulsive Disorders, Children and Adolescents](#)).

1.2 Geriatrics

Geriatrics (≥ 65 years): Evidence from clinical studies and experience suggests that use in the geriatric population is associated with differences in safety or effectiveness (see [4.2 Recommended Dose and Dosage Adjustment, Depression, Elderly and Debilitated Patients](#); [4.2 Recommended Dose and Dosage Adjustment, Obsessive Compulsive Disorder, Elderly and Debilitated Patients](#) and [7.1.4 Geriatrics](#)).

2 CONTRAINDICATIONS

ANAFRANIL (clomipramine hydrochloride) is contraindicated:

- in patients who are hypersensitive to this drug or to other tricyclic antidepressants (TCAs) belonging to the dibenzazepine group or to any ingredient in the formulation, including any non-medicinal ingredient, or component of the container. For a complete listing, see [6 DOSAGE FORMS, STRENGTHS, COMPOSITION AND PACKAGING](#).
- in conjunction with, or within 14 days before or after treatment with a monoamine oxidase inhibitor (MAOI) (see [9.1 Serious Drug Interactions](#)). Hypertensive crises, hyperactivity, hyperpyrexia, spasticity, severe convulsions or coma, and death have been reported in patients receiving MAO inhibitors and TCAs (see [4.1 Dosing Considerations](#); [7 WARNINGS AND PRECAUTIONS, Neurologic, Serotonin toxicity/Serotonin syndrome](#); and [9.1 Serious Drug Interactions](#)).

Starting ANAFRANIL in a patient who is being treated with linezolid or intravenous methylene blue is also contraindicated because of an increased risk of serotonin syndrome (see [4.1 Dosing Considerations](#) and [7 WARNINGS AND PRECAUTIONS, Neurologic, Serotonin toxicity/Serotonin syndrome](#)).

- during the acute recovery phase following myocardial infarction and in the presence of acute congestive heart failure- (see [7 WARNINGS AND PRECAUTIONS, Cardiovascular](#); [Peri-Operative Considerations](#) and [8.2 Clinical Trial Adverse Reactions, Cardiac disorders](#)).
- in patients with existing liver or kidney damage, or in patients with a history of blood dyscrasias.
- in patients with glaucoma, as the condition may be aggravated due to the atropine-like effects of the drug.

3 SERIOUS WARNINGS AND PRECAUTIONS BOX

Serious Warnings and Precautions

Cardiovascular: Tricyclic antidepressants (TCAs), particularly in high doses, have been reported to produce sinus tachycardia, changes in conduction time and arrhythmias. A few instances of unexpected death have been reported in patients with cardiovascular disorders. Myocardial infarction and stroke have also been reported with drugs of this class. Therefore, ANAFRANIL should be administered with extreme caution to patients with a history of cardiovascular disorders, especially those with cardiovascular insufficiency, conduction disorders (e.g., atrioventricular block grades I to III) or other arrhythmias, those with circulatory lability and elderly patients (see [7 WARNINGS AND PRECAUTIONS, Cardiovascular](#)).

Anticholinergic effects: Because of its anticholinergic properties, ANAFRANIL should be used with caution in patients with increased intraocular pressure, narrow angle glaucoma or urinary retention, particularly in the presence of prostatic hypertrophy (see [7](#)

[WARNINGS AND PRECAUTIONS, Ophthalmologic](#)).

Endocrine and Metabolism: Caution should be observed in prescribing ANAFRANIL for hyperthyroid patients or for patients receiving thyroid medication. Transient cardiac arrhythmias have occurred in rare instances in patients who have been receiving other tricyclic compounds concomitantly with thyroid medication (see [7 WARNINGS AND PRECAUTIONS, Endocrine and Metabolism](#)).

Seizures: Tricyclic agents are known to lower the convulsive threshold and ANAFRANIL should, therefore, be used with extreme caution in patients with a history of convulsive disorders and other predisposing factors, e.g., brain damage of varying etiology, concomitant use of neuroleptics, alcoholism and withdrawal from alcohol, and concomitant use with other drugs that lower the seizure threshold. It appears that the occurrence of seizures is dose dependent (see [7 WARNINGS AND PRECAUTIONS, Neurologic](#)).

Clinical Worsening and Suicide Risk: Increased risk of self-harm, harm to others, suicidal thinking and behavior with antidepressant use. Closely monitor all antidepressant-treated patients for clinical worsening and for emergence of agitation-type and/or suicidal thoughts and behaviors (see [7 WARNINGS AND PRECAUTIONS, Psychiatric, Clinical Worsening and Suicide Risk](#)).

4 DOSAGE AND ADMINISTRATION

4.1 Dosing Considerations

- The dosage of ANAFRANIL (clomipramine hydrochloride) should be individualized according to the requirements of each patient.
- Treatment should be initiated at the lowest recommended dose and increased gradually, noting carefully the clinical response and any evidence of intolerance.
- During the initial dose titration phase, the total daily dose of ANAFRANIL should be divided and administered with meals to reduce gastrointestinal side effects.
- Owing to the long elimination half-lives of ANAFRANIL and its active metabolite, desmethylclomipramine, steady-state plasma levels may not be achieved until 2 to 3 weeks after a dosage adjustment. It may thus be advisable to wait 2 to 3 weeks after the initial dose titration phase, before attempting further dosage adjustments. Plasma ANAFRANIL measurement would constitute the optimal guide to dosage monitoring.
- It should be kept in mind that a lag in therapeutic response usually occurs at the onset of therapy, lasting from several days to a few weeks. Increasing the dosage does not normally shorten this latent period and may increase the incidence of side effects.

Cardiovascular

- Prior to initiating treatment with ANAFRANIL, a cardiac evaluation, including blood pressure and electrocardiogram (ECG) examinations, should be performed, particularly in patients with a history of cardiovascular disorders (see [7 WARNINGS AND PRECAUTIONS, Cardiovascular](#)).

Signs of intolerance and toxicity

- Clinical symptoms of intolerance, especially drowsiness, dizziness, and postural hypotension, should alert the healthcare professional to the need for reduction in dosage- (see [5 OVERDOSAGE](#)).
- The best available evidence of impending toxicity from very high doses of ANAFRANIL is prolongation of the QRS or QT intervals on the ECG. Prolongation of the PR interval is also significant, but less closely correlated with plasma levels.
- If serious adverse events occur, the dosage should be reduced or treatment altered.

Switching a Patient To or From a Monoamine Oxidase Inhibitor (MAOI) Intended to Treat Psychiatric Disorders

- At least 14 days should elapse between discontinuation of a MAOI intended to treat psychiatric disorders and initiation of therapy with ANAFRANIL. Conversely, at least 14 days should be allowed after stopping ANAFRANIL before starting a MAOI intended to treat psychiatric disorders (see [2 CONTRAINDICATIONS](#)).

Use of ANAFRANIL With Other MAOIs, Such as Linezolid or Methylene Blue

- Do not start ANAFRANIL in a patient who is being treated with linezolid or intravenous methylene blue because there is increased risk of serotonin syndrome. In a patient who requires more urgent treatment of a psychiatric condition, other interventions, including hospitalization, should be considered (see [2 CONTRAINDICATIONS](#)).
- In some cases, a patient already receiving ANAFRANIL therapy may require urgent treatment with linezolid or intravenous methylene blue. If acceptable alternatives to linezolid or intravenous methylene blue treatment are not available and the potential benefits of linezolid or intravenous methylene blue treatment are judged to outweigh the risks of serotonin syndrome in a particular patient, ANAFRANIL should be stopped promptly, and linezolid or intravenous methylene blue can be administered. The patient should be monitored for symptoms of serotonin syndrome for two weeks or until 24 hours after the last dose of linezolid or intravenous methylene blue, whichever comes first. Therapy with ANAFRANIL may be resumed 24 hours after the last dose of linezolid or intravenous methylene blue (see [7 WARNINGS AND PRECAUTIONS, Neurologic, Serotonin toxicity / Serotonin syndrome](#)).
- The risk of administering methylene blue by non-intravenous routes (such as oral tablets or by local injection) or in intravenous doses much lower than 1 mg/kg with ANAFRANIL is unclear. The clinician should, nevertheless, be aware of the possibility of emergent symptoms of serotonin syndrome with such use (see [7 WARNINGS AND PRECAUTIONS, Neurologic, Serotonin toxicity / Serotonin syndrome](#)).

Monitor for agitation, suicidal tendencies.

- Rigorous clinical monitoring for suicidal ideation or other indicators of potential for suicidal behaviour is advised in patients of all ages, especially when initiating therapy or during any change in dose or dosage regimen. This includes monitoring for agitation-type emotional and behavioural changes (see [7 WARNINGS AND PRECAUTIONS, Psychiatric, Clinical Worsening and Suicide Risk](#)).

4.2 Recommended Dose and Dosage Adjustment

Depression

Initial Dosage

Adults

ANAFRANIL therapy should be initiated at daily doses of 25 mg. Dosage may be increased by 25 mg increments, as tolerated, at 3 to 4 day intervals up to a total daily dose of 150 mg by the end of 2 weeks. Thereafter, the dose may be gradually increased over a period of several weeks to 200 mg. Doses in excess of 200 mg daily are not recommended for outpatients. Occasionally, in more severely depressed hospitalized patients, dosages up to 300 mg daily may be required.

Elderly and Debilitated Patients

In general, lower dosages are recommended for these patients. Initially, 20 to 30 mg daily in divided doses is suggested, with very gradual increments, depending on tolerance and response. Blood pressure and cardiac rhythm should be checked frequently, particularly in patients who have unstable cardiovascular function.

Pediatrics (<18 years)

Health Canada has not authorized an indication for pediatric use (<18 years).

Maintenance Dosage

Dosage during maintenance therapy should be kept at the lowest effective level. To minimize daytime sedation during maintenance treatment, the total daily dosage may be given as a single dose at bedtime. Medication should be continued for the expected duration of the depressive episode in order to minimize the possibility of relapse following clinical improvement.

Obsessive Compulsive Disorders

Initial Dosage

Adults

ANAFRANIL therapy in adult Obsessive Compulsive patients should be initiated at daily doses of 25 mg. Dosage may be increased by 25 mg increments, as tolerated, at 3 to 4 day intervals up to a total daily dose of 100 or 150 mg by the end of 2 weeks. Thereafter, the dose may be gradually increased over a period of several weeks to 200 mg. Doses in excess of 200 mg per day are not generally recommended for outpatients. However, in the treatment of severe cases of Obsessive Compulsive Disorder (OCD), daily doses of up to 250 mg may be required.

Children and Adolescents (10 to 17 years)

Limited information is available for the use of ANAFRANIL in children aged 10 to 17 years. For this age group, an initial dose of 25 mg per day is recommended. Dosage may be increased by 25 mg increments, as tolerated, at 3 to 4 day intervals. By the end of 2 weeks, patients may be titrated up to 100 to 150 mg per day or 3 mg/kg, whichever is lower. Thereafter, the dose may be gradually increased to 200 mg or 3 mg/kg whichever is lower. A total daily dose above 200 mg should not be used in children or adolescents.

Pediatrics (<10 years)

Health Canada has not authorized an indication for pediatric use (<10 years).

Elderly and Debilitated Patients

In general, lower dosages are recommended for these patients. Initially, 20 to 30 mg daily in divided doses is suggested, with very gradual increments, depending on tolerance and response. Blood pressure and cardiac rhythm should be checked frequently, particularly in patients who have unstable cardiovascular function.

Maintenance Dosage (Adults, Children and Adolescents (10 to 17 years))

Double-blind extension phase studies of ANAFRANIL therapy in patients with OCD have followed patients for up to 52 weeks. Although placebo enrollment in these studies was inadequate to permit a controlled comparison, data suggest that ANAFRANIL therapy can be continued for up to a year without loss of efficacy.

Dosage adjustments may be made during maintenance therapy with the objective of maintaining the patient at the lowest effective dose. To minimize daytime sedation during maintenance treatment, the total daily dosage may be given as a single dose at bedtime. If symptoms recur, the dosage should be increased until the symptoms are controlled. Patients should be reassessed periodically to determine the need for continued treatment. To avoid withdrawal symptoms upon discontinuation of therapy, a gradual decrease in dosage and careful patient monitoring are recommended.

Pediatrics (<10 years)

Health Canada has not authorized an indication for pediatric use (<10 years).

Concomitant use with inhibitors of cytochrome P450 enzymes

Concomitant use of agents in the tricyclic antidepressant (TCA) class (which includes ANAFRANIL) with drugs that can inhibit cytochrome P450 2D6 (ex: fluoxetine) may require lower doses than usually prescribed for either the tricyclic antidepressant agent or the other drug. Furthermore, whenever one of these drugs is withdrawn from co-therapy, an increased dose of a TCA may be required. It is desirable to monitor TCA plasma levels whenever an agent of the TCA class, including ANAFRANIL, is going to be co-administered with another drug known to be an inhibitor of P450 2D6 (and/or P450 1A2) (see [9.2 Drug Interactions Overview](#)).

CYP3A and CYP2C inducers

CYP3A and CYP2C inducers, such as rifampicin or anticonvulsants (e.g. barbiturates, carbamazepine, phenobarbital and phenytoin), may decrease clomipramine concentrations as concomitant administration of drugs known to induce cytochrome P450 enzymes, particularly CYP3A4, CYP2C19 may accelerate the metabolism and decrease the efficacy of ANAFRANIL.

4.4 Administration

ANAFRANIL tablets should be swallowed whole orally. Do not break, chew or crush. ANAFRANIL can be administered with or without food.

4.5 Missed Dose

If the patient misses a dose, instruct the patient to take the dose as soon as they remember. If it is almost time for the next dose, inform the patient to skip the missed dose and continue the regular dosing schedule.

5 OVERDOSAGE

Since children may be more sensitive than adults to acute overdosage with tricyclic antidepressants (TCAs), and since fatalities in children have been reported, effort should be made to avoid potential overdose particularly in this age group.

Deaths by deliberate or accidental overdosage have occurred with this class of drugs. Since the propensity for suicide is high in depressed patients, a suicide attempt by other means may occur during the recovery phase. The possibility of simultaneous ingestion of other drugs should also be considered.

Signs and Symptoms of Overdosage

These may vary in severity depending on various factors such as the amount of drug absorbed, the interval between drug ingestion and start of treatment, and the age of the patient. Accidental ingestion in children should be regarded as serious and potentially fatal.

Signs and symptoms generally appear within 4 hours of ingestion and reach maximum severity after 24 hours. Owing to delayed absorption (increased anticholinergic effect due to overdose), long half-life and enterohepatic recycling of the drug, the patient may be at risk for up to 4 to 6 days.

Signs and symptoms may include drowsiness, stupor, ataxia, vomiting, cyanosis, restlessness, agitation, delirium, severe perspiration, hyperactive reflexes, muscle rigidity, athetoid and choreiform movements, and convulsions. Hyperpyrexia, mydriasis, oliguria or anuria, bowel and bladder paralysis, and respiratory depression may occur.

Hypotension and initial hypertension may occur. However, the usual finding is increasing hypotension which may lead eventually to shock. Serious cardiovascular disturbances are frequently present, including tachycardia, cardiac arrhythmias (flutter, atriofibrillation, premature ventricular beats and ventricular tachycardia) as well as impaired myocardial conduction, atrioventricular and intraventricular block, electrocardiogram (ECG) abnormalities

(such as widened QRS complexes and marked S-T shifts, QTc prolongation), signs of congestive heart failure and cardiac arrest. Coma may ensue.

Treatment of Overdosage

Patients in whom overdosage is suspected should be admitted to hospital without delay. No specific antidote is available and treatment is essentially symptomatic and supportive.

Administration of activated charcoal may help to reduce absorption of the drug. Do not induce vomiting. As ANAFRANIL (clomipramine hydrochloride) is largely protein bound, forced diuresis, peritoneal dialysis and hemodialysis are unlikely to be of value.

Treatment should be designed to ensure maintenance of the vital functions. An open airway should be maintained in comatose patients and assisted ventilation instituted, if necessary, but respiratory stimulants should not be used. Hyperpyrexia should be controlled by external measures, such as ice packs and cooling sponge baths. Acidosis may be treated by cautious administration of sodium bicarbonate. Adequate renal function should be maintained.

ECG monitoring in an intensive care unit is recommended in all patients, particularly in the presence of ECG abnormalities, and should be maintained for several days after the cardiac rhythm has returned to normal. Unexpected deaths attributed to cardiac arrhythmias have been reported several days following an apparent recovery from tricyclic antidepressant overdose. Correction of hypoxia and acidosis, if present, may be beneficial. Correction of metabolic acidosis and low potassium concentrations by means of bicarbonate I.V. and potassium substitution may also be effective for treatment of arrhythmias. If bradyarrhythmia or AV-block occur, consider temporary insertion of a cardiac pacemaker. Because of its effect on cardiac conduction, digitalis should be used only, with caution. If rapid digitalization is required for the treatment of congestive heart failure, special care should be exercised in using the drug.

External stimulation should be minimized to reduce the tendency to convulsions. If convulsions occur, anticonvulsants (preferably intravenous diazepam) should be administered. Barbiturates may intensify respiratory depression, particularly in children, and aggravate hypotension and coma. Paraldehyde may be used in some children to counteract muscular hypertonus and convulsions with less likelihood of causing respiratory depression. If the patient fails to respond rapidly to anticonvulsants, artificial ventilation should be instituted. Prompt control of convulsions is essential since they aggravate hypoxia and acidosis and may thereby precipitate cardiac arrhythmias and arrest.

Shock should be treated with supportive measures, such as intravenous fluids, plasma expanders and oxygen. The use of corticosteroids in shock is controversial and may be contraindicated in tricyclic antidepressant overdose. Hypotension usually responds to elevation of the foot of the bed. Pressor agents (but **not** epinephrine) should be given cautiously, if indicated. In the event of reduced myocardial function, consider recourse to treatment with dopamine or dobutamine by I.V. drip.

Since it has been reported that physostigmine may cause severe bradycardia, asystole and seizures, its use is not recommended in cases of overdosage with ANAFRANIL.

For management of a suspected drug overdose, contact your regional poison control centre.

6 DOSAGE FORMS, STRENGTHS, COMPOSITION AND PACKAGING

Table 1 – Dosage Forms, Strengths, Composition and Packaging

Route of Administration	Dosage Form / Strength/Composition	Non-medicinal Ingredients
Oral	Tablet 10 mg	Carnauba wax, colloidal silicon dioxide, croscarmellose sodium, hydroxypropyl methylcellulose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, titanium dioxide and yellow ferric oxide.
Oral	Tablet 25 mg	Carnauba wax, colloidal silicon dioxide, croscarmellose sodium, hydroxypropyl methylcellulose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, titanium dioxide and yellow ferric oxide.
Oral	Tablet 50 mg	Carnauba wax, colloidal silicon dioxide, croscarmellose sodium, hydroxypropyl methylcellulose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, and titanium dioxide.

ANAFRANIL (clomipramine hydrochloride) 10 mg Tablets

Pale-yellow, triangular, biconvex, film-coated tablet. Engraved 10 on one side and CP on the other side.

ANAFRANIL (clomipramine hydrochloride) 25 mg Tablets

Pale-yellow, round, biconvex, film-coated tablet. Engraved 25 on one side and CP on the other side.

ANAFRANIL (clomipramine hydrochloride) 50 mg Tablets

White, round, biconvex, film-coated tablet. Engraved CP over 50 on one side, other side plain. All strengths are available in bottles of 100 tablets.

7 WARNINGS AND PRECAUTIONS

Please see [3 SERIOUS WARNINGS AND PRECAUTIONS BOX](#).

Cardiovascular

ANAFRANIL is contraindicated in the acute recovery period following myocardial infarction or in cases of acute congestive heart failure.

Extreme caution should be used when ANAFRANIL is given to patients with:

- thyroid disease or those taking thyroid medication, because of the possibility of cardiovascular toxicity, including arrhythmias;
- cardiovascular disease.

Tricyclic antidepressants (TCAs), particularly in high doses, have been reported to produce sinus tachycardia, changes in conduction time, arrhythmias and severe hypotension. Cardiac arrhythmias and severe hypotension may also occur at normal doses in patients with pre-existing cardiovascular disease. ANAFRANIL also has a hypotensive action which may be detrimental in these circumstances. In such cases, treatment should be initiated at low doses with progressive increases only if required and tolerated, and the patients should be under close surveillance at all dosage levels. Monitoring of cardiac function and the electrocardiogram (ECG) is indicated in such patients as well as in the elderly (see [7 WARNINGS AND PRECAUTIONS, Monitoring and Laboratory test](#)).

A few instances of unexpected death have been reported in patients with cardiovascular disorders. Myocardial infarction and stroke have also been reported with drugs of this class (see [8.2 Clinical Trial Adverse Reactions, Cardiac disorders](#)). Therefore, ANAFRANIL should be administered with extreme caution to patients with a history of cardiovascular disorders, especially those with cardiovascular insufficiency, conduction disorders (e.g., atrioventricular block grades I to III) or other arrhythmias, those with circulatory lability and elderly patients (see [2 CONTRAINDICATIONS](#)).

There may be a risk of QTc prolongation at supra-therapeutic doses or supra-therapeutic plasma concentrations of clomipramine, as occur in the case of co-medication with selective serotonin reuptake inhibitors (SSRIs). It is established that hypokalemia is a risk-factor of QTc prolongation and Torsades de pointes. Therefore, hypokalemia should be treated before initiating treatment with ANAFRANIL and ANAFRANIL should be used with caution when combined with SSRIs or diuretics.

ECG abnormalities have been observed in patients treated with ANAFRANIL. The most common ECG changes were premature ventricular contractions (PVCs), ST-T wave changes, and abnormalities in intraventricular conduction. These changes were rarely associated with significant clinical symptoms. Nevertheless, caution is necessary when treating patients with heart disease, as well as elderly subjects. In these patients, cardiac function should be monitored and ECG examinations performed during long-term therapy. Gradual dose titration is also recommended.

Dental Effects

Lengthy treatment with TCAs can lead to an increased incidence of dental caries.

Dependence/Tolerance

Withdrawal Symptoms: A variety of withdrawal symptoms have been reported in association with abrupt discontinuation of ANAFRANIL, including dizziness, nausea, vomiting, headache, malaise, sleep disturbance, hyperthermia and irritability. In addition, such patients may experience a worsening of psychiatric status. While the withdrawal effects of ANAFRANIL have not been systematically evaluated in controlled trials, they are well known with closely related TCAs. It is recommended that the dosage be tapered gradually and the patient monitored carefully during discontinuation.

Driving and Operating Machinery

Since ANAFRANIL may produce sedation, particularly during the initial phase of therapy, patients should be cautioned about the danger of engaging in activities requiring mental alertness, judgement and physical coordination.

Endocrine and Metabolism

Caution should be observed in prescribing ANAFRANIL for hyperthyroid patients or for patients receiving thyroid medication (see [3 SERIOUS WARNINGS AND PRECAUTIONS BOX, Endocrine and Metabolism](#)). Transient cardiac arrhythmias have occurred in rare instances in patients who have been receiving other tricyclic compounds concomitantly with thyroid medication.

As with certain other psychotherapeutic drugs, ANAFRANIL elevates prolactin levels. Tissue culture experiments indicate that approximately one-third of human breast cancers are prolactin dependent *in vitro*, a factor of potential importance if the prescription of ANAFRANIL is contemplated in a patient with a previously detected breast cancer. Although disturbances such as galactorrhea, amenorrhea, gynecomastia, and impotence have been reported, the clinical significance of elevated serum prolactin levels is unknown for most patients. An increase in mammary neoplasms has been found in rodents after chronic administration of neuroleptic drugs. Neither clinical studies nor epidemiologic studies conducted to date, however, have shown an association between chronic administration of these drugs and mammary tumorigenesis; the available evidence is considered too limited to be conclusive at this time.

TCAs have been associated with porphyrinogenicity in susceptible patients.

Hyponatremia: Hyponatremia has occurred as a result of treatment with clomipramine (see [8.5 Post-Market Adverse Reactions, Metabolism and Nutrition Disorders](#)). In many cases, hyponatremia appears to be the result of the syndrome of inappropriate antidiuretic hormone secretion (SIADH). Elderly patients may be at greater risk of developing hyponatremia with a serotonergic antidepressant. Also, patients taking diuretics or who are otherwise volume-depleted can be at greater risk. Discontinuation of ANAFRANIL in patients with symptomatic hyponatremia and appropriate medical intervention should be instituted. Signs and symptoms of hyponatremia include headache, difficulty concentrating, memory impairment, confusion, weakness, and unsteadiness, which can lead to falls. More severe and/or acute cases have included hallucination, syncope, seizure, coma, respiratory arrest, and death.

Lactose: ANAFRANIL contains lactose monohydrate. Patients with hereditary problems of galactose intolerance, the Lapp lactase deficiency or glucose-galactose malabsorption should

not take this product.

Gastrointestinal

The more common adverse reactions of ANAFRANIL involve anticholinergic effects, including effects on the gastrointestinal system, such as dry mouth, and constipation.

TCAs may give rise to paralytic ileus, particularly in the elderly and in hospitalized patients. Therefore, appropriate measures should be taken if constipation occurs.

Hematologic

Isolated cases of bone marrow depression with agranulocytosis have been reported. Leukocyte and differential blood cell counts are recommended in patients receiving treatment with ANAFRANIL over prolonged periods, and should be performed for patients who develop fever, an influenzal infection, or sore throat. In the event of an allergic skin reaction, ANAFRANIL should be withdrawn.

Because ANAFRANIL is highly bound to serum protein, the administration of ANAFRANIL to patients taking other drugs that are highly bound to protein (e.g., warfarin, digoxin) may cause an increase in plasma concentrations of these drugs, potentially resulting in adverse effects (see [9.4 Drug-Drug Interactions](#)). Conversely, adverse effects may result from displacement of protein bound ANAFRANIL by other highly bound drugs. Therefore, caution is advised in patients with a history of a bleeding disorder.

Hepatic/Biliary/Pancreatic

ANAFRANIL is contraindicated in patients with existing liver damage (see [2 CONTRAINDICATIONS](#)).

ANAFRANIL has occasionally been associated with elevations in SGOT (AST) and SGPT (ALT) of potential clinical significance (e.g., values greater than 3 times the upper limit of normal).

In the majority of cases, these enzyme elevations were not associated with other clinical findings suggestive of hepatic injury.

Isolated cases of obstructive jaundice have been reported. Caution is indicated in treating patients with known liver disease and periodic monitoring of hepatic function is recommended in such patients.

Immune

In the event of an allergic skin reaction, ANAFRANIL should be withdrawn.

Monitoring and Laboratory Tests

Before initiating treatment, it is advisable to check the patient's blood pressure, because individuals with hypotension or a labile circulation may react to the drug with a fall in blood pressure. Regular measurements of blood pressure should be performed in susceptible patients. Postural hypotension may be controlled by reducing the dosage or administering circulatory stimulants.

Cardiac function, including blood pressure and ECG, should be periodically monitored before and during treatment with ANAFRANIL, particularly in patients with a history of cardiovascular disorders (see [7 WARNINGS AND PRECAUTIONS, Cardiovascular](#)).

The blood count should be monitored during treatment with ANAFRANIL (especially if the patient develops fever, sore throat or other symptoms which are associated with influenza infection), since isolated cases of agranulocytosis have been associated with the use of TCAs (see [7 WARNINGS AND PRECAUTIONS, Hematologic](#)). This is particularly relevant during the first few months of therapy and during prolonged treatment.

In patients with hepatic and renal disease or a history of liver disease, periodic monitoring of the hepatic enzyme levels and renal function is recommended (see [7 WARNINGS AND PRECAUTIONS, Hepatic/Biliary/Pancreatic](#); [Renal](#); and [8.2 Clinical Trial Adverse Reactions](#)).

Musculoskeletal

Epidemiological studies, mainly conducted in patients 50 years of age and older, show an increased risk of bone fractures in patients receiving SSRIs and TCAs. The mechanism leading to this risk is unknown. Elderly patients may be at greater risk of developing hyponatremia with a serotonergic antidepressant, which is associated with weakness, and unsteadiness, and can lead to falls (see [7 WARNINGS AND PRECAUTIONS, Hyponatremia](#)).

Neurologic

Seizures: Tricyclic agents are known to lower the convulsive threshold and ANAFRANIL (clomipramine hydrochloride) should, therefore, be used with extreme caution in patients with a history of convulsive disorders and other predisposing factors, e.g., brain damage of varying etiology, concomitant use of neuroleptics, alcoholism and withdrawal from alcohol, and concomitant use with other drugs that lower the seizure threshold. It appears that the occurrence of seizures is dose dependent. Therefore, the recommended total daily doses should not be exceeded (see [4.2 Recommended Dose and Dosage Adjustment](#)).

Electroconvulsive Therapy (ECT): Concurrent administration of electroconvulsive therapy and ANAFRANIL may be hazardous and such treatment should be limited to patients for whom it is essential. Healthcare professionals should discuss with patients the risk of taking ANAFRANIL while engaging in activities in which a sudden loss of consciousness could result in serious injury to the patient or others e.g., the operation of complex machinery, driving, swimming, or climbing.

Central Nervous System: Hyperthermia cases occurred when ANAFRANIL was used in combination with other drugs. Cases were considered to be neuroleptic malignant syndrome when ANAFRANIL and a neuroleptic were used concomitantly. (see [8.5 Post-market Adverse Reactions, Central Nervous System](#)).

Serotonin toxicity / Serotonin syndrome: Serotonin toxicity, also known as serotonin syndrome, is a potentially life-threatening condition and has been reported with TCAs including ANFRANIL, particularly during combined use with other serotonergic drugs (see [9 DRUG INTERACTIONS](#)).

Serotonin toxicity is characterised by neuromuscular excitation, autonomic stimulation (e.g.

tachycardia, flushing) and altered mental state (e.g. anxiety, agitation, hypomania). In accordance with the Hunter Criteria, serotonin toxicity diagnosis is likely when, in the presence of at least one serotonergic agent, one of the following is observed:

- Spontaneous clonus
- Inducible clonus or ocular clonus with agitation or diaphoresis
- Tremor and hyperreflexia
- Hypertonia and body temperature >38°C and ocular clonus or inducible clonus

If concomitant treatment with ANAFRANIL and other serotonergic agents, is clinically warranted, careful observation of the patient is advised, particularly during treatment initiation and dose increases- (see [2 CONTRAINDICATIONS](#), [4.1 Dosing Considerations](#), and [9.4 Drug-Drug Interactions](#)). If serotonin toxicity is suspected, discontinuation of the serotonergic agents should be considered.

Concomitant Use with Monoamine Oxidase Inhibitors (MAOI) including Linezolid and Intravenous Methylene Blue:

The concomitant use of ANAFRANIL with MAOIs intended to treat psychiatric disorders is contraindicated. ANAFRANIL should also not be started in a patient who is being treated with MAOIs such as linezolid or intravenous methylene blue (see [9.1 Serious Drug Interactions](#)). All reports with methylene blue that provided information on the route of administration involved intravenous administration in the dose range of 1 mg/kg to 8 mg/kg. No reports involved the administration of methylene blue by other routes (such as oral tablets or local tissue injection) or at lower doses. There may be circumstances when it is necessary to initiate treatment with a MAOI such as linezolid or intravenous methylene blue in a patient taking ANAFRANIL. ANAFRANIL should be discontinued before initiating treatment with the MAOI (see [2 CONTRAINDICATIONS](#) and [4.1 Dosing Considerations](#)).

Ophthalmologic

Angle-closure Glaucoma: As with other antidepressants, ANAFRANIL can cause mydriasis, which may trigger an angle closure attack in a patient with anatomically narrow ocular angles (see [8.2 Clinical Trial Adverse Reactions](#)). Patients should be examined to determine whether they are susceptible to angle-closure and be informed to seek immediate medical assistance if they experience eye pain, changes in vision or swelling or redness in or around the eye.

Anticholinergic effects: Because of its anticholinergic properties, ANAFRANIL should be used with caution in patients with increased intraocular pressure, narrow angle glaucoma or disturbances of visual accommodation (see [3 SERIOUS WARNINGS AND PRECAUTIONS BOX](#)).

Lacrimation: Decreased lacrimation and accumulation of mucoid secretions, due to the anticholinergic properties of TCAs, may cause damage to the corneal epithelium in patients with contact lenses.

Peri-Operative Considerations

ANAFRANIL is contraindicated in the acute recovery period following myocardial infarction (see

[2 CONTRAINDICATIONS](#)). ANAFRANIL should be discontinued as soon as possible prior to elective surgery because of possible cardiovascular effects. Hypertensive episodes have been observed during surgery in patients taking ANAFRANIL (see [7 WARNINGS AND PRECAUTIONS, Cardiovascular](#)).

Psychiatric

Clinical Worsening and Suicide Risk:

It is important that ANAFRANIL be dispensed in the least possible quantities to depressed outpatients, since suicide has been accomplished with this class of drug.

Pediatrics, Placebo-Controlled Clinical Trial Data: Analyses of placebo-controlled clinical trial safety databases from SSRIs and other newer antidepressants suggest that use of these drugs in patients under the age of 18 may be associated with behavioural and emotional changes, including an increased risk of suicidal ideation and behaviour over that of placebo. ANAFRANIL is not indicated for use in pediatric patients under 10 years of age (see [1.1 Pediatrics](#)).

Adults and Pediatrics: Rigorous clinical monitoring for suicidal ideation or other indicators of potential for suicidal behaviour is advised in patients of all ages given any antidepressant drug. This includes monitoring for emotional and behavioural changes. Patients with major depressive disorder (MDD), both adult and pediatric, may experience worsening of their depression and/or the emergence of suicidal ideation and behavior (suicidality) or unusual changes in behavior, whether or not they are taking antidepressant medications, and this risk may persist until significant remission occurs. Suicide is a known risk of depression and certain other psychiatric disorders, and these disorders themselves are the strongest predictors of suicide. There has been a longstanding concern, however, that antidepressants may have a role in inducing worsening of depression and the emergence of suicidality in certain patients during the early phases of treatment.

The following symptoms, anxiety, agitation, panic attacks, insomnia, irritability, hostility, aggressiveness, impulsivity, akathisia (psychomotor restlessness), hypomania, and mania, have been reported in adult and pediatric patients being treated with antidepressants for major depressive disorder as well as for other indications, both psychiatric and nonpsychiatric. Although a causal link between the emergence of such symptoms and either the worsening of depression and/or the emergence of suicidal impulses has not been established, there is concern that such symptoms may represent precursors to emerging suicidality.

All patients being treated with antidepressants for any indication should be monitored appropriately and observed closely for clinical worsening, suicidality, and unusual changes in behavior, especially during the initial few months of a course of drug therapy, or at times of dose changes, either increases or decreases.

Consideration should be given to changing the therapeutic regimen, including possibly discontinuing the medication, in patients whose depression is persistently worse, or who are experiencing emergent suicidality or symptoms that might be precursors to worsening depression or suicidality, especially if these symptoms are severe, abrupt in onset, or were not part of the patient's presenting symptoms.

Patients, families and caregivers of patients being treated with antidepressants for major depressive disorder or other indications, both psychiatric and nonpsychiatric, should be alerted about the emergence of agitation, irritability, unusual changes in behavior, and the other symptoms described above, as well as the emergence of suicidality, and to report such symptoms immediately to healthcare professionals. Such monitoring should include daily observation by families and caregivers.

Screening Patients for Bipolar Disorder: A major depressive episode may be the initial presentation of bipolar disorder. Prior to initiating treatment with an antidepressant, patients with depressive symptoms should be adequately screened to determine if they are at risk for bipolar disorder; such screening should include a detailed psychiatric history, including a family history of suicide, bipolar disorder, and depression. It should be noted that ANAFRANIL is not approved for use in treating bipolar depression.

Psychosis, Confusion, and Other Neuropsychiatric Phenomena: Patients treated with ANAFRANIL have been reported to show a variety of neuropsychiatric signs and symptoms including delusions, hallucinations, psychotic episodes, confusion, and paranoia. Because of the uncontrolled nature of many of the studies, it is impossible to provide a precise estimate of the extent of risk imposed by treatment with ANAFRANIL. In patients treated with TCAs, activation of latent schizophrenia or aggravation of existing psychotic manifestations in schizophrenic patients may occur. Hyperactive or agitated patients may become over-stimulated. A reduction in dose or discontinuation of ANAFRANIL should be considered under these circumstances.

In predisposed and elderly patients, TCAs may, particularly at night, provoke pharmacogenic (delirious) psychoses that disappear within a few days of withdrawing the drug.

Mania/Hypomania: Patients with manic-depressive tendencies may experience hypomanic or manic shifts.

Renal

ANAFRANIL is contraindicated in patients with existing liver or kidney damage (see [2 CONTRAINDICATIONS](#)).

It is advisable to monitor renal function during long-term therapy with TCAs.

Caution is called for when employing ANAFRANIL in patients with tumors of the adrenal medulla (e.g., pheochromocytoma, neuroblastoma), in whom the drug may provoke hypertensive crisis.

Because of its anticholinergic properties, ANAFRANIL should be used with caution in patients with urinary retention.

Reproductive Health: Female and Male Potential

- **Teratogenic Risk**

No teratogenic effects were observed in rats and mice at doses up to 20 times the maximum daily human dose. Slight nonspecific fetotoxic effects were seen in the offspring of pregnant mice given doses 10 times the maximum daily human dose. Slight nonspecific

embryotoxicity was observed in rats given doses 5 to 10 times the maximum daily human dose.

Skin

DRESS: Rare cases of drug rash with eosinophilia and systemic symptoms (DRESS), a potentially life-threatening adverse drug reaction, have been reported with the use of clomipramine. Symptoms include skin rashes or eruption, fever, eosinophilia, atypical lymphocytosis, swollen lymph nodes, inflammation of internal organs. In the event of severe acute reactions such as DRESS, discontinue clomipramine therapy immediately and institute appropriate treatment.

7.1 Special Populations

7.1.1 Pregnant Women

The safety of use in pregnant women has not been established. Therefore, ANAFRANIL should not be administered to women of childbearing potential, or during pregnancy, unless, in the opinion of the healthcare professionals, the expected benefit to the patient outweighs the potential risk to the fetus. Withdrawal symptoms including tremors, dyspnea, lethargy, colic, irritability, hypotonia/ hypertonia, convulsions and respiratory depression have been reported in neonates whose mothers received TCAs during the third trimester of pregnancy. To avoid such symptoms, ANAFRANIL should, if possible, be gradually withdrawn at least 7 weeks before the calculated date of confinement.

7.1.2 Breast-feeding

Since clomipramine passes into breast milk, ANAFRANIL should be gradually withdrawn or the infant weaned if the patient is breast-feeding.

7.1.3 Pediatrics

Depression

Pediatrics (<18 years): No data are available to Health Canada; therefore, Health Canada has not authorized an indication for pediatric use in children under 10 years of age.

Obsessive Compulsive Disorder

Pediatrics (<10 years): No data are available to Health Canada; therefore, Health Canada has not authorized an indication for pediatric use in children under 10 years of age.

Pediatrics (10 to 17 years): Limited information is available for the use of ANAFRANIL in children aged 10 to 17 years, therefore, ANAFRANIL is not recommended for use in this population. The long-term effects of ANAFRANIL on childhood growth and development have not been determined. (see [4.2 Recommended Dose and Dosage Adjustment, Obsessive Compulsive Disorders, Children and Adolescents](#)).

7.1.4 Geriatrics

Evidence from clinical studies and experience suggests that use of ANAFRANIL in the geriatric population is associated with differences in safety or effectiveness. Lower dosages are recommended for elderly and debilitated patients (see [4.2 Recommended Dose and Dosage Adjustment, Depression, Elderly and Debilitated Patients](#) and [4.2 Recommended Dose and Dosage Adjustment, Obsessive Compulsive Disorder, Elderly and Debilitated Patients](#)). Cardiovascular side effects may be reflected in ECG changes, which are seen most frequently in elderly patients, as is postural hypotension. The elderly are more prone to confusional states (see [8.2 Clinical Trial Adverse Reactions, Psychiatric disorders](#)).

ANAFRANIL has been associated with cases of clinically significant hyponatremia. Elderly patients may be at greater risk for this adverse reaction (see [7 WARNINGS AND PRECAUTIONS, Endocrine and Metabolism, Hyponatremia](#)).

8 ADVERSE REACTIONS

8.1 Adverse Reactions Overview

The most commonly observed adverse events associated with the use of ANAFRANIL (clomipramine hydrochloride) and not seen at an equivalent incidence among placebo-treated patients were gastrointestinal complaints, including dry mouth, constipation, nausea, dyspepsia, and anorexia; nervous system complaints, including somnolence, tremor, dizziness, nervousness and myoclonus; genitourinary complaints including changed libido, ejaculatory failure, impotence and micturition disorder; and other miscellaneous complaints, including fatigue, sweating, increased appetite, weight gain, and visual changes.

If severe neurological or psychiatric reactions occur, ANAFRANIL should be withdrawn.

Elderly patients are particularly susceptible to anticholinergic, psychiatric, neurological and cardiovascular effects.

8.2 Clinical Trial Adverse Reactions

Clinical trials are conducted under very specific conditions. The adverse reaction rates observed in the clinical trials, therefore, may not reflect the rates observed in practice and should not be compared to the rates in the clinical trials of another drug. Adverse reaction information from clinical trials may be useful in identifying and approximating rates of adverse drug reactions in real-world use.

The following adverse reactions have also been reported with clomipramine or other tricyclic antidepressants (TCAs).

(Frequency estimates: Very common >10%; Common >1 to 10%; Rare >0.01 to 1%; Very rare <0.01%)

Blood and lymphatic system disorders

Very rare: agranulocytosis, eosinophilia, leukopenia, purpura and thrombocytopenia may occur as an idiosyncratic response. One case of pancytopenia has been reported.

Cardiac disorders

Very common: sinus tachycardia.

Common: arrhythmia, palpitation, syncope.

Very rare: congestive heart failure, myocardial infarction, heart block, asystole, stroke, disturbances in cardiac conduction (e.g., widening of QRS complex, PQ changes, bundle-branch block, prolonged QT interval, Torsade de points in hypokalemia).

Ear and labyrinth disorders

Very rare: tinnitus.

Endocrine disorders

Very rare: gynecomastia in the male, galactorrhea in the female, inappropriate antidiuretic hormone (SIADH) secretion syndrome, menstrual irregularity.

Eye disorders

Very common: disturbances of visual accommodation.

Very rare: mydriasis, glaucoma.

General disorders and administration site conditions

Very common: hot flushes.

Very rare: drug fever, weakness, hyperpyrexia.

Gastrointestinal disorders

Very common: dry mouth and rarely associated sublingual adenitis.

Common: vomiting, abdominal cramps.

Rare: diarrhea.

Very rare: bitter taste, stomatitis, epigastric distress, black tongue, dysphagia, increased salivation, paralytic ileus.

Hepatobiliary disorders

Very rare: hepatitis with or without jaundice, obstructive jaundice.

Immune system disorders

Very rare: edema (general or of face and tongue), allergic alveolitis (pneumonia) with or without eosinophilia, systemic anaphylactic/anaphylactoid reactions including hypotension.

Investigations

Very common: electrocardiogram (ECG) changes (including flattening or inversion of T wave, depressed S-T segments) in patients of normal cardiac status.

Rare: elevated transaminases.

Very rare: alterations in EEG patterns, elevation or depression of blood sugar levels, increase in prolactin levels, weight loss.

Nervous system disorders

Common: headache, paresthesia (numbness, tingling sensation, symptoms suggestive of peripheral neuropathy), muscle hypertonia, muscle weakness.

Rare: epileptic seizures.

Very rare: incoordination, ataxia, extrapyramidal symptoms, speech disorders.

Psychiatric disorders

Common: delirium, drowsiness, insomnia, confusional states with hallucinations (particularly in geriatric patients and patients suffering from Parkinson's disease), anxiety, agitation, restlessness, sleep disturbances, nightmares, aggravated depression, hypomania, mania, decrease in memory, feeling of unreality, depersonalization, disorientation.

Rare: activation of latent psychosis.

Very rare: aggressiveness.

Renal and urinary disorders

Common: dilation of the urinary tract.

Reproductive system and breast disorders

Very rare: breast enlargement and galactorrhea in the female, testicular swelling, menstrual irregularity.

Respiratory thoracic and mediastinal disorders

Common: yawning.

Very rare: bronchospasm, nasal congestion, allergic alveolitis (pneumonia) with or without eosinophilia.

Skin and subcutaneous tissue disorders

Common: skin rash, urticarial.

Very rare: petechiae, itching, photosensitization (avoid excessive exposure to sunlight), alopecia.

Vascular disorders

Very common: hypotension, particularly orthostatic hypotension with associated vertigo.

Very rare: hypertension, peripheral vasospastic reactions.

Withdrawal Symptoms

Abrupt cessation of treatment with TCAs after prolonged administration may occasionally produce nausea, vomiting, abdominal pain, diarrhea, insomnia, nervousness, anxiety, headache and malaise. These symptoms are not indicative of addiction.

8.5 Post-Market Adverse Reactions

The following adverse drug reactions have been reported during post-approval use of ANAFRANIL. Because this reaction is reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate frequency.

Central Nervous System: Cases of hyperthermia have been recorded by nondomestic post-marketing surveillance systems. Most cases occurred when ANAFRANIL was used in combination with other drugs. When ANAFRANIL and a neuroleptic were used concomitantly, the cases were sometimes considered to be examples of a neuroleptic malignant syndrome.

Endocrine Disorders – Syndrome of inappropriate antidiuretic hormone secretion (SIADH).

Eye Disorders – Angle-closure glaucoma.

Immune System Disorders – Drug Rash with Eosinophilia and Systemic Symptoms (DRESS).

Metabolism and Nutrition Disorders – Hyponatremia.

9 DRUG INTERACTIONS

9.1 Serious Drug Interactions

Serious Drug Interactions

- **Monoamine oxidase inhibitors (MAOIs)**

ANAFRANIL should not be administered for a period of at least 14 days after the discontinuation of treatment with MAO-inhibitors due to the potential for severe interactions (see [2 CONTRAINDICATIONS](#)). The same caution should also be observed when administering a MAO-inhibitor after previous treatment with ANAFRANIL.

- **Thyroid Medication**

Prescribe ANAFRANIL with extreme caution for hyperthyroid patients or for patients receiving thyroid medication. Transient cardiac arrhythmias have occurred in rare instances in patients who have been receiving other tricyclic compounds concomitantly with thyroid medication: (see [3 SERIOUS WARNINGS AND PRECAUTIONS BOX, Endocrine and Metabolism](#)).

- **Linezolid or intravenous methylene blue**

Do not start ANAFRANIL in a patient who is being treated with linezolid or intravenous methylene blue (see [2 CONTRAINDICATIONS](#); [4.1 Dosing Considerations](#); [7 WARNINGS AND PRECAUTIONS, Serotonin Syndrome](#)).

9.2 Drug Interactions Overview

Metabolism of clomipramine

The primary route of clomipramine metabolism is demethylation to form the active metabolite, N-desmethylclomipramine, which can be formed by several P450 enzymes, primarily CYP3A4, CYP2C19 and CYP1A2.

Drugs Metabolized by P450 2D6 – The biochemical activity of the drug metabolizing isozyme cytochrome P450 2D6 (debrisoquin hydroxylase) is reduced in a subset of the Caucasian population (about 7% to 10% of Caucasians are so-called “poor metabolizers”); reliable estimates of the prevalence of reduced P450 2D6 isozyme activity among Asian, African and other populations are not yet available. Poor metabolizers have higher than expected plasma concentrations of tricyclic antidepressants (TCAs) when given usual doses. Depending on the fraction of drug metabolized by P450 2D6, the increase in plasma concentration may be small, or quite large (8- fold increase in plasma AUC of the TCA). In addition, certain drugs inhibit the activity of this isozyme and make normal metabolizers resemble poor metabolizers. An individual who is stable on a given dose of TCA may become abruptly toxic when given one of these inhibiting drugs as concomitant therapy. The drugs that inhibit cytochrome P450 2D6 include some that are not metabolized by the enzyme (quinidine; cimetidine) and many that are substrates for P450 2D6 (many other antidepressants, phenothiazines, and the Type 1C antiarrhythmics propafenone and flecainide). While all the selective serotonin reuptake inhibitors (SSRIs), e.g., fluoxetine, sertraline, paroxetine, and fluvoxamine, inhibit P450 2D6, they may vary in the extent of inhibition. Fluvoxamine has also been shown to inhibit P450 1A2, an isoform also involved in TCA metabolism. The extent to which SSRI-TCA interactions may pose clinical problems will depend on the degree of inhibition and the pharmacokinetics of the SSRI involved. Nevertheless, caution is indicated in the co-administration of TCAs with any of the SSRIs and also in switching from one class to the other. Of particular importance, sufficient time must elapse before initiating TCA treatment in a patient being withdrawn from fluoxetine, given the long half-life of the parent and active metabolite (at least 5 weeks may be necessary). Concomitant use of agents in the tricyclic antidepressant class (which includes ANAFRANIL) with drugs that can inhibit cytochrome P450 2D6 may require lower doses than usually prescribed for either the tricyclic antidepressant agent or the other drug. Furthermore, whenever one of these drugs is withdrawn from co-therapy, an increased dose of tricyclic antidepressant agent may be required. It is desirable to monitor TCA plasma levels whenever an agent of the tricyclic antidepressant class including ANAFRANIL is going to be co-administered with another drug known to be an inhibitor of P450 2D6 (and/or P450 1A2).

Because ANAFRANIL is highly bound to serum protein, the administration of ANAFRANIL to patients taking other drugs that are highly bound to protein (e.g., warfarin, digoxin) may cause an increase in plasma concentrations of these drugs, potentially resulting in adverse effects. Conversely, adverse effects may result from displacement of protein-bound ANAFRANIL by other highly bound drugs.

9.3 Drug-Behavioural Interactions

Alcohol: Patients should be warned that, while taking ANAFRANIL, their responses to alcoholic beverages may be exaggerated.

Smoking: Known inducers of CYP1A2 (e.g. nicotine/components in cigarette smoke) decrease plasma concentrations of tricyclic drugs. In cigarette smokers, clomipramine steady-state plasma concentrations were decreased 2-fold compared to non-smokers (no change in N-desmethyldclomipramine).

9.4 Drug-Drug Interactions

The drugs listed in this table are based on either drug interaction case reports or studies, or potential interactions due to the expected magnitude and seriousness of the interaction (i.e., those identified as contraindicated).

Table 2 - Established or Potential Drug-Drug Interactions

Proper/Common name	Source of Evidence	Effect	Clinical comment
Anticholinergics or Neuroleptics	T	Hyperexcitation states, delirium glaucoma	When TCAs are given in combination with anticholinergics or neuroleptics with an anticholinergic action, hyperexcitation states or delirium may occur, as well as attacks of glaucoma.
Antihypertensives	T	↓antihypertensive effects	Since ANAFRANIL may diminish or abolish the antihypertensive effects of guanethidine, bethanidine, clonidine, reserpine, or alpha-methyldopa, patients requiring concomitant treatment for hypertension should be given antihypertensives of a different type (e.g., vasodilators, beta-blockers).

Proper/Common name	Source of Evidence	Effect	Clinical comment
Arrhythmic agents (quinidine type)	T		TCAs should not be employed in combination with anti-arrhythmic agents of the quinidine type (see 7 WARNINGS AND PRECAUTIONS, Cardiovascular).
Barbiturates, carbamazepine, phenytoin, and oral contraceptives	T	↓ANAFRANIL	Substances which activate the hepatic mono-oxygenase enzyme system (e.g., barbiturates, carbamazepine, phenytoin and oral contraceptives) may lower plasma concentrations of TCAs and so reduce their antidepressive effects. In addition, ANAFRANIL may increase plasma levels of phenytoin and carbamazepine, therefore, it may be necessary to adjust the dosage of these drugs.
Buspirone	T	↑Risk for serotonin toxicity	Concomitant use of TCAs, such as clomipramine hydrochloride and buspirone may increase the risk of serotonin toxicity, a potentially life-threatening condition. See 7 WARNINGS AND PRECAUTIONS, Neurologic, Serotonin toxicity / Serotonin Syndrome

Proper/Common name	Source of Evidence	Effect	Clinical comment
Cimetidine or Methylphenidate	C	↑ANAFRANIL	Caution should be exercised if ANAFRANIL is administered together with cimetidine or methylphenidate since these drugs have been shown to inhibit the metabolism of several TCAs. Clinically significant increases in plasma levels of ANAFRANIL may occur, necessitating a dosage reduction.
CNS depressants or Anticholinergic agents	T	↑ANAFRANIL	<p>Patients should be warned that, while taking ANAFRANIL, their responses to CNS depressants (e.g., barbiturates, benzodiazepines or general anesthetics) or anticholinergic agents (e.g., atropine, antihistamines, biperiden, levodopa) may be exaggerated.</p> <p>Close supervision and careful adjustment of dosage are required when ANAFRANIL is administered with anticholinergic drugs.</p>
Coumarin drugs	T	↑anticoagulant effect	TCAs may potentiate the anticoagulant effect of coumarin drugs by inhibiting hepatic metabolism of these drugs. Careful monitoring of plasma prothrombin is therefore advised.

Proper/Common name	Source of Evidence	Effect	Clinical comment
Diuretics	T	Hypokalemia	Comedication with diuretics may lead to hypokalemia, which should be treated prior to administration of ANAFRANIL.
Estrogens	T	↑clomipramine	If administered concomitantly with estrogens, the dose of clomipramine should be reduced since steroid hormones inhibit the metabolism of clomipramine.
Fluoxetine, fluvoxamine and other selective serotonin reuptake inhibitors (SSRIs)	T	<p>↑ANAFRANIL</p> <p>↑Risk for serotonin toxicity</p>	<p>Fluoxetine, fluvoxamine and other SSRIs may increase the activity and plasma concentrations of TCAs, such as ANAFRANIL, with corresponding adverse effects. Comedication with SSRIs may lead to additive effects on the serotonergic system.</p> <p>If concomitant treatment with ANAFRANIL and other serotonergic agents is clinically warranted, careful observation of the patient is advised, particularly during treatment initiation and dose increases- (see 7 WARNINGS AND PRECAUTIONS, Neurologic, Serotonin toxicity / Serotonin Syndrome).</p>

Proper/Common name	Source of Evidence	Effect	Clinical comment
General anesthetics	T		ANAFRANIL should be discontinued prior to elective surgery for as long as is clinically feasible, since little is known about the interaction with general anesthetics.
Lithium	T	↑Risk for serotonin toxicity	Concomitant use of TCAs, such as clomipramine hydrochloride, and lithium may increase the risk of serotonin toxicity, a potentially life-threatening condition.
Noradrenaline or adrenaline, amphetamine, sympathomimetics	T	↑cardiovascular effects	ANAFRANIL may potentiate the cardiovascular effects of noradrenaline or adrenaline, amphetamine, as well as nasal drops and local anesthetics containing sympathomimetics (e.g., isoprenaline, ephedrine, phenylephrine). Close supervision and careful adjustment of dosage are required when ANAFRANIL is administered with sympathomimetic drugs.

Proper/Common name	Source of Evidence	Effect	Clinical comment
Opioids (e.g. tramadol, fentanyl)	T	<p>↑Risk for serotonin toxicity</p> <p>↑ Risk for seizures and opioid toxicity with concomitant use of tramadol.</p>	<p>Concomitant use of TCAs, such as clomipramine hydrochloride and triptans may increase the risk of serotonin toxicity, a potentially life-threatening condition. See 7 WARNINGS AND PRECAUTIONS, Neurologic, Serotonin toxicity / Serotonin Syndrome</p> <p>Concomitant use of TCAs, such as clomipramine and tramadol also increases the risk for seizures. In addition, this combination can inhibit the metabolism of tramadol to the active metabolite and thereby increasing tramadol concentrations, potentially causing opioid toxicity.</p>
Phenobarbital	T	↑Phenobarbital	Administration of ANAFRANIL has been reported to increase the plasma levels of phenobarbital, if given concomitantly.

Proper/Common name	Source of Evidence	Effect	Clinical comment
Phenothiazines, butyrophenones (haloperidol), thioridazine and diazepam	T	↑ANAFRANIL	Concomitant treatment with neuroleptic agents (e.g., phenothiazines and butyrophenones) may result in increased plasma concentrations of ANAFRANIL a lowered convulsion threshold and seizures. Combination with thioridazine may produce cardiac arrhythmias. No such effects are known to occur in combination with diazepam but it might be necessary to lower the dosage of ANAFRANIL if administered concomitantly with alprazolam or disulfiram.
Triptans	T	↑Risk for serotonin toxicity	Concomitant use of TCAs, such as clomipramine hydrochloride, and triptans may increase the risk of serotonin toxicity, a potentially life-threatening condition. See 7 WARNINGS AND PRECAUTIONS, Neurologic, Serotonin toxicity / Serotonin Syndrome

Legend: C = Case Study; CT = Clinical Trial; T = Theoretical

Interactions resulting in decreased effect of ANAFRANIL

CYP3A and CYP2C inducers, such as rifampicin or anticonvulsants (e.g. barbiturates, carbamazepine, phenobarbital and phenytoin), may decrease clomipramine concentrations as concomitant administration of drugs known to induce cytochrome P450 enzymes, particularly CYP3A4, CYP2C19 may accelerate the metabolism and decrease the efficacy of ANAFRANIL.

9.5 Drug-Food Interactions

Concomitant administration of ANAFRANIL with grapefruit, grapefruit juice, or cranberry juice may increase the plasma concentrations of clomipramine. Caution is therefore required when prescribing ANAFRANIL to patients taking these products.

9.6 Drug-Herb Interactions

Concomitant administration of St. John's Wort may reduce the plasma levels of clomipramine. Caution is therefore required when prescribing ANAFRANIL to patients taking St. John's Wort.

9.7 Drug-Laboratory Test Interactions

Interactions with laboratory tests have not been established.

10 CLINICAL PHARMACOLOGY

10.1 Mechanism of Action

ANAFRANIL (clomipramine hydrochloride) is a tricyclic agent with both antidepressant and antiobsessional properties. Like other tricyclics, clomipramine inhibits norepinephrine and serotonin uptake into central nerve terminals, possibly by blocking the membrane-pump of neurons. Clomipramine thereby increases the concentration of transmitter monoamines at receptor sites. Clomipramine is presumed to influence depression and obsessive and compulsive behavior through its effects on serotonergic neurotransmission. The actual neurochemical mechanism is unknown, but clomipramine's capacity to inhibit serotonin reuptake is thought to be important. Clomipramine appears to have a mild sedative effect that may be helpful in alleviating the anxiety component often accompanying depression.

As with other tricyclic compounds, ANAFRANIL possesses anticholinergic properties which are responsible for certain side effects. It also has weak antihistamine and antiserotonergic properties, lowers the convulsive threshold, potentiates the effect of norepinephrine and other drugs acting on the central nervous system, has a quinidine-like effect on the heart and may impair cardiac conduction.

The action of ANAFRANIL on the human electroencephalogram is one of desynchronization. Clomipramine causes a persistent increase in the frequency of shifts into stage I sleep and produces marked reduction or suppression of rapid eye movement sleep (REM or paradoxical sleep). Partial recovery occurs within 3 to 4 weeks as does a rebound after drug withdrawal which appears to last approximately the same time. In normal human volunteers tricyclic antidepressants (TCAs) tend to produce a sedative effect accompanied by atropine-like symptoms and may produce some difficulty in concentrating and thinking.

10.2 Pharmacodynamics

The pharmacological properties of clomipramine are similar to those of other TCAs, the main differences being quantitative rather than qualitative. The pharmacological profile of clomipramine includes reversal of reserpine and tetrabenazine effects, slight depressant effects on the central nervous system as manifested by behavioral, motor, electrocortical and visceral activity, anticholinergic and antihistaminic effects, and potentiation of adrenergic and serotonergic functions.

Clomipramine has a weak anticholinergic action demonstrated in laboratory animals by attenuation of the effect of acetylcholine on blood pressure and electrical stimulation of the

vagus, and slight counteraction of pilocarpine-induced salivation. The ED₅₀ for inhibition of tremorine-induced tremor in the mouse was 3.3 mg/kg. At 50 mg/kg, there was only incomplete inhibition of cholinergic intoxication due to tremorine (25 mg/kg I.P.). Clomipramine also has an antihistaminic effect demonstrated by inhibition of the fall in blood pressure following histamine injection in the cat.

As with other tricyclic agents, clomipramine produces a depression of spontaneous motor activity in laboratory animals (ED₅₀ approximately 40 mg/kg I.P.). Clomipramine can also produce irritability and aggressiveness. Clomipramine was considerably less effective than amitriptyline in depressing locomotor activity and similar in effectiveness to imipramine. However, on the rotating cylinder and wire traction tests, clomipramine was almost inactive, while imipramine and amitriptyline were active at relatively high doses. As with imipramine, clomipramine demonstrated only slight cataleptic activity and potentiated slightly bulbocapnine catalepsy at 50 mg/kg. However, unlike imipramine and amitriptyline, it exhibited no anticataleptic activity in chlorpromazine-induced catalepsy at the same dosage. Clomipramine also exhibited antiserotonin action, but was about 2½ times less effective than chlorpromazine in protecting against serotonin contraction of the guinea pig ileum.

Clomipramine has a depressant effect on behavioral and electrocortical arousal. Unlike the neuroleptic agents, this effect is more pronounced on electrocortical activity than on behavior. Clomipramine is as active as amitriptyline in producing slow waves of high voltage in the EEG of rabbits and in blocking the reaction induced by stimulation of the mesencephalic reticular formation. In low doses (1.25 mg/kg), clomipramine increased the duration and amplitude of after-discharges evoked by stimulation of the amygdala and had no effect on the recruiting response induced by stimulation of the anteromedian thalamic nucleus. In cats, clomipramine was found to suppress 'fast sleep' with progressive recovery. Performance on several conditioned tests was not affected significantly by clomipramine, imipramine or amitriptyline at doses up to 20 mg/kg. At doses of 10 mg/kg, clomipramine and imipramine pressed responding during the acquisition period of a conditioned avoidance test. Clomipramine is significantly less effective in inhibiting aggressive behavior in fighting mice (ED₅₀ 28 mg/kg) than imipramine (ED₅₀ 10 mg/kg) or amitriptyline (ED₅₀ 8 mg/kg). Clomipramine exerts a partial protective action against electroshock and pentylenetetrazol-induced tonic seizures in the rat with no effect in doses up to 50 mg/kg against strychnine convulsions in the mouse. The drug appears to be devoid of analgesic activity and produced only a slight depression of respiration in the non-anesthetized rabbit.

Antiarrhythmic effects of clomipramine in the dog were similar, but of shorter duration than those resulting from the quinidine-like action also observed with imipramine. As with imipramine, low doses (below 3 mg/kg I.V.), caused myocardial stimulation; cardiac depression occurred at higher doses. Clomipramine and amitriptyline were more active than imipramine in increasing the duration of barbiturate sleep. Clomipramine also demonstrated anti-emetic activity in dogs at doses of 10 mg/kg S.C.

Clomipramine occupies an intermediate position in reversing reserpine and tetrabenazine-induced catalepsy and ptosis. The anticholinergic effect of the drug combined with the potentiation of catecholamines may account for counteraction of ptosis. As with other

tricyclics, it potentiates sympathetic functions. Clomipramine was found to potentiate amphetamine-induced hyperthermia at 10 mg/kg S.C. and to block amphetamine toxicity in crowded mice at 75 mg/kg. It also potentiates the effect of adrenaline and noradrenaline on the blood pressure and the nictitating membrane of the anesthetized cat and inhibits the pressure effect of tyramine.

In histochemical and biochemical studies, clomipramine appeared more potent than imipramine in blocking serotonin uptake and in preventing 4-methyl- μ -ethyl-metatyramine induced depletion of serotonin in rat brain. Clomipramine was also more effective than amphetamine hyperthermia at 10 mg/kg S.C. and to block amphetamine toxicity in crowded mice at 75 mg/kg. It also potentiates the effect of adrenaline and noradrenaline on the blood pressure and the nictitating membrane of the anesthetizes cat and inhibits the pressure effect of tyramine.

Clomipramine was also more effective than imipramine in potentiating the effects of serotonin, tryptophan and nialamide on the extensor hind limb reflex in rats. Its effect on noradrenergic neurons was less pronounced.

10.3 Pharmacokinetics

Absorption

ANAFRANIL is rapidly and completely absorbed after oral administration in humans. Peak plasma levels are usually reached two hours after dosage, but much individual variation occurs. The plasma half-life after a single oral dose is approximately 21 hours. After 28 days of oral administration to patients in a daily dosage of 75 mg, plasma concentrations of clomipramine ranged from 17 to 70 ng/mL, (mean = 35.7 ng/mL). The concentration of the active metabolite, desmethylclomipramine, was about twice as high.

Distribution

The binding of ANAFRANIL to serum proteins is very high at 96 to 97% and is practically concentration-independent within the therapeutic range. Clomipramine has a volume of distribution of approximately 12 L/kg.

Metabolism

Clomipramine is extensively metabolized in the body with hydroxylation, demethylation and N-oxidation being the quantitatively more important routes of metabolism.

Elimination

As expected, the metabolites of ANAFRANIL are quite similar to those of imipramine, all retaining the benzazepine structure. Two-thirds of ANAFRANIL is excreted as water-soluble conjugates in the urine and approximately one-third in the feces. After a 25 mg radiolabeled dose of clomipramine in 2 subjects, the urinary recoveries of clomipramine and desmethylclomipramine were about 2% and 0.5% of the total radioactivity, respectively.

Animal Pharmacokinetics

Clomipramine is rapidly absorbed after oral administration to rabbits and rats and distributes to several organs, particularly liver and lungs, without exceeding blood plasma concentrations of 0.1 %. While clomipramine and imipramine follow similar distribution patterns in the rat after oral administration, clomipramine remains in various organs of the rabbit longer than imipramine. In the rabbit, the pattern of the breakdown products of clomipramine differs from imipramine, clomipramine giving rise to fewer conjugated metabolites. Clomipramine and imipramine are both catabolized via demethylation, hydroxylation of the ring structure, N-oxidation and removal of the side chain. In rabbit urine, only about 2% of the amount administered was found (by chromatography) unchanged.

Special Populations and Conditions

- **Pediatrics (< 18 years of age):** Health Canada has not authorized an indication for use in depression in pediatric patients (see [1.1 Pediatrics](#)).
- **Pediatrics (<10 years):** Health Canada has not authorized an indication for pediatric use in obsessive compulsive disorder (OCD) in children under 10 years of age.
- **Pediatrics (10 to 17 years of age):** Limited information is available for the use of ANAFRANIL for OCD in children aged 10 to 17 years, therefore, ANAFRANIL is not recommended for use in this population. The long-term effects of ANAFRANIL on childhood growth and development have not been determined.
- **Geriatrics:** Owing to the lower clearance of clomipramine in plasma, elderly patients require lower doses of ANAFRANIL than patients in younger age groups.
- **Pregnancy and Breast-feeding:** Clomipramine passes into breast milk, ANAFRANIL should be gradually withdrawn or the infant weaned if the patient is breast-feeding.
- **Genetic Polymorphism:** TCAs including clomipramine are primarily metabolised by the hepatic cytochrome P450 isozymes CYP2D6 and CYP2C19, which are polymorphic in the population. Patients known to be poor CYP2D6 or CYP2C19 metabolisers may have higher plasma levels of clomipramine (see [9.2 Drug Interactions Overview](#)). Dosage adjustments should be considered (see [4.2 Recommended Dose and Dosage Adjustment, Concomitant use with inhibitors of cytochrome P450 enzymes; CYP3A and CYP2C inducers](#)).

11 STORAGE, STABILITY AND DISPOSAL

10 mg, 25 mg and 50 mg tablets:

Store at controlled room temperature between 15°C to 30°C in tight containers. Protect from light.

Keep out of reach and sight of children.

12 SPECIAL HANDLING INSTRUCTIONS

None


PART II: SCIENTIFIC INFORMATION

13 PHARMACEUTICAL INFORMATION

Drug Substance

Proper name:	Clomipramine hydrochloride
Chemical name:	3-chloro-5-[3-(dimethylamino)propyl]-10,11-dihydro-5H-dibenz[b,f]azepine monohydrochloride
Molecular formula and molecular mass:	C ₁₉ H ₂₃ ClN ₂ • HCl and 351.3 g/mol

Structural formula:



Physicochemical properties:	White to off-white crystalline powder freely soluble in water, methanol and methylene chloride, and insoluble in ethyl ether and hexane, with a pKa of ca. 9.5 and a melting point of ca. 191 to 194°C
-----------------------------	--

14 CLINICAL TRIALS

This information is not available for this drug product.

15 MICROBIOLOGY

No microbiological information is required for this drug product.

16 NON-CLINICAL TOXICOLOGY

General Toxicology:

Acute Toxicity

The acute toxicity of clomipramine administered by oral, subcutaneous, intravenous and intraperitoneal routes has been studied in the mouse, rat, guinea pig, rabbit, and dog. Equal numbers of male and female animals were used and, in the case of mouse and rat, the number of animals per dose was ten. Clomipramine was given orally in a gum arabic suspension and, by the other routes, in aqueous solution. The animals were observed for a period of 8 days. The

LD₅₀ for each route of administration was determined by the method of Litchfield-Wilcoxon. Toxic manifestations included drowsiness, ventrolateral position, respiratory disturbances, ataxia and tonic-clonic convulsions.

Table 3 - Acute LD₅₀ Values for Clomipramine

Species	Route	LD ₅₀ mg/kg
Mouse	P.O.	630
	S.C.	245
	I.V.	44
	I.P.	98
Rat	P.O.	1450
	S.C.	1000
	I.V.	26
	I.P.	102
Guinea Pig	P.O.	575
	I.V.	30
Rabbit	P.O.	700
	I.V.	17
Dog	I.V.	40

Chronic Toxicity

One-year toxicity studies were performed on rats and dogs.

Rat

Doses of 0, 12.5, 25, 50 and 100 mg/kg of clomipramine were administered daily to Sprague Dawley rats (thirty-five males and thirty-five females per group). There was an increase in spontaneous mortality in animals in the highest dosage group only. No clinical or pathological alterations were noticed, except that histological examination revealed disturbance of spermatogenesis in male rats at higher dosages.

Dog

Doses of 0, 12.5, 50 and 100 mg/kg of clomipramine were administered daily to pedigree Pembrokehire Corgi dogs (four males and four females per group). Spontaneous death occurred only in the highest dosage group. Clinical and pathological studies, autopsy findings and measurement of organ weight gave no indication of a toxic effect of clomipramine, except that testicular damage was again apparent at higher doses.

One dog in the high dose group (100 mg/kg/day) showed no evidence of any mature spermatozoa. Spermatogenesis in this animal did not appear to extend beyond the secondary spermatocyte or spermatid stage. The histological picture did not suggest immaturity. In 2 of the intermediate dose level animals (50 mg/kg/day) there was evidence of bilateral inhibition of spermatogenesis associated with atrophy of some of the cells of the seminiferous tubules. In one dog (50 mg/kg/day), there is a possibility of some reduction in cellularity of some of the seminiferous tubules, although mature cell forms were present in this animal. The testes of low dose (12.5 mg/kg/day) and control animals were within normal limits and active spermatogenesis with mature cell forms was seen.

A 29-day intramuscular toxicity study was also performed in Beagle dogs. The dogs received doses of 0, 1 or 2 mg/kg clomipramine (two males and two females per group). No significant clinical or pathological changes were observed.

Other Chronic Toxicity Studies

As with other tricyclic compounds, ANAFRANIL has been associated with changes in testicular and lung tissue in long-term animal toxicology studies. In 1 and 2 year studies in rats, a dose 4 times the maximum daily human dose was associated with phospholipidosis in the lungs and changes in the testes (atrophy, aspermatogenesis, and calcification). In a 1 year toxicity study in dogs, testicular atrophy was detected in animals receiving 10 times the maximum recommended daily human dose.

Genotoxicity:

No studies have been performed to evaluate the mutagenic potential of ANAFRANIL.

Carcinogenicity:

No animal studies have been performed to evaluate the carcinogenic potential of ANAFRANIL.

Reproductive and Developmental Toxicology:

Teratogenicity

Tests of the teratogenic effect of clomipramine were performed on Swiss White Mice, Wistar Rats, and a strain of New Zealand White Rabbits, known to be susceptible to the teratogenic effect of thalidomide.

At doses of 0, 15, 30 and 60 mg/kg/day in rabbits and 0, 12.5, 25, 50 and 100 mg/kg/day in rats and mice, there was no evidence to suggest that clomipramine produced fetal abnormality. Doses of 12 and 24 mg/kg/day administered to male rats for two months and to female rats for 2 weeks before mating caused only a reduction of male activity.

PATIENT MEDICATION INFORMATION

READ THIS FOR SAFE AND EFFECTIVE USE OF YOUR MEDICINE

PrANAFRANIL

Clomipramine Hydrochloride Tablets

Read this carefully before you start taking **ANAFRANIL** and each time you get a refill. This leaflet is a summary and will not tell you everything about this drug. Talk to your healthcare professional about your medical condition and treatment and ask if there is any new information about **ANAFRANIL**.

Serious Warnings and Precautions

Heart Problems: Serious, and sometimes fatal, heart problems have been reported in patients taking tricyclic antidepressants, like ANAFRANIL. ANAFRANIL can cause a rapid heartbeat, **heart rhythm problems**, **heart attack** or **stroke** that can lead to death. You should talk to your healthcare professional before you take ANAFRANIL if you have a history of heart problems, especially problems with your heart rhythm, or if you are elderly. See the [Serious side effects and what to do about them](#) table, below, for more information on these and other serious side effects.

Anticholinergic Effects: ANAFRANIL can have an effect on the way chemical signals are passed between cells. This can cause problems in certain areas of the body, including the eyes and urinary tract. If you have increased pressure in your eye, glaucoma, trouble passing urine or an enlarged prostate gland your healthcare professional will have to monitor you closely as ANAFRANIL can make these conditions worse.

Thyroid Problems: If you have problems with your thyroid gland (hyperthyroidism) or are taking thyroid medication your healthcare professional will need to monitor you closely. Heart rhythm problems have been seen in patients taking tricyclic antidepressants, like ANAFRANIL, together with thyroid medication.

Seizures: Tricyclic antidepressants, like ANAFRANIL, can make you more likely to have seizures or fits. This risk is higher in patients who have a history of seizures, have had brain damage, are taking other medicines, particularly antipsychotics (used to treat mental health problems) or are suffering from alcoholism or withdrawal from alcohol. Talk to your healthcare professional about your risk of having seizures while you are taking ANAFRANIL.

New or worsened emotional or behavioural problems:

- When you first start taking ANAFRANIL or when your dose is adjusted, you may feel worse instead of better. You may feel new or worsened feelings of agitation, hostility, anxiety, or impulsivity.
- During your treatment with ANAFRANIL, it is important that you and your healthcare professional talk regularly about how you are feeling. They will closely monitor you

for signs of new or worsened emotions or behaviours while you are taking ANAFRANIL.

- You may find it helpful to tell a relative or close friend that you are depressed. Ask them to read this leaflet. You might ask them to tell you if they:
 - think your depression is getting worse, or
 - are worried about changes in your behaviour.
- If your depression worsens or you experience changes in your behaviour, tell your healthcare professional **right away**. Do not stop taking your medicine as it takes time for ANAFRANIL to work.

Self-harm or Suicide:

- Antidepressants, such as ANAFRANIL, may increase the risk of suicidal thoughts and actions for some patients.
- If you have thoughts of harming or killing yourself at any time, tell your healthcare professional or go to a hospital right away. Close observation by a healthcare professional is necessary in this situation.

What is ANAFRANIL used for?

ANAFRANIL is not for use in children under 10 years of age for the treatment of OCD and not for use in children under 18 years of age for the treatment of depression.

ANAFRANIL is used in adults to treat:

- Depression (feeling sad, a change in appetite or weight, difficulty concentrating or sleeping, feeling tired, headaches, unexplained aches and pain).
- obsessions and compulsions in patients (aged 10 and older) with obsessive compulsive disorder (OCD) (recurrent and intrusive thoughts, feelings, ideas, or sensations; recurrent pattern of behaviour, or unwanted thoughts or actions).

How does ANAFRANIL work?

ANAFRANIL belongs to a group of medicines called tricyclic antidepressants. It works by increasing the levels of two naturally occurring chemicals within the brain, noradrenaline and serotonin. This helps relieve the symptoms of depression and obsessive compulsive disorder.

What are the ingredients in ANAFRANIL?

Medicinal ingredients: Clomipramine hydrochloride

Non-medicinal ingredients: Carnuba wax, colloidal silicon dioxide, croscarmellose sodium, ferric oxide yellow (10 mg and 25mg tablets only), hydroxypropyl methylcellulose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, and titanium dioxide.

ANAFRANIL comes in the following dosage forms:

Tablets: 10 mg, 25 mg and 50 mg

Do not use ANAFRANIL if:

- you are allergic to clomipramine or any of the non-medicinal ingredients in ANAFRANIL (see [What are the ingredients in ANAFRANIL?](#)).
- you are allergic to any other tricyclic antidepressants.
- you are taking, or have taken within the last 14 days medicines for depression called monoamine oxidase inhibitors (MAOIs) including linezolid (an antibiotic) or methylene blue (a dye injected into a vein during surgery, x-rays or other imaging procedures).
- you have recently had a heart attack.
- you are in heart failure.
- you have any serious liver or kidney problems.
- you have or have had a blood disorder.
- you have glaucoma (increased eye pressure).

To help avoid side effects and ensure proper use, talk to your healthcare professional before you take ANAFRANIL. Talk about any health conditions or problems you may have, including if you:

- suffer from epilepsy (fits) or seizures.
- have suffered brain damage.
- are going to have electroconvulsive therapy (ECT).
- are elderly.
- have been told you have a low level of potassium in your blood (hypokalemia). Your healthcare professional will need to treat this before you start taking ANAFRANIL.
- have an overactive thyroid gland or are taking thyroid medication.
- have low levels of sodium in your blood.
- have trouble passing urine.
- have an enlarged prostate gland.
- have paralytic ileus (blocked intestine).
- have a tumour (cancer) of the adrenal gland (such as pheochromocytoma or neuroblastoma).
- have schizophrenia or any other mental health problems.
- have low blood pressure or other problems with your blood circulation.
- have dental problems.
- wear contact lenses.
- have or have had breast cancer.
- are taking medicines for depression called selective-serotonin reuptake inhibitors (SSRIs).
- are taking diuretics or “water pills”, used to treat high blood pressure.
- have been told you have enzymes that do not work well (such as “CYP2D6 poor metabolizer” or “CYP2C19 poor metabolizer”).

- had a recent bone fracture or were told you have osteoporosis or risk factors for osteoporosis.
- are lactose intolerant or have one of the following rare hereditary diseases:
 - Galactose intolerance
 - Lapp lactase deficiency
 - Glucose-galactose malabsorption

Because lactose is a non-medicinal ingredient in ANAFRANIL.

Other warnings you should know about:

Withdrawal symptoms: Do not stop taking ANAFRANIL without talking to your healthcare professional. You may need to lower your dose gradually and careful monitoring by your healthcare professional is required. Stopping ANAFRANIL suddenly may cause withdrawal symptoms including dizziness, nausea, vomiting, headache, malaise (general discomfort), sleep disturbance, increased body temperature, irritability and changes in behavior.

Bone Fracture: Taking ANAFRANIL may increase your risk of breaking a bone if you are elderly, have osteoporosis, or have other major risk factors for breaking a bone. You should take extra care to avoid falls, especially if you get dizzy or have low blood pressure.

ANAFRANIL can cause serious side effects, including:

- **Angle-closure glaucoma:** ANAFRANIL can cause angle-closure glaucoma (sudden eye pain). Having your eyes examined before you take ANAFRANIL could help identify if you are at risk of having angle-closure glaucoma. Talk to your healthcare professional right away if you have:
 - eye pain;
 - changes in vision;
 - swelling or redness in or around the eye.
- **Serotonin toxicity (also known as Serotonin syndrome):** ANAFRANIL can cause serotonin toxicity, rare but potentially life-threatening conditions. It can cause serious changes in how your brain, muscles and digestive system work. You may develop serotonin toxicity if you take ANAFRANIL with certain anti-depressants or migraine medications

Symptoms of serotonin toxicity include:

- fever, sweating, shivering, diarrhea, nausea, vomiting;
- muscle shakes, jerks, twitches or stiffness, overactive reflexes, loss of coordination;
- fast heartbeat, changes in blood pressure;
- confusion, agitation, restlessness, hallucinations, mood changes, unconsciousness, and coma.

See the [Serious side effects and what to do about them table](#), below, for more information on

these and other serious side effects

Pregnancy and Breastfeeding:

- You should not take ANAFRANIL if you are pregnant or thinking of becoming pregnant.
- Babies born to mothers that took medicines similar to ANAFRANIL while they were pregnant have experienced withdrawal symptoms after birth. Get immediate medical help for your baby if you took ANAFRANIL while you were pregnant and if they have any of the following symptoms:
 - breathing problems, bluish skin
 - seizures or fits
 - body temperature changes
 - stiff or floppy muscles
 - jitteriness, irritability, lethargy
 - drowsiness
 - constant crying
- Do not breastfeed while you are taking ANAFRANIL. ANAFRANIL passes into breastmilk.

Blood tests and monitoring: ANAFRANIL can cause abnormal blood test results. Your healthcare professional will decide when to perform blood tests and will interpret the results. Your healthcare professional will also monitor your blood pressure and the health of your heart while you are taking ANAFRANIL.

Driving and using machines: ANAFRANIL can cause you to feel relaxed and sleepy, especially when you first start taking it. Give yourself time after taking ANAFRANIL to see how you feel before driving a vehicle or using machinery.

Surgery: If you have a planned surgery, talk to your healthcare professional as soon as possible. They may ask you to stop taking ANAFRANIL.

Episodes of mania: Some patients with manic-depressive illness may enter into a manic phase. This is characterized by profuse and rapidly changing ideas, exaggerated gaiety and excessive physical activity. In such cases, it is important to contact your healthcare professional who probably will change your medication.

Drug reaction with eosinophilia and systemic symptoms (DRESS): ANAFRANIL may cause a serious skin reaction that can lead to death and may affect one or more organs. DRESS is characterized by some or all of the following:

- fever
- severe rash
- peeling skin
- swelling of the face
- swollen lymph glands
- flu-like feeling

- yellow skin or eyes
- shortness of breath
- swelling of the legs
- dry cough
- chest pain or discomfort
- feeling thirsty
- urinating less often, less urine.

Tell your healthcare professional about all the medicines you take, including any drugs, vitamins, minerals, natural supplements or alternative medicines.

Serious Drug Interactions

- Do **not** take ANAFRANIL if you are taking a monoamine oxidase inhibitor (MAOI), or if you have taken one in the last 14 days as this can cause serious side effects.
- Do not start ANAFRANIL if you are taking monoamine oxidase inhibitors such as the antibiotic linezolid and the intravenous dye methylene blue.
- Taking ANAFRANIL and thyroid medication can cause heart rhythm problems.

The following may interact with ANAFRANIL:

- Alcohol.
- Medicines used to treat anxiety and help you sleep called barbiturates and benzodiazepines, such as diazepam, alprazolam.
- General anesthetics, used during surgery to put you to sleep.
- Atropine, used as eye drops to dilate the pupil.
- Antihistamines, used to treat allergies.
- Medicines used to treat Parkinson’s Disease, such as biperiden, levodopa.
- Medicines used to treat heart rhythm problems, such as quinidine, propafenone and flecainide.
- Medicines used to treat high blood pressure, such as guanethidine, bethanidine, clonidine, reserpine, alpha-methyldopa.
- Diuretics or “water pills”, used to treat high blood pressure.
- Noradrenaline, used to treat low blood pressure.
- Adrenaline, used to treat serious allergic reactions.
- Nasal drops, used to relieve nasal congestion, and local anesthetics, used to numb the skin, such as isoprenaline, ephedrine, phenylephrine.
- Other medicines used to treat depression called selective serotonin reuptake inhibitors (SSRIs), such as fluoxetine and fluvoxamine, sertraline, paroxetine.

- Cimetidine, used to treat stomach ulcers or heartburn.
- Medicines used to treat attention-deficit/hyperactivity disorder (ADHD), such as methylphenidate, amphetamine.
- Medicines used to prevent seizures, such as carbamazepine, phenytoin, phenobarbital.
- Nicotine, this includes if you smoke or are using nicotine replacement therapy.
- Medicines used to treat mental health problems, like schizophrenia, such as phenothiazines, butyrophenones (e.g., haloperidol), lithium.
- Disulfiram, used to help you stop drinking alcohol.
- Coumarin, and other similar medicines, used to thin the blood (e.g., warfarin).
- Digoxin, used to treat heart failure.
- Medicines that contain estrogens, such as birth control pills or hormone replacement therapy.
- Opioids such as morphine, tramadol, buprenorphine.
- St. John's Wort (*hypericum perforatum*) – a herbal remedy used for depression.
- Grapefruit juice.

How to take ANAFRANIL:

- Always take ANAFRANIL exactly as your healthcare professional has told you. Check with your healthcare professional if you are not sure.
- Swallow ANAFRANIL tablets whole with water. Do not break, chew or crush the tablets.
- ANAFRANIL can be taken with or without food. To reduce stomach upset, take ANAFRANIL with food.
- Do not stop taking ANAFRANIL or change your dose without talking to your healthcare professional.

Usual dose:

Depression

Adults:

The recommended starting dose is 25 mg daily. Your healthcare professional may increase your dose by 25 mg increments up to 150 mg to 200 mg daily over a period of several weeks depending on your condition.

Elderly:

Lower doses are recommended for elderly patients. The recommended starting dose is 20 mg to 30 mg daily in divided doses.

Obsessive Compulsive Disorder

Adults:

The recommended starting dose is 25 mg daily. Your healthcare professional may increase your dose by 25 mg increments up to 100 mg to 200 mg daily by the end of 2 weeks depending on your condition.

Children and Adolescents (10 to 17 years of age):

Limited information is available for the use of ANAFRANIL in children aged 10 to 17 years, therefore, ANAFRANIL is not recommended for use in this population. For this age group the recommended starting dose may be 25 mg daily. The dose may be increased by 25 mg increments, up to 100 mg to 150 mg per day or 3 mg/kg of their body weight, whichever is lower.

Elderly:

Lower doses are recommended for elderly patients. The recommended starting dose is 20 mg to 30 mg daily in divided doses.

Overdose:

You may have the following symptoms if you take more ANAFRANIL than you should:

- drowsiness,
- stupor (when you are unable to move but still conscious),
- irregular muscle contractions,
- vomiting,
- blue skin,
- restlessness,
- agitation,
- enhanced reflexes,
- muscle stiffness,
- twisting and writhing movements of the hands and feet, convulsions (fits).

Other signs include fever, abnormally dilated pupil, a decrease or absence of urine production, constipation, shortness of breath, changes in blood pressure, rapid or irregular heartbeat, changes in electrocardiogram (ECG), heart attack and coma.

If you think you, or a person you are caring for, have taken too much ANAFRANIL, contact a healthcare professional, hospital emergency department, or regional poison control centre immediately, even if there are no symptoms.

Missed Dose:

If you forget to take ANAFRANIL, take it as soon as you remember. If it is almost time for your next dose, skip the missed dose. Do not take two doses at the same time to make up for a missed dose.

What are possible side effects from using ANAFRANIL?

These are not all the possible side effects you may have when taking ANAFRANIL. If you experience any side effects not listed here, tell your healthcare professional.

Side effects may include:

- headache
- dry mouth
- constipation
- diarrhea
- nausea, vomiting
- indigestion
- stomach pain, stomach cramps
- change in appetite
- weight gain
- sleepiness, fatigue
- trouble sleeping, nightmares
- shaking
- muscle spasms, muscle weakness
- dizziness
- nervousness
- sweating
- hot flashes
- increased sensitivity of the skin to sunlight
- yawning
- vision changes, blurred vision
- change in libido
- inability to have or maintain an erection
- inability to ejaculate or delay in ejaculation
- hair loss

Serious side effects and what to do about them			
Symptom / effect	Talk to your healthcare professional		Stop taking drug and get immediate medical help
	Only if severe	In all cases	
VERY COMMON			
Low blood pressure: dizziness, fainting, lightheadedness. May occur with you go from lying down or sitting to standing up.	✓		
COMMON			
Peripheral neuropathy: numbness or tingling sensation in the hands or feet		✓	

Serious side effects and what to do about them			
Symptom / effect	Talk to your healthcare professional		Stop taking drug and get immediate medical help
	Only if severe	In all cases	
Changes in feelings and behaviors: confusion, hallucinations (seeing or hearing things that are not there), anxiety, agitation, restlessness, worsening of depression, thoughts of suicide, extremely elevated and excitable mood memory problems, feeling of unreality, changes in your perception of reality, disorientation, thoughts or speech that do not make sense			✓
Heart rhythm problems: irregular heartbeat, fast heartbeat, palpitations, shortness of breath, fainting, loss of consciousness			✓
Mania: elevated or irritated mood, decreased need for sleep, racing thoughts, uneasiness, excessive physical activity.		✓	
RARE			
Seizures or fits: uncontrollable shaking with or without loss of consciousness			✓
VERY RARE			
Movement disorders: continuous muscle spasms and contractions, rigid muscles, restlessness, slowness of movement, tremor, jerky or irregular movements			✓
Angle-closure Glaucoma: increased pressure in the eye, pupil dilation, blurred vision, eye pain		✓	
Heart attack: chest pain or discomfort, lightheadedness, shortness of breath, pain in the			✓

Serious side effects and what to do about them			
Symptom / effect	Talk to your healthcare professional		Stop taking drug and get immediate medical help
	Only if severe	In all cases	
jaw, neck or back, pain in the arm or shoulder, cold sweat, nausea, vomiting			
Stroke: sudden numbness or weakness in the face, arm or leg, especially on one side of the body, confusion, trouble speaking, vision problems, weakness, dizziness, loss of coordination			✓
Liver problems: abdominal pain, nausea, vomiting, loss of appetite, yellowing of skin and eyes, dark urine, pale stool		✓	
Bronchospasm: coughing, tightness in the chest, wheezing, shortness of breath, difficulty breathing			✓
Hormonal changes: breast enlargement in men, breast enlargement and abnormal milk production in women, testicular swelling, irregular menstrual periods		✓	
Allergic reaction: rash, hives, tiny purple, red, or brown spots on the skin, itching, swelling of the face, lips and tongue, trouble swallowing or breathing, fever			✓
Withdrawal symptoms: nausea, vomiting, abdominal pain, diarrhea, sleeplessness, nervousness, anxiety, headache, generally feeling unwell, increased body temperature, irritability, behavioural changes		✓	

Serious side effects and what to do about them			
Symptom / effect	Talk to your healthcare professional		Stop taking drug and get immediate medical help
	Only if severe	In all cases	
Bone marrow depression: easy bruising, bleeding, nose bleeds, bleeding gums, red spots on the skin, fever and chills, rash, extreme fatigue, pale skin and lips			✓
UNKNOWN FREQUENCY			
Hyponatremia (low level of sodium in your blood): loss of energy, tiredness, muscle weakness or cramps, seizures		✓	
Self-harm or suicide: thoughts or actions about hurting or killing yourself or other people			✓
Serotonin Toxicity: a reaction which may cause feelings of agitation or restlessness, flushing, muscle twitching, involuntary eye movements, heavy sweating, high body temperature (>38 °C), or rigid muscles.			✓
Syndrome of Inappropriate Antidiuretic Hormone Secretion (SIADH): concentrated urine (dark in colour), feel or are sick, muscle cramps, confusion and fits (seizures) which may be due to inappropriate secretion of ADH (antidiuretic hormone)			✓
Drug reaction with eosinophilia and systemic symptoms (DRESS) (a serious skin reaction that can lead to death, it that may affect one or more organs): fever, severe rash, peeling skin, swollen lymph glands, flu-like feeling, yellow skin or eyes, shortness of breath, dry cough, chest pain or			✓

Serious side effects and what to do about them			
Symptom / effect	Talk to your healthcare professional		Stop taking drug and get immediate medical help
	Only if severe	In all cases	
discomfort, feel thirsty, urinating less often, less urine.			

If you have a troublesome symptom or side effect that is not listed here or becomes bad enough to interfere with your daily activities, tell your healthcare professional.

Reporting Side Effects

You can report any suspected side effects associated with the use of health products to Health Canada by:

- Visiting the Web page on Adverse Reaction Reporting (<https://www.canada.ca/en/health-canada/services/drugs-health-products/medeffect-canada/adverse-reaction-reporting.html>) for information on how to report online, by mail or by fax; or
- Calling toll-free at 1-866-234-2345.

NOTE: Contact your health professional if you need information about how to manage your side effects. The Canada Vigilance Program does not provide medical advice.

Storage:

Store at room temperature between 15°C to 30°C in a tight container.

Protect from light.

Keep out of reach and sight of children.

If you want more information about ANAFRANIL:

- Talk to your healthcare professional.
- Find the full product monograph that is prepared for healthcare professionals and includes this Patient Medication Information by visiting the Health Canada website: (<https://www.canada.ca/en/health-canada/services/drugs-health-products/drug-products/drug-product-database.html>); the manufacturer's website (<http://www.apotex.ca/products>), or by calling 1-800-667-4708.

This leaflet was prepared by Apotex Inc. 150, Signet Drive, Toronto, Ontario, M9L 1T9.

Last Revised: MAY 02, 2023