

PRODUCT MONOGRAPH
INCLUDING PATIENT MEDICATION INFORMATION

 EVUSHELD™

tixagevimab and cilgavimab injection

solution, 100 mg/mL (tixagevimab) and 100 mg/mL (cilgavimab), intramuscular use

Anti-SARS-CoV-2 spike protein monoclonal antibodies

AstraZeneca Canada Inc.
1004 Middlegate Road
Mississauga, Ontario
L4Y 1M4
www.astrazeneca.ca

Date of Initial Authorization:
April 14, 2022

Date of Revision:
May 30, 2023

Submission Control No: 273952

EVUSHELD™ is a trademark of AstraZeneca UK Ltd., used under license by AstraZeneca Canada Inc.

© AstraZeneca Canada Inc. 2023

RECENT MAJOR LABEL CHANGES

1 INDICATIONS	11/2022
4 DOSAGE AND ADMINISTRATION, 4.2 Recommended Dose and Dosage Adjustment	11/2022
4 DOSAGE AND ADMINISTRATION, 4.4 Administration	11/2022
7 WARNINGS AND PRECAUTIONS, Sensitivity/Resistance	11/2022
7 WARNINGS AND PRECAUTIONS, Immune	03/2023

TABLE OF CONTENTS

Sections or subsections that are not applicable at the time of authorization are not listed.

RECENT MAJOR LABEL CHANGES	2
TABLE OF CONTENTS	2
PART I: HEALTH PROFESSIONAL INFORMATION	4
1 INDICATIONS	4
1.1 Pediatrics	4
1.2 Geriatrics	4
2 CONTRAINDICATIONS	5
4 DOSAGE AND ADMINISTRATION	5
4.2 Recommended Dose and Dosage Adjustment	5
4.4 Administration	6
5 OVERDOSAGE	7
6 DOSAGE FORMS, STRENGTHS, COMPOSITION AND PACKAGING	7
7 WARNINGS AND PRECAUTIONS	7
7.1 Special Populations	9
7.1.1 Pregnant Women	9
7.1.2 Breast-feeding	9
7.1.3 Pediatrics	9
7.1.4 Geriatrics	9
8 ADVERSE REACTIONS	9
8.1 Adverse Reaction Overview	9
8.2 Clinical Trial Adverse Reactions	10
8.2.1 Clinical Trials adverse reactions: Pediatrics	11
8.3 Less Common Clinical Trial Adverse Reactions	11
8.5 Post-Market Adverse Reactions	11
9 DRUG INTERACTIONS	12
9.2 Drug Interactions Overview	12
9.4 Drug-Drug Interactions	12
9.5 Drug-Food Interactions	12
9.6 Drug-Herb Interactions	12
9.7 Drug-Laboratory Test Interactions	12
10 CLINICAL PHARMACOLOGY	12
10.1 Mechanism of Action	12
10.2 Pharmacodynamics	13

10.3	Pharmacokinetics.....	13
11	STORAGE, STABILITY AND DISPOSAL.....	15
PART II: SCIENTIFIC INFORMATION		16
13	PHARMACEUTICAL INFORMATION.....	16
14	CLINICAL TRIALS.....	16
14.1	Clinical Trials by Indication.....	16
	Pre-Exposure Prophylaxis of COVID-19.....	16
	Treatment of mild to moderate COVID-19	19
14.4	Immunogenicity.....	20
15	MICROBIOLOGY.....	21
16	NON-CLINICAL TOXICOLOGY.....	26
PATIENT MEDICATION INFORMATION		27

PART I: HEALTH PROFESSIONAL INFORMATION

1 INDICATIONS

Pre-exposure Prophylaxis of Coronavirus Disease 2019 (COVID-19)

EVUSHELD (tixagevimab and cilgavimab) is indicated for the pre-exposure prophylaxis of coronavirus disease 2019 (COVID-19) in adults and adolescents (≥ 12 years of age weighing at least 40 kg), who have not had a known recent exposure to an individual infected with SARS-CoV-2 and:

- Who are immune compromised and unlikely to mount an adequate immune response to COVID-19 vaccination **or**
- For whom COVID-19 vaccination is not recommended.

Pre-exposure prophylaxis with EVUSHELD is not a substitute for vaccination in individuals for whom COVID-19 vaccination is recommended.

Circulating SARS-CoV-2 viral variants may be associated with resistance to monoclonal antibodies. Healthcare professionals should routinely review the Antiviral Resistance information in 15 MICROBIOLOGY, in conjunction with literature and local guidelines, for details regarding specific variants and resistance, which may be updated regularly.

Treatment of Mild to Moderate COVID-19

EVUSHELD is indicated for the treatment of mild to moderate COVID-19 in adults and adolescents (≥ 12 years of age weighing at least 40 kg) [see also 14.1 Clinical Trials by Indication, [TACKLE](#)].

Circulating SARS-CoV-2 viral variants may be associated with resistance to monoclonal antibodies. Healthcare professionals should routinely review the Antiviral Resistance information in 15 MICROBIOLOGY, in conjunction with literature and local guidelines, for details regarding specific variants and resistance, which may be updated regularly.

1.1 Pediatrics

Pediatrics (<18 years of age): The safety and efficacy of EVUSHELD in children <18 years of age has not been established. No data are available. See 4.2 Recommended Dose and Dosage Adjustment. However, the recommended dose (see 4.2 Recommended Dose and Dosage Adjustment) is expected to result in comparable serum exposures of tixagevimab and cilgavimab in adolescents ≥ 12 years of age and weighing at least 40 kg as observed in adults based on an allometric scaling approach (which accounted for the effect of body weight changes associated with age on clearance and volume of distribution). See 10.2 Pharmacodynamics and 10.3 Pharmacokinetics. Close observation in adolescents is highly recommended.

1.2 Geriatrics

Geriatrics (≥ 65 years of age): Of the participants in the pooled pharmacokinetic (PK) analysis, 21% (N=534) were ≥ 65 years of age and 4.2% (N=107) were ≥ 75 years of age. There is no clinically meaningful difference in the PK of tixagevimab and cilgavimab in geriatric subjects

(≥65 years) compared to younger individuals. See 10.2 Pharmacodynamics and 10.3 Pharmacokinetics.

2 CONTRAINDICATIONS

EVUSHELD is contraindicated in individuals who have a history of severe hypersensitivity reactions, including anaphylaxis, to this drug or to any ingredient in the formulation, including any non-medicinal ingredient, or component of the container. For a complete listing, see 6 DOSAGE FORMS, STRENGTHS, COMPOSITION AND PACKAGING.

4 DOSAGE AND ADMINISTRATION

4.2 Recommended Dose and Dosage Adjustment

Prophylaxis

The recommended dose is 600 mg of EVUSHELD, administered as two separate 3.0 mL, sequential, injections of:

- 300 mg of tixagevimab
- 300 mg of cilgavimab

For individuals who require repeat dosing for ongoing prevention of COVID-19, subsequent doses of 600 mg of EVUSHELD (300 mg of tixagevimab and 300 mg of cilgavimab) should be given once every 6 months.

The dose recommendations for prophylaxis are based on the totality of the available data including clinical pharmacology, pharmacokinetics (PK), antiviral activity, and clinical trial data (see 8 ADVERSE REACTIONS, 10.2 Pharmacodynamics, and 10.3 Pharmacokinetics). EVUSHELD has been studied for the prophylaxis of COVID-19 at the 300 mg dose. The clinical safety of 600 mg EVUSHELD for prophylaxis use is supported by safety data from the Phase III multicentre study TACKLE in patients with mild to moderate COVID-19 (see 8 ADVERSE REACTIONS).

Treatment

The recommended dose is 600 mg of EVUSHELD, administered as two separate 3.0 mL, sequential, injections of:

- 300 mg of tixagevimab
- 300 mg of cilgavimab

EVUSHELD should be given as soon as possible after a positive viral test for SARS-CoV-2 and within 7 days after the onset of symptoms (see 14.1 Clinical Trials by Indication, Treatment of mild to moderate COVID-19).

Pediatrics (<18 years of age): The safety and efficacy of EVUSHELD in children <18 years of age have not been established.

4.4 Administration

For intramuscular injection

Tixagevimab and cilgavimab must be administered by a qualified healthcare provider using aseptic technique.

Visually inspect the vials for particulate matter and discoloration. Both tixagevimab and cilgavimab are clear to opalescent, colourless to slightly yellow solutions. Discard the vials if the solution is cloudy, discoloured or visible particles are observed. Do not shake the vials.

Withdraw the appropriate amount of tixagevimab solution and the appropriate amount of cilgavimab solution into separate syringes. Discard unused portion in vials.

Administer tixagevimab and cilgavimab as two separate, sequential intramuscular injections at different injection sites, preferably one in each of the gluteal muscles.

Each carton of EVUSHELD contains two vials:

- tixagevimab solution for injection (dark grey vial cap)
- cilgavimab solution for injection (white vial cap)

Indication	EVUSHELD dose (tixagevimab and cilgavimab)	Antibody dose	Number of vials needed ^a	Volume to withdraw from vial
Pre-exposure prophylaxis of COVID-19	600 mg (2 cartons)	tixagevimab 300 mg	2 vials	3.0 mL
		cilgavimab 300 mg	2 vials	3.0 mL
Treatment of mild to moderate COVID-19	600 mg (2 cartons)	tixagevimab 300 mg	2 vials	3.0 mL
		cilgavimab 300 mg	2 vials	3.0 mL

^a Each vial contains an overfill to allow the withdrawal of 150 mg (1.5 mL)

The tixagevimab and cilgavimab solutions for injection are preservative-free and therefore, the prepared syringes should be administered immediately.

If immediate administration is not possible, and the prepared tixagevimab and cilgavimab syringes need to be stored, the total time from vial puncture to administration should not exceed 4 hours, either:

- in a refrigerator at 2°C to 8°C or
- at room temperature up to 25°C.

Any unused solution should be discarded.

5 OVERDOSAGE

For management of a suspected drug overdose, contact your regional poison control centre.

There is no specific treatment for overdose with EVUSHELD. Treatment of overdose should consist of general supportive measures, including monitoring of vital signs and observation of the clinical status of the patient.

In clinical trials, doses up to 600 mg intramuscular (300 mg each of tixagevimab and cilgavimab) and 3000 mg intravenously (1500 mg each of tixagevimab and cilgavimab) have been administered without dose-limiting toxicity.

6 DOSAGE FORMS, STRENGTHS, COMPOSITION AND PACKAGING

To help ensure the traceability of biologic products, health professionals should recognise the importance of recording both the brand name and the non-proprietary (active ingredient) name as well as other product-specific identifiers such as the Drug Identification Number (DIN) and the batch/lot number of the product supplied.

Table 1 – Dosage Forms, Strengths, and Composition

Route of Administration	Dosage Form / Strength / Composition	Non-medicinal Ingredients
Intramuscular use	Solution for injection <ul style="list-style-type: none">100 mg/mL / 150 mg of tixagevimab100 mg/mL / 150 mg of cilgavimab	L-Histidine L-Histidine hydrochloride monohydrate Polysorbate 80 Sucrose Water for injection

Each carton of EVUSHELD contains two vials:

- **Tixagevimab**
1.5 mL of solution for injection in a clear glass vial closed by chlorobutyl elastomeric stopper sealed with a dark-grey aluminium flip-off top.
- **Cilgavimab**
1.5 mL of solution for injection in a clear glass vial closed by chlorobutyl elastomeric stopper sealed with a white aluminium flip-off top.

7 WARNINGS AND PRECAUTIONS

Cardiovascular

Cardiac adverse events

In PROVENT, a higher proportion of subjects who received EVUSHELD versus placebo reported serious cardiovascular events (0.7% versus 0.3%) notably myocardial infarction and

cardiac failure. A smaller imbalance was observed for serious thromboembolic events (0.5% versus 0.2%). Most subjects had cardiac risk factors and/or a prior history of cardiovascular disease, and there was no clear temporal pattern. A causal relationship between EVUSHELD and these events has not been established.

Consider the risks and benefits prior to initiating EVUSHELD in individuals at high risk for cardiovascular or thromboembolic events, and advise individuals to seek immediate medical attention if they experience any signs or symptoms suggestive of a cardiovascular or thromboembolic event.

Hematologic

Clinically significant bleeding disorders

As with any other intramuscular injections, EVUSHELD should be given with caution to patients with thrombocytopenia or any coagulation disorder.

Immune

Hypersensitivity and Anaphylaxis

Serious hypersensitivity reactions, including anaphylaxis, have been reported following administration of EVUSHELD (see 8.5 Post-Market Adverse Reactions). If signs and symptoms of a clinically significant hypersensitivity reaction or anaphylaxis occur, immediately discontinue administration and initiate appropriate medicinal products and/or supportive therapy.

Reproductive Health: Female and Male Potential

- **Fertility**

There are no data on the effects of tixagevimab and cilgavimab on human fertility.

Sensitivity/Resistance

Potential Risk of Prophylaxis or Treatment Failure due to Antiviral Resistance

Circulating SARS-CoV-2 viral variants may be associated with resistance to monoclonal antibodies such as EVUSHELD. Healthcare professionals should routinely review the Antiviral Resistance information in 15 MICROBIOLOGY, in conjunction with literature and local guidelines, for details regarding specific variants and resistance, which may be updated regularly.

Patients who receive EVUSHELD prophylactically should be informed of the potential for breakthrough infections to occur. Patients should be instructed to promptly seek medical advice if signs or symptoms of COVID-19 occur (the most common symptoms include fever, cough, tiredness and loss of taste or smell; the most serious symptoms include difficulty breathing or shortness of breath, loss of speech or mobility, or confusion and chest pain).

Decisions regarding the use of EVUSHELD for the treatment of COVID-19 should take into consideration what is known about the characteristics of the circulating SARS-CoV-2 viral variants, including geographical prevalence.

7.1 Special Populations

7.1.1 Pregnant Women

There are insufficient data from the use of tixagevimab and cilgavimab in pregnant women. Non-clinical reproductive toxicity studies have not been performed with tixagevimab and cilgavimab. In a tissue cross reactivity study with tixagevimab and cilgavimab using human fetal tissues, no binding of clinical concern was detected. Human immunoglobulin G1 (IgG1) antibodies are known to cross the placental barrier; therefore, tixagevimab and cilgavimab have the potential to be transferred from the mother to the developing fetus.

EVUSHELD should only be used during pregnancy if the potential benefit outweighs the potential risk for the mother and the fetus.

7.1.2 Breast-feeding

There are no data available on whether tixagevimab and cilgavimab are excreted in human milk. Exposure to the breast-fed child cannot be excluded.

The developmental and health benefits of breast-feeding should be considered along with the mother's clinical need for EVUSHELD and any potential adverse effects on the breast-fed child from EVUSHELD or from the underlying maternal condition.

7.1.3 Pediatrics

Pediatrics (<18 years of age): The safety and efficacy of EVUSHELD in children <18 years of age has not been established. No data are available. However, the recommended adult dose is expected to result in comparable serum exposures of tixagevimab and cilgavimab in children ≥ 12 years of age and weighing at least 40 kg. EVUSHELD is not authorized for use in pediatrics under 12 years of age or weighing less than 40 kg. See 10.2 Pharmacodynamics and 10.3 Pharmacokinetics.

7.1.4 Geriatrics

Geriatrics (≥ 65 years of age): Of the participants in the pooled PK analysis, 21% (N=534) were ≥ 65 years of age and 4.2% (N=107) were ≥ 75 years of age. There is no clinically meaningful difference in the PK of tixagevimab and cilgavimab in geriatric subjects (≥ 65 years) compared to younger individuals. See 10.2 Pharmacodynamics and 10.3 Pharmacokinetics.

8 ADVERSE REACTIONS

8.1 Adverse Reaction Overview

A total of 4210 adult participants received 300 mg of EVUSHELD (150 mg of tixagevimab and 150 mg of cilgavimab) via intramuscular injection, in the Phase III prophylaxis studies PROVENT, D8850C00002 (a double-blind, placebo-controlled clinical trial for the pre-exposure prophylaxis of COVID-19) and STORM CHASER, D8850C00003 (a double-blind, placebo-controlled clinical trial for the post-exposure prophylaxis of COVID-19, an indication for which EVUSHELD is not approved). The median duration for safety follow-up was 137 days for PROVENT and 121 days for STORM CHASER.

The most frequently reported adverse reaction in the pooled analysis of PROVENT and STORM CHASER was injection site reaction (1.3%).

In PROVENT, adverse events were reported in 1417 (41%) subjects receiving EVUSHELD and 698 (40%) receiving placebo. Serious adverse events were reported in 92 (3%) subjects receiving EVUSHELD and 42 (2%) receiving placebo. Of the participants with at least one adverse event, the majority were mild (23%) or moderate (14%) in intensity and balanced between treatment arms.

In STORM CHASER, adverse events were reported in 229 (31%) subjects receiving EVUSHELD and 150 (40%) receiving placebo. Serious Adverse Events (SAEs) were reported in 9 (1%) subjects receiving EVUSHELD and 7 (2%) receiving placebo. Of the participants with at least one adverse event, the majority were mild (EVUSHELD 19% versus placebo 25%) or moderate (EVUSHELD 10% versus placebo 12%) in intensity.

A total of 452 non-hospitalized adult patients with mild to moderate COVID-19 received 600 mg of EVUSHELD (300 mg of tixagevimab and 300 mg of cilgavimab) via intramuscular injection within ≤7 days of symptom onset, in the Phase III treatment study, TACKLE (D8851C00001). TACKLE is an ongoing Phase III, double-blind, placebo-controlled multicentre study for the treatment of adult patients with mild to moderate COVID-19. The median duration for safety follow-up was 170 days.

The most frequently reported adverse reaction in TACKLE was injection site reaction (2.4%).

In TACKLE, during a median follow-up of 170 days, adverse events were reported in 174 (38.5%) subjects receiving EVUSHELD and 196 (43.5%) receiving placebo.

During a median follow-up of 84 days, serious adverse events were reported in 33 (7.3%) subjects receiving EVUSHELD and 54 (12%) receiving placebo. Of the participants who received EVUSHELD and experienced at least one adverse event, the majority were mild (14.8%) or moderate (7.5%) in intensity and balanced between treatment arms. Four subjects reported cardiac SAEs. Acute myocardial infarction was reported for two subjects who received EVUSHELD (one of whom also experienced cardiac failure leading to death) and sudden cardiac death was reported for one subject who received EVUSHELD. One subject who received placebo reported arrhythmia. All subjects who experienced cardiac SAEs had cardiac risk factors and/or a prior history of cardiovascular disease at baseline. There were no reports of anaphylaxis or serious hypersensitivity reactions.

The overall safety profile in patients who received 600 mg IM EVUSHELD was generally similar to that reported in participants who received 300 mg IM EVUSHELD.

8.2 Clinical Trial Adverse Reactions

Clinical trials are conducted under very specific conditions. The adverse reaction rates observed in the clinical trials; therefore, may not reflect the rates observed in practice and should not be compared to the rates in the clinical trials of another drug. Adverse reaction information from clinical trials may be useful in identifying and approximating rates of adverse drug reactions in real-world use.

Adverse Reactions organized by MedDRA System Organ Class (SOC) is summarized in Table 2 below.

Table 2 – Adverse Reactions (Pooled PROVENT and STORM CHASER studies, Safety Analysis Set, and TACKLE)

MedDRA SOC	Preferred Term	EVUSHELD 300 mg (PROVENT and STORM CHASER)		EVUSHELD 600 mg (TACKLE)	
		EVUSHELD (N = 4210)	Placebo (N = 2108)	EVUSHELD (N = 452)	Placebo (N = 451)
Immune system disorders	Hypersensitivity ^a	1.0%	0.9%	0.4%	0.7%
Injury, poisoning and procedural complications	Injection site reaction ^a	1.3%	1.2%	2.4%	2.4%

^a Grouped terms: Hypersensitivity (including Rash and Urticaria); Injection site reaction (including Injection site pain, Injection site erythema, Injection site pruritus, Injection site reaction and Injection site induration).

Repeat dosing

There are limited safety data available with repeat dosing.

In an open-label sub-study, 305 participants who had received an initial dose of 300 mg EVUSHELD in PROVENT received a second dose of 300 mg EVUSHELD 10 to 14 months after administration of the initial dose. The median duration of follow-up after administration of the second dose was 17 days. The overall adverse event profile for participants who received a second EVUSHELD dose remained similar when compared to the initial dose.

8.2.1 Clinical Trials adverse reactions: Pediatrics

No data are available for pediatric individuals <18 years old. See 4.2 Recommended Dose and Dosage Adjustment and 10.3 Pharmacokinetics.

8.3 Less Common Clinical Trial Adverse Reactions

Less common clinical trial adverse reactions organized by MedDRA System Organ Class (SOC) are summarized below:

General disorders and administration site conditions: injection-related reaction

8.5 Post-Market Adverse Reactions

The following adverse reaction has been identified during post-marketing use of EVUSHELD. It is generally not possible to reliably determine the frequency because such reactions have been reported spontaneously from a population of uncertain size. The frequency of these adverse reactions is therefore 'not known' (cannot be estimated from available data).

Immune system disorders:

- Serious hypersensitivity including anaphylaxis (see Hypersensitivity and Anaphylaxis)

9 DRUG INTERACTIONS

9.2 Drug Interactions Overview

No interaction studies have been conducted with EVUSHELD.

EVUSHELD is not expected to undergo metabolism by hepatic enzymes or renal elimination. See 10.3 Pharmacokinetics.

An interaction with COVID-19 vaccinations has not been studied and can therefore not be excluded. Healthcare professionals should refer to local guidance regarding the timing of COVID-19 vaccination prior to or following administration of EVUSHELD.

9.4 Drug-Drug Interactions

Tixagevimab and cilgavimab are not renally excreted or metabolised by cytochrome P450 enzymes; therefore, interactions with concomitant medications that are renally excreted or that are substrates, inducers, or inhibitors of cytochrome P450 enzymes are unlikely.

Based on PK modelling, COVID-19 vaccination following EVUSHELD administration had no clinically relevant impact on the clearance of EVUSHELD.

In the absence of compatibility studies, EVUSHELD should not be mixed with other medicinal products.

9.5 Drug-Food Interactions

Interactions with food have not been established.

9.6 Drug-Herb Interactions

Interactions with herbal products have not been established.

9.7 Drug-Laboratory Test Interactions

Interactions with laboratory tests have not been established.

10 CLINICAL PHARMACOLOGY

10.1 Mechanism of Action

Tixagevimab and cilgavimab, two SARS-CoV-2 spike protein-directed attachment inhibitors, are two recombinant human IgG1 κ monoclonal antibodies, with amino acid substitutions to extend antibody half-life (YTE) and to reduce antibody effector function and potential risk of antibody-dependent enhancement of disease (TM). Tixagevimab and cilgavimab bind to non-overlapping regions of the spike protein receptor binding domain (RBD) of SARS-CoV-2. Tixagevimab, cilgavimab and their combination bind to the spike protein RBD with equilibrium dissociation constants of $K_D = 2.76$ pM, 13.0 pM and 13.7 pM, respectively, blocking its interaction with the human ACE2 receptor, the SARS-CoV-2 receptor, required for virus attachment. Tixagevimab,

cilgavimab and their combination blocked RBD binding to the human ACE2 receptor with IC₅₀ values of 0.32 nM (48 ng/mL), 0.53 nM (80 ng/mL) and 0.43 nM (65 ng/mL), respectively.

10.2 Pharmacodynamics

See 15 MICROBIOLOGY.

In a Phase I study, following a single 300 mg IM dose of EVUSHELD in healthy volunteers (N= 10) neutralizing antibodies geometric mean titers (GMT) were evaluated using an authentic virus neutralization assay (PRNT80) at 7, 30, 60, 90, 150, 210 and 270 days post-dose. GMT results were 689.2, 852.8, 656.8, 533.7, 290.1, 297.5 and 98.6 respectively.

In PROVENT, following a single 300 mg IM dose of EVUSHELD, neutralizing antibody GMT at 7, 28, 57, and 91 days post-dose were similar to those observed in the Phase I healthy volunteer study and were 16, 22, 17 and 12-fold higher, respectively, than the GMT measured in convalescent plasma from COVID-19 patients (GMT= 30.8).

In TACKLE, following a single 600 mg IM dose of EVUSHELD, neutralizing antibody GMT at Day 6, 15, 29, 85, and 169, were 16-fold, 14-fold, 22-fold, 18-fold, and 5-fold, respectively, over placebo.

10.3 Pharmacokinetics

The pharmacokinetics of tixagevimab and cilgavimab are comparable, linear, and dose-proportional between 300 to 600 mg following a single IM administration.

Absorption:

After a single 300 mg IM dose (150 mg each antibody) in healthy volunteers, the estimated absolute bioavailability was 68.5% for tixagevimab and 65.8% for cilgavimab.

After a single 600 mg IM treatment dose (300 mg each antibody) in COVID-19 participants from TACKLE, the mean (%CV) maximum concentration (C_{max}) was 21.9 (61.7%) and 20.3 (63.6%) µg/mL for tixagevimab and cilgavimab respectively, which were reached at a median T_{max} of 15 days.

Distribution:

Based on PK modelling, the central volume of distribution was 2.72 L for tixagevimab and 2.48 L for cilgavimab. The peripheral volume of distribution was 2.64 L for tixagevimab and 2.57 L for cilgavimab.

Metabolism:

Tixagevimab and cilgavimab are expected to be degraded into small peptides and component amino acids via catabolic pathways in the same manner as endogenous IgG antibodies.

Elimination:

The clearance (CL) was 0.041 L/day for tixagevimab and 0.041 L/day for cilgavimab with between subject variability of 21% and 29% respectively. The estimated population median terminal elimination half-life was 89 days for tixagevimab and 84 days for cilgavimab.

In PROVENT, following a single 300 mg IM dose of EVUSHELD, the geometric mean serum concentration was 23.4 µg/mL (geoSD: 1.9) on Day 29 and 12.2 µg/mL (geoSD: 1.4) on Day 183.

In the PROVENT sub-study, following a second IM dose of 300 mg EVUSHELD administered 10 to 14 months after the initial dose, the geometric mean serum concentration was 26.4 µg/mL (geoSD: 1.5) on Day 29. The PK of the total antibodies following a second dose of EVUSHELD are similar to the first dose.

In TACKLE, following a single 600 mg IM dose of EVUSHELD, the geometric mean serum concentration was 37.2 µg/mL (geoSD: 2.1) on Day 29. There was no clinically relevant difference in the clearance of tixagevimab or cilgavimab between participants with COVID-19 enrolled in TACKLE and those enrolled in the prophylaxis studies.

Special Populations and Conditions

- **Pediatrics (12 years of age and older weighing at least 40 kg):** The recommended dosing regimen is expected to result in comparable serum exposures of tixagevimab and cilgavimab in adolescents 12 years of age and older and weighing at least 40 kg as observed in adults, since adults with similar body weight have been included in Trial D8850C00002 (PROVENT) and D8851C00001 (TACKLE).
- **Geriatrics:** Of the participants in the pooled PK analysis, 21% (N= 534) were 65 years of age or older and 4.2% (N= 107) were 75 years of age or older. There is no clinically meaningful difference in the PK of tixagevimab and cilgavimab in geriatric subjects (≥65 years) compared to younger individuals.
- **Hepatic Insufficiency:** No specific studies have been conducted to examine the effects of hepatic impairment on the PK of tixagevimab and cilgavimab. The impact of hepatic impairment on the PK of tixagevimab and cilgavimab has not been established and it is unknown whether a dose adjustment is needed in these individuals.
- **Renal Insufficiency:** No specific studies have been conducted to examine the effects of renal impairment on the pharmacokinetics of tixagevimab and cilgavimab.

Tixagevimab and cilgavimab are not eliminated intact in the urine, since monoclonal antibodies with molecular weight >69 kDa do not undergo renal elimination, thus renal impairment is not expected to significantly affect the exposure of tixagevimab and cilgavimab. Similarly, dialysis is not expected to impact the PK of tixagevimab and cilgavimab.

Based on population PK analysis, there is no difference in the clearance of tixagevimab and cilgavimab in patients with mild (N= 978) or moderate (N= 174) renal impairment compared to patients with normal renal function. In the population PK model there were insufficient participants with severe renal impairment (N= 21) to draw conclusions.

- **Other Special Populations:** The PK profile of tixagevimab and cilgavimab were not affected by sex, age, race or ethnicity. Bodyweight had no clinically relevant effect on the PK of tixagevimab and cilgavimab in adults over the range of 36 kg to 177 kg.

11 STORAGE, STABILITY AND DISPOSAL

Store in a refrigerator (2°C to 8°C). See 4.4 Administration.

Do not freeze. Do not shake.

Keep the vials in the original carton to protect from light.

Any unused product or waste material should be disposed of in accordance with local requirements.

PART II: SCIENTIFIC INFORMATION

13 PHARMACEUTICAL INFORMATION

Drug Substance

Proper name: tixagevimab and cilgavimab injection

Chemical name: tixagevimab and cilgavimab

Molecular formula and molecular mass: tixagevimab: 149 kDa, cilgavimab: 152 kDa

Physicochemical properties: Both tixagevimab and cilgavimab are clear to opalescent, colourless to slightly yellow, pH 6.0 solutions.

Product Characteristics: Tixagevimab and cilgavimab are produced in Chinese hamster ovary (CHO) cells by recombinant DNA technology.

14 CLINICAL TRIALS

14.1 Clinical Trials by Indication

Pre-Exposure Prophylaxis of COVID-19

Trial Design and Study Demographics

Table 3 – Summary of patient demographics: PROVENT for pre-exposure prophylaxis of COVID-19

Study #	Study design	Dosage, route of administration and duration	Study subjects (n)	Mean age (Range)	Sex
D8850C00002 (PROVENT)	Phase III randomized (2:1), double-blind, placebo-controlled, multicentre study to determine the safety and efficacy of EVUSHELD for the pre-exposure prophylaxis of COVID-19 in adults ≥18 years of age	Single dose (administered as two intramuscular injections) of EVUSHELD 300 mg (150 mg of tixagevimab and 150 mg of cilgavimab administered separately) or saline placebo	EVUSHELD: 3441 Placebo: 1731	53.5 (18-99) years	Male: 54% Female: 46%

PROVENT

In PROVENT, all participants were individuals considered to be at increased risk for inadequate response to active immunization (due to age ≥60 years, co-morbidity, pre-existing chronic illness, immunocompromised, or intolerant of vaccination) or at increased risk of SARS-CoV-2 infection (due to their location or circumstances at time of enrolment). Subjects could not have previously received a COVID-19 vaccine. Participants received either a single dose (administered as two intramuscular injections) of EVUSHELD 300 mg (150 mg of tixagevimab

and 150 mg of cilgavimab administered separately) or placebo. The study excluded participants with a history of laboratory-confirmed SARS-CoV-2 infection or SARS-CoV-2 antibody positivity at screening. Once COVID-19 vaccines were locally available, subjects were permitted on request to unblind to make an informed decision on vaccine timing and to receive COVID-19 vaccination.

The baseline demographics were well balanced across the EVUSHELD and placebo arms (Table 3). The median age was 57 years (with 43% of participants aged 60 years or older), 46% of participants were female, 73% were White, 3.3% were Asian, 17% were Black/African American, and 15% were Hispanic/Latino. Of the 5197 participants, 78% had baseline co-morbidities or characteristics associated with an increased risk for severe COVID-19, including obesity (42%), diabetes (14%), cardiovascular disease (8%), cancer, including a history of cancer (7%), chronic obstructive pulmonary disease (5%), chronic kidney disease (5%), chronic liver disease (5%), immunosuppressive medications (3%) and immunosuppressive disease (<1%).

Study Results

PROVENT

The primary endpoint was a COVID-19 case defined as SARS-CoV-2 RT-PCR-positive symptomatic illness occurring between EVUSHELD administration and Day 183. The primary analysis included 5172 participants who were SARS-CoV2 RT-PCR-negative at baseline, of which 3441 received EVUSHELD and 1731 received placebo. Only events that occurred prior to unblinding or vaccine receipt were included. The results of the primary endpoint are shown in Table 4. The median follow-up time post-administration was 83 days.

Table 4 – Incidence of COVID-19 (Full Pre-Exposure Analysis Set)

	N	Number of events ^a , n (%)	Relative Risk Reduction, % (95% CI)
EVUSHELD 300 mg ^b	3441	8 (0.2%)	77% (46 - 90) p-value <0.001
Placebo	1731	17 (1.0%)	

CI = Confidence Interval, N = number of participants in analysis.

^a Primary endpoint, a participant was defined as a COVID-19 case if their first case of SARS-CoV-2 RT-PCR-positive symptomatic illness occurred after administration and prior to Day 183.

^b 300 mg (150 mg tixagevimab and 150 mg cilgavimab).


In a supportive analysis of the incidence of SARS-CoV-2 RT-PCR-positive symptomatic illness or death from any cause for participants who had received EVUSHELD (12/3441) compared with placebo (19/1731), the relative risk reduction was 69% (95% CI: 36, 85).

Among participants who received EVUSHELD there were no severe/critical COVID-19 events (defined as SARS-CoV-2 RT-PCR-positive symptomatic illness characterised by a minimum of either pneumonia [fever, cough, tachypnea or dyspnea, and lung infiltrates] or hypoxemia [SpO₂ <90% in room air and/or severe respiratory distress] and a WHO Clinical Progression Scale score of 5 or higher) compared to one event (0.1%) among participants who received placebo.

An additional data cut-off was conducted to provide post-hoc updated safety and efficacy analyses; the median follow-up was 6.5 months for participants in both the EVUSHELD and

placebo arms. The relative risk reduction of SARS-CoV-2 RT-PCR-positive symptomatic illness was 83% (95% CI 66-91), with 11/3441 [0.3%] events in the EVUSHELD arm and 31/1731 [1.8%] events in the placebo arm (see Figure 1).

Figure 1 – Kaplan Meier: Cumulative Incidence of Symptomatic COVID-19



The predominant SARS-CoV-2 variants in circulation for the time period represented in Figure 1 were Alpha, Beta, Gamma, Epsilon and Delta. Based on the incidence of primary endpoint events, the duration of efficacy was 6 months.

Treatment of mild to moderate COVID-19

Study Design and Demographics

Table 5 – Summary of patient demographics: TACKLE for treatment of mild to moderate COVID-19

Study #	Study design	Dosage, route of administration and duration	Study subjects (n)	Mean age (Range)	Sex
D8851C00001 (TACKLE)	Phase III, randomized (1:1), double-blind, placebo-controlled, multicentre study assessing EVUSHELD for the treatment of adult patients with mild to moderate COVID-19	Single dose (administered as two separate, sequential intramuscular injections) of EVUSHELD 600 mg (300 mg of tixagevimab and 300 mg of cilgavimab administered separately) or saline placebo	Modified Full Analysis Set: EVUSHELD: 413 Placebo: 421	46 (18-86)	Male: 50% Female: 50%

TACKLE

TACKLE is an ongoing Phase III, randomized (1:1), double-blind, placebo-controlled, multicentre study assessing EVUSHELD for the treatment of adult patients with mild to moderate COVID-19.

The study enrolled individuals who had not received COVID-19 vaccination, were not hospitalized for COVID-19 treatment, and had at least 1 or more COVID-19 symptom that was at least mild in severity. The majority of participants (84%) were seronegative at baseline, and 90% were considered at higher risk of progressing to severe COVID-19, defined as either individuals aged 65 years and older at randomization or individuals aged <65 years and having at least one medical condition or other factor that placed them at higher risk for progression to severe COVID-19. High risk co-morbidities included: obesity (BMI ≥ 30) (43%), smoking (current or former) (40%), hypertension (28%), chronic lung disease or moderate to severe asthma (12%), diabetes (12%), cardiovascular disease (including history of stroke) (9%), immunocompromised state (from solid organ transplant, blood or bone marrow transplant, immune deficiencies, HIV, use of corticosteroids, or use of other immunosuppressive medicines) (5%), cancer (4%), chronic kidney disease (2%), chronic liver disease (2%), sickle cell disease (0%). At baseline, 88% of patients had a WHO Clinical Progression Scale score of 2 and 12% had a score of 3. The median duration of symptoms prior to treatment was 5 days.

Demographics and disease characteristics were well balanced across the treatment and placebo groups. At baseline, 13% of subjects were aged 65 years or older, 50% of the subjects were female, 62% were White, 5.6% were Asian, 4.0% were Black and 52% were Hispanic/Latino.

Treatment was initiated within 3 days of obtaining the sample for a positive SARS-CoV-2 viral infection and within ≤ 7 days of COVID-19 symptom onset. Patients received standard of care

treatment and a single dose of either EVUSHELD 600 mg (N=413) or placebo (N=421). Participants were stratified by time from symptom onset (≤ 5 days versus >5 days) and risk of progression to severe COVID-19 (high risk versus low risk).

Study Results

TACKLE

The primary efficacy endpoint was a composite of either severe COVID-19 or death from any cause by Day 29, in subjects who received treatment within 7 days from symptom onset and were not hospitalized at baseline. Severe COVID-19 was defined as characterized by either pneumonia or hypoxemia ($\text{SpO}_2 < 90\%$ and/or severe respiratory distress) and a WHO Clinical Progression Scale score of 5 or higher.

The results of the primary endpoint are shown in Table 6 and demonstrate a statistically significant reduction in severe COVID-19 or death from any cause compared to placebo. The results of the primary composite endpoint were driven by the incidence of severe COVID-19. Up to Day 29, 7 deaths had been reported, 3 in the EVUSHELD arm and 4 in the placebo arm. Of the 7 deaths, 2 were not COVID-19 related. Both of these were in the EVUSHELD arm and contributed to the primary composite endpoint.

Table 6 – Incidence of Severe COVID-19 or death from any cause (Modified Full Analysis Set)

	N	Number of events ^a , n (%)	Relative Risk Reduction, % (95% CI)
EVUSHELD 600 mg ^b	407	18 (4.4%)	50% (15 - 71) p-value = 0.010
Placebo	415	37 (8.9%)	

CI = Confidence Interval, N = number of participants in analysis.

^a Primary endpoint, severe COVID-19 or death from any cause through Day 29.

^b 600 mg (300 mg tixagevimab and 300 mg cilgavimab).

The relative risk reduction in the incidence of severe COVID-19 or death from any cause was 67% (95% CI of 31, 84) in non-hospitalised patients dosed within 5 days of symptom onset.

Viral load was measured as the change from baseline for SARS-CoV-2 RNA (Log_{10} copies/mL) from nasal swabs specimens through Day 29. Treatment with EVUSHELD in comparison with placebo resulted in reductions in viral load at Day 3 (LS mean difference -0.28 [95% CI: -0.64, 0.09]) and Day 6 (LS mean difference -0.48 [95% CI: -0.76, -0.20]).

14.4 Immunogenicity

In PROVENT through Day 183, following a single 300 mg EVUSHELD dose, treatment-emergent anti-tixagevimab, anti-cilgavimab and anti-EVUSHELD antibodies were detected in 3.2% (101/3152), 3.7% (113/3068) and 4.9% (156/3158) anti-drug antibody (ADA)-evaluable participants, respectively. In the PROVENT repeat dose sub-study, participants received a second dose of 300 mg EVUSHELD 10 to 14 months after administration of the initial dose. Up to Day 29 post-administration of the second dose, treatment-emergent anti-tixagevimab, anti-

cilgavimab and anti-EVUSHELD antibodies were detected in 0% (0/49), 10.2% (5/49) and 10.2% (5/49) ADA-evaluable participants, respectively.

No evidence of an association of ADA with any impact on efficacy or safety has been observed.

In TACKLE through Day 169, following a single 600 mg EVUSHELD dose, treatment-emergent anti-tixagevimab, anti-cilgavimab and anti-EVUSHELD antibodies were detected in 5.2% (14/271), 10.7% (33/307), and 10.7% (37/346) ADA-evaluable participants, respectively.

15 MICROBIOLOGY

Viral variants are a potential risk for treatment failure of monoclonal antibodies used for the prevention or treatment of coronavirus disease 2019 (COVID-19) caused by the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2). The marketing authorization for this product is based on the review of clinical or neutralization data supporting efficacy against the variants of concern listed in Table 7. In the absence of data against other variants, the efficacy of the product for these other variants cannot be confirmed.

Antiviral activity

In a SARS-CoV-2 virus neutralization assay on Vero E6 cells, tixagevimab, cilgavimab and their combination neutralized SARS-CoV-2 (USA-WA1/2020 isolate) with EC₅₀ values of 60.7 pM (9 ng/mL), 211.5 pM (32 ng/mL) and 65.9 pM (10 ng/mL), respectively.

Tixagevimab, cilgavimab and the tixagevimab and cilgavimab combination showed reduced or no antibody-dependent cellular phagocytosis (ADCP) antibody-dependent cell-mediated cytotoxicity (ADCC) or antibody-dependent natural killer cell activation (ADNKA) in cell culture studies. Tixagevimab, cilgavimab and the tixagevimab and cilgavimab combination did not mediate ADCC activity with guinea pig complement proteins.

Antibody dependent enhancement (ADE) of infection

The potential of tixagevimab and cilgavimab to mediate antibody-dependent viral entry was assessed in FcγRII-expressing Raji cells co-incubated with recombinant virus pseudotyped with SARS-CoV-2 spike protein, with antibody concentrations at a range of 6.6 nM (1 µg/mL) to 824 pM (125 ng/mL). Tixagevimab, cilgavimab and their combination did not mediate entry of pseudovirus into these cells.

The potential for ADE was also evaluated in a non-human primate model of SARS-CoV-2 using EVUSHELD. Intravascular administration prior to virus inoculation resulted in a dose-dependent improvement in all measured outcomes (total viral RNA in the lungs or nasal mucosae, infectious virus levels in the lungs based on TCID₅₀ measurements, and lung injury and pathology based on histology measurements). No evidence of enhancement of disease was observed at any dose evaluated, including sub-neutralizing doses down to 0.04 mg/kg.

Antiviral resistance

There is a risk of resistance to monoclonal antibodies such as EVUSHELD due to the development of viral variants that are resistant to tixagevimab and cilgavimab. Prescribing healthcare professionals should consider the prevalence of SARS-CoV-2 variants in their area, in combination with antiviral resistance information, literature, and local guidelines when considering pre-exposure prophylaxis with EVUSHELD.

SARS-CoV-2 or recombinant vesicular stomatitis virus encoding SARS-CoV-2 spike protein were serially passaged in cell cultures in the presence of cilgavimab or tixagevimab individually, or tixagevimab and cilgavimab in combination. Escape variants were identified following passage with cilgavimab, but not with tixagevimab or tixagevimab and cilgavimab in combination. Variants which showed reduced susceptibility to cilgavimab alone included spike protein amino acid substitutions R346I (>200-fold), K444E (>200-fold), and K444R (>200-fold). All variants maintained susceptibility to tixagevimab alone, and tixagevimab and cilgavimab in combination.

Evaluation of neutralization susceptibility of variants identified through global surveillance and in participants who received tixagevimab and cilgavimab is ongoing.

In neutralization assays using recombinant SARS-CoV-2 pseudoviruses harbouring individual spike substitutions identified in circulating SARS-CoV-2, variants with reduced susceptibility to tixagevimab alone included those with F486S (>600-fold) and F486V (121- to 149-fold); variants with reduced susceptibility to cilgavimab alone included those with R346I (>200-fold), K444E (>200-fold), K444Q (>200-fold), K444R (>200-fold), and V445A (21- to 51-fold).

Pseudovirus SARS-CoV-2 spike variant strains with moderate reduced susceptibility to tixagevimab alone included those harbouring E484K (Alpha, 18.5-fold; Beta, 3.5- to 15-fold) and variants with moderate reduced susceptibility to cilgavimab alone included those with R346K:E484K:N501Y (Mu, 21-fold), as indicated above. Similar results were observed, where data was available, in neutralization assays using authentic SARS-CoV-2 variants strains.

Neutralization activity of EVUSHELD against pseudovirus and/or live virus SARS-CoV-2 variant strains are shown in Table 7. Data collection is ongoing to better understand how small reductions in activity seen in authentic SARS-CoV-2 or pseudotyped VLP assays may correlate with clinical outcomes.

Table 7 – Pseudovirus and Authentic SARS-CoV-2 Neutralization Data for SARS-CoV-2 Variant Substitutions with Tixagevimab and Cilgavimab Together

Lineage with Spike Protein Substitutions		Characteristic RBD Substitutions Tested	Fold Reduction in Susceptibility ^a		IC ₅₀ (ng/mL)	
Pango Lineage	WHO Label		Pseudo virus ^b	Authentic SARS-CoV-2 ^c	Pseudo virus ^b	Authentic SARS-CoV-2 ^c
Variants of Concern						
B.1.1.7	Alpha	N501Y	1.0-5.2	0.5-1.4	1.1-9.0	4-39.5
B.1.351	Beta	K417N:E484K: N501Y	2.5-5.5	0.9-3.8	5.6-11.4	6.5-256
P.1	Gamma	K417T:E484K: N501Y	0.8-1.7	0.4-2.0	1.8-2.7	3.2-8
B.1.617.2	Delta	L452R:T478K	1.0-1.2	0.6-1.0	1.9-2.2	3-7.5
AY.1/AY.2	Delta [+K417N]	K417N:L452R: T478K	1.0	ND	1.9	ND
B.1.1.529	Omicron BA.1	A67V, H69-, V70-, T95I, G142D, V143-, Y144-, Y145-, N211-, L212I, ins214EPE,	132- to 183-fold	12- to 30-fold	51-277	147-278

Lineage with Spike Protein Substitutions		Characteristic RBD Substitutions Tested	Fold Reduction in Susceptibility ^a		IC ₅₀ (ng/mL)	
Pango Lineage	WHO Label		Pseudo virus ^b	Authentic SARS-CoV-2 ^c	Pseudo virus ^b	Authentic SARS-CoV-2 ^c
		G339D, S371L, S373P, S375F, K417N, N440K, G446S, S477N, T478K, E484A, Q493R, G496S, Q498R, N501Y, Y505H, T547K, D614G, H655Y, N679K, P681H, N764K, D796Y, N856K, Q954H, N969K, L981F.				
BA.1.1	Omicron BA.1.1	G339D:R346K: S371L:S373P: S375F:K417N: N440K:G446S: S477N:T478K: E484A:Q493R: G496S:Q489R: N501Y:Y505H	424-fold	176-fold	466	1147
BA.2	Omicron BA.2	G339D:S371F: S373P:S375F: T376A:D405N: R408S:K417N: N440K:S477N: T478K:E484A: Q493R:Q498R: N501Y:Y505H:	3.2	5.4	9.8	35
BA.2.12.1	Omicron BA.2.12.1	G339D:S371F:S373P: S375F:T376A:D405N: R408S:K417N:N440K: L452Q:S477N:T478K: E484A:Q493R:Q498R: :N501Y:Y505H	5-fold	ND	10.7	ND
BA.2.75	Omicron BA.2.75	G339H:S371F:S373P: S375F:T376A:D405N: R408S:K417N:N440K: G446S:N460K:S477N: T478K:E484A:Q498R: N501Y:Y505H	2.4- to 15-fold	ND	1.2-14	ND
BA.2.75.2	Omicron BA.2.75.2	BA.2.75+ R346T:F486S	>5000-fold ^d	ND	>10000 ^d	ND
BA.3	Omicron BA.3	G339D: S371F:S373P: S375F:D405N:K417N: N440K:G446S:S477N:	16-fold	ND	34.5	ND

Lineage with Spike Protein Substitutions		Characteristic RBD Substitutions Tested	Fold Reduction in Susceptibility ^a		IC ₅₀ (ng/mL)	
Pango Lineage	WHO Label		Pseudo virus ^b	Authentic SARS-CoV-2 ^c	Pseudo virus ^b	Authentic SARS-CoV-2 ^c
		T478K:E484A:Q493R:Q498R:N501Y:Y505H				
BA.4	Omicron BA.4	G339D:S371F:S373P:S375F:T376A:D405N:R408S:K417N:N440K:L452R:S477N:T478K:E484A:F486V:Q498R:N501Y:Y505H	33- to 65-fold	ND	65-69.4	ND
BA.4.6	Omicron BA.4.6	G339D:R346T:S371F:S373P:S375F:T376A:D405N:R408S:K417N:N440K:L452R:S477N:T478K:E484A:F486V:Q498R:N501Y:Y505H	>1000-fold ^d	ND	>1000 ^d	ND
BA.5	Omicron BA.5	G339D:S371F:S373P:S375F:T376A:D405N:R408S:K417N:N440K:L452R:S477N:T478K:E484A:F486V:Q498R:N501Y:Y505H	33- to 65-fold	2.8- to 16-fold	65-69.4	56.6-229
BF.7	Omicron BF.7	BA.4+ R346T	>5000-fold ^d	ND	>10000 ^d	ND
BJ.1	Omicron BJ.1	G339H:R346T:L368I:S371F:S373:S375F:T376A:D405N:R408S:K417N:N440K:V445P:G446S:S477N:T478K:V483A:E484A:F490V:Q493R:Q498R:N501Y:Y505H	228 to 424-fold	ND	228-848	ND
BN.1	Omicron BN.1	G339D:R346T:K356T:S371F:S373P:S375F:D405N:R408S:K417N:N440K:G446S:N460K:S477N:T478K:E484A:F490S:Q493R:Q498R:Y505H	68-fold	ND	61-68	ND
BQ.1	Omicron BQ.1	BA.5+ K444T:N460K	>2000-fold ^d	ND	>10000 ^d	ND
BQ.1.1	Omicron BQ.1.1	BA.5+ R346T:K444T:N460K	>2000-fold ^d	ND	>10000 ^d	ND
XBB	Omicron XBB	G339H:R346T:L368I:S371F:S373P:S375F:T376A:D405N:R408S:K417N:N440K:V445P:G446S:N460K:S477N:	>1400-fold ^d	ND	>1000 ^d	ND

Lineage with Spike Protein Substitutions		Characteristic RBD Substitutions Tested	Fold Reduction in Susceptibility ^a		IC ₅₀ (ng/mL)	
Pango Lineage	WHO Label		Pseudo virus ^b	Authentic SARS-CoV-2 ^c	Pseudo virus ^b	Authentic SARS-CoV-2 ^c
		T478K:E484A:F486S: F490S:Q498R:N501Y: Y505H				
XBB.1.5	Omicron XBB.1.5	G339H+R346T+ L368I+S371F+S373+ S375F+T376A+ D405N+R408S+ K417N+N440K+ V445P+G446S+ N460K+S477N+ T478K+E484A+ F486P+F490S+ Q498R+N501Y +Y505H	>5000 fold ^d	ND	>10000 ^d	ND
Variants of Interest						
B.1.525	Eta	E484K	1.8-3.1	ND	5-9.5	ND
B.1.526	Iota	E484K	0.8-3.4	0.3-1.8	1.9-5.2	1.0-7.0
B.1.617.1	Kappa	L452R:E484Q	0.9-3.4	0.5-1.3	2.5-5.1	2.0-5.0
C.37	Lambda	L452Q:F490S	0.7	ND	1.1	ND
B.1.621	Mu	R346K:E484K: N501Y	7.5	ND	17.3	ND
B.1.427 / B.1.429	Epsilon	L452R	0.8-2.9	1.3-3.5	1.0-4.5	5.0-14.0
P.2	Zeta	E484K	2.9	ND	10.4	ND
Additional SARS-CoV-2 Variants						
R.1	-	E484K	3.5	ND	4.6	ND
B.1.1.519	-	T478K	1.0-1.4	ND	2.3	ND
C.36.3	-	R346S:L452R	2.3	ND	3.9	ND
B.1.214.2	-	Q414K:N450K	0.8	ND	1.6	ND
B.1.619.1	-	N440K:E484K	3.3	ND	7.6	ND
B.1.616	-	V483A	0.4-0.5	ND	1.1-1.2	ND
A.23.1	-	V367F	0.4	ND	0.5	ND
A.27	-	L452R:N501Y	0.8	ND	1.8	ND
AV.1	-	N439K:E484K	5.9	ND	13.0	ND

^a Range of reduced *in vitro* potency across multiple sets of co-occurring substitutions and/or testing labs using research-grade assays; mean fold change in half maximal inhibitory concentration (IC₅₀) of monoclonal antibody required for a 50% reduction in infection compared to wild type reference strain.

^b Pseudoviruses expressing the entire SARS-CoV-2 spike variant protein and individual characteristic spike substitutions except L452Q were tested including Alpha (+L455F, E484K, F490S, Q493R, and/or S494P), and Delta (+K417N) harbouring additional indicated RBD substitutions that are no longer detected or detected at extremely low levels within these lineages.

- c Authentic SARS-CoV-2 expressing the entire variant spike protein were tested including Alpha (+E484K or S494P) harbouring additional indicated RBD substitutions that are no longer detected or detected at extremely low levels within these lineages.
 - d Tixagevimab and cilgavimab together are unlikely to be active against this variant.
- ND, not determined; RBD, receptor binding domain.

It is not known how pseudovirus or authentic SARS-CoV-2 neutralization susceptibility data correlate with clinical outcome.

16 NON-CLINICAL TOXICOLOGY

General Toxicology: In a single-dose toxicology study in cynomolgus monkeys, EVUSHELD administered via an intramuscular injection of 150 mg/kg (75 mg/kg of each antibody) had no adverse effects.

In tissue cross-reactivity studies using human adult and fetal tissues no specific binding was detected.

Carcinogenicity: studies have not been conducted.

Genotoxicity: studies have not been conducted.

Reproductive and Developmental Toxicology: studies have not been conducted.

PATIENT MEDICATION INFORMATION

READ THIS FOR SAFE AND EFFECTIVE USE OF YOUR MEDICINE

EVUSHELD™ tixagevimab and cilgavimab injection, intramuscular use

Read this carefully before you start taking **EVUSHELD** and each time you get a refill. This leaflet is a summary and will not tell you everything about this drug. Talk to your healthcare professional about your medical condition and treatment and ask if there is any new information about **EVUSHELD**.

What is EVUSHELD used for?

EVUSHELD is used:

- For the pre-exposure prophylaxis (prevention) of Coronavirus Disease 2019 (COVID-19) illness in adults and adolescents (aged 12 years and older weighing at least 40 kg) who:
 - have a weaker immune system and are unlikely to be protected by a COVID-19 vaccine
 - or when vaccination is not recommended.
- To treat adults and adolescents (aged 12 years and older weighing at least 40 kg) with confirmed mild to moderate COVID-19.

EVUSHELD is not a substitute for vaccination in individuals for whom COVID-19 vaccination is recommended.

How does EVUSHELD work?

EVUSHELD contains the active substances tixagevimab and cilgavimab. Tixagevimab and cilgavimab are types of protein called 'monoclonal antibodies'. EVUSHELD work specifically against the SARS-CoV-2 virus (the virus that causes COVID-19 illness) by preventing the virus from infecting healthy cells in your body. This can help prevent you from getting COVID-19 illness or help your body to overcome the infection and reduce the risk of you developing severe illness.

What are the ingredients in EVUSHELD?

Medicinal ingredients: tixagevimab and cilgavimab

Non-medicinal ingredients: L-histidine, L-histidine hydrochloride monohydrate, polysorbate 80, sucrose and water.

EVUSHELD comes in the following dosage forms:

A carton containing two clear glass vials:

- 1 vial of tixagevimab (100 mg/mL) solution for injection (dark grey vial cap)
- 1 vial of cilgavimab (100 mg/mL) solution for injection (white vial cap)

Both solutions are a clear to opalescent, colourless to slightly yellow solution.

Do not use EVUSHELD if:

- if you are allergic to tixagevimab, cilgavimab or any of the other ingredients of this medicine (see What are the ingredients in EVUSHELD).

To help avoid side effects and ensure proper use, talk to your healthcare professional before you take EVUSHELD. Talk about any health conditions or problems you may have, including if you:

- have low numbers of blood platelets (which help blood clotting), a bleeding disorder or are taking an anticoagulant medicine (to prevent blood clots).
- are pregnant, think you may be pregnant or are planning to have a baby, ask your healthcare professional for advice before receiving this medicine. There is not enough information to be sure that EVUSHELD is safe for use in pregnancy. EVUSHELD will only be given if the potential benefits of use outweigh the potential risks to you and your unborn child.
- are breast-feeding, ask your healthcare professional for advice before receiving this medicine. It is unknown if EVUSHELD or the COVID-19 virus pass into human breast milk. You will need to consider the potential benefits of use for you, compared with the health benefits and risks of breast-feeding for your baby.
- have a history of severe allergic reaction to this drug.
- have had a heart attack or stroke, have other heart problems, or are at high-risk of cardiac (heart) events.

Serious allergic reaction

Tell your healthcare professional or seek medical help right away if you notice any signs of serious allergic reaction during or following administration of EVUSHELD including: difficulty breathing or swallowing, swelling of the face, lips, tongue or throat, severe itching of the skin, with a red rash or raised bumps.

Other warnings you should know about:

EVUSHELD should not be given to children below 12 years of age or weighing less than 40 kg.

COVID-19 is caused by different strains (variants) of the SARS-CoV-2 virus that change over time. EVUSHELD may be less effective at preventing COVID-19 caused by some strains than others. Contact your healthcare professional right away if you get symptoms of COVID-19. COVID-19 affects different people in different ways:

- the most common symptoms include fever, cough, tiredness and loss of taste or smell;
- the most serious symptoms include difficulty breathing or shortness of breath, loss of speech or mobility, or confusion and chest pain.

Tell your healthcare professional about all the medicines you take, including any drugs, vitamins, minerals, natural supplements or alternative medicines.

How to take EVUSHELD:

EVUSHELD will be given to you by a healthcare professional as two intramuscular injections, usually one into each of your buttocks.

Usual dose:

EVUSHELD consists of two medicines, tixagevimab and cilgavimab. You will receive 2 injections one after the other.

The recommended dose for pre-exposure prophylaxis (prevention) of COVID-19 is 600 milligrams (mg) given as two 3 mL injections:

- 300 mg of tixagevimab

- 300 mg of cilgavimab

For continuous prevention of COVID-19 you will need to receive repeat doses of 600 mg EVUSHELD (300 mg of tixagevimab and 300 mg of cilgavimab) every 6 months.

The recommended dose for treatment of mild to moderate COVID-19 is 600 mg, given as two 3 mL injections:

- 300 mg of tixagevimab
- 300 mg of cilgavimab

Overdose:

If you think you, or a person you are caring for, have taken too much EVUSHELD, contact a healthcare professional, hospital emergency department, or regional poison control centre immediately, even if there are no symptoms.

What are possible side effects from using EVUSHELD?

These are not all the possible side effects you may have when taking EVUSHELD. If you experience any side effects not listed here, tell your healthcare professional.

Allergic reactions may be serious and occur during or following administration of monoclonal antibodies (see Serious allergic reaction above).

A higher percentage of people who received EVUSHELD compared to people who did not receive EVUSHELD reported serious cardiac adverse events in a clinical trial. It is not known if these events are related to EVUSHELD or underlying medical conditions. Contact your healthcare professional or get medical help right away if you get any symptoms of cardiac events, including pain, pressure, or discomfort in the chest, arms, neck, back, stomach or jaw, as well as shortness of breath, feeling tired or weak (fatigue), feeling sick (nausea), or swelling in your ankles or lower legs.

If you notice any side effects, please tell your healthcare professional right away:

Common: may affect up to 1 in 10 people

- hypersensitivity reaction (rash or hives - an itchy red rash or raised bumps)
- injection site reaction (pain, redness, itching, swelling where the injection was given)

Uncommon: may affect up to 1 in 100 people

- injection related reaction (examples of these include headache, chills and redness, discomfort or soreness near where the injection was given)

Frequency not known: frequency cannot be estimated from the available data

- serious allergic reaction (serious hypersensitivity, including anaphylaxis - difficulty breathing or swallowing, swelling of the face, lips, tongue or throat, severe itching of the skin, with a red rash or raised bumps)

If you have a troublesome symptom or side effect that is not listed here or becomes bad enough to interfere with your daily activities, tell your healthcare professional.

Reporting Side Effects

You can report any suspected side effects associated with the use of health products to Health Canada by:

- Visiting the Web page on Adverse Reaction Reporting (<https://www.canada.ca/en/health-canada/services/drugs-health-products/medeffect-canada/adverse-reaction-reporting.html>) for information on how to report online, by mail or by fax; or
- Calling toll-free at 1-866-234-2345.

NOTE: Contact your health professional if you need information about how to manage your side effects. The Canada Vigilance Program does not provide medical advice.

Storage:

The following information about storage, expiry and use and handling is intended for the healthcare professional:

- Keep out of reach and sight of children.
- Do not use this medicine after the expiry date which is stated on the label and carton after EXP. The expiry date refers to the last day of that month.
- Store in a refrigerator (2°C to 8°C).
- Do not freeze. Do not shake.
- Store in the original package in order to protect from light.
- Prepared syringes should be used immediately. If necessary, store the prepared syringes for no more than 4 hours either: at 2°C to 8°C, or at room temperature up to 25°C.
- Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

If you want more information about EVUSHELD:

- Talk to your healthcare professional.
- Find the full product monograph that is prepared for healthcare professionals and includes this Patient Medication Information by visiting the Health Canada website: <https://www.canada.ca/en/health-canada/services/drugs-health-products/drug-products/drug-product-database.html>; the manufacturer's website: www.astrazeneca.ca, or by calling 1-800-668-6000.
- This Patient Medication Information is current at the time of printing. The most up-to-date version can be found at www.astrazeneca.ca.

This leaflet was prepared by AstraZeneca Canada Inc., Mississauga, Ontario L4Y 1M4

EVUSHELD™ is a trademark of AstraZeneca UK Ltd., used under license by AstraZeneca Canada Inc. The AstraZeneca logo is a registered trademark of AstraZeneca AB, used under license by AstraZeneca Canada Inc.

© AstraZeneca Canada Inc. 2023

Last Revised May 30, 2023

