

PRODUCT MONOGRAPH  
INCLUDING PATIENT MEDICATION INFORMATION

<sup>Pr</sup>**Osetamivir Powder for Oral Suspension**

oseltamivir powder for oral suspension  
6 mg/mL oseltamivir (as oseltamivir phosphate) when reconstituted

House Standard

Antiviral Agent

Strides Pharma Canada Inc.  
1565, Boul. Lionel-Boulet,  
Varenes, Quebec  
Canada J3X 1P7

Date of Initial Authorization:

December 30, 2020

Date of Revision:

June 1, 2023

Submission Control Number: 270702

**RECENT MAJOR LABEL CHANGES**

4 Dosage and Administration	06/2023
7 Warnings and Precautions, 7.1.1 Pregnant Women	06/2023

**TABLE OF CONTENTS**

Sections or subsections that are not applicable at the time of authorization are not listed.

**RECENT MAJOR LABEL CHANGES ..... 2**

**TABLE OF CONTENTS ..... 2**

**PART I: HEALTH PROFESSIONAL INFORMATION ..... 4**

**1 INDICATIONS..... 4**

    1.1 Pediatrics.....5

    1.2 Geriatrics .....5

**2 CONTRAINDICATIONS ..... 5**

**4 DOSAGE AND ADMINISTRATION ..... 5**

    4.1 Dosing Considerations .....5

    4.2 Recommended Dose and Dosage Adjustment.....5

    4.3 Reconstitution .....8

    4.4 Administration..... 11

    4.5 Missed Dose ..... 11

**5 OVERDOSAGE..... 11**

**6 DOSAGE FORMS, STRENGTHS, COMPOSITION AND PACKAGING ..... 12**

**7 WARNINGS AND PRECAUTIONS ..... 13**

    7.1 Special Populations..... 15

        7.1.1 Pregnant Women .....15

        7.1.2 Breast-feeding ..... 15

        7.1.3 Pediatrics .....15

        7.1.4 Geriatrics ..... 15

**8 ADVERSE REACTIONS ..... 16**

    8.1 Adverse Reaction Overview .....16

    8.2 Clinical Trial Adverse Reactions ..... 16

        8.2.1 Clinical Trial Adverse Reactions – Pediatrics .....18

8.3	Less Common Clinical Trial Adverse Reactions .....	21
8.4	Abnormal Laboratory Findings: Hematologic, Clinical Chemistry and Other Quantitative Data .....	21
8.5	Post-Market Adverse Reactions .....	21
<b>9</b>	<b>DRUG INTERACTIONS .....</b>	<b>21</b>
9.2	Drug Interactions Overview .....	21
9.3	Drug-Behavioural Interactions.....	21
9.4	Drug-Drug Interactions .....	21
9.5	Drug-Food Interactions.....	22
9.6	Drug-Herb Interactions.....	22
9.7	Drug-Laboratory Test Interactions.....	22
<b>10</b>	<b>CLINICAL PHARMACOLOGY .....</b>	<b>23</b>
10.1	Mechanism of Action.....	23
10.2	Pharmacodynamics .....	24
10.3	Pharmacokinetics .....	24
<b>11</b>	<b>STORAGE, STABILITY AND DISPOSAL .....</b>	<b>25</b>
<b>12</b>	<b>SPECIAL HANDLING INSTRUCTIONS .....</b>	<b>26</b>
<b>PART II: SCIENTIFIC INFORMATION .....</b>		<b>27</b>
<b>13</b>	<b>PHARMACEUTICAL INFORMATION .....</b>	<b>27</b>
<b>14</b>	<b>CLINICAL TRIALS .....</b>	<b>27</b>
14.1	Clinical Trials by Indication .....	27
14.2	Comparative Bioavailability Studies.....	31
<b>15</b>	<b>MICROBIOLOGY .....</b>	<b>31</b>
<b>16</b>	<b>NON-CLINICAL TOXICOLOGY.....</b>	<b>33</b>
<b>17</b>	<b>SUPPORTING PRODUCT MONOGRAPHS.....</b>	<b>34</b>
<b>PATIENT MEDICATION INFORMATION.....</b>		<b>35</b>

## PART I: HEALTH PROFESSIONAL INFORMATION

### 1 INDICATIONS

#### Treatment of Influenza

Oseltamivir Powder for Oral Suspension (oseltamivir phosphate) is indicated for:

- The treatment of uncomplicated acute illness due to influenza infection in adults and adolescents ( $\geq 13$  years) who have been symptomatic for no more than 2 days.

The treatment indication is based on two Phase III clinical studies of naturally occurring influenza in adults in which the predominant infection was influenza A (95%) and a limited number with influenza B (3%) and influenza of unknown type (2%), reflecting the distribution of these strains in the community. The indication is also supported by influenza A and B challenge studies. No data are available to support the safety and efficacy of oseltamivir phosphate in adult patients who commenced treatment after 40 hours of onset of symptoms.

- The treatment of uncomplicated acute illness due to influenza in pediatric patients 1 year and older who have been symptomatic for no more than 2 days.

The pediatric indication is based on one Phase III clinical study of naturally occurring influenza in pediatric patients aged 1 to 12 years in which 67% of influenza infected patients were infected with influenza A and 33% with influenza B.

Oseltamivir Powder for Oral Suspension, when taken as recommended for the treatment of influenza, alleviates the symptoms and reduces their duration (see [14 CLINICAL TRIALS](#)).

#### Prevention/Prophylaxis of Influenza

The decision to administer Oseltamivir Powder for Oral Suspension for prophylaxis to close contacts should be based on the knowledge that influenza is circulating in the area and the index case demonstrates characteristic symptoms of influenza. Oseltamivir Powder for Oral Suspension is not effective in providing prophylaxis for respiratory infections other than influenza, therefore a proper diagnosis of the index case is important.

Oseltamivir Powder for Oral Suspension is not a substitute for influenza vaccination. Vaccination is the preferred method of prophylactic prevention against influenza. The use of Oseltamivir Powder for Oral Suspension should not affect the evaluation of individuals for annual influenza vaccination, in accordance to "Health Canada. An Advisory Committee Statement on Influenza Vaccination for the Current Year/Season."

The use of antivirals for the treatment and prevention of influenza should be determined on the basis of official recommendations taking into consideration variability of epidemiology and the impact of the disease in different geographical areas and patient populations.

Oseltamivir Powder for Oral Suspension is indicated for:

- The prevention of influenza illness in adults and adolescents 13 years and older following close contact with an infected individual (the index case).

The prevention indication is based on a Phase III clinical study programme consisting of 4 Phase III clinical trials.

- The prevention of influenza illness in pediatric patients 1 year and older following close contact with an infected individual (the index case).

This indication is based on a sub-study of pediatric patients in a Phase III clinical trial.

## 1.1 Pediatrics

**Pediatrics (< 1 year of age):** The efficacy of oseltamivir phosphate in infants younger than 1 year of age have not been established (see [7 WARNINGS AND PRECAUTIONS, 7.1.3 Pediatrics](#) and [14 CLINICAL TRIALS](#)); therefore, Health Canada has not authorized an indication for pediatric use.

## 1.2 Geriatrics

**Geriatrics (≥ 65 years of age):** Efficacy of oseltamivir phosphate in the treatment of elderly patients has not been evaluated. Safety data in 372 elderly patients (≥ 65 years old) showed no overall difference between these subjects and younger adults. Based on drug exposure and tolerability, dosage adjustments are not anticipated for elderly patients (see [7 WARNINGS AND PRECAUTIONS, 7.1.4 Geriatrics](#) and [10 CLINICAL PHARMACOLOGY, 10.3 Pharmacokinetics, Special Populations and Conditions, Geriatrics](#)).

## 2 CONTRAINDICATIONS

- Oseltamivir Powder for Oral Suspension (oseltamivir phosphate) is contraindicated in patients who are hypersensitive to this drug or to any ingredient in the formulation, including any non-medicinal ingredient, or component of the container. For a complete listing, see [6 DOSAGE FORMS, STRENGTHS, COMPOSITION AND PACKAGING](#).

## 4 DOSAGE AND ADMINISTRATION

### 4.1 Dosing Considerations

- **Hepatic Impairment:** The safety, efficacy and pharmacokinetics in patients with severe hepatic impairment have not been studied. No studies have been carried out in pediatric patients with hepatic impairment.
- **Infants:** The efficacy of oseltamivir phosphate in infants younger than 1 year of age has not been established. Oseltamivir Powder for Oral Suspension should not be used in children under 1 year of age (see [16 NON-CLINICAL TOXICOLOGY, General Toxicology, Multiple Dose Toxicity](#)).
- For information on renal impairment and elderly patients, see 4.2 Recommended Dose and Dosage Adjustment, Dosage Adjustment.

### 4.2 Recommended Dose and Dosage Adjustment

#### **Recommended Dose – Treatment of Influenza**

Treatment should begin no more than two days after the onset of symptoms of influenza.

**Adults and Adolescents (≥ 13 years):** The recommended oral dose of Oseltamivir Powder for Oral Suspension for the treatment of influenza in adults and adolescents 13 years and older is 75 mg (12.5 mL oral suspension) twice daily, for 5 days.

**Pediatrics (1 to 12 years):** The recommended oral dose of Oseltamivir Powder for Oral Suspension for pediatric patients 1 year and older is shown in the table below.

Body Weight in kg	Body Weight in lbs	Recommended Dose for 5 Days	Amount of Oral Suspension (6 mg/mL)
-------------------	--------------------	-----------------------------	-------------------------------------

≤ 15 kg	≤ 33 lbs	30 mg twice daily	5.0 mL twice daily
> 15 kg to 23 kg	> 33 lbs to 51 lbs	45 mg twice daily	7.5 mL twice daily
> 23 kg to 40 kg	> 51 lbs to 88 lbs	60 mg twice daily	10.0 mL twice daily
> 40 kg	> 88 lbs	75 mg twice daily	12.5 mL twice daily

Use appropriate oral dosing dispenser for volume dispensed with the oral suspension. In the event that the dispenser provided is lost or damaged, another dosing dispenser may be used to deliver the volumes.

### **Recommended Dose – Prevention of Influenza**

Therapy should begin within 2 days of exposure after the onset of symptoms in the index case and continue for at least ten days. Viral shedding may continue for up to 14 days in children and elderly after the onset of influenza illness. Therefore, if the index case is a child or an elderly person, therapy with Oseltamivir Powder for Oral Suspension for prevention may continue for up to 14 days.

Patients should be instructed to complete the entire course of therapy.

**Adults and adolescents (≥ 13 years):** The recommended oral dose of Oseltamivir Powder for Oral Suspension for the prevention of influenza in adults and adolescents 13 years and older following close contact with an infected individual (the index case) is 75 mg (12.5 mL oral suspension) once daily.

Safety has been demonstrated for up to 12 weeks in immunocompromised patients.

**Pediatrics (1 to 12 years):** The recommended dose of Oseltamivir Powder for Oral Suspension for prevention in pediatric patients 1 year and older is shown in the table below.

Body Weight in kg	Body Weight in lbs	Recommended Dose for at least 10 Days	Amount of Oral Suspension (6 mg/mL)
≤ 15 kg	≤ 33 lbs	30 mg once daily	5.0 mL once daily
> 15 kg to 23 kg	> 33 lbs to 51 lbs	45 mg once daily	7.5 mL once daily
> 23 kg to 40 kg	> 51 lbs to 88 lbs	60 mg once daily	10.0 mL once daily
> 40 kg	> 88 lbs	75 mg once daily	12.5 mL once daily

Use appropriate oral dosing dispenser for volume dispensed with the oral suspension. In the event that the dispenser provided is lost or damaged, another dosing dispenser may be used to deliver the volumes.

### **Dosage Adjustment**

**Hepatic Impairment:** No dose adjustment is required in adult patients with mild or moderate hepatic impairment (see [10 CLINICAL PHARMACOLOGY, 10.3 Pharmacokinetics, Special Populations and Conditions, Hepatic Insufficiency](#)).

**Renal Impairment:** No dose adjustment is necessary for patients with creatinine clearance above 60 mL/min. Dosage adjustments for adult patients with creatinine clearance ≤ 60 mL/min are as indicated below. Note that the dosage adjustments are based primarily on population pharmacokinetic modeling data. Clinical safety and efficacy studies at these doses have not been performed.

Treatment of Influenza: In adult patients with a creatinine clearance of > 30-60 mL/min, it is recommended that the dose be reduced to 30 mg of Oseltamivir Powder for Oral Suspension twice daily for 5 days.

In adult patients with a creatinine clearance of 10-30 mL/min, it is recommended that the dose be reduced to 30 mg of Oseltamivir Powder for Oral Suspension once daily for 5 days.

Treatment of Influenza for patients undergoing routine hemodialysis: In adult patients, an initial dose of 30 mg of Oseltamivir Powder for Oral Suspension can be administered prior to the start of dialysis if influenza symptoms develop during the 48 hours between dialysis sessions. To maintain plasma concentrations at a therapeutic level, a dose of 30 mg of Oseltamivir Powder for Oral Suspension should be administered after every hemodialysis session, over a period of 5 days.

For adult patients on continuous ambulatory peritoneal dialysis (CAPD), a single 30 mg dose of Oseltamivir Powder for Oral Suspension administered prior to the start of dialysis is recommended for treatment.

Prevention of influenza: In adult patients with a creatinine clearance of > 30-60 mL/min, it is recommended that the dose be reduced to 30 mg of Oseltamivir Powder for Oral Suspension once daily for 10-14 days.

In adult patients with a creatinine clearance of 10-30 mL/min, it is recommended that the dose be reduced to 30 mg of Oseltamivir Powder for Oral Suspension every other day for a period of 10-14 days.

Prevention of Influenza for patients undergoing routine hemodialysis: In adult patients, an initial dose of 30 mg of Oseltamivir Powder for Oral Suspension can be administered prior to the start of dialysis. To maintain plasma concentrations at a therapeutic level, a dose of 30 mg should be administered after every alternate hemodialysis session for a period of 10-14 days.

For adult patients on peritoneal dialysis, an initial dose of 30 mg of Oseltamivir Powder for Oral Suspension administered prior to the start of dialysis followed by further 30 mg doses administered every 7 days, for a period of 10-14 days, is recommended for prophylaxis.

Note that the pharmacokinetics of oseltamivir have not been studied in patients with end-stage renal disease (i.e., creatinine clearance of < 10 mL/min) not undergoing dialysis. Hence, dosing recommendation cannot be provided for this group.

**Geriatrics:** No dose adjustment is required for elderly patients with normal renal function (see [7 WARNINGS AND PRECAUTIONS, 7.1 Special Populations, 7.1.4 Geriatrics](#)).

### 4.3 Reconstitution

#### Reconstitution of Oral Suspension:

It is recommended that Oseltamivir Powder for Oral Suspension be reconstituted by the pharmacist prior to dispensing to the patient.

1. Tap the closed bottle several times to loosen the powder.
2. Measure 55 mL of water in a graduated cylinder\*.
3. Add the total amount of water for reconstitution to the bottle and shake the closed bottle well for 15 seconds. (The bottle containing 360 mg oseltamivir will contain 60 mL of suspension at a final concentration of 6 mg/mL of oseltamivir (as oseltamivir phosphate) when reconstituted.)
4. Remove the child-resistant cap and push bottle adapter into neck of bottle.

5. Close bottle with child-resistant cap tightly. This will assure the proper seating of the bottle adapter in the bottle and child-resistant status of the cap.

\*Please ensure the overcap provided is not dispensed to the patient if Oseltamivir Powder for Oral Suspension is reconstituted by the pharmacist.

Dispense with patient information leaflet and oral dispenser. It is recommended to write the date of expiration of the reconstituted suspension on the bottle label. (The shelf life of the reconstituted suspension is 10 days if stored at room temperature (15-30°C) or 17 days if stored in a refrigerator (2-8°C)).

#### 4.4 Administration

Oseltamivir Powder for Oral Suspension may be taken with or without food (see [10 CLINICAL PHARMACOLOGY, 10.3 Pharmacokinetics, Absorption](#)). However, taking with food may enhance tolerability in some patients.

Shake the Oseltamivir Powder for Oral Suspension oral suspension well before each use.

#### 4.5 Missed Dose

The missed dose should be taken as soon as remembered, then the regular dosing schedule should be continued. Two doses of Oseltamivir Powder for Oral Suspension should not be taken at the same time.

### 5 OVERDOSAGE

Activated charcoal may be administered to aid in the removal of unabsorbed drug. General supportive measures are recommended.

Reports of overdoses with oseltamivir phosphate have been received from clinical trials and during post-marketing experience. These overdose cases included single or repeated overdose, accidental or intended. The majority of overdose cases were reported in children.

A subset of 114 overdose cases was evaluated in detail, where overdose was defined as receiving more than twice the recommended dose. From the 114 patients, 31% ingested oseltamivir phosphate in an overdose amount ranging from > 2 to < 4 fold the recommended dose (on single or multiple occasions), 49% experienced an overdose between 4 and < 9 fold the recommended dose, and 17% experienced a 9- fold or higher overdose. In the remaining 3% of patients, enough information was not provided to determine the exact amount of overdose.

The most frequently reported adverse events were gastrointestinal disorders, followed by psychiatric and nervous disorders. In the majority of cases reporting overdose, no adverse events were reported. Adverse events reported following overdose were similar in nature and distribution to those observed with therapeutic doses of oseltamivir phosphate, described in [8 ADVERSE REACTIONS](#).

For management of a suspected drug overdose, contact your regional poison control centre.

### 6 DOSAGE FORMS, STRENGTHS, COMPOSITION AND PACKAGING

**Table 1 – Dosage Forms, Strengths, Composition and Packaging**



Route of Administration	Dosage Form / Strength/Composition	Non-medicinal Ingredients
Oral	Powder for Oral Suspension / 6 mg/mL oseltamivir (as oseltamivir phosphate) when reconstituted	Monosodium citrate, saccharin sodium, sodium benzoate, sorbitol, titanium dioxide, tutti-frutti flavouring and xanthan gum.

### **Oseltamivir Powder for Oral Suspension (6 mg/mL)**

Oseltamivir Powder for Oral Suspension contains 360 mg oseltamivir as oseltamivir phosphate per bottle, which when reconstituted contains 60 mL of suspension at a final concentration of 6 mg/mL oseltamivir.

A bottle of 13 g Oseltamivir Powder for Oral Suspension contains 11.345 g of sorbitol. One dose of 75 mg oseltamivir administered twice daily delivers 2.364 g of sorbitol which is unsuitable for subjects with hereditary fructose intolerance (see 7 WARNINGS AND PRECAUTIONS, Endocrine and Metabolism).

Oseltamivir Powder for Oral Suspension is available as a white powder blend for reconstitution to a white tutti-frutti-flavoured suspension. Available in 100 mL glass bottles with a bottle adapter and 1 oral dispenser. Net contents after reconstitution: 60 mL containing oseltamivir phosphate equivalent to 360 mg oseltamivir base.

## **7 WARNINGS AND PRECAUTIONS**

### **General**

No increased efficacy was demonstrated in adult subjects receiving 150 mg oseltamivir phosphate twice daily for 5 days compared to those receiving 75 mg twice daily for the treatment of influenza.

There is no evidence for efficacy of oseltamivir phosphate in any illness caused by agents other than influenza viruses Types A and B. Data on treatment of influenza B are limited.

Efficacy of oseltamivir phosphate in patients who begin treatment after 48 hours of symptoms has not been established.

Efficacy of oseltamivir phosphate in the treatment of subjects with chronic cardiac disease and/or respiratory disease has not been established. No difference in the incidence of complications was observed between the treatment and placebo groups in this population. No information is available regarding treatment of influenza in patients with any medical condition sufficiently severe or unstable to be considered at imminent risk of requiring hospitalization.

Safety and efficacy of repeated treatment or prevention courses have not been studied.

Efficacy of oseltamivir phosphate for treatment or prevention of influenza in immunocompromised patients has not been established.

### **Endocrine and Metabolism**

A bottle of 13 g Oseltamivir Powder for Oral Suspension contains 11.345 g of sorbitol. One dose of 75 mg oseltamivir administered twice daily delivers 2.364 g of sorbitol, which is unsuitable for subjects with hereditary fructose intolerance.

## Hepatic/Biliary/Pancreatic

There have been post-marketing reports of elevated liver enzymes and hepatotoxicity including fulminant hepatitis/hepatic failure, in some cases with fatal outcome, where a causal relationship with oseltamivir could not be excluded, especially in patients with pre-existing liver disease.

The safety, efficacy and pharmacokinetics of oseltamivir in patients with severe hepatic impairment have not been studied (see [4 DOSAGE AND ADMINISTRATION, 4.2 Recommended Dose and Dosage Adjustment, Dosage Adjustment, Hepatic Impairment](#)).

## Neuropsychiatric

There have been post-marketing reports of delirium and self-injury, in some cases resulting in fatal outcomes, in patients with influenza who were receiving oseltamivir phosphate. Because these events were reported voluntarily during clinical practice, estimate of frequency cannot be made but they appear to be uncommon based on oseltamivir phosphate usage data. These events were reported primarily among pediatric patients. The contribution of oseltamivir phosphate to these events has not been established. Patients with influenza should be closely monitored for signs of abnormal behaviour. If neuropsychiatric symptoms occur, the risks and benefits of continuing treatment should be evaluated for each patient (see [8 ADVERSE REACTIONS, 8.5 Post-Market Adverse Drug Reactions, Neurologic, and Psychiatric](#)).

Influenza can be associated with a variety of neurologic and behavioural symptoms, which can include events such as hallucinations, delirium, and abnormal behaviour, in some cases resulting in fatal outcomes. These events may occur in the setting of encephalitis or encephalopathy, but can occur without obvious severe disease.

## Renal

*Renal Impairment:* For dose adjustments in patients with renal impairment (for both treatment and prevention) - see [4 DOSAGE AND ADMINISTRATION, 4.2 Recommended Dose and Dosage Adjustment, Dosage Adjustment, Renal Impairment](#).

The pharmacokinetics of oseltamivir have not been studied in patients with end-stage renal disease (i.e., creatinine clearance of < 10 mL/min) not undergoing dialysis. Hence, dosing recommendation cannot be provided for this group.

## Sensitivity/Resistance

In clinical studies of naturally acquired infection, the cumulative incidence of oseltamivir-resistant virus by phenotyping alone or by phenotyping and genotyping was 0.32% (4/1245) or 0.4% (5/1245) respectively in adult/adolescent patients. In children with naturally acquired influenza virus infection, resistance was determined in 6 clinical studies WV15731 (0%; 0/5), WV15758 (8%; 15/183), WV15759/WV15871 (0%, 0/60), JV16284 (19%, 8/43), WV16193 (0%, 0/147), NV16871 (8%, 2/26). From the data obtained in these studies, the cumulative incidence of oseltamivir resistance in pediatric patients aged 1 to 12 years was 4.1% (19/464) based on phenotyping and 5.4% (25/464) based on phenotyping and genotyping (full genotyping was not performed on all patients). The patients cleared the virus normally and showed no clinical deterioration.

There has been no evidence for emergence of drug resistance associated with the use of oseltamivir phosphate in clinical studies conducted to date in post-exposure (7 days), post-exposure within the household groups (10 days) and seasonal (42 days) prophylaxis of influenza in immunocompetent persons (see [15 MICROBIOLOGY, Resistance](#)).

Insufficient information is available to fully characterize the risk of emergence of resistance to oseltamivir phosphate in clinical use (see [15 MICROBIOLOGY, Resistance](#)).

## Skin and Hypersensitivity Reactions

Severe skin and hypersensitivity reactions have been reported since marketing in patients treated with oseltamivir phosphate (see [8 ADVERSE REACTIONS, 8.5 Post-Market Adverse Reactions](#)).

### 7.1 Special Populations

#### 7.1.1 Pregnant Women

Oseltamivir Powder for Oral Suspension should be used in pregnancy only if the potential benefits outweigh the potential risks to the fetus, taking into consideration the pathogenicity of the circulating influenza virus strain and the underlying condition of the pregnant woman.

A large amount of data from pregnant women exposed to oseltamivir (more than 1000 exposed outcomes during the first trimester) from post-marketing reports and observational studies indicate no direct or indirect harmful effects with respect to pregnancy. Results from a pooled pharmacokinetic analysis indicate a lower exposure to the active metabolite in pregnant women compared to non-pregnant women; however, this predicted exposure is expected to have clinical benefits and there are insufficient pharmacokinetic and safety data to recommend a dose adjustment in pregnant women who are justified to use Oseltamivir Powder for Oral Suspension.

Studies for effects on embryo-fetal development were conducted in rats (50, 250 and 1500 mg/kg/day) and rabbits (50, 150 and 500 mg/kg/day) by the oral route. Relative exposures at these doses were, respectively, 2, 13 and 100 times human exposure in the rat and 4, 8 and 50 times human exposure in the rabbit. Pharmacokinetic studies indicated that fetal exposure was seen in both species. In the rat study, minimal maternal toxicity was reported in the 1500 mg/kg/day group. In the rabbit study, slight and marked maternal toxicities were observed, respectively, in the 150 and 500 mg/kg/day groups. An increased incidence of abortion was seen in the 500 mg/kg/day group. There was a dose-dependent increase in the incidence rates of a variety of minor skeletal individual abnormalities and variants in the exposed offspring in these studies. However, the individual incidence rate of each skeletal abnormality or variant remained within the background rates of occurrence in the species studied (see [16 NON-CLINICAL TOXICOLOGY](#)).

#### 7.1.2 Breast-feeding

Very limited information is available on children breast-fed by mothers taking oseltamivir and on excretion of oseltamivir in breast milk. In lactating rats, oseltamivir and the active metabolite are excreted in milk. Limited data demonstrated that oseltamivir and the active metabolite were detected in breast milk, however the levels were low, which would result in a sub-therapeutic dose to the infant. Based on this information, and the pathogenicity of the circulating influenza virus strain, administration of oseltamivir may be considered where the potential benefit to the lactating mother justifies the potential risk to the nursing infant.

#### 7.1.3 Pediatrics

**Pediatrics (< 1 year of age):** Oseltamivir Powder for Oral Suspension should not be used in children under 1 year of age (see [16 NON-CLINICAL TOXICOLOGY, General Toxicology, Multiple Dose Toxicity](#)). The efficacy of oseltamivir phosphate in infants younger than 1 year of age have not been established (see [14 CLINICAL TRIALS](#)).

#### 7.1.4 Geriatrics

**Geriatrics (≥ 65 years of age):** Efficacy of oseltamivir phosphate in the treatment of elderly patients has not been evaluated. Safety data in 372 elderly patients (≥ 65 years old) showed no overall difference between these subjects and younger adults. Based on drug exposure and tolerability, dosage adjustments are not anticipated for elderly patients (see [10 CLINICAL PHARMACOLOGY, 10.3 Pharmacokinetics, Special Populations and Conditions, Geriatrics](#)).

Safety has been demonstrated in elderly residents of nursing homes who took oseltamivir phosphate for the prevention of influenza. Many of these individuals had cardiac and/or respiratory disease, and most had received vaccine that season (see [14 CLINICAL TRIALS](#)).

### 8 ADVERSE REACTIONS

#### 8.1 Adverse Reaction Overview

In adult/adolescent treatment studies with oseltamivir phosphate, the most frequently reported adverse events were nausea and vomiting. In the prevention studies, adverse events were qualitatively very similar to those seen in the treatment studies. In the pediatric treatment and prophylaxis studies, the most frequently reported adverse event was vomiting.

#### 8.2 Clinical Trial Adverse Reactions

Clinical trials are conducted under very specific conditions. The adverse reaction rates observed in the clinical trials; therefore, may not reflect the rates observed in practice and should not be compared to the rates in the clinical trials of another drug. Adverse reaction information from clinical trials may be useful in identifying and approximating rates of adverse drug reactions in real-world use.

##### Adult Treatment Studies

In a total of 4624 patients, including patients on placebo and 75 mg b.i.d. oseltamivir phosphate, in adult/adolescent Phase III studies in the treatment of influenza, the most frequently reported adverse events were nausea and vomiting. These events were transient and generally occurred with first dosing. These events did not lead to patient discontinuation of study drug in the vast majority of instances. At the recommended dose of 75 mg twice daily, the most frequently reported adverse events that led to patient withdrawal were nausea and vomiting.

In adult/adolescent Phase III treatment studies, some adverse events occurred more frequently in patients taking oseltamivir phosphate compared to those taking placebo. The adverse events that occurred with an incidence of ≥ 1% at the recommended dose, either for treatment or prophylaxis, are shown in Table 2. This summary includes healthy young adults and “at risk” patients (patients at higher risk of developing complications associated with influenza e.g. elderly patients and patients with chronic cardiac or respiratory disease). Those events with an incidence of ≥ 1% and which were reported more frequently in patients taking oseltamivir phosphate compared with placebo, irrespective of causality, were nausea, vomiting, abdominal pain and headache.

**Table 2 - Most Frequent Adverse Events in Studies in Naturally Acquired Influenza for Adults/Adolescents for Treatment or Prophylaxis**

	Treatment*	Prevention
--	------------	------------

<b>Adverse Events System Organ Class (MedDRA)</b>	<b>Oseltamivir phosphate 75 mg twice daily for 5 days N=2647</b>		<b>Placebo N=1977</b>		<b>Oseltamivir phosphate 75 mg once daily N=1945</b>		<b>Placebo N=1588</b>	
<b>Ear and Labyrinth Disorders</b>								
Dizziness (incl. Vertigo)*	54	(2.0%)	49	(2.5%)	28	(1.4%)	24	(1.5%)
<b>Gastrointestinal Disorders</b>								
Nausea (without vomiting)	263	(9.9%)	118	(6.0%)	156	(8.0%)	66	(4.2%)
Vomiting	205	(7.7%)	63	(3.2%)	46	(2.4%)	19	(1.2%)
Diarrhea*	149	(5.6%)	147	(7.4%)	66	(3.4%)	66	(4.2%)
Abdominal pain (incl. upper abdominal pain)*	62	(2.3%)	51	(2.6%)	47	(2.4%)	31	(2.0%)
Dyspepsia*	12	(0.5%)	9	(0.5%)	29	(1.5%)	23	(1.4%)
<b>General Disorders and Administration Site Reactions</b>								
Fatigue*	14	(0.5%)	9	(0.5%)	142	(7.3%)	107	(6.7%)
Pain*	8	(0.3%)	5	(0.3%)	70	(3.6%)	43	(2.7%)
Pyrexia*	6	(0.2%)	9	(0.5%)	33	(1.7%)	33	(2.1%)
Pain in limb*	5	(0.2%)	1	(0.1%)	20	(1.0%)	5	(0.3%)
Influenza like illness*	–	–	1	(0.1%)	21	(1.1%)	24	(1.5%)
<b>Infections and Infestations</b>								
Bronchitis	70	(2.6%)	72	(3.6%)	10	(0.5%)	14	(0.9%)
Sinusitis*	35	(1.3%)	22	(1.1%)	15	(0.8%)	12	(0.8%)
Herpes simplex*	27	(1.0%)	23	(1.2%)	11	(0.1%)	9	(0.6%)
Nasopharyngitis*	4	(0.2%)	2	(0.1%)	80	(4.1%)	68	(4.3%)
Upper respiratory tract infections*	5	(0.2%)	1	(0.1%)	60	(3.1%)	51	(3.2%)
Influenza*	–	–	–	–	46	(2.4%)	41	(2.6%)
<b>Nervous System Disorders</b>								
Headache	45	(1.7%)	27	(1.4%)	335	(17.2%)	260	(16.4%)
Insomnia*	31	(1.2%)	17	(0.9%)	22	(1.1%)	14	(0.9%)
<b>Respiratory, Thoracic and Mediastinal Disorders</b>								
Cough*	42	(1.6%)	38	(1.9%)	94	(4.8%)	90	(5.7%)
Nasal congestion*	27	(1.0%)	22	(1.1%)	134	(6.9%)	113	(7.1%)
Sore throat*	25	(0.9%)	14	(0.7%)	100	(5.1%)	86	(5.4%)

Rhinorrhea*	6	(0.02%)	5	(0.3%)	29	(1.5%)	19	(1.2%)
<b>Musculoskeletal, Connective Tissue and Bone Disorders</b>								
Back pain*	13	(0.5%)	9	(0.5%)	41	(2.1%)	41	(2.6%)
Arthralgia*	4	(0.2%)	3	(0.2%)	28	(1.4%)	39	(2.5%)
Myalgia*	7	(0.3%)	4	(0.2%)	20	(1.0%)	21	(1.3%)
<b>Disorders of Reproductive System and Breast</b>								
Dysmenorrhea*	–	–	–	–	53	(2.7%)	51	(3.2%)

\* These events may be related to the underlying influenza infection because they occurred either more frequently in patients on placebo compared to patients on oseltamivir phosphate, or the difference in frequency between the two arms was less than 1%.

### Adult Prevention Studies

A total of 3533 subjects (adolescents, healthy adults and elderly) participated in 3 Phase III prevention studies, of whom 1480 received the recommended dose of 75 mg once daily. Adverse events were qualitatively similar to those seen in the treatment studies (see Table 2). There were no clinically relevant differences in the safety profile of the 942 subjects 65 years of age and older, who received oseltamivir phosphate or placebo, compared with the younger population (aged up to 65 years).

In another study, an additional 399 subjects received 75 mg of oseltamivir phosphate once daily for 10 days following the identification of a household index case. Similar to previous studies, nausea (8.3%), vomiting (4.5%), diarrhea (0.8%) and headache (7.8%) were among the most commonly reported adverse events.

### 8.2.1 Clinical Trial Adverse Reactions – Pediatrics

#### Pediatric Treatment Studies

A total of 1480 pediatric patients aged 1 to 12 years (including 698 otherwise healthy pediatric patients aged 1 to 12 and 334 asthmatic pediatric patients aged 6 to 12) participated in clinical studies of oseltamivir phosphate given for the treatment of influenza. A total of 858 pediatric patients received treatment with oseltamivir phosphate oral suspension.

Adverse events occurring in  $\geq 1\%$  of pediatric patients receiving oseltamivir phosphate are listed in Table 3. The most frequently reported adverse event was vomiting. Other events reported more frequently by pediatric patients treated with oseltamivir phosphate included abdominal pain, epistaxis, ear disorder and conjunctivitis.

These events generally occurred once and resolved despite continued dosing. They did not cause discontinuation of drug in the vast majority of cases.

The adverse event profile in adolescents is similar to that described for adult patients and pediatric patients aged 1 to 12 years.

### Table 3 - Most Frequent Adverse Events Occurring in Children Aged 1 to 12 Years in Studies in Naturally Acquired Influenza

Adverse Events System Organ Class (MedDRA)	Treatment <sup>a</sup>		Prophylaxis <sup>b</sup>	
	Oseltamivir phosphate 2 mg/kg twice daily N= 858	Placebo  N= 622	Prophylaxis with Oseltamivir phosphate <sup>c</sup>  N=148	No Prophylaxis <sup>b</sup>  N=87
<b>Nervous System Disorders</b>				
Headaches	3 (0.3%)	5 (0.8%)	5 (3.4%)	1 (1.1%)
<b>Blood and Lymphatic System Disorders</b>				
Lymphadenopathy*	5 (0.6%)	8 (1.3%)	–	–
<b>Ear and Labyrinth Disorders</b>				
Earache*	10 (1.2%)	4 (0.6%)	1 (0.7%)	–
Tympanic membrane disorder*	4 (0.5%)	6 (1.0%)	–	–
<b>Eye Disorders</b>				
Conjunctivitis (incl. red eyes, eye discharge and eye pain)*	9 (1.0%)	2 (0.3%)	–	–
<b>Gastrointestinal Disorders</b>				
Vomiting	140 (16.3%)	51 (8.2%)	12 (8.1%)	2 (2.3%)
Diarrhea*	63 (7.3%)	49 (7.9%)	1 (0.7%)	–
Abdominal pain (incl. upper abdominal pain)*	29 (3.4%)	21 (3.4%)	3 (2.0%)	–
Nausea	32 (3.7%)	27 (4.3%)	6 (4.1%)	1 (1.1%)
Dyspepsia*	2 (0.2%)	–	3 (2.0%)	–
<b>Infections and Infestations</b>				
Otitis media*	43 (5.0%)	51 (8.2%)	3 (2.0%)	2 (2.3%)
Pneumonia*	29 (3.4%)	19 (3.1%)	–	1 (1.1%)
Bronchitis*	14 (1.6%)	15 (2.4%)	–	2 (2.3%)
Sinusitis*	11 (1.3%)	13 (2.1%)	1 (0.7%)	–
Nasopharyngitis*	2 (0.2%)	2 (0.3%)	2 (1.4%)	4 (4.6%)
Upper respiratory tract infection*	3 (0.3%)	3 (0.5%)	2 (1.4%)	3 (3.4%)
<b>Respiratory, Thoracic and Mediastinal Disorders</b>				
Cough*	8 (0.9%)	4 (0.6%)	18 (12.2%)	23 (26.4%)
Nasal congestion*	3 (0.3%)	–	16 (10.8%)	17 (19.5%)

Adverse Events System Organ Class (MedDRA)	Treatment <sup>a</sup>		Prophylaxis <sup>b</sup>	
	Oseltamivir phosphate 2 mg/kg twice daily N= 858	Placebo  N= 622	Prophylaxis with Oseltamivir phosphate <sup>c</sup>  N=148	No Prophylaxis <sup>b</sup>  N=87
Asthma (including aggravated)	22 (2.6%)	25 (4.0%)	2 (1.4%)	1 (1.1%)
Epistaxis*	18 (2.1%)	14 (2.3%)	1 (0.7%)	–
Rhinorrhoea*	2 (0.2%)	2 (0.3%)	2 (1.4%)	1 (1.1%)
<b>Skin and Subcutaneous Tissue Disorders</b>				
Dermatitis*	9 (1.0%)	11 (1.8%)	–	–
<b>General Disorders and Administration Site Reactions</b>				
Pyrexia*	2 (0.2%)	2 (0.3%)	3 (2.0%)	6 (6.9%)

<sup>a</sup> Pooled data from Phase III trials of oseltamivir phosphate treatment of naturally acquired influenza.

<sup>b</sup> Subjects participating in an uncontrolled household transmission study (in which household contacts received either oseltamivir phosphate for prophylaxis (once-daily dosing for 10 days) or no prophylaxis but oseltamivir phosphate treatment if they became ill), who remained on no prophylaxis and did not receive treatment with oseltamivir phosphate.

<sup>c</sup> Pooled data from pediatric prophylaxis studies. Unit dose = age-based dosing (see 4 DOSAGE AND ADMINISTRATION).

\* These events may be related to the underlying influenza infection because they occurred either more frequently in patients on placebo compared to patients on oseltamivir phosphate, or the difference in frequency between the two arms was less than 1%.

### Pediatric Prevention Studies

Amongst the 148 pediatric patients aged 1 to 12 years who received the recommended dose of oseltamivir phosphate once daily, in a post-exposure prophylaxis study in households (N=99), and in a separate 6 week pediatric prophylaxis study (n=49), gastrointestinal events, particularly vomiting was the most frequently reported adverse event. Other events reported more frequently by pediatric patients who received oseltamivir phosphate for prophylaxis are nausea, abdominal pain, dyspepsia and earaches. Oseltamivir phosphate was well tolerated in these studies and the adverse events noted were consistent with those previously observed in pediatric treatment studies (see Table 3).

**Immunocompromised Patients:** In a 12-week prophylaxis study in 475 immunocompromised subjects, including 18 children 1-12 years of age, the safety profile in the 238 subjects receiving oseltamivir phosphate was consistent with that previously observed in oseltamivir phosphate prophylaxis clinical trials.



### 8.3 Less Common Clinical Trial Adverse Reactions

#### Adult Treatment Studies

Additional adverse events occurring in < 1% of patients receiving oseltamivir phosphate for treatment included unstable angina, anemia, pseudomembranous colitis, humerus fracture, pneumonia, pyrexia, and peritonsillar abscess.

### 8.4 Abnormal Laboratory Findings: Hematologic, Clinical Chemistry and Other Quantitative Data

Not applicable.

### 8.5 Post-Market Adverse Reactions

The following adverse reactions have been identified during post-marketing use of oseltamivir phosphate. Because these reactions are reported voluntarily from a population of uncertain size, it is not possible to reliably estimate their frequency or establish a causal relationship to oseltamivir phosphate exposure.

**Skin and hypersensitivity reactions:** dermatitis, rash, eczema, urticaria, erythema multiforme, Stevens-Johnson-Syndrome, toxic epidermal necrolysis, anaphylactic/anaphylactoid reactions and face edema.

**Hepatobiliary disorders:** elevated liver enzymes, hepatotoxicity including fulminant hepatitis/hepatic failure, in some cases with fatal outcome.

**Gastro-intestinal disorders:** gastro-intestinal bleeding, hemorrhagic colitis.

**Cardiac disorders:** arrhythmia.

**Neurologic:** seizure.

**Metabolism and nutrition disorders:** aggravation of diabetes.

**Psychiatric:** delirium, including symptoms such as altered level of consciousness, confusion, abnormal behavior leading to self-injury, delusions, hallucinations, agitation, anxiety, nightmares (see 7 WARNINGS AND PRECAUTIONS).

## 9 DRUG INTERACTIONS

### 9.2 Drug Interactions Overview

Oseltamivir is extensively converted to oseltamivir carboxylate by esterases, located predominantly in the liver. Drug interactions involving competition for esterases have not been extensively reported in literature. Low protein binding of oseltamivir and oseltamivir carboxylate suggests that the probability of drug displacement interactions is low.

*In vitro* studies demonstrated that neither oseltamivir nor the active metabolite are good substrates for P450 mixed-function oxidases or for glucuronyl transferases.

### 9.3 Drug-Behavioural Interactions

No drug-behavioural studies have been performed with oseltamivir phosphate.

### 9.4 Drug-Drug Interactions

Cimetidine, a non-specific inhibitor of cytochrome P450 isoforms and competitor for renal tubular secretion of basic or cationic drugs, has no effect on plasma levels of oseltamivir or its active metabolite.

Clinically important drug interactions involving competition for renal tubular secretion are unlikely due to the known safety margin for most of these drugs, the elimination characteristics of the active metabolite (glomerular filtration and anionic tubular secretion) and the excretion capacity of these pathways. Co-administration of probenecid results in an approximate two-fold increase in exposure to the active metabolite due to a decrease in active anionic tubular secretion in the kidney. However, due to the wide safety margin of the active metabolite, no dose adjustments are required when co-administering with probenecid. Other drugs excreted via anionic tubular secretion have not been evaluated.

Co-administration with amoxicillin does not alter plasma levels of either compound, indicating that competition for the anionic secretion pathway is weak.

In six subjects, co-administration with acetaminophen did not alter plasma levels of oseltamivir, its active metabolite, or acetaminophen.

Co-administration with paracetamol does not alter plasma levels of oseltamivir, its active metabolite, or paracetamol.

No pharmacokinetic interactions between oseltamivir or its major metabolite have been observed when co-administering oseltamivir with paracetamol, acetyl-salicylic acid, warfarin, rimantadine, amantadine, cimetidine or with antacids (magnesium and aluminum hydroxides and calcium carbonates).

In Phase III treatment and prophylaxis clinical studies, oseltamivir phosphate has been administered with commonly used drugs such as ACE inhibitors (enalapril, captopril), thiazide diuretics (bendrofluazide), antibiotics (penicillin), H<sub>2</sub>-receptor blockers (cimetidine), and analgesic agents (acetylsalicylic acid, ibuprofen and paracetamol). No change in adverse event profile or frequency has been observed as a result of co-administration of oseltamivir phosphate with these compounds.

## **9.5 Drug-Food Interactions**

Interactions with food have not been established.

## **9.6 Drug-Herb Interactions**

Interactions with herbal products have not been established.

## **9.7 Drug-Laboratory Test Interactions**

Interactions with laboratory tests have not been established.

## 10 CLINICAL PHARMACOLOGY

### Animal Pharmacology

Oseltamivir phosphate produced effects in the non-clinical safety pharmacology studies only at oral doses well in excess of any clinically relevant therapeutic levels. These effects in the rat, were reduced gastrointestinal transit and gastric emptying at 1000 mg/kg. In the rodent toxicology studies these effects were not reported as any sign of gastrointestinal disturbance. Additionally, there were increases in excretion of electrolytes at 100 and 1000 mg/kg and increased urine production at 1000 mg/kg. Increases in electrolyte excretion were reported in a 27-week rat toxicology study at 1000 mg/kg/day and attributed to a high phosphate intake due to the salt of the test material. In the same study less pronounced effects were seen at 200 mg/kg/day, while in another rat toxicology study no significant effects were seen at 100 mg/kg/day. A statistically significant increase in response to a painful stimulus was seen but this was neither time nor dose related and therefore not thought to be of pharmacological significance.

The intravenous infusion of the active metabolite at 2, 15 and 100 mg/kg cumulatively produced statistically significant changes in heart rate, QT and QTc interval, QRS duration and pCO<sub>2</sub> when compared with time matched controls in the anaesthetized dog. The effects on heart rate and pCO<sub>2</sub> were at isolated time points and the decrease in QRS duration was not accompanied by any other relevant physiological changes and so not likely to be due to the drug treatment. The statistical differences in QT and QTc interval between the active metabolite and vehicle treated groups was seen in the predose and just after the start of the infusion suggesting no pharmacological significance. Because of this, significance was tested for by comparing percentage changes from predose values within the active metabolite treatment group. There were no significant differences detected. However, on comparison with the absolute values significant differences were seen during the infusion of the 100 mg/kg/dose. To clarify this situation a further test was performed in an isolated sheep Purkinje fibre study where no significant effects were observed on cardiac action potential parameters. Other than these findings, no additional effects were seen on the cardiovascular and respiratory dynamics of the anaesthetized dog.

In conclusion, oseltamivir phosphate produced significant pharmacological effects only at doses much greater than would be of clinical relevance. It is therefore concluded that oseltamivir phosphate and the active metabolite produced no clinically relevant pharmacological effects on the central nervous, cardiovascular, respiratory, gastrointestinal, smooth muscle, renal, hepatic and immune systems tested.

### Human Pharmacology

*QT/QTc:* A retrospective analysis of ECGs from 8 clinical pharmacology studies (n=182 subjects including 30 placebo) concluded that oseltamivir phosphate does not cause prolongation of QT intervals in humans. Although some individuals were found to have some alterations in QTc measurements, none were of clinical significance and the frequency was similar among placebo and subjects treated with oseltamivir phosphate.

In a study on ECG intervals in which healthy volunteers received daily doses of either 75, 225 or 450 mg oseltamivir phosphate b.i.d. orally for 5 days, treatment with oseltamivir phosphate had no impact on any ECG parameters.

#### 10.1 Mechanism of Action

Oseltamivir phosphate is an ethyl ester prodrug requiring ester hydrolysis for conversion to the active metabolite, oseltamivir carboxylate. The active metabolite is a selective inhibitor of influenza A and B virus neuraminidase enzymes which are glycoproteins found on the virion surface. Viral neuraminidase

is primarily important for the release of recently formed virus particles from infected cells and the further spread of infectious virus in the body. The proposed mechanism of action of oseltamivir is via inhibition of influenza virus neuraminidase with the possibility of alteration of virus particle aggregation and release. It has also been suggested that neuraminidase can play a role in viral entry into uninfected cells.

Oseltamivir is readily absorbed after oral administration and converted by hepatic esterases to its active metabolite. The mean volume of distribution ( $V_{ss}$ ) of the active metabolite is approximately 23 L. The active metabolite is not further metabolized and is eliminated in the urine. The half-life of elimination of this metabolite is 6 to 10 hours. Renal clearance (18.8 L/h) exceeds glomerular filtration rate (7.5 L/h), indicating that tubular secretion in addition to glomerular filtration occurs. Also, the prodrug which reaches the systemic circulation (less than 5%) is eliminated by renal excretion. The binding of oseltamivir to human plasma protein is 42% and that of the active metabolite is negligible, approximately 3%.

Exposure to the active metabolite is inversely proportional to declining renal function.

## **10.2 Pharmacodynamics**

Refer to Section 10.1 Mechanism of Action. Additional information in the Product Monograph not included at the time of authorization.

## **10.3 Pharmacokinetics**

### **Absorption**

Oseltamivir is readily absorbed from the gastrointestinal tract after oral administration of oseltamivir phosphate and is extensively converted predominantly by hepatic esterases to the active metabolite. At least 75% of an oral dose reaches the systemic circulation as the active metabolite. Exposure to the prodrug is less than 5% relative to the active metabolite. Plasma concentrations of active metabolite are proportional to dose and are not significantly affected by co-administration with food (see 4 DOSAGE AND ADMINISTRATION).

### **Distribution**

The mean volume of distribution ( $V_{ss}$ ) of the active metabolite is approximately 23 litres in humans.

The binding of oseltamivir to human plasma protein is 42% and that of the active metabolite is negligible, approximately 3%.

### **Metabolism**

Oseltamivir is extensively converted to the active metabolite by esterases located predominantly in the liver. Neither oseltamivir nor the active metabolite are substrates for, or inhibitors of, cytochrome P450 isoforms.

### **Elimination**

Absorbed oseltamivir is primarily (> 90%) eliminated by conversion to the active metabolite. The active metabolite is not further metabolized and is eliminated in the urine. Peak plasma concentrations of the active metabolite decline with a half-life of 6 to 10 hours in most subjects. The active drug is eliminated entirely (> 99%) by renal excretion. Renal clearance (18.8 L/h) exceeds glomerular filtration rate (7.5 L/h), indicating that tubular secretion in addition to glomerular filtration occurs. Less than 20% of an oral radiolabelled dose is eliminated in feces.

## Special Populations and Conditions

- **Pediatrics:** The pharmacokinetics of oseltamivir have been evaluated in single dose pharmacokinetic studies in pediatric patients aged 1 to 16 years. Multiple dose pharmacokinetics were studied in a small number of pediatric patients aged 3-12 years enrolled in a clinical trial. Younger pediatric patients cleared both the prodrug and the active metabolite faster than adults resulting in lower exposure for a given mg/kg dose. Doses of 2 mg/kg give oseltamivir carboxylate exposures comparable to those achieved in adults receiving a single 75 mg capsule dose (approximately 1 mg/kg). The pharmacokinetics of oseltamivir in pediatric patients over 12 years of age is similar to those in adults.

Oseltamivir Powder for Oral Suspension should not be used in children under 1 year of age (see 16 NON-CLINICAL TOXICOLOGY, General Toxicology, Multiple Dose Toxicity).

- **Geriatrics:** Exposure to the active metabolite at steady-state was 25% to 35% higher in elderly patients (age range 65 to 78) compared to young adults given comparable doses of oseltamivir phosphate. Half-lives observed in the elderly patients were similar to those seen in young adults. On the basis of drug exposure and tolerability, dosage adjustments are not required for elderly patients for either treatment or prevention (see 4 DOSAGE AND ADMINISTRATION, 4.2 Recommended Dose and Dosage Adjustment, Dosage Adjustment).
- **Hepatic Insufficiency:** The safety, efficacy and pharmacokinetics of oseltamivir phosphate in patients with severe hepatic impairment have not been studied (see 7 WARNINGS AND PRECAUTIONS, Hepatic/Biliary/Pancreatic). In a clinical study of adult patients with moderate hepatic impairment (N=11), compared with healthy volunteers (N=23), metabolic conversion of oseltamivir into the active metabolite oseltamivir carboxylate was not significantly altered (see 4 DOSAGE AND ADMINISTRATION, 4.2 Recommended Dose and Dosage Adjustment, Dosage Adjustment, Hepatic Impairment). No studies have been carried out in pediatric patients with hepatic impairment.
- **Renal Insufficiency:** Administration of 100 mg of oseltamivir phosphate twice daily for five days to patients with various degrees of renal impairment showed that exposure to the active metabolite is inversely proportional to declining renal function. For dosage information, see 4 DOSAGE AND ADMINISTRATIONS, Dosage Adjustment.

## 11 STORAGE, STABILITY AND DISPOSAL

**Oseltamivir Powder for Oral Suspension:** Store dry powder at 15-30°C. Store reconstituted suspension:

- a) At room temperature (15-30°C). Discard unused portion within 10 days of reconstitution.
- or
- b) In a refrigerator (2-8°C). Discard unused portion within 17 days of reconstitution.

Do not freeze reconstituted suspension.

### Disposal of unused/expired medicines

The release of pharmaceuticals in the environment should be minimized. Medicines should not be disposed of via wastewater and disposal through household waste should be avoided.

Any unused medicinal product or waste materials should be disposed of in accordance with local requirements.

## **12 SPECIAL HANDLING INSTRUCTIONS**


Not applicable.

## PART II: SCIENTIFIC INFORMATION

### 13 PHARMACEUTICAL INFORMATION

#### Drug Substance

Proper Name:	oseltamivir phosphate
Chemical name:	Ethyl (3R, 4R, 5S)-4-acetamino-5-amino-3-(1-ethylpropoxy)-1-cyclohexene-1-carboxylate, phosphate (1:1)
Molecular formula and molecular mass:	$C_{16}H_{28}N_2O_4 \cdot H_3PO_4$ 410.4 g/mol for oseltamivir phosphate salt
Structural formula:	



Physicochemical properties:	<ul style="list-style-type: none"><li>• Physical Form: White to off-white powder</li><li>• Solubility: Freely soluble in water and methanol</li><li>• pKa and pH values: pKa: 7.75</li><li>• Melting Point: 195-205°C with degradation</li></ul>
-----------------------------	--

## 14 CLINICAL TRIALS

### 14.1 Clinical Trials by Indication

#### Treatment of Influenza

**Adult Patients:** Phase III clinical trials evaluated the safety and efficacy of oseltamivir phosphate for the treatment of naturally occurring influenza during a period when influenza virus was known to be circulating in the community. A total of 1418 patients received any treatment (oseltamivir phosphate or placebo) of whom 476 patients received 75 mg oseltamivir phosphate twice daily for 5 days. Patients started treatment with oseltamivir phosphate within 40 hours after reported onset of symptoms. The primary efficacy parameter was the time to alleviation of all symptoms. The population used in the primary analyses was the intent-to-treat-infected (ITTI) population. This population included only subjects who received at least one dose of study treatment and who had laboratory confirmed influenza. An intent-to-treat (ITT) population included all subjects who took at least one dose of study medication, regardless of whether they proved to have influenza.

The results for two pivotal studies (WV15670 and WV15671) are shown in Table 4.

**Table 4 - Median Time (Hours) to Alleviation of All Symptoms in the ITTI and ITT Populations**

Study	Population	Placebo (95% CI)	Oseltamivir Phosphate 75 mg twice daily (95% CI)	p-value*
WV15670	ITTI	n=161 116.5 (101.5 to 137.8)	n=157 87.4 (73.3 to 104.7)	0.017
	ITT	n=235 116.1 (99.8 to 129.5)	n=240 97.6 (79.1 to 115.3)	0.051
WV15671	ITTI	n=128 103.3 (92.6 to 118.7)	n=121 71.5 (60.0 to 83.2)	< 0.0001
	ITT	n=200 97.0 (86.3 to 113.6)	n=204 76.3 (66.3 to 89.2)	0.004

ITT intent-to-treat

ITTI intent-to-treat infected

\* difference between medians

Treatment with oseltamivir phosphate significantly reduced the duration by 1.3 days, of clinically relevant symptoms of influenza. The seven symptoms assessed were: feverish feeling, muscle aches or myalgia, headache, sore throat, cough, overall discomfort, and nasal stuffiness or runny nose.

**Pediatric Patients:** One double-blind placebo controlled treatment trial was conducted in pediatric patients, aged 1 to 12 years (mean age 5.3), who had fever (> 100°F) plus one respiratory symptom (cough or coryza) when influenza virus was known to be circulating in the community. In this study, 67% of influenza-infected patients were infected with influenza A and 33% with influenza B.



Treatment with oseltamivir phosphate, started within 48 hours of onset of symptoms, significantly reduced the duration of illness by 1.5 days compared to placebo. Duration of illness was defined as time to alleviation of cough, alleviation of coryza, resolution of fever, and return to normal health and activity.

### **Prevention of Influenza**

**Adult Patients:** The efficacy of oseltamivir phosphate in preventing naturally occurring influenza illness has been proven in four separate trials which are summarized below and in Table 5.

In a Phase III study (WV15799) in adult and adolescent contacts of a household case of influenza, administered 75 mg of oseltamivir phosphate once daily, starting within 2 days of onset of symptoms in the household case and continued for 7 days, significantly reduced the incidence of influenza illness occurring in the contacts by 92% (p value = < 0.001).

In a double blind placebo controlled study (WV15673) conducted in unvaccinated otherwise healthy adults 18-65 years of age, 75 mg of oseltamivir phosphate administered once daily significantly reduced the incidence of clinical influenza by 76% (p value = 0.0006) during a community outbreak of influenza. The subjects in this study received oseltamivir phosphate for a period of 42 days. No additional benefit was demonstrated in this study using 75 mg of oseltamivir phosphate twice daily.

In a double blind placebo controlled study (WV15825) in elderly residents of nursing homes, many of whom had chronic cardiac disease and/or respiratory disease, 80% received vaccine in the season of the study. The vaccine was a good match for circulating strains. The administration of 75 mg of oseltamivir phosphate once daily significantly reduced the incidence of clinical influenza illness in these patients by 92% (p value = 0.0015). In the same study, oseltamivir phosphate significantly reduced the incidence of influenza associated bronchitis, pneumonia and sinusitis by 86% (p value = 0.037). The subjects in this study received oseltamivir phosphate for a period of 42 days.

In all three of these clinical trials (WV15799, WV15673 and WV15825), approximately 1% of subjects taking oseltamivir phosphate for prevention developed influenza during the dosing period.

In a fourth Phase III study (WV16193), it was demonstrated that oseltamivir phosphate effectively prevents the secondary spread of influenza within households. In this study, the index case was treated with oseltamivir phosphate and household contacts were randomized (by household) to receive either prophylaxis (P) with oseltamivir phosphate or treatment (T) with oseltamivir phosphate upon emergence of influenza-like illness. In households with infected index cases where subjects who were already shedding virus at baseline were excluded (ITTIINAB population) there was a 78.8% (p = 0.0008) reduction in the incidence of laboratory-confirmed influenza in P versus T. Amongst contacts, the outcome was analogous to that seen for households with a significantly lower number of infected contacts in P versus T (84.5% reduction, p = 0.0002, ITTIINAB population). No virus shedding was detected in any subject in the prophylaxis group while 7% of contact in the treatment group (ITTIINAB) shed virus.

Oseltamivir phosphate also significantly reduced the incidence of virus shedding and successfully prevented virus transmission in families.

**Table 5 - Clinical Summary of Prevention Studies**

Study	Number of Subjects	Dose	Reduction in Clinical Influenza (Protective Efficacy)
<b>Seasonal Studies</b>			

			ITT Population
WV15673/WV15697 Adults	1559	Placebo 75 mg o.d. 75 mg b.i.d. 42 days	76%, $p = 0.00055$ 72%, $p = 0.00125$
WV15825 Elderly	548	Placebo 75 mg o.d. 42 days	92%, $p = 0.00153$
Post-exposure Studies			
			ITTIINAB Population
WV15799 Adult/adolescent contacts, index case not treated	405	Placebo 75 mg o.d., 7 days	92%, $p = 0.000076$
WV16193 Index case treated, age $\geq 1$ year	89**	Prophylaxis: 75 mg o.d., 10 days* Treatment: 75 mg b.i.d., 5 days*	78.8%, $p = 0.0008$
WV16193 (children 1-12 years)	117	Prophylaxis: 30 mg (1-2 years) 45 mg (3-5 years) 60 mg (6-12 years) o. d., 10 days Treatment: 30 mg (1-2 years) 45 mg (3-5 years) 60 mg (6-12 years) b. i. d., 5 days	80.1% (22.0-94.9), $p = 0.0206$

\*Pediatric dosing adjusted according to age

\*\* Number of households

ITT=Intent to treat

ITTIINAB=Intent to treat, index infected, not infected at baseline

### Pediatric Patients

In the post-exposure prophylaxis study in the family (WV16193; see 'Adult Prevention Studies') there were 215 pediatric contact cases (> 1 to 12 years of age). There was an even distribution of boys and girls with the majority being Caucasian. The mean age was 8 years (range 1 to 12). The data from this subset of pediatric patients was examined to determine if oseltamivir was effective in the prevention of influenza infection in this setting. When subjects who were already shedding virus at baseline were excluded (ITTIINAB population, 117), 17 pediatric contacts became infected, 2 in the prophylaxis group and 15 in the treatment group (see Table 5). The protective efficacy in the pediatric contacts was similar to that achieved in the overall population in this study.

The dosing schedule in this study was by age. The majority of children received the now recommended schedule of treatment by weight in children (see 4 DOSAGE AND ADMINISTRATION). There were, however, some children who were under- or over-dosed (23% and 9%, respectively) in this study.

### Prophylaxis of Influenza in Immunocompromised Patients

A double-blind, placebo-controlled study was conducted for seasonal prophylaxis of influenza in 475

immunocompromised subjects (including 18 pediatric subjects 1 to 12 years of age) who had received solid organ (n=388; liver, kidney, liver and kidney) or hematopoietic stem cell transplants (n=87). Median time since transplant for solid organ transplant recipients was 1105 days for the placebo group and 1379 days for the oseltamivir group. Median time since transplant for hematopoietic stem cell transplant recipients was 424 days for the placebo group and 367 days for the oseltamivir group. Approximately 40% of subjects received influenza vaccine prior to entering the study. The primary efficacy endpoint for this study was the incidence of confirmed, clinical influenza, defined as oral temperature > 99.0°F/37.2°C plus cough and/or coryza, all recorded within 24 hours, plus either a positive virus culture or a four-fold increase in virus antibody titers from baseline. The incidence of confirmed clinical influenza was 3% (7/238) in the group not receiving oseltamivir phosphate compared with 2% (5/237) in the group receiving oseltamivir phosphate; this difference was not statistically significant. A secondary analysis was performed using the same clinical symptoms and RT-PCR for laboratory confirmation of influenza. Among subjects who were not already shedding virus at baseline, the incidence of RT-PCR confirmed clinical influenza was 3% (7/231) in the group not receiving oseltamivir phosphate and < 1% (1/232) in the group receiving oseltamivir phosphate.

#### 14.2 Comparative Bioavailability Studies

An open label, randomized, single-dose, two-treatment, two-sequence, two-period crossover oral bioequivalence study of Oseltamivir Powder for Oral Suspension, 6 mg/mL oseltamivir (as oseltamivir phosphate) (Strides Pharma Canada Inc.) and TAMIFLU<sup>®</sup>, 6 mg/mL oseltamivir (as oseltamivir phosphate) (Genentech USA, Inc.) was performed in 59 healthy, adult, Asian male subjects under fasting conditions. A summary of the data from 55 subjects who completed the study is presented in the following table.

**SUMMARY TABLE OF THE COMPARATIVE BIOAVAILABILITY DATA**

Oseltamivir (12.5 mL x 6 mg/mL Oseltamivir Oral Suspension) From measured data				
Geometric Mean Arithmetic Mean (CV %)				
Parameter	Test <sup>1</sup>	Reference <sup>2</sup>	% Ratio of Geometric Means	90 % Confidence Interval
AUC <sub>T</sub> (ng·hr/mL)	169.56 173.93 (22.02)	183.86 187.70 (19.29)	92.2	90.1 - 94.4
AUC <sub>I</sub> (ng·hr/mL)	173.05 177.37 (21.67)	187.18 191.02 (19.07)	92.5	90.4 - 94.6
C <sub>max</sub> (ng/mL)	49.90 51.62 (26.64)	54.62 56.93 (28.77)	91.3	87.0 - 95.9
T <sub>max</sub> <sup>3</sup> (hr)	2.00 (0.33 - 3.50)	1.75 (0.33 - 3.50)		
T <sub>1/2</sub> <sup>4</sup> (hr)	3.32 (49.65)	2.95 (42.08)		

<sup>1</sup> Oseltamivir Powder for Oral Suspension (oseltamivir as oseltamivir phosphate) 6 mg/mL Oral Suspension (Strides Pharma Canada Inc.)

<sup>2</sup> TAMIFLU® (oseltamivir as oseltamivir phosphate) 6 mg/mL Oral Suspension (Genentech USA, Inc.) was purchased in the USA

<sup>3</sup> Expressed as the median (range) only

<sup>4</sup> Expressed as the arithmetic mean (% CV) only

## 15 MICROBIOLOGY

**Virology:** Oseltamivir was also tested for its effect on human T cell proliferation *in vitro*. Both antigen specific T cell lines and peripheral blood lymphocytes were isolated from whole blood. There was a slight but significant inhibition of influenza specific T cell line proliferation in the presence of 1 and 10 µM active metabolite, while there was no effect on antigen stimulation of peripheral blood lymphocytes. This slight effect (< 20%) on T cell proliferation is unlikely to compromise the long-term immune status of the patient with respect to subsequent influenza infection.

The active metabolite inhibits neuraminidases of influenza viruses of both types A and B. Inhibitory concentrations *in vitro* are in the low nanomolar range. The 50% inhibitory concentration (IC<sub>50</sub>) was in the range of 0.1 to 2.6 nM. The relationship between the *in vitro* antiviral activity in cell culture and the inhibition of influenza virus replication in humans has not been established. The active metabolite also inhibits influenza virus infection and replication *in vitro* and inhibits influenza virus replication and pathogenicity in animal models.

### Resistance:

#### *In vitro*

Extensive *in vitro* work has been completed with the active metabolite. Resistance to this compound does not arise readily *in vitro*. Several different resistance mutations in the viral neuraminidase have been selected *in vitro* in Roche studies or reported in the published literature. Resistance mutations tend to be viral sub-type specific. The degree of reduced sensitivity differs markedly for different mutations from 2 fold for I222V in N1 to 30,000 fold for R292K in N2. Influenza A virus H1N1 strains are associated with a histidine to tyrosine change at position 274 (H274Y) on the enzyme. In H3N2 subtypes the genetic alteration of interest is an arginine to lysine at position 292 (R292K) on the enzyme. *In vitro* these mutant viruses exhibit reduced growth potential compared to wild-type virus.

#### *In vivo*

*In vivo* experiments of infectivity and pathogenicity have been conducted with mutated viruses in mice and ferrets. These experiments have demonstrated that the H274Y H1N1 mutant and the R292K H3N2 mutant have reduced ability to infect susceptible animals compared to wild-type virus and that infection is not associated with clinical evidence of pathogenicity in the ferret. Correlation of *in vitro* resistance patterns to resistance *in vivo* is not known. Viruses with resistant neuraminidase genotypes have varying degrees of loss of fitness compared to wild-type.

### *Treatment of Influenza*

#### *Clinical Studies*

The risk of emergence of influenza viruses with reduced susceptibility or resistance to oseltamivir has been examined during Roche-sponsored clinical studies. Patients who were found to carry oseltamivir-resistant virus generally did so transiently, and showed no worsening of the underlying symptoms. In some pediatric patients, oseltamivir-resistant virus was detected for a prolonged period compared to patients carrying oseltamivir-sensitive virus; however these patients showed no prolongation of influenza symptoms. In immunocompromised adults and pediatrics (1 year of age and older), selection

of influenza viruses resistant oseltamivir can occur at higher frequencies than in the otherwise healthy population. In a treatment study of immunocompromised subjects, treatment-associated genotypic resistance was detected in 27% (8/30), 12% (6/52), and 0% (0/42) of influenza A/H1N1, A/H3N2, and B virus infections, respectively. Treatment-emergent resistance was observed at a higher frequency in hematopoietic stem cell transplant recipients (32%; 6/19).

**Table 6 - Incidence of Oseltamivir Resistance in Clinical Studies**

Patient Population	Patients with Resistance Mutations (%)	
	Phenotyping	Geno- and Phenotyping*
Adults and adolescents	4/1245 (0.32%)	5/1245 (0.4%)
Children (1-12 years)	19/464 (4.1%)	25/464 (5.4%)

\* Full genotyping was not performed in all studies.

Insufficient information is available to fully characterize the risk of emergence of resistance to oseltamivir phosphate in clinical use.

#### *Prevention of Influenza*

There has been no evidence for emergence of drug resistance associated with the use of oseltamivir phosphate in clinical studies conducted to date in post-exposure (7 days), post-exposure within household contacts (10 days) and seasonal (42 days) prevention of influenza in immunocompetent subjects.

Clinical and surveillance data: Natural mutations associated with reduced susceptibility to oseltamivir *in vitro* have been detected in influenza A and B viruses isolated from patients without exposure to oseltamivir. For example, in 2008 the oseltamivir resistance-associated substitution H275Y was found in > 99% of circulating 2008 H1N1 influenza isolates in Europe, while the 2009 H1N1 influenza (“swine flu”) was almost uniformly susceptible to oseltamivir. Resistant strains have also been isolated from both immunocompetent and immunocompromised patients treated with oseltamivir. The susceptibility to oseltamivir and the prevalence of such viruses appears to vary seasonally and geographically. Oseltamivir resistance has also been reported in patients with pandemic H1N1 influenza in connection with both therapeutic and prophylactic regimens.

The rate of emergence of resistance may be higher in the youngest age groups, and in immunocompromised patients. Oseltamivir-resistant viruses isolated from oseltamivir-treated patients and oseltamivir-resistant laboratory strains of influenza viruses have been found to contain mutations in N1 and N2 neuraminidases. Resistance mutations tend to be viral sub-type specific.

Prescribers should consider available information on influenza virus drug susceptibility patterns for each season when deciding whether to use Oseltamivir Powder for Oral Suspension (for latest information, please refer to WHO and/or local government websites).

#### **Cross resistance:**

Cross-resistance between zanamivir-resistant influenza mutants and oseltamivir-resistant influenza mutants has been observed *in vitro*.

Due to the limitations in the assays available to detect drug-induced shifts in virus susceptibility due to mutations in the viral hemagglutinin, an estimate of the incidence of oseltamivir resistance and possible cross-resistance to zanamivir in clinical isolates cannot be made. However, one of the three

oseltamivir-induced mutations in the viral neuraminidase from clinical isolates is the same as one of the three mutations in the viral neuraminidase from clinical isolates observed in zanamivir-resistant virus.

Insufficient information is available to fully characterize the risk of emergence of resistance or cross-resistance to oseltamivir phosphate in clinical use.

## 16 NON-CLINICAL TOXICOLOGY

### General Toxicology:

#### Acute Dose Toxicity

Acute oral administration was well tolerated by male and female adult rodents (mice and rats) and unweaned 14-day old male and female rats at 2000 mg/kg (~1000-fold the highest clinical dose). Single oral administration of 500 mg/kg (free base, corresponding to 657 mg/kg phosphate salt dose) or higher to juvenile 7-day old rats resulted in treatment-related mortality together with functional observation battery findings (FOB) and clinical signs indicative of general toxicity and imminent mortality (including low arousal, tremors, convulsions, alterations in general body posture, respiration, mucous membrane and skin colouration, and/or hypoactivity) and reduced body weight gain. The no effect level was 300 mg/kg (free base, corresponding to 394 mg/kg phosphate salt dose; ~150-fold the highest clinical dose) in juvenile rats in that study.

An intravenous range-finding study in mice (n=1/sex/dose) produced convulsions immediately after intravenous dosing with 250 mg/kg. The male died and the female recovered after 40 minutes. The maximum non-lethal dose of 100 mg/kg was confirmed in a further five males and five females observed for two weeks. Other than some evidence of a local reaction in the tail of two females, there were no significant adverse effects in this group.

#### Multiple Dose Toxicity

In multiple dose rat studies, doses up to 500 mg/kg/day (2 weeks), 650 mg/kg/day (4 weeks), and 200 mg/kg/day (27 weeks) were generally well tolerated, with no significant toxicologic effects.

A dose of 1000 mg/kg/day in a two week range finding rat study in unweaned 7-9 day old rats resulted in a high rate of mortality (18/24). At 500 mg/kg, no adverse effects were seen in the 7-9 day old rats or repeated treatment (500 mg/kg/day administered from 7 to 21 days post-partum).

In multiple dose rat studies, the highest doses examined ( $\geq 1000$  mg/kg/day) also induced two renal changes. One consisted of cortico-medullary mineralisation in the proximal tubules due to the imbalance of the calcium/phosphate ratio in the diet caused by dosing high levels of a phosphate salt. The second was a mild enhancement of chronic progressive nephropathy; rats are specifically sensitive to both these changes. A dose of 1000 mg/kg/day in the rat results in approximately 70 and 520 times the clinical exposure in humans, to the active metabolite and prodrug, respectively. In clinical studies, there was no biochemical evidence of renal effects in humans.

Marked gastrointestinal irritation was observed in marmosets at 2000 mg/kg/day, but not in four- and 39-week studies at 2 x 500 mg/kg/day. Emesis occurred at 500 mg/kg/day and above in marmosets, probably related to the concentration of the oral formulation. A reduction in incidence was associated with dividing the doses and halving the concentrations administered. This effect was seen at approximately 100 and 200 times the exposure values following clinical use in human, of the active metabolite and prodrug, respectively.

In the 39-week marmoset study, one 2 x 25 mg/kg/day group and two 2 x 100 mg/kg/day group animals were sacrificed prematurely. All three showed evidence of osteomalacia before dosing commenced, at autopsy and at the histopathological examination of the bones. No animal in the 2 x 500 mg/kg/day group was affected. Review of the clinical safety database, including the elderly, failed to reveal any biochemical evidence of skeletal effects in humans.

**Genotoxicity:** There was no evidence of mutagenic potential in any study (doses up to 5000 µg/plate), with or without metabolic activation. Separate bacterial cell gene mutation (Ames) tests were conducted for the pro drug and active metabolite. A mouse lymphoma cell mutation test examined the active metabolite. The pro drug was tested in a chromosome analysis assay with human lymphocytes, and in an *in vivo* micronucleus test in mice (oral dose of up to 2000 mg/kg). All the study systems were verified as sensitive by positive controls, and all the results were negative.

Two year rat and mouse studies and a six month transgenic Tg: AC mouse assay performed with the active metabolite were negative.

**Reproductive and Developmental Toxicology:** Fertility, teratology and pre- and post-natal studies were conducted to cover all phases of the reproductive process. There was no evidence of adverse effects on fertility or embryo-foetal development up to the highest dose of 1500 mg/kg/day in rats, or for teratogenicity testing in rabbits up to 500 mg/kg/day. These dose levels were associated with maternal toxicity. In rabbits mortalities occurred at 750 and 1500 mg/kg/day during a non-pregnant tolerance study. Some rabbits were sacrificed in the teratology range-finding and main studies at 500 mg/kg/day due to abortions associated with maternal toxicity. In the regulatory pre- and post-natal study in rats, maternal deaths occurred (9/25) at or immediately prior to delivery in the 1500 mg/kg/day group; prolonged parturition was also observed. Two further studies were therefore undertaken; although only 1/125 maternal deaths at parturition were seen in the combined 1500 mg/kg/day groups, extension of parturition was confirmed by these studies. It was concluded that the drug alone was not responsible for the maternal deaths in the first pre- and post-natal study.

At 1500 mg/kg/day in the rat teratology study, there was a slightly increased incidence of incomplete ossification of the 3<sup>rd</sup> sternebra in the exposed offspring, when compared to controls. Statistical significance was achieved, however, the majority of incidences occurred in one litter, where a general reduction in ossification was observed. In view of the isolated nature of this finding it was considered to be of doubtful toxicological significance.

## 17 SUPPORTING PRODUCT MONOGRAPHS

1. TAMIFLU® (oseltamivir powder for oral suspension, 6 mg/mL), submission control 255281, Product Monograph, Hoffmann-La Roche Limited. (SEP 09, 2022).

## **PATIENT MEDICATION INFORMATION**

### **READ THIS FOR SAFE AND EFFECTIVE USE OF YOUR MEDICINE**

#### **PrOseltamivir Powder for Oral Suspension**

oseltamivir powder for oral suspension

Read this carefully before you start taking Oseltamivir Powder for Oral Suspension and each time you get a refill. This leaflet is a summary and will not tell you everything about this drug. Talk to your healthcare professional about your medical condition and treatment and ask if there is any new information about Oseltamivir Powder for Oral Suspension.

#### **What is Oseltamivir Powder for Oral Suspension used for?**

Oseltamivir Powder for Oral Suspension is used to treat or prevent the onset of the flu in adults and children 1 year and older.

#### **How does Oseltamivir Powder for Oral Suspension work?**

The flu (influenza) is an infection caused by influenza viruses. It is most common in the fall and winter. The flu is highly contagious. The virus usually enters the body through the mouth, nose or eyes. When a person with the flu coughs or sneezes, the virus goes into the air and can be inhaled by anyone nearby.

Oseltamivir Powder for Oral Suspension works by stopping the flu virus from spreading through your body.

When Oseltamivir Powder for Oral Suspension is taken within 2 days of you feeling symptoms of the flu, it can help you feel better sooner.

When Oseltamivir Powder for Oral Suspension is taken within 2 days after coming into contact with someone with flu symptoms, it can help prevent you from getting sick with the flu.

#### **What are the ingredients in Oseltamivir Powder for Oral Suspension?**

Medicinal ingredients: oseltamivir (as oseltamivir phosphate)

Non-medicinal ingredients: monosodium citrate, saccharin sodium, sodium benzoate, sorbitol, titanium dioxide, tutti-frutti flavouring and xanthan gum.

#### **Oseltamivir Powder for Oral Suspension comes in the following dosage forms:**

Oseltamivir Powder for Oral Suspension is available as a 6 mg/mL liquid suspension (liquid form with a dosing dispenser called oral suspension). The suspension will be prepared by your pharmacist and will be a thick white fruit-flavoured liquid.

#### **Do not use Oseltamivir Powder for Oral Suspension if:**

You are allergic or sensitive to the medicinal ingredient of Oseltamivir Powder for Oral Suspension (oseltamivir phosphate) or any other ingredient in Oseltamivir Powder for Oral Suspension (see “What are the ingredients in Oseltamivir Powder for Oral Suspension?”).

**To help avoid side effects and ensure proper use, talk to your healthcare professional before you take Oseltamivir Powder for Oral Suspension. Talk about any health conditions or problems you may have, including if you:**

- Have hereditary fructose intolerance, as the Oseltamivir Powder for Oral Suspension oral suspension contains sorbitol and you should not use the suspension if you have this condition
- Are allergic to other medicines, food and dyes
- Have any type of kidney disease



- Have liver disease
- Are pregnant or plan on becoming pregnant
- Are breast-feeding or planning to breast-feed

**Tell your healthcare professional about all the medicines you take or plan on taking, including any drugs, vitamins, minerals, natural supplements or alternative medicines.**

How to take Oseltamivir Powder for Oral Suspension:

- Your doctor has prescribed Oseltamivir Powder for Oral Suspension after diagnosing your flu condition. Other people may not benefit from taking this medicine, even though their problems may seem similar to yours. Do not give your Oseltamivir Powder for Oral Suspension to anyone else.
- Oseltamivir Powder for Oral Suspension does not replace the flu shot. If you are at high risk of developing complications from the flu, you should continue to get your flu shot as recommended by your doctor. You may still get the flu even though you got your annual flu shot.
- It is important that you start taking Oseltamivir Powder for Oral Suspension as soon as possible from the first appearance of your flu symptoms. If your flu symptoms, most notably fever, do not begin to improve in the first day or two after you start Oseltamivir Powder for Oral Suspension, speak with your doctor.
- Oseltamivir Powder for Oral Suspension can be taken with food. As with many medicines, if you take Oseltamivir Powder for Oral Suspension with a light snack, milk, or a meal, you may reduce possible stomach upset.
- Shake the medicine well before each use.
- You must complete the entire treatment recommended by your doctor, even if you are feeling better.
- Take this medicine only as directed by your doctor. Do not take more of it, do not take it more often, and do not take it for a longer time than prescribed by your doctor. Never share Oseltamivir Powder for Oral Suspension with anyone, even if they have the same symptoms.

**Usual dose:**

Treatment with Oseltamivir Powder for Oral Suspension (adults, adolescents and children (1 year and older))

You should start taking Oseltamivir Powder for Oral Suspension no more than two days after flu symptoms have started. Typical symptoms of the flu include sudden onset of fever, headache, tiredness, muscular weakness, runny or stuffy nose, sore throat and cough.

- Oseltamivir Powder for Oral Suspension should be taken twice a day (once in the morning and once in the evening) for five days.

Prevention with Oseltamivir Powder for Oral Suspension (adults, adolescents and children (1 year and older))

You should start taking Oseltamivir Powder for Oral Suspension within 2 days after coming into contact with someone with flu symptoms. Typically, about one in ten people get the flu during an outbreak, although this may change. If someone close to you (like someone living in your home) has the flu, your chance of getting it yourself is higher (around one in five people).

- Oseltamivir Powder for Oral Suspension should be taken once a day for 10 days or longer as

recommended by your doctor if you have come into contact with someone who has the flu. It is important that you start taking Oseltamivir Powder for Oral Suspension as soon as possible after you were in contact.

### Preparing Oseltamivir Powder for Oral Suspension (liquid form taken by mouth):

- Your pharmacist may have prepared the medicine for you when you collected your prescription. If not, you can do it easily yourself.




1. Tap the closed bottle gently several times to loosen the powder.
2. The overcap in the pack has a line marked to show you an exact amount (55 mL). Fill it with water to the fill line.
3. Pour all of the water from the overcap into the bottle, onto the powder. Always use 55 mL of water, whatever the dose you need. Put the child-resistant cap back on the bottle. Shake the bottle well for 15 seconds.
4. Open the bottle and press the bottle adapter hard into the neck of the bottle.
5. Screw the child-resistant cap tightly onto the top of the bottle, which now includes the adapter. This will make sure that the bottle adapter fits in the bottle in the right position.


You now have a bottle of Oseltamivir Powder for Oral Suspension oral suspension ready to measure out a dose. You will not need to prepare it again unless you start a new bottle.

### How to Measure and Give the Correct Dose


- Always use the oral dispenser (syringe) provided with your medicine to measure a correct dose.
- For doses above 5.0 mL to 10 mL, the 10 mL oral dispenser (transparent plunger) should be used.



1. Shake closed bottle well for about 5 seconds before each use.
2. Push the plunger completely down toward the tip of the dispenser.



3. Remove child-resistant cap from the bottle. Insert tip of the dispenser firmly into the opening of the bottle adapter.
4. Turn the entire unit (bottle and dispenser) upside down.



Pull the plunger out slowly until the desired amount of medication is withdrawn into the dispenser.

5. Turn the entire unit (bottle and dispenser) right side up and remove the oral dispenser slowly from the bottle.
6. Put the medicine directly into the mouth by pushing down the plunger of the dispenser. Make sure the medicine is swallowed.
7. Close bottle with child-resistant cap after each use.
8. Take apart the oral dispenser, rinse under running tap water and air dry prior to next

use. Do not boil the oral dispenser to avoid damage to the dispenser.

Contact your doctor or pharmacist if the dispenser is lost or damaged, and they will advise you on how to continue to take your medication.

**Overdose:**

If you think you, or a person you are caring for, have taken too much Oseltamivir Powder for Oral Suspension, contact a healthcare professional, hospital emergency department, or regional poison control centre immediately, even if there are no symptoms.

**Missed Dose:**

If you forget to take a dose of Oseltamivir Powder for Oral Suspension, take it as soon as possible, then continue with the regular times you take your medication. Do not take double the amount if you miss one dose.

If you have missed several doses, inform your doctor and follow the advice given to you. Do not change your dose of Oseltamivir Powder for Oral Suspension unless instructed to do so by your doctor.

**What are possible side effects from using Oseltamivir Powder for Oral Suspension?**

These are not all the possible side effects you may have when taking Oseltamivir Powder for Oral Suspension. If you experience any side effects not listed here, tell your healthcare professional.

Oseltamivir Powder for Oral Suspension is generally well tolerated. The most common/possible unwanted effects are:

- Nausea
- Vomiting
- Abdominal pain
- Headache

<b>Serious side effects and what to do about them</b>			
<b>Symptom / effect</b>	<b>Talk to your healthcare professional</b>		<b>Stop taking drug and get immediate medical help</b>
	<b>Only if severe</b>	<b>In all cases</b>	
<b>UNKNOWN</b>			
Allergic Reaction: dizziness, breathing problems			X
Liver Problems: yellowing of skin or eyes, dark urine		X	
Severe Skin Reaction: flushing, rash, itching, swelling		X	
Neurologic and Behavioural Issues (in pediatric patients): delirium (confused thinking), hallucinations, self-harm (sometimes fatal)		X	

If you have a troublesome symptom or side effect that is not listed here or becomes bad enough to interfere with your daily activities, tell your healthcare professional.

### **Reporting Side Effects**

You can report any suspected side effects associated with the use of health products to Health Canada by:

- Visiting the Web page on Adverse Reaction Reporting (<https://www.canada.ca/en/health-canada/services/drugs-health-products/medeffect-canada.html>) for information on how to report online, by mail or by fax; or
- Calling toll-free at 1-866-234-2345.

*NOTE: Contact your health professional if you need information about how to manage your side effects. The Canada Vigilance Program does not provide medical advice.*

### **Storage:**

- Keep out of reach and sight of children.
- Store away from heat.
- Store Oseltamivir Powder for Oral Suspension (dry powder) at room temperature at 15-30°C.
- After reconstitution, store at room temperature (15-30°C) for 10 days or in a refrigerator (2-8°C) for 17 days.
- Keep Oseltamivir Powder for Oral Suspension liquid suspension in its original labeled container in a refrigerator at 2-8°C. Do not freeze. The pharmacist will write the date of expiration on the bottle label when reconstituted by the pharmacist.
- Do not use this medicine after the expiry date on the package.
- Do not throw away medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. This will help to protect the environment.

### **If you want more information about Oseltamivir Powder for Oral Suspension:**

- Talk to your healthcare professional
- Find the full product monograph that is prepared for healthcare professionals and includes this Patient Medication Information by visiting the Health Canada website: (<https://www.canada.ca/en/health-canada/services/drugs-health-products/drug-products/drug-product-database.html>); the manufacturer's website ([www.strides.com](http://www.strides.com)), or by calling (905) 829-3838.

This leaflet was prepared by Strides Pharma Canada Inc.

Last Revised: June 1, 2023