

PRODUCT MONOGRAPH
INCLUDING PATIENT MEDICATION INFORMATION

PrIPG-PREGABALIN

Pregabalin Capsules

Capsules, 25, 50, 75, 100, 150, 200, 225, 300 mg, oral

Mfr. Std.

Analgesic Agent

Marcan Pharmaceuticals Inc.
2 Gurdwara Road, Suite 112
Ottawa, ON K2E 1A2

Date of Initial Authorization:
JAN 24, 2014

Date of Revision:
JUL 13, 2023

Submission Control Number: 272305

RECENT MAJOR LABEL CHANGES

7 WARNINGS AND PRECAUTIONS	07/2023
7 WARNINGS AND PRECAUTIONS, 7.1.1. Pregnant Women	
3 SERIOUS WARNINGS AND PRECAUTIONS BOX	06/2020
7 WARNINGS AND PRECAUTIONS, Neurologic	
7 WARNINGS AND PRECAUTIONS, General, Patient Counselling Information	06/2020

TABLE OF CONTENTS

Sections or subsections that are not applicable at the time of authorization are not listed.

RECENT MAJOR LABEL CHANGES	2
TABLE OF CONTENTS	2
PART I: HEALTH PROFESSIONAL INFORMATION	4
1 INDICATIONS	4
1.1 Pediatrics	4
1.2 Geriatrics.....	4
2 CONTRAINDICATIONS	4
3 SERIOUS WARNINGS AND PRECAUTIONS BOX	4
4 DOSAGE AND ADMINISTRATION	5
4.1 Dosing Considerations	5
4.2 Recommended Dose and Dosage Adjustment	5
4.4 Administration	7
5 OVERDOSAGE	7
6 DOSAGE FORMS, STRENGTHS, COMPOSITION AND PACKAGING	8
7 WARNINGS AND PRECAUTIONS	10
7.1 Special Populations.....	19
7.1.1 Pregnant Women	20
7.1.2 Breast-feeding.....	21
7.1.3 Pediatrics.....	21
7.1.4 Geriatrics.....	21

8	ADVERSE REACTIONS.....	22
8.2	Clinical Trial Adverse Reactions.....	22
8.3	Less Common Clinical Trial Adverse Reactions (< 2%).....	37
8.4	Abnormal Laboratory Findings: Hematologic, Clinical Chemistry and Other Quantitative Data.....	43
8.5	Post-Market Adverse Reactions	43
9	DRUG INTERACTIONS	47
9.2	Drug Interactions Overview	47
9.4	Drug-Drug Interactions	47
9.5	Drug-Food Interactions.....	49
9.6	Drug-Herb Interactions.....	49
9.7	Drug-Laboratory Test Interactions.....	49
10	CLINICAL PHARMACOLOGY	49
10.1	Mechanism of Action.....	49
10.2	Pharmacodynamics.....	49
10.3	Pharmacokinetics.....	50
11	STORAGE, STABILITY AND DISPOSAL.....	52
12	SPECIAL HANDLING INSTRUCTIONS.....	52
PART II: SCIENTIFIC INFORMATION.....		53
13	PHARMACEUTICAL INFORMATION.....	53
14	CLINICAL TRIALS	53
14.1	Clinical Trials by Indication	53
	Diabetic Peripheral Neuropathy.....	52
	Post-herpetic Neuralgia	56
	Spinal Cord Injury.....	59
	Fibromyalgia.....	60
14.3	Comparative Bioavailability Studies.....	64
15	MICROBIOLOGY	64
16	NON-CLINICAL TOXICOLOGY	64
17	SUPPORTING PRODUCT MONOGRAPHS.....	68
PATIENT MEDICATION INFORMATION.....		69

PART I: HEALTH PROFESSIONAL INFORMATION

1 INDICATIONS

Adults

IPG-PREGABALIN (Pregabalin capsules) is indicated for the management of neuropathic pain associated with:

- Diabetic peripheral neuropathy;
- Post-herpetic neuralgia;

IPG-PREGABALIN is indicated for the management of pain associated with fibromyalgia.

The efficacy of pregabalin in the management of pain associated with fibromyalgia for up to 6 months was demonstrated in a placebo-controlled trial in patients who had initially responded to pregabalin during a 6-week open-label phase.

1.1 Pediatrics

Pediatrics (< 18 years of age): The safety and efficacy of pregabalin in pediatric patients (< 18 years of age) have not been established and its use in this patient population is not indicated (see [7.1.3 Pediatrics](#)).

1.2 Geriatrics

Geriatrics: Pregabalin oral clearance tended to decrease with increasing age. This decrease in pregabalin oral clearance is consistent with age-related decreases in creatinine clearance. Reduction of pregabalin dose may be required in patients who have age-related compromised renal function (see [7.1.4 Geriatrics](#)).

2 CONTRAINDICATIONS

IPG-PREGABALIN is contraindicated in patients who are hypersensitive to this drug or to any ingredient in the formulation, including any non-medicinal ingredient, or component of the container. For a complete listing, see [6 DOSAGE FORMS, STRENGTHS, COMPOSITION AND PACKAGING](#).

3 SERIOUS WARNINGS AND PRECAUTIONS BOX

Life-threatening Respiratory Depression

Concomitant use of IPG-PREGABALIN with opioids may result in respiratory depression, profound sedation, syncope, and death.

- Limit dosages and durations to the minimum required.
- Follow patients for signs and symptoms of respiratory depression and sedation.

4 DOSAGE AND ADMINISTRATION

4.1 Dosing Considerations

In accordance with current clinical practice, if IPG-PREGABALIN has to be discontinued, it is recommended this should be done gradually over a minimum of 1 week (see [7 WARNINGS AND PRECAUTIONS, Abrupt or Rapid Discontinuation](#)).

Patients with Impaired Renal Function

Pregabalin is primarily eliminated from the systemic circulation by renal excretion as unchanged drug. In some elderly patients and those with a medical history of significant renal insufficiency, daily dosages should be reduced accordingly (see [4.2 Recommended Dose and Dosage Adjustment, Dosage Adjustment Based on Renal Function](#)).

4.2 Recommended Dose and Dosage Adjustment

Adults

Neuropathic Pain Associated with Diabetic Peripheral Neuropathy

The recommended starting dose for IPG-PREGABALIN is 150 mg/day, given in two or three divided doses (75 mg BID or 50 mg TID), with or without food in patients with a creatinine clearance rate of at least 60 mL/min. Efficacy of pregabalin has been demonstrated within the first week. Based on individual patient response and tolerability, the dose may be increased to 150 mg BID (300 mg/day) after one week.

For patients who experience significant and ongoing pain and can tolerate pregabalin 300 mg/day well, maximum daily dose of 600 mg (300 mg twice a day, BID) can be used. However, in clinical trials, pregabalin 600 mg/day did not provide additional significant efficacy and patients treated with this dose experienced markedly higher rates of adverse events and discontinued the trial more frequently (see [8 ADVERSE REACTIONS, Table 2](#) and [Table 6](#)). Doses above 600 mg/day have not been studied and are not recommended.

Neuropathic Pain Associated with Post-herpetic Neuralgia

The recommended starting dose for IPG-PREGABALIN is 150 mg/day, given in two or three divided doses (75 mg BID or 50 mg TID), with or without food in patients with a creatinine clearance rate of at least 60 mL/min. Efficacy of pregabalin has been demonstrated within the first week. Based on individual patient response and tolerability, the dose may be increased to 150 mg BID (300 mg/day) after one week.

For patients who experience significant and ongoing pain and can tolerate pregabalin 300 mg/day well, maximum daily dose of 600 mg (300 mg twice a day, BID) can be used. However, in clinical trials, pregabalin 600 mg/day did not provide additional significant efficacy and patients treated with this dose experienced markedly higher rates of adverse events and discontinued the trial more frequently (see [8 ADVERSE REACTIONS, Table 4](#) and [Table 7](#)). Doses above 600 mg/day have not been studied and are not recommended.

Pain Associated with Fibromyalgia

The recommended dosage is 300 to 450 mg/day, given in two divided doses. The recommended starting dose for IPG-PREGABALIN is 150 mg/day, given in two divided doses (75 mg BID), with or

without food in patients with a creatinine clearance rate of at least 60 mL/min. Based on individual response and tolerability, the dose may be increased to 150 mg BID (300 mg/day) after one week. Patients who do not experience sufficient benefit with 300 mg/day may be further increased to 225 mg BID (450 mg/day). In some patients, efficacy of pregabalin has been demonstrated within the first week.

For patients who experience significant and ongoing pain and can tolerate pregabalin 300 mg/day well, maximum daily dose of 600 mg (300 mg twice a day, BID) can be used. However, in clinical trials of fibromyalgia, pregabalin 600 mg/day did not provide additional significant efficacy and patients treated with this dose experienced significantly higher rates of adverse events and discontinued the trial more frequently (see [8 ADVERSE REACTIONS, Table 8](#) and [Table 11](#)). In view of the dose-related adverse events, the decision to treat patients with doses above 450 mg/day should be based on clinical judgment of the treating physician. Doses above 600 mg/day have not been studied and are not recommended.

Dosage Adjustment Based on Renal Function

Pregabalin is primarily eliminated by renal excretion. Therefore, the dose should be adjusted for patients with reduced renal function. Pregabalin clearance is directly proportional to creatinine clearance. Therefore, dosing adjustment should be based on creatinine clearance (CL_{Cr}), as indicated in Table 1.

To use this dosing table, an estimate of the patient's creatinine clearance (CL_{Cr}) in mL/min is needed. CL_{Cr} in mL/min may be estimated from serum creatinine (mg/dL) determination using the Cockcroft and Gault equation:

$$CL_{Cr} = \frac{[140 - \text{age (years)}] \times \text{weight (kg)}}{72 \times \text{serum creatinine (mg/dL)}} \quad (\times 0.85 \text{ for female patients})$$

Pregabalin is effectively removed from plasma by hemodialysis. Over a 4-hour hemodialysis treatment, plasma pregabalin concentrations are reduced by approximately 50%. For patients receiving hemodialysis, pregabalin daily dose should be adjusted based on renal function. In addition to the daily dose adjustment, a supplemental dose should be given immediately following every 4-hour hemodialysis treatment (see Table 1).

Table 1. Pregabalin Dosage Adjustment Based on Renal Function

Creatinine Clearance (CL _{cr}) (mL/min)	Total Pregabalin Daily Dose (mg/day) ^a				Dose Regimen
	Recommended Dose Escalation*				
	Starting Dose	→ up to		Maximum daily dose	
≥ 60	150	300	450	600	BID or TID
30 - 60	75	150	225	300	BID or TID
15 - 30	25-50	75	100-150	150	QD or BID
< 15	25	25-50	50-75	75	QD
Supplementary dosage following hemodialysis (mg)^b					
Patients on the 25 mg QD regimen: take one supplemental dose of 25 mg or 50 mg					
Patients on the 25 - 50 mg QD regimen: take one supplemental dose of 50 mg or 75 mg					
Patients on the 50 - 75 mg QD regimen: take one supplemental dose of 75 mg or 100 mg					
Patients on the 75 mg QD regimen: take one supplemental dose of 100 mg or 150 mg					

TID = Three divided doses; BID = Two divided doses; QD = Single daily dose.

* Based on individual patient response and tolerability.

^a Total daily dose (mg/day) should be divided as indicated by dose regimen to provide mg/dose.

^b Supplementary dose is a single additional dose.

Geriatrics (> 65 years): Pregabalin oral clearance tended to decrease with increasing age. This decrease in pregabalin oral clearance is consistent with age-related decreases in creatinine clearance. Reduction of pregabalin dose may be required in patients who have age-related compromised renal function.

Pediatrics (< 18 years of age): The safety and efficacy of pregabalin in pediatric patients (< 18 years of age) have not been established and its use in this patient population is not indicated.

4.4 Administration

IPG-PREGABALIN is given orally with or without food (see [9.5 Drug-Food Interactions](#)).

5 OVERDOSAGE

Signs, Symptoms and Laboratory Findings of Acute Overdosage in Humans

The highest known dose of pregabalin received in the clinical development program in which there was no fatal outcome was 15,000 mg in 1 patient. The types of adverse events experienced by patients who received an overdose were not clinically different from other patients receiving recommended doses of pregabalin.

In post-marketing experience, fatal outcomes in cases in which pregabalin has been taken in combination with other medications have been reported with a pregabalin overdose as low as 800 mg in a day. In none of these cases has pregabalin been established as the cause of death or in pregabalin monotherapy. The lowest fatal dose with pregabalin alone has not yet been identified.

The most commonly reported adverse events observed when pregabalin was taken in overdose (dose range from 800 mg/day up to 11,500 mg as a single dose) included affective disorder, somnolence, confusional state, depression, agitation, and restlessness. Seizures were also reported.

Treatment or Management of Overdose

There is no specific antidote for overdose with pregabalin. If indicated, elimination of unabsorbed drug may be attempted by emesis or gastric lavage; usual precautions should be observed to maintain the airway. General supportive care of the patient is indicated including monitoring of vital signs and observation of the clinical status of the patient. A Certified Poison Control Center should be contacted for up-to-date information on the management of overdose with pregabalin.

Hemodialysis

Standard hemodialysis procedures result in significant clearance of pregabalin (approximately 50% in 4 hours) and should be considered in cases of overdose. Although hemodialysis has not been performed in the few known cases of overdose, it may be indicated by the patient's clinical state or in patients with significant renal impairment.

For management of a suspected drug overdose, contact your regional poison control centre.

6 DOSAGE FORMS, STRENGTHS, COMPOSITION AND PACKAGING

Table – Dosage Forms, Strengths, Composition and Packaging

Route of Administration	Dosage Form / Strength/Composition	Non-medicinal Ingredients
Oral	Capsules, 25 mg, 50 mg, 75 mg, 100 mg, 150 mg, 200 mg, 225 mg, 300 mg	Pre-gelatinized starch and Talc. Capsule shells contain gelatin and titanium dioxide. In addition, orange capsule shells contain red iron oxide. The markings on the capsules are in black ink, which contains ammonia solution, black iron oxide, butyl alcohol, dehydrated alcohol, isopropyl alcohol, potassium hydroxide, propylene glycol, shellac and water.

Dosage Forms

IPG-PREGABALIN supplied as a hard gelatin capsule for daily oral administration.

25 mg capsules: Opaque White/Opaque white size '4' hard gelatin capsules, radially imprinted with 'PG' on cap and '25' on body with black ink, filled with white to off white powder.

50 mg capsules: Opaque White/Opaque white size '2' hard gelatin capsules, radially imprinted with 'PG' on cap and '50' on body and band on body with black ink, filled with white to off white powder.

75 mg capsules: Opaque Orange/Opaque White size '4' hard gelatin capsules, radially imprinted with 'PG' on cap and '75' on body with black ink, filled with white to off white powder.

100 mg capsules: Orange Opaque/ Orange Opaque size '3' hard gelatin capsules, radially imprinted with 'A' on cap and '143' on body with black ink, filled with white to off white powder.

150 mg capsules: Opaque White/Opaque white size '2' hard gelatin capsules, radially imprinted with 'PG' on cap and '150' on body with black ink, filled with white to off white powder.

200 mg capsules: Orange Opaque/ Orange Opaque size '1' hard gelatin capsules, radially imprinted with 'A' on cap and '145' on body with black ink, filled with white to off white powder.

225 mg capsules: Opaque White/ Light Orange size '1' hard gelatin capsules, radially imprinted with 'A' on cap and '146' on body with black ink, filled with white to off white powder.

300 mg capsules: Opaque White/Orange size '0' hard gelatin capsules, radially imprinted with 'A' on cap and '147' on body with black ink, filled with white to off white powder.

Packaging

25 mg, 50 mg, 75 mg and 150 mg capsules are available in the packs of HDPE bottles of 100's or 500's.

100 mg, 200 mg and 225 mg capsules are available in HDPE bottles of 1000's.

300 mg capsules are available in the packs of HDPE bottles of 500's.

7 WARNINGS AND PRECAUTIONS

Please see [3 SERIOUS WARNINGS AND PRECAUTIONS BOX](#).

General

- **Peripheral Edema**

Pregabalin may cause peripheral edema. In controlled peripheral neuropathic pain and fibromyalgia clinical trials, pregabalin treatment caused peripheral edema in 9% of patients compared with 3% of patients in the placebo group. In these studies, 0.7% of pregabalin patients and 0.3% of placebo patients withdrew due to peripheral edema (see [8 ADVERSE REACTIONS, Peripheral Edema](#)).

In controlled clinical trials of up to 13 weeks in duration of patients without clinically significant heart or peripheral vascular disease, there was no apparent association between peripheral edema and cardiovascular complications such as hypertension or congestive heart failure. In the same trials, peripheral edema was not associated with laboratory changes suggestive of deterioration in renal or hepatic function.

Higher frequencies of weight gain and peripheral edema were observed in patients taking both pregabalin and a thiazolidinedione antidiabetic agent compared to patients taking either drug alone. The majority of patients using thiazolidinedione antidiabetic agents in the overall safety database were participants in studies of pain associated with diabetic peripheral neuropathy. In this population, peripheral edema was reported in 3% (2/60) of patients who were using thiazolidinedione antidiabetic agents only, 8% (69/859) of patients who were treated with pregabalin only, and 19% (23/120) of patients who were on both pregabalin and thiazolidinedione antidiabetic agents. Similarly, weight gain was reported in 0% (0/60) of patients on thiazolidinediones only; 4% (35/859) of patients on pregabalin only; and 7.5% (9/120) of patients on both drugs.

As the thiazolidinedione class of antidiabetic drugs can cause weight gain and/or fluid retention, possibly exacerbating or leading to heart failure, care should be taken when co-administering pregabalin and these agents.

- **Abrupt or Rapid Discontinuation**

Following abrupt or rapid discontinuation of pregabalin, some patients reported symptoms including insomnia, nausea, headache, anxiety, hyperhidrosis, and diarrhea. Pregabalin should be tapered gradually over a minimum of one week rather than discontinued abruptly (see [8 ADVERSE REACTIONS, Adverse Events Following Abrupt or Rapid Discontinuation](#)).

Convulsions, including status epilepticus and grand mal convulsions, have occurred in non-epileptic patients during treatment with pregabalin or after abrupt discontinuation (see [8.5 Post-Market Adverse Reactions](#)).

- **Encephalopathy**

There have been serious post-marketing reports of encephalopathy, mostly in patients with underlying conditions that may precipitate encephalopathy. Some cases were reported in

patients with a history of kidney or liver disease. Since there have been rare reports of renal failure with pregabalin, specific caution should be exercised when prescribing pregabalin to the elderly with age-related compromised renal function and patients with kidney disease or risk factors for renal failure (see [7 WARNINGS AND PRECAUTIONS, Renal Failure](#) and [8.5 Post-Market Adverse Reactions](#)).

- **Patient Counselling Information**

Patients receiving IPG-Pregabalin should be given the following instructions by the physician:

- 1. Angioedema:** Patients should be advised that IPG-Pregabalin may cause angioedema, with swelling of the face, mouth (lip, gum, tongue) and neck (larynx and pharynx) that can lead to life-threatening respiratory compromise. Instruct patients to discontinue IPG-Pregabalin and immediately seek medical care if they experience these symptoms.
- 2. Hypersensitivity:** Patients should be advised that IPG-Pregabalin has been associated with hypersensitivity reactions such as wheezing, dyspnea, rash, hives, and blisters. Instruct patients to discontinue IPG-Pregabalin and immediately seek medical care if they experience these symptoms.
- 3. Suicidal Behaviour and Ideation:** Patients, their caregivers, and families should be counselled to monitor for signs of suicidal ideation and behaviours and should be encouraged to report any distressing thoughts or feelings at any time to their healthcare professional.
- 4. Dizziness and Somnolence:** Patients should be counseled that IPG-Pregabalin may cause dizziness, somnolence, blurred vision and other CNS signs and symptoms. Accordingly, they should be advised not to drive, operate complex machinery, or engage in other hazardous activities until they have gained sufficient experience on pregabalin to gauge whether or not it affects their mental, visual, and/or motor performance adversely.
- 5. Edema and Weight Gain:** Patients should be counseled that IPG-Pregabalin may cause edema and weight gain. Patients should be advised that concomitant treatment with IPG-Pregabalin and a thiazolidinedione antidiabetic agent may lead to an additive effect on edema and weight gain. For patients with preexisting cardiac conditions, this may increase the risk of heart failure.
- 6. Abrupt or Rapid Discontinuation:** Patients should be advised to take IPG-Pregabalin as prescribed. Abrupt or rapid discontinuation may result in insomnia, nausea, headache, anxiety, hyperhidrosis, or diarrhea.
- 7. Visual Disturbances:** Patients should be counseled that IPG-Pregabalin may cause visual disturbances. Patients should be informed that if changes in vision occur, they should notify their physician (see [7 WARNINGS AND PRECAUTIONS, Ophthalmological Effects](#)).
- 8. Muscle Pain, Tenderness or Weakness:** Patients should be instructed to promptly report unexplained muscle pain, tenderness, or weakness, particularly if

accompanied by malaise or fever.

- 9. Concomitant Treatment with CNS Depressants, Alcohol:** Patients who require concomitant treatment with central nervous system depressants such as opiates or benzodiazepines should be informed that they may experience additive CNS side effects, such as somnolence.

In post-marketing experience, there are reports of respiratory failure, coma and deaths in patients taking pregabalin alone or in combination with other CNS depressants, including in patients with substance use disorders.

Patients should be told to avoid consuming alcohol while taking IPG-Pregabalin, as IPG-Pregabalin may potentiate the impairment of motor skills and sedation of alcohol.

- 10. Pregnant Women:** Patients should be instructed to notify their physician if they become pregnant or intend to become pregnant during their therapy, and to notify their physician if they are breast-feeding or intend to breast-feed during therapy. Patients should be encouraged to enroll in the North American Antiepileptic Drug Pregnancy Registry (NAAED) if they become pregnant. This registry is collecting information about the safety of anticonvulsant medications that can be taken by women during pregnancy to treat disorders such as epilepsy, mood disorder, and chronic pain. To enroll, patients can call the toll free number 1-888-233-2334. Information on the registry can also be found at the website <http://www.aedpregnancyregistry.org/> (see [7.1.1 Pregnant Women](#)).

- 11. Animal Studies in Male Reproduction:** Men being treated with IPG-Pregabalin who plan to father a child should be informed of the potential risk of male-mediated teratogenicity. In preclinical studies in rats, pregabalin was associated with an increased risk of male-mediated teratogenicity (see [7 WARNINGS AND PRECAUTIONS, Reproductive Health: Female and Male Potential](#)). The clinical significance of this finding is uncertain.

- 12. Skin:** Diabetic patients should be instructed to pay particular attention to skin integrity while being treated with IPG-PREGABALIN. Some animals treated with pregabalin developed skin ulcerations, although no increased incidence of skin lesions associated with pregabalin was observed in clinical trials (see [16 NON-CLINICAL TOXICOLOGY, Preclinical Toxicology](#)).

Patients should be informed of the availability of a patient information leaflet, and they should be instructed to read the leaflet prior to taking IPG-Pregabalin.

Carcinogenesis and Mutagenesis

- **Tumorigenic Potential**

In standard preclinical *in vivo* lifetime carcinogenicity studies of pregabalin, a high incidence of hemangiosarcoma was identified in two different strains of mice (see [16 NON-CLINICAL TOXICOLOGY](#)). The clinical significance of this finding is uncertain.

Clinical experience during pregabalin's premarketing development provides no direct means to assess its potential for inducing tumors in humans.

In clinical studies across various patient populations, comprising 6396 patient-years of exposure in 8666 patients ranging in age from 12 to 100 years, new or worsening-preexisting tumors were reported in 57 patients. The most common malignant tumor diagnosed was skin carcinoma (17 patients) followed by breast carcinoma (8 patients), prostatic carcinoma (6 patients), carcinoma not otherwise specified (6 patients), and bladder carcinoma (4 patients). Without knowledge of the background incidence and recurrence in similar populations not treated with pregabalin, it is impossible to know whether the incidence seen in these cohorts is or is not affected by treatment.

Cardiovascular

- **Congestive Heart Failure**

In controlled clinical studies, events of congestive heart failure were reported at an infrequent rate (between 0.1% and 1%; see [8.3 Less Common Clinical Trial Adverse Reactions \(< 2%\)](#)).

There have been post-marketing reports of congestive heart failure in some patients receiving pregabalin (see [8.5 Post-Market Adverse Reactions](#)). Although this adverse reaction has mostly been observed in elderly cardiovascular compromised patients during pregabalin treatment for a neuropathic pain indication, some cases have occurred in patients without reported edema or previous history of cardiovascular disease. Pregabalin should be used with caution in these patients. Discontinuation of pregabalin may resolve the reaction.

- **ECG Changes, PR Interval Prolongation**

Pregabalin treatment was associated with mild PR interval prolongation. In analyses of clinical trial ECG data, the mean PR interval increase was 3-6 msec at pregabalin doses ≥ 300 mg/day. This mean change difference was not associated with an increased risk of PR increase $\geq 25\%$ from baseline, an increased percentage of subjects with on-treatment PR > 200 msec, or an increased risk of adverse events of second or third degree AV block.

Dependence, Abuse Potential or Misuse

Cases of misuse, abuse and dependence have been reported. Caution should be exercised in patients with a history of substance abuse and the patient should be monitored for symptoms of pregabalin misuse, abuse or dependence (development of tolerance, dose escalation, drug-seeking behaviour have been reported).

Driving and Operating Machinery

Patients should be advised not to drive or operate complex machinery or engage in other hazardous activities until they have gained sufficient experience on pregabalin to gauge whether or not it affects their mental and/or motor performance adversely (see [PATIENT MEDICATION INFORMATION](#)).

Endocrine and Metabolism

- **Weight Gain**

Pregabalin may cause weight gain. In pregabalin-controlled peripheral neuropathic pain and fibromyalgia clinical trials with durations of up to 14 weeks, a gain of 7% or more over baseline weight was observed in 8% of pregabalin-treated patients and 3% of placebo-treated patients. Few patients treated with pregabalin (0.6%) withdrew from controlled trials due to weight gain (see [8 ADVERSE REACTIONS, Weight Gain](#)).

Pregabalin-associated weight gain was related to dose and duration of exposure. Pregabalin-associated weight gain did not appear to be associated with baseline BMI, gender, or age. Weight gain was not limited to patients with edema and was not necessarily due to edema-related events (see [7 WARNINGS AND PRECAUTIONS, Peripheral Edema](#)).

Although weight gain was not associated with clinically important changes in blood pressure in short-term controlled studies, the long-term cardiovascular effects of pregabalin-associated weight gain are unknown.

Among diabetic patients, pregabalin-treated patients gained an average of 1.6 kg (range: -16 to 16 kg), compared to an average 0.3 kg (range: -10 to 9 kg) weight gain in placebo patients. In a cohort of 333 diabetic patients who received pregabalin for at least 2 years, the average weight gain was 5.2 kg.

In the controlled fibromyalgia clinical trials, 10.7% of pregabalin-treated patients experienced weight gain of 7% or greater compared to 4.9% of placebo-treated patients. Pregabalin-treated patients gained an average of 1.7 Kg compared to an average of 0.7 Kg weight gain in placebo patients.

While the effects of pregabalin-associated weight gain on glycemic control have not been systematically assessed, in controlled and longer-term open label clinical trials with diabetic patients, pregabalin treatment did not appear to be associated with loss of glycemic control (as measured by HbA1C).

Gastrointestinal

There have been post-marketing reports of events related to reduced lower gastrointestinal tract function (e.g., intestinal obstruction, paralytic ileus, and constipation) in patients, some without reported previous history/episode(s), during initial/acute and chronic treatment with pregabalin, primarily in combination with other medications that have the potential to produce constipation. Some of these events were considered serious and required hospitalization. In a number of instances, patients were taking opioid analgesics including tramadol.

Caution should be exercised when pregabalin and opioid analgesics are used in combination, and measures to prevent constipation may be considered, especially in female patients and elderly as they may be at increased risk of experiencing lower gastrointestinal-related events (see [8.5 Post-Market Adverse Reactions](#)).

Hematologic

- **Laboratory Changes, Decreased Platelet Count**

Pregabalin treatment was associated with a decrease in platelet count. Pregabalin-treated subjects experienced a mean maximal decrease in platelet count of $20 \times 10^3/\text{mL}$, compared to $11 \times 10^3/\text{mL}$ in placebo patients. In the overall database of controlled trials, 2% of placebo patients and 3% of pregabalin patients experienced a potentially clinically significant decrease in platelets, defined as 20% below baseline value and $< 150 \times 10^3/\text{mL}$.

In randomized controlled trials, pregabalin was not associated with an increase in bleeding related adverse events.

Immune

- **Angioedema**

There have been post-marketing reports of angioedema in patients, some without reported previous history/episode(s), during initial/acute and chronic treatment with pregabalin). Specific symptoms included swelling of the face, mouth (tongue, lips, and gums), neck, throat, and larynx/upper airway. There have been reports of life-threatening angioedema with respiratory compromise requiring emergency treatment. Some of these patients did not have reported previous history/episode(s) of angioedema. IPG-Pregabalin should be immediately discontinued in patients with these symptoms. During the pre-marketing assessment of pregabalin in clinical trials, angioedema was reported as a rare reaction (see [8.3 Less Common Clinical Trial Adverse Reactions \(< 2%\)](#) and [8.5 Post-Market Adverse Reactions](#)).

Caution should be exercised when prescribing IPG-Pregabalin to patients with previous history/episode(s) of angioedema and related events. In addition, patients who are taking other drugs associated with angioedema (e.g., ACE-inhibitors) may be at increased risk of developing this condition.

- **Hypersensitivity**

There have been post-marketing reports of hypersensitivity reactions (e.g., skin redness, blisters, hives, rash, dyspnea, and wheezing). Pregabalin should be discontinued immediately if such symptoms occur (see [8.5 Post-Market Adverse Reactions](#)).

Monitoring and Laboratory Tests

Routine therapeutic drug monitoring or clinical laboratory testing is not required for patients treated with IPG-Pregabalin (see [8 ADVERSE REACTIONS](#)).

Musculoskeletal

- **Creatine Kinase Elevations**

Pregabalin treatment was associated with creatine kinase elevations. Mean changes

in creatine kinase from baseline to the maximum value were 60 U/L for pregabalin-treated patients and 28 U/L for the placebo patients. In all controlled trials across multiple patient populations, 2% of patients on pregabalin and 1% of placebo patients had a value of creatine kinase at least three times the upper limit of normal. Three pregabalin-treated subjects had events reported as rhabdomyolysis in premarketing clinical trials. The relationship between these myopathy events and pregabalin is not completely understood because the cases had documented factors that may have caused or contributed to these events. Prescribers should instruct patients to promptly report unexplained muscle pain, tenderness, or weakness, particularly if these muscle symptoms are accompanied by malaise or fever. Pregabalin treatment should be discontinued if myopathy is diagnosed or suspected or if markedly elevated creatine kinase levels occur.

Neurologic

- **Respiratory Depression**

Pregabalin has been associated with central nervous system (CNS) depression including sedation, somnolence, loss of consciousness as well as serious cases of respiratory depression. Patients with compromised respiratory function, respiratory or neurological disease, renal impairment and the elderly are at higher risk of experiencing these severe adverse effects. Concomitant use of CNS depressants with pregabalin is also a contributing factor.

- **Concomitant Use With Opioids**

Caution is advised when prescribing IPG-Pregabalin concomitantly with opioids due to risk of CNS depression. Concomitant use of opioids with IPG-PREGABALIN potentiates the risk of respiratory depression, profound sedation, syncope, and death. In an observational study of opioid users, those patients who took pregabalin concomitantly with an opioid had an increased risk for opioid-related death compared to opioid use alone (adjusted odds ratio [aOR], 1.68 [95% CI, 1.19 to 2.36]). Patients who require concurrent treatment with opioids or other CNS depressants should be observed carefully for signs and symptoms of CNS depression, and the dose of pregabalin or opioid should be reduced accordingly (see [7 WARNINGS AND PRECAUTIONS, General](#) and [7 WARNINGS AND PRECAUTIONS, Patient Counselling Information](#)).

- **Dizziness and Somnolence**

IPG-Pregabalin may cause dizziness and somnolence. In controlled peripheral neuropathic pain and fibromyalgia studies, pregabalin caused dizziness in 32% of patients compared to 8% in placebo. Somnolence was experienced by 17% and 4% of the patients treated with pregabalin and placebo, respectively. These events begin shortly after the initiation of therapy and generally occur more frequently at higher doses. In these studies, dizziness and somnolence led to withdrawal of 5% (placebo: 0.5%) and 3% (placebo: 0.1%) of the pregabalin-treated patients, respectively. For the remaining patients who experienced these events, dizziness and somnolence persisted until the last dose of pregabalin in 35% and 49% of the patients, respectively (see [8 ADVERSE REACTIONS, Table 3, Table 5, and Table 12](#), and [8.5 Post-Market Adverse Reactions](#)).

Ophthalmologic

- **Ophthalmological Effects**

In controlled studies, pregabalin treatment was associated with vision-related adverse events such as blurred vision (amblyopia) [6% pregabalin and 2% placebo] and diplopia (2% pregabalin and 0.5% placebo). Approximately 1% of pregabalin-treated patients discontinued treatment due to vision-related adverse events (primarily blurred vision). Of the patients who did not withdraw, the blurred vision resolved with continued dosing in approximately half of the cases (see [8.5 Post-Market Adverse Reactions](#)).

Prospectively planned ophthalmologic testing, including visual acuity testing, formal visual field testing and dilated funduscopy examination, was performed in over 3600 patients. In these patients, visual acuity was reduced in 7% of patients treated with pregabalin, and 5% of placebo-treated patients. Visual field changes were detected in 13% of pregabalin-treated, and 12% of placebo-treated patients. Funduscopy changes were observed in 2% of pregabalin-treated and 2% of placebo-treated patients. At this time, clinical significance of the ophthalmologic findings is unknown.

Patients should be informed that if changes in vision occur, they should notify their physician. If visual disturbance persists, further assessment, including discontinuation of pregabalin, should be considered. More frequent assessments should be considered for patients who are already routinely monitored for ocular conditions.

Psychiatric

- **Suicidal Behaviour and Ideation**

There have been post-marketing reports of suicide-related events, including completed suicide, suicide attempt, and suicidal ideation in patients treated with pregabalin for a variety of indications such as neuropathic pain, fibromyalgia, etc. In some of these reports, underlying psychiatric disorders may have contributed to the event. The mechanism of this risk is not known. Patients should be monitored for signs of suicidal ideation and behaviours and appropriate treatment should be considered. Patients should be encouraged to report any distressing thoughts or feelings at any time to their healthcare professional (see [8.5 Post-Market Adverse Reactions](#)).

Renal

- **Renal Failure**

In both clinical trials of various indications and post-marketing database, there are reports of patients, with or without previous history, experiencing renal failure while receiving pregabalin alone or in combination with other medications. Discontinuation of pregabalin should be considered as it has shown reversibility of this event in some cases. Caution is advised when prescribing pregabalin to the elderly or those with any degree of renal impairment (see [7.1 Special Populations, Renal](#); [7 WARNINGS AND PRECAUTIONS, Abrupt or Rapid Discontinuation](#); [8.5 Post-Market Adverse Reactions](#); and [4 DOSAGE AND ADMINISTRATION](#)).

Reproductive Health: Female and Male Potential

- **Fertility**

Impairment of Male Fertility

Preclinical Data

In fertility studies in which male rats were orally administered pregabalin (50 to 2500 mg/kg) prior to and during mating with untreated females, a number of adverse reproductive and developmental effects were observed. These included decreased sperm counts and sperm motility, increased sperm abnormalities, reduced fertility, increased preimplantation embryo loss, decreased litter size, decreased fetal body weights, and an increased incidence of fetal abnormalities. Effects on sperm and fertility parameters were reversible in studies of this duration (3-4 months). The no-effect dose for male reproductive toxicity in these studies (100 mg/kg) was associated with a plasma pregabalin exposure (AUC) approximately 3 times human exposure at the maximum recommended dose (MRD) of 600 mg/day.

In addition, adverse effects on reproductive organ (testes, epididymides) histopathology were observed in male rats exposed to pregabalin (500 to 1250 mg/kg) in general toxicology studies of 4 weeks or greater duration. The no-effect dose for male reproductive organ histopathology in rats (250 mg/kg) was associated with a plasma exposure approximately 8 times human exposure at the MRD.

In a fertility study in which female rats were given pregabalin (500, 1250, or 2500 mg/kg) orally prior to and during mating and early gestation, disrupted estrous cyclicity and an increased number of days to mating were seen at all doses, and embryo lethality occurred at the highest dose. The low dose in this study produced a plasma exposure approximately 9 times that in humans receiving the MRD. A no-effect dose for female reproductive toxicity in rats was not established. The clinical significance of female fertility findings in animals is unknown.

Human Data

In a double-blind, placebo-controlled clinical trial to assess the effect of pregabalin on sperm motility, 30 healthy male subjects were exposed to pregabalin 600 mg/day for 3 months (one complete sperm cycle). Pregabalin did not exhibit significant detrimental effects on the reproductive function of healthy male subjects, as measured by semen analysis, when compared with placebo (n = 16). However, due to the small sample size and short-term exposure to pregabalin (only one complete sperm cycle), no conclusions can be made regarding possible reproductive effects of pregabalin during long-term exposure. Effects on other male reproductive parameters in humans have not been adequately studied.

- **Teratogenic Risk**

Pregabalin was not teratogenic in mice, rats, or rabbits. Pregabalin induced fetal toxicity in rats and rabbits at ≥ 39 times the mean human exposure at the maximum recommended clinical dose of 600 mg/day [AUC (0-24) of 123 mcg•hr/mL]. In the prenatal-postnatal toxicity study, pregabalin induced offspring developmental toxicity

in rats at ≥ 5 times the maximum recommended human exposure. No developmental effects occurred at 2 times the maximum recommended human exposure (see [16 NON-CLINICAL TOXICOLOGY](#)).

Skin

- **Serious Skin Reactions**

There have been very rare post-marketing reports of serious cutaneous reactions, including Stevens-Johnson Syndrome (SJS), Toxic Epidermal Necrolysis (TEN), dermatitis exfoliative, bullous skin reactions, and erythema multiforme in patients treated with pregabalin (see [8.5 Post-Market Adverse Reactions](#)). Post-market reporting rate is generally accepted to be an underestimate due to under-reporting. Most of the reports were in patients taking concomitant medications also associated with the potential development of these serious skin reactions. Therefore, in most cases, causality in relation to pregabalin could not be clearly established. Patients should be advised that if they experience a skin rash, they should discontinue IPG-Pregabalin treatment and contact their physician for assessment and advice.

7.1 Special Populations

- **Renal**

There have been reports of patients, with or without previous history, experiencing renal failure while receiving pregabalin alone or in combination with other medications. Discontinuation of pregabalin showed reversibility of this event in some cases (see [7 WARNINGS AND PRECAUTIONS](#); [8.5 Post-Market Adverse Reactions](#); and [4 DOSAGE AND ADMINISTRATION](#)). Because pregabalin is eliminated primarily by renal excretion, the dose of pregabalin should be adjusted as noted for elderly patients or those with renal impairment (see [10 CLINICAL PHARMACOLOGY](#) and [4 DOSAGE AND ADMINISTRATION](#)).

- **Adjustment of Dose in Renally-Impaired Patients**

In patients with a medical history of significant renal insufficiency, daily dosages should be reduced accordingly (see [Table 1 in 4.1 Dosing Considerations](#)).

7.1.1 Pregnant Women

Pregabalin should not be used during pregnancy unless the benefit of the drug to the mother clearly outweighs the potential risk to the fetus. If women decide to become pregnant while taking pregabalin, the use of this product should be carefully re-evaluated. If therapy with pregabalin is considered necessary during pregnancy, the lowest possible therapeutic dose should be used.

Women of childbearing potential must use effective contraception during treatment with IPG-PREGABALIN.

Major congenital malformations

IPG-PREGABALIN use in the first-trimester of pregnancy can cause major birth defects in the unborn child.

Data from a Nordic observational study of more than 2700 pregnancies exposed to IPG-PREGABALIN in the first trimester showed a higher prevalence of major congenital malformations (MCM) among the pediatric population (live or stillborn) exposed to IPG-PREGABALIN compared to the unexposed population (5.9% vs. 4.1%).

In the Nordic study, the risk of MCM among the pediatric population exposed to IPG-PREGABALIN in the first trimester was slightly higher compared to the unexposed population (adjusted prevalence ratio and 95% confidence interval: 1.14 (0.96-1.35).

The analyses on specific malformations showed higher risks for malformations of the nervous system, the eye, orofacial clefts, urinary and genital malformations, but numbers were small and estimates imprecise.

Birth and postnatal neurodevelopmental outcomes

In the Nordic study, there was a significantly higher prevalence of stillbirth and “small for gestational age” in the IPG-PREGABALIN-exposed population compared to the unexposed population (adjusted prevalence ratio and 95% confidence interval: 1.72 (1.02-2.91) and 1.21 (1.01-1.44), respectively.

There were no statistically significant findings for low birth weight, preterm birth, low Apgar score, and microcephaly.

Pregnancy Registry

Patients should be encouraged to enroll in the North American Antiepileptic Drug (NAAED) Pregnancy Registry if they become pregnant. This registry is collecting information about the safety of anticonvulsant medications that can be taken by women during pregnancy to treat disorders such as epilepsy, mood disorder, and chronic pain. The primary goal is to determine the frequency of major malformations, such as heart defects, spina bifida and cleft lip, in the infants exposed during pregnancy to anticonvulsant drugs. To enroll, patients can call the toll-free number, 1-888-233-2334. Information on the registry can be also found at the website <http://www.aedpregnancyregistry.org/> (see [7 WARNINGS AND PRECAUTIONS, General](#), and [7 WARNINGS AND PRECAUTIONS, Patient Counselling Information](#)).

Labour and Delivery

The effects of pregabalin on labour and delivery in pregnant women are unknown. In the prenatal-postnatal study in rats, pregabalin prolonged gestation and induced dystocia at exposures ≥ 47 times the mean human exposure [$AUC_{(0-24)}$ of 123 mcg•hr/mL] at the maximum recommended clinical dose of 600 mg/day (see [16 NON-CLINICAL TOXICOLOGY](#)).

7.1.2 Breast-feeding

Pregabalin is excreted in the milk of lactating women. As the safety of pregabalin in infants is not known, breast-feeding is not recommended during treatment with pregabalin. A decision must be made whether to discontinue breast-feeding or to discontinue from pregabalin therapy taking into account the benefit of breast feeding for the child and the benefit of therapy for the women (see [16 NON-CLINICAL TOXICOLOGY](#)). Patients should be advised to notify their physician if they are breast-feeding.

The pharmacokinetics of 150 mg pregabalin given every 12 hours (300 mg daily dose) was evaluated in 10 lactating women who were at least 12 weeks postpartum. Pregabalin was excreted into breast milk with average peak and steady-state concentrations approximately 53 and 76% of those in maternal plasma, respectively. The estimated average daily infant dose of pregabalin from breast milk (assuming mean milk consumption of 150 mL/kg/day) was 0.31 mg/kg/day, which on a mg/kg basis would be approximately 7% of the total maternal daily dose.

7.1.3 Pediatrics

Pediatrics (< 18 years of age): The safety and efficacy of pregabalin in pediatric patients (< 18 years of age) have not been established and its use in this patient population is not indicated.

7.1.4 Geriatrics

Of the 1831 patients who received pregabalin in neuropathic pain studies, 528 were 65 to 74 years of age, and 452 were 75 years of age or older. No significant differences in efficacy were observed between these patients and younger patients. Pregabalin oral clearance tended to decrease with increasing age. This decrease in pregabalin oral clearance is consistent with age-related decreases in creatinine clearance. Reduction of pregabalin dose may be required in patients who have age-related compromised renal function. In general, the incidence of adverse events did not increase with age.

8 ADVERSE REACTIONS

8.2 Clinical Trial Adverse Reactions

Clinical trials are conducted under very specific conditions. The adverse reaction rates observed in the clinical trials; therefore, may not reflect the rates observed in practice and should not be compared to the rates in the clinical trials of another drug. Adverse reaction information from clinical trials may be useful in identifying and approximating rates of adverse drug reactions in real-world use.

In all controlled and uncontrolled trials during the pre-marketing development of pregabalin, more than 8666 patients have received pregabalin, with 83% of exposure at dosages of 300 mg/day or above and 32% at dosages of 600 mg/day or higher. Approximately 4010 patients had at least 6 months of exposure, 2415 had at least 1 year of exposure, and 939 had at least 2 years of exposure to pregabalin. In controlled trials, 1831 patients with neuropathic pain received pregabalin. In all controlled and uncontrolled studies in fibromyalgia, 3446 patients have received pregabalin doses of 150-600 mg/day. A total of 969 patients had at least 6 months of exposure and 440 patients had at least 1 year exposure to pregabalin. Doses above 600 mg/day have not been studied.

Most Common Adverse Events in All Pre-marketing Controlled Clinical Studies of Neuropathic Pain

The most commonly observed adverse events ($\geq 5\%$ and twice the rate of that seen in placebo) in pregabalin-treated patients in pre-marketing studies were: dizziness, somnolence, peripheral edema, and dry mouth. Adverse events were usually mild to moderate in intensity.

Discontinuation Due to Adverse Events in Pre-marketing Controlled Clinical Studies

In all pre-marketing controlled studies, the discontinuation rate due to adverse events was 14% for patients receiving pregabalin and 7% for patients receiving placebo. The most common reasons for discontinuation due to adverse events ($\geq 2\%$) in the pregabalin treatment groups were dizziness and somnolence. Other adverse events that led to withdrawal more frequently in the pregabalin group than the placebo group were ataxia (1%), and asthenia, confusion, headache and nausea ($< 1\%$ each).

In pre-marketing controlled neuropathic pain studies, the discontinuation rate due to adverse events was 11% for pregabalin and 5% for placebo. The most common reasons for discontinuation due to adverse events ($\geq 2\%$) in the pregabalin treatment groups were dizziness and somnolence. Other adverse events that led to withdrawal more frequently in the pregabalin group than the placebo group were confusion (1%) and asthenia, peripheral edema and ataxia ($< 1\%$ each).

Incidence of Adverse Events in Pre-marketing Controlled Clinical Studies of Neuropathic Pain

In summaries of adverse events, investigator's terms for individual adverse events have been grouped into a smaller number of standardized categories using the COSTART IV dictionary. The prescriber should be aware that the percentages in Table 2 through Table 9 cannot be used to predict the frequency of adverse events in the course of usual medical practice where patient characteristics and other factors may differ from those prevailing during clinical studies. Similarly, the cited frequencies cannot be directly compared with figures obtained from other clinical investigations involving different treatments, uses or investigators. An inspection of these frequencies, however, does provide the prescriber with one basis to estimate the relative contribution of drug and non-drug factors to the adverse event incidences in the population studied.

Adverse Events From Pre-marketing Controlled Clinical Studies of Neuropathic Pain

Diabetic Peripheral Neuropathy

Table 2 lists all adverse events, regardless of causality, occurring in $\geq 2\%$ of patients with neuropathic pain associated with diabetic peripheral neuropathy receiving pregabalin for at least one of the pregabalin groups, and for which the incidence was greater than in the placebo group. A majority of pregabalin-treated patients in these studies had adverse events with a maximum intensity of mild or moderate. In these studies, 979 patients received pregabalin and 459 patients received placebo for up to 13 weeks.

Table 2. Incidence (%) of Treatment-Emergent Adverse Events in Placebo-Controlled Studies in Neuropathic Pain Associated with Diabetic Peripheral Neuropathy (Events in at Least 2% of Patients Receiving Pregabalin and More Frequent Than in Placebo-Treated Patients)

Body System Preferred Term	Placebo (n = 459) %	Pregabalin (mg/day)			
		75 (n = 77) %	150 (n = 212) %	300 (n = 321) %	600 (n = 369) %
Body as a whole					
Infection	6.1	3.9	7.5	8.4	4.6
Asthenia	2.4	3.9	1.9	4.4	7.3
Pain	3.9	5.2	4.2	2.5	4.9
Accidental injury	2.8	5.2	2.4	2.2	5.7
Back pain	0.4	0.0	2.4	1.2	1.9
Chest pain	1.1	3.9	1.4	1.2	1.6
Face edema	0.4	0.0	0.9	0.9	2.2
Digestive system					
Dry mouth	1.1	2.6	1.9	4.7	6.5
Constipation	1.5	0.0	2.4	3.7	6.0
Diarrhea	4.8	5.2	2.8	1.9	3.0
Flatulence	1.3	2.6	0	2.2	2.7
Vomiting	1.5	1.3	0.9	2.2	1.1
Hemic and lymphatic system					
Ecchymosis	0.2	2.6	0.5	0.6	0.3
Metabolic and nutritional disorders					
Peripheral edema	2.4	3.9	6.1	9.3	12.5
Weight gain	0.4	0.0	4.2	3.7	6.2
Edema	0.0	0.0	1.9	4.0	1.9
Hypoglycemia	1.1	1.3	3.3	1.6	1.1

Body System Preferred Term	Placebo (n = 459) %	Pregabalin (mg/day)			
		75 (n = 77) %	150 (n = 212) %	300 (n = 321) %	600 (n = 369) %
Nervous system					
Dizziness	4.6	7.8	9.0	23.1	29.0
Somnolence	2.6	3.9	6.1	13.1	16.3
Neuropathy	3.5	9.1	1.9	2.2	5.4
Ataxia	1.3	6.5	0.9	2.2	4.3
Vertigo	1.1	1.3	1.9	2.5	3.5
Confusion	0.7	0.0	1.4	2.2	3.3
Euphoria	0.0	0.0	0.5	3.4	1.6
Thinking abnormal ^a	0.0	1.3	0.0	0.9	3.0
Abnormal gait	0.0	1.3	0.0	0.6	2.7
Reflexes decreased	1.7	3.9	0.5	1.2	1.4
Amnesia	0.2	2.6	0.9	0.0	2.2
Hypesthesia	0.7	2.6	0.0	0.0	0.8
Hyperalgesia	0.2	2.6	0.0	0.0	0.3
Respiratory system					
Dyspnea	0.7	2.6	0.0	1.9	1.9
Skin and appendages					
Pruritus	1.3	2.6	0.0	0.9	0.0
Special senses					
Blurred vision ^b	1.5	2.6	1.4	2.8	5.7
Conjunctivitis	0.2	2.6	1.4	0.6	0.3

^a Thinking abnormal primarily consists of events related to difficulty with concentration/attention but also includes events related to cognition and language problems and slow thinking.

^b Investigator term; summary level term is amblyopia.

Discontinuation in Controlled Clinical Studies of Diabetic Peripheral Neuropathy

Approximately 9% of patients receiving pregabalin and 4% receiving placebo discontinued from controlled diabetic peripheral neuropathy studies due to adverse events. The adverse events most commonly leading to discontinuation are presented in Table 3.

Table 3. Adverse Events Most Frequently ($\geq 2\%$ of patients) Leading to Discontinuation in Placebo-Controlled Studies in Patients With Neuropathic Pain Associated with Diabetic Peripheral Neuropathy

COSTART Preferred Term	Number (%) of Patients				
	Placebo (n = 459)	Pregabalin (mg/day)			
		75 (n = 77)	150 (n = 212)	300 (n = 321)	600 (n = 369)
Dizziness	2 (0.4)	0 (0.0)	3 (1.4)	6 (1.9)	21 (5.7)
Somnolence	0 (0.0)	0 (0.0)	0 (0.0)	5 (1.6)	15 (4.1)

Post-herpetic Neuralgia

Table 4 lists all adverse events, regardless of causality, occurring in $\geq 2\%$ of patients with neuropathic pain associated with post-herpetic neuralgia receiving pregabalin for at least one of the pregabalin groups, and for which the incidence was greater than in the placebo group. A majority of pregabalin-treated patients in these studies had adverse events with a maximum intensity of mild or moderate. In these studies, 852 patients received pregabalin and 398 patients received placebo for up to 13 weeks.

Table 4. Incidence (%) of Treatment-Emergent Adverse Events in Placebo-Controlled Studies in Neuropathic Pain Associated with Post-herpetic Neuralgia (Events in at Least 2% of Patients Receiving Pregabalin and More Frequent Than in Placebo-Treated Patients)

Body System Preferred Term	Placebo (n = 398) %	Pregabalin (mg/day)			
		75 (n = 84) %	150 (n = 302) %	300 (n = 312) %	600 (n = 154) %
		Body as a whole			
Infection	3.5	14.3	8.3	6.4	2.6
Headache	5.3	4.8	8.9	4.5	8.4
Pain	3.8	4.8	4.3	5.4	4.5
Asthenia	4.0	3.6	5.0	2.6	5.2
Accidental injury	1.5	3.6	2.6	3.2	5.2
Flu syndrome	1.3	1.2	1.7	2.2	1.3
Face edema	0.8	0.0	1.7	1.3	3.2
Malaise	1.0	2.4	0.3	0.6	0.0

Body System Preferred Term	Placebo (n = 398) %	Pregabalin (mg/day)			
		75 (n = 84) %	150 (n = 302) %	300 (n = 312) %	600 (n = 154) %
Cardiovascular system					
Vasodilatation	1.3	2.4	1.0	0.6	0.0
Digestive system					
Dry mouth	2.8	7.1	7.0	6.1	14.9
Constipation	2.3	3.6	4.6	5.4	5.2
Diarrhea	4.0	2.4	4.3	3.5	4.5
Flatulence	1.0	2.4	1.3	1.6	3.2
Vomiting	0.8	1.2	0.7	2.9	2.6
Metabolic and nutritional disorders					
Peripheral edema	3.5	0.0	7.9	15.7	16.2
Weight gain	0.3	1.2	1.7	5.4	6.5
Edema	1.3	0.0	1.0	2.2	5.8
Hyperglycemia	0.8	2.4	0.3	0.0	0.0

Body System Preferred Term	Placebo (n = 398) %	Pregabalin (mg/day)			
		75 (n = 84) %	150 (n = 302) %	300 (n = 312) %	600 (n = 154) %
Nervous system					
Dizziness	9.3	10.7	17.9	31.4	37.0
Somnolence	5.3	8.3	12.3	17.9	24.7
Ataxia	0.5	1.2	2.0	5.4	9.1
Abnormal gait	0.5	0.0	2.0	3.8	7.8
Confusion	0.3	1.2	2.3	2.9	6.5
Thinking abnormal ^a	1.5	0.0	1.7	1.3	5.8
Incoordination	0.0	2.4	1.7	1.3	2.6
Amnesia	0.0	0.0	1.0	1.3	3.9
Speech disorder	0.0	0.0	0.3	1.3	3.2
Insomnia	1.8	0.0	0.7	2.2	0.0
Euphoria	0.0	2.4	0.0	1.3	1.3
Nervousness	0.5	0.0	1.0	0.3	2.6
Tremor	1.5	1.2	0.0	1.0	2.6
Hallucinations	0.0	0.0	0.3	0.3	3.2
Hyperesthesia	0.3	2.4	0.3	0.0	1.3
Respiratory system					
Bronchitis	0.8	0.0	1.3	1.0	2.6
Pharyngitis	0.8	0.0	2.6	0.6	0.6
Rhinitis	1.8	1.2	0.7	0.6	3.2
Skin and appendages					
Rash	3.0	2.4	2.0	2.9	5.2
Special senses					
Blurred vision ^b	2.5	1.2	5.0	5.1	9.1
Diplopia	0.0	0.0	1.7	1.9	3.9
Abnormal vision	0.3	0.0	1.0	1.6	5.2

Body System Preferred Term	Placebo (n = 398) %	Pregabalin (mg/day)			
		75 (n = 84) %	150 (n = 302) %	300 (n = 312) %	600 (n = 154) %
Urogenital system					
Urinary tract infection	1.5	0.0	2.3	1.6	3.2

^a Thinking abnormal primarily consists of events related to difficulty with concentration/attention but also includes events related to cognition and language problems and slow thinking.

^b Investigator term; summary level term is amblyopia.

Discontinuation in Controlled Clinical Studies of Post-herpetic Neuralgia

Approximately 14% of patients receiving pregabalin and 7% receiving placebo discontinued from controlled post-herpetic neuralgia studies due to adverse events. The adverse events most commonly leading to discontinuation are presented in Table 5.

Table 5. Adverse Events Most Frequently ($\geq 2\%$ of patients) Leading to Discontinuation in Placebo-Controlled Studies in Patients With Neuropathic Pain Associated with Post-herpetic Neuralgia

COSTART Preferred Term	Placebo (n = 398)	Number (%) of Patients			
		75 (n = 84)	150 (n = 302)	300 (n = 312)	600 (n = 154)
Dizziness	3 (0.8)	0 (0.0)	11 (3.6)	12 (3.8)	12 (7.8)
Somnolence	1 (0.3)	0 (0.0)	6 (2.0)	12 (3.8)	10 (6.5)
Confusion	1 (0.3)	0 (0.0)	2 (0.7)	5 (1.6)	8 (5.2)
Peripheral edema	1 (0.3)	0 (0.0)	2 (0.7)	5 (1.6)	5 (3.2)
Ataxia	0 (0.0)	0 (0.0)	1 (0.3)	5 (1.6)	4 (2.6)
Abnormal gait	0 (0.0)	0 (0.0)	0 (0.0)	4 (1.3)	4 (2.6)
Hallucinations	0 (0.0)	0 (0.0)	0 (0.0)	1 (0.3)	4 (2.6)
Dry mouth	1 (0.3)	0 (0.0)	0 (0.0)	0 (0.0)	4 (2.6)

Incidence of Most Common Dose-Related Treatment-Emergent Adverse Events

Most common dose-related treatment-emergent adverse events are presented in Table 6 (diabetic peripheral neuropathy), Table 7 (post-herpetic neuralgia), and Table 8 (fibromyalgia).

Table 6. Incidence (%) of Most Common Dose-Related Treatment-Emergent Adverse Events in Placebo-Controlled Studies in Neuropathic Pain Associated with Diabetic Peripheral Neuropathy

Adverse Event Preferred Term	Placebo (n = 459) %	Pregabalin (mg/day)			
		75 (n = 77) %	150 (n = 212) %	300 (n = 321) %	600 (n = 369) %
Dizziness	4.6	7.8	9.0	23.1	29.0
Somnolence	2.6	3.9	6.1	13.1	16.3
Peripheral edema	2.4	3.9	6.1	9.3	12.5
Asthenia	2.4	3.9	1.9	4.4	7.3
Dry mouth	1.1	2.6	1.9	4.7	6.5
Weight gain	0.4	0.0	4.2	3.7	6.2
Constipation	1.5	0.0	2.4	3.7	6.0
Blurred vision ^a	1.5	2.6	1.4	2.8	5.7

^a Investigator term; summary level term is amblyopia.

Table 7. Incidence (%) of Most Common Dose-Related Treatment-Emergent Adverse Events in Placebo-Controlled Studies in Neuropathic Pain Associated with Postherpetic Neuralgia

Adverse Event Preferred Term	Placebo (n = 398) %	Pregabalin (mg/day)			
		75 (n = 84) %	150 (n = 302) %	300 (n = 312) %	600 (n = 154) %
Dizziness	9.3	10.7	17.9	31.4	37.0
Somnolence	5.3	8.3	12.3	17.9	24.7
Peripheral edema	3.5	0.0	7.9	15.7	16.2
Dry mouth	2.8	7.1	7.0	6.1	14.9
Blurred vision ^a	2.5	1.2	5.0	5.1	9.1
Ataxia	0.5	1.2	2.0	5.4	9.1
Weight gain	0.3	1.2	1.7	5.4	6.5
Abnormal gait	0.5	0.0	2.0	3.8	7.8

^a Investigator term; summary level term is amblyopia.

Table 8. Incidence (%) of Most Common Dose-Related Treatment-Emergent Adverse Events in Placebo-Controlled Studies in Pain Associated with Fibromyalgia

System Organ Class and Preferred Term (MedDRA version 10.1)	Placebo (n = 689) %	Pregabalin (mg/day)			
		150 (n = 132) %	300 (n = 685) %	450 (n = 687) %	600 (n = 564) %
Dizziness	10.4	22.7	32.6	42.5	46.5
Somnolence	4.6	12.9	18.5	19.9	20.7
Weight increased	2.5	7.6	11.1	10.0	13.7
Peripheral edema	2.5	5.3	6.7	6.4	10.8
Dry mouth	1.7	6.8	6.7	9.2	9.4
Constipation	2.8	3.8	5.8	6.8	9.2
Fatigue	5.4	4.5	7.2	8.4	8.2
Balance disorder	0.1	1.5	3.2	4.9	6.9
Disturbance in attention	1.3	3.8	4.4	6.4	6.9
Increased appetite	1.3	3.8	3.4	4.5	5.5
Euphoria	0.9	1.5	4.1	4.8	5.1

Most Common Adverse Events in Controlled Clinical Studies in Fibromyalgia

The most commonly observed treatment-related adverse events ($\geq 5\%$ and twice the rate of that seen in placebo) in pregabalin-treated patients were: dizziness (37.5%), somnolence (18.6%), weight gain (10.6%), dry mouth (7.9%), blurred vision (6.7%), peripheral edema (6.1%), constipation (5.8%), and disturbance in attention (5.3%). Adverse events were usually mild to moderate in intensity.

Adverse Events from Controlled Clinical Studies in Fibromyalgia

Table 9 lists all adverse events occurring in $\geq 2\%$ of patients receiving pregabalin and for which the incidence was greater than in the placebo group. A majority of pregabalin-treated patients in these studies had adverse events with mild or moderate intensity. There was a dose-related increase in the incidence of adverse events (see [Table 8](#)). Severe adverse events occurred at roughly the same rate in low versus high doses of pregabalin.

Table 9. Incidence (%) of Treatment-Emergent Adverse Events in Controlled Fibromyalgia Studies (in at Least 2% of Pregabalin-Treated Patients and More Frequent Than in Placebo-Treated Patients)

System Organ Class and Preferred Term (MedDRA version 10.1)	Placebo (n = 689) %	Pregabalin (mg/day)			
		150 (n = 132) %	300 (n = 685) %	450 (n = 687) %	600 (n = 564) %
Ear and labyrinth disorders					
Vertigo	0.9	1.5	3.1	3.2	3.5
Eye disorders					
Blurred vision ^a	1.0	8.3	5.8	6.4	10.1
Gastrointestinal disorders					
Dry mouth	1.7	6.8	6.7	9.2	9.4
Constipation	2.8	3.8	5.8	6.8	9.2
Abdominal distention	1.5	2.3	2.2	1.9	2.0
Flatulence	1.0	0.8	0.9	2.0	2.3
General disorders and administration site conditions					
Fatigue	5.4	4.5	7.2	8.4	8.2
Peripheral edema	2.5	5.3	6.7	6.4	10.8
Feeling abnormal	0.3	0.8	1.9	1.9	2.0
Feeling drunk	0	0.8	2.3	1.5	2.1
Infections and infestations					
Sinusitis	3.0	3.8	3.6	5.2	4.1
Investigations					
Weight increased	2.5	7.6	11.1	10.9	13.7
Metabolism and nutritional disorders					
Increased appetite	1.3	3.8	3.4	4.5	5.5
Fluid retention	0.7	1.5	2.2	2.0	2.0
Musculoskeletal and connective tissue disorders					
Arthralgia	2.5	3.8	3.6	3.2	4.6
Muscle spasm	1.9	2.3	3.4	3.3	3.2

System Organ Class and Preferred Term (MedDRA version 10.1)	Placebo (n = 689) %	Pregabalin (mg/day)			
		150 (n = 132) %	300 (n = 685) %	450 (n = 687) %	600 (n = 564) %
Nervous system disorders					
Dizziness	10.4	22.7	32.6	42.5	46.5
Somnolence	4.6	12.9	18.5	19.9	20.7
Disturbance in attention	1.3	3.8	4.4	6.4	6.9
Balance Disorder	0.1	1.5	3.2	4.9	6.9
Memory Impairment	0.6	0.8	2.6	3.5	3.5
Hypoesthesia	0.6	1.5	2.0	2.8	2.3
Tremor	0.6	0	0.6	2.9	3.0
Lethargy	0.4	2.3	1.3	0.7	1.4
Psychiatric disorders					
Euphoria	0.9	1.5	4.1	4.8	5.1
Anxiety	0.9	1.5	1.9	2.5	1.8
Confusion	0.1	0	2.0	1.9	2.7

^a Investigator term; summary level term is amblyopia.

Discontinuation Due to Adverse Events in Controlled Clinical Studies in Fibromyalgia

Approximately 20% of patients receiving pregabalin and 11% receiving placebo discontinued due to adverse events. The adverse events most commonly leading to discontinuation were dizziness (6.1%) and somnolence (3.3%) as presented in Table 10. Other events leading to discontinuation in clinical trials of fibromyalgia included weight gain (1.1%), vision blurred (0.8%) and peripheral edema (0.6%). There was a dose-dependent increase in rate of discontinuation due to adverse events.

Table 10. Adverse Events Most Frequently (≥ 2% of patients) Leading to Discontinuation in Placebo-Controlled Studies in Patients With Pain Associated with Fibromyalgia

System Organ Class and Preferred Term (MedDRA version 10.1)	Number (%) of Patients				
	Placebo (n = 689)	150 (n=132)	300 (n = 675)	450 (n = 687)	600 (n = 564)
Dizziness	0.4%	1.5%	4.1%	6.6%	9.2%
Somnolence	0.1%	0.8%	2.9%	3.2%	4.6%

Adverse Events Following Abrupt or Rapid Discontinuation

Following abrupt or rapid discontinuation of pregabalin, some patients reported symptoms including insomnia, nausea, headache, anxiety, hyperhidrosis, and diarrhea. Pregabalin should be tapered gradually over a minimum of one week rather than discontinued abruptly (see [7 WARNINGS AND PRECAUTIONS, General](#) and [7 WARNINGS AND PRECAUTIONS, Abrupt or Rapid Discontinuation](#)).

Other Events Observed During the Premarketing Evaluation of IPG-PREGABALIN

Following is a list of treatment-emergent adverse events reported during premarketing assessment of pregabalin in clinical trials (over 8600 adult subjects) except those already listed in the previous tables or elsewhere in labeling. In the tabulations that follow, a COSTART-based dictionary of terminology has been used to classify reported adverse events. The frequencies presented, therefore, represent the proportion of the over 8600 adult individuals exposed to multiple doses of pregabalin who experienced an event of the type cited on at least 1 occasion while receiving pregabalin. It is important to emphasize that although the events reported occurred during treatment with IPG-PREGABALIN, they were not necessarily caused by it.

8.3 Less Common Clinical Trial Adverse Reactions (< 2%)

Events are further categorized by body system and listed in order of decreasing frequency according to the following definitions: frequent adverse events are those occurring on 1 or more occasions in at least 1/100 patients; infrequent adverse events are those occurring in 1/100 to 1/1000 patients; rare events are those occurring in fewer than 1/1000 patients.

Body System	Adverse Events
Body as a whole	
Frequent	Flu syndrome, back pain, allergic reaction, fever, generalized edema
Infrequent	Neck pain, neoplasm, cellulitis, cyst, chills, malaise, overdose, moniliasis, hernia, viral infection, photosensitivity reaction, pelvic pain, abdomen enlarged, abscess, neck rigidity, lab test abnormal, drug level increased, carcinoma, sepsis, suicide attempt, reaction unevaluable
Rare	Infection fungal, unexpected benefit, chills and fever, body odor, drug level decreased, halitosis, hangover effect, injection site reaction, hormone level altered, hypothermia, infection bacterial, injection site hemorrhage, intentional overdose, mucous membrane disorder, accidental overdose, adenoma, anaphylactoid reaction, ascites, chest pain substernal, death, sarcoidosis, sudden death, immune system disorder, increased drug effect, injection site pain, Lupus Erythematosus syndrome, medication error, sarcoma, shock, tolerance decreased

Body System	Adverse Events
Cardiovascular	
Frequent	Hypertension, vasodilatation
Infrequent	Palpitation, migraine, tachycardia, peripheral vascular disorder, electrocardiogram abnormal, cardiovascular disorder, angina pectoris, congestive heart failure, hemorrhage, myocardial infarct, hypotension, postural hypotension, ventricular extrasystoles, atrial fibrillation, coronary artery disorder, bradycardia, cerebrovascular accident, arrhythmia, cerebral ischemia, vascular disorder, sinus bradycardia, myocardial ischemia, bundle branch block, AV block first degree, arteriosclerosis, deep thrombophlebitis, phlebitis, arterial anomaly, heart failure, pulmonary embolus, retinal vascular disorder, varicose vein
Rare	Heart arrest, vascular anomaly, occlusion, supraventricular tachycardia, atrial arrhythmia, atrial flutter, cerebral infarct, coronary occlusion, thrombophlebitis, thrombosis, cardiomegaly, extrasystoles, pallor, AV block, AV block second degree, cardiomyopathy, peripheral gangrene, QT interval prolonged, retinal artery occlusion, supraventricular extrasystoles, cerebral hemorrhage, digitalis intoxication, ventricular arrhythmia, aortic stenosis, bigeminy, cerebrovascular disorder, left heart failure, ventricular tachycardia, AV block complete, carotid occlusion, carotid thrombosis, cor-pulmonale, embolus lower extremity, endocarditis, heart block, increased capillary fragility, intracranial aneurysm, nodal tachycardia, QT interval shortened, retinal vein thrombosis, ST elevated, T inverted, vascular headache, vasculitis
Digestive system	
Frequent	Nausea, diarrhea, anorexia, gastrointestinal disorder
Infrequent	Gastroenteritis, tooth disorder, periodontal abscess, colitis, gastritis, liver function tests abnormal, increased salivation, thirst, nausea and vomiting, rectal disorder, gingivitis, dysphagia, stomatitis, mouth ulceration, cholelithiasis, rectal hemorrhage, gastrointestinal hemorrhage, glossitis, tooth caries, abnormal stools, cholecystitis, melena, oral moniliasis, esophagitis, tongue disorder, cheilitis, tongue edema

Body System	Adverse Events
Rare	Eructation, pancreatitis, stomach ulcer, ulcerative stomatitis, esophageal stenosis, fecal incontinence, gum hemorrhage, intestinal obstruction, enteritis, peptic ulcer, enterocolitis, gum hyperplasia, hepatomegaly, liver fatty deposit, tenesmus, biliary pain, fecal impaction, jaundice, periodontitis, ulcerative colitis, aphthous stomatitis, cholestatic jaundice, gastrointestinal carcinoma, hemorrhagic gastritis, hepatitis, liver tenderness, nausea, vomiting and diarrhea, salivary gland enlargement, stomach atony, bloody diarrhea, cardiospasm, duodenal ulcer, gamma glutamyl transpeptidase increased, hematemesis, hepatoma, intestinal perforation, intestinal stenosis, intestinal ulcer, leukoplakia of mouth, necrotizing pancreatitis, pancreas disorder, pseudomembranous colitis, sialadenitis, stomach ulcer hemorrhage, tongue discoloration
Endocrine system	
Infrequent	Diabetes mellitus, hypothyroidism
Rare	Goiter, prolactin increased, thyroid disorder, gonadotropic follicle stim hormone increase, hyperthyroidism, thyroiditis, adrenal insufficiency, parathyroid disorder, thyroid carcinoma, thyroid neoplasia, virilism
Hemic and lymphatic	
Infrequent	Anemia, leukopenia, thrombocytopenia, lymphadenopathy, hypochromic anemia, leukocytosis, eosinophilia
Rare	Lymphocytosis, petechia, iron deficiency anemia, cyanosis, lymphedema, polycythemia, lymphoma like reaction, megaloblastic anemia, splenomegaly, purpura, thrombocythemia, thrombocytopenic purpura, chronic leukemia, coagulation disorder, erythrocytes abnormal, leukemoid reaction, lymphangitis, macrocytic anemia, neutropenia, pancytopenia, prothrombin decreased, rupture of spleen, sedimentation rate increased
Metabolic and nutritional	
Infrequent	Hyperglycemia, SGPT increased, hypoglycemia, hypokalemia, hypercholesteremia, SGOT increased, weight loss, hyperlipemia, amylase increased, hyperuricemia, alkaline phosphatase increased, creatinine increased, hyponatremia, gout, dehydration, BUN increased, healing abnormal
Rare	Hypercalcemia, hyperkalemia, hypocalcemia, bilirubinemia, alcohol intolerance, hypoglycemic reaction, ketosis, calcium disorder, hypochloremia, hypomagnesemia, hypoproteinemia, NPN increased, uremia, acidosis, avitaminosis, enzymatic abnormality, gamma globulins increased, hypernatremia, hypophosphatemia, lactic acidosis, obesity

Body System	Adverse Events
Musculoskeletal system	
Frequent	Arthralgia, myalgia, arthritis, leg cramps, myasthenia
Infrequent	Tendon disorder, arthrosis, joint disorder, bone disorder, tenosynovitis, bursitis, tendinous contracture, osteoporosis, tendon rupture, bone pain
Rare	Rheumatoid arthritis, osteomyelitis, rhabdomyolysis, myopathy, muscle atrophy, myositis, pyogenic arthritis, bone neoplasm, musculoskeletal congenital anomaly, pathological fracture
Nervous system	
Frequent	Insomnia, anxiety, libido decreased, depersonalization, hypertonia, neuropathy
Infrequent	Reflexes decreased, sleep disorder, abnormal dreams, hostility, hallucinations, hyperkinesia, personality disorder, dysarthria, hyperesthesia, hypokinesia, circumoral paresthesia, libido increased, neuralgia, vestibular disorder, aphasia, movement disorder, hyperalgesia, apathy, hypotonia, convulsion, facial paralysis, psychosis
Rare	Drug dependence, neuritis, paranoid reaction, CNS depression, CNS neoplasia, manic reaction, neurosis, extrapyramidal syndrome, meningitis, hemiplegia, reflexes increased, akathisia, delirium, paralysis, withdrawal syndrome, brain edema, CNS stimulation, dyskinesia, encephalopathy, foot drop, grand mal convulsion, hypalgesia, peripheral neuritis, psychotic depression, addiction, arachnoiditis, cerebellar syndrome, cogwheel rigidity, dementia, dystonia, Guillain-Barre syndrome, intracranial hemorrhage, multiple sclerosis, myelitis, schizophrenic reaction, subarachnoid hemorrhage, torticollis
Respiratory system	
Frequent	Sinusitis, rhinitis, dyspnea, cough increased, pneumonia, lung disorder
Infrequent	Asthma, epistaxis, laryngitis, voice alteration, respiratory disorder, sputum increased
Rare	Apnea, emphysema, aspiration pneumonia, hyperventilation, lung edema, pleural disorder, atelectasis, hemoptysis, hiccup, hypoxia, laryngismus, lung fibrosis, pleural effusion, lung function decreased, pulmonary hypertension, yawn, bronchiectasis, bronchiolitis, carcinoma of lung, hypoventilation, laryngeal neoplasia, nasal septum disorder, pneumothorax

Body System	Adverse Events
Skin and appendages	
Infrequent	Pruritus, sweating, skin disorder, acne, dry skin, alopecia, skin ulcer, herpes simplex, urticaria, nail disorder, eczema, herpes zoster, skin benign neoplasm, fungal dermatitis, maculopapular rash, vesiculobullous rash, skin carcinoma, furunculosis, skin discoloration, skin hypertrophy, psoriasis, seborrhea, hirsutism
Rare	Skin nodule, angioedema, cutaneous moniliasis, skin atrophy, exfoliative dermatitis, pustular rash, ichthyosis, skin melanoma, subcutaneous nodule, sweating decreased, hair disorder, lichenoid dermatitis, melanosis, miliaria, purpuric rash, skin necrosis, Stevens Johnson Syndrome
Special sense	
Frequent	Eye disorder, conjunctivitis, otitis media
Infrequent	Retinal disorder, tinnitus, eye pain, cataract specified, dry eyes, taste perversion, ear pain, lacrimation disorder, ear disorder, deafness, eye hemorrhage, photophobia, glaucoma, vitreous disorder, corneal lesion, otitis externa, refraction disorder, blepharitis, retinal edema, taste loss, abnormality of accommodation
Rare	Hyperacusis, keratitis, mydriasis, parosmia, ptosis, retinal hemorrhage, color blindness, retinal depigmentation, retinal detachment, corneal opacity, corneal ulcer, iritis, night blindness, optic atrophy, retinal degeneration, cataract NOS, scleritis, strabismus, anisocoria, blindness, exophthalmos, keratoconjunctivitis, ophthalmoplegia, papilledema
Urogenital system	
Frequent	Anorgasmia
Infrequent	Urinary frequency, urinary incontinence, cystitis, abnormal ejaculation, urination impaired, dysuria, metrorrhagia, hematuria, vaginal moniliasis, prostatic disorder, vaginitis, dysmenorrhea, urinary urgency, kidney calculus, breast pain, menstrual disorder, amenorrhea, menorrhagia, kidney function abnormal, nephritis, urine abnormality, vaginal hemorrhage, urinary retention, urinary tract disorder, leukorrhea, breast neoplasm, menopause, oliguria, polyuria, albuminuria, pyuria

Body System	Adverse Events
Rare	Breast carcinoma, penis disorder, papanicolau smear suspicious, fibrocystic breast, prostatic carcinoma, uterine fibroids enlarged, acute kidney failure, creatinine clearance decreased, nephrosis, nocturia, polycystic kidney, bladder carcinoma, breast enlargement, cervicitis, cervix disorder, female lactation, glycosuria, gynecomastia, hypomenorrhea, kidney pain, mastitis, pyelonephritis, kidney failure, breast abscess, epididymitis, orchitis, prostate neoplasia, prostatic specific antigen increase, salpingitis, urogenital disorder, urolithiasis, uterine disorder, vulvovaginal disorder, balanitis, bladder calculus, calcium crystalluria, cervix neoplasm, dyspareunia, endometrial carcinoma, endometrial disorder, glomerulitis, hydronephrosis, ovarian cancer, unintended pregnancy, urethral pain, urethritis, urogenital anomaly, urogenital neoplasia, uterine hemorrhage

Comparison of Gender and Race

The overall adverse event profile of pregabalin was similar between women and men. There are insufficient data to support a statement regarding the distribution of adverse experience reports by race.

Peripheral Edema

Incidence of peripheral edema in pre-marketing controlled peripheral neuropathic pain studies was 10.4% in the pregabalin group compared with 2.9% in the placebo group. The incidence in the controlled fibromyalgia studies was 7.6% in the pregabalin group compared with 2.5% in the placebo group. In clinical trials, these events of peripheral edema were dose-related, mostly mild to moderate in intensity and rarely led to withdrawal. Peripheral edema was not associated with cardiovascular complications such as hypertension or congestive heart failure and there was no evidence of hemodilution or changes in any laboratory parameters indicative of underlying organ dysfunction (see [7 WARNINGS AND PRECAUTIONS, General](#) and [7 WARNINGS AND PRECAUTIONS, Peripheral Edema](#)).

Weight Gain

In the pre-marketing controlled peripheral neuropathic pain studies, patients on pregabalin had a higher incidence (5.9%) of weight gain as defined by a $\geq 7\%$ increase from baseline weight as compared with the placebo group (1.6%). The mean change in the pregabalin group was an increase of 1.5 kg compared with 0.2 kg in the placebo group; few patients (0.1%) withdrew due to weight gain.

The incidence in the controlled fibromyalgia studies was 11.5% in the pregabalin group compared with 2.5% in the placebo group; approximately 1% of pregabalin-treated patients withdrew due to weight gain.

This weight gain was dose-related, and not associated with clinically important changes in blood pressure or cardiovascular adverse events. There was no relationship between baseline body mass index and the incidence of $\geq 7\%$ weight gain in the controlled trials.

Based on the results of a controlled study of reproductive function in healthy male volunteers, the $\geq 7\%$ weight gain on pregabalin appeared to be reversible. In this study, there were no reports of peripheral edema (see [7 WARNINGS AND PRECAUTIONS, General](#), and [7 WARNINGS AND PRECAUTIONS, Weight Gain](#)).

8.4 Abnormal Laboratory Findings: Hematologic, Clinical Chemistry and Other Quantitative Data

Clinical Trial Findings

In all pre-marketing controlled trials, 1.0% of patients on pregabalin and 0.5% of placebo patients had an increase in creatine kinase of $\geq 3X$ upper limit of normal. Renal dysfunction was generally not associated with the elevated creatine kinase in these patients. Mean changes in creatine kinase ranged from 9.6 to 26.3 U/L for pregabalin-treated patients and 4.8 U/L for the placebo patients (see [4.1 Dosing Considerations, Patients With Impaired Renal Function](#)). Routine therapeutic drug monitoring or clinical laboratory testing is not required for patients treated with IPG-PREGABALIN (see [7 WARNINGS AND PRECAUTIONS, Creatinine Kinase Elevations](#)).

8.5 Post-Market Adverse Reactions

Since the first global approval of IPG-PREGABALIN on 06 July 2004 through 31 March 2012, there has been an estimated 15,951,859 patient-years of exposure to IPG-PREGABALIN. Table 11 is based on post-market spontaneous adverse event reports. The percentages shown are calculated by dividing the number of adverse events reported to the company by the estimated number of patient years exposed to pregabalin. Because these adverse reactions are reported spontaneously from a population of uncertain size, it is not possible to reliably estimate their frequency. Furthermore, a causal relationship between pregabalin and the emergence of these events has not been clearly established.

Table 11. Post-Market Spontaneous Adverse Event Reports

Adverse Event	Frequency			
	Common $\geq 1\%$	Uncommon < 1% and $\geq 0.1\%$	Rare < 0.1% and $\geq 0.01\%$	Very rare < 0.01%
Cardiovascular				
congestive heart failure ^a				X
Eye disorders^b				
diplopia				X
vision blurred				X
visual disturbance				X
vision loss				X

Adverse Event	Frequency			
	Common ≥ 1%	Uncommon < 1% and ≥ 0.1%	Rare < 0.1% and ≥ 0.01%	Very rare < 0.01%
Psychiatric disorders				
aggression				X
confusional state				X
depression				X
euphoric mood				X
insomnia				X
psychotic disorder ^c				X
Gastrointestinal disorders				
diarrhea				X
dry mouth				X
nausea				X
vomiting				X
intestinal obstruction ^d				X
paralytic ileus ^d				X
General disorders and administration site conditions				
fatigue				X
feeling abnormal				X
pain				X
Immune system disorders				
angioedema ^e				X
hypersensitivity ^f				X
Nervous system disorders				
ataxia				X
coordination abnormal				X
dizziness ^g				X
dysarthria				X
headache				X

Adverse Event	Frequency			
	Common ≥ 1%	Uncommon < 1% and ≥ 0.1%	Rare < 0.1% and ≥ 0.01%	Very rare < 0.01%
memory impairment				X
paresthesia				X
somnolence ^g				X
speech disorder				X
tremor				X
coma				X
loss of consciousness				X
mental impairment				X
Renal and urinary disorders				
urinary retention				X
Respiratory, thoracic and mediastinal disorders				
dyspnea				X
pulmonary edema				X
Skin and subcutaneous tissue disorders				
hyperhidrosis				X
pruritus				X

^a These reactions are mostly seen in elderly cardiovascular compromised patients during pregabalin treatment for a neuropathic pain indication (see [7 WARNINGS AND PRECAUTIONS, Congestive Heart Failure](#))

^b There have also been rare reports of accommodation disorder, eyelid edema and eye redness (see [7 WARNINGS AND PRECAUTIONS, Ophthalmological Effects](#))

^c There have been rare reports of psychotic disorders in patients receiving pregabalin

^d See below and [7 WARNINGS AND PRECAUTIONS, Gastrointestinal](#)

^e see [7 WARNINGS AND PRECAUTIONS, Angioedema](#)

^f see [7 WARNINGS AND PRECAUTIONS, Hypersensitivity](#)

^g see [7 WARNINGS AND PRECAUTIONS, Dizziness and Somnolence](#)

Gastrointestinal: There have been post-marketing events related to reduced lower gastrointestinal tract function (e.g., intestinal obstruction, paralytic ileus, and constipation) primarily reported when pregabalin was given in combination with other medications that have the potential to produce constipation, such as opioid analgesics (see [7 WARNINGS AND PRECAUTIONS, Gastrointestinal](#)).

Urinary and Renal Disorder: renal failure. There have been rare post-marketing reports of patients, with or without previous history, experiencing renal failure while receiving pregabalin alone or in combination with other medications. Discontinuation of pregabalin showed reversibility of this event in some cases. Caution is advised when prescribing IPG-PREGABALIN to the elderly or those with impaired renal function (see [7.1 Special Populations, Renal](#); [7 WARNINGS and PRECAUTIONS, Renal Failure](#), and [4 DOSAGE AND ADMINISTRATION](#)).

Cardiovascular: congestive heart failure. These reactions are mostly seen in elderly cardiovascular compromised patients during pregabalin treatment for a neuropathic pain indication (see [7 WARNINGS AND PRECAUTIONS, Congestive Heart Failure](#)).

Respiratory, Thoracic and Mediastinal Disorder: pulmonary edema. There have been post-marketing reports of pulmonary edema in patients receiving pregabalin. Although this adverse reaction has mostly been observed in elderly patients with predisposing factors, some cases have occurred in patients with no known previous history or predisposing conditions.

Eye disorders: diplopia, vision blurred, visual disturbance and vision loss. There have also been rare reports of accommodation disorder, eyelid edema and eye redness (see [7 WARNINGS AND PRECAUTIONS, Ophthalmological Effects](#)).

Encephalopathy: There have been serious post-marketing reports of encephalopathy, mostly in patients with underlying conditions that may precipitate encephalopathy. Some cases were reported in patients with a history of kidney or liver disease. Since there have been rare reports of renal failure with pregabalin, specific caution should be exercised when prescribing IPG-PREGABALIN to the elderly with age-related compromised renal function and patients with kidney disease or risk factors for renal failure (see [7 WARNINGS AND PRECAUTIONS, Renal Failure](#) and [7 WARNINGS AND PRECAUTIONS, Encephalopathy](#)).

Convulsions: convulsions, including status epilepticus and grand mal convulsions, have been reported in non-epileptic patients during treatment with IPG-PREGABALIN or following abrupt discontinuation (see [7 WARNINGS AND PRECAUTIONS, Abrupt or Rapid Discontinuation](#)).

Suicidal Behaviour and Ideation: There have been post-marketing reports of suicide-related events, including completed suicide, suicide attempt, and suicidal ideation in patients treated with pregabalin for a variety of indications such as neuropathic pain, fibromyalgia, etc. In some of these reports, underlying psychiatric disorders may have contributed to the event. The mechanism of this risk is not known. Patients should be monitored for signs of suicidal ideation and behaviours and appropriate treatment should be considered. Patients should be encouraged to report any distressing thoughts or feelings at anytime to their healthcare professional (see [7 WARNINGS AND PRECAUTIONS, Suicidal Behaviour and Ideation](#)).

Serious Skin Reactions

There have been very rare post-marketing reports of serious cutaneous reactions, including Stevens-Johnson Syndrome (SJS), Toxic Epidermal Necrolysis (TEN), dermatitis exfoliative, bullous skin reactions, and erythema multiforme in patients treated with IPG-PREGABALIN (see [7 WARNINGS AND PRECAUTIONS](#)). Post-market reporting rate is generally accepted to be an underestimate due to under-reporting. Most of the reports were in patients taking concomitant medications also associated with the potential development of these serious skin reactions. Therefore, in most cases, causality in relation to pregabalin could not be clearly established.

Patients should be advised that if they experience a skin rash, they should discontinue IPG-PREGABALIN treatment and contact their physician for assessment and advice.

Drug Abuse and Dependence/Liability

In a study of recreational users (n = 15) of sedative/hypnotic drugs, including alcohol, a single dose of pregabalin 450 mg received subjective ratings of “good drug effect”, “high”, and “liking” to a degree that was similar to a single dose of diazepam 30 mg. In controlled clinical studies in over 5500 patients, 4% of pregabalin -treated patients and 1% of placebo-treated patients overall reported euphoria as an adverse event. However, in clinical trials of diabetic peripheral neuropathy, euphoria was reported as an adverse event by 1.8% of pregabalin-treated patients and 0 % of placebo-treated patients, and in clinical trials of post-herpetic neuralgia, euphoria was reported as an adverse event by 0.9% of pregabalin-treated patients and 0% of placebo- treated patients. In clinical studies, following abrupt or rapid discontinuation of pregabalin, some patients reported symptoms including insomnia, nausea, headache, anxiety, hyperhidrosis, or diarrhea suggestive of physical dependence (see [7 WARNINGS AND PRECAUTIONS, Abrupt or Rapid Discontinuation](#)).

Pregabalin is not known to be active at receptor sites associated with drugs of abuse. As with any CNS active drug, physicians should carefully evaluate patients for history of drug abuse and observe them for signs of pregabalin misuse or abuse (e.g., development of tolerance, dose escalation, drug-seeking behaviour).

9 DRUG INTERACTIONS

9.2 Drug Interactions Overview

Since pregabalin is predominately excreted unchanged in the urine, undergoes negligible metabolism in humans ($\leq 2\%$ of a dose recovered in urine as metabolites), does not inhibit drug metabolism *in vitro*, and is not bound to plasma proteins, pregabalin is unlikely to produce, or be subject to, pharmacokinetic interactions.

9.4 Drug-Drug Interactions

Pharmacokinetic

In Vitro Studies: *In vitro* drug metabolism studies revealed that pregabalin at concentrations which were, in general, 10-fold greater than observed in Phase 2/3 clinical trials, does not inhibit human CYP1A2, CYP2A6, CYP2C9, CYP2C19, CYP2D6, CYP2E1, and CYP3A4 enzyme systems.

In Vivo Studies: The drug interaction data described in this section were obtained from studies involving healthy adults, patients with epilepsy, and patients with chronic pain disorders.

Carbamazepine, valproic acid, lamotrigine, phenytoin, phenobarbital, and topiramate: *In vitro* and *in vivo* studies showed that pregabalin is unlikely to be involved in significant pharmacokinetic drug interactions. Specifically, there are no clinically significant pharmacokinetic interactions between pregabalin and the following antiepileptic drugs: carbamazepine, valproic acid, lamotrigine, phenytoin, phenobarbital, and topiramate. Important pharmacokinetic interactions would also not be expected to occur between pregabalin and commonly used antiepileptic drugs.

Tiagabine: The results of a population pharmacokinetic analysis indicated that in patients with partial seizures tiagabine had no clinically significant effect on pregabalin clearance.

Gabapentin: The pharmacokinetics of pregabalin and gabapentin were investigated in 12 healthy subjects following concomitant single dose administration of 100 mg pregabalin and 300 mg gabapentin, and in 18 healthy subjects following concomitant multiple dose administration of 200 mg pregabalin q8h and 400 mg gabapentin q8h. Gabapentin pharmacokinetics following single and multiple dose administration were unaltered by pregabalin co-administration. The rate of pregabalin absorption was reduced by approximately 26% (single dose administration) and 18% (multiple dose administration) based on lower C_{max} values; however, the extent of pregabalin absorption was unaffected by gabapentin co-administration.

Oral Contraceptives: Pregabalin co-administration (200 mg TID) had no effect on the steady state pharmacokinetics of norethindrone and ethinyl estradiol (1 mg/35 mcg, respectively) in healthy subjects.

Lorazepam: Multiple dose administration of pregabalin (300 mg BID) in healthy subjects had no effect on the rate and extent of lorazepam single dose pharmacokinetics and single dose administration of lorazepam (1 mg) had no clinically significant effect on the steady state pharmacokinetics of pregabalin.

Oxycodone: Multiple dose administration of pregabalin (300 mg BID) in healthy subjects had no effect on the rate and extent of oxycodone single dose pharmacokinetics. Single dose administration of oxycodone (10 mg) had no clinically significant effect on the steady state pharmacokinetics of pregabalin.

Ethanol: Multiple dose administration of pregabalin (300 mg BID) in healthy subjects had no effect on the rate and extent of ethanol single dose pharmacokinetics and single dose administration of ethanol (0.7 g/kg) had no clinically significant effect on the steady state pharmacokinetics of pregabalin.

Diuretics, Oral Hypoglycemics, and Insulin: A population pharmacokinetic analysis in patients with chronic pain showed no clinically significant effect on pregabalin clearance with the concomitant use of diuretics, oral hypoglycemics, and insulin.

Pharmacodynamic

Opioids, benzodiazepines and alcohol: Pregabalin appears to be additive in the impairment of cognitive and gross motor function caused by opioids, benzodiazepines and alcohol.

In post-marketing experience, there are reports of respiratory failure, coma and deaths in patients taking pregabalin alone or in combination with other CNS depressants, including in patients with substance use disorders.

Thiazolidinedione Antidiabetic Agents: Higher frequencies of weight gain and peripheral edema were observed in patients taking both IPG-PREGABALIN and a thiazolidinedione antidiabetic agent compared to patients taking either drug alone. The majority of patients using thiazolidinedione antidiabetic agents in the overall safety database were participants in studies of pain associated with diabetic peripheral neuropathy.

As the thiazolidinedione class of antidiabetic drugs or IPG-PREGABALIN can cause weight gain and/or fluid retention alone or together, possibly exacerbating or leading to heart failure, caution should be exercised when co-administering IPG-PREGABALIN and these agents (see [7 WARNINGS AND PRECAUTIONS, Peripheral Edema](#)).

9.5 Drug-Food Interactions

The rate of pregabalin absorption is decreased when given with food resulting in a decrease in C_{max} by approximately 25% to 30% and an increase in T_{max} to approximately 3 hours. However, administration of pregabalin with food has no clinically relevant effect on the total amount of pregabalin absorbed. Therefore, pregabalin can be taken with or without food.

9.6 Drug-Herb Interactions

Pregabalin has no known drug-herb interactions.

9.7 Drug-Laboratory Test Interactions

Pregabalin has no known drug-laboratory test interactions.

10 CLINICAL PHARMACOLOGY

10.1 Mechanism of Action

Pregabalin binds to the α_2 - δ subunit of the voltage-gated calcium channels in central nervous system tissues. *In vitro*, pregabalin reduces calcium influx at nerve terminals, which may inhibit the release of excitatory neurotransmitters such as glutamate. Through this mechanism, pregabalin may modulate nerve impulses involved in the transmission of pain. However, the clinical relevance of these findings in man is unknown.

10.2 Pharmacodynamics

Pregabalin binds with high affinity to the α_2 -delta protein (a calcium channel subunit) of brain tissues and has analgesic, antiepileptic, and anxiolytic activity. Pregabalin is known chemically as (S)-3-(aminomethyl)-5-methylhexanoic acid.

Although the mechanism of action of pregabalin is unknown, results with genetically modified mice and with compounds structurally-related to pregabalin indicate that selective binding to the α_2 -delta protein is required for analgesic, antiepileptic and anxiolytic action in animal models. *In vitro*, pregabalin reduces the release of several neurotransmitters, suggesting a modulatory action on calcium channel function.

Pregabalin does not mimic GABA at GABA_A or GABA_B receptors, nor does it augment GABA_A responses like benzodiazepines or barbiturates. In contrast to vascular calcium channel blockers, pregabalin does not alter systemic blood pressure or cardiac function. Various *in vitro* and *in vivo* results differentiate pregabalin from GABA uptake inhibitors or GABA transaminase inhibitors. In addition, pregabalin does not block sodium channels, it is not active at opiate receptors, it does not alter

cyclooxygenase enzyme activity, it is not a serotonin agonist, it is not a dopamine antagonist, and it is not an inhibitor of dopamine, serotonin or noradrenaline reuptake.

Pregabalin treatment reduces pain-related behavior in neuropathic animal models of diabetes, peripheral nerve damage or chemotherapeutic insult and in a model of musculoskeletal-associated pain. Pregabalin given intrathecally prevents pain-related behaviors and reduces pain-related behavior caused by spinally administered agents, suggesting that it acts directly on tissues of the spinal cord or brain.

10.3 Pharmacokinetics

All pharmacological actions following pregabalin administration are due to the activity of the parent compound; pregabalin is not appreciably metabolized in humans. Mean steady-state plasma pregabalin concentration-time profiles following 75, 300, and 600 mg/day given in equally divided doses every 8 hours (TID) and 600 mg/day given in equally divided doses every 12 hours (BID) are shown in Table 12. Pregabalin pharmacokinetics are linear over the recommended daily dose range. Inter-subject pharmacokinetic variability for pregabalin is low (< 20%).

Table 12. Pregabalin Mean (CV%^a) Steady-State Pharmacokinetic Parameter Values in Healthy Volunteers

Dose (mg)	Regimen	Daily Dose (mg/day)	n	C _{maxss} (mcg/mL)	t _{max} (hr)	C _{minss} (mcg/mL)	AUC _(0-t) (mcg•hr/mL)	t _½ (hr)	CL/F (mL/min)
25	TID ^b	75	8	1.39	0.9	0.45	6.7	5.9	64.1
				-19.5	-34.2	-25	-18.3	-17.3	-16.1
100	TID	300	6	5.03	0.8	1.94	25.2	6.3	68.9
				-21.3	-31	-33.6	-23	-19.6	-20.9
200	TID	600	11	8.52	0.9	3.28	41.7	6.3	81
				-14.8	-22.2	-29.2	-12.8	-13.6	-11.7
300	BID ^c	600	8	9.07	1.4	2.6	59	6.7	85.1
				-10.5	-57.1	-15.5	-6.4	-16.2	-6.4

C_{maxss}: Steady-state peak plasma concentration.

t_{max}: Time of peak plasma concentration at steady state.

C_{minss}: Steady-state trough plasma concentration

AUC_(0-t): Area under the plasma concentration-time curve during one dosing interval at steady state

t_½: Elimination half-life

CL/F: Oral clearance

^a: Percent coefficient of variation


^b: Total daily dose given in equally divided doses every 8 hours

^c: Total daily dose given in equally divided doses every 12 hours

Absorption

Pregabalin is rapidly absorbed when administered in the fasted state, with peak plasma concentrations occurring within 1.5 hours following both single- and multiple-dose administration. Pregabalin oral bioavailability is $\geq 90\%$ and is independent of dose. C_{max} (Figure 1) and AUC values increase proportionally following single- and multiple-dose administration. Following repeated administration, steady state is achieved within 24 to 48 hours. Multiple dose pharmacokinetics are predictable from single-dose data.

Figure 1. Individual and Mean Steady-State Pregabalin C_{max} Values Following 75, 300 and 600 mg/day Given in Equally Divided Doses TID (q8h) to Healthy Volunteers^a



a: Solid line is the regression line going through the origin; individual (○) and mean (◆) values.

Distribution

In preclinical studies, pregabalin has been shown to readily cross the blood brain barrier in mice, rats, and monkeys. Pregabalin is a substrate for system L transporter which is responsible for the transport of large amino acids across the blood-brain barrier. Pregabalin has been shown to cross the placenta in rats and is present in the milk of lactating rats. In humans, the apparent volume of distribution of pregabalin following oral administration is approximately 0.5 L/kg. Pregabalin is not bound to plasma proteins. At clinically efficacious doses of 150 and 600 mg/day, the average steady-state plasma pregabalin concentrations were approximately 1.5 and 6.0 mcg/mL, respectively.

Metabolism

Pregabalin undergoes negligible metabolism in humans. Following a dose of radiolabeled pregabalin, approximately 98% of the radioactivity recovered in the urine was unchanged pregabalin. The N-methylated derivative of pregabalin, the major metabolite of pregabalin found in urine, accounted for 0.9% of the dose. In preclinical studies, pregabalin (S- enantiomer) did not undergo racemization to the R-enantiomer in mice, rats, rabbits, or monkeys.

Elimination

Pregabalin is eliminated from the systemic circulation primarily by renal excretion as unchanged drug. Pregabalin mean $t_{1/2}$ is 6.3 hours. Pregabalin elimination is proportional to creatinine clearance. Pregabalin clearance is reduced in patients with impaired renal function (see [4 DOSAGE AND ADMINISTRATION](#)).

Special Populations and Conditions

Pregabalin undergoes negligible metabolism, is not bound to plasma proteins, and is eliminated predominately as unchanged drug by renal excretion. Clinically important differences in pregabalin pharmacokinetics due to race and gender have not been observed and are not anticipated.

- **Pediatrics** Pharmacokinetics of pregabalin have not been studied in pediatric patients.
- **Geriatrics** Pregabalin oral clearance tended to decrease with increasing age. This decrease in pregabalin oral clearance is consistent with age-related decreases in creatinine clearance. Reduction of pregabalin dose may be required in patients who have age-related compromised renal function (see [7 WARNINGS AND PRECAUTIONS](#) and [4 DOSAGE AND ADMINISTRATION](#)).
- **Gender** A population pharmacokinetic analysis of the Phase 2/3 clinical program showed that the relationship between daily dose and pregabalin drug exposure is similar between genders when adjusted for gender-related differences in creatinine clearance.
- **Race** A population pharmacokinetic analysis of the Phase 2/3 clinical program showed that the relationship between daily dose and pregabalin drug exposure is similar among Caucasians, Blacks, and Hispanics.
- **Renal Insufficiency** Because renal elimination is the major elimination pathway, dosage reduction in patients with renal dysfunction is necessary. Pregabalin is effectively removed from plasma by hemodialysis. Following a 4-hour hemodialysis treatment, plasma pregabalin concentrations are reduced by approximately 50%. For patients on hemodialysis, dosing must be modified (see [4DOSAGE AND ADMINISTRATION](#)).

11 STORAGE, STABILITY AND DISPOSAL

Store at 15°C – 30°C. Keep out of the reach and sight of children.

12 SPECIAL HANDLING INSTRUCTIONS

There are no special handling requirements for this drug product.

PART II: SCIENTIFIC INFORMATION

13 PHARMACEUTICAL INFORMATION


Drug Substance

Proper name: pregabalin

Chemical name: (S)-3-(aminomethyl)-5-methylhexanoic acid

Molecular formula and molecular mass: C₈H₁₇NO₂, 159.23 g/mol

Structural formula:



Physicochemical properties: Pregabalin is a white crystalline solid. It is soluble in water and in both basic and acidic aqueous solutions.

14 CLINICAL TRIALS

14.1 Clinical Trials by Indication

Diabetic Peripheral Neuropathy

Table 13. Summary of Patient Demographics for Clinical Trials in Diabetic Peripheral Neuropathy

Study #	Study Design	Dosage, Route of Administration and Duration	Study Subjects (n)	Mean Age (Range)	Sex
DPN1	Double-blind, fixed-dose, placebo-controlled, multicenter	150 or 600 mg/day (TID regimen), oral, 6 weeks	246	57 (46 – 67)	149M 97F
DPN2	Double-blind, fixed-dose, placebo-controlled, multicenter	75, 300, or 600 mg/day (TID regimen), oral, 5 weeks	337	59 (26 – 85)	202M 135F

Study #	Study Design	Dosage, Route of Administration and Duration	Study Subjects (n)	Mean Age (Range)	Sex
DPN3	Double-blind, fixed-dose, placebo-controlled, multicenter	300 mg/day (100 mg TID), oral, 8 weeks	146	59 (48 – 71)	82M 64F
DPN4	Double-blind, fixed-dose, placebo-controlled, multicenter	150, 300, or 300/600 mg/day For the 300/600 mg/day IPG-PREGABALIN group, to achieve equivalent exposures, patients received either 300 or 600 mg/day depending on the creatinine clearance rate (BID regimen)	395	58 (47 – 70)	219M 176F

Studies DPN1, DPN2, DPN3 and DPN4 enrolled a total of 1124 patients with type 1 or 2 diabetes mellitus who had painful distal symmetrical sensorimotor polyneuropathy for 1 to 5 years. Across all 4 studies, the baseline mean pain scores ranged from 6.3 to 6.7. The analysis population for all primary and secondary analyses for each study was the intent-to-treat population. Among the pregabalin-treated patients enrolled in these studies, the completion rate was 86%.

Table 14. Results of Study DPN1 in Diabetic Peripheral Neuropathy

Primary Endpoints	Associated Value and Statistical Significance for Drug at Specific Dosages	Associated Value and Statistical Significance for Placebo
Reduction in endpoint mean pain scores (mean of the last 7 daily pain scores while on study medication)	150 mg/day: not significantly better than placebo 600 mg/day: significantly better than placebo (p = 0.0002)	
Proportion of responders	600 mg/day: 39%	15% (p = 0.002)

The 600 mg/day arm was associated with higher reporting of adverse events and withdrawals due to adverse events (see [8 ADVERSE REACTIONS](#)).

Table 15. Results of Study DPN2 in Diabetic Peripheral Neuropathy

Primary Endpoints	Associated Value and Statistical Significance for Drug at Specific Dosages	Associated Value and Statistical Significance for Placebo
Reduction in endpoint mean pain scores (mean of the last 7 daily pain scores while on study medication)	300 mg/day: significantly better than placebo (p = 0.0001) 600 mg/day: significantly better than placebo (p = 0.0001)	
Proportion of responders	300 mg/day: 46% 600 mg/day: 48%	18% (p = 0.001 for each dose)

Compared to 300 mg/day treatment arm, patients treated with 600 mg/day experienced more adverse reactions and discontinued the study more frequently (see [Tables 2](#) and [3](#), [8 ADVERSE REACTIONS](#)).

Table 16. Results of Study DPN3 in Diabetic Peripheral Neuropathy

Primary Endpoints	Associated Value and Statistical Significance for Drug at Specific Dosages	Associated Value and Statistical Significance for Placebo
Reduction in endpoint mean pain scores (mean of the last 7 daily pain scores while on study medication)	300 mg/day: significantly better than placebo (p = 0.0001)	
Proportion of responders	300 mg/day: 40%	15% (p = 0.001)

Table 17. Results of Study DPN4 in Diabetic Peripheral Neuropathy

Primary Endpoints	Associated Value and Statistical Significance for Drug at Specific Dosages	Associated Value and Statistical Significance for Placebo
Reduction in endpoint mean pain scores (mean of the last 7 daily pain scores while on study medication)	150 or 300 mg/day: not significantly better than placebo 300/600 mg/day: significantly better than placebo (p = 0.0054)	
Proportion of responders	300/600 mg/day: 46%	30% (p = 0.036)

Patients recorded their pain on a daily diary using an 11-point numerical pain rating scale ranging from 0 = "no pain" to 10 = "worst possible pain." To enter the study, patients had to have moderate to severe pain, i.e., mean baseline score (mean of the last 7 daily diary pain score prior to study medication) of ≥ 4 . The primary measure of efficacy was reduction in endpoint mean pain scores (mean of the last 7 daily pain scores while on study medication). Supplemental analyses included mean pain scores computed for each week during the study, and the proportion of responders (those patients reporting at least 50% reduction in endpoint mean pain score compared to baseline). Secondary outcome variables included daily sleep interference scores, and the Patient Global Impressions of Change (PGIC). Mean sleep scores were computed at endpoint and for each week during the study. The PGIC was completed at the end of the study and measured the patient's overall status compared to baseline using a 7- point categorical scale ranging from "very much improved" to "very much worse."

Post-herpetic Neuralgia

Table 18. Summary of Patient Demographics for Clinical Trials in Post-herpetic Neuralgia

Study #	Study Design	Dosage, Route of Administration and Duration	Study Subjects (n)	Mean Age (Range)	Sex
PHN1	Double-blind, fixed-dose, placebo-controlled, multicenter	150 or 300 mg/day (TID regimen), oral, 8 weeks	238	72 (32 – 96)	107M 131F
PHN2	Double-blind, fixed-dose, placebo-controlled, multicenter	300 or 600 mg/day depending on patient creatinine clearance rate, oral, 8 weeks	173	71 (60 – 82)	81M 92F

Study #	Study Design	Dosage, Route of Administration and Duration	Study Subjects (n)	Mean Age (Range)	Sex
PHN3	Double-blind, fixed-dose, placebo-controlled, multicenter	150, 300, or 300/600 mg/day For the 300/600 mg/day IPG-PREGABALIN group, to achieve equivalent exposures, patients received either 300 or 600 mg/day depending on the creatinine clearance rate (BID regimen)	368	71 (18 – 92)	168M 200F

Studies PHN1, PHN2 and PHN3 enrolled 779 patients with post-herpetic neuralgia who continued to have pain for at least 6 months (Study PHN1) or 3 months (Study PHN2 and PHN3) after healing of the herpes zoster skin rash. Across all 3 studies, the baseline mean pain scores ranged from 6.4 to 6.8. The analysis population for all primary and secondary analyses for each study was the intent-to-treat population. Among the IPG-PREGABALIN-treated patients enrolled in these studies, the completion rate was 71%.

Patients were randomly assigned to one of the treatment arms depending on their creatinine clearance rate. Compared to patients with creatinine clearance rate ≥ 60 mL/min, those with decreased creatinine clearance rate (≤ 60 mL/min) experienced markedly higher rates of adverse events and discontinued the trial more frequently.

Table 19. Results of Study PHN1 in Postherpetic Neuralgia

Primary Endpoints	Associated Value and Statistical Significance for Drug at Specific Dosages	Associated Value and Statistical Significance for Placebo
Reduction in endpoint mean pain scores (mean of the last 7 daily pain scores while on study medication)	150 mg/day: significantly better than placebo (p = 0.0002) 300 mg/day: significantly better than placebo (p = 0.0002)	
Proportion of responders	150 mg/day: 26% 300 mg/day: 28%	10% (p = 0.006 for each dose)

This trial included patients with decreased creatinine clearance rate (30-60 mL/min) who were randomly assigned to one of the treatment arms.

Table 20. Results of Study PHN2 in Post-herpetic Neuralgia

Primary Endpoints	Associated Value and Statistical Significance for Drug at Specific Dosages	Associated Value and Statistical Significance for Placebo
Reduction in endpoint mean pain scores (mean of the last 7 daily pain scores while on study medication)	300/600 mg/day: significantly better than placebo (p = 0.0001)	
Proportion of responders	300/600 mg/day: 50%	20% (p ≤ 0.001)

Table 21. Results of Study PHN3 in Post-herpetic Neuralgia

Primary Endpoints	Associated Value and Statistical Significance for Drug at Specific Dosages	Associated Value and Statistical Significance for Placebo
Reduction in endpoint mean pain scores (mean of the last 7 daily pain scores while on study medication)	150 mg/day: significantly better than placebo (p = 0.0077) 300 mg/day: significantly better than placebo (p = 0.0016) 300/600 mg/day: significantly better than placebo (p = 0.0003)	
Proportion of responders	150 mg/day: 26% 300 mg/day: 27% 300/600 mg/day: 38%	8% (p = 0.001 for each dose)

Patients recorded their pain on a daily diary using an 11-point numerical pain rating scale ranging from 0 = "no pain" to 10 = "worst possible pain." To enter the study, patients had to have moderate to severe pain, i.e., mean baseline score (mean of the last 7 daily diary pain score prior to study medication) of ≥ 4 . The primary measure of efficacy was reduction in endpoint mean pain scores (mean of the last 7 daily pain scores while on study medication). Supplemental analyses included mean pain scores computed for each week during the study, and the proportion of responders (those patients reporting at least 50% reduction in endpoint mean pain score compared to baseline). Secondary outcome variables included daily sleep interference scores, and the Patient Global Impressions of Change (PGIC). Mean sleep scores were computed at endpoint and for each week during the study. The PGIC was completed at the end of the study and measured the patient's overall status compared to baseline using a 7-point categorical scale ranging from "very much improved" to "very much worse."

Overall Analysis of Diabetic Peripheral Neuropathy and Post-herpetic Neuralgia Studies

When endpoint mean pain scores are combined across all controlled diabetic neuropathic pain and post-herpetic neuralgia studies, no significant differences in efficacy based on gender, or race, were observed.

In controlled diabetic neuropathy and post-herpetic neuralgia clinical trials, pregabalin, at doses of 150, 300, and 600 mg/day, was statistically significantly more effective than placebo in reducing sleep disturbance and improving PGIC.

Fibromyalgia

Table 22. Summary of Patient Demographics for Clinical Trials in Fibromyalgia

Study #	Study Design	Dosage, Route of Administration and Duration	Study Subjects (n)	Mean Age (Range)	Sex
F1	Multicentre, parallel, double-blind, randomized, placebo-controlled	150, 300 or 450 mg/day (TID regimen), oral, 8 weeks	529	48.6 (20 to 78)	45M 484F
F2	Multicentre, parallel, double-blind, randomized, placebo-controlled	300, 450 or 600 mg/day (BID regimen), oral, 13 weeks	748	48.8 (18 to 82)	42M 706F
F3	Multicentre, parallel, double-blind, randomized, placebo-controlled, using enriched population, only	300, 450 or 600 mg/day (BID regimen), oral, 14 weeks	745	50.1 (18 to 81)	41M 704F
F4	Multicentre, parallel, double-blind, randomized, placebo-controlled, using enriched population, only	300, 450 or 600 mg/day (BID regimen), oral, 14 weeks	735	48.5 (20 to 81)	63M 672F

To enter the study, patients had to have moderate to severe pain, i.e., mean baseline score (mean of the last 7 daily diary pain scores prior to study medication) of ≥ 4 , and a diagnosis of fibromyalgia based on the American College of Rheumatology criteria (history of widespread

pain for 3 months, and pain present at 11 or more out of 18 specific tender points). During all studies described below, patients were allowed to take acetaminophen up to 4 g per day as needed for pain relief.

Table 23. Results of Study F1 in Fibromyalgia

Primary Endpoints	Associated Value and Statistical Significance for Drug at Specific Dosages	Associated Value and Statistical Significance for Placebo
Reduction in endpoint mean pain scores (mean of the last 7 daily pain scores while on study medication)	Significantly larger reduction from baseline, ($p < 0.005$) only at 150 mg TID.	
Proportion of responders	50 mg TID: 78% 100 mg TID: 83% 150 mg TID: 75%	74%

A dose-related increase in adverse events and withdrawals due to adverse events was observed (see [8 ADVERSE REACTIONS](#)).

Table 24. Results of Study F2 in Fibromyalgia

Primary Endpoints	Associated Value and Statistical Significance for Drug at Specific Dosages	Associated Value and Statistical Significance for Placebo
Reduction in endpoint mean pain scores (mean of the last 7 daily pain scores while on study medication)	Significantly larger reduction from baseline, ($p < 0.05$) at all doses studied.	
Proportion of responders	150 mg TID: 66% 225 mg TID: 66% 300 mg TID: 58%	68%

A dose-related increase in adverse events and withdrawals due to adverse events was observed (see [8 ADVERSE REACTIONS](#)).

Table 25. Results of Study F3 in Fibromyalgia

Primary Endpoints	Associated Value and Statistical Significance for Drug at Specific Dosages	Associated Value and Statistical Significance for Placebo
Reduction in endpoint mean pain scores (mean of the last 7 daily pain scores while on study medication)	Significantly larger reduction from baseline, ($p < 0.0005$) at all doses studied.	
Proportion of responders	150 mg BID: 67.2% 225 mg BID: 65.8% 300 mg BID: 60.1%	67.9%

A dose-related increase in adverse events and withdrawals due to adverse events was observed (see [8 ADVERSE REACTIONS](#)).

Table 26. Results of Study F4 in Fibromyalgia

Primary Endpoints	Associated Value and Statistical Significance for Drug at Specific Dosages	Associated Value and Statistical Significance for Placebo
Reduction in endpoint mean pain scores (mean of the last 7 daily pain scores while on study medication)	Significantly larger reduction from baseline, ($p < 0.05$) only at 225 mg BID.	
Proportion of responders	150 mg BID: 67% 225 mg BID: 73% 300 mg BID: 65%	76%

A dose-related increase in adverse events and withdrawals due to adverse events was observed (see [8 ADVERSE REACTIONS](#)).

Patients recorded their pain on a daily diary using an 11-point numerical pain rating (Likert) scale ranging from 0=“no pain” to 10=“worst possible pain”. Treatment differences, defined as the change in endpoint mean pain scores for pregabalin versus placebo (drug – placebo), were calculated.

The primary efficacy endpoint in the 26-week long-term relapse observation study was the time to loss of therapeutic response as determined from the Pain Visual Analog Scale (VAS) or from worsening of fibromyalgia symptoms necessitating an alternative treatment per the clinical judgment of the principal investigator. Secondary endpoints included the time to worsening of PGIC, FIQ, and MOS-SS scores.

A 26-week long-term relapse observation study compared pregabalin with placebo. Patients who met the American College of Rheumatology criteria for fibromyalgia (history of widespread pain for 3 months, and pain present at 11 or more out of 18 specific tender points) and with a score of ≥ 40 mm on the pain Visual Analog Scale (VAS) were eligible to enter a 6-week, open-label, dose-optimization phase. During this phase, patients were titrated up to a total daily dose of 300 mg, 450 mg, or 600 mg. Patients were considered to be pregabalin responders if they had both at least a 50% reduction in pain on the VAS and rated their overall improvement on the PGIC as “much improved” or “very much improved.” A total of 566 pregabalin responders were then randomized in the double-blind phase to either their optimized pregabalin dose (279 patients) achieved in the open-label phase (therapy continued) or to placebo (287 patients; therapy discontinued). Patients were treated for up to 26 weeks in the double-blind phase. Treatment with pregabalin resulted in a significantly longer time to loss of therapeutic response than treatment with placebo ($p < 0.0001$), with fewer relapses on pregabalin (32%) compared to placebo (61%). Of the pregabalin responders entering the double-blind phase assigned to remain on pregabalin, 38% completed 26 weeks of treatment, versus 19% of those assigned to receive placebo. All supplemental and secondary endpoints were supportive of the primary efficacy measure.

14.3 Comparative Bioavailability Studies

A double blinded, randomized, two-treatment, two period, single-dose, cross-over bioequivalence study comparing IPG-Pregabalin 25mg capsules, (Marcan Pharmaceuticals Inc.) to PrLYRICA® 25mg capsules (Pfizer Canada Inc.) was conducted in 31 healthy adult, human male subjects under fasting conditions. The results are presented in the table below.

SUMMARY TABLE OF THE COMPARATIVE BIOAVAILABILITY DATA

Pregabalin (1x 25 mg) From measured data Geometric Mean Arithmetic Mean (CV %)				
Parameter	Test*	Reference†	% Ratio of Geometric Means	Confidence Interval
AUC _{0t} (ng*hr/mL)	6200.84 6262.05 (14.10)	6095.23 6144.78 (12.83)	101.83	99.82-103.88
AUC _{0-∞} (ng*hr/mL)	6551.67 6612.57 (13.73)	6457.78 6505.830 (12.35)	101.54	99.65-103.46
C _{max} (ng/mL)	959.32 984.84 (23.76)	896.93 913.13 (19.25)	106.80	99.37-114.79
T _{max} [§] (h)	0.83 (0.50-1.83)	0.83 (0.50-2.25)		
t _{1/2} [€] (h)	6.33 (11.43)	6.42 (13.03)		

* IPG-Pregabalin 25 mg capsules (Marcan Pharmaceuticals Inc.)

† PrLYRICA® 25 mg capsules (Pfizer Canada Inc.,) purchased in Canada.

§ Expressed as the median (range).

€ Expressed as the arithmetic mean (CV %) only.

A double blinded, randomized, two-treatment, two period, single-dose, cross-over bioequivalence study comparing IPG-Pregabalin 300mg capsules, (Marcan Pharmaceuticals Inc.) to PrLYRICA® 300mg capsules (Pfizer Canada Inc.) was conducted in 29 healthy adult, human male subjects under fasting conditions. The results are presented in the table below.

SUMMARY TABLE OF THE COMPARATIVE BIOAVAILABILITY DATA

Pregabalin (1x 300 mg) From measured data Geometric Mean Arithmetic Mean (CV %)				
Parameter	Test*	Reference [†]	% Ratio of Geometric Means	Confidence Interval
AUC _{0t} (ng*hr/mL)	63901.95 64300.34 (11.27)	63738.14 64062.84 (10.07)	100.28	98.84-101.74
AUC _{0-∞} (ng*hr/mL)	66262.14 66747.69 (12.32)	65963.88 66332.38 (10.56)	100.48	98.94-102.05
C _{max} (ng/mL)	7724.77 7806.27 (14.44)	7907.71 8003.39 (15.69)	97.59	93.51-101.84
T _{max} [§] (h)	1.50 (0.67-3.00)	1.50 (0.50-2.25)		
t _{1/2} [€] (h)	6.09 (12.36)	5.96 (11.84)		

* IPG-Pregabalin 300 mg capsules (Marcan Pharmaceuticals Inc.).

† PrLyrica® 300 mg capsules (Pfizer Canada Inc.,) were purchased in Canada.

§ Expressed as the median (range)

€ Expressed as the arithmetic mean (CV %) only

15 MICROBIOLOGY

No microbiological information is required for this drug product.

16 NON-CLINICAL TOXICOLOGY

Acute and Repeated-Dose Toxicity

The acute toxicity of pregabalin is low. Animals given oral doses of pregabalin in acute and repeated-dose studies were hypoactive, hyperactive, and/or ataxic, signs commonly observed with CNS-active drugs. No significant target organ toxicity was observed in rats treated for up to 52 weeks or in monkeys treated for up to 69 weeks at exposures at least 8 times the mean human exposure at the maximum recommended dose of 600 mg/day. Effects of pregabalin in juvenile animals were similar to those in adults.

Acute Toxicity

Acute oral and IV studies were conducted in mice and rats. Hypoactivity in mice and rats, and diarrhea and urine staining in rats, but no deaths or changes in clinical biochemical parameters, occurred following administration of single oral doses of 5000 mg/kg or single IV doses of 300 mg/kg, the highest IV dose achievable based on solubility and dose volume limitations. No drug-related gross or histopathologic changes were observed in rodents given single oral or IV doses.

Repeated-Dose Toxicity

Repeated-dose studies in definitive toxicology species are listed in Table 16 and results are discussed in the sections below.

Table 27. Repeated-Dose Oral Studies with Pregabalin

Species	Duration (Weeks)	Dose Range (mg/kg)
Rat	2	500 – 2500
	4	500 – 5000
	4	50 – 250
	13	50 – 1250
	26/52	50 – 500
Monkey	Escalating-Dose	50 – 2000
	2	100 – 2000
	4	100 – 2000

Species	Duration (Weeks)	Dose Range (mg/kg)
	4-day (toxicokinetic)	500 – 1000
	4	25 - 500 BID
	13	10 – 500
	65-69	10 -500

Rat: Ataxia and hypoactivity were seen in rats given repeated oral doses ≥ 500 mg/kg with associated $AUC_{(0-24)} \geq 1280$ mcg•hr/mL; tail dermatopathy and urine staining were seen at ≥ 250 mg/kg with associated $AUC_{(0-24)} \geq 802$ mcg•hr/mL. Body weight gain decreased 18% to 70% in rats given pregabalin at 500 to 5000 mg/kg for 4, 13, or 52 weeks. No effects on body weight gain occurred at 50 or 250 mg/kg for 52 weeks with associated exposures ≤ 1210 mcg•hr/mL. Urinary bladder changes (dilatation, edema, and hemorrhage) and sporadic mortality associated with pyelonephritis and/or cystitis occurred in rats at ≥ 250 mg/kg; associated $AUC_{(0-24)}$ was ≥ 802 mcg•hr/mL. Red blood cell (RBC) parameters increased 6% to 16% in rats given pregabalin at 500 to 5000 mg/kg for 4 weeks, and were reversible. RBC count only increased 5% to 11% at 50 to 1250 mg/kg in Weeks 13, 26, and 52 with associated exposures ≥ 228 mcg•hr/mL. Platelet count decreased 14% to 36% in male rats given ≥ 50 mg/kg and in female rats given ≥ 100 mg/kg for up to 52 weeks with associated $AUC_{(0-24)} \geq 228$ mcg•hr/mL. Changes in platelet count were reversed during a 4-week withdrawal period after 4 weeks of dosing. Total nucleated cells in the bone marrow decreased 18% to 44% in rats given pregabalin at 250 to 1250 mg/kg for 13, 26, or 52 weeks. No drug-related effects on bone marrow occurred at 50 mg/kg for up to 52 weeks with associated exposures ≤ 228 mcg•hr/mL. In rats given pregabalin for 4 weeks, epididymal hypospermia occurred at ≥ 500 mg/kg and spermatogenic epithelial degeneration occurred at 1250 mg/kg with associated $AUC_{(0-24)} \geq 1690$ mcg•hr/mL. Similar changes were not seen in the 52-week study at doses up to 500 mg/kg. No adverse effects were observed in rats given 50 mg/kg for 13 weeks, with associated combined-sex $AUC_{(0-24)}$ of 188 mcg•hr/mL; effects at this dose in the 52-week study were minimal.

Monkeys: Nasal discharge and soft feces/diarrhea occurred in monkeys given repeated oral doses ≥ 100 mg/kg for up to 13 weeks, with associated $AUC_{(0-24)} \geq 398$ mcg•hr/mL. Hypoactivity occurred at ≥ 500 mg/kg; $AUC_{(0-24)}$ was at least 974 mcg•hr/mL. Deaths occurred within 3 days after treatment initiation in monkeys given 500 mg/kg BID or 1000 or 2000 mg/kg. Although toxicokinetic data were not available for all these monkeys, the lowest $AUC_{(0-24)}$ in an animal that died was 1640 mcg•hr/mL. Myocardial effects seen in monkeys at ≥ 50 mg/kg in 4-week studies were not seen in sub-chronic and chronic studies and, therefore, not considered drug-related. There were no effects on body weight gain, hematology, or bone marrow parameters in monkeys given pregabalin at 10 to 500 mg/kg for up to 69 weeks. Sperm count, motility, and morphology were not affected in monkeys given up to 500 mg/kg for 69 weeks. Tail dermatopathy was observed in monkeys at ≥ 25 mg/kg. With the exception of tail dermatopathy, no significant effects were observed in monkeys given up to 500 mg/kg up to 69 weeks with associated plasma concentrations up to 76.2 mcg/mL. Based on the 4-week toxicokinetic profiles and single time point samples in Week 52, combined-sex $AUC_{(0-24)}$ was estimated to be 1040 mcg•hr/mL.

The etiology of these skin lesions seen in both rats and monkeys in repeated-dose toxicology studies is unknown. At the maximum recommended human dose (MRD) of 600 mg/day, there is a 2-fold safety margin for the dermatological lesions. The more severe dermatopathies involving necrosis were associated with pregabalin exposures (as expressed by plasma AUCs) of approximately 3 to 8 times those achieved in humans given the MRD. No increase in incidence of skin lesions was observed in clinical studies.

Genetic Toxicity

The genotoxic potential of pregabalin was evaluated in an extensive battery of tests. Pregabalin was not mutagenic in bacteria up to 5000 mcg/plate with or without metabolic activation provided by post mitochondrial supernatant fractions from livers of B6C3F1 or CD-1 mice or Wistar rats treated with Arochlor 1254. In mammalian cells *in vitro*, no evidence of mutagenicity or structural chromosome aberrations was observed up to 1600 mcg/mL with or without metabolic activation. Pregabalin did not induce unscheduled DNA synthesis in hepatocytes of B6C3F1 or CD-1 mice or Wistar rats given single oral doses up to 2000 mg/kg. Micronucleus frequency was not increased in bone marrow from B6C3F1 or CD-1 mice or Wistar rats given single oral doses of pregabalin up to 2000 mg/kg.

These results demonstrate pregabalin is not genotoxic *in vitro* or *in vivo*.

Mutagenesis

Pregabalin is not genotoxic based on results of a battery of *in vitro* and *in vivo* tests. Pregabalin was not mutagenic in bacteria or in mammalian cells *in vitro*, was not clastogenic in mammalian systems *in vitro* and *in vivo*, and did not induce unscheduled DNA synthesis in mouse or rat hepatocytes.

Carcinogenesis

Dose-dependent increase in the incidence of malignant vascular tumors (hemangiosarcomas) was observed in two strains of mice (B6C3F1 and CD-1) given pregabalin (200, 1000, or 5000 mg/kg) in the diet for two years. Plasma pregabalin exposure (AUC) in mice receiving the lowest dose that increased hemangiosarcomas was approximately equal to the human exposure at the maximum recommended dose (MRD) of 600 mg/day. A no-effect dose for induction of hemangiosarcomas in mice was not established. In an investigative study in female B6C3F1 mice, chronic treatment (24 months) with pregabalin at 1000 mg/kg caused an increased incidence of hemangiosarcoma, consistent with previous studies, but not at 50 or 200 mg/kg. Discontinuation of treatment after 12 months at 1000 mg/kg did not significantly reduce the incidence of hemangiosarcoma at 24 months. Evidence of carcinogenicity was not seen in two studies in Wistar rats following dietary administration of pregabalin for two years at doses (50, 150, or 450 mg/kg in males and 100, 300, or 900 mg/kg in females) that were associated with plasma exposures in males and females up to approximately 14 and 24 times, respectively, human exposure at the MRD. The clinical significance in humans of this finding in mice is unknown.

Ocular lesions (characterized by retinal atrophy [including loss of photoreceptor cells] and/or corneal inflammation/mineralization) were observed in two lifetime carcinogenicity studies in Wistar rats. These findings were observed at plasma pregabalin exposures (AUC) ≥ 2 times those achieved in humans given the maximum recommended dose of 600 mg/day. A no-effect

dose for ocular lesions was not established. Similar lesions were not observed in lifetime carcinogenicity studies in two strains of mice or in monkeys treated for 1 year. The clinical significance of this finding in rats is unknown.

Reproductive and Developmental Toxicology

In female rats, although estrus and diestrus stages were prolonged at 1250 and 2500 mg/kg, no effects on fertility were observed in females given 500 to 2500 mg/kg prior to mating with untreated males. In male rats, pregabalin administration resulted in decreased sperm motility and decreased fertility at exposures \geq 27 times the mean human exposure at the maximum recommended clinical dose and were reversible. There were no drug-related effects on sperm parameters in monkeys treated for 69 weeks with exposures up to 8 times the maximum human exposure.

In a double-blind, placebo-controlled clinical trial to assess the effect of pregabalin on sperm motility, 30 healthy male subjects were exposed to pregabalin at a dose of 600 mg/day. After 3 months of treatment (one complete sperm cycle), the difference between placebo- and pregabalin-treated subjects in mean percent sperm with normal motility was $<$ 4% and neither group had a mean change from baseline of more than 2%. Effects on other male reproductive parameters in humans have not been adequately studied (see [7 WARNINGS AND PRECAUTIONS](#)).

Pregabalin induced maternal toxicity in embryo-fetal development studies in rats at \geq 500 mg/kg and rabbits at \geq 250 mg/kg, and fetal toxicity in rats at 2500 mg/kg and in rabbits at 1250 mg/kg. Pregabalin was not teratogenic in mice, rats, or rabbits at exposures 31 to 77 times the mean human exposure at the maximum recommended clinical dose of 600 mg/day. In the prenatal-postnatal toxicity study in rats, pregabalin induced offspring developmental toxicity at exposures \geq 5 times the mean human exposure. No effects occurred at exposures twice the mean human exposure at the maximum recommended clinical dose (see [7 WARNINGS AND PRECAUTIONS](#)).

Pregabalin is excreted in milk of lactating rats. Because of the potential exposure in breast-feeding infants, breast feeding is not recommended (see [7 WARNINGS AND PRECAUTIONS](#)).

17 SUPPORTING PRODUCT MONOGRAPHS

1. ^{Pr}LYRICA[®], Capsules, 25, 50, 75, 150, 225, 300 mg, Submission Control 274588, Product Monograph, BGP Pharma ULC. (May 09, 2023).

PATIENT MEDICATION INFORMATION

READ THIS FOR SAFE AND EFFECTIVE USE OF YOUR MEDICINE

PrIPG-PREGABALIN

Pregabalin Capsules

Read this carefully before you start taking **IPG-PREGABALIN** and each time you get a refill. This leaflet is a summary and will not tell you everything about this drug. Talk to your healthcare professional about your medical condition and treatment and ask if there is any new information about **IPG-PREGABALIN**.

Serious Warnings and Precautions

Taking IPG-PREGABALIN with opioid medicines, benzodiazepines, alcohol, or other central nervous system depressants (including street drugs) can cause severe drowsiness, decreased awareness, breathing problems, coma, and death.

What is IPG-PREGABALIN used for?

IPG-PREGABALIN is used in adults to treat pain caused by nerve damage due to:

- Diabetes
- Shingles

IPG-PREGABALIN is also used to treat pain associated with a condition called fibromyalgia (widespread pain).

How does IPG-PREGABALIN work?

IPG-PREGABALIN works by decreasing the number of nerve signals in your body. This helps to calm down oversensitive nerve cells, which helps to relieve pain.

What are the ingredients in IPG-PREGABALIN?

Medicinal ingredients: Pregabalin

Non-medicinal ingredients: The non-medicinal ingredients that make up the capsule core are pre-gelatinized starch and talc.

The capsule shells contain gelatin and titanium dioxide. In addition, the orange capsule shells contain iron oxide red.

The markings on the capsules are in black ink, which contains ammonia solution, black iron oxide, butyl alcohol, dehydrated alcohol, isopropyl alcohol, potassium hydroxide, propylene glycol, shellac, and water.

IPG-PREGABALIN comes in the following dosage forms:

Capsules: 25 mg, 50 mg, 75 mg, 100 mg, 150 mg, 200 mg, 225 mg and 300 mg.

Do not use IPG-PREGABALIN if:

- You are allergic to pregabalin, or any other ingredient in IPG-PREGABALIN.

To help avoid side effects and ensure proper use, talk to your healthcare professional before you take IPG-PREGABALIN. Talk about any health conditions or problems you may have, including if you:

- have any kidney problems.
- have any breathing problems or respiratory disease.
- are pregnant, plan to become pregnant, or think you might be pregnant.
- are breastfeeding. IPG-PREGABALIN passes into breast milk and it is not known if it can harm your baby. You and your healthcare provider should discuss whether you should take IPG-PREGABALIN or breast-feed, but you should not do both.
- have ever had an allergic reaction to any other drug.
- have a history of heart disease or heart failure.
- have a history of lower gastrointestinal problems (e.g., constipation, blocked or paralysed bowel), or you are taking medication(s) that may cause constipation.
- have a history of addiction or substance abuse.

Other warnings you should know about:

Pregnancy: IPG-PREGABALIN should not be taken during pregnancy. If you become pregnant while taking IPG-PREGABALIN, tell your healthcare professional **right away**. If you are taking IPG-PREGABALIN and are of childbearing age, you must use an effective method of birth control. If you take IPG-PREGABALIN during your first trimester of pregnancy, it can cause major birth defects in your unborn child.

Pregnancy Registry: If you become pregnant while taking IPG-PREGABALIN, talk to your healthcare professional about registering with the North American Antiepileptic Drug Pregnancy Registry. You can enroll in this registry by calling 1-888-233-2334. The purpose of this registry is to collect information about the safety of anticonvulsant medicines during pregnancy. Information on the registry can also be found at the website <http://www.aedpregnancyregistry.org/>.

Eye Problems: If you experience any changes in your vision while taking IPG-PREGABALIN, tell your healthcare professional right away. If you already have eye problems, your healthcare professional may recommend that you have more frequent eye exams.

Stopping Your Medication: Do NOT stop taking IPG-PREGABALIN without discussing it with your healthcare professional first. Stopping your treatment abruptly may cause you to experience unwanted side effects. If you have any concerns with your treatment, talk to your healthcare professional.

Driving and Using Machines: IPG-PREGABALIN may cause you to feel dizzy or sleepy. Wait until you know how IPG-PREGABALIN affects you before performing tasks that require special attention.

Tell your healthcare professional about all the medicines you take, including any drugs, vitamins, minerals, natural supplements or alternative medicines.

The following may interact with IPG-PREGABALIN:

- Alcohol

- Opioid medicines
- Benzodiazepines
- Medications used to treat diabetes, such as rosiglitazone, pioglitazone

How to take IPG-PREGABALIN:

- Take IPG-PREGABALIN exactly as your healthcare professional has told you to.
- Do NOT stop taking IPG-PREGABALIN abruptly or you may experience unwanted side effects. If you have concerns with your treatment, talk to your healthcare professional first.
- You can take IPG-PREGABALIN with or without food.

Usual dose:

Your healthcare professional has decided the best dose for you and will tell you how much to take and when to take it.

Overdose:

If you think you, or a person you are caring for, have taken too much IPG-PREGABALIN, contact a healthcare professional, hospital emergency department, or regional poison control centre immediately, even if there are no symptoms.

Missed Dose:

If you miss a dose by a few hours, take it as soon as you remember. If it is close to your next dose, do NOT take the missed capsules. Simply wait and restart IPG-PREGABALIN with your next scheduled dose.

What are possible side effects from using IPG-PREGABALIN?

These are not all the possible side effects you may have when taking IPG-PREGABALIN. If you experience any side effects not listed here, tell your healthcare professional.

Side effects may include:

- Dizziness
- Sleepiness
- Headache
- Nausea
- Dry mouth
- Increased cough
- Weight gain
- Trouble concentrating
- Forgetfulness
- Lack of energy
- Muscle weakness
- Constipation; talk with your healthcare professional about ways to prevent constipation when you start using IPG-PREGABALIN

Serious side effects and what to do about them			
Symptom / effect	Talk to your healthcare professional		Stop taking drug and get immediate medical help
	Only if severe	In all cases	
COMMON			
Blurred vision		✓	
UNCOMMON			
Edema: unusual swelling of the arms, hands, legs, feet and ankles, face or airway passages		✓	
Angiodema: swelling of the face, mouth, lips, gums, neck or throat, trouble breathing, diarrhea, nausea or vomiting			✓
Allergic Reaction: difficulty swallowing or breathing, wheezing, feeling sick to your stomach and throwing up, hives or rash, swelling of the face, lips, tongue or throat			✓
Decrease in the amount of urine		✓	
Suicidal thoughts or actions		✓	
RARE			
Kidney Failure: confusion, itchiness or rashes, puffiness in your face and hands, swelling in your feet or ankles, decreased amount of urine, weight gain			
VERY RARE			
Seizures (fits): uncontrollable shaking with or without loss of consciousness.			✓
Severe Skin Reactions: fever, severe rash, swollen lymph glands, flu-like feeling, blisters and peeling skin that may start in and around the mouth, nose, eyes and genitals and spread to other areas of the body, yellow skin or eyes, shortness of breath, dry cough, chest pain or discomfort, feeling thirsty, urinating less often, less urine			✓

If you have a troublesome symptom or side effect that is not listed here or becomes bad enough to interfere with your daily activities, tell your healthcare professional.

Reporting Side Effects

You can report any suspected side effects associated with the use of health products to Health Canada by:

- Visiting the Web page on Adverse Reaction Reporting (<https://www.canada.ca/en/health-canada/services/drugs-health-products/medeffect-canada/adverse-reaction-reporting.html>) for information on how to report online, by mail or by fax; or
- Calling toll-free at 1-866-234-2345.

NOTE: Contact your health professional if you need information about how to manage your side effects. The Canada Vigilance Program does not provide medical advice.

Storage:

Store at room temperature (15°C to 30°C) in the package it came in.

Keep out of reach and sight of children.

If you want more information about IPG-PREGABALIN:

- Talk to your healthcare professional
 - Find the full product monograph that is prepared for healthcare professionals and includes this Patient Medication Information by visiting the Health Canada website: (<https://www.canada.ca/en/health-canada/services/drugs-health-products/drug-products/drug-product-database.html>); the manufacturer's website www.marcanpharma.com, or by calling 1-855-627-2261.

This leaflet was prepared by Marcan Pharmaceuticals Inc.

Last Revised: July 13, 2023