

PRODUCT MONOGRAPH  
INCLUDING PATIENT MEDICATION INFORMATION

**PrLINESSA<sup>®</sup> 21 and PrLINESSA<sup>®</sup> 28**

desogestrel and ethinyl estradiol tablets USP

0.100 mg, 0.025 mg

0.125 mg, 0.025 mg

0.150 mg, 0.025 mg

Oral Contraceptive  
G03AB05

Aspen Pharmacare Canada Inc.  
8 – 1155 North Service Road West  
Oakville, ON  
L6M 3E3

Date of Initial Approval:  
June 13, 2006

Date of Preparation:  
September 12, 2023

Trademarks are owned by or licensed to the Aspen Group of companies.  
©2023 Aspen Group of companies or its licensor. All rights reserved.

Submission Control No: 273119

## RECENT MAJOR LABEL CHANGES

|                              |         |
|------------------------------|---------|
| Contraindications (2)        | 09/2023 |
| Warnings and Precautions (7) | 09/2023 |

## TABLE OF CONTENTS

|   |           |
|---|-----------|
| <b>RECENT MAJOR LABEL CHANGES</b> .....                           | <b>2</b>  |
| <b>TABLE OF CONTENTS</b> .....                                    | <b>2</b>  |
| <b>PART I: HEALTH PROFESSIONAL INFORMATION</b> .....              | <b>4</b>  |
| <b>1 INDICATIONS</b> .....  | <b>4</b>  |
| 1.1 Pediatrics.....   | 4         |
| 1.2 Geriatrics.....   | 4         |
| <b>2 CONTRAINDICATIONS</b> .....                                  | <b>4</b>  |
| <b>3 SERIOUS WARNINGS AND PRECAUTIONS BOX</b> .....               | <b>5</b>  |
| <b>4 DOSAGE AND ADMINISTRATION</b> .....                          | <b>6</b>  |
| 4.1 Dosing Considerations .....                                   | 6         |
| 4.2 Recommended Dose and Dosage Adjustment .....                  | 6         |
| 4.3 Administration .....  | 6         |
| 4.4 Missed Dose .....   | 8         |
| <b>5 OVERDOSAGE</b> .....   | <b>9</b>  |
| <b>6 DOSAGE FORMS, STRENGTHS, COMPOSITION AND PACKAGING</b> ..... | <b>10</b> |
| <b>7 WARNINGS AND PRECAUTIONS</b> .....                           | <b>11</b> |
| 7.1 Special Populations .....                                     | 21        |
| 7.1.1 Pregnant Women .....  | 21        |
| 7.1.2 Breast-feeding.....   | 21        |
| 7.1.3 Pediatrics.....   | 21        |
| 7.1.4 Geriatrics .....  | 21        |
| <b>8 ADVERSE REACTIONS</b> .....                                  | <b>21</b> |
| 8.1 Adverse Reaction Overview .....                               | 21        |
| 8.2 Clinical Trial Adverse Reactions .....                        | 23        |
| 8.3 Less Common Clinical Trial Adverse Reactions.....             | 26        |
| 8.4 Post-Market Adverse Reactions.....                            | 27        |
| <b>9 DRUG INTERACTIONS</b> .....                                  | <b>28</b> |
| 9.1 Drug Interactions Overview .....                              | 28        |
| 9.2 Drug-Drug Interactions .....                                  | 28        |
| 9.3 Drug-Food Interactions.....                                   | 34        |
| 9.4 Drug-Herb Interactions .....                                  | 34        |
| 9.5 Drug-Laboratory Test Interactions.....                        | 34        |

|           |   |           |
|-----------|---|-----------|
| 9.6       | Drug-Lifestyle Interactions.....            | 35        |
| <b>10</b> | <b>CLINICAL PHARMACOLOGY.....</b>           | <b>36</b> |
| 10.1      | Mechanism of Action .....                   | 36        |
| 10.2      | Pharmacodynamics .....                      | 36        |
| 10.3      | Pharmacokinetics .....                      | 37        |
| <b>11</b> | <b>STORAGE, STABILITY AND DISPOSAL.....</b> | <b>39</b> |
| <b>12</b> | <b>SPECIAL HANDLING INSTRUCTIONS.....</b>   | <b>40</b> |
| <b>13</b> | <b>PHARMACEUTICAL INFORMATION .....</b>     | <b>41</b> |
| <b>14</b> | <b>CLINICAL TRIALS .....</b>                | <b>42</b> |
| <b>15</b> | <b>MICROBIOLOGY .....</b>                   | <b>47</b> |
| <b>16</b> | <b>NON-CLINICAL TOXICOLOGY .....</b>        | <b>47</b> |
|           | <b>PATIENT MEDICATION INFORMATION .....</b> | <b>55</b> |

## PART I: HEALTH PROFESSIONAL INFORMATION

### 1 INDICATIONS

LINESSA<sup>®</sup> (desogestrel and ethinyl estradiol tablets, USP) is indicated for:

- prevention of pregnancy

#### 1.1 Pediatrics

**Pediatrics (<18 years of age):** based on the data submitted and reviewed by Health Canada, the safety and efficacy of LINESSA in pediatric patients has not been established; therefore, Health Canada has not authorized an indication for pediatric use (see [7 WARNINGS AND PRECAUTIONS, 7.1.3 Pediatrics](#)).

#### 1.2 Geriatrics

**Geriatrics (> 65 years of age):** based on the data submitted and reviewed by Health Canada, LINESSA is not indicated for use in postmenopausal women.

### 2 CONTRAINDICATIONS

Combined hormonal contraceptives (CHCs) should not be used in the presence of any of the conditions listed below. Should any of the conditions appear for the first time during CHC use, the product should be stopped immediately.

- LINESSA is contraindicated in patients who are hypersensitive to this drug or to any ingredient in the formulation, including any non-medicinal ingredient, or component of the container. For a complete listing, see the [6 DOSAGE FORMS, STRENGTHS, COMPOSITION AND PACKAGING](#).
- Presence or history of venous thrombosis (deep vein thrombosis, pulmonary embolism);
- A history of actual cerebrovascular disorders;
- Presence or history of arterial thrombosis (myocardial infarction, cerebrovascular accident) or prodromal conditions (e.g., transient ischaemic attack, angina pectoris);
- valvular heart disease with complications
- Presence or history of severe hepatic disease as long as liver function values have not returned to normal
- Presence or history of liver tumours (benign or malignant);
- known or suspected sex steroid-influenced malignancies (e.g. of the genital organs or of the breast);
- undiagnosed abnormal vaginal bleeding;
- Steroid-dependent jaundice, cholestatic jaundice, history of jaundice of pregnancy;
- any ocular lesion arising from ophthalmic vascular disease, such as partial or

- complete loss of vision or defect in visual fields;
- known or suspected pregnancy;
- current or history of migraine with focal aura;
- history of actual pancreatitis if associated with severe hypertriglyceridemia.
- Known predisposition for arterial or venous or thrombosis:
  - severe hypertension (persistent values of  $\geq 160/100$  mmHg)
  - hereditary or acquired predisposition for venous or arterial thrombosis, such as Factor V Leiden mutation and activated protein C (APC-) resistance, antithrombin-III-deficiency, protein C deficiency, protein S deficiency, hyperhomocysteinemia (eg, due to MTHFR C677T, A1298 mutations), prothrombin mutation G20210A, and antiphospholipid-antibodies (anticardiolipin antibodies, lupus anticoagulant)
  - severe dyslipoproteinemia
  - smoking and over age 35
  - diabetes mellitus with vascular involvement
  - major surgery associated with an increased risk of postoperative thromboembolism (see [7 WARNINGS AND PRECAUTIONS](#))
  - prolonged immobilization (see [7 WARNINGS AND PRECAUTIONS](#))
  - Major surgery with prolonged immobilisation
- LINESSA is contraindicated for concomitant use with the medicinal products containing ombitasvir/paritaprevir/ritonavir with or without dasabuvir, and medicinal products containing glecaprevir/pibrentasvir or sofosbuvir/velpatasvir/voxilaprevir (see section [9 DRUG INTERACTIONS](#)).

### 3 SERIOUS WARNINGS AND PRECAUTIONS BOX

#### Serious Warnings and Precautions

- Cigarette smoking increases the risk of serious adverse effects on the heart and blood vessels. This risk increases with age and becomes significant in oral contraceptive users older than 35 years of age, and with the number of cigarettes smoked. For this reason, combination oral contraceptives, including LINESSA, should not be used by women who are over 35 years of age and smoke (see [Cardiovascular](#) section below).
- Patients should be counseled that birth control pills **DO NOT PROTECT** against sexually transmitted infections including HIV/AIDS. For protection against STIs, patients should be counseled to use condoms **IN COMBINATION WITH** birth control pills.

## 4 DOSAGE AND ADMINISTRATION

### 4.1 Dosing Considerations

Patients should be instructed to read the package insert prior to starting LINESSA and any time they are unsure of administration. If they have additional questions, they should call their doctor or clinic.

LINESSA tablets are prescribed as a 21-day or 28-day regimen. LINESSA tablets must be taken at approximately the same time every day until the pack is empty. The patient may begin taking LINESSA on Day 1 of her menstrual cycle (i.e. the first day of menstrual flow) or on the first Sunday after her period begins. If the period starts on Sunday, she should start that same day.

### 4.2 Recommended Dose and Dosage Adjustment

**LINESSA 21 (21-Day Regimen):** One coloured tablet is to be taken as follows for 21 consecutive days (three weeks): light yellow for 7 days; orange for 7 days and red for 7 days. Tablets are then discontinued for one week. The patient must not be off the pill for more than seven consecutive days. A new pack will be started on the eighth day. The patient will have a period during the seven days off the pill (the bleeding may be lighter and shorter than their usual period).

**LINESSA 28 (28-Day Regimen):** Tablets are taken sequentially following the arrows marked on the dispenser. One light yellow tablet is taken daily for 7 days; followed by one orange tablet for 7 days then one red tablet daily for 7 days. On the fourth week the patient will take one “inactive” green pill daily for the next seven consecutive days. A new pack will be started on the eighth day following completion of the green tablets. The patient will have a period during the seven days on the green pill. On this regimen the patient must not go a day without taking a pill.

### 4.3 Administration

It is recommended that LINESSA be taken at the same time each day. The patient should be counseled to associate the pill with some regular activity like eating a meal or going to bed.

The first-time user may wish to use a second method of birth control (e.g. condoms and spermicidal foam or gel) for the first seven days of the first cycle of pill use. This will provide a back-up in case pills are forgotten while they are getting used to taking them.

If spotting, light bleeding or feeling sick to their stomach occurs during the first three months the woman should be counseled to not stop taking the pill. The problem will usually go away. If it does not subside, the patient should consult with her doctor or clinic.

The dosage regimen should not be altered (i.e. the pill should not be stopped) even if the woman does not have sex very often.

When receiving any medical treatment, patients should tell their doctor that they are using birth control pills.

***Advice in case of vomiting***

In case of severe gastro-intestinal disturbance, absorption may not be complete and additional contraceptive measures should be taken. If vomiting occurs within 3-4 hours after tablet-taking, the advice concerning missed tablets ([4 DOSAGE AND ADMINISTRATION, 4.4 Missed Dose](#)) is applicable. If the woman does not want to change her normal tablet-taking schedule, she has to take the extra tablet(s) needed from another pack.

***Special Notes on Administration***

When to start LINESSA

**No hormonal contraceptive use in the preceding cycle:** Tablet taking should start on Day 1 of the woman's menstrual cycle or on the first Sunday after her period begins.

**Switching from another combination hormonal contraceptive (combined oral contraceptive (COC), vaginal ring, or transdermal patch):** The woman should start LINESSA preferably on the day after the last active tablet of her previous COC, but at the latest on the day following the usual tablet-free or inactive tablet of her previous COC. In case a vaginal ring or transdermal patch has been used, the woman should start using LINESSA preferably on the day of removal, but at the latest when the next application would have been due.

**Switching from a progestogen-only-method (mini-pill, injection, implant) or from a progestogen-releasing intrauterine system (IUS):** The woman may switch from the mini-pill to LINESSA on any day of her cycle. Patients using a progestogen injection should start LINESSA on the day the next injection is due. Patients using an implant or an IUS should start LINESSA on the day it is removed. In all cases, the woman should be advised to use an additional barrier method for the first 7 days of LINESSA use.

**Following first-trimester abortion:** The woman may start LINESSA immediately. When doing so, she need not take additional contraceptive measures.

**Following delivery or second-trimester abortion:** Women should be advised to start LINESSA at day 21 to 28 after delivery or second trimester abortion, after consulting with their physician. When starting later, the woman should be advised to additionally use a barrier method for the first 7 days of tablet-taking. However, if intercourse has already occurred, pregnancy should be excluded before the actual start of use or the woman should be advised to wait for her first menstrual period prior to starting LINESSA.

The increased risk of VTE during the postpartum period should be considered when restarting LINESSA (see [7 WARNINGS AND PRECAUTIONS](#)).

For breastfeeding women see [7 WARNINGS AND PRECAUTIONS, 7.1.2 Breast-feeding](#).

**4.4 Missed Dose**

The patient should be instructed to use the following chart if she misses one or more birth control pills (light yellow, orange, or red). She should be told to match the number of pills missed with the appropriate starting time for her dosing regimen.

| <b>Sunday Start</b>   | <b>Day One Start</b>   |
|---|--|
| <b>Miss One Pill</b>  | <b>Miss One Pill</b>   |
| Take it as soon as you remember and take the next pill at the usual time. This means that you might take 2 pills in one day.  | Take it as soon as you remember and take the next pill at the usual time. This means that you might take 2 pills in one day.   |
| <b>Miss Two Pills in a Row</b>  | <b>Miss Two Pills in a Row</b>   |
| <b>First Two Weeks:</b> <ol style="list-style-type: none"> <li>1. Take 2 pills the day you remember and 2 pills the next day.</li> <li>2. Then take 1 pill a day until you finish the pack.</li> <li>3. Use a back-up method of birth control if you have sex in the 7 days after you miss the pills.</li> </ol>  | <b>First Two Weeks:</b> <ol style="list-style-type: none"> <li>1. Take 2 pills the day you remember and 2 pills the next day.</li> <li>2. Then take 1 pill a day until you finish the pack.</li> <li>3. Use a back-up method of birth control if you have sex in the 7 days after you miss the pills.</li> </ol>   |
| <b>Third Week:</b> <ol style="list-style-type: none"> <li>1. Keep taking 1 pill a day until Sunday.</li> <li>2. On Sunday, safely discard the rest of the pack and start a new pack that day.</li> <li>3. Use a back-up method of birth control if you have sex in the 7 days after you miss the pills.</li> <li>4. You may not have a period this month.</li> </ol> <b>If you miss two periods in a row, call your doctor or clinic.</b> | <b>Third Week:</b> <ol style="list-style-type: none"> <li>1. Safely dispose of the rest of the pill pack and start a new pack that same day.</li> <li>2. Use a back-up method of birth control if you have sex in the 7 days after you miss the pills.</li> <li>3. You may not have a period this month.</li> </ol> <b>If you miss two periods in a row, call your doctor or clinic.</b> |
| <b>Miss Three or More Pills in a Row</b>  | <b>Miss Three or More Pills in a Row</b>   |



|   |  |
|---|--|
| <p><b>Anytime in the Cycle:</b></p> <ol style="list-style-type: none"> <li>1. Keep taking 1 pill a day until Sunday.</li> <li>2. On Sunday, safely discard the rest of the pack and start a new pack that day.</li> <li>3. Use a back-up method of birth control if you have sex in the 7 days after you miss the pills.</li> <li>4. You may not have a period this month.</li> </ol> <p><b>If you miss two periods in a row, call your doctor or clinic.</b></p> | <p><b>Anytime in the Cycle:</b></p> <ol style="list-style-type: none"> <li>1. Safely dispose of the rest of the pill pack and start a new pack that same day.</li> <li>2. Use a back-up method of birth control if you have sex in the 7 days after you miss the pills.</li> <li>3. You may not have a period this month.</li> </ol> <p><b>If you miss two periods in a row, call your doctor or clinic.</b></p> |
|---|--|

Missing pills can cause spotting or light bleeding, even if the missed pills are made up. The woman may also feel a little sick to her stomach on the days she takes two pills to make up for missed pills.

If a woman misses pills at any time, she could get pregnant. The greatest risks for pregnancy are starting a pack late or missing a pill(s) at the beginning or at the very end of the pack.

The patient should be counseled to always have another kind of birth control (such as condoms and spermicidal foam or gel) to use as a back-up in case they miss pills, and an extra full pack of pills available.

If the patient forgets more than one pill, two months in a row, they should be instructed to talk to their doctor or clinic. The patient may require further counseling about ways to make pill-taking easier or about using another method of birth control.

**NOTE to patients on the 28-day regimen (LINESSA 28):** If the patient forgets any of the 7 green pills (inactive pills) in Week 4, she should be advised to safely dispose of the pills she missed and then keep taking one pill each day until the pack is empty. A back-up method is not needed.

## 5 OVERDOSAGE

Serious ill effects have not been reported following acute ingestion of large doses of oral contraceptives by young children. Overdosage may cause nausea, and withdrawal bleeding may occur in females. There are no antidotes and further treatment should be symptomatic.

For management of a suspected drug overdose, contact your regional poison control centre.

## 6 DOSAGE FORMS, STRENGTHS, COMPOSITION AND PACKAGING

**Table – Dosage Forms, Strengths, Composition and Packaging.**

| Route of Administration | Dosage Form / Strength/Composition                              | Non-medicinal Ingredients  |
|-------------------------|---|--|
| Oral                    | Tablets/<br>0.100 mg desogestrel and 0.025 mg ethinyl estradiol | Hydroxypropyl, lactose monohydrate, methylcellulose, polyethylene glycol, starch, stearic acid, titanium dioxide, talc and vitamin E.  |
|                         | 0.125 mg desogestrel and 0.025 mg ethinyl estradiol             | The yellow and orange tablets also contain yellow ferric oxide. The orange and red tablets also contain red ferric oxide.  |
|                         | 0.150 mg desogestrel and 0.025 mg ethinyl estradiol             | LINESSA 28 also contains 7 green tablets containing the following non-medicinal ingredients: corn starch, FD&C Blue No.2 aluminum lake, hydroxypropyl methylcellulose, lactose monohydrate, magnesium stearate, polyethylene glycol, talc, titanium dioxide and yellow ferric oxide. |

### ***Dosage Forms***

**LINESSA 21:** Each sachet contains a blister card dispenser with 21 round film coated tablets:

- 7 light yellow containing 0.100 mg desogestrel and 0.025 mg ethinyl estradiol;
- 7 orange, containing 0.125 mg desogestrel and 0.025 mg ethinyl estradiol;
- 7 red, containing 0.150 mg desogestrel and 0.025 mg ethinyl estradiol.

**LINESSA 28:** Each sachet contains a blister card dispenser with 28 round film coated tablets consisting of the same three dosing phases as the 21–day pack and an additional seven (7) green tablets which do not contain any active ingredients.

### ***Composition***

LINESSA is a triphasic oral contraceptive containing two active components, desogestrel and ethinyl estradiol.

Each treatment cycle pack consists of three active dosing phases:

- light yellow tablets containing 0.100 mg desogestrel and 0.025 mg ethinyl estradiol
- orange tablets containing 0.125 mg desogestrel and 0.025 mg ethinyl estradiol, and
- red tablets containing 0.150 mg desogestrel and 0.025 mg ethinyl estradiol.

Inactive ingredients include vitamin E, pregelatinized starch, stearic acid, lactose monohydrate, hydroxypropyl methylcellulose, polyethylene glycol, titanium dioxide, talc, yellow ferric oxide, (in light yellow and orange tablets), and red ferric oxide (in orange and red tablets).

The inactive green tablets contained in the 28-day treatment cycle pack contain the following inert ingredients: lactose monohydrate, corn starch, magnesium stearate, hydroxypropyl methylcellulose, polyethylene glycol, titanium dioxide, FD&C Blue No. 2 aluminum lake, yellow ferric oxide, and talc.

### ***Packaging***

**LINESSA 21:** Push-through strips contain 21 (as listed below) round, biconvex, film coated tablets which are 5mm in diameter:

- Seven yellow are coded TR above 0 on one side and plain on the reverse side
- Seven orange tablets are coded TR above 6 on one side and plain on the reverse side
- Seven red tablets are coded TR above 1 on one side and plain on the reverse side

**LINESSA 28:** Push-through strips contain 28 (as listed below) round, biconvex, film coated tablets which are 5mm in diameter:

- Seven yellow are coded TR above 0 on one side and plain on the reverse side
- Seven orange tablets are coded TR above 6 on one side and plain on the reverse side
- Seven red tablets are coded TR above 1 on one side and plain on the reverse side
- Seven green tablets are coded KH above 2 on one side and plain on the reverse side

The push-through strip is a polyvinylchloride (PVC)/aluminum blister consisting of PVC film backed by aluminum foil with a heat-seal coating. Each blister is packed in a sealed aluminum laminated sachet. One sachet is packed in a printed cardboard box together with the package insert.

## **7 WARNINGS AND PRECAUTIONS**

Please see the [3 Serious Warnings and Precautions Box](#) at the beginning of Part I: Health Professional Information.

### **General**

#### **Discontinue Medication at the Earliest Manifestation of:**

- A. Thromboembolic and cardiovascular disorders, such as thrombophlebitis, pulmonary embolism, cerebrovascular disorders, myocardial ischemia, mesenteric thrombosis, and retinal thrombosis.

- B. Conditions which predispose to venous stasis and to vascular thrombosis (e.g., immobilization after accidents or confinement to bed during long-term illness). Other non-hormonal methods of contraception should be used until regular activities are resumed. For use of oral contraceptives when surgery is contemplated, see [Peri-operative Considerations](#).
- C. Visual defects- partial or complete
- D. Papilledema or ophthalmic vascular lesions
- E. Severe headache of unknown etiology or worsening of pre-existing migraine headache
- F. Increase in epileptic seizures

Throughout this section the general term combined hormonal contraceptives (CHC) is used when data exist for oral and non-oral contraceptives. The term combined oral contraceptives (COC) is used when data exist only for oral contraceptives.

The following information is provided from studies of combination oral contraceptives (COCs).

The use of combination hormonal contraceptives is associated with increased risks of several serious conditions including myocardial infarction, thromboembolism, stroke, hepatic neoplasia and gallbladder disease, although the risk of serious morbidity and mortality is small in healthy women without underlying risk factors. The risk of morbidity and mortality increases significantly if associated with the presence of other risk factors such as hypertension, hyperlipidemias, obesity and diabetes. Other medical conditions which have been associated with adverse circulatory events include systemic lupus erythematosus, hemolytic uremic syndrome, chronic inflammatory bowel disease (Crohn's disease or ulcerative colitis), sickle cell disease, valvular heart disease and atrial fibrillation.

The following conditions have been reported to occur or deteriorate with both pregnancy and COC use, although a direct association with COC's has not been firmly established: jaundice and/or pruritus related to cholestasis; gallstone formation; porphyria, systemic lupus erythematosus, hemolytic uremic syndrome, Sydenham's chorea, herpes gestationis, and otosclerosis-related hearing loss, (hereditary or acquired) angioedema.

The information contained in this section is principally from studies carried out in women who used combination oral contraceptives with higher formulations of estrogens and progestogens than those in common use today. The effect of long-term use of combination oral contraceptives with lower doses of both estrogen and progestogen remains to be determined.

## **Carcinogenesis and Mutagenesis**

### ***Breast Cancer***

Increasing age and a strong family history are the most significant risk factors for the development of breast cancer. Other established risk factors include obesity, nulliparity and late age at first full-term pregnancy. The identified groups of women that may be at increased risk of developing breast cancer before menopause are long-term users of oral contraceptives (more than eight years) and starters at early age. In a few women, the use of oral contraceptives may accelerate the growth of an existing but undiagnosed breast cancer. Since any potential increased risk related to oral contraceptive use is small, there is no reason to change prescribing habits at present.

Women receiving oral contraceptives should be instructed in self-examination of their breasts. Their physicians should be notified whenever any masses are detected. A yearly clinical breast examination is also recommended because, if a breast cancer should develop, estrogen-containing drugs may cause a rapid progression.

### ***Cervical Cancer***

The most important risk factor for cervical cancer is persistent human papillomavirus (HPV) infection. Some epidemiological studies have indicated that long-term use of Combination Oral Contraceptives (COCs) may further contribute to this increased risk but there continues to be controversy about the extent to which this finding is attributable to the confounding effects, e.g., cervical screening and sexual behavior including use of barrier contraceptives.

### ***Hepatocellular Carcinoma***

Hepatocellular carcinoma may be associated with oral contraceptives. The risk appears to increase with duration of hormonal contraceptive use. However, the attributable risk (the excess incidence) of liver cancers in oral contraceptive users is extremely small.

## **Cardiovascular**

### ***Predisposing Factors for Coronary Artery Disease***

Cigarette smoking increases the risk of serious cardiovascular side effects and mortality. Birth control pills increase this risk, especially with increasing age. Convincing data are available to support an upper age limit of 35 years for oral contraceptive use in women who smoke.

Other women who are independently at high risk for cardiovascular disease include those with diabetes, hypertension, abnormal lipid profile, or a family history of these. Whether oral contraceptives accentuate this risk is unclear.

In low risk, non-smoking women of any age, the benefits of oral contraceptive use outweigh the possible cardiovascular risks associated with low-dose formulations. Consequently, oral contraceptives may be prescribed for these women up to the age of menopause.

### ***Hypertension***

Patients with essential hypertension whose blood pressure is well-controlled may be given hormonal contraceptives but only under close supervision. If a significant elevation of blood pressure in previously normotensive or hypertensive subjects occurs at any time during the administration of the drug, cessation of medication is necessary.

### **Driving and Operating Machinery**

Due caution should be exercised when driving or operating a vehicle or potentially dangerous machinery.

### **Endocrine and Metabolism**

#### ***Diabetes***

Current low-dose oral contraceptives exert minimal impact on glucose metabolism. Diabetic patients, or those with a family history of diabetes, should be observed closely to detect any worsening of carbohydrate metabolism. Patients predisposed to diabetes who can be kept under close supervision may be given oral contraceptives. Young diabetic patients whose disease is of recent origin, well-controlled, and not associated with hypertension or other signs of vascular disease such as ocular fundal changes, should be monitored more frequently while using oral contraceptives.

#### ***Lipid and Other Metabolic Effects***

A small proportion of women will have adverse lipid changes while on oral contraceptives. Alternative contraception should be used in women with uncontrolled dyslipidemias. (See also [2 CONTRAINDICATIONS](#)). Elevations of plasma triglycerides may lead to pancreatitis and other complications.

#### ***Lactose Intolerance***

LINESSA contains < 65 mg lactose per tablet. Patients with lactase deficiency including Lapp lactase deficiency or glucose-galactose malabsorption on lactose-free diet, should consider this excipient when considering LINESSA for contraception.

### **Gastrointestinal**

Published epidemiological studies indicate a possible association of COC use and the development of Crohn's disease and ulcerative colitis, although this has not been firmly established.

### **Genitourinary**

#### ***Vaginal Bleeding***

Persistent irregular vaginal bleeding requires assessment to exclude underlying pathology.

#### ***Fibroids***


Patients with fibroids (leiomyomata) should be carefully observed. Sudden enlargement, pain, or tenderness requires discontinuation of the use of oral contraceptives.

## Hematologic

Epidemiological studies have shown an association between the use of CHCs and an increased risk of arterial and venous thrombotic and thromboembolic diseases such as myocardial infarction, stroke, deep venous thrombosis, and pulmonary embolism.

Epidemiological studies have shown that the incidence of venous thromboembolism (VTE) in users of CHC with low estrogen content (<50 mcg ethinyl estradiol) ranges from about 3 to 12 cases per 10,000 women-years, but this risk estimate varies according to the progestogen. This compares with 1 to 5 cases per 10,000 women-years for non-CHC users.

The use of CHC carries an increased risk of VTE compared with no use. The excess risk of VTE is highest during the first year a woman ever uses a CHC. The risk is also increased when patients initially start a CHC or restart the same or different CHC after a break in use of 4 weeks or more. The increased risk is less than the risk of VTE associated with pregnancy, which is estimated as 5 to 20 cases per 10,000 women-years or the risk in the postpartum period which is estimated as 40-65 cases per 10,000 women-years. VTE is fatal in 1-2% of cases. The figure below shows the risk of developing a VTE for women who are not pregnant and do not use CHCs, for women who use CHCs, for pregnant women, and for women in the postpartum period.



\*CHC=combined hormonal contraception

\*\*Pregnancy data based on actual duration of pregnancy in the reference studies. Based on a model assumption that pregnancy duration is nine months, the rate is 7 to 27 per 10 000 WY.

Several epidemiological studies indicate that third generation oral contraceptives, including those containing desogestrel, are associated with a higher risk of venous thromboembolism than certain second generation oral contraceptives. These studies indicate an approximate 2-fold difference in risk, which corresponds to 1-2 cases of venous thromboembolism per 10,000 women-years of use. However, data from additional studies have not shown this difference in risk.

Extremely rarely, thrombosis has been reported to occur in other blood vessels, e.g., hepatic, mesenteric, renal, cerebral or retinal veins and arteries, in CHC users.

Symptoms of venous or arterial thrombotic/thromboembolic events or of a cerebrovascular accident can include: unilateral leg pain and/ or swelling; sudden severe pain in the chest, whether or not it radiates to the left arm; sudden breathlessness; sudden onset of coughing; any unusual, severe, prolonged headache; sudden partial or complete loss of vision; diplopia; slurred speech or aphasia; vertigo; collapse with or without focal seizure; weakness or very marked numbness suddenly affecting one side or one part of the body; motor disturbances; 'acute' abdomen.

#### ***Other Risk Factors for Venous Thromboembolism***

risk of venous thromboembolism increases with:

- increasing age;
- a personal history,
- a positive family history ( i.e. venous thromboembolism ever in a sibling or parent at a relatively early age). If a hereditary or acquired predisposition for venous thromboembolism is suspected, the woman should be referred to a specialist for advice before deciding on any CHC use.
- obesity (body mass index  $>30 \text{ kg/m}^2$ )
- smoking
- systemic lupus erythematosus.
- and possibly also with superficial thrombophlebitis and varicose veins and;
  
- The risk of VTE may be temporarily increased with prolonged immobilization, major surgery, any surgery to the legs or major trauma. In these situations, it is advisable to discontinue CHC use (in the case of elective surgery at least four weeks in advance) and not to resume until two weeks after complete remobilisation (see [2 CONTRAINDICATIONS](#)).

Patients with superficial thrombophlebitis and varicose veins and leg cast should be closely supervised.



### ***Other Risk Factors for Arterial Thromboembolism***

The risk of arterial thromboembolic complications increases with:

- increasing age;
- smoking (with heavier smoking and increasing age the risk further increases, especially in women over 35 years of age);
- dyslipoproteinaemia;
- obesity (body mass index over 30 kg/m<sup>2</sup>);
- hypertension;
- migraine;
- valvular heart disease;
- atrial fibrillation;
- a positive family history (i.e. arterial thrombosis ever in a sibling or parent at a relatively early age). If a hereditary predisposition is suspected, the woman should be referred to a specialist for advice before deciding about any hormonal contraceptive use.

The increased risk of thromboembolism in the puerperium must be considered when LINESSA is being considered for contraception.

An increase in frequency or severity of migraine during COC use (which may be prodromal of a cerebrovascular event) may be a reason for immediate discontinuation of the COC.

Biochemical factors that may be indicative of hereditary or acquired predisposition for venous or arterial thrombosis include Activated Protein C (APC) resistance, hyperhomocysteinaemia, antithrombin-III deficiency, protein C deficiency, protein S deficiency, antiphospholipid antibodies (anticardiolipin antibodies, lupus anticoagulant).

When considering risk/benefit, the healthcare professional should take into account that adequate treatment of a condition may reduce the associated risk of thrombosis. Other medical conditions which have been associated with adverse circulatory events include diabetes mellitus, systemic lupus erythematosus, haemolytic uraemic syndrome, chronic inflammatory bowel disease (Crohn's disease or ulcerative colitis) and sickle cell disease.

### **Hepatic/Biliary/Pancreatic**

Acute or chronic disturbances of liver function may necessitate the discontinuation of COC use until markers of liver function return to normal.

### ***Jaundice***

Patients who have had jaundice should be given oral contraceptives only with great care and under close observation. Oral contraceptive-related cholestasis has been described in women with a history of pregnancy-related cholestasis. Women with a history of cholestasis may have the condition recur with subsequent hormonal contraceptive use.

The development of severe generalized pruritus or icterus requires that the medication be withdrawn until the problem is resolved.

If a patient develops jaundice that proves to be cholestatic in type, the use of oral contraceptives should not be resumed. In patients taking hormonal contraceptives, changes in the composition of the bile may occur and an increased incidence of gallstones has been reported.

### ***ALT elevations***

During clinical trials with patients treated for hepatitis C virus infections (HCV) with the medicinal products containing ombitasvir /paritaprevir/ritonavir and dasabuvir with or without ribavirin, transaminase (ALT) elevations greater than 5 times the upper limit of normal (ULN) occurred significantly more frequent in women using ethinylestradiol-containing medications such as combined hormonal contraceptives (CHCs) (see section [CONTRAINDICATIONS](#) and [DRUG INTERACTIONS](#)). Additionally, in patients treated with glecaprevir/pibrentasvir or sofosbuvir/velpatasvir/voxilaprevir, ALT elevations were also observed in women using ethinylestradiol-containing medications such as CHCs.

LINESSA should be discontinued prior to starting therapy with the following combination drug regimens 1) ombitasvir/partaprevir/ritonavir and dasabuvir with or without ribavirin, 2) glecaprevir/pibrentasvir, 3) sofosbuvir/velpatasvir/voxilaprevir. LINESSA can be restarted 2 weeks following completion of therapy for HCV. The contraceptive efficacy of LINESSA may be reduced in patients taking HCV or other hepatically cleared medication.

### ***Gallbladder Disease***

Patients taking oral contraceptives have a greater risk of developing gallbladder disease requiring surgery within the first year of use. The risk may double after four or five years.

### ***Hepatic Nodules***

Hepatic nodules (adenoma and focal nodular hyperplasia) have been reported, particularly in long-term users of oral contraceptives. Although these lesions are extremely rare, they have caused fatal intra-abdominal hemorrhage and should be considered in women presenting with an abdominal mass, acute abdominal pain, or evidence of intra-abdominal bleeding.

### ***Immune***

#### ***Angioedema***

Exogenous estrogens may induce or exacerbate symptoms of angioedema, in particular in women with hereditary or acquired angioedema.

## **Monitoring and Laboratory Tests**

### ***Physical Examination and Follow-up***

Prior to the initiation or reinstatement of LINESSA a complete medical history (including family history) should be taken and pregnancy must be ruled out. Blood pressure should be measured and if clinically indicated physical examination should be performed, guided by the contra-indications and warnings.

In addition, disturbances of the clotting system must be ruled out if any members of the family have suffered from thromboembolic diseases (e.g., deep vein thrombosis, stroke, myocardial infarction) at a young age. Breasts, liver, extremities and pelvic organs should be examined and a Papanicolaou (PAP) smear should be taken if the patient has been sexually active.

The woman should also be instructed to carefully read the user leaflet and to adhere to the advice given. The frequency and nature of further periodic checks should be based on established practice guidelines and be adapted to the individual woman.

The first follow-up visit should be done three months after oral contraceptives are prescribed. Thereafter, examinations should be performed at least once a year or more frequently if indicated. At each annual visit, examination should include those procedures that were done at the initial visit as outlined above or per recommendations of the Canadian Task Force on the Periodic Health Examination.

## **Neurologic**

### ***Migraine and Headache***

The onset or exacerbation of migraine or the development of headache of a new pattern that is recurrent, persistent or severe, requires discontinuation of hormonal contraceptives and evaluation of the cause. Women with migraine headaches who take oral contraceptives may be at increased risk of stroke (see [2 CONTRAINDICATIONS](#)).

## **Ophthalmologic**

Patients who are pregnant or are taking oral contraceptives, may experience corneal edema that may cause visual disturbances and changes in tolerance to contact lenses, especially of the rigid type. Soft contact lenses usually do not cause disturbances. If visual changes or alterations in tolerance to contact lenses occur, temporary or permanent cessation of wear may be advised.

## **Peri-Operative Considerations**

There is an increased risk of thromboembolic complications in oral contraceptive users after major surgery. If feasible, oral contraceptives should be discontinued and an alternative method substituted at least one month prior to **MAJOR** elective surgery. Oral contraceptives should not be resumed until the first menstrual period after hospital discharge following surgery.

## **Psychiatric**

Patients with a history of emotional disturbances, especially the depressive type, may be more prone to have a recurrence of depression while taking oral contraceptives. In cases of a serious recurrence, a trial of an alternate method of contraception should be made which may help to clarify the possible relationship. Women with premenstrual syndrome (PMS) may have a varied response to oral contraceptives, ranging from symptomatic improvement to worsening of the condition.

## **Renal**

### ***Fluid Retention***

Hormonal contraceptives may cause some degree of fluid retention. They should be prescribed with caution, and only with careful monitoring in patients with conditions which might be aggravated by fluid retention.

## **Sexual Health Function/ Reproduction**

### ***Return to Fertility***

After discontinuing oral contraceptive therapy, the patient should delay pregnancy until at least one normal spontaneous cycle has occurred in order to date the pregnancy. An alternate contraceptive method should be used during this time.

### ***Amenorrhea***

In some women, withdrawal bleeding may not occur during the tablet-free interval. If the COC has been taken according to directions, it is unlikely that the woman is pregnant. However, if the COC has not been taken according to directions prior to the first missed withdrawal bleed, or if two withdrawal bleeds are missed, pregnancy must be ruled out before COC use is continued.

Women having a history of oligomenorrhea, secondary amenorrhea, or irregular cycles may remain anovulatory or become amenorrheic following discontinuation of estrogen-progestin combination therapy.

Amenorrhea, especially if associated with breast secretion that continues for six months or more after withdrawal warrants a careful assessment of hypothalamic-pituitary function.

### ***Reduced Efficacy***

The efficacy of COCs may be reduced in the event of missed tablets, gastro-intestinal disturbances or concomitant medication (see [9 DRUG INTERACTIONS](#)).

## **Skin**

Chloasma may occasionally occur with use of COCs, especially in women with a history of chloasma gravidarum. Women with a tendency to chloasma should avoid exposure to the sun or ultraviolet radiation while taking COCs.

## 7.1 Special Populations

### 7.1.1 Pregnant Women

Oral contraceptives should not be taken by pregnant women. If pregnancy occurs during treatment with LINESSA, further intake should be stopped. However, if conception accidentally occurs while taking the pill, there is no conclusive evidence that the estrogen and progestin contained in the oral contraceptive will damage the developing child.

### 7.1.2 Breast-feeding

In breastfeeding women, the use of oral contraceptives results in the hormonal components being excreted in breast milk and may reduce its quantity and quality. Published studies have indicated that during lactation, 0.1% of the daily maternal dose of levonorgestrel and 0.02% of the daily maternal dose of ethinyl estradiol could be transferred to the newborn via milk. Adverse effects on the child have been reported, including jaundice and breast enlargement. The nursing mother should be advised not to use oral contraceptives but to use other forms of contraception until she has completely weaned her child.

### 7.1.3 Pediatrics

**Pediatrics (<18 years of age):** based on the data submitted and reviewed by Health Canada, the safety and efficacy of LINESSA in pediatric patients has not been established; therefore, Health Canada has not authorized an indication for pediatric use (see [1 INDICATIONS, 1.1 Pediatrics](#)).

Use of this product before menarche is not indicated.

### 7.1.4 Geriatrics

LINESSA is not indicated for use in postmenopausal women.

## 8 ADVERSE REACTIONS

### 8.1 Adverse Reaction Overview

An increased risk of the following serious adverse reactions has been associated with the use of combination hormonal contraceptives:

- arterial and venous thromboembolism
- benign and malignant hepatic tumours
- cerebral hemorrhage
- cerebral thrombosis
- congenital anomalies
- gallbladder disease
- hypertension
- mesenteric thrombosis

- myocardial infarction
- neuro-ocular lesions, (e.g., retinal thrombosis)
- pulmonary embolism
- thrombophlebitis

The following adverse reactions also have been reported in patients receiving combination hormonal contraceptives: nausea and vomiting, usually the most common adverse reaction, occurs in approximately 10% or less of patients during the first cycle. The following other reactions, as a general rule, are seen less frequently or only occasionally:

- abdominal pain
- amenorrhea during and after treatment
- angioedema (exogenous estrogens may induce or exacerbate symptoms of angioedema in women with hereditary or acquired angioedema)<sup>a</sup>
- auditory disturbances
- breakthrough bleeding
- breast changes: tenderness, enlargement, and secretion
- cataracts
- changes in appetite
- change in corneal curvature (steepening)
- changes in glucose tolerance or effect on peripheral insulin resistance
- changes in libido
- change in menstrual flow
- change in weight (increase or decrease)
- chloasma or melasma which may persist
- cholestatic jaundice
- chorea
- Crohn's disease
- cystitis-like syndrome
- diarrhea
- dizziness
- dysmenorrhea
- edema
- endocervical hyperplasias
- erythema multiforme
- erythema nodosum
- gallstone formation<sup>a</sup>
- gastrointestinal symptoms (such as abdominal cramps and bloating)
- headache
- hemolytic uremic syndrome
- hemorrhagic eruption
- herpes gestationis<sup>a</sup>
- hirsutism
- hypersensitivity
- hypertension<sup>a</sup>

- hypertriglyceridemia (increased risk of pancreatitis when using COCs)
- impaired renal function
- increase in size of uterine leiomyomata
- intolerance to contact lenses
- jaundice related to cholestasis<sup>a</sup>
- liver function disturbances
- loss of scalp hair
- mental depression
- migraine
- nervousness
- optic neuritis
- otosclerosis-related hearing loss<sup>a</sup>
- pancreatitis
- porphyria
- possible diminution in lactation when given immediately post-partum
- premenstrual-like syndrome
- pruritus related to cholestasis<sup>a</sup>
- rash (allergic)
- Raynaud's phenomenon
- reduced tolerance to carbohydrates
- retinal thrombosis
- rhinitis
- spotting
- Sydenham's chorea<sup>a</sup>
- Systemic lupus erythematosus<sup>a</sup>
- temporary infertility after discontinuance of treatment
- ulcerative colitis
- urticaria
- vaginal candidiasis
- vaginal discharge
- vaginitis

<sup>a</sup>Occurrence or deterioration of conditions for which association with COC use is not conclusive.

## 8.2 Clinical Trial Adverse Reactions

*Because clinical trials are conducted under very specific conditions the adverse drug reaction rates observed in the clinical trials may not reflect the rates observed in practice and should not be compared to the rates in the clinical trials of another drug. Adverse drug reaction information from clinical trials is useful for identifying drug-related adverse events and for approximating rates.*

Two multicenter 6-cycle controlled efficacy and safety studies were conducted in 5,552 women.

A list of adverse events experienced by > 1% of the subjects is listed in Table 1.

**Table 1: Incidence of All Adverse Experiences (>1%) (All Subjects Treated Group) to LINESSA**

| Preferred (WHOART) Term                               | Incidence During Study |      |
|---|------------------------|------|
|   | n                      | %    |
| <b>LINESSA (Number of Subjects)</b>                   | <b>( n=2768)</b>       |      |
| <b><u>Body as a Whole-General</u></b>                 |                        |      |
| Influenza-like symptoms                               | 97                     | 3.5  |
| Back pain   | 80                     | 2.9  |
| Allergy   | 59                     | 2.1  |
| Fatigue   | 48                     | 1.7  |
| Fever   | 29                     | 1.0  |
| <b><u>Central &amp; Peripheral Nervous System</u></b> |                        |      |
| Headache  | 420                    | 15.2 |
| Migraine  | 34                     | 1.2  |
| <b><u>Gastrointestinal System</u></b>                 |                        |      |
| Nausea  | 225                    | 8.1  |
| Diarrhea  | 61                     | 2.2  |
| Flatulence  | 53                     | 1.9  |
| Dyspepsia   | 46                     | 1.7  |
| Vomiting  | 43                     | 1.6  |
| Abdominal pain  | 33                     | 1.2  |
| <b><u>Metabolic &amp; Nutritional</u></b>             |                        |      |
| Weight increase                                       | 64                     | 2.3  |
| <b><u>Musculoskeletal system</u></b>                  |                        |      |
| Myalgia   | 40                     | 1.4  |
| Arthralgia  | 37                     | 1.3  |
| <b><u>Psychiatric</u></b>                             |                        |      |
| Emotional Lability                                    | 117                    | 4.2  |
| Depression  | 45                     | 1.6  |
| Insomnia  | 30                     | 1.1  |
| Nervousness   | 28                     | 1.0  |



| Preferred (WHOART) Term              | Incidence During Study |      |
|--------------------------------------|------------------------|------|
|                                      | n                      | %    |
| <b>LINESSA (Number of Subjects)</b>  | <b>( n=2768)</b>       |      |
| <b><u>Reproductive, Female</u></b>   |                        |      |
| Intermenstrual bleeding              | 181                    | 6.5  |
| Dysmenorrhoea                        | 150                    | 5.4  |
| Breast pain female                   | 139                    | 5.0  |
| Moniliasis genital                   | 132                    | 4.8  |
| Vaginitis                            | 57                     | 2.1  |
| Pelvic cramping                      | 36                     | 1.3  |
| <b><u>Resistance Mechanism</u></b>   |                        |      |
| Herpes simplex                       | 30                     | 1.1  |
| <b><u>Respiratory System</u></b>     |                        |      |
| Upper respiratory tract infection    | 328                    | 11.8 |
| Sinusitis                            | 227                    | 8.2  |
| Pharyngitis                          | 134                    | 4.8  |
| Bronchitis                           | 70                     | 2.5  |
| Coughing                             | 50                     | 1.8  |
| Rhinitis                             | 51                     | 1.8  |
| <b><u>Secondary Terms-Events</u></b> |                        |      |
| Cervical smear test PAP II           | 51                     | 1.8  |
| Inflicted injury                     | 47                     | 1.7  |
| <b><u>Skin &amp; Appendages</u></b>  |                        |      |
| Acne                                 | 80                     | 2.9  |
| Rash                                 | 35                     | 1.3  |
| <b><u>Urinary System</u></b>         |                        |      |
| Urinary tract infection              | 112                    | 4.0  |
| Cystitis                             | 30                     | 1.1  |

**Notes:** This table contains all adverse events which occurred during treatment, including those deemed to be not related or unlikely related, in addition to those which were deemed to be possibly related, probably related and related.

This table contains counts of subjects. Within each treatment group, percentages based on the number of subjects with an event or without an event divided by the total number of subjects in each demographic subgroup. Adverse experiences that stopped before first dose date or started after last dose date were excluded from this table.

### 8.3 Less Common Clinical Trial Adverse Reactions

Rare adverse events (<1%) which were observed in clinical trials and deemed to be at least possibly related to LINESSA are as follows:

*Body as a whole-general:* Chest pain, crying abnormal, hot flushes, leg pain, edema, peripheral edema, pain, rigors, syncope, vertebral disk disorder.

*Cardiovascular, general:* Hypertension

*Central & peripheral nervous system:* leg Cramps, dizziness, involuntary muscle contractions, tremor

*Gastrointestinal system:* Constipation, eructation, irritable bowel syndrome

*Hearing & vestibular:* Ear disorder, earache, motion sickness

*Liver & biliary system:* Bilirubinaemia, cholecystitis, cholelithiasis, increased hepatic enzymes, abnormal hepatic function, SGOT increased, SGPT increased

*Metabolic & nutritional:* Hypercholesterolaemia, hyperglycaemia, hypertriglyceridaemia, LDH increased, generalised edema, edema of the legs, peripheral edema

*Neoplasms:* Breast fibroadenosis, female breast neoplasm, cervical uterine polyp, ovarian cyst, uterine fibroid

*Platelet bleeding & clotting:* Epistaxis, gingival bleeding, purpura

*Psychiatric:* Agitation, anorexia, anxiety, increased appetite, impaired concentration, confusion, dyspareunia, decreased libido, neurosis, somnolence, suicide attempt

*Red blood cell:* Anaemia

*Female reproductive system:* Amenorrhea, breast discharge, breast engorgement, breast enlargement, cervicitis, cervix lesion, non-puerperal lactation, leukorrhea, menorrhagia, menstrual disorder, ovarian pain, female perineal pain, premenstrual tension, uterine contractions, uterine haemorrhage, vaginal bleeding, vaginal discomfort, vaginal haemorrhage, vulva disorder

*Resistance mechanism:* Infection, viral infection

*Secondary terms, events:* Ectropion

*Skin & appendages:* Alopecia, chloasma, dermatitis, eczema, erythema nodosum, folliculitis, abnormal hair texture, hypertrichosis, melanosis, abnormal pigmentation, pruritus, genital pruritus, skin discolouration, dry skin, increased sweating

*Urinary system:* Dysuria, haematuria, micturition frequency, abnormal urine

*Vascular:* Flushing, thrombophlebitis, deep vein disorders

*Vision:* Photophobia, abnormal vision, xerophthalmia

*White cell & reticular endothelial system:* Lymphadenopathy

## 8.4 Post-Market Adverse Reactions

Additional adverse events which have been reported occasionally since the introduction of LINESSA to the market include: peripheral edema, cyst, hypoesthesia, menorrhagia, metrorrhagia, mood swings, abdominal distention, bleeding tendency, angioneurotic edema, drug exposure during pregnancy, pruritic rash, emotional lability, back pain, pollakiuria, fluid retention, pruritis, frequent bowel movements. These adverse events are compiled from spontaneous reports and are listed regardless of whether or not there was a possible causal relation to LINESSA.

The most serious undesirable effects associated with the use of COCs are listed in [7 WARNINGS AND PRECAUTIONS](#). Other side effects that have been reported in users of COCs but for which the association has been neither confirmed nor refuted are found in Table 2.

**Table 2: Post Market Adverse Events occurring with Combined Oral Contraceptives**

| Body system                            | Common (more than / equal to 1/100)                      | Common/Uncommon (more than/ equal to 1/1000 and less than 100) | Rare (less than 1/1000)  |
|--|--|--|--|
| Immune system disorders                |  |  | Hypersensitivity   |
| Metabolism and nutrition disorders     |  | Fluid retention  |  |
| Psychiatric disorders                  | Headache, Depressed mood, mood altered, Libido decreased |  | Libido increased   |
| Nervous system disorders               | Headache, Migraine                                       |  |  |
| Eye disorders                          |  |  | Contact lens intolerance   |
| Vascular disorders                     |  |  | Venous thromboembolism <sup>2</sup><br>Arterial thromboembolism <sup>2</sup> |
| Gastrointestinal disorders             | Nausea, , abdominal pain, Vomiting, diarrhea             |  |  |
| Skin and subcutaneous tissue disorders |  | Rash, urticaria  | Erythema nodosum, erythema multiforme  |

| Body system                              | Common (more than / equal to 1/100) | Common/Uncommon (more than/ equal to 1/1000 and less than 100) | Rare (less than 1/1000)             |
|--|-------------------------------------|--|-------------------------------------|
| Reproductive system and breast disorders | Breast pain, breast tenderness,     | Breast enlargement   | Vaginal discharge, breast discharge |
| Investigations                           | Weight increased                    |  | Weight decreased                    |

<sup>1</sup>The most appropriate MedDRA term (version 6.1) to describe a certain adverse reaction is listed. Synonyms or related conditions are not listed, but should be taken into account as well.

<sup>2</sup> Incidence in observational cohort studies of  $\geq 1/10000$  to  $< 1/1000$  women-years.

## 9 DRUG INTERACTIONS

### 9.1 Drug Interactions Overview

Interactions between desogestrel/ethinyl estradiol and other medicinal products have been reported in the literature which may alter the response to either agent (see [9.2 Drug-Drug Interactions](#)). No formal drug-drug interaction studies were conducted with LINESSA.

*Hepatic metabolism:* Interactions can occur with medicinal or herbal products that induce microsomal enzymes, specifically cytochrome P450 enzymes (CYP), which can result in increased clearance reducing plasma concentrations of sex hormones and may decrease the effectiveness of combined oral contraceptives, including LINESSA.

Enzyme induction can occur after a few days of treatment. Maximum enzyme induction is generally observed within a few weeks. After drug therapy is discontinued, enzyme induction can last for about 28 days.

A barrier contraceptive method should be used in addition to LINESSA during administration of the hepatic enzyme-inducing medicinal product, and for 28 days after discontinuation of the hepatic enzyme-inducing medicinal product.

For women on long-term therapy with enzyme-inducing medicinal products an alternative method of contraception unaffected by enzyme-inducing medicinal products should be considered.

The prescribing information of concomitant medications should be consulted to identify potential interactions.

### 9.2 Drug-Drug Interactions

The concurrent administration of oral contraceptives with other drugs may result in an altered response to either agent (Table 3 and 4). Reduced effectiveness of the oral

contraceptive, should it occur, is more likely with the low-dose formulations. It is important to ascertain all drugs that a patient is taking, both prescription and non-prescription, before oral contraceptives are prescribed.

**Table 3: Drugs Which May Decrease the Efficacy of Oral Contraceptives**

| Class of Compound           | Drug  | Proposed Mechanism  | Suggested Management  |
|-----------------------------|---|---|---|
| Antacids                    |   | Decreased intestinal absorption of progestins.  | Dose two hours apart  |
| Antibiotics (30)            | Ampicillin<br>Cotrimoxazole<br>Penicillin   | Enterohepatic circulation disturbance, intestinal hurry.  | For short course, use additional method or use another drug. For long course, use another method. |
|                             | Rifampicin  | Increased metabolism of progestins. Suspected acceleration of estrogen metabolism.  | Use another method.   |
|                             | Chloramphenicol<br>Metronidazole<br>Neomycin<br>Nitrofurantoin<br>Sulfonamides<br>Tetracyclines                                     | Induction of hepatic microsomal enzymes. Also disturbance of enterohepatic circulation.   | For short course, use additional method or use another drug. For long course, use another method. |
|                             | Troleandomycin  | May retard metabolism of oral contraceptives, increasing the risk of cholestatic jaundice.  |   |
| Anticonvulsants (31-33)     | Carbamazepine<br>Ethosuximide<br>Felbamate<br>Lamotrigine<br>Oxcarbazepine<br>Phenobarbital<br>Phenytoin<br>Primidone<br>Topiramate | Induction of hepatic microsomal enzymes. Rapid metabolism of estrogen and increased binding of progestin and ethinyl estradiol to SHBG. | Use higher dose oral contraceptives (50 mcg ethinyl estradiol) another drug or another method.    |
| Antifungals                 | Griseofulvin  | Stimulation of hepatic metabolism of contraceptive steroids may occur.  | Use another method.   |
| Cholesterol Lowering Agents | Clofibrate  | Reduces elevated serum triglycerides and  | Use another method.   |

| <b>Class of Compound</b>                        | <b>Drug</b>  | <b>Proposed Mechanism</b>  | <b>Suggested Management</b>   |
|---|--|--|---|
|   |  | cholesterol; this reduces oral contraceptive efficacy.                             |   |
| HCV Protease Inhibitors                         | Boceprevir<br>Telaprevir   | Remains to be confirmed  | Use another drug or another non-hormonal method of contraception.   |
| HIV protease inhibitors                         | Ritonavir  | Induction of hepatic microsomal enzymes.   | Use another drug or another method.   |
| Non-nucleoside reverse transcriptase inhibitors | Nevirapine   | Induction of hepatic microsomal enzymes.   | Use another drug or another method.   |
| Sedatives and Hypnotics                         | Barbiturates<br>Benzodiazepines<br>Chloral Hydrate<br>Glutethimide<br>Meprobamate        | Induction of hepatic microsomal enzymes.   | For short course, use additional method or another drug. For long course use another method or higher dose oral contraceptives. |
| Other Drugs                                     | Analgesics<br>Antihistamines<br>Antimigraine preparations<br>Phenylbutazone<br>Vitamin E | Reduced oral contraceptive efficacy has been reported.<br>Remains to be confirmed. |   |

Oral contraceptives may interfere with the metabolism of other drugs. Accordingly, plasma and tissue concentrations may either increase (e.g., cyclosporine) or decrease (e.g., lamotrigine).

**Table 4: Drugs Which May Decrease the Efficacy of Oral Contraceptives**

| <b>Class of Compound</b>       | <b>Drug</b>                    | <b>Modification of Other Drug Action</b>  | <b>Suggested Management</b>  |
|--------------------------------|--------------------------------|---|--|
| Alcohol                        |                                | Possible increased levels of ethanol or acetaldehyde.   | Use with caution   |
| Alpha-II Adrenoreceptor Agents | Clonidine                      | Sedation effect increased.  | Use with caution.  |
| Anticoagulants                 | All                            | oral contraceptives increase clotting factors, decrease efficacy. However oral contraceptives may potentiate action in some patients. | Use another method.  |
| Anticonvulsants                | All                            | Estrogens may increase risk of seizures.  | Use another method.  |
|                                | Lamotrigine                    | Decrease lamotrigine levels, may lead to breakthrough seizures.   | Use another method.  |
| Antidiabetic drugs             | Oral hypoglycemics and insulin | Oral contraceptives may impair glucose tolerance and increase blood glucose.  | Use low-dose estrogen and progestin oral contraceptive or another method. Monitor blood glucose. |
| Antihypertensive agents        | Guanethidine and Methyldopa    | Estrogen component cause sodium retention, progestin has no effect.   | Use low estrogen oral contraceptive or use another method.                                       |
|                                | Beta Blockers                  | Increased drug effect (decreased metabolism).   | Adjust dose of drug if necessary. Monitor cardiovascular status.                                 |
| Antipyretics                   | Acetaminophen                  | Increased metabolism and renal clearance.   | Dose of drug may have to be increased.   |
|                                | Antipyridine                   | Impaired metabolism.  | Decrease dose of drug.   |
|                                | ASA                            | Effects of ASA may be decreased by the short-term use of oral contraceptives.   | Patients on chronic ASA therapy may require an increase in ASA dosage.                           |

| <b>Class of Compound</b>    | <b>Drug</b>   | <b>Modification of Other Drug Action</b>  | <b>Suggested Management</b>  |
|-----------------------------|---------------|---|--|
| Aminocaproic Acid           |               | Theoretically, a hypercoagulable state may occur because oral contraceptives augment clotting factors.                  | Avoid concomitant use.   |
| Betamimetic Agents          | Isoproterenol | Estrogen causes decreased response to these drugs.  | Adjust dose of drug as necessary. Discontinuing oral contraceptives can result in excessive drug activity. |
| Caffeine                    |               | The actions of caffeine may be enhanced as oral contraceptives may impair the hepatic metabolism of caffeine.           | Use with caution.  |
| Cholesterol Lowering Agents | Clofibrate    | Their action may be antagonized by oral contraceptives. Oral contraceptives may also increase metabolism of clofibrate. | May need to increase dose of clofibrate.   |
| Corticosteroids             | Prednisone    | Markedly increased serum levels.  | Possible need for decrease in dose.  |
| Cyclosporine                |               | May lead to an increase in cyclosporine levels and hepatotoxicity.  | Monitor hepatic function. The cyclosporine dose may have to be decreased.                                  |
| Folic Acid                  |               | oral contraceptives have been reported to impair folate metabolism.   | May need to increase dietary intake, or supplement.  |
| Meperidine                  |               | Possible increased analgesia and CNS depression due to decreased metabolism of meperidine.                              | Use combination with caution.  |



| Class of Compound              | Drug  | Modification of Other Drug Action  | Suggested Management   |
|--------------------------------|---|--|--|
| Phenothiazine<br>Tranquilizers | All phenothiazines, reserpine and similar drugs.      | Estrogen potentiates the hyperprolactinemia effect of these drugs.                         | Use other drugs or lower dose oral contraceptives. If galactorrhea or hyperprolactinemia occurs, use other method. |
| Sedatives and Hypnotics        | Chlordiazepoxide<br>Lorazepam<br>Oxazepam<br>Diazepam | Increased effect (increased metabolism)  | Use with caution   |
| Theophylline                   | All   | Decreased oxidation, leading to possible toxicity.   | Use with caution. Monitor theophylline levels.   |
| Tricyclic Antidepressants      | Clomipramine (possibly others)                        | Increased side effects; i.e. depression.   | Use with caution   |
| Vitamin B <sub>12</sub>        |   | oral contraceptives have been reported to reduce serum levels of Vitamin B <sub>12</sub> . | May need to increase dietary intake, or supplement.  |

Several of the anti-HIV/HCV protease inhibitors (e.g., ritonavir, telaprevir, boceprevir) and non-nucleoside reverse transcriptase inhibitors (e.g., nevirapine) have been studied with coadministration of oral combination hormonal contraceptives; significant changes (increase and decrease) in the mean AUC of the estrogen and progestin and the potential to affect hepatic metabolism have been noted in some cases. The efficacy and safety of oral contraceptive products may be affected. Healthcare providers should refer to the label of the individual anti-HIV/HCV protease inhibitors for further drug-drug interaction information.

Concomitant administration of strong (e.g. ketoconazole, itraconazole, clarithromycin) or moderate (e.g. fluconazole, diltiazem, erythromycin) CYP3A4 inhibitors may increase the serum concentrations of estrogens or progestins, including etonogestrel, the active metabolite of desogestrel.

### **Pharmacodynamic interactions**

Concomitant use with the medicinal products containing ombitasvir/paritaprevir/ritonavir and dasabuvir, with or without ribavirin, or glecaprevir/pibrentasvir, or sofosbuvir/velpatasvir/voxilaprevir, may increase the risk of ALT elevations (see [CONTRAINDICATIONS, WARNINGS AND PRECAUTIONS](#)).

Therefore, LINESSA users must switch to an alternative method of contraception (e.g., progestagen-only contraception or non-hormonal methods) prior to starting therapy with these combination drug regimens. LINESSA can be restarted 2 weeks following completion of treatment with these combination drug regimens.

### **9.3 Drug-Food Interactions**

Interactions with food have not been established.

### **9.4 Drug-Herb Interactions**

Herbal products containing St. John's Wort (*hypericum perforatum*) may induce hepatic enzymes (cytochrome P450) and p-glycoprotein transporter and may reduce the effectiveness of contraceptive steroids. This may also result in breakthrough bleeding. Physicians and other health care providers should be made aware of the non-prescription products concomitantly used by the patient, including herbal and natural products.

### **9.5 Drug-Laboratory Test Interactions**

Results of laboratory tests should be interpreted in the light that the patient is on oral contraceptives. The following laboratory tests are modified.

Enzyme induction:

Enzyme induction can occur after a few days of treatment. Maximum enzyme induction is generally observed within a few weeks. After drug therapy is discontinued, enzyme induction can last for about 28 days.

When co-administered with hormonal contraceptives, many combinations of HIV protease inhibitors (e.g., nelfinavir) and non-nucleoside reverse transcriptase inhibitors (e.g., nevirapine), and/or combinations with Hepatitis C virus (HCV) medicinal products (e.g., boceprevir, telaprevir), can increase or decrease plasma concentrations of progestins, including etonogestrel, the active metabolite of desogestrel, or oestrogens. The net effect of these changes may be clinically relevant.

Women receiving any of the above mentioned hepatic enzyme-inducing medicinal or herbal products should be advised that the efficacy of LINESSA may be reduced. A barrier contraceptive method should be used in addition to LINESSA during administration of the hepatic enzyme-inducing medicinal product and for 28 days after discontinuation of the hepatic enzyme-inducing medicinal product.

For women on long-term therapy with enzyme-inducing medicinal products an alternative method of contraception unaffected by enzyme-inducing medicinal products should be considered.

Concomitant administration of strong (e.g., ketoconazole, itraconazole, clarithromycin) or moderate (e.g., fluconazole, diltiazem, erythromycin) CYP3A4 inhibitors may increase the serum concentrations of oestrogens or progestins, including etonogestrel, the active metabolite of desogestrel.

#### ***Liver function tests***

Aspartate Serum Aminotransferase (AST) - variously reported elevations. Alkaline phosphatase and gamma-glutamyl transferase (GGT) - slightly elevated.

#### ***Coagulation tests***

Minimal elevation of test values reported for such parameters as prothrombin and Factors VII, VIII, IX and X.

#### ***Thyroid function tests***

Thyroxin-binding Protein is increased as indicated by increased total serum thyroxine concentrations and decreased T<sub>3</sub> resin uptake.

#### ***Lipoproteins***

Small changes of unproven clinical significance may occur in lipoprotein cholesterol fractions.

#### ***Gonadotropins***

LH and FSH levels are suppressed by the use of oral contraceptives. Wait two weeks after discontinuing the use of oral contraceptives before measurements are made.

#### ***Glucose tolerance***

Oral glucose tolerance remained unchanged or was slightly decreased.

#### ***Tissue Specimens***

Pathologists should be advised of oral contraceptive therapy when specimens obtained from surgical procedures and Pap smears are submitted for examination.

### **9.6 Drug-Lifestyle Interactions**

Several health advantages other than contraception have been reported.

1. Combination oral contraceptives reduce the incidence of cancer of the endometrium and ovaries.
2. Oral contraceptives reduce the likelihood of developing benign breast disease and as a result decrease the incidence of breast biopsies.
3. Oral contraceptives reduce the likelihood of development of functional ovarian cysts.

4. Pill-users have less menstrual blood loss and have more regular cycles, thereby reducing the chance of developing iron-deficiency anemia.
5. The use of oral contraceptives may decrease the severity of dysmenorrhea and premenstrual syndrome and may improve acne vulgaris, hirsutism and other androgen-mediated disorders.
6. Oral contraceptives decrease the incidence of acute pelvic inflammatory disease and thereby reduce as well the incidence of ectopic pregnancy.
7. Oral contraceptives have potential beneficial effects on endometriosis.

## **10 CLINICAL PHARMACOLOGY**

### **10.1 Mechanism of Action**

Combination oral contraceptives act by the suppression of gonadotropins. The primary mechanism of action is inhibition of ovulation, but other alterations include impaired sperm penetration and “spinnbarkeit” of the cervical mucus, and changes to the endometrium to reduce the likelihood of implantation. Receptor binding studies, as well as studies in animals and humans, have shown etonogestrel, the biologically active metabolite of desogestrel, combines high progestational activity with minimal intrinsic androgenicity. DSG in combination with EE, does not counteract the estrogen-induced increase in SHBG resulting in lower serum levels of free testosterone.

### **10.2 Pharmacodynamics**

The contraceptive effect of COCs is based on the interaction of various factors, the most important of which are seen as the inhibition of ovulation and the changes in the cervical secretion.

Desogestrel, the progestogen component of LINESSA, displays low androgenic activity in relation to its progestogenic effects and may increase the HDL/LDL ratio. Like other oral contraceptives, these changes in lipid profile are associated with an increase in triglycerides.

#### **Animal and in vitro pharmacology**

Animal pharmacology and in vitro receptor binding studies indicate that etonogestrel, the biologically active metabolite of desogestrel, is a highly selective progestational agent (see Table 9) with no estrogenic effects, and only residual androgenicity.

**Table 5: Comparison Of Relative Binding Affinity Of Desogestrel, Etonogestrel And Progesterone For The Progesterone Receptor In Uterine Cytosol.\***

|              | RABBIT MYOMETRIUM | HUMAN MYOMETRIUM |
|--------------|-------------------|------------------|
| Desogestrel  | 5                 | 2                |
| Etonogestrel | 111               | 113              |
| Progesterone | 32                | 18               |

\*Binding affinities were determined at 4°C using the reference standard 16 $\alpha$ -ethyl-21-hydroxy-9-nor-pregn-4-ene-3,20-dione.

Desogestrel and its metabolites, other than etonogestrel and 3-keto-5 $\alpha$ -H-desogestrel, display minimal binding affinity for the androgen receptor with respect to dihydrotestosterone, as studied in intact MCF-7 cells. The binding affinity of both etonogestrel and 3-keto-5 $\alpha$ -H-desogestrel is approximately 1/10 of 5 $\alpha$ -dihydrotestosterone; suggesting a low androgenic activity. The binding affinity for the androgen receptor in intact MCF-7 cells as displayed by etonogestrel was also significantly lower than that of other progestogens.

The "selectivity index" (progestogen/androgen receptor binding affinity ratio) for etonogestrel in intact MCF-7 cells is higher than any other oral progestagen used in contraceptives.

Oral desogestrel displays weak androgenic activity, approximately 0.05 the activity of 17 $\alpha$ -methyl-testosterone, in orchidectomized rats, using the Herschberger test.

### Human pharmacology

After oral administration of desogestrel, typical anti-gonadotropic and progestational effects are observed; these include suppression of the hypothalamic-pituitary-gonadal axis; secretory transformation of an estrogen primed endometrium; impaired sperm penetration and "spinnbarkeit" of the cervical mucus.

### 10.3 Pharmacokinetics

**Absorption:** Desogestrel (DSG) is rapidly and almost completely absorbed and converted into etonogestrel, (ENG), its biologically active metabolite. Following oral administration, the relative bioavailability of desogestrel, based on the lowest and highest tablet strengths, 0.100 mg desogestrel/0.025 mg ethinyl estradiol and 0.150 mg desogestrel/0.025 mg ethinyl estradiol, compared to solution, as measured by serum levels of etonogestrel, is approximately 100%. Ethinyl estradiol is rapidly and almost completely absorbed. When the lowest and highest tablet strengths, 0.100 mg desogestrel/0.025 ethinyl estradiol and 0.150 mg desogestrel/0.025 mg ethinyl estradiol, were compared to solution, the relative bioavailability of ethinyl estradiol was 92% and 98% respectively. The effect of food on the bioavailability of LINESSA tablets following oral administration has not been evaluated.

The pharmacokinetics of etonogestrel and ethinyl estradiol following multiple dose administration of LINESSA tablets was determined during the third cycle in 21 subjects. After multiple dosing with LINESSA, plasma concentrations of etonogestrel reached steady-state after four days of treatment during dosing Phases 1 and 3. During dosing Phase 2, steady-state was reached after five days of treatment. The dose-normalized AUC<sub>0-24</sub> for etonogestrel was increased approximately 20% from Phase 1 to Phase 2 and approximately 10% from Phase 2 to Phase 3, indicating a possibility of time-dependent kinetics. Time dependency may be explained by a decreased clearance presumably due to increased binding of etonogestrel to sex hormone-binding globulin (SHBG). SHBG concentrations were shown to be induced by the daily administration of ethinyl estradiol. Steady state for ethinyl estradiol was reached after four days of dosing in all dosing phases. The pharmacokinetic parameters of etonogestrel and ethinyl estradiol during the third cycle following multiple dose administration of LINESSA tablets are summarized in Table 6.

**Table 6: Mean (SD) Pharmacokinetic Parameters of LINESSA Over a 28-Day Dosing Period in the Third Cycle (n=21)**

| <b>Etonogestrel</b>      |                      |                              |                        |                                   |                          |
|--------------------------|----------------------|------------------------------|------------------------|-----------------------------------|--------------------------|
| Phase (days)             | Dose <sup>H</sup> mg | C <sub>max</sub> pg/mL       | t <sub>max</sub> hr    | n-AUC <sub>0-24</sub> pgxhr/mL/mg | CL/F L/hr                |
| 1(1-7)                   | 0.1                  | 2163.3 (856.4)               | 1.6 (0.7)              | 196.0 (75.4)                      | 6.1 (2.3)                |
| 2(8-14)                  | 0.125                | 3241.5 (1296.5) <sup>a</sup> | 1.1 (0.3) <sup>a</sup> | 234.4 (85.0) <sup>a</sup>         | 5.1 (1.9) <sup>a</sup>   |
| 3(15-21)                 | 0.15                 | 3855.7 (1273.1)              | 1.5 (0.8)              | 256.6 (104.0)                     | 4.6 (1.6)                |
| <b>Ethinyl Estradiol</b> |                      |                              |                        |                                   |                          |
| 1(1-7)                   | 0.025                | 85.4 (51.7)                  | 1.5 (0.8)              | 26.4 (11.5)                       | 43.5 (15.0)              |
| 2(8-14)                  | 0.025                | 91.3 (52.2) <sup>a</sup>     | 1.2 (1.2) <sup>a</sup> | 29.0 (15.5) <sup>a</sup>          | 41.7 (15.5) <sup>a</sup> |
| 3(15-21)                 | 0.025                | 90.1 (48.2)                  | 1.2 (0.7)              | 28.3 (13.2)                       | 42.5 (18.7)              |

H= Desogestrel

a= n=20

C<sub>max</sub>= maximum serum drug concentration

t<sub>max</sub>= time at which maximum serum drug concentration occurs

n-AUC<sub>0-24</sub>= area under the concentration-vs time curve -0 to 24 hours normalized to 1 mcg administered

CL/F- apparent clearance

Note: for information on t<sub>1/2</sub> for Day 21, see the [Excretion](#).

**Distribution:** Etonogestrel, the active metabolite of desogestrel, was found to be 98% protein bound, primarily to sex hormone-binding globulin (SHBG). Ethinyl estradiol is primarily bound to plasma albumin. Ethinyl estradiol does not bind to SHBG but induces SHBG synthesis. Desogestrel, in combination with ethinyl estradiol, does not counteract the estrogen-induced increase in SHBG, resulting in lower serum levels of free testosterone.

**Metabolism: Desogestrel:** Desogestrel is rapidly and completely metabolized by hydroxylation in the intestinal mucosa and on first pass through the liver to etonogestrel. *In vitro* data suggest an important role for the cytochrome P450 CYP2C9 in the bioactivation of desogestrel. Further metabolism of etonogestrel into 6 $\beta$ -hydroxy, etonogestrel and 6 $\beta$ -13ethyl-dihydroxylated as major metabolites is catalyzed by CYP3A4. Other metabolites (i.e. 3 $\alpha$ -OH-desogestrel, 3 $\beta$ -OH-desogestrel, and 3 $\alpha$ -OH-5 $\alpha$ -H-desogestrel) also have been identified and these metabolites may undergo glucuronide and sulfate conjugation.

**Ethinyl estradiol:** Ethinyl estradiol is subject to a significant degree of presystemic conjugation (phase II metabolism). Ethinyl estradiol, escaping gut wall conjugation, undergoes phase I metabolism and hepatic conjugation (phase II metabolism). Major phase I metabolites are 2-OH-ethinyl estradiol and 2-methoxy-ethinyl estradiol. Sulfate and glucuronide conjugates of both ethinyl estradiol and phase I metabolites, which are excreted in bile, can undergo enterohepatic circulation.

**Excretion:** Etonogestrel and ethinyl estradiol are primarily eliminated in urine, bile and feces. At steady state, on Day 21, the elimination half-lives of etonogestrel and ethinyl estradiol are 37.1 $\pm$ 14.8 hours and 28.2 $\pm$ 10.5 hours, respectively.

### Special Populations and Conditions

**Ethnic origin:** There is no information to determine the effect of race on the pharmacokinetics of LINESSA tablets.

**Hepatic Insufficiency:** No formal studies were conducted to evaluate the effect of hepatic disease on the disposition of LINESSA. However, steroid hormones may be poorly metabolized in patients with impaired liver function (see [7 WARNINGS & PRECAUTIONS – Hepatic/Biliary/Pancreatic](#)).

**Renal Insufficiency:** No formal studies were conducted to evaluate the effect of renal disease on the disposition of LINESSA.

## 11 STORAGE, STABILITY AND DISPOSAL

Store between 15-30°C.

Keep in a safe place out of the reach of children and pets.

## **12 SPECIAL HANDLING INSTRUCTIONS**

Any unused portion or waste material should be disposed of in accordance with local requirements.



## PART II: SCIENTIFIC INFORMATION

### 13 PHARMACEUTICAL INFORMATION

#### Drug Substance


##### I. Progestogen

Proper name: Desogestrel

Chemical name: 17 ( $\alpha$ ) -13- ethyl-11-methylene 18,19-dinor-pregn-4-en-20-yn-17-ol

Molecular formula and molecular mass:  $C_{22}H_{30}O$  and 310.48 g/mol

Structural formula:



Physical Form: White, crystalline powder

Solubility: Solubility at 20°C: n-Hexane: 40 mg/mL  
Ethanol (96%): > 200 mg/mL  
Ethyl acetate: > 150 mg/mL  
Water: Practically insoluble

Melting Point: 111 - 113 °C


## II. Estrogen

Proper Name: Ethinyl Estradiol

Chemical Name: 19-nor-17 $\alpha$ -pregna-1,3,5(10)-trien-20-yne-3,17-diol

Molecular Formula and molecular mass: C<sub>20</sub>H<sub>24</sub>O<sub>2</sub> and 296.4 g/mol

Structural Formula:



Physical Form: White, crystalline powder

Solubility:

- Ethanol: Approximately 170 mg/mL
- Acetone: Approximately 200 mg/mL
- Chloroform: Approximately 50 mg/mL
- Dioxane: Approximately 250 mg/mL
- Diethyl ether: Approximately 250 mg/mL
- Water: Practically insoluble.

Melting Point: 180-186°C

### **14 CLINICAL TRIALS**

Data is provided from 2,768 subjects treated with LINESSA who provided 14,526.8 cycles of exposure to LINESSA, including 2,168 women who completed six cycles of exposure. The combined results of the two pivotal studies provide data to determine the Pearl Index and Life Table estimate values.

#### **a) Pearl Index**

The observed Pearl Index among LINESSA users compares favourably to what has been reported for other low-dose oral contraceptives. Twelve of the 2,752 subjects using

LINESSA became pregnant. The Pearl Index for total pregnancies was 1.08 per 100 women-years calculated from 14,456 cycles and included 2,643 women.

**b) Life Table Estimates**

The six-cycle cumulative life-table pregnancy rate is estimated as 0.0051 women-years.

**c) Cycle Control**

Table 6 presents the incidence of intermenstrual breakthrough bleeding (IMB), early withdrawal bleeding (EWB) and absence of withdrawal bleeding (AWB), by treatment group for all subjects in the Cycle Control Analysis Group. Breakthrough spotting occurred more frequently than breakthrough bleeding.

Table 7 summarizes the duration of withdrawal bleeding, which included early withdrawal bleeding and continued withdrawal bleeding, if any, by cycle and treatment group. The overall mean length of withdrawal bleeding, defined as “any bleeding-spotting episode that began during or continued into the active tablet-free interval” was 5.1 days for the LINESSA Group and 4.8 days for the Triphasic Norethindrone/Ethinyl Estradiol (Net/EE) group.

**Table 7: Incidence of Intermenstrual Bleeding, Early Withdrawal Bleeding and Absence of Withdrawal Bleeding by Cycle and Treatment Group**

| Total  |                | IMB            |      | EWB            |     | AWB            |     |
|--|----------------|----------------|------|----------------|-----|----------------|-----|
| Cycle  | N <sup>a</sup> | n <sup>b</sup> | %    | n <sup>b</sup> | %   | n <sup>b</sup> | %   |
| LINESSA  |                |                |      |                |     |                |     |
| 1  | 2475           | 333            | 13.5 | 184            | 7.4 | 82             | 3.3 |
| 2  | 2401           | 275            | 11.5 | 156            | 6.2 | 70             | 2.9 |
| 3  | 2319           | 258            | 11.1 | 127            | 5.5 | 56             | 2.4 |
| 4  | 2257           | 210            | 9.3  | 108            | 4.8 | 52             | 2.3 |
| 5  | 2216           | 209            | 9.4  | 108            | 4.9 | 70             | 3.2 |
| 6  | 2093           | 234            | 11.2 | 108            | 5.2 | 67             | 3.2 |
| Total  | 13761          | 1519           | 11   | 785            | 5.7 | 317            | 2.9 |
| Triphasic Norethindrone/Ethinyl Estradiol (Net/EE) |                |                |      |                |     |                |     |
| 1  | 2525           | 567            | 22.5 | 210            | 8.3 | 133            | 5.3 |
| 2  | 2450           | 397            | 16.2 | 183            | 7.5 | 121            | 4.9 |
| 3  | 2358           | 361            | 15.3 | 149            | 6.4 | 127            | 5.4 |
| 4  | 2291           | 277            | 12.1 | 127            | 5.5 | 110            | 4.8 |
| 5  | 2228           | 253            | 11.4 | 103            | 4.6 | 114            | 5.1 |
| 6  | 2114           | 303            | 14.3 | 128            | 6.1 | 113            | 5.3 |
| Total  | 13966          | 2158           | 15.5 | 900            | 6.4 | 718            | 5.1 |

a Number of subjects with a valid cycle

b Number of subjects with the event within each cycle

Notes: Absence of withdrawal bleeding is defined as no bleeding-spotting episode during the placebo tablet period. Early withdrawal bleeding (EWB) is defined as that portion of the withdrawal bleeding that occurred before the intake of placebo tablets.

Intermenstrual bleeding (IMB) is defined as any bleeding-spotting event that occurred during the active tablet period that was neither part of an early nor continued withdrawal bleeding. IMB consists of breakthrough bleeding and breakthrough spotting.

**Table 8: Duration of Withdrawal Bleeding by Treatment Group and Cycle**

| Total   |      |     |        |
|---------|------|-----|--------|
| n       | Mean | SD  | Median |
| LINESSA |      |     |        |
| 2393    | 5.4  | 2.8 | 5      |
| 2331    | 5.2  | 2.3 | 5      |
| 2263    | 5.2  | 2.2 | 5      |
| 2205    | 5.1  | 2.1 | 5      |
| 2146    | 5.1  | 2   | 5      |
| 2026    | 4.4  | 2   | 4      |
| 13364   | 5.1  | 2.3 | 5      |
| NET/EE  |      |     |        |
| 2392    | 5.1  | 2.4 | 5      |
| 2329    | 4.9  | 2   | 5      |
| 2231    | 4.9  | 2   | 5      |
| 2181    | 4.8  | 1.8 | 5      |
| 2114    | 4.7  | 1.8 | 5      |
| 2001    | 4.6  | 1.7 | 5      |
| 13248   | 4.8  | 2   | 5      |

The results indicate that LINESSA cycle control is generally excellent which was also reflected in the low number of dropouts due to irregular bleeding or absence of withdrawal bleeding. These results are very similar to those obtained with other oral contraceptives.

Based on these results LINESSA demonstrated comparable cycle control to another triphasic preparation using a 7/7/7 regimen (Net/EE).

**Tolerance**

Among 2,768 LINESSA subjects in the two pivotal studies 124 (4.4%) discontinued the study due to drug-related adverse experiences and 161 (5.8%) discontinued due to any adverse experience.

The most common system-organ class for which adverse experiences resulting in discontinuation were reported was the female reproductive (1.8%).

**Table 9: Overall Assessment of Adverse Experiences by Treatment Period and Treatment Group-Pivotal Clinical Studies (All Subjects Treated Group)**

|   | LINESSA<br>(n=2768) |       | (Net/EE)<br>(N=2784) |       |
|---|---------------------|-------|----------------------|-------|
|   | n                   | %     | n                    | %     |
| <b>Adverse Experiences that Occurred Prior to Start of Study Drug</b>     |                     |       |                      |       |
| Total of all subjects treated group                                       | 2768                | 100.1 | 2784                 | 100.0 |
| Subjects with an AE   | 126                 | 4.6   | 105                  | 3.8   |
| Subjects with a serious AE  | 4                   | 0.1   | 2                    | 0.1   |
| Subjects with AE as a cause for discontinuation                           | 0                   | 0.0   | 0                    | 0.0   |
| Subjects with a drug-related AE <sup>a</sup>                              | N/A                 | N/A   | N/A                  | N/A   |
| Subjects with a severe AE   | 17                  | 0.6   | 13                   | 0.5   |
| <b>Adverse Experiences that Occurred During the Treatment Period</b>      |                     |       |                      |       |
| Total of all subjects treated group                                       | 2768                | 100.0 | 2784                 | 100.0 |
| Subjects with an AE   | 1891                | 68.3  | 1830                 | 65.7  |
| Subjects with a serious AE  | 32                  | 1.2   | 34                   | 1.2   |
| Subjects with AE as a cause for discontinuation                           | 159                 | 5.7   | 150                  | 5.4   |
| Subjects with a drug-related AE <sup>a</sup>                              | 936                 | 33.8  | 903                  | 32.4  |
| Subjects with a severe AE   | 329                 | 11.9  | 302                  | 10.8  |
| <b>Adverse Experiences that Occurred During the Post-Treatment Period</b> |                     |       |                      |       |
| Total of all subjects treated group                                       | 2768                | 100.0 | 2784                 | 100.0 |
| Subjects with an AE   | 353                 | 12.8  | 322                  | 11.6  |
| Subjects with a serious AE  | 3                   | 0.1   | 10                   | 0.4   |
| Subjects with AE as a cause for discontinuation                           | 2                   | 0.1   | 0                    | 0.0   |
| Subjects with a drug-related AE <sup>a</sup>                              | 56                  | 2.0   | 38                   | 1.4   |
| Subjects with a severe AE   | 38                  | 1.4   | 39                   | 1.4   |

<sup>a</sup> Adverse experiences classified as related were judged as possibly, probably or definitely related.

The incidence of discontinuations due to intermenstrual bleeding was 0.8% for LINESSA and 0.7% for (Net/EE).

### **Vital Signs and Weight Gain**

In the two controlled clinical studies, mean systolic and diastolic blood pressure remained relatively stable; there were no clinically significant differences at any time point between LINESSA and Net/EE. Clinically significant high pulse rates and low respiratory rates were incidental in both treatment groups. There were minimal changes in body weight and Body Mass Index (BMI) over the course of the controlled clinical studies. Mean changes in body weight ranged between a mean loss of 0.1 kg to a mean gain of 0.4 kg in the LINESSA group while in the Net/EE group, mean changes in body weight ranged between a mean loss of 0.2 kg and a mean gain of 0.4 kg. An increase in body weight was reported by 2.3% of subjects in the LINESSA group and 1.8% of subjects in the Net/EE group. There was a slight decrease in the mean BMI from baseline to last measurement for subjects in the LINESSA Group (-0.1 kg/m<sup>2</sup>) and no change for subjects receiving Net/EE.

A total of 20 (0.7%) LINESSA - treated subjects and 10 (0.4%) subjects who received Net/EE were reported to have mild to moderate hypertension, while one subject in the LINESSA group showed severe hypertension. Eight subjects (0.8%) in the LINESSA group and one subject (<0.10%) in the Net/EE group discontinued due to hypertension, each of these was considered to be drug-related by the investigator. These figures should be considered in light of the fact that mild to moderate hypertension is a common condition and its prevalence is 7-13 % among women aged 20-44 years, and 20 to 50% in women aged 30-65 years.

### **Lipid Metabolism**

The analysis of lipids included data from 2026 subjects who received LINESSA. The observed changes in total cholesterol and triglycerides were mostly within the normal range. Results of the six-cycles exposure showed statistically significant mean percent increases in triglycerides in subjects who received LINESSA (29.5%) when compared to Net/EE (25.0%). The weighted mean difference between LINESSA and Net/EE was 5.37 mg/dL which represented approximately 5% of the starting values of plasma triglyceride or 3% of the normal range. Potentially clinically significant increases in cholesterol and triglycerides were seen in 0.3% and 0.2% of LINESSA subjects, respectively, and 0.2% and 0% of NET/EE subjects, respectively.

Other metabolic parameters were minimally affected (i.e. carbohydrate metabolism).

### **Clinical Laboratory Results**

As a class, oral contraceptives are known to be associated with decreased glucose tolerance (in pre-diabetic and diabetic women).

Pooled data from two controlled clinical studies (over 2,000 women, 6 cycles) as well as the results of two smaller clinical pharmacology studies indicate that LINESSA has no observable adverse effects on fasting serum glucose.

In addition, there were no observable adverse effects of LINESSA on hepatic or renal parameters, red or white blood cell indices or urinalysis tests in the two controlled clinical studies, or in the smaller phase II and clinical pharmacology studies. The incidences of adverse events related to abnormalities of these analytes during the two major trials were, in general, low and similar between the two treatment groups. Data from the other studies support these conclusions.

## **15 MICROBIOLOGY**

No microbiological information is required for this drug product.

## **16 NON-CLINICAL TOXICOLOGY**

### **General Toxicology**

#### ***Acute Toxicity Studies***

Acute single-dose studies were conducted in both rats and mice, with desogestrel + ethinyl estradiol and desogestrel alone, to determine the upper limits of tolerance and to assess specific signs of toxicity. Both compounds were dosed orally by gavage or intraperitoneal as aqueous suspensions. The oral dosage level of 2000 mg/kg was about  $6 \times 10^5$  times the projected human clinical dose. The intraperitoneal dosage was 500 mg/kg. Groups of 10 males and 10 females were tested with desogestrel + ethinyl estradiol and groups of 6 males and 6 females with desogestrel alone. The animals were observed for 7 days and then necropsied.

None of the test animals died during the oral or intraperitoneal studies. The oral dosed mice and rats had temporary signs of reduced activity, some motor incoordination, diminished food consumption, and other nonspecific signs related to the large dose of the test material. Likewise, mice and rats dosed intraperitoneal showed similar signs. Some evidence of serositis (localized peritoneal irritation) was associated with the test substances.

These data are consistent with published information on other contraceptive steroids which indicate that steroids in general have a low level of toxicity in single-dose acute animal studies.

#### ***Multidose Toxicity Studies***

The objective of the multidose toxicity studies was to determine whether the chronic oral administration of either desogestrel + ethinyl estradiol or desogestrel alone to mice, rats, dogs, and monkeys would induce either reversible or irreversible systemic adverse effects or cause the development of benign or malignant neoplasms. Desogestrel + ethinyl estradiol, in a ratio of 2.5:1, was employed in most multidose toxicity and multidose tumorigenicity toxicity studies and in a ratio of 5:1 in 52-, 104-week and 3-year studies in dogs and monkeys. The test compounds were administered orally by gavage to mice and rats, orally by tablet or capsule to dogs, and orally by soft drink or by intubation to monkeys.

The protocol for each of these studies was typical of that used for multidose toxicity tests in general. The doses were multiples of the human dose and generally calculated to be 2, 20, and 200 times the expected human usage levels in most multidose and tumorigenicity studies in mouse, rat and dog. In shorter studies, the duration of treatment was 26 or 52 weeks with a 4 to 13-week recovery period incorporated into the study design. In the 52-, 104-week and 3-year dog and monkey studies dose levels were 1, 10, 25 and 2, 10, and 50 times the human dose respectively.

The following table lists the study duration, species tested, and the test compounds:



**Table 10: Multidose Toxicity Studies**

| Multidose Toxicity Studies |          |           |   |       |
|----------------------------|----------|-----------|---|-------|
| Duration                   | Species  | Drugs     | Dose(mg/kg)   | n     |
| 52 weeks                   | rat, dog | DSG + EE* | 0.005+0.002 <sup>a</sup><br>0.05 +0.02<br>0.5 +0.2      | 70,14 |
|                            | dog      | DSG + EE  | 0.003+0.0006 <sup>b</sup><br>0.03 +0.006<br>0.075+0.015 | 20    |
|                            | monkey   | DSG + EE  | 0.006+0.0012 <sup>c</sup><br>0.03 +0.006<br>0.15 +0.03  | 20    |
| 80 weeks                   | mouse    | DSG + EE  | see <sup>a</sup>  | 112   |
| 104 weeks                  | rat      | DSG + EE  | see <sup>a</sup>  | 110   |
|                            | dog      | DSG + EE  | see <sup>b</sup>  | 20    |
|                            | monkey   | DSG + EE  | see <sup>c</sup>  | 20    |
| 3 years                    | dog      | DSG + EE  | see <sup>b</sup>  | 20    |
|                            | monkey   | DSG + EE  | see <sup>c</sup>  | 20    |
| 26 weeks                   | rat, dog | DSG       | 0.00625<br>0.0625<br>0.625                              | 64,14 |
| 52 weeks                   | rat, dog | DSG       | 0.005 <sup>d</sup><br>0.05<br>0.5                       | 60,12 |
| 81 weeks                   | mouse    | DSG       | see <sup>d</sup>  | 112   |
| 104 weeks                  | rat      | DSG       | see <sup>d</sup>  | 110   |

\*DSG = desogestrel, EE = ethinyl estradiol

The 52-week study with desogestrel + ethinyl estradiol in rats revealed no direct treatment-associated effect on mortality. Clinical signs of treatment included alopecia and reduction of testicle size, primarily in high-dose animals, which were reversible on treatment cessation. Depressed weight gain and/or decreased food consumption was present in both sexes of the intermediate- and high-dose animals. There was an alteration in APTT, Hb, and PCV noted along with lowered neutrophil and lymphocyte counts. These changes are known to occur in these type of studies and were found to be reversible upon treatment cessation. No unusual changes were found in blood chemistry or urinalysis. Dose-related lower protein content of the urine in males may be attributed to the atrophic change in secondary sex organs.

Organ weight changes were consistent with those noted with other combination oral contraceptives. The liver weight was increased at 26 and 52 weeks in primarily intermediate-dose (ID) and high-dose animals; testes, epididymides, prostate, seminal vesicles, ovaries, uterus, adrenals, and the pituitary gland were also affected by treatment.

Microscopic tissue changes included the following: Hepatocytic vacuolation and occasional foci of hepatocellular hyperplasia, especially in high-dose animals; a dose-related increase in yellowish pigment in the kidney cortical tubule epithelium, and increased mineralized concretions in high-dose males; atrophy of the testes, epididymides, prostate, and seminal vesicles; reduction or absence of corpora lutea in the ovaries; hyalinization or endometrial hyperplasia of the uterus; increased keratinization of the vagina in high-dose females; hypertrophy and hyperplasia of the adrenal cortex with sinusoidal telangiectasis; and hypertrophy/hyperplasia of the anterior lobe of the pituitary, especially at 52 weeks in high-dose animals.

The 8-week withdrawal period used in this study resulted in a partial reversal of the prior changes. All would have probably reverted to normal with a longer recovery period. There was an increased incidence of benign mammary neoplasms in all drug-treated groups.

The 52-week dog study was conducted with oral dosed desogestrel + ethinyl estradiol tablets in a ratio of 2.5:1. Three high-dose mortalities occurred during the study. Two females died and the other was killed in extremis. The cause of death or morbidity was peritonitis, secondary to perforating pyometra. Clinical signs included typical skin thickening and folding with alopecia, interruption of the estrous cycle with swelling of external genitalia in females, vaginal discharge in high-dose females, pendulous penile sheath in males with reduction in testicle size, enlarged and/or secretory mammary tissue in females, and 2 transient (1 intermediate-dose) and 1 transient and 1 persistent nodule (1-high-dose) of the mammary gland. The persistent nodule was an area of hyperplasia.

Changes in certain hematological, coagulation, blood chemistry and urinalysis parameters were neither unusual nor unexpected for this type of compound. Changes either in weight or histomorphological characteristics were noted in the primary and secondary sex organs and liver, primarily in high-dose animals. All were associated with the hormonal attributes of the drug.

The multidose toxicity study in the monkey was performed at a 5:1 ratio of desogestrel to ethinyl estradiol with dosing for 21 days followed by a 7-day drug-free period. The 12-month data revealed no unexpected clinical, clinicopathological, or histomorphological findings. Typical hormonal dose-related changes occurred, such as decreased corpora lutea, secretory mammary glands, increased endocervical mucus, decreased thickness of the endometrium with secretory changes, a dose-related decrease in the thickness of the vaginal epithelium and increased pituitary weight.

The multidose studies in rats and dogs with desogestrel alone resulted in fewer alterations in the primary and secondary sex organs and other peripheral hormonally sensitive tissues.

In rats, the absence of ethinyl estradiol in the test compound resulted in expected progestational changes at 26 and 52 weeks, such as secretory changes in the uterine endometrium, mucification of the vaginal epithelium, mild glandular hyperplasia of the mammary glands, and reduced pituitary weights. In the 52-week portion of the study, a small number of benign or malignant neoplasms were observed, but none of these were causally related to the test compound.

The toxicity of multidoses of desogestrel alone in dogs resulted in no unusual or unexpected changes at 26 weeks. The liver weight in high-dose animals was increased but this was due primarily to the progestogenic effect of increased glycogen storage. The uterus was increased in both size and weight due to hormonal stimulation of the endometrium and the ovaries had a lack of mature follicles and an absence of corpora lutea. The prostate weight was slightly reduced in high-dose males. Lobular development of the mammary glands was increased in intermediate and high-dose females.

The 52-week segment of the dog study with desogestrel alone resulted in changes similar to those seen at 26 weeks; however, occasional small mammary nodules (5 mm or less) were present in 1 control (C), 1 low-dose (LD), 1 ID, and 4 high-dose animals. They disappeared in the 1 C and 2 high-dose animals. The remaining nodules were found to be nonneoplastic and proved to be either smaller superficial lymph nodes or dilated ducts. The uterine stimulation was increased at 52 weeks but did not result in the death of any animal.

Four multidose toxicity studies of up to 2 years in duration were conducted in rats, dogs, and monkeys. Desogestrel + ethinyl estradiol was studied in rats, monkeys, and dogs, and desogestrel alone was studied in rats.

In rats, there was no evidence of a neoplastic response when desogestrel was administered alone, however, increased evidence of benign mammary neoplasms were evident in all desogestrel + ethinyl estradiol-treated groups. Other clinical, clinicopathological, and histopathological changes were attributable to the hormonal influences of either desogestrel or its combination with ethinyl estradiol.

The 2-year dog study utilized a 5:1 desogestrel + ethinyl estradiol ratio. The test compound was dosed at 1, 10 and 25 times the human dosage levels for 21 days with a 7-day drug-free period. There was evidence of the following: suppression of the estrous cycle in intermediate- and high-dose animals, an increased incidence of mammary gland development and secretory activity similar to those observed in the normal metestrous phase of the cycle; decreased AP in high-dose dogs, and a single focus of ductal epithelial hyperplasia in 1 low-dose dog. No tumorigenic effect was present.

The 2-year study of desogestrel + ethinyl estradiol in monkeys caused the expected pattern of hormonally-mediated changes. Menstrual and ovarian activities were reduced in high-dose animals. Secretory activity of the mammary glands was increased in a dose-related manner in intermediate- and high-dose animals. Other hormonally-associated changes included: an increased fibrinogen and APTT; decreased PPT; reduced AP; increased triglycerides and cholesterol levels; and lowered albumen in intermediate- and high-dose animals; endometrium which was either stimulated (ID and HD) or lacked activity (some high-dose animals); and increased acidophils and decreased basophils in the pituitary in intermediate- and high-dose animals. All of these findings are consistent with contraceptive steroid effects in the monkey.

Multidose tumorigenicity studies were conducted in the mouse (80-81 months) and rat (2 years) with either desogestrel + ethinyl estradiol or desogestrel alone, respectively. Desogestrel + ethinyl estradiol in mice resulted in a higher mortality rate; this was primarily due to the increased incidence of pituitary tumors in treated mice, especially high-dose animals. Other nonneoplastic alterations occurred, but were within expected limits for a compound of this type. Desogestrel alone in mice did not remarkably affect the mortality rate and had no influence on tumorigenicity.

Desogestrel + ethinyl estradiol in the rat resulted in slightly increased mortality at the high-dose level and contributed to a dose-dependent increase in the number of pituitary and mammary neoplasms; this increase was largely attributable to the ethinyl estradiol component.

Desogestrel alone in the rat had no influence on mortality and possibly was responsible for a slight lowering effect. Incidences of mammary and pituitary tumors were slightly lessened at the high-dose level. This is in contrast to the 104-week rat study with desogestrel + ethinyl estradiol, where the differences noted were considered to have been attributable to the ethinyl estradiol component.

Three-year studies were conducted in both Beagle dogs and Rhesus monkeys with desogestrel + ethinyl estradiol with a 1- and 2-year interim sacrifice in monkey and a 2-year interim sacrifice in dogs. No tumorigenic response was noted. Mammary glands of dogs had lobulo-alveolar development with limited secretory change, an expected hormonal effect. Other tissue changes as described under the 2-year interim report, limited to the primary and secondary sex organs, were associated with the hormonal activities of the combination OC. The monkey study conducted for 3 years, with a 1- and 2- year interim sacrifice, revealed no evidence of a tumorigenic effect. The changes observed, as described at the 2-year interim studies, were typical of the hormonal activities of the combination OC and included effects on the menstrual cycle, cervical mucus and endometrial morphology.

### **Mutagenicity Studies**

The Ames test and the rat Micronucleus test were conducted on desogestrel, either alone or in combination with ethinyl estradiol. Both assays demonstrated that neither desogestrel alone nor in combination with ethinyl estradiol exert any mutagenic effect.

### **Reproductive and Developmental Toxicology**

Nonclinical reproductive toxicity studies included 11 studies conducted in rats and 2 studies conducted in rabbits. Desogestrel, both alone and in combination with ethinyl estradiol, was tested. These studies were conducted to assess what effect, if any, the test substance might have on the reproductive process, including; fertility and reproductive performance, teratogenicity and embryotoxicity, and perinatal and postnatal effects in the offspring.

Four segment I reproductive toxicity studies were conducted in rats; 1 study with desogestrel + ethinyl estradiol and 3 studies with desogestrel alone. The desogestrel + ethinyl estradiol study, conducted using doses of 0.5 mg desogestrel + 0.2 mg ethinyl estradiol/kg/day, demonstrated that the test compound had no adverse effect on mating and pregnancy performance in F<sub>0</sub> females or on the number, anatomical features, development and fertility of the offspring.

Desogestrel alone was studied in both Sprague Dawley and CFY rats. An additional study in Sprague Dawley rats was conducted after microphthalmia was increased in CFY offspring of the desogestrel -treated dams. No increase in microphthalmia was seen in the second Sprague Dawley study. The defect was thus thought to be strain-related. In all 3 studies the contraceptive effect of desogestrel was reversible. Treatment at contraceptive and subcontraceptive dose levels did not cause any serious after-effects on the dams or their offspring.

A fertility and embryotoxicity study with desogestrel + ethinyl estradiol at levels causing complete infertility, slight infertility, and no infertility, were conducted in rats. Uninterrupted daily administration of desogestrel + ethinyl estradiol, at subcontraceptive doses before and during pregnancy, reduced the number of offspring but had no effect on the quality of the F<sub>1</sub> generation.

Segment II embryotoxicity studies following the classical design with dosage exclusively during pregnancy and organogenesis were performed in both the rat and rabbit. A total of 5 embryotoxicity studies were conducted; 3 studies with desogestrel alone and 2 studies with desogestrel + ethinyl estradiol.

Desogestrel + ethinyl estradiol tested at high-dose levels in rats and rabbits caused maternal toxicity and embryoletality, but at lower doses had no untoward reaction in the dams and no detectable effect on the course of pregnancy, embryonic mortality, or fetal morphology.

Desogestrel alone was tested in both Sprague Dawley and CFY rats and in rabbits. High dosages of desogestrel caused maternal toxicity (2-8 mg/kg) in rats, while doses of 2 to 4 mg/kg caused abortion in rabbits. Lower dosages in rats and rabbits caused no discernible effect on the course of pregnancy, embryonic mortality, or on fetal morphology.

The effects of desogestrel alone, when dosed during late pregnancy, were assessed in rats. Dose levels up to 4 mg/kg/day from days 14-20 of pregnancy caused neither masculinization of female fetuses nor feminization of male fetuses.

Segment III studies, to evaluate the possible effects on peri- and postnatal development due to transfer of drug through the milk, were conducted with desogestrel, either alone or in combination with ethinyl estradiol. Desogestrel + ethinyl estradiol caused reduced food consumption in intermediate and high-dose dams. Retarded pup growth persisted until weaning in the high-dose group, but there was no effect on the pre- or post-weaning physical development. Fertility of the F<sub>1</sub> offspring was not affected. Desogestrel alone had no effect on the treated dams, weight gain in the pups, or physical development of the pups. Fertility of the F<sub>1</sub> treated animals was comparable to that of the F<sub>1</sub> control females.

## PATIENT MEDICATION INFORMATION

### READ THIS FOR SAFE AND EFFECTIVE USE OF YOUR MEDICINE

#### **LINESSA 21 and LINESSA 28 desogestrel and ethinyl estradiol tablets**

Read this carefully before you start taking **LINESSA** and each time you get a refill. This leaflet is a summary and will not tell you everything about this drug. Talk to your healthcare professional about your medical condition and treatment and ask if there is any new information about **LINESSA**.

#### **Serious Warnings and Precautions**

- Cigarette smoking increases the risk of serious adverse effects on the heart and blood vessels. This risk increases with age particularly in women older than 35 years of age, who use hormonal birth control. The risk also increases with the number of cigarettes smoked. For this reason, women who smoke and are over 35 years of age should not use LINESSA.
- Birth control pills DO NOT PROTECT against sexually transmitted infections (STIs), including HIV/AIDS. For protection against STIs, you must use latex or polyurethane condoms AND take your birth control pills.

#### **What is LINESSA used for?**

LINESSA is used to prevent pregnancy in women 18 to 65 years old.

#### **How does LINESSA work?**

LINESSA is a birth control pill. It is considered to be a combination oral contraceptive. This is because it contains two female sex hormones: desogestrel and ethinyl estradiol. Combination birth control pills work in two ways:

- They stop the monthly release of an egg by the ovaries.
- They change the mucus produced by the cervix. This slows the movement of the sperm through the mucus and into the uterus (womb).

#### Effectiveness of Birth Control Pills

Combination birth control pills are more than 99 percent effective in preventing pregnancy when:

- the pill is **taken as directed**, and
- the amount of estrogen is 20 micrograms or more.

The chance of becoming pregnant increases with incorrect use.

### Other Ways to Prevent Pregnancy

Other methods of birth control are available to you. They are usually less effective than birth control pills. When used properly, however, other methods of birth control are effective enough for many women.

The following table gives reported pregnancy rates for various forms of birth control, including no birth control. The reported rates represent the number of women out of 100 who would become pregnant in one year.

#### **Reported Pregnancies per 100 Women per Year:**

|   |                  |
|---|------------------|
| Combination pill                        | less than 1 to 2 |
| Intrauterine device (IUD)               | less than 1 to 6 |
| Condom with spermicidal foam or gel     | 1 to 6           |
| Mini-pill                               | 3 to 6           |
| Condom                                  | 2 to 12          |
| Diaphragm with spermicidal foam or gel  | 3 to 18          |
| Spermicide                              | 3 to 21          |
| Sponge with spermicide                  | 3 to 28          |
| Cervical cap with spermicide            | 5 to 18          |
| Periodic abstinence (rhythm), all types | 2 to 20          |
| No birth control                        | 60 to 85         |

Pregnancy rates vary widely because people differ in how carefully and regularly they use each method. This does not apply to IUDs since they are implanted in the uterus. Regular users may achieve pregnancy rates in the lower ranges. Others may expect pregnancy rates more in the middle ranges.

#### **What are the ingredients in LINESSA?**

Medicinal ingredients: desogestrel and ethinyl estradiol

Non-medicinal ingredients:

LINESSA 21 and LINESSA 28:

- Light yellow tablets: hydroxypropyl, lactose monohydrate, methylcellulose, polyethylene glycol, starch, stearic acid, titanium dioxide, talc, vitamin E and yellow ferric oxide.
- Orange tablets: hydroxypropyl, lactose monohydrate, methylcellulose, polyethylene glycol, red ferric oxide, starch, stearic acid, titanium dioxide, talc, vitamin E and yellow ferric oxide.
- Red tablets: hydroxypropyl, lactose monohydrate, methylcellulose, polyethylene glycol, red ferric oxide, starch, stearic acid, titanium dioxide, talc and vitamin E.

LINESSA 28:

- Green tablets: corn starch, FD&C Blue No.2 aluminum lake, hydroxypropyl methylcellulose, lactose monohydrate, magnesium stearate, polyethylene glycol, talc, titanium dioxide and yellow ferric oxide.



**LINESSA comes in the following dosage forms:**

LINESSA 21 (21-Day Pack) and LINESSA 28 (28-Day Pack): Each sachet contains a blister card with 21 tablets for oral administration. The 21 tablets are divided into three different dosing phases;

- Seven (7) light yellow tablets: 0.100 mg desogestrel and 0.025 mg ethinyl estradiol
- Seven (7) orange tablets: 0.125 mg desogestrel and 0.025 mg ethinyl estradiol
- Seven (7) red tablets: 0.150 mg desogestrel and 0.025 mg ethinyl estradiol.

LINESSA 28 (28-Day Pack): The blister pack contains the same three dosing phases as LINESSA 21 with an additional seven (7) green “reminder” tablets that do not contain hormones.

**Do not use LINESSA if:**

- You are allergic to any ingredients in LINESSA or component of the packaging;
- You have or have had a blood clot in the legs (deep vein thrombosis), lung (pulmonary embolism), or somewhere else in your body;
- You have blood vessel disease of the eye that has caused loss of vision;
- You have the following risk factors for blood clots:
  - Severe high blood pressure or high blood pressure that is not under control (hypertension);
  - Blood clot disorders such as:
    - Abnormal Factor V Leiden mutation
    - Activated protein C (APC) resistance
    - Antithrombin-III-deficiency
    - Protein C deficiency
    - Protein S deficiency
    - hyperhomocysteinemia
    - Prothrombin mutation G20210A
    - Antiphospholipid-antibodies
  - You have an unusual amount of lipoproteins in your blood;
  - You have diabetes with complications;
  - Increasing age such as a woman older than 50 years old;
  - You have too much body fat (you are obese);
  - A family history of blood clot disorders;
  - You had or will have a major surgery (including to the legs, pelvis or nervous system);
  - You cannot stand or move for long periods of time, including prolonged bed rest;
  - You are a woman over age 35 and smoke;
- You have or had stroke or heart attack;
- You have or had coronary artery disease (including angina) or a condition that may be a first sign of stroke (such as mini stroke, small reversible stroke, chest pains);
- You have disease of the heart valves with complications;
- You have or had liver disease;
- You have or had jaundice (yellowing of the eyes or skin). This may have been related to other medicines you were taking or may have happened during pregnancy;
- You have or have a history of liver tumors (cancerous or non-cancerous);
- You have or might have breast cancer;

- You have a cancer of the uterus or a cancer that is sensitive to hormones;
- You have unusual vaginal bleeding without a known reason;
- You have or have had migraine headaches with or without focal aura (flashes of light, blind spots and other vision changes);
- You have or have had inflammation of the pancreas (pancreatitis) and high levels of fat in your blood (triglycerides);
- You are pregnant or think you might be pregnant;
- You have hepatitis C and are taking combination medication such as ombitasvir/paritaprevir/ritonavir with or without dasabuvir, glecaprevir/pibrentasvir or sofosbuvir/velpatasvir/voxilaprevir

**To help avoid side effects and ensure proper use, talk to your healthcare professional before you take LINESSA. Talk about any health conditions or problems you may have, including if you:**

- have a history of depression;
- have or have a history of kidney problems;
- have a history of liver problems;
- are obese;
- have heart disease;
- have had or will have a major surgery;
- have high blood pressure;
- have or have a family history of diabetes;
- have a history of breast disease or family history of breast cancer;
- have migraine headaches;
- have a family history of blood clots;
- have uterine fibroid tumours (benign tumours of the uterus);
- have cholestasis. This is a condition where the bile flow from the liver is decreased;
- have porphyria;
- are in menopause;
- have inflammatory bowel disease such as Crohn's disease or ulcerative colitis;
- have sickle cell disease. This is a disease that affects hemoglobin, a molecule in red blood cells that delivers oxygen throughout the body;
- have haemolytic uremic syndrome. This is when there is an abnormal breakdown of blood cells, which clogs the kidney;
- have systemic lupus erythematosus. This is a disease of the immune system that affects many organs of the body;
- have a history of seizures/epilepsy;
- have been told that you have a condition called hereditary or acquired angioedema or if you have had episodes of swelling in body parts such as hands, feet, face or airway passages;
- have problems with the valves in your heart and/or have an irregular heart rhythm, or other heart problems;
- have high cholesterol levels;

- wear contact lenses;
- have a history of a skin condition called chloasma (hyperpigmentation);
- are unable to digest lactose or milk products, are on a lactose-free diet or have any of the following diseases:
  - Galactose intolerance
  - Lapp lactase deficiency
  - Glucose-galactose malabsorption

**Other warnings you should know about:**

**LINESSA may not work as well as it should to prevent pregnancy if you:**

- miss pills,
- don't take your pills as directed by your healthcare professional,
- have gastrointestinal problems, or
- are taking certain medicines.

If this happens, you should use another method of birth control, like condoms (barrier method). Talk to your healthcare professional on when you can start LINESSA again.

**Blood clot in legs, lungs, heart, eyes or brain:**

Women who use birth control that contains hormones are more likely to develop blood clots. Blood clots are the most common serious side effects of birth control pills. The risk for blood clots is highest during the first year a woman uses a hormonal birth control. The risk is also high if you restart a hormonal birth control (the same product or a different product) after a break of 4 weeks or more. Clots may occur in many areas of the body can lead to blindness or impaired vision as well as damage to or loss of a limb and death.

While you are taking LINESSA, if you have any of the following symptoms, contact your healthcare professional right away. These are signs of blood clots:

- Sharp pain in the chest, coughing blood, or sudden shortness of breath. These symptoms could indicate a possible blood clot in the lung.
- Crushing chest pain or heaviness, discomfort radiating to your back, jaw, throat or stomach, feeling full. These symptoms could indicate a possible heart attack.
- Sudden severe or worsening headache, vomiting, dizziness, fainting or seizures, trouble walking, changes in vision or speech, weakness or numbness in your face, arm or leg. These symptoms could indicate a possible stroke.
- Pain and/or swelling, redness or slight blue discoloration, skin feeling “warm to the touch” in the calf, irregular heartbeat, confusion. These symptoms could indicate a possible blood clot in the leg.
- Sudden partial or complete loss of vision. This symptom could indicate a possible blood clot in the eye.

## **Cancer:**

Using birth control pills may increase the risk of certain cancers including cancer of the breast, cervix and liver.

### **Breast cancer**

The risk of breast cancer in women increases as you get older. It also increases if there is a family history of breast cancer (mother or sister). Other factors include obesity, never having children or having your first full-term pregnancy at a late age.

Some women who use birth control pills may be at a higher risk of developing breast cancer before menopause. These women may have used birth control pills for a long time (more than eight years) or may have started using birth control pills at an early age.

In a few women, the use of birth control pills may speed up the growth of a breast cancer that has not been found. Finding breast cancer early can reduce the effect of breast cancer on a woman's life expectancy. The risks for breast cancer related to birth control pills seem to be small. You should, however, have a healthcare professional check your breasts examination at least once per year.

While you are taking LINESSA, check your breasts often. See your healthcare professional if you notice any changes, such as:

- Dimpling or sinking of the skin,
- Changes in the nipple, or
- Any lumps you can see or feel.

### **Cervical cancer**

Women who use birth control pills may have a higher chance of getting cervical cancer. However, this may be due to other reasons including infection with the Human Papilloma Virus (HPV). HPV is an important risk factor for cervical cancer. However, it is possible that oral birth control pills may also cause such cancers.

### **Liver cancer**

Liver cancer (hepatocellular carcinoma) and liver tumours may be linked to oral birth control pills. The risk for liver cancer increases the longer these pills are used. However liver tumours are extremely rare.

If you have yellowing of the skin or eyes, dark urine, nausea, vomiting, severe pain or a lump in the abdomen, contact your healthcare professional right away.

### **Gallbladder disease**

The risk for gallbladder disease that needs surgery is higher in women using birth control pills. The risk is highest in the first year of use and increases the longer these pills are used. The risk may double after four or five years of use.

### **Vaginal bleeding**

Breakthrough bleeding or spotting sometimes happens in women using birth control pills including LINESSA. This is blood coming from the vagina between periods. It is most likely to happen in the first months of starting a birth control pill. If the bleeding is heavy or does not stop, contact your healthcare professional.

While you are taking LINESSA you may not get your period each month. If you were not taking LINESSA as directed by your healthcare professional, you should have a pregnancy test. This will rule out if the missed period is because you are pregnant.

If you go more than 6 months without a period contact your healthcare professional. This will be especially important if you also notice secretions from your breasts.

### **Pregnancy, Breastfeeding, Miscarriage and Abortions:**

#### **Use in pregnancy**

Birth control pills should not be taken by pregnant women. Stop taking LINESSA if you get pregnant. You should check with your healthcare professional about risks to your unborn child from any medication taken during pregnancy.

#### **Use after pregnancy, miscarriage or an abortion**

Your healthcare professional will tell you when to start using LINESSA after childbirth, miscarriage or an abortion.

#### **Pregnancy after stopping LINESSA**

You will have a menstrual period when you stop using LINESSA. Wait until after your next period before getting pregnant. You should not get pregnant until another menstrual period occurs within four to six weeks. In this way, the pregnancy can be more accurately dated. Speak to your healthcare professional about other forms of birth control you can use during this time.

#### **Breast feeding**

If you are breast-feeding, talk to your healthcare professional before starting LINESSA. Side effects in the child have been reported, including yellowing of the skin (jaundice) and breast growth. You should use another type of birth control while you are breast-feeding. The use of birth control pills should not be used until breast-feeding has stopped.

### **Skin Conditions**

Chloasma may develop while you are using LINESSA. This appears as yellowish-brown patches on the skin, particularly of the face. It is more likely to happen if you have previously had chloasma gravidarum. This is when these patches appear on the skin of the face during pregnancy. This is commonly known as “the mask of pregnancy”. If you have or had chloasma, avoid too much exposure to the sun while using LINESSA.

## **Surgery**

Tell your healthcare professional if you are scheduled for surgery. You may need to stop using LINESSA one month before surgery and during prolonged bedrest. You may need to wait until after your first period, after hospital discharge, before restarting LINESSA. Talk to your healthcare professional about stopping the use of LINESSA one month before surgery and not using LINESSA for a period of time after surgery or during bed rest.

**Driving and Using Machines:** Drive or use machines with caution when taking LINESSA.

## **Check-ups and tests**

Before starting LINESSA, you will need to have examinations and tests. Your healthcare professional will conduct a physical exam. They will examine your breasts, liver, arms and legs. They will conduct a pelvic exam which includes a PAP smear. Your healthcare professional will also ask you some questions about your personal health history and that of your close relatives. They will also measure your blood pressure and do blood tests.

While you are taking LINESSA, you will need to have regular check-ups with your healthcare professional. Your first check up should be about three months after starting LINESSA. Afterward, you will see your healthcare professional about once per year. At these visits, your healthcare professional will conduct physical and internal exams. He or she will also measure your blood pressure and do blood tests.

If you are scheduled for any laboratory tests, be sure to tell your healthcare professional that you are taking LINESSA. This is because birth control pills can affect some blood tests.

**Tell your healthcare professional about all the medicines you take, including any drugs, vitamins, minerals, natural supplements or alternative medicines.**

## **The following may interact with LINESSA:**

- Medicines used for the treatment of epilepsy, such as primidone, phenytoin, barbiturates, carbamazepine, oxcarbazepine, topiramate, felbamate, ethosuximide, phenobarbital, lamotrigine;
- Medicines used to treat tuberculosis, such as rifampicin, rifabutin;
- Medicines used to treat HIV infections, such as ritonavir
- Medicines used to treat hepatitis C Virus such as boceprevir, telaprevir, ombitsavir/paritaprevir/ritonavir, dasabuvir with or without ribavirin, glecaprevir/pibrentasvir, sofosbuvir/velpatasvir/voxilaprevir;
- Alpha-II adrenoreceptor agents including clonidine;
- Medicines used to treat bacterial infections, such as penicillins, tetracyclines, metronidazole, erythromycin, ampicillin, cotrimoxazole, chloramphenicol, neomycin, nitrofurantoin, sulfonamides, tetracyclines, troleandomycin;
- Medicines used to treat fungal infections, such as griseofulvin, fluconazole, itraconazole, ketoconazole, voriconazole, clarithromycin;

- Medicines used to lower cholesterol, such as clofibrate;
- Medicines used to prevent blood clots;
- St. John’s wort, an herbal product used to treat depression;
- Medicines used to treat diabetes including insulin and others that lower blood sugar;
- Medicines used to help you relax or sleep, such as benzodiazepines, barbiturates, chloral hydrate, glutethimide, meprobamate, chlordiazepoxide, lorazepam, oxazepam, diazepam, phenothiazines;
- Medicines used to treat depressants, such as clomipramine;
- Medicines used to treat fever, pain or inflammation, such as phenylbutazone, acetaminophen, ASA, antipyrine, meperidine, prednisone;
- Medicines used to help prevent organ rejection, such as cyclosporine;
- Medicines used to treat migraine headaches;
- Medicines used to treat allergies;
- Nutritional supplements, such as vitamin E, vitamin B12, folic acid;
- Medicines used to help prevent organ rejection including cyclosporine;
- A medicine used to help treat bleeding called aminocaproic acid;
- Medicines used to treat lung diseases such as asthma and COPD (bronchitis, emphysema) including theophylline;
- Medicines used to slow the heart rate including isoproterenol;
- Medicines used to treat high blood pressure including guanethidine, methyldopa, beta blockers, reserpine, diltiazem and verapamil;

Antacids may affect how LINESSA is absorbed in your body. If you need to use antacids, like TUMS, take them 2 hours before or 2 hours after taking LINESSA.


The effects of caffeine and alcohol may also be increased. This is because birth control pills affect how these are metabolized.

**How to take LINESSA:**

- **Read these directions:**
  - before you start taking LINESSA, and
  - any time you are not sure what to do.
- **Look at your pill pack** to see if it has 21 or 28 pills:
  - LINESSA 21 (21-Day Pack):
    - 21 active pills (with hormones): 7 light yellow, 7 orange and 7 red. These are taken daily for three weeks, and then no pills for one week;
  - or
  - LINESSA 28 (28-Day Pack):
    - 21 active pills (with hormones): 7 light yellow, 7 orange and 7 red. These are taken daily for three weeks, then
    - 7 green “reminder” pills (without hormones): taken daily for one week.

- Decide with your healthcare professional what is the best day for you to start taking your first pill. Pick a time of day that will be easy to remember (like when eating a meal or going to bed).
- Label the pill pack by selecting the appropriate day label strip: **Day 1** or **Sunday** start (see below for explanation). Place the day label strip in the space where you see the words "Start/ Début". Having the pill pack labelled with the days of the week will help remind you to take your pill every day.
- **Taking LINESSA:**

#### A. LINESSA 21 - Day Pack




- With this type of birth control pill, you are on pills for 21 days and off pills for 7 days. You must not be off the pill for more than 7 days in a row.

#### If you have not used hormonal birth control in the past month:

1. **The first day of your menstrual period (bleeding) is Day 1 of your cycle.** Your healthcare professional may tell you to start taking LINESSA on Day 1 or on the first Sunday after your period begins. If your period starts on Sunday, start that same day.
2. Take one pill at about the same time every day for 21 days. Follow the arrows marked on the pill pack (one light yellow tablet daily for 7 days, one orange tablet daily for 7 days, and one red tablet daily for 7 days).
3. **Then, do NOT take any pills for 7 days.** You will probably have a period during the 7 days you do not take LINESSA. This bleeding may be lighter and shorter than your usual period.
4. Start a new pack on the eighth day.



## B. LINESSA 28 - Day Pack



- With this type of birth control pill, you take 21 pills that contain hormones and seven “reminder” pills that contain no hormones.

### If you have not used hormonal birth control in the past month:

1. **The first day of your menstrual period (bleeding) is Day 1 of your cycle.** Your healthcare professional may advise you to start taking LINESSA on Day 1 or on the first Sunday after your period begins. If your period starts on Sunday, start that same day.
2. Take one pill at about the same time every day for 28 days. Follow the arrows marked on the pill pack (one light yellow tablet daily for 7 days, one orange tablet daily for 7 days, one red tablet daily for 7 days, and one green “reminder” tablet daily for 7 days).
3. Your period should occur during the last 7 days of using the pill pack (i.e. while you are taking the green “reminder” pills).
4. Begin a new pack the next day. **DO NOT miss any days.**

### Other instructions for LINESSA 21 and LINESSA 28:

- You may wish to use a second method of birth control (e.g. condoms and spermicidal foam or gel) for the first seven days of the first cycle of pill use. This will provide a back-up in case pills are forgotten while you are getting used to taking them.
- When receiving any medical treatment, be sure to tell your healthcare professional that you are using birth control pills.

- Many women have spotting or light bleeding, or may feel sick to their stomach during the first three months taking birth control pills. If you do feel sick, do not stop taking LINESSA. The problem will usually go away. If it does not go away, check with your healthcare professional or clinic.
- Do NOT stop taking LINESSA or skip any pills even if you are sick to your stomach, have bleeding between your periods or do not have sex very often.
- If you have vomiting or diarrhea within 3-4 hours of taking a LINESSA pill, LINESSA may not work as well. Use a back-up method of birth control until you can check with your healthcare professional or clinic. See "Missed Dose" below for more instructions.

If your questions are not answered here, ask your healthcare professional or clinic.

### **Switching to LINESSA 21 or LINESSA 28:**

- Talk to your healthcare professional about when to start taking LINESSA.

### **If you are switching from another combined birth control (combined birth control pill, vaginal ring or transdermal patch):**

- Switching from another combined birth control pill: Start LINESSA on the day after the last active pill (the last pill containing hormones). If this is not possible, you should start LINESSA on the day after your usual pill-free interval or after the last "reminder" pill.
- If a vaginal ring or transdermal patch has been used, start using LINESSA on the day of removal. If this is not possible, start LINESSA when the next application would have been due.

### **If you are switching from a progestogen-only-method (mini-pill, injection, implant) or from a progestogen-releasing intrauterine system (IUS):**

- Switching from a mini-pill: You may start LINESSA any day after the mini-pill.
- Switching from an injectable: Start LINESSA when the next injection would be due.
- Switching from an implant or an IUS: Start LINESSA on the day of removal.

In all of these cases you should use a back-up barrier birth control method (like condoms) for the first 7 days of taking LINESSA.

### **If you are starting following a first-trimester abortion:**

- You may start LINESSA right away. You do not need to use any back-up birth control.

### **If you are starting following delivery of a baby or a second-trimester abortion:**

- You should start LINESSA between days 21 and 28 after delivery or second trimester abortion.
- When starting later, you should use a back-up method for the first 7 days you take LINESSA.
- If you already had sex or intercourse, you must make sure you are not pregnant or wait for your first menstrual period before starting LINESSA.
- If you are breastfeeding, or planning to breastfeed, talk to your healthcare professional.

**Usual Adult Dose:**

- **LINESSA 21:** Take 1 tablet each day for 21 days and off pills for 7 days. Begin a new pack the next day.
- **LINESSA 28:** Take 1 tablet each day for 28 days. Begin a new pack the next day.

**Overdose:**

If too many birth control pills are taken at one time, nausea, vomiting, breast tenderness, dizziness, abdominal pain, fatigue/drowsiness and withdrawal bleeding may happen.

If you think you, or a person you are caring for, have taken too much LINESSA, contact a healthcare professional, hospital emergency department or regional Poison Control Centre immediately, even if there are no symptoms.

**Missed Dose:**

**If you miss pills at any time, you could get pregnant.** The greatest risks for pregnancy are:

- when you start a pack late
- when you miss pills at the beginning or at the very end of the pack.

Missing pills also can cause some spotting or light bleeding, even if you make up the missed pills. You also could feel a little sick to your stomach on the days you take two pills to make up for missed pills.

If you forget more than one pill two months in a row, talk to your healthcare professional or clinic about how to make pill-taking easier or about using another method of birth control.

The following chart explains what you should do if you miss one or more birth control pills. Match the number of pills missed with the appropriate starting time for your pill pack.

| Sunday Start  | Day 1 Start |
|---|-------------|
| <b>Miss 1 Pill</b>  |             |
| Take it as soon as you remember, and take the next pill at the usual time. This means that you might take 2 pills in one day. |             |

| Sunday Start  | Day 1 Start   |
|---|---|
| <b>Miss 2 Pills in a Row</b>  |   |
| <p><b>First 2 weeks</b></p> <ol style="list-style-type: none"> <li>1. Take 2 pills the day you remember and 2 pills the next day.</li> <li>2. Then take 1 pill a day until you finish the pack.</li> <li>3. Use a non-hormonal back-up method of birth control if you have sex in the 7 days after you miss the pills.</li> </ol>   |   |
| <p><b>Third Week:</b></p> <ol style="list-style-type: none"> <li>1. Keep taking 1 pill a day until Sunday.</li> <li>2. On Sunday, safely throw away the rest of the pack and start a new pack that day.</li> <li>3. Use a non-hormonal back-up method of birth control if you have sex in the 7 days after you miss the pills.</li> <li>4. You may not have a period this month.</li> </ol> <p><b>If you miss 2 periods in a row, call your healthcare professional or clinic.</b></p>          | <p><b>Third Week:</b></p> <ol style="list-style-type: none"> <li>1. Safely throw away the rest of the pill pack and start a new pack that same day.</li> <li>2. Use a non-hormonal back-up method of birth control if you have sex in the 7 days after you miss the pills.</li> <li>3. You may not have a period this month.</li> </ol> <p><b>If you miss 2 periods in a row call your healthcare professional or clinic.</b></p>           |
| <b>Miss 3 or More Pills in a Row</b>  |   |
| <p><b>Anytime in the Cycle:</b></p> <ol style="list-style-type: none"> <li>1. Keep taking 1 pill a day until Sunday.</li> <li>2. On Sunday, safely throw away the rest of the pack and start a new pack that day.</li> <li>3. Use a non-hormonal back-up method of birth control if you have sex in the 7 days after you miss the pills.</li> <li>4. You may not have a period this month.</li> </ol> <p><b>If you miss 2 periods in a row call your healthcare professional or clinic.</b></p> | <p><b>Anytime in the Cycle:</b></p> <ol style="list-style-type: none"> <li>1. Safely throw away the rest of the pill pack and start a new pack that same day.</li> <li>2. Use a non-hormonal back-up method of birth control if you have sex in the 7 days after you miss the pills.</li> <li>3. You may not have a period this month.</li> </ol> <p><b>If you miss 2 periods in a row call your healthcare professional or clinic.</b></p> |

**LINESSA 28 - Day Pack:** If you forget any of the 7 green “reminder” pills in Week 4, just safely throw away the pills you missed. Then keep taking one pill each day until the pack is empty. You do not need to use a back-up method.

**Always be sure you have ready:**

- An extra full pack of pills;
- Another kind of birth control (such as condoms and spermicidal foam or gel) to use as a back-up in case you miss pills. You will need back-up birth control if you miss pills and in some other situations. Always talk to your healthcare professional if you are not sure whether you need to use back-up birth control.

## What are possible side effects from using LINESSA?

These are not all the possible side effects you may feel when taking LINESSA. If you experience any side effects not listed here, contact your healthcare professional.

- migraine, severe headaches
- acne
- rash
- skin colour changes, red skin lumps
- hair loss or increase in growth
- nausea, vomiting
- abdominal or back pain
- sleep disorder like insomnia
- dizziness
- diarrhea
- urinary tract infections or inflammation
- painful period cramps
- flu-like symptoms
- bronchitis, runny or stuffy nose, sore throat, common cold
- loss of strength, weakness, fatigue
- feeling of physical discomfort or uneasiness
- cough
- fever
- indigestion
- weight gain
- difficulty wearing contact lenses
- nervousness, anxiety, mood swings

| Serious side effects and what to do about them   |                                      |              |   |
|--|--------------------------------------|--------------|---|
| Symptom / effect   | Talk to your healthcare professional |              | Stop taking drug and get immediate medical help |
|  | Only if severe                       | In all cases |   |
| <b>UNCOMMON</b>  |                                      |              |   |
| <b>Arterial thromboembolism, Myocardial infarction</b> (blood clot in the artery, heart attack): sudden pain, discomfort, pressure, heaviness, sensation of squeezing or fullness in the shoulder, chest, arm, or below the breastbone; discomfort radiating to the back, jaw, throat, arm, stomach, feeling of being full, having indigestion or choking; sweating, nausea, vomiting or dizziness; extreme weakness, anxiety, or shortness of breath; rapid or irregular heartbeats, cold sweat, heart burn |                                      |              | √   |
| <b>Behavior and mood changes:</b> agitation including aggressive behavior or hostility, changes in sexual desire or sexual activity, increased eating, stress  | √                                    |              |   |
| <b>Breast changes (breast lumps/breast cancer):</b> pain and tenderness, lumps, nipple discharge   |                                      | √            |   |
| <b>Blood clot on the eye:</b> sudden partial or complete loss of vision or double vision   |                                      |              | √   |
| <b>Depression:</b> persistent sad mood accompanied by difficulty in sleeping, weakness, lack of energy, fatigue  |                                      | √            |   |
| <b>Reproductive System Disorders:</b> Pelvic pain, painful intercourse, abdominal bloating or swelling, pain during bowel movements; cysts usually disappear on their own within a few months and may not show symptoms; serious cysts are uncommon  |                                      | √            |   |

|  |   |   |   |
|--|---|---|---|
| <b>Deep vein thrombosis</b> (blood clot in the deep veins of legs or arms): swelling of one leg or one foot, pain or tenderness in the leg, difficulty standing or walking, feeling of warmth in the leg, red or discolored skin on the leg, sudden pain, swelling and slight blue discoloration of an extremity |   |   | ✓ |
| <b>Hypersensitivity</b> (allergic reaction): rash, hives, swelling of the face, lips, tongue or throat, difficulty swallowing or breathing, feeling sick to your stomach and throwing up; wheezing, drop in blood pressure   |   |   | ✓ |
| <b>Liver problems including liver tumour, jaundice:</b> abnormal liver test, yellowing of the skin or eyes, dark urine, nausea, vomiting, severe pain or lump in the abdomen, loss of appetite, fever, light-coloured bowel movements  |   |   | ✓ |
| <b>Pulmonary embolism (blood clot in the lung):</b> sharp pain in the chest, coughing blood, sudden shortness of breath  |   |   | ✓ |
| <b>Stroke:</b> sudden severe or worsening headache, vomiting, dizziness, fainting, vision or speech problems, weakness or numbness in the arm or leg   |   |   | ✓ |
| <b>Edema:</b> swelling of the arms or legs   |   | ✓ |   |
| <b>Vaginal bleeding changes:</b> increased or decreased menstrual bleeding, spotting, infrequent periods or absence of bleeding  | ✓ |   |   |
| <b>Vaginal infection</b> (inflammation of the vagina or surrounding area): itching, or unusual or increased vaginal discharge  | ✓ |   |   |

If you have a troublesome symptom or side effect that is not listed here or becomes bad enough to interfere with your daily activities, talk to your healthcare professional.

## Reporting Side Effects

You can report any suspected side effects associated with the use of health products to Health Canada by:

- Visiting the Web page on Adverse Reaction Reporting (<https://www.canada.ca/en/health-canada/services/drugs-health-products/medeffect-canada/adverse-reaction-reporting.html>) for information on how to report online, by mail or by fax; or
- Calling toll-free at 1-866-234-2345.

*NOTE: Contact your health professional if you need information about how to manage your side effects. The Canada Vigilance Program does not provide medical advice.*

## Storage:

- Store at room temperature (15°C – 30°C).
- Keep in a safe place out of reach and sight of children and pets.
- Do not throw away any drugs via wastewater or household waste. Ask your pharmacist how to throw away drugs you no longer use. These measures will help to protect the environment.

## If you want more information about LINESSA:

- Talk to your healthcare professional
- Find the full product monograph that is prepared for healthcare professionals and includes this Patient Medication Information by visiting the Health Canada website (<https://www.canada.ca/en/health-canada/services/drugs-health-products/drug-products/drug-product-database.html>); or by contacting Aspen Pharmacare Canada Inc. at [www.aspenpharma.ca](http://www.aspenpharma.ca) or at 1-844-330-1213.

This leaflet was prepared by:  
Aspen Pharmacare Canada Inc.  
8 – 1155 North Service Road West  
Oakville, ON  
L6M 3E3

Trademarks are owned by or licensed to the Aspen Group of companies.  
© 2023 Aspen Group of companies or its licensor. All rights reserved.

Last Revised: September 12, 2023