

**PRODUCT MONOGRAPH**

**PrAPO-TIZANIDINE**

**Tizanidine Hydrochloride Tablets USP**

**2 mg and 4 mg Tizanidine**

**Antispastic Agent**

**APOTEX INC  
150 Signet Drive  
Toronto, Ontario  
Canada M9L 1T9**

**DATE OF PREPARATION:  
OCT 25, 2023**

**Control No.: 279972**

## Table of Contents

<b>PART I: HEALTH PROFESSIONAL INFORMATION .....</b>	<b>3</b>
SUMMARY PRODUCT INFORMATION.....	3
INDICATIONS AND CLINICAL USE.....	3
CONTRAINDICATIONS.....	3
WARNINGS AND PRECAUTIONS .....	3
ADVERSE REACTIONS .....	8
DRUG INTERACTIONS .....	12
DOSAGE AND ADMINISTRATION.....	13
OVERDOSAGE.....	14
ACTION AND CLINICAL PHARMACOLOGY .....	14
STORAGE AND STABILITY .....	16
DOSAGE FORMS, COMPOSITION AND PACKAGING.....	16
<b>PART II: SCIENTIFIC INFORMATION.....</b>	<b>17</b>
PHARMACEUTICAL INFORMATION.....	17
CLINICAL TRIALS .....	18
DETAILED PHARMACOLOGY .....	21
TOXICOLOGY .....	22
REFERENCES.....	24
<b>PART III: CONSUMER INFORMATION .....</b>	<b>25</b>

**Pr**APO-TIZANIDINE  
Tizanidine Hydrochloride Tablets USP

**PART I: HEALTH PROFESSIONAL INFORMATION**

**SUMMARY PRODUCT INFORMATION**

<b>Route of Administration</b>	<b>Dosage Form / Strength</b>	<b>Clinically Relevant Nonmedicinal Ingredients</b>
Oral	Tablet 2 mg , 4 mg	colloidal silicon dioxide, stearic acid, microcrystalline cellulose and anhydrous lactose <i>For a complete listing see Dosage Forms, Composition and Packaging section.</i>

**INDICATIONS AND CLINICAL USE**

**Adults:**

APO-TIZANIDINE (tizanidine HCl) is a short-acting drug for the management of spasticity.

**Geriatrics:**

Evidence from clinical studies and experience suggests that use in the geriatric population may be associated with differences in safety or effectiveness and a brief discussion can be found in the appropriate sections (**WARNINGS AND PRECAUTIONS, ACTION AND CLINICAL PHARMACOLOGY**).

**Pediatrics (< 18 years of age):**

No data are available.

**CONTRAINDICATIONS**

Patients who are hypersensitive to this drug or to any ingredient in the formulation or component of the container. For a complete listing, see the **DOSAGE FORMS, COMPOSITION AND PACKAGING** section of the product monograph.

Co-administration of tizanidine with moderate and potent CYP1A2 inhibitors such as fluvoxamine or ciprofloxacin is contra-indicated (see **WARNINGS AND PRECAUTIONS**).

APO-TIZANIDINE is contraindicated in patients for whom spasticity is needed to maintain function, such as maintenance of upright posture and balance in locomotion.

**WARNINGS AND PRECAUTIONS**

**General**

### **Hypotension**

Tizanidine HCl is an  $\alpha_2$ -adrenergic agonist (like clonidine) and can produce hypotension. In a single dose study where blood pressure was monitored closely after dosing, two thirds of patients treated with 8 mg of tizanidine had a 20% reduction in either the diastolic or systolic BP. The reduction was seen within 1 hour after dosing, peaked 2 to 3 hours after dosing and was associated, at times, with bradycardia, orthostatic hypotension, lightheadedness/dizziness and rarely syncope. The hypotensive effect is dose related and has been measured following single doses of  $\geq 2$  mg.

The chance of significant hypotension may possibly be minimized by titration of the dose and by focusing attention on signs and symptoms of hypotension prior to dose advancement. In addition, patients moving from a supine to a fixed upright position may be at increased risk for hypotensive and orthostatic effects.

Caution is advised when APO-TIZANIDINE is to be used in patients who have a history of orthostatic hypotension or labile blood pressure or who are receiving concurrent antihypertensive therapy. APO-TIZANIDINE should not be used with other  $\alpha_2$ -adrenergic agonists.

### **Risk of Liver Injury**

Tizanidine use occasionally causes drug induced liver injury, most often hepatocellular in type. In controlled clinical studies, approximately 5% of patients treated with tizanidine had elevations of liver function tests (ALT/SGPT, AST/SGOT) to greater than 3 times the upper limit of normal (or 2 times if baseline levels were elevated). The patients usually remain asymptomatic despite increased aminotransferases. In occasional symptomatic cases, nausea, vomiting, anorexia and jaundice have been reported. The onset of the elevated liver enzymes typically occurred within the first 6 months of treatment with tizanidine and most resolved rapidly upon drug withdrawal with no reported residual problems. In postmarketing experience, three deaths associated with liver failure have been reported in patients treated with tizanidine, including one case of fatal fulminant hepatitis.

Monitoring of aminotransferase levels is recommended during the first 6 months of treatment (e.g., baseline, 1, 3 and 6 months) and periodically thereafter, based on clinical status. Because of the potential toxic hepatic effect of tizanidine, the drug should be used only with extreme caution in patients with impaired hepatic function.

### **Sedation**

In the multiple dose, controlled clinical studies, 48% of patients receiving any dose of tizanidine reported sedation as an adverse event. In 10% of these cases, the sedation was rated as severe compared to <1% in the placebo treated patients. Sedation may interfere with every day activity.

The effect appears to be dose related. In a single dose study, 92% of the patients receiving 16 mg, when asked, reported that they were drowsy during the 6-hour study. This compares to 76% of the patients on 8 mg and 35% of the patients on placebo. Patients began noting this effect 30 minutes following dosing. The effect peaked 1.5 hours following dosing. Of the patients who received a single dose of 16 mg, 51% continued to report drowsiness 6 hours following dosing compared to 13% in the patients receiving placebo or 8 mg of tizanidine.

In the multiple dose studies, the prevalence of patients with sedation peaked following the first week of titration and then remained stable for the duration of the maintenance phase of the study.

### **Hallucinations**

Tizanidine use has been associated with hallucinations. Formed, visual hallucinations or delusions have been reported in 5 of 170 patients (3%) in two North American controlled clinical studies. Most of the patients were aware that the events were unreal. One patient developed psychoses in association with the hallucinations. One patient continued to have problems for at least 2 weeks following discontinuation of tizanidine. Dosage reduction or discontinuation should be considered for patients who experience hallucinations while receiving APO-TIZANIDINE. Particular caution should be observed if APO-TIZANIDINE is administered to patients with a prior history of psychotic illness.

### **Limited Database for Chronic Use of Single Doses Above 8 mg and Multiple Doses Above 24 mg per day**

Clinical experience with long-term use of tizanidine at single doses of 8 to 16 mg or total daily doses of 24 to 36 mg is limited. Approximately 75 patients have been exposed to individual doses of 12 mg or more for at least one year and approximately 80 patients have been exposed to total daily doses of 30 to 36 mg/day for at least one year. There is essentially no long-term experience with single, daytime doses of 16 mg. Because long-term clinical study experience at high doses is limited, only those adverse events with a relatively high incidence are likely to have been identified.

### **Discontinuation of Treatment with APO-TIZANIDINE**

If therapy needs to be discontinued, especially in patients who have been receiving high doses for long periods, the dose should be decreased slowly to minimize the risk of withdrawal and rebound hypertension, tachycardia, and hypertonia.

**The following additional precautions are listed alphabetically.**

#### **Carcinogenesis and Mutagenesis**

Only animal carcinogenesis data and mutagenesis data from *in vitro* and *in vivo* assays are available (see **TOXICOLOGY**).

#### **Cardiovascular**

Prolongation of the QT interval and bradycardia were noted in chronic toxicity studies in dogs at doses equal to the maximum human dose on a mg/m<sup>2</sup> basis. ECG evaluation was not performed in the controlled clinical studies. There have been post-market reports of QT prolongation and a small number of reports of Torsades de Pointes, none of them fatal, during tizanidine treatment.

#### **Caution should be exercised when APO-TIZANIDINE is prescribed with drugs known to prolong the QT interval.**

Tizanidine HCl can produce hypotension associated, at times, with bradycardia and orthostatic hypotension, dizziness and rarely syncope (See **WARNINGS AND PRECAUTIONS, General**).

#### **Dependence/Tolerance**

Monkeys were shown to self-administer tizanidine in a dose-dependent manner, and abrupt cessation of tizanidine produced transient signs of withdrawal at doses > 35 times the maximum recommended human dose on a mg/m<sup>2</sup> basis. These transient withdrawal signs (increased

locomotion, body twitching, and aversive behavior toward the observer) were not reversed by naloxone administration. Tizanidine is closely related to clonidine, which is often abused in combination with narcotics and is known to cause symptoms of rebound upon abrupt withdrawal. There have been cases of rebound symptoms reported on sudden withdrawal of tizanidine. Some of the case reports suggest that these patients were also misusing opioids. Withdrawal symptoms included but were not limited to: hypertension, tachycardia, hypertonia, convulsions, tremor, and anxiety. As with clonidine, withdrawal is expected to be more likely in cases where high doses are used, especially for prolonged periods. There have been also reports of tizanidine abuse and dependence, most of them with concomitant use of opioids, benzodiazepines, other hypnotics or multiple analgesics. The potential for tizanidine abuse should be monitored, especially in patients simultaneously using opioids or benzodiazepines.

### **Drug Interaction with CYP1A2 inhibitors**

Concomitant use of tizanidine and moderate or potent CYP450 1A2 inhibitors is contraindicated (see **CONTRAINDICATIONS**). Concomitant use of tizanidine with fluvoxamine, a potent CYP450 1A2 inhibitor in man, resulted in a 33-fold increase in the tizanidine AUC by fluvoxamine; concomitant use of tizanidine with ciprofloxacin, another CYP1A2 inhibitor, resulted in a 10-fold increase in tizanidine AUC; in both studies, clinically significant hypotension resulted along with somnolence, dizziness and decreased psychomotor performance (see **DRUG INTERACTIONS**). Co-administration of tizanidine with other inhibitors of CYP1A2 such as antiarrhythmics (amiodarone, mexiletine, propafenone), cimetidine, other fluoroquinolones (norfloxacin, moxifloxacin), and ticlopidine should be avoided or used with caution.

### **Hepatic/Biliary/Pancreatic**

Tizanidine use occasionally causes drug induced liver injury, most often hepatocellular in type (See **WARNINGS AND PRECAUTIONS, General**).

### **Information to be Provided to the Patients**

Patients should be advised of the limited clinical experience with APO-TIZANIDINE both in regard to duration of use and the higher doses required to reduce muscle tone (see **WARNINGS AND PRECAUTIONS, General**).

Because of the possibility of APO-TIZANIDINE lowering blood pressure, patients should be warned about the risk of clinically significant orthostatic hypotension (see **WARNINGS AND PRECAUTIONS, General**).

Because of the possibility of sedation, patients should be warned about performing activities requiring alertness, such as driving a vehicle or operating machinery (See **WARNINGS AND PRECAUTIONS, General**). Patients should also be instructed that the sedation may be additive when APO-TIZANIDINE is taken in conjunction with drugs (baclofen, benzodiazepines) or substances (e.g., alcohol) that act as CNS depressants.

### **Neurologic**

Tizanidine use has been associated with sedation (See **WARNINGS AND PRECAUTIONS, General**).

Tizanidine use has been associated with hallucinations (See **WARNINGS AND PRECAUTIONS, General**).

### **Ophthalmologic**

Dose-related retinal degeneration and corneal opacities have been found in animal studies at doses equivalent to approximately the maximum recommended dose on a mg/m<sup>2</sup> basis. There have been no reports of corneal opacities or retinal degeneration in the clinical studies.

### **Renal**

APO-TIZANIDINE should be used with caution in patients with renal insufficiency (Cl<sub>cr</sub> <25 mL/min), as clearance is reduced by more than 50%. In these patients, during titration, the individual doses should be reduced. If higher doses are required, individual doses rather than dosing frequency should be increased. These patients should be monitored closely for onset or increase in severity of the common adverse events (dry mouth, somnolence, asthenia and dizziness) as indicators of potential overdose.

### **Special Populations**

#### **Use in Women Taking Oral Contraceptives**

APO-TIZANIDINE should be used with caution in women taking oral contraceptives; as clearance of tizanidine is reduced by approximately 50% in such patients. In these patients, during titration, the individual doses should be reduced.

**Pregnant Women:** The effect of APO-TIZANIDINE on labor and delivery in humans is unknown.

Reproduction studies performed in rats at a dose of 3 mg/kg, equal to the maximum recommended human dose on a mg/m<sup>2</sup> basis and in rabbits at 30 mg/kg, 16 times the maximum recommended human dose on a mg/m<sup>2</sup> basis did not show evidence of teratogenicity. Tizanidine at doses equal to and up to 8 times the maximum recommended human dose on a mg/m<sup>2</sup> basis increased gestation duration in rats. Prenatal and postnatal pup loss was increased and developmental retardation occurred. Post-implantation loss was increased in rabbits at doses of 1 mg/kg or greater, equal to or greater than 0.5 times the maximum recommended human dose on a mg/m<sup>2</sup> basis. APO-TIZANIDINE has not been studied in pregnant women. APO-TIZANIDINE should be given to pregnant women only if the potential benefit clearly outweighs the potential risk to mother and child.

**Nursing Women:** It is not known whether APO-TIZANIDINE is excreted in human milk, although as a lipid soluble drug, it might be expected to pass into breast milk.

**Pediatrics** (< 18 years of age): There are no adequate and well-controlled studies to document the safety and efficacy of APO-TIZANIDINE in children under 18 years in age.

**Geriatrics:** APO-TIZANIDINE should be used with caution in elderly patients because clearance is decreased four-fold.

## ADVERSE REACTIONS

### Adverse Drug Reaction Overview

In multiple dose, placebo-controlled clinical studies, 264 patients were treated with tizanidine and 261 with placebo. Adverse events, including severe adverse events, were more frequently reported with tizanidine than with placebo.

### **Common adverse events leading to discontinuation**

Forty five of 264 (17%) patients receiving tizanidine and 13 of 261 (5%) patients receiving placebo in three multiple dose, placebo-controlled clinical studies discontinued treatment for adverse events. When patients withdrew from the study, they frequently had more than one reason for discontinuing. The adverse events most frequently leading to withdrawal of tizanidine treated patients in the controlled clinical studies were asthenia (weakness, fatigue and/or tiredness) (3%), somnolence (3%), dry mouth (3%), increased spasm or tone (2%) and dizziness (2%).

### **Most Frequent Adverse Clinical Events Seen in Association with the Use of APO-Tizanidine**

In multiple-dose, placebo-controlled clinical studies involving 264 patients with spasticity, the most frequent adverse events were dry mouth, somnolence/sedation, asthenia (weakness, fatigue and/or tiredness) and dizziness. Three quarters of the patients rated the events as mild to moderate and one quarter of the patients rated the events as being severe. These events appeared to be dose related.

### Clinical Trial Adverse Drug Reactions

*Because clinical trials are conducted under very specific conditions the adverse reaction rates observed in the clinical trials may not reflect the rates observed in practice and should not be compared to the rates in the clinical trials of another drug. Adverse drug reaction information from clinical trials is useful for identifying drug-related adverse events and for approximating rates.*

The events cited reflect experience gained under closely monitored conditions of clinical studies in a highly selected patient population. In actual clinical practice or in other clinical studies, these frequency estimates may not apply, as the conditions of use, reporting behavior, and the kinds of patients treated may differ. Table 1 lists treatment emergent signs and symptoms that were reported in greater than 2% of patients in three multiple dose, placebo-controlled studies who received tizanidine where the frequency in the tizanidine group was at least as common as in the placebo group. These events are not necessarily related to tizanidine treatment. For comparison purposes, the corresponding frequency of the event (per 100 patients) among placebo treated patients is also provided.

**Table 1 - Multiple Dose, Placebo-Controlled Studies (Frequent (> 2%) Adverse Events Reported for Which Tizanidine Incidence is Greater Than Placebo)**

	<b>Tizanidine n= 264 (%)</b>	<b>placebo n= 261 (%)</b>
Dry mouth	49	10
Somnolence	48	10
Asthenia*	41	16
Dizziness	16	4

	<b>Tizanidine n= 264 (%)</b>	<b>placebo n= 261 (%)</b>
UTI	10	7
Infection	6	5
Constipation	4	1
Liver function tests abnormal	3	<1
Vomiting	3	0
Speech disorder	3	0
Amblyopia (blurred vision)	3	<1
Urinary frequency	3	2
Flu syndrome	3	2
SGPT/ALT increased	3	<1
Dyskinesia	3	0
Nervousness	3	<1
Pharyngitis	3	1
Rhinitis	3	2

\* weakness, fatigue and/or tiredness

In the single dose, placebo-controlled study involving 142 patients with spasticity, the patients were specifically asked if they had experienced any of the four most common adverse events dry mouth, somnolence (drowsiness), asthenia (weakness, fatigue and/or tiredness), and dizziness.

In addition, hypotension and bradycardia were observed. The occurrence of these adverse events is summarized in Table 2. Other events were, in general, reported at a rate of 2% or less.

	<b>Tizanidine 8 mg n= 45 (%)</b>	<b>Tizanidine 16 mg n= 49 (%)</b>	<b>placebo n= 48 (%)</b>
Somnolence	78	92	31
Dry Mouth	76	88	35
Asthenia*	67	78	40
Dizziness	22	45	4
Hypotension	16	33	0
Bradycardia	2	10	0

\* weakness, fatigue and/or tiredness

### **Other Adverse Events Observed During the Evaluation of Tizanidine**

Tizanidine was administered to 1187 patients in additional clinical studies where adverse event information was available. The conditions and duration of exposure varied greatly, and included (in overlapping categories) double-blind and open-label studies, uncontrolled and controlled studies, inpatient and outpatient studies, and titration studies. Untoward events associated with this exposure were recorded by clinical investigators using terminology of their own choosing.

Consequently, it is not possible to provide a meaningful estimate of the proportion of individuals experiencing adverse events without first grouping similar types of untoward events into a smaller number of standardized event categories.

In the tabulations that follow, reported adverse events were classified using a standard COSTART-based dictionary terminology. The frequencies presented, therefore, represent the proportion of the 1187 patients exposed to tizanidine who experienced an event of the type cited on at least one occasion while receiving tizanidine. All reported events are included except those already listed in Table 1. If the COSTART term for an event was so general as to be uninformative, it was replaced with a more informative term. It is important to emphasize that, although the events reported occurred during treatment with tizanidine, they were not necessarily caused by it.

Events are further categorized by body system and listed in order of decreasing frequency according to the following definitions: frequent adverse events are those occurring on one or more occasions in at least 1/100 patients (only those not already listed in the tabulated results from placebo-controlled studies appear in this listing); infrequent adverse events are those occurring in 1/100 to 1/1000 patients.

**Body as a Whole:** *Frequent:* fever; *Infrequent:* allergic reaction, moniliasis, malaise, abscess, neck pain, sepsis, cellulitis, death, overdose; *Rare:* carcinoma, congenital anomaly, suicide attempt.

**Cardiovascular System:** *Infrequent:* vasodilatation, postural hypotension, syncope, migraine, arrhythmia; *Rare:* angina pectoris, coronary artery disorder, heart failure, myocardial infarct, phlebitis, pulmonary embolus, ventricular extrasystoles, ventricular tachycardia.

**Digestive System:** *Frequent:* abdomen pain, diarrhea, dyspepsia; *Infrequent:* dysphagia, cholelithiasis, fecal impaction, flatulence, gastrointestinal hemorrhage, hepatitis, melena; *Rare:* gastroenteritis, hematemesis, hepatoma, intestinal obstruction, liver damage.

**Hemic and Lymphatic System:** *Infrequent:* ecchymosis, hypercholesteremia, anemia, hyperlipemia, leukopenia, leukocytosis, sepsis; *Rare:* petechia, purpura, thrombocythemia, thrombocytopenia.

**Metabolic and Nutritional System:** *Infrequent:* edema, hypothyroidism, weight loss; *Rare:* adrenal cortex insufficiency, hyperglycemia, hypokalemia, hyponatremia, hypoproteinemia, respiratory acidosis.

**Musculoskeletal System:** *Frequent:* myasthenia, back pain; *Infrequent:* pathological fracture, arthralgia, arthritis, bursitis.

**Nervous System:** *Frequent:* depression, anxiety, paresthesia; *Infrequent:* tremor, emotional lability, convulsion, paralysis, thinking abnormal, vertigo, abnormal dreams, agitation, depersonalization, euphoria, migraine, stupor, dysautonomia, neuralgia; *Rare:* dementia, hemiplegia, neuropathy.

**Respiratory System:** *Infrequent:* sinusitis, pneumonia, bronchitis; *Rare:* asthma.

**Skin and Appendages:** *Frequent:* rash, sweating, skin ulcer; *Infrequent:* pruritus, dry skin, acne, alopecia, urticaria; *Rare:* exfoliative dermatitis, herpes simplex, herpes zoster, skin carcinoma.

**Special Senses:** *Infrequent:* ear pain, tinnitus, deafness, glaucoma, conjunctivitis, eye pain, optic neuritis, otitis media, retinal hemorrhage, visual field defect; *Rare:* iritis, keratitis, optic atrophy.

**Urogenital System:** *Infrequent:* urinary urgency, cystitis, menorrhagia, pyelonephritis, urinary retention, kidney calculus, uterine fibroids enlarged, vaginal moniliasis, vaginitis; *Rare:* albuminuria, glycosuria, hematuria, metrorrhagia.

### Post-Market Adverse Drug Reactions

Table 3 includes events determined to be medically significant and/or potentially life threatening and assessed as associated with the use of Tizanidine or the possible relationship to Tizanidine cannot be completely excluded.

**Table 3 – Tizanidine Post-Market Spontaneous Adverse Event Reports**

Adverse Event	Frequency			
	≥1%	<1% and ≥0.1%	<0.1% and ≥0.01%	<0.01%
<b>Blood and lymphatic system disorders</b>				
Agranulocytosis				X
Disseminated intravascular coagulation				X
<b>Cardiac disorders</b>				
Tachycardia				X
Pulmonary oedema				X
Cardiac arrest				X
Torsade de pointes				X
Cardio-respiratory arrest				X
Ventricular fibrillation				X
<b>General disorders and administration site conditions</b>				
Pyrexia				X
<b>Hepatobiliary Disorders</b>				
Hepatic function abnormal				X
Hepatitis				X
Hepatic disorder				X
Jaundice				X
Hepatic failure				X
Hepatic necrosis				X
Hepatitis fulminant				X
Hepatic fibrosis				X
Hepatic cirrhosis				X
<b>Immune system disorders</b>				
Anaphylactic shock				X
<b>Investigations</b>				
Electrocardiogram QT prolonged				X
<b>Musculoskeletal and connective tissue disorders</b>				
Rhabdomyolysis				X
<b>Nervous system disorders</b>				
Loss of consciousness				X
Cerebrovascular accident				X
Cerebral infarction				X
<b>Psychiatric disorders</b>				
Confusional state				X
<b>Renal and urinary disorders</b>				
Renal failure acute				X

Adverse Event	Frequency			
	≥1%	<1% and ≥0.1%	<0.1% and ≥0.01%	<0.01%
Renal failure				X
<b>Respiratory, thoracic and mediastinal disorders</b>				
Dyspnea				X
<b>Skin and subcutaneous tissue disorders</b>				
Erythema multiforme				X
<b>Vascular disorders</b>				
Shock				X
Circulatory collapse				X

## DRUG INTERACTIONS

### Serious Drug Interactions

- **Moderate or potent CYP1A2 inhibitors such as fluvoxamine and ciprofloxacin (see CONTRAINDICATIONS).**

### Overview

*In vitro* studies of cytochrome P450 isoenzymes using human liver microsomes indicate that neither tizanidine nor its major metabolites are likely to affect the metabolism of other drugs metabolized by cytochrome P450 isoenzymes. There are reports of drug interaction of tizanidine and CYP1A2 inhibitors, such as oral contraceptives, fluvoxamine, fluoroquinolones, and others.

### Acetaminophen:

Tizanidine delayed the  $T_{max}$  of acetaminophen by 16 minutes. Acetaminophen did not affect the pharmacokinetics of Tizanidine.

### Alcohol:

Alcohol increased the AUC of Tizanidine by approximately 20% while also increasing its  $C_{max}$  by approximately 15%. This was associated with an increase in side effects of Tizanidine. The CNS depressant effects of Tizanidine and alcohol are additive.

### Oral Contraceptives:

No specific pharmacokinetic study was conducted to investigate interaction between oral contraceptives and tizanidine, but retrospective analysis of population pharmacokinetic data following single and multiple dose administration of 4 mg tizanidine showed that women concurrently taking oral contraceptives had 50% lower clearance of tizanidine than women not on oral contraceptives.

### Antihypertensives:

In placebo-controlled clinical trials, tizanidine has been administered concomitantly with antihypertensive medications in 30 patients. The addition of tizanidine to antihypertensive therapy was associated with a 20-30% increase in the incidence of clinically significant decreases in systolic or diastolic blood pressure compared with both placebo plus antihypertensive (N=36) and tizanidine alone (N=226).

Concurrent use of antihypertensive and tizanidine therapy also resulted in an increase in reports of orthostatic hypotension. Lower initial doses and cautious dose titration should be considered when

APO-TIZANIDINE is to be administered to patients receiving antihypertensive therapy or if antihypertensive therapy is to be initiated in a patient receiving APO-TIZANIDINE.

**Fluvoxamine and other CYP1A2 inhibitors:**

APO-TIZANIDINE should not be used together with moderate and potent CYP1A2 inhibitors such as fluvoxamine and ciprofloxacin (see **CONTRAINDICATIONS**). Concomitant use of tizanidine with fluvoxamine, a potent CYP450 1A2 inhibitor in man, resulted in a 33-fold increase in the tizanidine AUC by fluvoxamine in 10 healthy male subjects. Concomitant use of tizanidine with ciprofloxacin resulted in a 10-fold increase in tizanidine AUC in 10 healthy male subjects. In both studies, clinically significant hypotension resulted along with somnolence, dizziness and decreased psychomotor performance. Co-administration of tizanidine with other inhibitors of CYP1A2 such as antiarrhythmics (amiodarone, mexiletine, propafenone), cimetidine, other fluoroquinolones (norfloxacin, moxifloxacin), oral contraceptives, and ticlopidine should be avoided or used in caution (see also **WARNINGS and PRECAUTIONS, Drug Interaction with CYP1A2 inhibitors**).

**Drugs known to prolong the QT interval:**

Prolongation of the QT interval and bradycardia were noted in chronic toxicity studies in dogs at doses equal to the maximum human dose on a mg/m<sup>2</sup> basis. ECG evaluation was not performed in the controlled clinical studies. There have been post-market reports of QT prolongation and a small number of reports of Torsades de Pointes, none of them fatal, during tizanidine treatment.

**APO-TIZANIDINE should be used with caution in patients taking drugs known to prolong the QT interval.**

**Drug-Food Interactions**

Administering tizanidine with food increases the C<sub>max</sub>, the time to peak concentration, and the extent of absorption of tizanidine (see **ACTION and CLINICAL PHARMACOLOGY**). These pharmacokinetic differences may result in clinically significant differences when switching between fed or fasted states, such as changed incidence of adverse events or delayed/more rapid onset of activity, depending on the nature of the switch (see **DOSAGE and ADMINISTRATION**).

**DOSAGE AND ADMINISTRATION**

**Dosing Considerations**

- Administering tizanidine with food increases the C<sub>max</sub>, the time to peak concentration, and the extent of absorption of tizanidine. These pharmacokinetic differences may result in clinically significant differences when switching between fed or fasted states, such as changed incidence of adverse events or delayed/more rapid onset of activity, depending on the nature of the switch. For this reason, the prescriber should recommend patients to always take tizanidine the same way with regard to fed and fasted state (see **ACTION and CLINICAL PHARMACOLOGY**).
- If therapy needs to be discontinued, especially in patients who have been receiving high doses for long periods, the dose should be decreased slowly to minimize the risk of withdrawal and rebound hypertension, tachycardia, and hypertonia.

**Recommended Dose and Dosage Adjustment**

A single oral dose of 8 mg of APO-TIZANIDINE (tizanidine HCl) reduces muscle tone in patients with spasticity for a period of several hours. The effect peaks at approximately 1 to 2 hours and dissipates between 3 to 6 hours. APO-TIZANIDINE dosing should be scheduled such that the peak effect coincides with activities for which relief of spasticity is most desirable. Effects are dose-related.

Although single doses of less than 8 mg have not been demonstrated to be effective in controlled clinical studies, the dose-related nature of tizanidine's common adverse events, particularly blood pressure reduction, make it prudent to begin treatment with single oral doses of 2 mg. Increase the dose gradually (2 to 4 mg steps) to optimum effect (satisfactory reduction of muscle tone at a tolerated dose).

The dose can be repeated at 6 to 8 hour intervals, as needed, to a maximum of three doses in 24 hours. The total daily dose should not exceed 36 mg.

Experience with single doses exceeding 8 mg and daily doses exceeding 24 mg is limited. There is essentially no experience with repeated, single, daytime doses greater than 12 mg or total daily doses greater than 36 mg (see **WARNINGS AND PRECAUTIONS**).

## **OVERDOSAGE**

For management of a suspected drug overdose, contact your regional Poison Control Centre.

There have been cases of tizanidine overdose reported in post-marketing experience. Most of these were intentional overdoses, about a quarter have resulted in fatality, and in at least half of these cases, other CNS depressants were involved. The majority of cases involved depressed consciousness (somnolence, stupor, or coma), depressed cardiovascular function (bradycardia, hypotension), and depressed respiratory function (respiratory depression or failure).

Should overdosage occur, basic steps to ensure the adequacy of an airway and the monitoring of cardiovascular and respiratory systems should be undertaken. For the most recent information concerning the management of overdose, contact a poison control center.

## **ACTION AND CLINICAL PHARMACOLOGY**

### **Mechanism of Action**

Tizanidine is an agonist at  $\alpha_2$ -adrenergic receptor sites and presumably reduces spasticity by increasing presynaptic inhibition of motor neurons. In animal models, tizanidine has no direct effect on skeletal muscle fibers or the neuromuscular junction, and no major effect on monosynaptic spinal reflexes. The effects of tizanidine are greatest on polysynaptic pathways.

The overall effect of these actions is thought to reduce facilitation of spinal motor neurons.

The imidazoline chemical structure of tizanidine is related to that of the anti-hypertensive drug clonidine and other  $\alpha_2$ -adrenergic agonists. Pharmacological studies in animals show similarities

between the two compounds, but tizanidine was found to have one-tenth to one-fiftieth (1/50) of the potency of clonidine in lowering blood pressure.

### **Pharmacokinetics**

#### **Absorption:**

Following oral administration, tizanidine is essentially completely absorbed and has a half-life of approximately 2 hours. Following administration of tizanidine peak plasma concentrations occurred at approximately 1 hour after dosing. Food increases the mean  $C_{max}$  by approximately 30% and increases the median time to peak concentration by approximately 25 minutes, from 1 hour to 1 hour and 25 minutes. Food increases the extent of absorption of tizanidine by approximately 30%. Tizanidine has linear pharmacokinetics over a dose of 1 to 20 mg.

#### **Metabolism:**

The absolute oral bioavailability of tizanidine is approximately 40%, due to extensive first-pass metabolism in the liver; approximately 95% of an administered dose is metabolized. Tizanidine metabolites are not known to be active; their half-lives range from 20 to 40 hours.

#### **Distribution:**

Tizanidine is widely distributed throughout the body; mean steady state volume of distribution is 2.4 L/kg following intravenous administration in healthy adult volunteers. Tizanidine is approximately 30% bound to plasma proteins, independent of concentration over the therapeutic range.

#### **Excretion:**

Following single and multiple oral dosing of  $^{14}C$ -tizanidine, an average of 60% and 20% of total radioactivity was recovered in the urine and feces, respectively.

### **Special Populations and Conditions**

**Age Effects:** No specific pharmacokinetic study was conducted to investigate age effects. Cross study comparison of pharmacokinetic data, following single dose administration of 6 mg tizanidine showed that younger subjects cleared the drug four times faster than the elderly subjects. Tizanidine has not been evaluated in children (see **WARNINGS AND PRECAUTIONS**).

**Gender:** No specific pharmacokinetic study was conducted to investigate gender effects. Retrospective analysis of pharmacokinetic data, however, following single and multiple dose administration of 4 mg tizanidine showed that gender had no effect on the pharmacokinetics of tizanidine.

**Race:** Pharmacokinetic differences due to race have not been studied.

**Hepatic Insufficiency:** Pharmacokinetic differences due to hepatic impairment have not been studied (see **WARNINGS AND PRECAUTIONS**).

**Renal Insufficiency:** Tizanidine clearance is reduced by more than 50% in elderly patients with renal insufficiency (creatinine clearance < 25 mL/min) compared to healthy elderly subjects; this

would be expected to lead to a longer duration of clinical effect. Tizanidine should be used with caution in renally impaired patients (see **WARNINGS AND PRECAUTIONS**).

## **STORAGE AND STABILITY**

Stored at room temperature 15 - 30°C (59 - 86°F).

## **DOSAGE FORMS, COMPOSITION AND PACKAGING**

APO-TIZANIDINE 2 mg: each white to off-white, round tablet, engraved “APO” over “TI-2” on one side, plain with a bisect score on the other, contains tizanidine hydrochloride equivalent to 2 mg tizanidine. Available in bottles of 100.

APO-TIZANIDINE 4 mg: each white to off-white, round tablet, engraved “APO” over “TI-4” on one side, plain with a quadrisect score on the other, contains tizanidine hydrochloride equivalent to 4 mg tizanidine. Available in bottles of 100.

### **Tablet Composition**

APO-TIZANIDINE (tizanidine hydrochloride) tablets are composed of the active ingredient, tizanidine hydrochloride (2.29 mg equivalent to 2 mg tizanidine base or 4.58 mg equivalent to 4 mg tizanidine base), and the non-medicinal ingredients, colloidal silicon dioxide, stearic acid, microcrystalline cellulose and anhydrous lactose.

APO-TIZANIDINE meets USP Dissolution Test 2.

## PART II: SCIENTIFIC INFORMATION

### PHARMACEUTICAL INFORMATION


#### Drug Substance

Proper Name: tizanidine hydrochloride (USAN)

Chemical Name: 5-chloro-4-(2-imidazolylamino)-2,1,3-benzothiazole hydrochloride

Molecular formula and molecular weight: C<sub>9</sub>H<sub>9</sub>Cl<sub>2</sub>N<sub>5</sub>S

Structural Formula:



Physicochemical properties:

Appearance: Almost white to slightly yellow crystalline powder, odorless or faint characteristic odor

Solubility: Approximately 5% soluble in water and methanol; solubility in water decreases as the pH increases

pK<sub>a</sub> value: 7.35 determined potentiometrically

pH: 3.50 - 5.00

Partition coefficient: 3.6:1

Melting point: 286 – 290°C

## CLINICAL TRIALS

### Comparative Bioavailability Studies

A comparative bioavailability study was performed using healthy adult male volunteers. The rate and extent of absorption of tizanidine was measured and compared following administration of 4 mg tizanidine (one tablet of either APO-TIZANIDINE or Zanaflex®). The results from measured data are summarized as follows:

**Table 4: Summary Table of the Comparative Bioavailability Data**

Summary Table of the Comparative Bioavailability Data Tizanidine (Dose: 1 x 4 mg) From Measured Data – Under Fasting Conditions				
Parameter	Geometric Mean Arithmetic Mean (CV%)		Ratio of Geometric Means (%)**	90% Geometric Confidence Intervals
	APO-TIZANIDINE	Zanaflex®†		
AUC <sub>T</sub> (ng•hr/mL)	5.62	5.54	101.4	(92.4 - 111.1)
	6.18 (43)	6.08 (43)		
AUC <sub>I</sub> (ng•hr/mL)	5.79	5.70	101.6	(92.9 - 111.2)
	6.35 (42)	6.24 (42)		
C <sub>max</sub> (ng/mL)	2.26	2.21	102.3	(90.9 - 115.1)
	2.51 (47)	2.40 (40)		
T <sub>max</sub> (hr)*	1.14 (45)	1.09 (52)	-	
t <sub>1/2</sub> (hr)*	1.41 (18)	1.33 (18)	-	

\* Arithmetic means (CV%).  
 \*\* Based on the least squares estimate.  
 † Zanaflex® was manufactured by Novartis Pharma AG, Basel, Switzerland for Elan Pharmaceuticals Inc., South San Francisco, CA 94080 USA, distributed by Draxis Health Inc., and was purchased in Canada.


The capacity of tizanidine to reduce increased muscle tone associated with spasticity was demonstrated in two adequate and well controlled studies in patients with multiple sclerosis or spinal injury.

In one study, patients with multiple sclerosis were randomized to receive single oral doses of drug or placebo.<sup>4</sup> Patients and assessors were blind to treatment assignment and efforts were made to reduce the likelihood that assessors would become aware indirectly of treatment assignment (e.g., they did not provide direct care to patients and were prohibited from asking questions about side effects). In all, 140 patients received either placebo, 8 mg or 16 mg of tizanidine.

Response was assessed by physical examination; muscle tone was rated on a 5 point scale (Ashworth score), with a score of 0 used to describe normal muscle tone. A score of 1 indicated a slight spastic catch while a score of 2 indicated more marked muscle resistance. A score of 3 was used to describe considerable increase in tone, making passive movement difficult. A muscle immobilized by spasticity was given a score of 4. Spasm counts were also collected.

Assessments were made at 1, 2, 3 and 6 hours after treatment. A statistically significant reduction of the Ashworth score for tizanidine compared to placebo was detected at 1, 2 and 3 hours after treatment. Figure 1 below shows a comparison of the mean change in muscle tone from baseline as measured by the Ashworth scale. The greatest reduction in muscle tone was 1 to 2 hours after treatment. By 6 hours after treatment, muscle tone in the 8 and 16 mg tizanidine groups was indistinguishable from muscle tone in placebo treated patients. Within a given patient, improvement in muscle tone was correlated with plasma concentration. Plasma concentrations were variable from patient to patient at a given dose. Although 16 mg produced a larger effect, adverse events including hypotension were more common and more severe than in the 8 mg group. There were no differences in the number of spasms occurring in each group.

**FIGURE 1: Single Dose Study - Mean Change in Muscle Tone from Baseline as Measured by the Ashworth Scale  $\pm$  95% Confidence Interval**  
 (A Negative Ashworth Score Signifies an Improvement in Muscle Tone from Baseline)




In a multiple dose study, 118 patients with spasticity secondary to spinal cord injury were randomized to either placebo or tizanidine.<sup>5</sup> Steps similar to those taken in the first study were employed to ensure the integrity of blinding.

Patients were titrated over 3 weeks up to a maximum tolerated dose or 36 mg daily given in three unequal doses (e.g., 10 mg given in the morning and afternoon and 16 mg given at night). Patients were then maintained on their maximally tolerated dose for 4 additional weeks (i.e., maintenance phase). Throughout the maintenance phase, muscle tone was assessed on the Ashworth scale within a period of 2.5 hours following either the morning or afternoon dose and counts of spasms were collected by patient diary.


At endpoint (the protocol specified time of outcome assessment), there were statistically significant reductions in muscle tone and spasms in the tizanidine treated group compared to placebo. The reduction in muscle tone was not associated with a reduction in muscle strength (a desirable outcome) but also did not lead to any consistent advantage of tizanidine treated patients on

measures of activities of daily living. Figures 2 and 3 below show a comparison of the mean change in muscle tone from baseline as measured by the Ashworth scale and a comparison of the mean change in daytime spasms as recorded in patient diaries, respectively.

**FIGURE 2: Multiple Dose Study - Mean Change in Muscle Tone 0.5-2.5 Hours after Dosing as Measured by the Ashworth Scale  $\pm$  95% Confidence Interval (A Negative Ashworth Score Signifies an Improvement in Muscle Tone from Baseline)**



**FIGURE 3: Multiple Dose Study - Mean Change in Response Ratio of Daytime Spasms  $\pm$  95% Confidence Interval (A Negative Response Ratio Signifies an Improvement in Spasms from Baseline)**



In a second multiple dose study, 187 patients with spasticity secondary to multiple sclerosis were randomized to either placebo or tizanidine.<sup>6</sup> Patients were titrated over 3 weeks up to a maximum tolerated dose or 36 mg daily given in three equal doses. Patients were then maintained on their maximally tolerated dose for 9 additional weeks (i.e., maintenance phase).

Throughout the maintenance phase, muscle tone was assessed on the Ashworth scale and global efficacy was assessed by both patient and investigator. There was a statistically significant reduction in muscle tone in the tizanidine treated group as compared to placebo at the last maintenance phase measurement of muscle tone (the protocol specified time of outcome assessment) and throughout the maintenance phase. The reduction in muscle tone was not associated with a reduction in muscle strength.

## DETAILED PHARMACOLOGY

In calf cerebral cortex, tizanidine exhibited a selective affinity for  $\alpha_2$ -adrenergic receptors ( $IC_{50}=2.1$  nM) over  $\alpha_1$  receptors ( $IC_{50}= 4,687$  nM). In rat kidney membrane, tizanidine exhibited a higher affinity for imidazoline receptors ( $K_i= 4$  nM) than for  $\alpha_2$ -adrenergic receptors ( $K_i= 91$  nM).

The  $\alpha_2$ -adrenoreceptor sub-type specificity of tizanidine was investigated using tissues and cell lines which express only one of the three receptor sub-types,  $\alpha_2A$ -,  $\alpha_2B$ -, and  $\alpha_2C$ . The  $K_i$  values for tizanidine were 65, 167, and 107 nM at the  $\alpha_2A$ -,  $\alpha_2B$ -, and  $\alpha_2C$ -receptors, respectively.

Tizanidine at concentration of 3 nM to 1  $\mu$ M exhibited  $\alpha_2$ -agonist activity in two *in vitro* peripheral smooth muscle tissues, rat vas deferens and guinea pig ileum. Tizanidine was demonstrated to inhibit the contractile response to electrical stimulation, an effect which was antagonized by the  $\alpha_2$ -antagonist, yohimbine, but not by the  $\alpha_1$ -antagonist, prazosin.

Unlike dantrolene, tizanidine showed little or no activity in two models for demonstration of direct effects on muscle (direct stimulation of the peroneal nerve in chloralose-anaesthetized cats and of an *in vitro* rat diaphragm-phrenic nerve preparation).

Tizanidine was, however, found to be active in whole animal models for effects on reflex activity. In conscious rabbits, tizanidine inhibited hind limb tonic stretch reflex activity with a potency which was 12 times greater than that of diazepam and 16 times greater than that of baclofen. The inhibitory effect of tizanidine on the extensor reflex was maximal at 5 minutes and exhibited a duration of activity with a half-life of 21 minutes. Tolerance to this effect did not develop over a 14-day course of treatment.

The flexor reflex in rats was used as a model for polysynaptic reflex. In intact, anaesthetized rats and unanaesthetized decerebrate rats, tizanidine ablated the electromyogram response to flexor reflex stimulation, an effect which could be prevented by pretreatment with the  $\alpha_2$ -antagonist, yohimbine. However, in acutely spinalized rats, tizanidine was observed to facilitate reflex activity, an effect which was reversible in the presence of the  $\alpha_1$ -antagonist, prazosin, but not the  $\alpha_2$ -antagonist, yohimbine.

Tizanidine exhibited activity in a model of droperidol/fentanyl-induced rigidity in rats.

Tizanidine was more potent in this regard than either baclofen or diazepam (ED50 values for tizanidine, diazepam, and baclofen were 0.5, 4, and 5.2 mg/kg p.o., respectively).

Tizanidine, baclofen, and diazepam were likewise efficacious in abolishing the rigidity observed in a decerebrate model of gamma-rigidity in rats (ED50 values of 0.5 mg/kg, 1.3 mg/kg, and 3.8 mg/kg i.v.). The two major metabolites of tizanidine identified in human blood and urine, metabolites 3 and 4, were without pharmacological activity in this model.

Reduced motor activity, disturbed gait, and ataxia were observed in mice after 10 mg/kg oral doses of tizanidine and in rats after 2.5 mg/kg oral doses of tizanidine. These doses were 5 to 20 fold higher than the oral ED50 for inhibition of droperidol/fentanyl-induced muscle rigidity in rats (0.5 mg/kg).

Sedative effects were observed in rhesus monkeys at oral doses of 6 to 10 mg/kg and in baboons after oral doses of 8 to 16 mg/kg and an intravenous dose of 1 mg/kg.

In normotensive rats, 0.3 and 1.0 mg/kg oral doses of tizanidine were associated with an initial small decrease in blood pressure (maximum 12%), followed by an increase in blood pressure at 1.5 to 7 hours (maximum 12%). At an oral dose of 3.0 mg/kg, statistically significant decreases in both systemic blood pressure and heart rate were observed (approximately 13% decrease) which were sustained for 5 and 2 hours, respectively. In a hypertensive rat model, an oral dose of 5 mg/kg tizanidine was associated with a maximum 25% reduction in mean arterial blood pressure.

In both anaesthetized rats and dogs, intravenously administered tizanidine at doses of 0.03 and 0.1 mg/kg produced an initial, transient pressor response, followed by sustained hypotension and bradycardia. The  $\alpha_1$ - and  $\alpha_2$ -adrenergic antagonist, phenoxybenzamine, inhibited the initial pressor response to tizanidine in anaesthetized rats. In anaesthetized cats, tizanidine at a cumulative intravenous dose of 0.7 mg/kg, decreased blood pressure by a maximum of 35%.

Tizanidine appears to influence blood pressure through  $\alpha$ -adrenoreceptor-mediated vasoconstriction in the peripheral vasculature and by activities on the centrally located  $\alpha$ -adrenoreceptors regulating peripheral sympathetic activity.

## **TOXICOLOGY**

Slight, reversible increases in SGPT/ALT were observed after the oral administration of tizanidine in subchronic (13 week) studies in rats and dogs and a chronic (52 week) study in dogs. Mild, reversible histopathological changes in the liver were also reported in the subchronic studies in rats and dogs at doses representing about 55 and 6 times the maximum recommended human dose, respectively. For this and all subsequent comparisons the reference weight for a small human is 50 kg.

In the 13-week subchronic toxicity studies, the maximum dose at which no toxic effect was observed was 1.7 mg/kg in rats and 0.3 mg/kg in dogs.

In dogs receiving doses of 1.5 mg/kg during the 52-week chronic toxicity study, the heart rate was reduced by 40% and the QT interval was prolonged. Slight sedation or unstable equilibrium,

emesis, and salivation were also observed at this dose. The maximum dose at which no toxic effects were observed in this study was 0.45 mg/kg.

### **Teratology Studies**

Reproduction studies in rats performed at oral doses of tizanidine up to approximately 4 times the maximum recommended human dose did not provide evidence of embryotoxic or teratogenic effects. However, at dose levels producing exaggerated pharmacodynamic effects in rats (from 5 to over 130 times the maximum recommended human dose), the duration of pregnancy was prolonged, prenatal and perinatal mortality was increased, and development retardation (reduced fetal weight, increased incidence of skeletal retardation) was observed.

Tizanidine has been found to pass into the milk of nursing rats with a milk to blood concentration ratio of 1.8:1. In young nursing rats, abnormal results were obtained in tests indicative of central nervous system function. Several changes were noted in the development of the pups that may have been attributable either to the toxicity of the drug on the dams or else to direct drug effects on the offspring, following ingestion through milk.

Reproduction studies in rabbits with tizanidine doses over 40 times the maximum recommended human dose showed no evidence of embryotoxicity or teratogenicity.

### **Mutagenesis**

Tizanidine was not mutagenic or clastogenic in the following in vitro assays: the bacterial Ames test and the mammalian gene mutation test and chromosomal aberration test in Chinese hamster cells.

It was also negative in the following in vivo assays: the bone marrow micronucleus test in mice, the bone marrow micronucleus and cytogenicity test in Chinese hamsters, the dominant lethal mutagenicity test in mice, and the unscheduled DNA synthesis (UDS) test in mice.

### **Carcinogenesis**

No evidence for carcinogenicity was seen in two dietary studies in rodents. Tizanidine was administered to mice for 78 weeks at doses up to 16 mg/kg, which is equivalent to 2 times the maximum recommended human dose on a mg/m<sup>2</sup> basis. Tizanidine was also administered to rats for 104 weeks at doses up to 9 mg/kg which is equivalent to 2.5 times the maximum recommended human dose on a mg/m<sup>2</sup> basis. There was no statistically significant increase in tumors in either species.

### **Impairment of Fertility**

Tizanidine did not affect fertility in male rats at doses of 10 mg/kg, approximately 2.7 times the maximum recommended human dose on a mg/m<sup>2</sup> basis and in females at doses of 3 mg/kg, approximately equal to the maximum recommended human dose on a mg/m<sup>2</sup> basis; fertility was reduced in males receiving 30 mg/kg (8 times the maximum recommended human dose on a mg/m<sup>2</sup> basis) and in females receiving 10 mg/kg (2.7 times the maximum recommended human dose on a mg/m<sup>2</sup> basis). At these doses, maternal behavioral effects and clinical signs were observed including marked sedation, weight loss, and ataxia.

## REFERENCES

- 1) Coward DM. Tizanidine: Neuropharmacology and mechanism of action. *Neurology* 1994; 44 (Suppl 9): 6-11.
- 2) Coward DM, et al. *Pharmacological Properties of Tizanidine (DS 103-282)*. Stuttgart, New York: FK Schattauer Verlag; 1994: 61-71.
- 3) Delwaide PJ, et al. Tizanidine and electrophysiologic analysis of spinal control mechanisms in humans with spasticity. *Neurology* 1994; 44 (Suppl 9) : 21-28.
- 4) Nance PW, et al. Relationship of the anti-spasticity effect of tizanidine to plasma concentration in patients with multiple sclerosis. *Archives of Neurology* 1997; 54: 731-736.
- 5) Nance PW, et al. Efficacy and safety of tizanidine in the treatment of spasticity in patients with spinal cord injury. *Neurology* 1994; 44 (Suppl 9): 44-52.
- 6) Barnes MP, et al. A double-blind, placebo-controlled trial of tizanidine in the treatment of spasticity caused by multiple sclerosis. *Neurology* 1994; 44 (Suppl 9): 70-78.
- 7) Product Monograph - ZANAFLEX® 2 and 4 mg Tablets. Paladin Labs Inc. Date of Preparation July 7, 2009 (Control Number: 131114).

## IMPORTANT: PLEASE READ

### PART III: CONSUMER INFORMATION

Pr APO-TIZANIDINE

#### Tizanidine Hydrochloride Tablets USP

This leaflet is part III of a three-part “Product Monograph” published when APO-TIZANIDINE was approved for sale in Canada and is designed specifically for Consumers. This leaflet is a summary and will not tell you everything about APO-TIZANIDINE. Contact your doctor or pharmacist if you have any questions about the drug.

### ABOUT THIS MEDICATION

#### What the medication is used for:

APO-TIZANIDINE (tizanidine hydrochloride) is a drug used to reduce the spasticity which may be caused by medical conditions such as spinal cord injury or multiple sclerosis.

#### What it does:

Tizanidine presumably reduces spasticity by decreasing the activity of nerves that cause muscle spasticity. The overall effect is muscle relaxation.

#### When it should not be used:

Do not use APO-TIZANIDINE if you:

- are allergic to it or to any of the nonmedicinal ingredients (see below). Stop taking the drug and contact your doctor immediately if you experience an allergic reaction or any severe or unusual side effects.
- are currently taking fluvoxamine or ciprofloxacin.
- need spasticity to maintain upright posture and balance for your movements.

#### What the medicinal ingredient is:

The medicinal ingredient in APO-TIZANIDINE is: Tizanidine hydrochloride

#### What the important nonmedicinal ingredients are:

The non-medicinal ingredients in APO-TIZANIDINE are: colloidal silicon dioxide, stearic acid, microcrystalline cellulose and anhydrous lactose.

#### What dosage forms it comes in:

2 mg and 4 mg tablets

### WARNINGS AND PRECAUTIONS

BEFORE you use APO-TIZANIDINE, talk to your doctor or pharmacist if any of the following conditions apply to you:

- history of an unusual or allergic reaction to APO-TIZANIDINE or to any other substances, such as foods, preservatives or dyes
- pregnancy or breast-feeding

- liver disease
- kidney disease
- low blood pressure
- presently receiving treatment with antihypertensives (high blood pressure medicine), oral contraceptives (birth control pills), fluvoxamine, ciprofloxacin, or phenytoin.

Do not drive or operate hazardous machinery if you are unsure how this medication will affect you or if you experience drowsiness, coordination problems, or blurred vision.

It is important to tell your doctor or pharmacist about all medicines that you are taking including other medicines that a doctor has prescribed, medicines that you buy yourself without a prescription, and any herbal remedies that you are taking. While on APO-TIZANIDINE, do not start taking a new medicine or herbal remedy before checking with your doctor.

### INTERACTIONS WITH THIS MEDICATION

Do not use APO-TIZANIDINE if you are taking fluvoxamine or ciprofloxacin. Alcohol and other central nervous system depressants (medicines causing drowsiness or decreased alertness such as antihistamines, sedatives, tranquilizers, sleeping medicines, prescription pain medicines, narcotics, anti-epileptics, other muscle relaxants, and anaesthetics) may increase the drowsiness experienced with APO-TIZANIDINE.

Other drugs that may interact with APO-TIZANIDINE include: acetaminophen, oral contraceptives, blood pressure lowering drugs, antiarrhythmic heart medications (e.g. amiodarone, mexiletine, propafenone), cimetidine (to reduce stomach acid), fluoroquinolone-type antibiotics (e.g. norfloxacin, moxifloxacin), and ticlopidine (to prevent blood clotting).

### PROPER USE OF THIS MEDICATION

#### Usual dose:

Take APO-TIZANIDINE as directed by your doctor. Do not take larger or more frequent doses than your doctor has prescribed. During the first weeks of treatment, your doctor may want to adjust your dose to meet your individual needs.

If you need to stop taking APO-TIZANIDINE, follow your doctor's instruction on how to gradually reduce your dose.

**IMPORTANT: PLEASE READ**

Food can affect the absorption of APO-TIZANIDINE in your body - so it is important to take it in the same way all the time. For example, if you normally take APO-TIZANIDINE with a meal, you should take it with food each time. If you normally take APO-TIZANIDINE on an empty stomach, then you should always take it before you eat.

**Overdose:**

The majority of the cases of tizanidine overdose reported in postmarketing experience involved sleepiness, very long or deep sleeplike state from which a person can be awakened only briefly by vigorous stimulation (stupor) or cannot be at all (coma), heart beat at an unusually slow rate, fall in blood pressure, and difficulty breathing.

In case of drug overdose, contact a health care practitioner, hospital emergency department or regional Poison Control Centre immediately, even if there are no symptoms.

**SIDE EFFECTS AND WHAT TO DO ABOUT THEM**

APO-TIZANIDINE may cause dry mouth, drowsiness, weakness, fatigue, hallucinations, or dizziness in some patients.

APO-TIZANIDINE may cause a decrease in your blood pressure. In some cases, decreased blood pressure may result in dizziness, lightheadedness, or fainting when a person rises suddenly from a sitting or lying position. Rising slowly may decrease the risk of these problems.

Rare cases of liver damage have been reported in patients receiving APO-TIZANIDINE. Monitoring of liver function tests (ALT/SGPT and AST/SGOT) during treatment with APO-TIZANIDINE may be required by your doctor.

Serious side effects and what to do about them				
Symptom / effect		Talk with your doctor or pharmacist		Stop taking drug and immediate medical help
		Only if severe	In all cases	
	appetite combined with itching, yellowing of the skin or eyes, dark urine]			
	Allergic reactions (red and lumpy skin, rash, hives, swelling, trouble breathing)			✓*
	Breakdown of muscle fibers [symptoms include abnormal urine color (dark, red, or cola colored), muscle tenderness, weakness of the affected muscle(s), generalized weakness or muscle stiffness or aching]		✓	

\*If you think you have these side effects, it is important that you seek medical advice from your doctor straight away.

*This is not a complete list of side effects. For any unexpected effects while taking APO-TIZANIDINE, contact your doctor or pharmacist.*

**HOW TO STORE IT**

APO-TIZANIDINE should be stored out of the reach of children and away from heat and direct light.

Serious side effects and what to do about them				
Symptom / effect		Talk with your doctor or pharmacist		Stop taking drug and immediate medical help
		Only if severe	In all cases	
Common	Hallucination		✓	
	Hypotension (feeling dizzy, lightheaded, or faint)		✓	
Uncommon	Liver disorder [symptoms include nausea, vomiting, loss of		✓*	

### **Reporting Side Effects**

You can report any suspected side effects associated with the use of health products to health Canada by:

- Visiting the Web page on Adverse Reaction Reporting (<https://www.canada.ca/en/health-canada/services/drugs-health-products/medeffect-canada/adverse-reaction-reporting.html>) for information on how to report online, by mail or by fax; or
- Calling toll-free at 1-866-234-2345

*NOTE: Contact your health professional if you need information about how to manage your side effects. The Canada Vigilance Program does not provide medical advice.*

### **MORE INFORMATION**

- Talk to your healthcare professional
- Find the full product monograph that is prepared for healthcare professionals and includes this Consumer Information by visiting the Health Canada website: (<https://www.canada.ca/en/health-canada/services/drugs-health-products/drug-products/drug-product-database.html>); the manufacturer's website <http://www.apotex.ca/products>, or by calling 1-800-667-4708

This leaflet was prepared by Apotex Inc., Toronto, Ontario, M9L 1T9.

Last revised: October 25, 2023