

PRODUCT MONOGRAPH
INCLUDING PATIENT MEDICATION INFORMATION

Pr WAYMADE-TRIENTINE

Trientine Hydrochloride Capsules

Capsules, 250 mg, Oral

USP

Copper-chelating agent

Waymade PLC
Sovereign House
Miles Gray Road
Basildon, Essex
United Kingdom SS14 3FR

Date of Initial Authorization:
April 19, 2021

Date of Revision:
November 14, 2023

Canadian Importer:
Waymade Canada Inc.
4150 Sainte-Catherine West
Montreal, QC
H3Z 2Y5

Submission Control Number: 276415

RECENT MAJOR LABEL CHANGES

Not applicable.

TABLE OF CONTENTS

Sections or subsections that are not applicable at the time of authorization are not listed.

RECENT MAJOR LABEL CHANGES	2
TABLE OF CONTENTS	2
PART I: HEALTH PROFESSIONAL INFORMATION.....	4
1 INDICATIONS.....	4
1.1 Pediatrics.....	4
1.2 Geriatrics	4
2 CONTRAINDICATIONS	4
3 SERIOUS WARNINGS AND PRECAUTIONS BOX	4
4 DOSAGE AND ADMINISTRATION	5
4.1 Dosing Considerations.....	5
4.2 Recommended Dose and Dosage Adjustment	5
4.4 Administration.....	5
4.5 Missed Dose	6
5 OVERDOSAGE.....	6
6 DOSAGE FORMS, STRENGTHS, COMPOSITION AND PACKAGING	6
7 WARNINGS AND PRECAUTIONS	6
7.1 Special Populations.....	8
7.1.1 Pregnant women	8
7.1.2 Breast-feeding	8
7.1.3 Pediatrics	8
7.1.4 Geriatrics	8
8 ADVERSE REACTIONS	8
8.1 Adverse Reaction Overview.....	8
8.2 Clinical Trial Adverse Reactions	9
8.3 Post-Market Adverse Reactions	9
9 DRUG INTERACTIONS	9

9.4	Drug-Drug Interactions.....	9
9.5	Drug-Food Interactions.....	9
9.6	Drug-Herb Interactions.....	9
9.7	Drug-Laboratory Test Interactions.....	9
10	CLINICAL PHARMACOLOGY	9
10.1	Mechanism of Action.....	9
10.2	Pharmacodynamics	10
10.3	Pharmacokinetics	10
11	STORAGE, STABILITY AND DISPOSAL	11
12	SPECIAL HANDLING INSTRUCTIONS	12
PART II: SCIENTIFIC INFORMATION		13
13	PHARMACEUTICAL INFORMATION	13
14	CLINICAL TRIALS	13
14.1	Clinical Trials by Indication	13
	The Treatment Of Patients With Wilson's Disease Who Are Intolerant To Penicillamine.	
	13
14.2	Comparative Bioavailability Studies.....	15
15	MICROBIOLOGY	15
16	NON-CLINICAL TOXICOLOGY	16
PATIENT MEDICATION INFORMATION.....		17

PART I: HEALTH PROFESSIONAL INFORMATION

1 INDICATIONS

WAYMADE-TRIENTINE (trientine hydrochloride capsules) is indicated for:

- The treatment of patients with Wilson's Disease who are intolerant to penicillamine.

WAYMADE-TRIENTINE should only be initiated by physicians experienced in the management of Wilson's Disease.

1.1 Pediatrics

Pediatrics (5-17 years of age): Controlled studies of the safety and effectiveness of trientine hydrochloride in pediatric patients have not been conducted. Data are lacking to establish the safety and effectiveness of trientine hydrochloride in children less than 5 years of age.

Pediatrics (<5 years of age): Based on the data submitted and reviewed by Health Canada, the safety and efficacy of WAYMADE-TRIENTINE in these pediatric patients have not been established; therefore, Health Canada has not authorized an indication for children less than 5 years of age.

1.2 Geriatrics

Geriatrics (> 65 years of age): Clinical studies of trientine hydrochloride did not include sufficient numbers of patients aged 65 and over to determine whether they respond differently than patients less than 65 years of age. WAYMADE-TRIENTINE should be used with caution in geriatric patients (see [4.2 Recommended Dose and Dosage Adjustment](#) and [7.1.4 Geriatrics](#)).

2 CONTRAINDICATIONS

Trentine hydrochloride is contraindicated in:

- Patients who are hypersensitive to this drug or to any ingredient in the formulation, including any non-medicinal ingredient, or component of the container. For a complete listing, see [6 DOSAGE FORMS, STRENGTHS, COMPOSITION AND PACKAGING](#).

3 SERIOUS WARNINGS AND PRECAUTIONS BOX

Serious Warnings and Precautions

- WAYMADE-TRIENTINE (trientine hydrochloride) should only be initiated by physicians experienced in the management of Wilson's Disease.
- Worsening of neurologic or neurocognitive functioning, which can be irreversible, may occur in patients with pre-existing neurologic and/or neuropsychiatric impairment due to Wilson's Disease and treated with WAYMADE-TRIENTINE (see [7 WARNINGS AND PRECAUTIONS, Neurologic](#)).

4 DOSAGE AND ADMINISTRATION

4.1 Dosing Considerations

- The starting dose of WAYMADE-TRIENTINE would usually be the lowest recommended dose. The dose should subsequently be adjusted according to the patient's clinical response.
- Patients receiving WAYMADE-TRIENTINE require regular and ongoing monitoring of their clinical condition and response to treatment throughout the period of drug administration (see [General](#)).
- Overtreatment carries the risk of copper deficiency. Ongoing regular monitoring for manifestations of overtreatment should be undertaken (see [General](#) and [Monitoring and Laboratory Tests](#)).
- Studies in animals have shown that trientine hydrochloride is teratogenic (see [16 NON-CLINICAL TOXICOLOGY, Reproductive and Developmental Toxicology](#)). Some data have also shown that trientine hydrochloride has a genotoxic potential (see [16 NON-CLINICAL TOXICOLOGY, Genotoxicity](#)). If used during pregnancy, serum free copper concentrations should be closely monitored and a dosage adjustment to the lowest dose of WAYMADE-TRIENTINE that maintains free copper concentrations within acceptable limits should be considered (see [7.1.1 Pregnant women](#)).
- If iron deficiency develops and iron administration is required, doses of WAYMADE-TRIENTINE and iron should be separated by at least two hours (see [Hematologic](#) and [9.4 Drug-Drug Interactions](#)).
- WAYMADE-TRIENTINE should be taken at least one hour apart from any other drug, food, or milk (see [9.5 Drug-Food Interactions](#)).

4.2 Recommended Dose and Dosage Adjustment

Adults: The recommended starting dose of WAYMADE-TRIENTINE is 750 mg/day, given in divided doses 2 to 4 times a day. Subsequently, if required, the daily dose may be increased gradually to a maximum of 2,000 mg/day. Most adult patients can be maintained at a daily dose of 1,250 mg/day or less.

Pediatrics (5-17 years old): The recommended starting dose is 500 mg/day, given in divided doses 2 to 4 times a day. Subsequently, if required, the daily dose may be increased gradually to a maximum of 1,500 mg/day for patients aged 12 years or under.

The daily dose of WAYMADE-TRIENTINE should be increased only when the clinical response is not adequate or the concentration of serum free copper is persistently above 3.1 mcmol/L (20 mcg/dL). The optimal long-term maintenance dosage should generally be determined at 6 to 12 month intervals.

Geriatrics (> 65 years of age): Dose selection should be cautious in patients over the age of 65 years, usually starting at the low end of the dosing range (see [7.1.4 Geriatrics](#)).

Hepatic impairment: Clinical studies evaluating patients with hepatic impairment have not been conducted. No dosage adjustment is required, however it is advised that patients be monitored every two to three weeks after initiation of treatment with WAYMADE-TRIENTINE.

Renal impairment: Clinical studies evaluating patients with renal impairment have not been conducted. There is limited information in patients with renal impairment. No dosage adjustment is required in these patients.

4.4 Administration

WAYMADE-TRIENTINE should be given on an empty stomach, at least one hour before meals or two hours after meals and at least one hour apart from any other drug, food, or milk. The capsules should

be swallowed whole with water and should not be opened or chewed (see [9.4 Drug-Drug Interactions](#) and [9.5 Drug-Food Interactions](#)).

4.5 Missed Dose

A missed dose should be administered as soon as possible. However, two doses of WAYMADE-TRIENTINE should not be taken at the same time or very close to each other.

If you missed a dose of this medication, take it as soon as you remember. But if it is almost time for your next dose, skip the missed dose and continue with your next scheduled dose. Go back to the regular dosing schedule. Do not take two doses at the same time.

5 OVERDOSAGE

There is a single report of an adult woman who ingested 30 grams of trientine hydrochloride without apparent ill effects. No other data on overdosage are available.

For management of a suspected drug overdose, contact your regional poison control centre.

6 DOSAGE FORMS, STRENGTHS, COMPOSITION AND PACKAGING

Table 1: Dosage Forms, Strengths, Composition and Packaging

Route of Administration	Dosage Form/ Strength/Composition	Non-medicinal Ingredients
Oral	Capsules, 250 mg	Black ink, gelatin, purified water, stearic acid, sunset yellow FCF, titanium dioxide

Description

Cylindrical size “1” hard gelatin capsule with opaque orange colour cap printed with “NAV” in black ink and opaque white colour body printed with “101” in black ink. Capsule filled with white to off white colour powder.

Primary packaging system for trientine hydrochloride capsules 250 mg, is 150 cc white HDPE round bottle, 38 mm screw cap, and 1 g silica gel as desiccant. Pack size is 100's count.

7 WARNINGS AND PRECAUTIONS

Please see [3 SERIOUS WARNINGS AND PRECAUTIONS BOX](#).

General

Patient experience with trientine hydrochloride is limited (see [14 CLINICAL TRIALS](#)). Patients receiving WAYMADE-TRIENTINE require regular and ongoing monitoring of their clinical condition and response to treatment throughout the period of drug administration.

Overtreatment carries the risk of copper deficiency, which is especially harmful for children and pregnant women, since copper is required for proper growth and mental development. Therefore, ongoing regular monitoring for manifestations of overtreatment should be undertaken.

Carcinogenesis and Mutagenesis

Data on carcinogenesis, and impairment of fertility are not available. For genotoxicity potential of trientine, see [16 NON-CLINICAL TOXICOLOGY, Genotoxicity](#).

Hematologic

Trentine hydrochloride is a chelating agent that has been found to reduce serum iron levels, which may lead to iron deficiency anemia. Oral iron supplementation may be necessary, and if required, it should be administered at a different time of day than WAYMADE-TRIENTINE (see [9.4 Drug-Drug Interactions](#)).

Immune

There are no reports of hypersensitivity in patients who have been administered WAYMADE-TRIENTINE for Wilson's Disease. However, there have been reports of asthma, bronchitis and dermatitis occurring after prolonged environmental exposure in workers who use trientine hydrochloride as a hardener of epoxy resins. Patients should be observed closely for signs of possible hypersensitivity.

Monitoring and Laboratory Tests

The most reliable index for monitoring treatment is the determination of free copper in the serum, which equals the difference between quantitatively determined serum total copper and ceruloplasmin-bound copper. Adequately treated patients will usually have less than 3.1 mcmol/L (20 mcg/dL) of serum free copper.

Chelation therapy may be monitored with a 24-hour urinary copper analysis periodically, e.g., every 6 to 12 months. Urine must be collected in copper-free glassware. Since a low copper diet should maintain copper absorption to less than one milligram per day, the patient will probably be in the desired state of negative copper balance if 0.5 to 1.0 mg of copper is present in a 24-hour collection of urine.

Neurologic

Worsening of neurologic or neurocognitive functioning, which can be irreversible, may occur in patients with pre-existing neurologic and/or neuropsychiatric impairment at the time of initiation of chelator therapy with trientine hydrochloride. Initiation of WAYMADE-TRIENTINE should be carried out with caution in these patients, and only after careful consideration of all available treatment options and the determination of an acceptable benefit-risk for the patient. Close monitoring of clinical response in these patients is required in the first few months following initiation of WAYMADE-TRIENTINE.

Reproductive Health: Female and Male Potential

See [7.1.1 Pregnant women](#).

- Fertility**

No human fertility studies have been conducted. Based on animal studies, there are no available data regarding effects on fertility (see [16 NON-CLINICAL TOXICOLOGY](#)).

- Teratogenic Risk**

Trentine hydrochloride may cause fetal harm when administered to a pregnant woman based on results from animal reproduction studies (see [16 NON-CLINICAL TOXICOLOGY, Reproductive and Developmental Toxicology](#)). Some available data have showed that trientine hydrochloride has a genotoxic potential (see [16 NON-CLINICAL TOXICOLOGY, Genotoxicity](#)).

WAYMADE-TRIENTINE should only be used during pregnancy after carefully considering the potential benefits of treatment compared to its risks and when the benefits to women of childbearing potential are judged to outweigh the risks (see [7.1.1 Pregnant women](#)).

Skin

Because of the potential for contact dermatitis, any exposure to the capsule contents should be washed promptly with water.

7.1 Special Populations

7.1.1 Pregnant women

No well-controlled studies of the use of trientine hydrochloride in pregnant women have been conducted. There is only a limited amount of experience with data from the use of trientine hydrochloride in pregnant women. WAYMADE-TRIENTINE should be used during pregnancy only after careful consideration of the potential benefits for the patient, compared with its risks. Factors to consider when using trientine hydrochloride in pregnancy include the risks associated with uncontrolled disease, maintenance of serum iron concentrations, the risks of alternative treatments, and the possible teratogenic effects of trientine hydrochloride.

Studies in animals have shown that trientine hydrochloride is teratogenic in rats at systemic exposures similar to those seen in humans, which may have been the result of trientine-induced copper deficiency. The frequencies of both resorptions and fetal abnormalities of the brain increased, while fetal copper concentrations decreased, when maternal rats were fed trientine hydrochloride in the diet. Some data are available with respect to the genotoxic potential of trientine hydrochloride (see [16 NON-CLINICAL TOXICOLOGY, Genotoxicity](#)). Closely monitor serum free copper concentrations and consider a dosage adjustment to the lowest dose of WAYMADE-TRIENTINE that maintains free copper concentrations within acceptable limits.

7.1.2 Breast-feeding

It is not known whether trientine hydrochloride is excreted in human milk. Limited clinical data suggest that trientine hydrochloride is not excreted in breast milk, however, a risk to newborns and infants cannot be excluded. Because many drugs are excreted in human milk, caution should be exercised when WAYMADE-TRIENTINE is administered to breastfeeding women.

7.1.3 Pediatrics

Pediatrics (< 18 years of age): Controlled studies of the safety and effectiveness of trientine hydrochloride in pediatric patients have not been conducted.

7.1.4 Geriatrics

Geriatrics (> 65 years of age): Reported clinical experience with trientine hydrochloride has not included sufficient numbers of patients over the age of 65 years to determine whether these patients respond differently than patients less than 65 years of age. In general, dose selection should be cautious in these patients, usually starting at the low end of the dosing range, reflecting the greater frequency of concomitant disease or other drug therapy.

8 ADVERSE REACTIONS

8.1 Adverse Reaction Overview

Clinical experience with trientine hydrochloride has been limited.

8.2 Clinical Trial Adverse Reactions

Clinical trials are conducted under very specific conditions. The adverse reaction rates observed in the clinical trials; therefore, may not reflect the rates observed in practice and should not be compared to the rates in the clinical trials of another drug. Adverse reaction information from clinical trials may be useful in identifying and approximating rates of adverse drug reactions in real-world use.

Nausea occurs commonly following treatment initiation with trientine hydrochloride. The occurrence of skin rash has also been reported. Severe duodenitis and/or colitis have been observed on occasion. Sideroblastic anemia and aplastic anemia have been reported rarely.

Iron deficiency has been observed with trientine hydrochloride treatment (see [7 WARNINGS AND PRECAUTIONS, Hematologic](#)).

8.3 Post-Market Adverse Reactions

In addition to the adverse reactions reported during pre-marketing use, the following adverse reactions have been reported in post-marketed use: Neurological deterioration, with signs including dystonia, rigidity, tremor, and dysarthria.

9 DRUG INTERACTIONS

9.4 Drug-Drug Interactions

No interaction studies have been performed with WAYMADE-TRIENTINE. Mineral supplements should be avoided as they may block the absorption of WAYMADE-TRIENTINE. If iron deficiency develops, particularly in children and menstruating or pregnant women, or as a result of the low copper diet recommended for Wilson's Disease, iron may be administered in short courses. As iron and WAYMADE-TRIENTINE inhibit the absorption of each other, doses of WAYMADE-TRIENTINE and iron should be separated by at least two hours.

9.5 Drug-Food Interactions

Trentine hydrochloride is poorly absorbed following oral intake and food further inhibits its absorption. It is important that WAYMADE-TRIENTINE be taken on an empty stomach, at least one hour before meals or two hours after meals and at least one hour apart from any other drug, food, or milk. This permits maximum absorption and reduces the likelihood of inactivation of the drug by metal binding in the gastrointestinal tract.

9.6 Drug-Herb Interactions

Interactions with herbal products have not been established.

9.7 Drug-Laboratory Test Interactions

Interactions with laboratory tests have not been established.

10 CLINICAL PHARMACOLOGY

10.1 Mechanism of Action

Trentine hydrochloride is a copper chelating agent that aids in the elimination of copper from the body by forming a stable complex that is readily excreted by the kidneys. Trentine hydrochloride may also chelate copper in the intestinal tract, and thus, inhibit copper absorption.

10.2 Pharmacodynamics

Trientine is a chelator with a polyamine-like structure and copper is chelated by forming a stable complex with the four constituent nitrogens in a planar ring. Thus, the pharmacodynamic action of trientine is dependent on its chemical property of chelating copper and not on its interaction with receptors, enzyme systems, or any other biological system that might differ between species.

Preclinical Studies

Studies in animals have shown that trientine hydrochloride has cupriuretic activities in both normal and copper-loaded rats without significant changes in serum copper. Urinary copper excretion was greater and accelerated in copper-loaded rats compared with normal rats. In general, the effects of trientine hydrochloride on urinary copper excretion are similar to those of penicillamine.

Human Studies

Renal clearance studies were carried out with penicillamine and trientine hydrochloride in selected patients with confirmed Wilson's Disease, treated with penicillamine for at least one year. Six-hour excretion rates of copper were determined pre-treatment (basal) and after a single dose of 500 mg of penicillamine or 1.2 g of trientine. The mean urinary excretion rates of copper were as follows:

No. of Patients	Single Dose Treatment	Basal Excretion Rate (mcg Cu ++ /6 h)	Post-Test Dose Excretion Rate (mcg Cu ++ /6 h)
6	Trientine, 1.2 g	19	234
4	Penicillamine, 500 mg	17	320

In patients not previously treated with chelating agents, a similar comparison was made:

No. of Patients	Single Dose Treatment	Basal Excretion Rate (mcg Cu ++ /6 h)	Post-Test Dose Excretion Rate (mcg Cu ++ /6 hr)
8	Trientine, 1.2 g	71	1326
7	Penicillamine, 500 mg	68	1074

Trientine hydrochloride is effective as a cupriuretic agent in patients with Wilson's Disease, although on a molar basis, the effect may be smaller than with penicillamine. The difference in cupriuretic effect between these two drugs could be due to a difference in selectivity of the drugs for different copper pools within the body.

10.3 Pharmacokinetics

Data on the pharmacokinetics of trientine hydrochloride are limited. Dosage adjustment recommendations are based upon clinical use of the drug (see [4 DOSAGE AND ADMINISTRATION](#)).

Table 2: Summary of Trientine Hydrochloride Pharmacokinetic Parameters in Healthy Adult Subject

	C _{max} (mg/L)	T _{max} (h)	t _½ (h)	AUC _{0..∞} (mg.h/L)
Multiple dose mean (Day 14, 1.2 g/d)	1.55	0.83	10.37	11.14

C_{max} = maximum observed concentration, t_{max} = time to maximum observed concentration; t_½ = elimination half-life; AUC = area under the curve.

Absorption

The absorption of trientine hydrochloride following oral administration is low and variable in patients with Wilson's Disease. Trientine is absorbed with t_{max} occurring between 0.5 and 4 hours after dosing and shows biexponential decline in serum concentrations. Gender and body weight do not seem to influence pharmacokinetic parameters.

Distribution

No data are available for the tissue distribution of trientine in humans. However, trientine is distributed in most rat tissues, particularly in the liver and kidney. In healthy volunteers, the central and peripheral volumes were shown to be large, indicating that trientine is probably widely distributed in human body tissues.

Metabolism

Trientine is acetylated into two major metabolites, 1-N-acetyltriethylene tetramine (MAT) and N1,N10-diacetyltriethylenetetramine (DAT). MAT may participate in the overall activity of trientine, however the extent to which MAT effects copper concentrations has not been determined.

Elimination

Trientine and its metabolites are rapidly excreted in the urine, although traces could still be detected after 24 hours. Unabsorbed trientine is eliminated through fecal excretion.

Special Populations and Conditions

Pediatrics: A non-compartmental pharmacokinetic analysis of four pediatric patients with confirmed Wilson's Disease, who were aged 12 years or older and who were receiving a stable regimen of daily trientine (0.3-0.9 g/day), demonstrated similarity with pharmacokinetic parameters in adult patients with Wilson's Disease. There are no pharmacokinetic data available for pediatric patients less than 12 years of age.

Health Canada has not authorized an indication for pediatric use in children less than 5 years of age (see [1.1 Pediatrics](#)).

Geriatrics: There are no data available for the pharmacokinetic parameters of trientine in the elderly.

11 STORAGE, STABILITY AND DISPOSAL

Keep container tightly closed.

Store refrigerated between 2° to 8°C. Do not freeze. After opening, the bottle can be stored at room temperature for 2 months, then discarded.

Store in the original container and retain the silica gel sachet in the bottle in order to protect from moisture.

12 SPECIAL HANDLING INSTRUCTIONS

Because of the potential for contact dermatitis, any site of exposure to the capsule contents should be washed with water promptly.

PART II: SCIENTIFIC INFORMATION

13 PHARMACEUTICAL INFORMATION


Drug Substance

Proper name: Trientine Hydrochloride

Chemical name: 1,2-Ethanediamine,N,N'-bis (2-aminoethyl)- dihydrochloride or Triethylenetetramine dihydrochloride

Molecular formula and molecular mass: C₆H₁₈N₄•2HCl, 219.16 g/mole

Structural formula:



Physicochemical properties: White to pale yellow crystalline powder. Freely soluble in water. Soluble in methanol. Slightly soluble in alcohol. Insoluble in chloroform and in ether.

14 CLINICAL TRIALS

14.1 Clinical Trials by Indication

The Treatment Of Patients With Wilson's Disease Who Are Intolerant To Penicillamine.

Summary of patient demographics for clinical trials in patients with Wilson's Disease

A total of 405 patients with Wilson's Disease were evaluated in a retrospective cohort study, which included 380 patients followed at tertiary health care centres in Central Europe by specialist physicians responsible for their care. There were 25 additional patients from the EUROWILSON patient registry who were included in this retrospective analysis, with all of these registry patients receiving trientine hydrochloride monotherapy as treatment for Wilson's Disease.

Data were collected in this study over a mean of 13.3 years after treatment initiation for Wilson's Disease. Patients in this study were categorised based on disease status at the time of disease diagnosis, as either asymptomatic, with hepatic disease, with neurologic disease, or with combined hepatic or neurologic presentations. The presence of Kayser–Fleischer rings at slit-lamp examination and hepatic cirrhosis was recorded at study entry.

Patients whose disease course was considered stable were followed yearly. However, upon treatment initiation or following a change in drug treatment, patients were followed more intensively, i.e., at 3, 6 and 12 months, until clinically stable.

Patients received initial treatment with chelation therapy, if symptomatic. The choice of initial chelating agent for each patient, whether penicillamine or trientine hydrochloride, was left to the discretion of attending medical staff. Patients who were treated with zinc salts were excluded from study analysis.

For purposes of study analysis, two treatment regimens in patients were identified for this retrospective analysis, either penicillamine monotherapy, or trientine hydrochloride monotherapy. Stable drug treatment periods of at least six (6) consecutive months were included in the study

analyses. Efficacy analyses were conducted on initial and/or subsequent treatment regimens of penicillamine and trientine hydrochloride. In addition, analyses of the cause of treatment discontinuations were conducted.

From patient records, hepatic and neurologic outcomes were assessed at 6, 12, 24, 36 and to a maximum of 48 months, following initiation of each treatment regimen, with outcome measures stratified by first-line and second-line treatment use. Hepatic outcome measures were assessed using clinical symptoms, as well as liver function tests. The presence and/or progress of neurologic disease were evaluated by physician assessment. Both hepatic and neurologic outcomes were categorised as unchanged, improved to normal, improved but not normal, deteriorated, or asymptomatic, over the period of time assessed.

Results of study in patients with Wilson's Disease

Treatment regimens from a total of 405 patients were reviewed for the analysis. Of these patients, 207 patients (51.1%) presented with hepatic symptoms only, 92 patients (22.7%) presented with neurologic symptoms only, 52 (12.8%) presented with both hepatic and neurologic symptoms, and 54 (13.3%) were asymptomatic. At the time of diagnosis, 21 patients (5.2%) presented in hepatic failure with a fulminant disease course.

Changes in study treatments were common in the study dataset. In total, 467 chelator-based treatment regimens of a duration of at least six months were identified and analysed. Of these, 326 were penicillamine treatment regimens, while 141 were trientine regimens. Of the 326 penicillamine treatment regimens identified for analysis, 294 were first-line therapies, whereas the majority of trientine monotherapies, i.e., 105 of 141, were second-line therapies. None of these patients received penicillamine and trientine concomitantly at any time.

Efficacy was determined for both first-line and second-line study treatments for the last available follow-up evaluation of a 6-month to 48-month follow-up period.

In symptomatic hepatic patients treated with chelator therapy as second-line treatment when first-line treatment failed or was not tolerated, 75.0% of penicillamine treatment regimens and 68.9% of trientine hydrochloride regimens were associated with improvement in hepatic signs and symptoms. For symptomatic hepatic patients, the observed incidence of patients who maintained stable hepatic disease was comparable for penicillamine and trientine hydrochloride treatment in second-line settings. A non-significant trend to increased rates of hepatic deterioration was seen when trientine hydrochloride was used as second-line treatment, compared to second-line penicillamine treatment, however, the majority of second-line trientine hydrochloride-treated patients, at 91.1%, were not observed to have such deterioration. On the other hand, in patients who initially presented free of hepatic symptoms, no worsening in hepatic status was observed for any of these patients who received either of the chelation monotherapy treatment regimens evaluated.

With respect to neurologic outcomes in patients with Wilson's Disease having symptomatic neurologic disease at baseline, rates of improvement following second-line penicillamine treatment were noted at 23.1%, compared to 51.0% with second-line trientine hydrochloride. Stable neurologic disease was reported in 69.2% of second-line penicillamine treatment regimens, compared to 33.3% of second-line trientine hydrochloride treatments. In this study, neurologic worsening was reported in neurologically symptomatic patients at study entry treated with second-line trientine hydrochloride at 15.7%, compared with second-line penicillamine treatment at 7.3%.

Overall, these efficacy analyses demonstrate the utility of trientine hydrochloride in the treatment of patients with Wilson's Disease as second-line therapy. In patients with symptomatic hepatic disease,

69% given trientine hydrochloride as second-line therapy, due to intolerance or inadequate response to penicillamine, were observed to have shown hepatic improvement with trientine hydrochloride, while 51% of patients with symptomatic neurological disease at study entry were observed to have shown improvement in these symptoms with trientine hydrochloride when used as second-line treatment.

Over a median follow-up period of over 13 years, study treatment regimens that were discontinued for any reason were 43.6% of penicillamine treatment regimens, compared with 25.5% of trientine hydrochloride treatment regimens.

14.2 Comparative Bioavailability Studies

An open label, randomized, single-dose, two-way, crossover comparative bioavailability study of WAYMADE-TRIENTINE 250 mg capsules (Waymade PLC) and SYPRINE® 250 mg capsules (Valeant Pharmaceuticals North America LLC, USA) was conducted in healthy, adult, male subjects under fasting conditions. Comparative bioavailability data from 38 subjects that were included in the statistical analysis are presented in the following table:

Table 3: Summary Table of the Comparative Bio-Availability Data

Trientine (1 x 250 mg) From measured data Geometric Mean Arithmetic Mean (CV %)				
Parameter	Test ¹	Reference ²	% Ratio of Geometric Means	90 % Confidence Interval
AUC _T (ng.h/mL)	3632.17 4298.19 (56.25)	3787.00 4444.25 (60.64)	95.9	87.1 – 105.6
AUC _I (ng.h/mL)	3804.38 4475.81 (55.74)	3968.54 4616.42 (58.67)	95.9	87.2 – 105.4
C _{max} (ng/mL)	817.15 936.71 (47.52)	826.50 933.561 (50.93)	98.9	90.3 – 108.2
T _{max} ³ (h)	1.50 (0.33 – 4.50)	1.50 (0.50 – 3.50)		
T _½ ⁴ (h)	13.52 (59.65)	13.44 (40.48)		

¹ WAYMADE-TRIENTINE (trientine dihydrochloride) capsules, 250 mg (Waymade PLC)

² Syprine® (trientine hydrochloride) capsules, 250 mg (Valeant Pharmaceuticals North America LLC, USA)

³ Expressed as the median (range) only

⁴ Expressed as the arithmetic mean (CV %) only

15 MICROBIOLOGY

No microbiological information is required for this drug product.

16 NON-CLINICAL TOXICOLOGY

General Toxicology

Mice administered trientine in drinking water exhibited more inflammation of the lung interstitium, including alveolar histiocytic infiltration, liver periportal fatty infiltration, and hematopoietic cell proliferation in the spleen. Body and kidney weights as well as renal cytoplasmic vacuolization were reduced in males. The NOAEL was considered to be 92 and 99 mg/kg/day in males and females, respectively.

Rats administered oral doses of trientine up to 600 mg/kg/day for 26 weeks displayed dose-related incidence and severity of focal chronic interstitial pneumonitis with fibrosis of the alveolar walls, indicative of persistent inflammation or a toxic effect on alveolar cells. As trientine has irritating properties, it was suggested that the accumulation of trientine in bronchial epithelial cells and alveolar pneumocytes had a cytotoxic effect resulting in chronic interstitial pneumonitis. The NOAEL was considered to be 50 mg/kg/day for females and less than 50 mg/kg/day for males as no NOAEL was established.

Dogs administered oral doses of trientine up to 200 mg/kg/day in repeat-dose toxicity studies demonstrated reversible underactivity, tremors, abnormal gait, limited use of limb, and prone posture. The NOAEL was established at 50 mg/kg/day.

Carcinogenicity

No carcinogenicity studies have been conducted with trientine.

Genotoxicity

In vitro, trientine was mutagenic in the Ames test, the *Salmonella typhimurium* mutation assay, and the sister-chromatid exchange assay in Chinese hamster ovary cells. In hepatocytes, it also produced unscheduled DNA synthesis activity, indicating the induction of primary damage in this assay.

In vivo, trientine was negative in the mouse micronucleus test.

Reproductive and Developmental Toxicology

A study of oral prenatal trientine exposure (doses of 0, 3000, 6000, and 12000 ppm in drinking water) in mice demonstrated that trientine had stronger fetal than maternal adverse effects. Fetal tissues demonstrated low dose-dependent copper concentrations at doses higher than 3000 ppm, which was calculated as approximately 500 mg/kg/day in pregnant mice, and morphological abnormalities in the brain. Fetal resorption increased and fetal viability was reduced in a dose-dependent manner and serum copper concentrations were lower in dams that experienced total resorption at 6000 or 12000 ppm. The effects observed may be due, at least in part, to induction of copper deficiency.

PATIENT MEDICATION INFORMATION

READ THIS FOR SAFE AND EFFECTIVE USE OF YOUR MEDICINE

Pr WAYMADE-TRIENTINE

Trentine Hydrochloride Capsules

Read this carefully before you start taking **WAYMADE-TRIENTINE** and each time you get a refill. This leaflet is a summary and will not tell you everything about this drug. Talk to your healthcare professional about your medical condition and treatment and ask if there is any new information about **WAYMADE-TRIENTINE**.

Serious Warnings and Precautions

WAYMADE-TRIENTINE will only be given to you under the care of a healthcare professional who is experienced in the treatment of Wilson's disease.

Starting treatment with WAYMADE-TRIENTINE may result in permanent worsening of your existing nervous system problems. If you experience any worsening of the following symptoms while taking WAYMADE-TRIENTINE, stop taking the drug and get immediate medical help:

- shaking or twisting movements that can not be controlled
- lack of coordination
- muscle stiffness
- muscle spasms
- slurred speech or changes in your voice

Your healthcare professional will monitor your response for these symptoms for the first few months when you start taking WAYMADE-TRIENTINE.

What is WAYMADE-TRIENTINE used for?

- WAYMADE-TRIENTINE is used for the treatment of Wilson's disease. It is used in adult patients who cannot take the drug penicillamine.

How does WAYMADE-TRIENTINE work?

WAYMADE-TRIENTINE works by attaching to the copper, and then passing it from your body. WAYMADE-TRIENTINE may also work by attaching to the copper in your stomach and stopping it from being absorbed.

What are the ingredients in WAYMADE-TRIENTINE?

Medicinal ingredients: trientine hydrochloride

Non-medicinal ingredients: black ink, gelatin, purified water, stearic acid, sunset yellow FCF, titanium dioxide

WAYMADE-TRIENTINE comes in the following dosage forms:

WAYMADE-TRIENTINE is available as 250 mg capsules.

Do not use WAYMADE-TRIENTINE if:

- You are allergic to trientine hydrochloride or any of the other ingredients in this drug or the container.

To help avoid side effects and ensure proper use, talk to your healthcare professional before you take WAYMADE-TRIENTINE. Talk about any health conditions or problems you may have, including if you:

- have breathing problems such as asthma or bronchitis (swelling of the bronchial tubes, the airways that carry air to your lungs).
- have dermatitis (red and itchy dry skin)
- have anemia (low iron in your blood)
- are pregnant, think you are pregnant, or plan to become pregnant
- are breast feeding
- are older than 65 years of age.

Other warnings you should know about:

Monitoring and Testing

Your healthcare professional will order lab tests for you during your treatment. This will help determine copper levels in your body and how your body responds to the treatment with WAYMADE-TRIENTINE.

Tell your healthcare professional about all the medicines you take, including any drugs, vitamins, minerals, natural supplements or alternative medicines.

The following may interact with WAYMADE-TRIENTINE:

- Mineral and iron supplements

How to take WAYMADE-TRIENTINE:

- Take WAYMADE-TRIENTINE exactly as your healthcare professional tells you. Your healthcare professional will determine the right dose for you.
- Take WAYMADE-TRIENTINE on an empty stomach, at least 1 hour before meals or 2 hours after meals.
- Take WAYMADE-TRIENTINE at least 1 hour apart from any other drug, food, or milk.
- Your healthcare professional may give you an iron supplement if needed. Take the iron at least 2 hours apart from WAYMADE-TRIENTINE.
- Swallow the capsules whole with water, and do not open or chew the capsules.
- Touching the contents of the WAYMADE-TRIENTINE capsules may cause your skin to become red and itchy. Wash your hands with water if you accidentally touch the contents of the WAYMADE-TRIENTINE capsules.

Usual dose:

Adults: the starting dose is 3 capsules (750 mg) per day, given in divided doses 2 – 4 times a day. The maximum dose is 8 capsules (2,000 mg) per day.

Children and Adolescents (5 – 17 years old): the starting dose is 2 capsules per day (500 mg / day) given in divided doses 2 – 4 times a day. The maximum dose is 6 capsules a day (1,500 mg / day) for patients 12 years old or under.

Overdose:

If you think you, or a person you are caring for, have taken too much WAYMADE-TRIENTINE, contact a healthcare professional, hospital emergency department, or regional poison control centre immediately, even if there are no symptoms.

Missed Dose:

If you missed a dose of WAYMADE-TRIENTINE, take it as soon as you remember. But if it is almost time for your next dose, skip the missed dose and continue with your next scheduled dose. Go back to the regular dosing schedule. Do not take two doses at the same time or very close to each other.

What are possible side effects from using WAYMADE-TRIENTINE?

These are not all the possible side effects you may have when taking WAYMADE-TRIENTINE. If you experience any side effects not listed here, tell your healthcare professional.

Symptoms may include:

- Nausea
- Rash, itchy or red skin

Serious side effects and what to do about them			
Symptom / effect	Talk to your healthcare professional		Stop taking drug and get immediate medical help
	Only if severe	In all cases	
UNCOMMON			
Duodenitis and/or colitis (bowel inflammation): nausea, fatigue, vomiting, diarrhea, bloating, abdominal pain, fever		✓	
RARE			
Anemia (low levels of red blood cells): pale skin, feeling tired or weak, shortness of breath, irregular heartbeat	✓		
UNKNOWN			
Asthma, bronchitis (lung or airway problems): difficulty breathing and coughing, shortness of breath, chest tightness, wheezing or whistling sound when breathing, fever, runny nose		✓	
Neurological deterioration (nervous system problems):			✓

Serious side effects and what to do about them			
Symptom / effect	Talk to your healthcare professional		Stop taking drug and get immediate medical help
	Only if severe	In all cases	
<p>symptoms include:</p> <ul style="list-style-type: none"> • Dystonia (twisting movements that you cannot control and can affect posture or the face including eyes, mouth, tongue or jaw, tightness of the throat, difficulty swallowing or breathing which may lead to choking). • Rigidity (stiff muscles) • Tremor (shaking that can't be controlled) • Dysarthria (slurred speech, changes in your voice) 			

If you have a troublesome symptom or side effect that is not listed here or becomes bad enough to interfere with your daily activities, tell your healthcare professional.

Reporting Side Effects

You can report any suspected side effects associated with the use of health products to Health Canada by:

- Visiting the Web page on Adverse Reaction Reporting (<https://www.canada.ca/en/health-canada/services/drugs-health-products/medeffect-canada/adverse-reaction-reporting.html>) for information on how to report online, by mail or by fax; or
- Calling toll-free at 1-866-234-2345.

NOTE: Contact your health professional if you need information about how to manage your side effects. The Canada Vigilance Program does not provide medical advice.

Storage:

- Store WAYMADE-TRIENTINE at 2° to 8°C. Keep refrigerated. Do not freeze.
- After the first opening, the bottle can be stored at room temperature for 2 months.. Dispose after 2 months.
- Store the capsules in their original bottle. Keep the bottle tightly closed and leave the silica gel sachet in the bottle to protect from humidity.
- Keep out of reach and sight of children.

If you want more information about WAYMADE-TRIENTINE:

- Talk to your healthcare professional

- Find the full product monograph that is prepared for healthcare professionals and includes this Patient Medication Information by visiting the Health Canada website: (<https://www.canada.ca/en/health-canada/services/drugs-health-products/drug-products/drug-product-database.html>); the manufacturer's website (www.waymade.co.uk), or by calling 1-800-970-8841.

This leaflet was prepared by Waymade PLC.

Last Revised: November 14, 2023