

PRODUCT MONOGRAPH
INCLUDING PATIENT MEDICATION INFORMATION

PrTeva-Tofacitinib

Tofacitinib Tablets
Tablets, 5 mg and 10 mg tofacitinib (as tofacitinib citrate), Oral

Selective Immunosuppressant

Teva Canada Limited
30 Novopharm Court
Toronto, Ontario
M1B 2K9
www.tevacanada.com

Date of Authorization: December 29, 2023

Submission Control Number: 260906

RECENT MAJOR LABEL CHANGES

Sections or subsections that are not applicable at the time of authorization are not listed.

TABLE OF CONTENTS

| | |
|---|-----------|
| RECENT MAJOR LABEL CHANGES | 2 |
| PART I: HEALTH PROFESSIONAL INFORMATION | 4 |
| 1 INDICATIONS | 4 |
| 1.1 Pediatrics | 4 |
| 1.2 Geriatrics | 4 |
| 2 CONTRAINDICATIONS | 4 |
| 3 SERIOUS WARNINGS AND PRECAUTIONS BOX | 5 |
| 4 DOSAGE AND ADMINISTRATION | 6 |
| 4.1 Dosing Considerations | 6 |
| 4.2 Recommended Dose and Dosage Adjustment | 6 |
| 4.4 Administration | 9 |
| 4.5 Missed Dose | 9 |
| 5 OVERDOSE | 9 |
| 6 DOSAGE FORMS, STRENGTHS, COMPOSITION AND PACKAGING | 10 |
| 7 WARNINGS AND PRECAUTIONS | 10 |
| 7.1 Special Populations | 18 |
| 7.1.1 Pregnant Women | 18 |
| 7.1.2 Breast-feeding | 18 |
| 7.1.3 Pediatrics | 18 |
| 7.1.4 Geriatrics | 18 |
| 7.1.5 Asian Patients | 18 |
| 8 ADVERSE REACTIONS | 19 |
| 8.1 Adverse Reaction Overview | 19 |
| 8.2 Clinical Trial Adverse Reactions | 20 |
| Rheumatoid Arthritis | 20 |
| Psoriatic Arthritis | 28 |
| Ulcerative Colitis | 28 |
| 8.3 Less Common Clinical Trial Adverse Reactions | 31 |
| 8.4 Abnormal Laboratory Findings: Hematologic, Clinical Chemistry and Other Quantitative Data 32 | |
| 8.5 Post-Market Adverse Reactions | 36 |
| 9 DRUG INTERACTIONS | 36 |
| 9.2 Drug Interactions Overview | 36 |
| 9.4 Drug-Drug Interactions | 36 |
| 9.5 Drug-Food Interactions | 40 |
| 9.6 Drug-Herb Interactions | 40 |
| 9.7 Drug-Laboratory Test Interactions | 41 |
| 10 CLINICAL PHARMACOLOGY | 41 |
| 10.1 Mechanism of Action | 41 |
| 10.2 Pharmacodynamics | 41 |
| 10.3 Pharmacokinetics | 42 |
| 11 STORAGE, STABILITY AND DISPOSAL | 46 |
| 12 SPECIAL HANDLING INSTRUCTIONS | 46 |
| PART II: SCIENTIFIC INFORMATION | 47 |

| | | |
|------|---|-----------|
| 13 | PHARMACEUTICAL INFORMATION | 47 |
| 14 | CLINICAL TRIALS..... | 47 |
| 14.1 | Clinical Trials by Indication | 47 |
| | Rheumatoid Arthritis | 47 |
| | Psoriatic Arthritis..... | 56 |
| | Ulcerative Colitis..... | 61 |
| 14.2 | Comparative Bioavailability Studies | 66 |
| 15 | MICROBIOLOGY..... | 67 |
| 16 | NON-CLINICAL TOXICOLOGY..... | 67 |
| 17 | SUPPORTING PRODUCT MONOGRAPHS..... | 77 |
| | PATIENT MEDICATION INFORMATION | 78 |

PART I: HEALTH PROFESSIONAL INFORMATION

1 INDICATIONS

Rheumatoid Arthritis

Teva-Tofacitinib (tofacitinib), in combination with methotrexate (MTX), is indicated for reducing the signs and symptoms of rheumatoid arthritis (RA) in adult patients with moderately to severely active RA who have had an inadequate response to MTX and to one or more disease-modifying anti-rheumatic drugs (DMARDs).

In cases of intolerance to MTX and other DMARDs, physicians may consider the use of Teva-Tofacitinib (tofacitinib) as monotherapy.

Psoriatic Arthritis

Teva-Tofacitinib (tofacitinib), in combination with methotrexate (MTX) or another conventional synthetic disease-modifying antirheumatic drug (DMARD), is indicated for reducing the signs and symptoms of psoriatic arthritis (PsA) in adult patients with active PsA when the response to previous DMARD therapy has been inadequate.

Ulcerative Colitis

Teva-Tofacitinib (tofacitinib) is indicated for the treatment of adult patients with moderately to severely active ulcerative colitis (UC) with an inadequate response, loss of response or intolerance to either conventional UC therapy or a TNF α inhibitor.

Limitations of Use

Teva-Tofacitinib should not be used in combination with other Janus kinase (JAK) inhibitors, immunomodulating biologics (e.g., biologic DMARDs), or with potent immunosuppressants such as azathioprine and cyclosporine.

1.1 Pediatrics

Pediatrics (<18 years of age): No data are available to Health Canada; therefore, Health Canada has not authorized an indication for pediatric use.

1.2 Geriatrics

Geriatrics (>65 years of age): Evidence from clinical studies and experience suggests that use in the geriatric population is associated with differences in safety or effectiveness. The frequency of adverse events including serious infections, all-cause mortality, cardiovascular events, malignancies, non-melanoma skin cancer, gastrointestinal perforations, interstitial lung disease, venous thromboembolism, and arterial thromboembolism in tofacitinib-treated subjects 65 years of age and older was higher than among those under the age of 65. Therefore, caution should be used when treating geriatric patients with Teva-Tofacitinib (see [7 WARNINGS AND PRECAUTIONS](#), [4 DOSAGE AND ADMINISTRATION](#) and [10 CLINICAL PHARMACOLOGY](#))

2 CONTRAINDICATIONS

Teva-Tofacitinib (tofacitinib) is contraindicated:

- In patients with known hypersensitivity to tofacitinib or ingredient in the formulation, including any non-medicinal ingredient, or component of the container. For a complete listing, see

[6 DOSAGE FORMS, STRENGTHS, COMPOSITION AND PACKAGING.](#)

- In patients with severe hepatic impairment (see [7 WARNINGS AND PRECAUTIONS](#)).
- During pregnancy and breastfeeding (see [7.1.1 Pregnancy](#) and [7.1.2 Breast-feeding](#)).

3 SERIOUS WARNINGS AND PRECAUTIONS BOX

Serious Warnings and Precautions

SERIOUS INFECTIONS

Patients treated with Teva-Tofacitinib (tofacitinib) are at increased risk for developing serious infections that may lead to hospitalization or death (see [7 WARNINGS AND PRECAUTIONS](#) and [8.2 Clinical Trial Adverse Reactions](#)). Most patients who developed these infections were taking concomitant immunosuppressants such as methotrexate or corticosteroids.

If a serious infection develops, interrupt Teva-Tofacitinib until the infection is controlled.

Reported infections include:

- Active tuberculosis, which may present with pulmonary or extrapulmonary disease. Patients should be tested for latent tuberculosis before Teva-Tofacitinib use and during therapy. Treatment for latent infection should be initiated prior to Teva-Tofacitinib use.
- Invasive fungal infections, including cryptococcosis and pneumocystosis. Patients with invasive fungal infections may present with disseminated, rather than localized disease.
- Bacterial, viral, and other infections due to opportunistic pathogens.

Treatment with Teva-Tofacitinib should not be initiated in patients with active infections including chronic or localized infection.

Patients should be closely monitored for the development of signs and symptoms of infection during and after treatment with Teva-Tofacitinib, including the possible development of tuberculosis in patients who tested negative for latent tuberculosis infection prior to initiating therapy (see [8 ADVERSE REACTIONS](#)).

MALIGNANCIES

Lymphoma and other malignancies have been observed in patients treated with tofacitinib. Epstein Barr Virus-associated post-transplant lymphoproliferative disorder has been observed at an increased rate in renal transplant patients treated with tofacitinib and concomitant immunosuppressive medications. An increase in malignancies, including lung cancer, were observed in rheumatoid arthritis patients 50 years or older with at least one additional cardiovascular (CV) risk factor who were taking tofacitinib compared with TNF inhibitors (see [8.2 Clinical Trial Adverse Reactions](#)). Caution should be applied when using Teva-Tofacitinib in geriatric patients, patients who are current or past smokers, and patients with other malignancy risk factors (see [7 WARNINGS AND PRECAUTIONS](#)).

THROMBOSIS

Rheumatoid arthritis patients with at least one CV risk factor had a higher rate of all-cause mortality and thrombosis, including pulmonary embolism, deep venous thrombosis, and arterial thrombosis, with tofacitinib 10 mg twice daily (BID) compared to those treated with 5 mg BID or TNF blockers. Many of these adverse events were serious and some resulted in death. Avoid Teva-Tofacitinib in patients at risk of thrombosis. Discontinue Teva-Tofacitinib and promptly evaluate patients with symptoms of thrombosis (see [7 WARNINGS AND PRECAUTIONS](#)).

For patients with ulcerative colitis (UC), use Teva-Tofacitinib at the lowest effective dose and for the shortest duration needed to achieve/maintain therapeutic response (see [4 DOSAGE AND ADMINISTRATION](#)).

MAJOR ADVERSE CARDIOVASCULAR EVENTS

Major adverse cardiovascular events, including non-fatal myocardial infarction, were observed more frequently with tofacitinib compared to TNF inhibitors in rheumatoid arthritis patients who were 50 years or older with at least one additional CV risk factor (see [8.2 Clinical Trial Adverse Reactions](#)). Caution should be applied when using Teva-Tofacitinib in geriatric patients, patients who are current or past smokers, and patients with other CV risk factors (see [7 WARNINGS AND PRECAUTIONS](#)).

4 DOSAGE AND ADMINISTRATION

4.1 Dosing Considerations

- Use of Teva-Tofacitinib with other potent systemic immunosuppressants should be avoided. Combined use of tofacitinib with potent immunosuppressants or biologic DMARDs (tumor necrosis factor (TNF) antagonists, interleukin 1 receptor (IL-1R) antagonists, IL-6R antagonists, anti- CD20 monoclonal antibodies, IL-17 antagonists, IL-12/IL-23 antagonists and selective co-stimulation modulators) has not been studied in RA, PsA and UC patients. There is a risk of added immunosuppression when tofacitinib is coadministered with potent immunosuppressive drugs (e.g. azathioprine, tacrolimus, cyclosporine).
- Teva-Tofacitinib should not be initiated in patients with an absolute neutrophil count (ANC) less than 1×10^9 cells/L, hemoglobin (Hgb) levels <90 g/L, or with a lymphocyte count less than 0.5×10^9 cells/L (see [7 WARNINGS AND PRECAUTIONS](#)).
- Teva-Tofacitinib is contraindicated in patients with severe hepatic impairment

4.2 Recommended Dose and Dosage Adjustment

Rheumatoid Arthritis

Teva-Tofacitinib is to be used in combination with methotrexate.

Teva-Tofacitinib monotherapy may be considered in cases of intolerance to methotrexate and to one or more DMARDs.

The recommended dose of Teva-Tofacitinib is 5 mg administered twice daily (BID).

A dosage of Teva-Tofacitinib 10 mg BID is not recommended for the treatment of rheumatoid arthritis (see [7 WARNINGS AND PRECAUTIONS](#)).

Psoriatic Arthritis

The recommended dose of Teva-Tofacitinib is 5 mg administered BID in combination with MTX or another csDMARD.

A dosage of Teva-Tofacitinib 10 mg BID is not recommended for the treatment of psoriatic arthritis (see [7 WARNINGS and PRECAUTIONS, Thrombosis](#)).

Ulcerative Colitis

The recommended dose is 10 mg given orally BID for induction for at least 8 weeks and 5 mg given BID for maintenance.

Depending on therapeutic response; 10 mg BID may also be used for maintenance in some patients. However, the lowest effective dose possible should be used for maintenance therapy to minimize adverse effects (see [7 WARNINGS AND PRECAUTIONS](#)).

Teva-Tofacitinib induction therapy should be discontinued in patients who show no evidence of adequate therapeutic benefit by Week 16.

In patients who have responded to treatment with Teva-Tofacitinib, corticosteroids may be cautiously reduced and/or discontinued in accordance with standard of care.

Dose Interruption or Discontinuation due to Serious Infections and Cytopenias

- Avoid use of Teva-Tofacitinib if a patient develops a serious infection until the infection is controlled.
- Dose interruption is recommended for management of anemia, lymphopenia, and neutropenia as described in Table 1 (see [7 WARNINGS AND PRECAUTIONS](#) and [8 ADVERSE REACTIONS](#)).

Table 1: Laboratory measures and dose adjustment recommendations

| Laboratory Measure | Lab Value | Recommendation |
|---------------------------------|--|---|
| Hemoglobin | <20 g/L decrease and ≥90 g/L | Maintain dose |
| | ≥20 g/L decrease or <80 g/L (Confirmed by repeat testing) | Interrupt the administration of Teva-Tofacitinib until hemoglobin values have normalized (above 80 g/L) |
| Absolute Neutrophil Count (ANC) | >1 x 10 ⁹ cells/L | Maintain dose |
| | 0.5-1 x 10 ⁹ cells/L | For persistent decreases in this range, interrupt or reduce administration with Teva-Tofacitinib until ANC is >1x 10 ⁹ cells/L <ul style="list-style-type: none"> • For patients receiving Teva-Tofacitinib 5 mg BID, interrupt Teva-Tofacitinib dosing. When ANC is >1 x 10⁹ cells/L, resume Teva-Tofacitinib 5 mg BID. UC patients: <ul style="list-style-type: none"> • For patients receiving Teva-Tofacitinib 10 mg BID, reduce dose to Teva-Tofacitinib 5 mg BID. When ANC is >1 x 10⁹ cells/L, increase to Teva-Tofacitinib 10 mg BID based on clinical response. |
| | <0.5 x 10 ⁹ cells/L (Confirmed by repeat testing) | Discontinue treatment with Teva-Tofacitinib |
| Absolute Lymphocyte Count | ≥ 0.5 x 10 ⁹ cells/L | Maintain dose |
| | < 0.5 x 10 ⁹ cells/L (Confirmed by repeat testing) | Discontinue Teva-Tofacitinib |

Dose Modification in Patients with Renal or Hepatic Impairment, or Due to Drug Interactions

- Use Teva-Tofacitinib with caution in patients with moderate (CLcr ≥30 and <60 mL/min) or severe (CLcr ≥15 and <30 mL/min) renal insufficiency (including patients with ESRD but not limited to those

undergoing hemodialysis). Modified dosing is indicated in Table 2.

- For patients undergoing hemodialysis, dose should be administered after the dialysis session on dialysis days. If a dose was taken before the dialysis procedure, supplemental doses are not recommended in patients after dialysis.
- Patients with severe renal insufficiency should remain on a reduced dose even after hemodialysis.
- Use Teva-Tofacitinib with caution in patients with moderate hepatic impairment. Modified dosing is indicated in Table 2.
- Modified dosing of Teva-Tofacitinib is recommended with concomitant CYP inhibitors as indicated in Table 3.
- Coadministration of potent inducers of CYP3A4 with Teva-Tofacitinib is not recommended. Coadministration of potent inducers of CYP3A4 (e.g. rifampin) with Teva-Tofacitinib may result in loss of efficacy or reduced clinical response to Teva-Tofacitinib (see [9.4 Drug-Drug Interactions](#)).

Table 2: Recommended dose adjustment Teva-Tofacitinib in patients with renal insufficiency or hepatic impairment

| | | Teva-Tofacitinib | |
|--|--|------------------|-----------------|
| Indicated dose (in normal renal/hepatic function) | | 5 mg BID | 10 mg BID |
| Modified dosing | Moderate Renal insufficiency (CLcr \geq 30 and <60 mL/min) | 5 mg once daily | 5 mg BID |
| | Severe Renal insufficiency (CLcr \geq 15 and <30 mL/min) | | |
| | Moderate hepatic impairment | | |
| | Severe hepatic impairment | Contraindicated | Contraindicated |

Table 3: Recommended dose adjustment of Teva-Tofacitinib in patients with CYP modifiers

| | | Teva-Tofacitinib | |
|-----------------|---|------------------|-----------------|
| Indicated dose | | 5 mg BID | 10 mg BID |
| Modified dosing | Patients receiving: <ul style="list-style-type: none"> ● Potent CYP3A4 inhibitors (e.g. ketoconazole), or ● a moderate CYP3A4 inhibitor and a potent CYP2C19 inhibitor (e.g. fluconazole) | 5 mg once daily | 5 mg BID |
| | Patients receiving: <ul style="list-style-type: none"> ● Potent CYP3A4 inducers (e.g. rifampin) | Not recommended | Not recommended |

Special Populations

Geriatrics (>65 years): No dosage adjustment is required in patients aged 65 years and older (see [7.1.4 Geriatrics](#) and [10 CLINICAL PHARMACOLOGY](#)).

Pediatrics (<18 years of age): Health Canada has not authorized an indication for pediatric use. No data are available regarding the safety and efficacy of tofacitinib in children aged from neonates to less than 18 years of age. Therefore Teva-Tofacitinib should not be used in this patient population (see [1.1 Pediatrics, 7.1.3 Pediatrics](#) and [10 CLINICAL PHARMACOLOGY](#)).

4.4 Administration

Teva-Tofacitinib is to be taken orally with or without food.

4.5 Missed Dose

For a missed dose, resume at the next scheduled dose.

5 OVERDOSE

There is no experience with overdose of tofacitinib. There is no specific antidote for overdose with tofacitinib. Treatment should be symptomatic and supportive. In case of an overdose, it is recommended that the patient be monitored for signs and symptoms of adverse reactions. Patients who develop adverse reactions should receive appropriate treatment.

Pharmacokinetic data up to and including a single dose of 100 mg in healthy volunteers indicates that more than 95% of the administered dose is expected to be eliminated within 24 hours.

| |
|---|
| For management of a suspected drug overdose, contact your regional poison control centre. |
|---|

6 DOSAGE FORMS, STRENGTHS, COMPOSITION AND PACKAGING

Table 4: Dosage Forms, Strengths, Composition and Packaging

| Route of Administration | Dosage Form / Strength/ Composition | Non-medicinal Ingredients |
|-------------------------|---|---|
| Oral | Teva-Tofacitinib 5 mg Tablets (as tofacitinib citrate) | The core contains croscarmellose sodium, lactose monohydrate, magnesium stearate, microcrystalline cellulose. The film coat contains HPMC/hypromellose, macrogol/PEG, titanium dioxide. |
| | Teva-Tofacitinib 10 mg Tablets (as tofacitinib citrate) | The core contains croscarmellose sodium, lactose monohydrate, magnesium stearate, microcrystalline cellulose. The film coat contains FD&C blue #2/indigo carmine aluminum lake, HPMC/hypromellose, macrogol/PEG, titanium dioxide. |

Teva-Tofacitinib

Tablet: 5 mg tofacitinib (as tofacitinib citrate) are white, round, film-coated tablets, debossed with “TV” on one side and “E71” on the other side. The tablets are available in HDPE bottles with desiccant and child-resistant caps containing 30, 60 or 100 film-coated tablets.

Tablet: 10 mg tofacitinib (as tofacitinib citrate) are blue, round, film-coated tablets, debossed with “TV” on one side and “E77” on the other side. The tablets are available in HDPE bottles with desiccant and child-resistant caps containing 30 or 60 film-coated tablets.

7 WARNINGS AND PRECAUTIONS

Please see [3 SERIOUS WARNINGS AND PRECAUTIONS BOX](#).

Carcinogenesis and Mutagenesis

In patients treated with tofacitinib, malignancies were observed in clinical studies and the post-marketing setting including but not limited to: lymphomas, lung cancer, breast cancer, colorectal cancer, gastric cancer, melanoma, prostate cancer, pancreatic cancer, thyroid cancer and renal cell carcinoma (see [3 SERIOUS WARNINGS AND PRECAUTIONS BOX](#) and [8.2 Clinical Trial Adverse Reactions](#)).

An increase in malignancies (excluding NMSC) was observed in patients treated with tofacitinib compared with TNF inhibitors in a post-authorization safety study (see [8.2 Clinical Trial Adverse Reactions](#)). Malignancies (excluding NMSC) were more common in geriatric patients and in patients who were current or past smokers.

Lung cancers were observed in patients treated with tofacitinib and an increased rate was observed in patients treated with tofacitinib 10 mg BID compared with TNF inhibitors in a post-authorization safety study. Patients with rheumatoid arthritis and taking tofacitinib may be at higher risk than the general

population for the development of lung cancer.

Lymphomas were also observed in patients treated with tofacitinib in a post-authorization safety study (see [8.2 Clinical Trial Adverse Reactions](#)).

Caution should be used in treating geriatric patients, patients who are current or past smokers, and patients with other malignancy risk factors.

Consider the risks and benefits of tofacitinib treatment prior to initiating therapy in patients with current or a history of malignancy other than a successfully treated non-melanoma skin cancer (NMSC) or when considering continuing tofacitinib in patients who develop a malignancy. Recommendations for NMSC are presented below.

Rheumatoid Arthritis

In the 5 controlled clinical studies, 5 malignancies (excluding NMSC) were diagnosed in patients receiving tofacitinib 5 mg BID, and 8 malignancies (excluding NMSC) were diagnosed in patients receiving tofacitinib 10 mg BID, compared to 0 malignancies (excluding NMSC) in patients in the placebo/placebo plus DMARD group during the first 12 months. Lymphomas and solid cancers have also been observed in the long-term extension study in patients treated with tofacitinib (see [8.2 Clinical Trial Adverse Reactions](#)). Patients with RA particularly those with highly active disease, may be at a higher risk (several fold) than the general population for the development of lymphoma.

In Phase 2B, controlled dose-ranging trials in de-novo renal transplant patients, all of whom received induction therapy with basiliximab, high dose corticosteroids, and mycophenolic acid products, Epstein Barr Virus-associated post-transplant lymphoproliferative disorder was observed in 5 out of 218 patients treated with tofacitinib (2.3%) compared to 0 out of 111 patients treated with cyclosporine.

Psoriatic Arthritis

In the 2 controlled PsA clinical trials, there were 3 malignancies (excluding NMSC) in 474 patients receiving tofacitinib plus csDMARD (6 to 12 months exposure) compared with 0 malignancies in 236 patients in the placebo plus csDMARD group (3 months exposure) and 0 malignancies in 106 patients in the adalimumab plus DMARD group (12 months exposure). Malignancies have also been observed in the long-term extension study in PsA patients treated with tofacitinib.

Ulcerative Colitis

In the 4 controlled clinical studies for ulcerative colitis (up to 52-week treatment), no malignancies (excluding NMSC) were reported with tofacitinib. In the long-term extension open-label study, malignancies (excluding NMSC) have been observed in patients treated with tofacitinib 10 mg BID, including solid cancers and lymphoma.

Non-Melanoma Skin Cancer: Non-melanoma skin cancers (NMSCs) have been reported in patients treated with tofacitinib. NMSC is a dose related adverse reaction, with a greater risk in patients treated with 10 mg BID of tofacitinib than in patients treated with 5 mg BID. An increase in overall NMSCs, including cutaneous squamous cell carcinomas was observed in patients treated with tofacitinib compared to TNF inhibitors in a post-authorization safety study (see [8.2 Clinical Trial Adverse Reactions](#)). Caution should be used when treating geriatric patients and patients with a prior history of NMSC, where a higher incident of NMSC was observed. Periodic skin examination is recommended.

In the UC 52-week maintenance study, NMSC was reported in 3 patients (1.5%) treated with 10 mg BID, as compared with no reported events in patients treated with 5 mg BID and 1 patient (0.5%) treated with placebo. In the long-term open label extension study, NMSC was reported in 6 patients in the 10 mg BID group and 2 patients in the 5 mg BID group.

Cardiovascular

Heart Rate Decrease and PR Interval Prolongation: Tofacitinib caused a decrease in heart rate and a prolongation of the PR interval (see [7 WARNINGS AND PRECAUTIONS – Monitoring and Laboratory Tests](#) and [8 ADVERSE REACTIONS](#)). Caution should be observed in patients with a low heart rate at baseline (<60 beats per minute), a history of syncope or arrhythmia, sick sinus syndrome, sinoatrial block, atrioventricular (AV) block, ischemic heart disease, or congestive heart failure. Concomitant medications that result in a decrease in heart rate and/or PR interval prolongation should be avoided to the extent possible during treatment with Teva-Tofacitinib (see [9 DRUG INTERACTIONS](#)).

Thrombosis

Thrombosis, including pulmonary embolism, deep venous thrombosis, and arterial thrombosis, was observed at an increased incidence in patients treated with tofacitinib in a post-authorization safety study. In this post-authorization safety study, patients treated with tofacitinib 10 mg BID had a higher rate of all-cause mortality, including sudden CV death, and thrombosis compared to those treated with tofacitinib 5 mg given BID or TNF inhibitors. Many of these events were serious and some resulted in death (see [3 SERIOUS WARNINGS AND PRECAUTIONS BOX](#)).

In a long-term extension study in patients with ulcerative colitis (UC), four cases of pulmonary embolism were reported in patients taking tofacitinib 10 mg BID, including one death in a patient with advanced cancer.

A dosage of Teva-Tofacitinib 10 mg BID is not recommended for the treatment of RA or PsA (see [4 DOSAGE AND ADMINISTRATION](#)).

For the treatment of UC, use Teva-Tofacitinib at the lowest effective dose and for the shortest duration needed to achieve/maintain therapeutic response (see [4 DOSAGE AND ADMINISTRATION](#)).

Avoid Teva-Tofacitinib in patients that may be at increased risk of thrombosis. Discontinue Teva-Tofacitinib and promptly evaluate patients with symptoms of thrombosis.

Major Adverse Cardiovascular Events (including Myocardial Infarction)

Major adverse cardiovascular events (MACE), including events of myocardial infarction, were observed in patients treated with tofacitinib 5 mg BID, tofacitinib 10 mg BID or TNF inhibitors in a post-authorization safety study. An increase in non-fatal myocardial infarctions was observed in patients treated with tofacitinib compared to TNF inhibitor (see [8.2 Clinical Trial Adverse Reactions](#)). MACE, including events of myocardial infarction, were more common in geriatric patients and in patients who were current or past smokers (see [3 SERIOUS WARNINGS AND PRECAUTIONS BOX](#)). Caution should be used in treating geriatric patients, patients who are current or past smokers, and patients with other CV risk factors.

Retinal venous thrombosis

Retinal venous thrombosis (RVT) has been reported in patients treated with tofacitinib (see [8.2 Clinical Trial Adverse Reactions, Thromboembolism, Table 10](#) and [14.1 Clinical Trials by Indication, Rheumatoid Arthritis, Study RA-VI](#)). Patients should be advised to promptly seek medical care if they experience symptoms suggestive of RVT.

Driving and Operating Machinery

No formal studies have been conducted on the effects on the ability to drive and use machines.

Fractures

Fractures of multiple types, including osteoporotic fractures, have been observed in patients treated with tofacitinib in clinical studies and the postmarketing setting (see [8.2 Clinical Trial Adverse Reactions](#)).

Caution should be applied when using Teva-Tofacitinib in patients with known risk factors for fractures such as geriatric patients, female patients, and patients using corticosteroids.

Gastrointestinal

Events of gastrointestinal perforation have been reported with tofacitinib in RA patients, in clinical trials and in the post-market setting. The role of JAK inhibition in these events is not known. Many patients who developed gastrointestinal perforations were taking concomitant nonsteroidal anti-inflammatory drugs (NSAIDs) and/or corticosteroids. The relative contribution of these concomitant medications versus tofacitinib to the development of gastrointestinal perforations is not known.

There was no discernable difference in frequency of gastrointestinal perforation between the placebo and the tofacitinib arms in clinical trials of patients with UC, and many of them were receiving background corticosteroids.

Teva-Tofacitinib should be used with caution in patients who may be at increased risk for gastrointestinal perforation (e.g., use of concomitant NSAIDs and/or corticosteroids, patients with a history of diverticulitis). Patients presenting with new onset abdominal symptoms should be evaluated promptly for early identification of gastrointestinal perforation (see [8 ADVERSE REACTIONS](#)).

Hematologic

Anemia: Treatment with tofacitinib has been associated with decreases in hemoglobin levels. Evaluate hemoglobin prior to initiation of Teva-Tofacitinib (see [7 WARNINGS AND PRECAUTIONS – Monitoring and Laboratory Tests](#) and [4.2 Recommended Dose and Dosage Adjustment](#)). Avoid initiation of Teva-Tofacitinib treatment in patients with low hemoglobin values (i.e., <90 g/L). Treatment with Teva-Tofacitinib should be interrupted in patients who develop hemoglobin levels <80 g/L or whose hemoglobin level drops >20 g/L on treatment.

For recommended monitoring and dose modification based on hemoglobin results, see [7 WARNINGS AND PRECAUTIONS – Monitoring and Laboratory Tests](#) and [4.2 Recommended Dose and Dosage Adjustment](#).

Lymphopenia: Treatment with tofacitinib was associated with initial lymphocytosis at one month of exposure followed by a gradual decrease in mean lymphocyte counts below the baseline of approximately 10% during 12 months of therapy. Lymphocyte counts less than 0.5×10^9 cells/L were associated with an increased incidence of treated and serious infections. Evaluate lymphocyte count prior to initiation of Teva-Tofacitinib approximately 4-8 weeks after initiation with Teva-Tofacitinib treatment, and every 3 months thereafter.

Avoid initiation of Teva-Tofacitinib treatment in patients with a low lymphocyte count (i.e., less than 0.5×10^9 cells/L). In patients who develop a confirmed absolute lymphocyte count less than 0.5×10^9 cells/L, Teva-Tofacitinib should be discontinued.

For recommended monitoring and dose modifications based on lymphocyte counts see [7 WARNINGS AND PRECAUTIONS – Monitoring and Laboratory Tests](#) and [4.2 Recommended Dose and Dosage Adjustment](#).

Neutropenia: Treatment with tofacitinib was associated with an increased incidence of neutropenia (< 2×10^9 cells/L) compared to placebo. Evaluate neutrophil count prior to initiation of Teva-Tofacitinib

approximately 4-8 weeks after initiation with Teva-Tofacitinib, and every 3 months thereafter.

Avoid initiation of Teva-Tofacitinib treatment in patients with a low neutrophil count (i.e., ANC (absolute neutrophil count) $<1 \times 10^9$ cells/L). For patients who develop a persistent ANC of 0.5 to 1×10^9 cells/L, interrupt dosing until ANC is $>1 \times 10^9$ cells/L. In patients who develop an absolute neutrophil count $<0.5 \times 10^9$ cells/L, discontinue treatment.

For recommended monitoring and dose modification based on ANC, see [7 WARNINGS AND PRECAUTIONS – Monitoring and Laboratory Tests](#) and [4.2 Recommended Dose and Dosage Adjustment](#).

Lipid Elevations: Treatment with tofacitinib was associated with increases in lipid parameters including total cholesterol, low-density lipoprotein (LDL) cholesterol, and high-density lipoprotein (HDL) cholesterol (see **8 ADVERSE REACTIONS**).

Maximum effects were generally observed within 6 weeks. The effect of these lipid parameter elevations on cardiovascular morbidity and mortality has not been determined.

Assessment of lipid parameters should be performed at baseline and approximately 4-8 weeks following initiation of Teva-Tofacitinib therapy, and every 6 months thereafter (see [7 WARNINGS AND PRECAUTIONS – Monitoring and Laboratory Tests](#)). Patients should be managed according to local clinical guidelines for the management of hyperlipidemia.

Hepatic/Biliary/Pancreatic

Teva-Tofacitinib is contraindicated in patients with severe hepatic impairment.

Treatment with tofacitinib was associated with an increased incidence of liver enzyme elevation compared to placebo (see [8 ADVERSE REACTIONS](#)).

Evaluate liver enzymes before initiating Teva-Tofacitinib and thereafter according to routine patient management (see [7 WARNINGS AND PRECAUTIONS – Monitoring and Laboratory Tests](#)). Prompt investigation of the causes of liver enzyme elevations is recommended to identify potential cases of drug-induced liver injury (DILI). If increases in ALT (alanine transaminase) or AST (aspartate transaminase) are observed and DILI is suspected, the administration of Teva-Tofacitinib should be interrupted until the diagnosis is excluded.

Most of the liver enzyme abnormalities in RA and PsA patients occurred in studies with background DMARD (primarily methotrexate) therapy.

One case of DILI was reported in a RA patient treated with tofacitinib 10 mg BID for approximately 2.5 months. The patient developed symptomatic elevations of AST and ALT with values greater than 3x ULN associated concurrently with total bilirubin value greater than 2x ULN, which required hospitalization and a liver biopsy.

In UC patients, tofacitinib treatment with 5 and 10 mg BID was also associated with an increased incidence of liver enzyme elevation compared to placebo, with a trend for higher incidence with the 10 mg BID as compared to the 5 mg BID (see [8 ADVERSE REACTIONS](#)).

One patient treated with tofacitinib 10 mg BID in the maintenance UC study experienced an increase in liver enzymes which decreased upon discontinuation of treatment. The case was adjudicated as possible DILI, while noting ultrasound findings of fatty liver.

The impact of tofacitinib on chronic viral hepatitis reactivation is unknown. Tofacitinib has not been studied in patients with positive hepatitis B virus or hepatitis C virus serology and should therefore not be used in these populations.

Tofacitinib has not been studied in patients with severe hepatic impairment and should not be used in

these patients. Dose adjustment of Teva-Tofacitinib is recommended for patients with moderate hepatic impairment (see [4 DOSAGE AND ADMINISTRATION](#) and [10 CLINICAL PHARMACOLOGY](#)).

Immune

Hypersensitivity Reactions: Reactions such as angioedema and urticaria that may reflect drug hypersensitivity have been observed in patients treated with tofacitinib. Some events were serious. If a hypersensitivity reaction is suspected, promptly discontinue tofacitinib while evaluating the potential cause or causes of the reaction (see [2 CONTRAINDICATIONS](#) and [8 ADVERSE REACTIONS](#)).

Immunocompromised Patients: Tofacitinib can increase the risk of infections and immunosuppression when co-administered with potent immunosuppressants such as cyclosporine, azathioprine and tacrolimus. Combined use of tofacitinib with potent immunosuppressive drugs has not been studied and is not recommended (see [9.4 Drug-Drug Interactions](#)).

Immunizations: No data are available on the secondary transmission of infection by live vaccines to patients receiving tofacitinib. It is recommended that all patients be brought up to date with all immunizations in agreement with current immunization guidelines prior to initiating Teva-Tofacitinib therapy and that live vaccines not be given concurrently with Teva-Tofacitinib. The interval between live vaccinations and initiation of tofacitinib therapy should be in accordance with current vaccination guidelines regarding immunomodulatory agents.

In patients being considered for Teva-Tofacitinib therapy, live zoster vaccine should only be administered to patients with a known history of chickenpox or those that are seropositive for varicella zoster virus. Vaccination should occur at least 2 weeks but preferably 4 weeks before initiating immunomodulatory agents such as Teva-Tofacitinib.

In a clinical trial, a varicella naïve patient treated with tofacitinib and methotrexate developed disseminated infection with the vaccine strain of the varicella zoster virus 16 days after vaccination. A satisfactory immune response to the vaccine was developed 6 weeks post-vaccination.

Antibody levels after vaccination may be lower in patients treated with tofacitinib (see [10.2 Pharmacodynamics](#)).

Infections:

Serious and sometimes fatal infections due to bacterial, mycobacterial, invasive fungal, viral, or other opportunistic pathogens have been reported in RA patients receiving immunomodulatory agents, including tofacitinib. The most common serious infections reported with tofacitinib included pneumonia, urinary tract infection, cellulitis, herpes zoster, bronchitis, septic shock, diverticulitis, gastroenteritis, appendicitis and sepsis. Among opportunistic infections, tuberculosis and other mycobacterial infections, cryptococcus, histoplasmosis, esophageal candidiasis, pneumocystosis, multidermatomal herpes zoster, cytomegalovirus infections, BK virus infections, listeriosis and aspergillosis were reported with tofacitinib (see [8 ADVERSE REACTIONS](#)). Some patients have presented with disseminated rather than localized disease and were often taking concomitant immunomodulating agents such as methotrexate or corticosteroids. Other serious infections that were not reported in clinical studies may also occur (e.g., coccidioidomycosis).

A dose dependent increase in serious infections was observed in patients treated with tofacitinib compared to TNF inhibitors in a post-authorization safety study (see [8 ADVERSE REACTIONS](#)). Some of these serious infections resulted in death. Opportunistic infections were also reported in the study.

Patients treated with tofacitinib 10 mg BID are at higher risk of serious infections, and herpes zoster infections compared to those treated with 5 mg BID. The incidence rate per 100 person-years (PYs) for

herpes zoster opportunistic infections in the UC 52-week maintenance study was higher in patients treated with tofacitinib 10 mg BID (6.64) as compared to tofacitinib 5 mg BID (2.05) or placebo (0.97) (see [8 ADVERSE REACTIONS](#)).

Teva-Tofacitinib should not be administered in patients with an active infection, including localized infections. The risks and benefits of treatment should be considered prior to initiating Teva-Tofacitinib in patients:

- with chronic or recurrent infections,
- who have been exposed to tuberculosis,
- with a history of a serious or an opportunistic infection,
- who have resided or travelled in areas of endemic tuberculosis or endemic mycoses; or
- with underlying conditions that may predispose them to infection.

Patients should be closely monitored for the development of signs and symptoms of infection during and after treatment with Teva-Tofacitinib. Teva-Tofacitinib should be interrupted if a patient develops a serious infection, an opportunistic infection, or sepsis. A patient who develops a new infection during treatment with Teva-Tofacitinib should undergo prompt and complete diagnostic testing appropriate for an immunocompromised patient, appropriate antimicrobial therapy should be initiated, and the patient should be closely monitored.

As there is a higher incidence of infections in the geriatric and in the diabetic populations in general, caution should be used when treating geriatric patients and patients with diabetes. Caution is also recommended in patients with a history of chronic lung disease as they may be more prone to infections. Events of interstitial lung disease (some of which had a fatal outcome) have been reported in RA patients treated with tofacitinib in clinical trials and in the post-marketing setting.

Risk of infection may be higher with increasing degrees of lymphopenia and consideration should be given to lymphocyte counts when assessing individual patient risk of infection (see [7 WARNINGS AND PRECAUTIONS – Monitoring and Laboratory Tests](#)).

Treatment with tofacitinib was associated with increased rates of infections in Asian patients compared to other races (see [7.1.5 Asian Patients](#) and [8 ADVERSE REACTIONS](#)). Teva-Tofacitinib should be used with caution in this population.

Tuberculosis

Patients should be evaluated and tested for latent or active tuberculosis (TB) infection prior to administration of Teva-Tofacitinib and periodically (e.g., annually) while taking Teva-Tofacitinib.

Teva-Tofacitinib should not be given to patients with active TB.

Antituberculosis therapy should also be considered prior to administration of Teva-Tofacitinib in patients with a past history of latent or active tuberculosis in whom an adequate course of treatment cannot be confirmed, and for patients with a negative test for latent tuberculosis but have risk factors for tuberculosis infection.

Patients with latent tuberculosis should be treated with standard antimycobacterial therapy before administering Teva-Tofacitinib.

Patients should be closely monitored for the development of signs and symptoms of tuberculosis, including patients who tested negative for latent tuberculosis infection prior to initiating therapy.

Viral Reactivation

Viral reactivation, including cases of herpes virus reactivation (e.g., herpes zoster) were observed in clinical studies with tofacitinib. An increase in herpes zoster events was observed in patients treated with

tofacitinib compared to TNF inhibitors in a post-authorization safety study (see [8 ADVERSE REACTIONS](#)). Post-marketing cases of hepatitis B reactivation have been reported in patients treated with tofacitinib (see [8 ADVERSE REACTIONS](#)). The impact of tofacitinib on chronic viral hepatitis reactivation is unknown. Patients who screened positive for hepatitis B or C were excluded from clinical trials. Screening for viral hepatitis should be performed in accordance with clinical guidelines before starting therapy with Teva-Tofacitinib.

Monitoring and Laboratory Tests

Lipid tests should be performed at baseline, approximately 4-8 weeks after initiation with Teva-Tofacitinib and every 6 months thereafter. Patients should be managed according to clinical guidelines for the management of hyperlipidemia (see [4.2 Recommended Dose and Dosage Adjustment](#) and [7 WARNINGS AND PRECAUTIONS](#)).

Liver enzyme tests are recommended before initiating Teva-Tofacitinib treatment and thereafter according to routine patient management. If increases in ALT or AST are observed during routine patient management and DILI is suspected, the administration of Teva-Tofacitinib should be interrupted until this diagnosis has been excluded (see [4.2 Recommended Dose and Dosage Adjustment](#) and [7 WARNINGS AND PRECAUTIONS](#)).

Assessment of renal function is recommended prior to initiation of Teva-Tofacitinib (see [4.2 Recommended Dose and Dosage Adjustment](#) and [7 WARNINGS AND PRECAUTIONS](#)).

Lymphocyte, neutrophil and hemoglobin tests should be performed at baseline, approximately 4-8 weeks after initiation with Teva-Tofacitinib treatment, and every 3 months thereafter (see [4.2 Recommended Dose and Dosage Adjustment](#) and [7 WARNINGS AND PRECAUTIONS](#)).

Vital signs: Patients should be monitored for pulse rate and blood pressure at baseline and periodically during treatment with Teva-Tofacitinib (see [7 WARNINGS AND PRECAUTIONS – Cardiovascular](#), [8 ADVERSE REACTIONS](#), and [9.4 Drug-Drug Interactions](#)).

Musculoskeletal

Treatment with tofacitinib was associated with increases in creatine kinase (CK). Maximum effects were generally observed within 6 months. Rhabdomyolysis was reported in one patient treated with tofacitinib. Creatine kinase levels should be checked in patients with symptoms of muscle weakness and/or muscle pain to evaluate for evidence of rhabdomyolysis. Increases in CK were reported more frequently in patients treated with tofacitinib 10 mg as compared to those treated with 5 mg BID (see [8 ADVERSE REACTIONS](#)).

Renal

Dosage adjustment of Teva-Tofacitinib is recommended in patients with moderate and severe renal impairment (see, [4.2 Recommended Dose and Dosage Adjustment](#), and [10 CLINICAL PHARMACOLOGY](#)). In clinical trials, tofacitinib was not evaluated in patients with baseline creatinine clearance values (estimated by the Cockcroft-Gault equation) less than 40 mL/min.

Reproductive Health: Female and Male Potential

- **Fertility:** Based on findings in animal studies, tofacitinib may cause decreased fertility when administered to females (see [16 NON-CLINICAL TOXICOLOGY](#)).
- **Teratogenic Risk:** Based on findings in animal studies, tofacitinib may cause fetal harm when administered to a pregnant woman (see [2 CONTRAINDICATIONS](#)). Administration of tofacitinib to rats and rabbits during organogenesis caused increases in fetal malformations (see [16 NON-CLINICAL](#)

[TOXICOLOGY](#)). Pregnant women should be advised of the potential risk to a fetus. Females of reproductive potential should be advised to use effective contraception during treatment with Teva-Tofacitinib and for 4 to 6 weeks following completion of therapy (see [7.1.1 Pregnant Women](#)).

Respiratory

Interstitial Lung Disease: Events of interstitial lung disease (ILD) have been reported in RA clinical trials with tofacitinib, although the role of JAK inhibition in these events is not known. All patients who developed ILD were taking concomitant methotrexate, corticosteroids and/or sulfasalazine, which have been associated with ILD. Asian patients had an increased risk of ILD (see [7.1.5 Asian Patients](#)).

Teva-Tofacitinib should be used with caution in patients with a risk or history of ILD.

7.1 Special Populations

7.1.1 Pregnant Women

Teva-Tofacitinib is contraindicated during pregnancy (see [2 CONTRAINDICATIONS](#)). There are no adequate and well-controlled studies on the use of tofacitinib in pregnant women. Tofacitinib has been shown to be teratogenic in rats and rabbits, and have effects in rats on female fertility, parturition, and peri/postnatal development (see [16 NON-CLINICAL TOXICOLOGY](#)).

Women of reproductive potential should be advised to use effective contraception during Teva-Tofacitinib treatment and for 4 to 6 weeks after the last dose.

7.1.2 Breast-feeding

Teva-Tofacitinib is contraindicated in women who breastfeed (see [2 CONTRAINDICATIONS](#)). Tofacitinib was secreted in milk of lactating rats. It is not known whether tofacitinib is excreted in human milk. (see [16 NON-CLINICAL TOXICOLOGY](#)).

7.1.3 Pediatrics

Pediatrics (<18 years of age): No data are available to Health Canada; therefore, Health Canada has not authorized an indication for pediatric use.

7.1.4 Geriatrics

Geriatrics (>65 years of age): The frequency of adverse events including serious infections, all-cause mortality, cardiovascular events, malignancies, non-melanoma skin cancer, gastrointestinal perforations, interstitial lung disease, venous thromboembolism, and arterial thromboembolism among tofacitinib treated subjects 65 years of age and older was higher than among those under the age of 65. Caution should be used when treating geriatric patients with Teva-Tofacitinib (see [4 DOSAGE AND ADMINISTRATION](#) and [10 CLINICAL PHARMACOLOGY](#)).

7.1.5 Asian Patients

Asian patients have an increased risk of herpes zoster and opportunistic infections. Asian patients with RA also have an increased risk of ILD. An increased incidence of some adverse events such as elevated transaminases (ALT, AST) and decreased white blood cells (WBCs) were also observed. Therefore, Teva-Tofacitinib should be used with caution in Asian patients (see [8 ADVERSE REACTIONS](#)).

8 ADVERSE REACTIONS

8.1 Adverse Reaction Overview

Rheumatoid Arthritis, Psoriatic arthritis

The most common serious adverse reactions (SAEs) in rheumatoid arthritis clinical trials were osteoarthritis and serious infections, including pneumonia, cellulitis, herpes zoster, and urinary tract infection.

The most commonly reported adverse reactions during the first 3 months in controlled clinical trials in rheumatoid arthritis (occurring in $\geq 2\%$ of patients treated with tofacitinib monotherapy or in combination with DMARDs) were upper respiratory tract infections, headache, nasopharyngitis, and diarrhea. Additionally, bronchitis, urinary tract infection, herpes zoster, RA, back pain and hypertension were also reported in the 5 mg BID tofacitinib group in the long-term extension trial.

The most common adverse reactions in rheumatoid arthritis clinical trials that resulted in discontinuation of tofacitinib were infections. Pneumonia was the most common adverse reactions leading to discontinuation of therapy, followed by blood creatinine increased and herpes zoster.

Asian patients had higher rates of herpes zoster, opportunistic infections, elevated transaminases (ALT, AST) and decreased WBCs. Asian patients with RA also have an increased risk of ILD (see [7.1.5 Asian Patients](#)). Therefore, Teva-Tofacitinib should be used with caution in Asian patients.

Ulcerative Colitis

In the induction studies, the most common categories of serious adverse events were gastrointestinal disorders and infections. The most common serious adverse events (excluding events reported as UC) were abdominal pain, anal abscess, and drug hypersensitivity. The most common adverse events ($\geq 5\%$) were headache and nasopharyngitis.

In the maintenance study, the most common categories of serious adverse events were gastrointestinal disorders, infections, injuries, and nervous system disorders. All serious adverse events were single reports (excluding events reported as UC). The most common adverse events ($\geq 5\%$) (excluding events reported as UC) in patients treated with 5 mg BID were nasopharyngitis, arthralgia, headache, and upper respiratory tract infection. In patients treated with 10 mg BID, the most common adverse events were nasopharyngitis, arthralgia, blood creatine phosphokinase increased, upper respiratory tract infection, rash, hypercholesterolemia, and herpes zoster.

In induction studies, adverse events were reported in 515 subjects (54.9%) treated with 10 mg BID and 155 subjects (55.0%) treated with placebo. In the maintenance study, adverse events were reported in 143 subjects (72.2%) treated with 5 mg BID, 156 subjects (79.6%) treated with 10 mg BID, and 149 subjects (75.3%) treated with placebo.

In induction and maintenance studies, the most frequent reason for study discontinuation was worsening of UC. Excluding discontinuations due to worsening of UC, the proportion of patients who discontinued due to adverse reactions was less than 5% in any of the tofacitinib or placebo treatment groups in these studies.

Four cases of pulmonary embolism were reported in patients taking tofacitinib 10 mg BID.

Overall, the safety profile observed in UC patients treated with tofacitinib was consistent with the safety profile of tofacitinib across indications. Dose-dependent risks seen in patients treated with tofacitinib 10 mg BID in comparison with 5 mg BID include the following: herpes zoster infections, serious infections, and NMSC.

8.2 Clinical Trial Adverse Reactions

Clinical trials are conducted under very specific conditions. The adverse reaction rates observed in the clinical trials; therefore, may not reflect the rates observed in practice and should not be compared to the rates in the clinical trials of another drug. Adverse reaction information from clinical trials may be useful in identifying and approximating rates of adverse drug reactions in real-world use.

Rheumatoid Arthritis

During controlled clinical trials, 8.0% (11.0 events/100 patient-years) of patients in the 5 mg BID in the tofacitinib group were hospitalized due to serious adverse reactions compared to 7.8% (9.1 events/100 patient-years) and 3.8% (13.0 events/100 patient-years) of patients in the adalimumab and placebo group, respectively.

During the first 3 months of Phase 3 studies, serious infections (those requiring parenteral antibiotics or hospitalization) were reported in 0.7% (2.8 events/100 patient-years) and 0.2% (0.6 events/100 patient-years) of patients treated with tofacitinib or placebo, respectively. From 0-12 months, serious infections were reported in 2.4% (3.2 events/100 patient-years) of tofacitinib treated patients (see [7 WARNINGS AND PRECAUTIONS](#)). In a post-authorization safety study, the frequency of pulmonary embolism was increased in patients treated with 10 mg BID tofacitinib (1.65%) compared to the TNF inhibitor (0.21%) and 5 mg BID tofacitinib (0.62%).

Deaths occurred in 0.4% (0.6 events/100 patient-years) of patients in the 5 mg BID tofacitinib group, compared to 0.5% (0.6 events/100 patient-years) and 0.2% (0.5 events/100 patient-years) of patients in the adalimumab and placebo groups, respectively. In a post-authorization safety study, all cause mortality was increased in patients treated with 10 mg BID tofacitinib (2.7%) compared to the TNF inhibitor (1.2%) and 5 mg BID tofacitinib treatment arms (1.8%).

The proportion of patients who discontinued treatment due to any adverse reactions during the first 3 months in double-blind placebo-controlled studies was 7.8% for patients taking 5 mg BID of tofacitinib and 3.7% for placebo-treated patients. In the long-term extension trial, the proportion of patients who discontinued treatment due to any adverse reaction was 24.8% (6.78 events/100 patient-years) for all patients, 27.9% (6.67 events/100 patient-years) for patients taking 5 mg BID of tofacitinib, and 23.8% (6.83 events/100 patient-years) for patients taking 10 mg BID of tofacitinib.

Following completion of the Phase 2/3, open-label, uncontrolled, long-term extension follow-up trial (up to 114 months) from the Phase 2 studies and Phase 3 clinical program, there were 4040 subjects with 16113 patient-years of exposure to tofacitinib. The design of the long-term safety studies allowed for modification of tofacitinib doses according to clinical judgment. This limits the interpretation of the long-term safety data with respect to dose. Tofacitinib 10 mg BID is not recommended in RA patients. Overall, the safety profile of tofacitinib 5 mg BID in the long-term extension study was comparable to what was seen in the controlled clinical trials

Table 5 below lists the adverse events (regardless of causality) occurring in $\geq 1\%$ of patients treated with tofacitinib during the double-blind, placebo-controlled portion of the phase 3 RA studies.

Table 5: Summary of Adverse Events Reported by $\geq 1\%$ of RA Patients Treated with Tofacitinib (All Causalities) - All Phase 3 Studies (up to 3 months)

| Body System/Adverse Event | Tofacitinib 5mg BID N=1216 (%) | Placebo N=681 (%) | Adalimumab 40 mg SC q2w N=204 (%) |
|---|---|------------------------------|--|
| Blood and lymphatic system disorders | | | |
| Anemia | 15 (1.2) | 8 (1.2) | 0 |
| Gastrointestinal disorders | | | |
| Diarrhoea | 45 (3.7) | 16 (2.3) | 2 (1.0) |
| Nausea | 32 (2.6) | 18 (2.6) | 3 (1.5) |
| Dyspepsia | 19 (1.6) | 11 (1.6) | 3 (1.5) |
| Abdominal pain upper | 23 (1.9) | 5 (0.7) | 3 (1.5) |
| Vomiting | 21 (1.7) | 10 (1.5) | 0 |
| Constipation | 16 (1.3) | 6 (0.9) | 2 (1.0) |
| Gastritis | 12 (1.0) | 7 (1.0) | 0 |
| Gastroenteritis | 12 (1.0) | 5 (0.7) | 0 |
| General disorders and administration site conditions | | | |
| Oedema peripheral | 17 (1.4) | 16 (2.3) | 3 (1.5) |
| Pyrexia | 13 (1.1) | 5 (0.7) | 1 (0.5) |
| Infections and infestations | | | |
| Upper respiratory tract infection | 53 (4.4) | 23 (3.4) | 7 (3.4) |
| Nasopharyngitis | 48 (3.9) | 19 (2.8) | 7 (3.4) |
| Urinary tract infection | 25 (2.1) | 12 (1.8) | 7 (3.4) |
| Bronchitis | 14 (1.2) | 10 (1.5) | 4 (2.0) |
| Investigations | | | |
| Alanine aminotransferase increased | 14 (1.2) | 7 (1.0) | 1 (0.5) |
| Metabolism and nutrition disorders | | | |
| Hypercholesterolaemia | 12 (1.0) | 3 (0.4) | 1 (0.5) |
| Musculoskeletal and connective tissue disorders | | | |
| Rheumatoid arthritis | 17 (1.4) | 17 (2.5) | 1 (0.5) |
| Back pain | 18 (1.5) | 5 (0.7) | 1 (0.5) |
| Arthralgia | 13 (1.1) | 16 (2.3) | 4 (2.0) |
| Nervous system disorders | | | |
| Headache | 54 (4.4) | 15 (2.2) | 5 (2.5) |
| Dizziness | 13 (1.1) | 8 (1.2) | 3 (1.5) |
| Vascular disorders | | | |
| Hypertension | 20 (1.6) | 7 (1.0) | 0 |

Overall Infections

In the five controlled trials, during 0 to 3 months exposure, the overall frequency of infections was 20% in the 5 mg BID tofacitinib group, and 18% in the placebo group.

In the long-term extension trial, overall frequency of infections was 67.7% (39.63 events/100 patient-years) in all tofacitinib group; 65.5% of patients (33.22 events/100 patient-years) and 68.4% of patients (42.24 events/100 patient-years) in the 5 mg and 10 mg BID of tofacitinib, respectively.

Infections were also reported in a post-authorization safety study in RA patients who were 50 years or older with at least one additional cardiovascular risk factor.

The most commonly reported infections were upper respiratory tract infections, nasopharyngitis, bronchitis, herpes zoster, and urinary tract infections.

Serious Infections

In the five controlled trials, during the 0 to 3 months exposure, serious infections were reported in 1 patient (0.6 events/100 patient-years) who received placebo and 8 patients (2.8 events/100 patient-years) who received 5 mg BID of tofacitinib.

During 0 to 12 months exposure, the overall frequencies of serious infections were 2.4% (3.2 events/100 patient-years) for the 5 mg BID tofacitinib group.

In the long-term extension trial, the most common serious infections reported with tofacitinib included pneumonia, cellulitis, appendicitis, diverticulitis, gastroenteritis, urinary tract infection, and herpes zoster (see [7 WARNINGS AND PRECAUTIONS](#)).

Serious infections were more frequently reported in subjects taking tofacitinib compared to TNF inhibitors (TNFi), and in patients treated with tofacitinib 10 mg BID compared to those treated with tofacitinib 5 mg BID in a post-authorization safety study (Study RA-VI), as shown in Table 6.

Table 6. Serious Infections in [Study RA-VI](#)

| | Tofacitinib 5 mg BID N = 1455 | Tofacitinib 10 mg BID* N = 1456 | All Tofacitinib N=2911 | TNFi N = 1451 |
|---|--|--|-----------------------------------|--------------------------|
| n (%) | 141 (9.69) | 169 (11.61) | 310 (10.65) | 119 (8.20) |
| IR per 100 PY (95% CI) | 2.86 (2.41, 3.37) | 3.64 (3.11, 4.23) | 3.24 (2.89, 3.62) | 2.44 (2.02, 2.92) |
| Tofacitinib vs TNFi HR (95% CI) | 1.17 (0.92, 1.50) | 1.48 (1.17, 1.87) | 1.32 (1.07, 1.63) | -- |
| * The tofacitinib 10 mg BID treatment group consists of data from patients that were switched from tofacitinib 10 mg BID to tofacitinib 5 mg BID during the trial as a result of a study modification following a recommendation by the Data and Safety Monitoring Board in February 2019. Abbreviations: IR=incidence rate, CI=confidence interval, PY=patient years, HR=hazard ratio | | | | |

Tuberculosis

Cases of tuberculosis have been reported with treatment with tofacitinib.

In the five controlled Phase 3 trials, during 0 to 3 months exposure, no cases of tuberculosis were reported in patients who received placebo or 5 mg BID of tofacitinib.

During 0 to 12 months of exposure, tuberculosis was reported in 0 patients who received 5 mg BID of tofacitinib.

In the long-term extension trial, adjudicated tuberculosis events were reported in 0.6% patients (0.15 events/100 patient-years) who received tofacitinib; 0.4% of patients (0.10 events/100 patient-years) and 0.6% of patients (0.17 events/100 patient-years) in the 5 mg and 10 mg BID of tofacitinib, respectively.

Cases of disseminated tuberculosis were also reported. The median tofacitinib exposure prior to diagnosis of tuberculosis was 10 months (range from 152 to 960 days) (see [7 WARNINGS AND PRECAUTIONS](#)).

Opportunistic Infections (excluding tuberculosis)

In the five controlled Phase 3 trials, during 0 to 3 months exposure, opportunistic infections were reported in 0 patients who received placebo and 2 (0.2%) patients (0.7 events/100 patient-years) who received 5 mg BID of tofacitinib.

During 0 to 12 months of exposure, opportunistic infections were reported in 3 (0.3%) patients (0.3 events/100 patient-years) who received 5 mg BID of tofacitinib.

The median tofacitinib exposure prior to diagnosis of an opportunistic infection was 8 months (range from 41 to 698 days).

The similar frequency of opportunistic infections was observed in the long-term extension trial with tofacitinib treatment up to 114 months.

Malignancy (excluding non-melanoma skin cancer)

In the five Phase 3 controlled trials, during 0 to 3 months exposure, malignancies (excluding non-melanoma skin cancer) were reported in 0 patients who received placebo and 2 (0.2%) patients (0.7 events/100 patient-years) who received 5 mg BID of tofacitinib.

During 0 to 12 months of exposure, malignancies (excluding non-melanoma skin cancer) were reported in 5 (0.4%) patients (0.6 events/100 patient-years) who received 5 mg BID of tofacitinib.

In the long-term extension trial, overall frequency of malignancies (excluding non-melanoma skin cancer) was 3.1% (0.83 events/100 patient-years) in all tofacitinib-treated patients; 3.4% of patients (0.8 events/100 patient-years) and 3% of patients (0.84 events/100 patient-years) in the 5 mg and 10 mg BID of tofacitinib, respectively.

The most common types of malignancy (excluding non-melanoma skin cancer), including malignancies observed during the long-term extension, were lung and breast cancer, followed by gastric, colorectal, renal cell, prostate cancer, lymphoma and malignant melanoma (see [7 WARNINGS AND PRECAUTIONS](#)).

In a post-authorization safety study (Study RA-VI), malignancies (excluding NMSC) were observed more frequently in patients taking tofacitinib compared with patients taking TNFi (Table 7). Frequency of lung cancer was higher in patients taking tofacitinib 10 mg BID compared with patients taking tofacitinib 5 mg BID. Thyroid cancer was observed in 5, 2, and 0 subjects taking tofacitinib 5 mg BID, taking tofacitinib 10 mg BID, and TNFi, respectively.

Table 7: Malignancies (Excluding NMSC), Lymphoma, and Lung Cancer in [Study RA-VI](#)

| | Tofacitinib 5 mg BID N = 1455 | Tofacitinib 10 mg BID* N = 1456 | All Tofacitinib N=2911 | TNFi N = 1451 |
|------------------------------------|--|--|-----------------------------------|--------------------------|
| Malignancies excluding NMSC | | | | |
| n (%) | 62 (4.26) | 60 (4.12) | 122 (4.19) | 42 (2.89) |
| IR (95% CI) per 100 PY | 1.13 (0.87, 1.45) | 1.13 (0.86, 1.45) | 1.13 (0.94, 1.35) | 0.77 (0.55, 1.04) |
| Tofacitinib vs TNFi HR (95% CI) | 1.47 (1.00, 2.18) | 1.48 (1.00, 2.19) | 1.48 (1.04, 2.09)† | -- |
| Lymphoma | | | | |
| n (%) | 4 (0.27) | 6 (0.41) | 10 (0.34) | 1 (0.07) |
| IR (95% CI) per 100 PY | 0.07 (0.02, 0.18) | 0.11 (0.04, 0.24) | 0.09 (0.04, 0.17) | 0.02 (0.00, 0.10) |
| Tofacitinib vs TNFi HR (95% CI) | 3.99 (0.45, 35.70) | 6.24 (0.75, 51.86) | 5.09 (0.65, 39.78) | -- |
| Lung Cancer | | | | |
| n (%) | 13 (0.89) | 17 (1.17) | 30 (1.03) | 7 (0.48) |
| IR (95% CI) per 100 PY | 0.23 (0.12, 0.40) | 0.32 (0.18, 0.51) | 0.28 (0.19, 0.39) | 0.13 (0.05, 0.26) |
| Tofacitinib vs TNFi HR (95% CI) | 1.84 (0.74, 4.62) | 2.50 (1.04, 6.02) | 2.17 (0.95, 4.93) | -- |

* The tofacitinib 10 mg BID treatment group consists of data from patients that were switched from tofacitinib 10 mg BID to tofacitinib 5 mg BID during the trial as a result of a study modification following a recommendation by the Data and Safety Monitoring Board in February 2019.

† The non-inferiority criterion was not met for the primary comparison of the combined tofacitinib doses to TNFi since the upper limit of the 95% CI exceeded the pre-specified non-inferiority criterion of 1.8.

Abbreviations: IR=incidence rate, CI=confidence interval, PY=patient years, HR=hazard ratio

Non-Melanoma Skin Cancer

NMSC is a dose related adverse reaction, with a greater risk in patients treated with 10 mg BID of tofacitinib than in patients treated with 5 mg BID.

In the five Phase 3 controlled trials, during the 0 to 3 months exposure, NMSC was reported in 1 (0.2%) patient (0.6 events/100 patient-years) who received placebo and 2 (0.2%) patients (0.7 events/100 patient-years) who received 5 mg BID of tofacitinib.

During 0 to 12 months exposure, NMSC was reported in 3 (0.3%) patients (0.3 events/100 patient-years) who received 5 mg BID of tofacitinib.

In the long-term extension trial, overall frequency of NMSC was 2.6% (0.71 events/100 patient-years) in all tofacitinib-treated patients; 2.5% of patients (0.6 events/100 patient-years) and 2.6% of patients (0.75 events/100 patient-years) in the 5 mg and 10 mg BID of tofacitinib, respectively.

In a post-authorization safety study ([Study RA-VI](#)), NMSC, including cutaneous squamous cell carcinoma, was more frequently observed in patients taking tofacitinib compared with patients taking TNFi (Table 8).

Table 8: NMSC in [Study RA-VI](#)

| | Tofacitinib 5 mg BID N = 1455 | Tofacitinib 10 mg BID* N = 1456 | All Tofacitinib N=2911 | TNFi N = 1451 |
|--|--|--|-----------------------------------|--------------------------|
| Non-melanoma Skin Cancer (NMSC) | | | | |

| | Tofacitinib 5 mg BID N = 1455 | Tofacitinib 10 mg BID* N = 1456 | All Tofacitinib N=2911 | TNFi N = 1451 |
|--|--|--|-----------------------------------|--------------------------|
| n (%) | 31 (2.13) | 33 (2.27) | 64 (2.20) | 16 (1.10) |
| IR (95% CI) per 100 PY | 0.61 (0.41, 0.86) | 0.69 (0.47, 0.96) | 0.64 (0.50, 0.82) | 0.32 (0.18, 0.52) |
| Tofacitinib vs TNFi HR (95% CI) | 1.90 (1.04, 3.47) | 2.16 (1.19, 3.92) | 2.02 (1.17, 3.50) | -- |
| Basal Cell Carcinoma | | | | |
| n (%) | 19 (1.31) | 16 (1.10) | 35 (1.20) | 13 (0.90) |
| IR (95% CI) per 100 PY | 0.37 (0.22, 0.58) | 0.33 (0.19, 0.54) | 0.35 (0.24, 0.49) | 0.26 (0.14, 0.44) |
| Tofacitinib vs TNFi HR (95% CI) | 1.43 (0.71, 2.90) | 1.28 (0.61, 2.66) | 1.36 (0.72, 2.56) | -- |
| Cutaneous Squamous Cell Carcinoma | | | | |
| n (%) | 15 (1.03) | 22 (1.51) | 37 (1.27) | 8 (0.55) |
| IR (95% CI) per 100 PY | 0.29 (0.16, 0.48) | 0.45 (0.29, 0.69) | 0.37 (0.26, 0.51) | 0.16 (0.07, 0.31) |
| Tofacitinib vs TNFi HR (95% CI) | 1.82 (0.77, 4.30) | 2.86 (1.27, 6.43) | 2.32 (1.08, 4.99) | -- |
| * The tofacitinib 10 mg BID treatment group consists of data from patients that were switched from tofacitinib 10 mg BID to tofacitinib 5 mg BID during the trial as a result of a study modification following a recommendation by the Data and Safety Monitoring Board in February 2019. Abbreviations: IR=incidence rate, CI=confidence interval, PY=patient years, HR=hazard ratio | | | | |

Mortality

In a post-authorization safety study ([Study RA-VI](#)), all-cause mortality was observed more frequently for patients taking tofacitinib (n=65/2911; 2.2%) compared with patients taking TNFi (n=17/1451; 1.2%). Related study data is presented in Table 9.

Table 9. Mortality in [Study RA-VI](#)

| Parameter | Tofacitinib 5mg BID N=1455 | Tofacitinib 10mg BID* N=1456 | All Tofacitinib N=2911 | TNFi N=1451 |
|---------------------------------------|---------------------------------------|---|-----------------------------------|------------------------|
| Deaths - Total | | | | |
| n (%) | 26 (1.79) | 39 (2.68) | 65 (2.23) | 17 (1.17) |
| IR (95% CI) per 100 PY | 0.50 (0.33, 0.74) | 0.80 (0.57, 1.09) | 0.65 (0.50, 0.82) | 0.34 (0.20, 0.54) |
| Tofacitinib vs TNFi HR (95% CI) | 1.49 (0.81, 2.74) | 2.37 (1.34, 4.18) | 1.91 (1.12, 3.27) | |
| Deaths - Infections | | | | |
| n (%) | 4 (0.27) | 9 (0.62) | 13 (0.45) | 3 (0.21) |
| IR (95% CI) per 100 PY | 0.08 (0.02, 0.20) | 0.18 (0.08, 0.35) | 0.13 (0.0, 0.22) | 0.06 (0.01, 0.17) |
| Tofacitinib vs TNFi HR (95% CI) | 1.30 (0.29, 5.79) | 3.10 (0.84, 11.45) | 2.17 (0.62, 7.62) | |
| Deaths - Cardiovascular Events | | | | |
| n (%) | 13 (0.89) | 20 (1.37) | 33 (1.13) | 10 (0.69) |
| IR (95% CI) per | 0.25 (0.13, 0.43) | 0.41 (0.25, 0.63) | 0.33 (0.23, 0.46) | 0.20 (0.10, 0.36) |

| Parameter | Tofacitinib 5mg BID N=1455 | Tofacitinib 10mg BID* N=1456 | All Tofacitinib N=2911 | TNFi N=1451 |
|--|-------------------------------|---------------------------------|---------------------------|-------------------|
| 100 PY | | | | |
| Tofacitinib vs TNFi HR (95% CI) | 1.26 (0.55, 2.88) | 2.05 (0.96, 4.39) | 1.65 (0.81, 3.34) | |
| Deaths - Malignancies | | | | |
| n (%) | 5 (0.34) | 0 | 5 (0.17) | 1 (0.07) |
| IR (95% CI) per 100 PY | 0.10 (0.03, 0.23) | 0.00 (0.00, 0.08) | 0.05 (0.02, 0.12) | 0.02 (0.00, 0.11) |
| Tofacitinib vs TNFi HR (95% CI) | 4.88 (0.57, 41.74) | 0 (0.00, Inf) | 2.53 (0.30, 21.64) | |
| Deaths - Other Causes | | | | |
| n (%) | 4 (0.27) | 10 (0.69) | 14 (0.48) | 3 (0.21) |
| IR (95% CI) per 100 PY | 0.08 (0.02, 0.20) | 0.21 (0.10, 0.38) | 0.14 (0.08, 0.23) | 0.06 (0.01, 0.17) |
| Tofacitinib vs TNFi HR (95% CI) | 1.30 (0.29, 5.81) | 3.45 (0.95, 12.54) | 2.34 (0.67, 8.16) | |
| * The tofacitinib 10 mg BID treatment group consists of data from patients that were switched from tofacitinib 10 mg BID to tofacitinib 5 mg BID during the trial as a result of a study modification following a recommendation by the Data and Safety Monitoring Board in February 2019. | | | | |

Thromboembolism

Venous thromboembolism, including pulmonary embolism, were observed more frequently in a post-authorization safety study ([Study RA-VI](#)), as shown in Table 10. Pulmonary embolism was observed more frequently with tofacitinib 10 mg BID than tofacitinib 5 mg BID. Deep vein thrombosis, and arterial thromboembolism were also observed in the study.

Table 10: Thromboembolism Adverse Reactions in [Study RA-VI](#)

| | Tofacitinib 5 mg BID N = 1455 | Tofacitinib 10 mg BID* N = 1456 | All Tofacitinib N=2911 | TNFi N = 1451 |
|------------------------------------|-------------------------------------|---------------------------------------|---------------------------|-------------------|
| Venous Thromboembolism** | | | | |
| n(%) | 17 (1.17) | 34 (2.34) | 51 (1.75) | 10 (0.69) |
| IR (95% CI) per 100 PY | 0.33 (0.19, 0.53) | 0.70 (0.49, 0.99) | 0.51 (0.38, 0.67) | 0.20 (0.10, 0.37) |
| Tofacitinib vs TNFi HR (95% CI) | 1.66 (0.76, 3.63) | 3.52 (1.74, 7.12) | 2.56 (1.30, 5.05) | -- |
| Pulmonary Embolism | | | | |
| n (%) | 9 (0.62) | 24 (1.65) | 33 (1.13) | 3 (0.21) |
| IR (95% CI) per 100 PY | 0.17 (0.08, 0.33) | 0.50 (0.32, 0.74) | 0.33 (0.23, 0.46) | 0.06 (0.01, 0.17) |
| Tofacitinib vs TNFi HR (95% CI) | 2.93 (0.79, 10.83) | 8.26 (2.49, 27.43) | 5.53 (1.70, 18.02) | -- |
| Deep Vein Thrombosis | | | | |
| n(%) | 11 (0.76) | 15 (1.03) | 26 (0.89) | 7 (0.48) |
| IR (95% CI) per 100 PY | 0.21 (0.11, 0.38) | 0.31 (0.17, 0.51) | 0.26 (0.17, 0.38) | 0.14 (0.06, 0.29) |
| Tofacitinib vs TNFi HR (95% CI) | 1.54 (0.60, 3.97) | 2.21 (0.90, 5.43) | 1.87 (0.81, 4.30) | -- |

| Arterial Thromboembolism | | | | |
|--|-------------------|-------------------|-------------------|-------------------|
| n(%) | 47 (3.23) | 45 (3.09) | 92 (3.16) | 41 (2.83) |
| IR (95% CI) per 100 PY | 0.92 (0.68, 1.22) | 0.94 (0.68, 1.25) | 0.93 (0.75, 1.14) | 0.82 (0.59, 1.12) |
| Tofacitinib vs TNFi HR (95% CI) | 1.12 (0.74, 1.70) | 1.14 (0.75, 1.74) | 1.13 (0.78, 1.63) | -- |
| * The tofacitinib 10 mg BID treatment group consists of data from patients that were switched from tofacitinib 10 mg BID to tofacitinib 5 mg BID during the trial as a result of a study modification following a recommendation by the Data and Safety Monitoring Board in February 2019. Abbreviations: IR=incidence rate, CI=confidence interval, PY=patient years, HR=hazard ratio | | | | |
| ** Venous thromboembolism (e.g., pulmonary embolism, deep vein thrombosis, retinal venous thrombosis). | | | | |

Major Adverse Cardiovascular Events (MACE), Including Myocardial Infarction

In a post-authorization study ([Study RA-VI](#)) the risk of MACE, including non-fatal myocardial infarction, was higher in patients treated with tofacitinib, compared to patients treated with TNFi (Table 11). In the tofacitinib 5 mg BID, tofacitinib 10 mg BID, All tofacitinib, and TNFi treatment arms, there were a total of 19, 19, 38, and 11 patients with MI events, respectively. Of these totals, the number of patients with fatal MI events was 0, 3, 3, and 3, respectively, whereas the number of patients with non-fatal MI events was 19, 16, 35, and 8, respectively.

Table 11: MACE (Including Myocardial Infarction) in [Study RA-VI](#)

| | Tofacitinib 5 mg BID N = 1455 | Tofacitinib 10 mg BID* N = 1456 | All Tofacitinib N=2911 | TNFi N = 1451 |
|--|-------------------------------|---------------------------------|--------------------------------|-------------------|
| Major Adverse Cardiovascular Events (MACE) ^α | | | | |
| n (%) | 47 (3.23) | 51 (3.50) | 98 (3.37) | 37 (2.55) |
| IR (95% CI) per 100 PY | 0.91 (0.67, 1.21) | 1.05 (0.78, 1.38) | 0.98 (0.79, 1.19) | 0.73 (0.52, 1.01) |
| Tofacitinib vs TNFi HR (95% CI) | 1.24 (0.81, 1.91) | 1.43 (0.94, 2.18) | 1.33 (0.91, 1.94) [‡] | -- |
| Non-fatal Myocardial Infarction | | | | |
| n (%) | 19 (1.31) | 16 (1.10) | 35 (1.20) | 8 (0.55) |
| IR (95% CI) per 100 PY | 0.37 (0.22, 0.57) | 0.33 (0.19, 0.53) | 0.35 (0.24, 0.48) | 0.16 (0.07, 0.31) |
| Tofacitinib vs TNFi HR (95% CI) | 2.32 (1.02, 5.30) | 2.08 (0.89, 4.86) | 2.20 (1.02, 4.75) | -- |
| * The tofacitinib 10 mg BID treatment group consists of data from patients that were switched from tofacitinib 10 mg BID to tofacitinib 5 mg BID during the trial as a result of a study modification following a recommendation by the Data and Safety Monitoring Board in February 2019. | | | | |
| [‡] The non-inferiority criterion was not met for the primary comparison of the combined tofacitinib doses to TNFi since the upper limit of the 95% CI exceeded the pre-specified non-inferiority criterion of 1.8. | | | | |
| ^α MACE includes nonfatal myocardial infarction, nonfatal stroke, and cardiovascular deaths excluding pulmonary embolism. Abbreviations: IR=incidence rate, CI=confidence interval, PY=patient years, HR=hazard ratio | | | | |

Gastrointestinal Perforations

In a post-authorization study ([Study RA-VI](#)), gastrointestinal perforations were observed in subjects treated with tofacitinib 5 mg BID, tofacitinib 10 mg BID, and TNF inhibitors (Table 12).

Table 12: Gastrointestinal Perforations in [Study RA-VI](#)

| | Tofacitinib 5 mg BID N = 1455 | Tofacitinib 10 mg BID* N = 1456 | All Tofacitinib N=2911 | TNFi N = 1451 |
|---|--|--|-----------------------------------|--------------------------|
| N (%) | 9 (0.62) | 5 (0.34) | 14 (0.48) | 4 (0.28) |
| IR (95% CI) per 100 PY | 0.17 (0.08, 0.33) | 0.10 (0.03, 0.24) | 0.14 (0.08, 0.23) | 0.08 (0.02, 0.20) |
| Tofacitinib vs TNFi HR (95% CI) | 2.20 (0.68, 7.15) | 1.29 (0.35, 4.80) | 1.76 (0.58, 5.34) | -- |
| * The tofacitinib 10 mg BID treatment group consists of data from patients that were switched from tofacitinib 10 mg BID to tofacitinib 5 mg BID during the trial as a result of a study modification following a recommendation by the Data and Safety Monitoring Board in February 2019. Abbreviations: BID=twice daily, TNFi=Tumor necrosis factor inhibitor, IR=incidence rate, CI=confidence interval, PY=patient years, HR=hazard ratio | | | | |

Fractures

In a post-authorization study ([Study RA-VI](#)) the incidence rate (IR) (95% CI) for fractures was 2.79 (95%CI: 2.34-3.30) for tofacitinib 5 mg BID, and 2.87 (95% CI: 2.40-3.40) for tofacitinib 10 mg BID, compared to 2.27 (95% CI: 1.87-2.74) for TNF inhibitors.

Psoriatic Arthritis

A total of 783 patients were treated with any dose of tofacitinib in PsA clinical studies resulting in 1238 patient-years of exposure. Of these, 635 patients were exposed to tofacitinib for at least one year.

The most common serious adverse reactions were serious infections. The most commonly reported adverse reactions (≥2%) in patients treated with tofacitinib 5 mg BID during the first 3 months in placebo-controlled clinical studies were bronchitis, diarrhea, dyspepsia, headache, nasopharyngitis, nausea.

The proportion of patients who discontinued treatment due to any adverse reactions during the first 3-months of the double-blind placebo-controlled studies was 3.2% for tofacitinib-treated patients and 2.5% for placebo-treated patients.

Overall, the safety profile observed in patients with active PsA treated with tofacitinib was consistent with the safety profile observed in patients with RA treated with tofacitinib. Incidence rates and types of adverse drug reactions, overall infections, serious infections, opportunistic infections in the two controlled Phase 3 PsA clinical studies were generally similar to those reported in RA Phase 3 clinical studies. The incidence rates of tuberculosis, malignancies (excluding NMSC), and NMSC in the two controlled Phase 3 PsA clinical studies were generally similar to those reported in RA Phase 3 clinical studies.

Ulcerative Colitis

Table 13 below lists adverse drug reactions reported by ≥1% of patients treated with tofacitinib – UC Phase 2 and Phase 3 Induction Studies

Table 13: Summary of Adverse Drug Reactions (adverse events for which there is evidence of causality) Reported by ≥1% of Patients Treated with Tofacitinib – UC Phase 2 and Phase 3 Induction Studies (up to 8 weeks)

| Body System[±]/Adverse Drug Reaction | Tofacitinib 10 mg BID N=938 (%) | Placebo N=282 (%) |
|--|--|--------------------------|
| Subjects with one or more ADR | 494 (52.7) | 130 (46.1) |
| Blood and lymphatic system disorders | 26 (2.8) | 10 (3.5) |

| | | |
|---|------------|----------|
| Anemia | 22 (2.3) | 9 (3.2) |
| Gastrointestinal disorders | 82 (8.7) | 26 (9.2) |
| Nausea | 28 (3.0) | 11 (3.9) |
| Abdominal pain | 25 (2.7) | 11 (3.9) |
| Vomiting | 9 (1.0) | 3 (1.1) |
| Dyspepsia | 12 (1.3) | 1(0.4) |
| General disorders and administration site conditions | 48 (5.1) | 13 (4.6) |
| Fatigue | 17 (1.8) | 5 (1.8) |
| Pyrexia | 24 (2.6) | 4 (1.4) |
| Infections and infestations | 111 (11.8) | 24 (8.5) |
| Nasopharyngitis | 56 (6.0) | 14 (5.0) |
| Influenza | 9 (1.0) | 3 (1.1) |
| Urinary tract infection | 11 (1.2) | 1 (0.4) |
| Pharyngitis | 10 (1.1) | 1 (0.4) |
| Investigations | 65 (6.9) | 4 (1.4) |
| Blood creatine phosphokinase increased | 25 (2.7) | 3 (1.1) |
| Elevated cholesterol levels* | 31 (3.3) | 0 |
| Musculoskeletal and connective tissue disorders | 33 (3.5) | 12 (4.3) |
| Arthralgia | 27 (2.9) | 12 (4.3) |
| Nervous system disorders | 77 (8.2) | 20 (7.1) |
| Headache | 73 (7.8) | 19 (6.7) |
| Respiratory | 14 (1.5) | 8 (2.8) |
| Cough | 13 (1.4) | 7 (2.5) |
| Skin and Subcutaneous Tissue Disorders | 18 (1.9) | 9 (3.2) |
| Rash | 12 (1.3) | 2 (0.7) |
| Vascular disorders | 9 (1.0) | 1 (0.4) |
| Hypertension | 9 (1.0) | 1 (0.4) |
| <p>* includes: hypercholesterolemia, hyperlipidemia, blood cholesterol increased, dyslipidemia, blood triglycerides increased, low density lipoprotein increased, low density lipoprotein abnormal, or lipids increased.</p> <p>± the total number of subjects with adverse reactions and the total number of subjects with adverse reactions for each body system include all adverse drug reactions (those reported by ≥1% of subjects treated with tofacitinib and those reported by <1% of subjects treated with tofacitinib); the total also includes some subjects who reported more than one adverse drug reaction (which inflates the percentage).</p> | | |

Table 14: Summary of Adverse Drug Reactions (adverse events for which there is evidence of causality) Reported by ≥1% of Patients Treated with Tofacitinib – UC Phase 3 Maintenance Study (up to 12 months)

| Body System [±] /Adverse Drug Reaction | Tofacitinib 5mg BID N=198 (%) | Tofacitinib 10mg BID N=196 (%) | Placebo N=198 (%) |
|--|----------------------------------|-----------------------------------|----------------------|
| Subjects with one or more ADR (%) | 166 (83.8) | 207 (100) | 153 (77.3) |
| Blood and lymphatic system disorders | 9 (4.5) | 5 (2.6) | 3 (1.5) |
| Anemia | 8(4.0) | 4 (2.0) | 3 (1.5) |
| Gastrointestinal disorders | 16 (8.1) | 32 (16.3) | 26 (13.1) |
| Diarrhea | 3 (1.5) | 9 (4.6) | 5 (2.5) |
| Nausea | 1 (0.5) | 8 (4.1) | 5 (2.5) |
| Abdominal pain | 5 (2.5) | 7 (3.6) | 11 (5.6) |
| Vomiting | 3 (1.5) | 6 (3.1) | 2 (1.0) |
| Dyspepsia | 4 (2.0) | 1 (0.5) | 2 (1.0) |
| General disorders and administration site | 12 (6.1) | 11 (5.6) | 17 (8.6) |

| Body System[±]/Adverse Drug Reaction | Tofacitinib 5mg BID N=198 (%) | Tofacitinib 10mg BID N=196 (%) | Placebo N=198 (%) |
|---|--|---|------------------------------|
| conditions | | | |
| Fatigue | 8 (4.0) | 4 (2.0) | 11 (5.6) |
| Pyrexia | 3 (1.5) | 6 (3.1) | 5 (2.5) |
| Infections and infestations | 51 (25.8) | 65 (33.2) | 37 (18.7) |
| Nasopharyngitis | 19 (9.6) | 27 (13.8) | 11 (5.6) |
| Herpes zoster | 3 (1.5) | 10 (5.1) | 1 (0.5) |
| Influenza | 4 (2.0) | 7 (3.6) | 7 (3.5) |
| Urinary tract infection | 5 (2.5) | 6 (3.1) | 4 (2.0) |
| Bronchitis | 5 (2.5) | 6 (3.1) | 3 (1.5) |
| Sinusitis | 6 (3.0) | 2 (1.0) | 2 (1.0) |
| Pharyngitis | 6 (3.0) | 1 (0.5) | 3 (1.5) |
| Gastroenteritis viral | 0 | 3 (1.5) | 2 (1.0) |
| Viral infection | 2 (1.0) | 1 (0.5) | 1 (0.5) |
| Injury, poisoning and procedural complications | 2 (1.0) | 2 (1.0) | 0 |
| Ligament sprain | 1 (0.5) | 2 (1.0) | 0 |
| Investigations | 19 (9.6) | 38 (19.4) | 7 (3.5) |
| Elevated cholesterol levels* | 9 (4.5) | 18 (9.2) | 3 (1.5) |
| Blood creatine phosphokinase increased | 6 (3.0) | 13 (6.6) | 4 (2.0) |
| Weight increased | 3 (1.5) | 4 (2.0) | 0 |
| Gamma glutamyltransferase increased, | 1 (0.5) | 3 (1.5) | 0 |
| Musculoskeletal and connective tissue disorders | 19 (9.6) | 19 (9.7) | 25 (12.6) |
| Arthralgia | 17 (8.6) | 17 (8.7) | 19 (9.6) |
| Musculoskeletal pain | 1 (0.5) | 2 (1.0) | 5 (2.5) |
| Neoplasms benign, malignant and unspecified (including cysts and polyps) | 0 | 2 (1.0) | 1 (0.5) |
| Non-melanoma skin cancers | 0 | 2 (1.0) | 1 (0.5) |
| Nervous system disorders | 18 (9.1) | 7 (3.6) | 12 (6.1) |
| Headache | 17 (8.6) | 6 (3.1) | 12 (6.1) |
| Psychiatric | 3 (1.5) | 1 (0.5) | 1 (0.5) |
| Insomnia | 3 (1.5) | 1 (0.5) | 1 (0.5) |
| Respiratory | 6 (3.0) | 8 (4.1) | 6 (3.0) |
| Cough | 6 (3.0) | 5 (2.6) | 5 (2.5) |
| Dyspnea | 0 | 2 (1.0) | 1 (0.5) |
| Skin and Subcutaneous Tissue Disorders | 7 (3.5) | 12 (6.1) | 17 (8.6) |
| Rash | 6 (3.0) | 11 (5.6) | 8 (4.0) |
| Vascular disorders | 4 (2.0) | 4 (2.0) | 1 (0.5) |
| Hypertension | 4 (2.0) | 4 (2.0) | 1 (0.5) |
| <p>* includes: hypercholesterolemia, hyperlipidemia, blood cholesterol increased, dyslipidemia, blood triglycerides increased, low density lipoprotein increased, low density lipoprotein abnormal, or lipids increased.</p> <p>± The total number of subjects with adverse reactions and the total number of subjects with adverse reactions for each body system include all adverse drug reactions (those reported by ≥1% of subjects treated with tofacitinib and those reported by <1% of subjects treated with tofacitinib); the total also includes some subjects who reported more than one adverse drug reaction (which inflates the percentage).</p> | | | |

Overall Infections

In the randomised 8-week Phase 2/3 induction studies, the proportions of patients with infections were 21.1%

(198 patients) in the tofacitinib 10 mg BID group compared to 15.2% (43 patients) in the placebo group. In the randomised 52-week Phase 3 maintenance study, the proportion of patients with infections were 35.9% (71 patients) in the 5 mg BID and 39.8% (78 patients) in the 10 mg BID tofacitinib groups, compared to 24.2% (48 patients) in the placebo group.

In the maintenance study, results suggested that the risk of opportunistic infection was possibly dose related: tofacitinib 10 mg BID (2.0%), tofacitinib 5 mg BID (1.0%), and placebo (0.5%). All opportunistic infections were herpes zoster infections. Herpes zoster was reported more frequently with tofacitinib 10 mg BID (5.1%), as compared to tofacitinib 5 mg BID (1.5%), or placebo (0.5%), indicating that the risk of herpes zoster is dose related.

In the entire treatment experience with tofacitinib, the most commonly reported infection was nasopharyngitis, occurring in 18.2% of patients (211 patients).

Serious Infections

The incidence rates and types of serious infections in the UC clinical trials were generally similar to those reported in RA Phase 3 clinical trials with tofacitinib.

Patients treated with tofacitinib 10 mg BID had a higher rate of serious infections compared to those treated with 5 mg BID.

Opportunistic infections (excluding tuberculosis)

In the maintenance study, herpes zoster was reported more frequently with tofacitinib 10 mg BID (5.1%), as compared to tofacitinib 5 mg BID (1.5%), or placebo (0.5%), indicating that the risk of herpes zoster is dose related.

Also, opportunistic herpes zoster infections (including serious cases, such as, disseminated, meningoencephalitis, ophthalmologic) were reported in patients treated with tofacitinib 10 mg BID.

Malignancies (excluding NMSC)

In the controlled clinical studies (up to 52-week treatment), no malignancies (excluding NMSC) were reported with tofacitinib.

In the long-term extension open-label study, malignancies (excluding NMSC) have been observed in patients treated with tofacitinib 10 mg BID, including solid cancers and lymphoma.

8.3 Less Common Clinical Trial Adverse Reactions

Rheumatoid Arthritis, Psoriatic Arthritis

Blood and Lymphatic System Disorders: leukopenia, lymphopenia, neutropenia

Cardiovascular: congestive heart failure, myocardial infarction

Gastrointestinal Disorders: abdominal pain, appendicitis, gastrointestinal perforation

General Disorders and Administration Site Conditions: influenza

Hepatobiliary Disorders: hepatic steatosis

Infections and Infestations: atypical mycobacterial infection, arthritis bacterial, bacteraemia, cellulitis, cytomegalovirus infection, disseminated tuberculosis, diverticulitis, encephalitis, gastroenteritis viral, herpes simplex, herpes zoster, meningitis cryptococcal, mycobacterium avium complex infection, necrotising fasciitis, pneumonia bacterial, pneumonia pneumococcal, pneumocystis jiroveci pneumonia, pyelonephritis, sepsis, staphylococcal bacteraemia, tuberculosis, tuberculosis of central nervous system, urosepsis, viral infection.

Injury, Poisoning and Procedural Complications: muscle strain, fall

Investigations: blood cholesterol increased, blood creatinine increased, blood creatine phosphokinase increased, gamma glutamyltransferase increased, hepatic enzyme increased, liver function test abnormal, low density lipoprotein increased, transaminases increased, weight increased,

Metabolism and Nutrition Disorders: dehydration, dyslipidemia, hyperlipidemia

Musculoskeletal and Connective Tissue Disorders: joint swelling, ligament sprain, musculoskeletal pain, tendonitis,

Neoplasm Benign, Malignant and Unspecified (Including Cysts and Polyps): lymphoma, non-melanoma skin cancers, solid tumours

Nervous System Disorders: paraesthesia

Psychiatric Disorders: insomnia

Respiratory, Thoracic and Mediastinal Disorders: cough, dyspnoea, sinus congestion,

Skin and Subcutaneous Tissue Disorders: erythema, pruritus

Vascular disorders: arterial thrombosis, deep vein thrombosis, pulmonary embolism.

Ulcerative Colitis

Blood and Lymphatic System Disorders: neutropenia, lymphopenia, leukopenia

Gastrointestinal Disorders: gastritis

General Disorders and Administration Site Conditions: oedema peripheral

Hepatobiliary Disorders: hepatic steatosis

Infections and Infestations: pneumonia, pyelonephritis, cellulitis, herpes simplex, tuberculosis, arthritis bacterial, cytomegalovirus infection, diverticulitis

Injury, Poisoning and Procedural Complications: muscle strain

Investigations: hepatic enzyme increased, transaminases increased, blood creatinine increased, liver function test abnormal, low density lipoprotein increased

Metabolism and Nutrition Disorders: dehydration

Musculoskeletal and Connective Tissue Disorders: tendonitis, joint swelling

Neoplasm Benign, Malignant and Unspecified (Including Cysts and Polyps): non-melanoma skin cancers, solid cancers, lymphomas

Nervous System Disorders: paraesthesia

Respiratory, Thoracic and Mediastinal Disorders: sinus congestion

Skin and Subcutaneous Tissue Disorders: erythema, pruritus

8.4 Abnormal Laboratory Findings: Hematologic, Clinical Chemistry and Other Quantitative Data

Clinical Trial Findings

Laboratory Tests– Rheumatoid Arthritis and Ulcerative Colitis

Creatine Kinase

Treatment with tofacitinib was associated with increases in creatine kinase (CK). Maximum effects were generally observed within 6 months. Rhabdomyolysis was reported in one patient treated with tofacitinib.

CK levels should be checked in patients with symptoms of muscle weakness and/or muscle pain to evaluate for evidence of rhabdomyolysis (see [7 WARNINGS AND PRECAUTIONS](#)).

ECG Findings

In placebo-controlled Phase 2 clinical trials, steady-state treatment with 5-10 mg BID tofacitinib was associated with statistically significant 4-7 bpm decreases in heart rate and 4-10 ms increases in the PR interval compared with placebo (see [7 WARNINGS AND PRECAUTIONS](#) and [9 DRUG INTERACTIONS](#)).

Lipids

Treatment with tofacitinib was associated with dose related increases in lipid parameters.

Elevations in lipid parameters (total cholesterol, LDL cholesterol, HDL cholesterol, triglycerides) generally reached maximal effects at 6 weeks following initiation of tofacitinib in the controlled RA double-blind clinical trials. Changes in lipid parameters from baseline through the end of the study (6-12 months) in the controlled clinical studies in RA are summarized below:

- Mean LDL cholesterol increased by 14% in the tofacitinib 5 mg BID arm.
- Mean HDL cholesterol increased by 16% in the tofacitinib 5 mg BID arm.
- Mean LDL/HDL ratios were essentially unchanged in tofacitinib-treated patients.

In the five controlled RA clinical trials, 4.4% of patients treated with 5 mg BID, initiated lipid-lowering medication while on study.

In the RA long-term safety population, elevations in the lipid parameters remained consistent with what was seen in the controlled clinical studies.

Increases of total cholesterol, LDL cholesterol, and HDL cholesterol were also reported in a post-authorization safety ([Study RA-VI](#); Table 15).

Table 15: Mean percent increase of cholesterol ([Study RA-VI](#))

| | | Tofacitinib 5 mg BID | Tofacitinib 10 mg BID* | TNFi |
|-------------------------------|-----------|-------------------------|---------------------------|------|
| LDL, mean percent increase | 12 months | 13.80 | 17.04 | 5.50 |
| | 24 months | 12.71 | 18.14 | 3.64 |
| HDL, mean percent increase | 12 months | 11.71 | 13.63 | 2.82 |
| | 24 months | 11.58 | 13.54 | 1.42 |

* The tofacitinib 10 mg BID treatment group consists of data from patients that were switched from tofacitinib 10 mg BID to tofacitinib 5 mg BID during the trial as a result of a study modification following a recommendation by the Data and Safety Monitoring Board in February 2019.

Liver Enzyme Tests

Confirmed increases in liver enzymes >3x upper limit of normal (ULN) were uncommonly observed. In those patients experiencing liver enzyme elevation, modification of treatment regimen, such as reduction in the dose of concomitant DMARD, interruption of tofacitinib, or reduction in tofacitinib dose, resulted in decrease or normalization of liver enzymes.

In the controlled portion of the RA Phase 3 monotherapy study (0-3 months), ALT elevations >3x ULN were observed in 1.65% and 0.41% of patients receiving placebo and 5 mg BID, respectively. In this study, AST elevations >3x ULN were observed in 1.65%, and 0.41% of patients receiving placebo and 5 mg BID, respectively.

In the controlled portion of the RA Phase 3 studies on background DMARDs (0-3 months), ALT elevations >3x ULN were observed in 0.9% and 1.24% of patients receiving placebo and 5 mg BID, respectively. In these studies, AST elevations >3x ULN were observed in 0.72% and 0.52% of patients receiving placebo and 5 mg BID, respectively.

In the RA long-term extension trial, ALT and AST elevations greater than 3x ULN were observed in 2.2% and 1.1% of all tofacitinib-treated patients, respectively. Overall, total bilirubin elevations greater than 2x ULN were observed in 3 (0.1%) patients. Increases to ≥5x and ≥10x ULN were observed for both ALT (0.5% and 0.2% of patients, respectively) and AST (0.3% and 0.1% of patients, respectively) in all patients treated with tofacitinib.

In RA patients taking 5 mg BID of tofacitinib, the ALT and AST elevations greater than 3x ULN were observed in 2.4% and 1.3% of patients, respectively. There was no subject who had the total bilirubin elevations greater than 2x ULN. Increases to ≥5 and ≥10x ULN were observed for both ALT (0.4% and 0.1% of patients, respectively) and AST (0.2% and 0% of patients, respectively).

In RA patients taking 10 mg BID of tofacitinib, the ALT and AST elevations greater than 3x ULN were observed in 2.1% and 1.1% of patients, respectively. The total bilirubin elevations greater than 2x ULN were observed in 3 (0.1%) patients. Increases to ≥5 and ≥10x ULN were observed for both ALT (0.5% and 0.2% of patients, respectively) and AST (0.3% and 0.1% of patients, respectively).

Two patients treated with 10 mg BID of tofacitinib in the RA long-term extension trial were assessed as probable DILI by the adjudication committee. One of the two patients had other possible causes of alcohol intake and methotrexate.

Elevations of ALT and AST were reported more frequently in patients taking tofacitinib compared with patients taking TNFi in a post-authorization safety study ([Study RA-VI](#); Table 16).

Table 16: Percentage of patients with at least one post-baseline elevation of liver enzymes ([Study RA-VI](#))

| | | Tofacitinib 5 mg BID | Tofacitinib 10 mg BID* | All Tofacitinib | TNFi |
|---------------------------------------|-----------|----------------------|------------------------|-----------------|-------|
| ALT elevation, percentage of patients | > 1 x ULN | 52.83 | 54.46 | 53.64 | 43.33 |
| | > 3 x ULN | 6.01 | 6.51 | 6.27 | 3.77 |
| | > 5 x ULN | 1.68 | 1.97 | 1.82 | 1.12 |
| AST elevation, percentage of patients | > 1 x ULN | 45.84 | 51.58 | 48.70 | 37.18 |
| | > 3 x ULN | 3.21 | 4.57 | 3.89 | 2.38 |
| | > 5 x ULN | 0.98 | 1.62 | 1.30 | 0.70 |

* The tofacitinib 10 mg BID treatment group consists of data from patients that were switched from tofacitinib 10 mg BID to tofacitinib 5 mg BID during the trial as a result of a study modification following a recommendation by the Data and Safety Monitoring Board in February 2019.

In the clinical studies in UC, changes in liver enzyme tests observed with tofacitinib 5 mg BID treatment were similar to the changes observed in clinical studies in RA.

In UC patients, tofacitinib treatment with 5 and 10 mg BID was also associated with an increased incidence of liver enzyme elevation compared to placebo, with a trend for higher incidence with the 10 mg BID as compared to the 5 mg BID dose.

One patient with tofacitinib 10 mg BID in the maintenance UC study experienced an increase in liver

enzymes which decreased upon discontinuation of treatment. The case was adjudicated as possible DILI, while noting ultrasound findings of fatty liver.

Lymphocytes

In the five controlled RA clinical trials, confirmed decreases in absolute lymphocyte counts below 0.5×10^9 cells/L occurred in 0.2% of patients for the 5 mg BID tofacitinib group during 12 months of exposure.

Confirmed lymphocyte counts less than 0.5×10^9 cells/L were associated with an increased incidence of treated and serious infections (see [7 WARNINGS AND PRECAUTIONS](#)).

In the RA long-term extension trial, cases of lymphopenia have been reported in 181 (4.0%) patients (1.11 events/100 patient-years) treated with tofacitinib; 4.5% of patients (1.07 events/100 patient-years) and 3.9% of patients (1.12 events/100 patient-years) in the 5 mg and 10 mg BID of tofacitinib, respectively. Confirmed decreased in absolute lymphocyte counts below 0.5×10^9 cells/L occurred in 1.3% of all tofacitinib-treated patients; 1.1% of patients for the 5 mg BID tofacitinib group, and 1.4% of patients for the 10 mg BID tofacitinib group.

In a post-authorization safety study ([Study RA-VI](#)) the median decrease in lymphocyte counts were greater in patients taking tofacitinib (-0.21) compared with patients taking TNFi (0.37).

In the 52-week maintenance study in UC, a single absolute lymphocyte count below 0.5×10^9 cells/L was reported in 2.6% (n=5) of patients treated with 10 mg BID, and was not reported in patients treated with 5 mg BID or placebo. No patients in any treatment group had confirmation of a lymphocyte count below 0.5×10^9 cells/L based on two sequential tests.

Neutrophils

In the controlled RA clinical studies, confirmed decreases in ANC below 1×10^9 cells/L occurred in 0.08% of patients in the 5 mg BID tofacitinib group during 12 months of exposure. There were no confirmed decreases in ANC below 0.5×10^9 cells/L observed in any treatment group.

There was no clear relationship between neutropenia and the occurrence of serious infections.

In the long-term extension trial, cases of neutropenia have been reported in 86 (1.9%) patients (0.52 events/100 patient-years) treated with tofacitinib; 4.0% of patients (0.97 events/100 patient-years) and 1.2% of patients (0.35 events/100 patient-years) in the 5 mg and 10 mg BID of tofacitinib, respectively. Confirmed decreased in ANC below 1×10^9 cells/L occurred in 0.2% in all tofacitinib-treated patients; 0.4% of patients for the 5 mg BID tofacitinib group, and 0.1% of patients for the 10 mg BID tofacitinib group.

In the clinical studies in UC, changes in neutrophils observed with tofacitinib treatment were similar to the changes observed in clinical studies in RA.

Serum Creatinine

In the controlled RA clinical trials, dose-related elevations in serum creatinine were observed with tofacitinib treatment. The mean increase in serum creatinine was <0.1 mg/dL in the 12-month pooled safety analysis; however, with increasing duration of exposure in the long-term extension trial, up to 6.9% of patients were discontinued from tofacitinib treatment due to the protocol-specified discontinuation criterion of an increase in creatinine by more than 50% of baseline. The clinical significance of the observed serum creatinine elevations is unknown.

In the UC studies, an increase of more than 50% in serum creatinine was reported in 1.6% of patients predominantly treated with tofacitinib 5 mg BID, and 3.4% of those predominantly treated with tofacitinib 10 mg BID.

Laboratory Tests— Psoriatic Arthritis

In the controlled clinical trials in PsA, changes in hematologic and clinical chemistry findings observed with tofacitinib treatment were similar to the changes observed in Phase 3 clinical trials in RA.

8.5 Post-Market Adverse Reactions

Because these reactions are reported voluntarily from a population of uncertain size, it is generally not possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

Immune system disorders: drug hypersensitivity reactions including angioedema and urticaria (see [2 CONTRAINDICATIONS](#) and [7 WARNINGS AND PRECAUTIONS](#))

Serious infections: viral reactivation (hepatitis B reactivation) (see [7 WARNINGS AND PRECAUTIONS](#))

9 DRUG INTERACTIONS

9.2 Drug Interactions Overview

In vitro studies indicate that tofacitinib does not significantly inhibit the activity of the major human drug metabolizing CYPs (CYP1A2, CYP2B6, CYP2C8, CYP2C9, CYP2C19, CYP2D6, and CYP3A4) at concentrations exceeding 80 times the steady state C_{max} of a 5 and 10 mg BID dose in patients treated with tofacitinib. In vitro studies also indicated a low risk of induction of CYP3A4 (2-fold mRNA at 6.25 μ M), CYP2B6 (2-fold mRNA at 12.5 μ M), and CYP1A2 (no enzyme changes) at clinically relevant concentrations (total C_{max} of 0.186 μ M).

In vitro, tofacitinib is a substrate for multidrug resistance (MDR) 1, but not for breast cancer resistance protein (BCRP), organic anion transporting polypeptide (OATP) 1B1/1B3, or organic cationic transporter (OCT) 1/2. In vitro data indicate that the potential for tofacitinib to inhibit transporters such as P-glycoprotein, MDR1, organic anion transporter (OAT) P1B1/1B3, OCT2, OAT1/3, cationic transporters or multidrug resistance-associated protein (MRP) at therapeutic concentrations is also low.

Tofacitinib exposure is increased when tofacitinib is coadministered with potent CYP3A4 inhibitors (e.g., ketoconazole) or when administration of one or more concomitant medications results in both moderate inhibition of CYP3A4 and potent inhibition of CYP2C19 (e.g., fluconazole). Tofacitinib exposure is decreased when tofacitinib is coadministered with potent CYP3A4 inducers (e.g., rifampin). Inhibitors of CYP2C19 or P-glycoprotein are unlikely to alter the PK of tofacitinib.

The in vitro results were confirmed by a human drug interaction study showing no changes in the PK of midazolam, a highly sensitive CYP3A4 substrate, when coadministered with tofacitinib.

In vitro studies indicate that tofacitinib does not significantly inhibit the activity of the major human drug-metabolizing uridine 5'-diphospho-glucuronosyltransferases (UGTs) [UGT1A1, UGT1A4, UGT1A6, UGT1A9, and UGT2B7] at concentrations exceeding 250 times the steady state C_{max} of a 5 and 10 mg BID dose in RA, PsA and UC patients.

The oral clearance of tofacitinib does not vary with time, indicating that tofacitinib does not normalize CYP enzyme activity in patients. Therefore, coadministration with tofacitinib is not expected to result in clinically relevant increases in the metabolism of CYP substrates.

9.4 Drug-Drug Interactions


Table 17: Summary of Drug-Drug Interactions

| Drug | Reference | Effect | Clinical Comment |
|-----------------------------|-----------|---|---|
| Methotrexate | CT | Coadministration with methotrexate (15-25 mg MTX once weekly) had no effect on the PK of tofacitinib and decreased methotrexate AUC (area under the curve) and C_{max} by 10% and 13% respectively. | No dose adjustment is required for either drug. |
| Ketoconazole | CT | Coadministration of ketoconazole, a strong CYP3A4 inhibitor, with a single dose of tofacitinib increased the AUC and C_{max} of tofacitinib by 103% and 16%, respectively | The recommended dose is half the daily dose indicated for patients not receiving strong CYP3A4 inhibitors concomitantly, i.e., in patients already taking: Teva-Tofacitinib 10 mg BID, reduce the dose to Teva-Tofacitinib 5 mg BID, or Teva-Tofacitinib 5 mg BID, reduce the dose to Teva-Tofacitinib 5 mg once daily. |
| Fluconazole | CT | Coadministration of fluconazole, a moderate inhibitor of CYP3A4 and a strong inhibitor of CYP2C19, increased the AUC and C_{max} of tofacitinib by 79% and 27%, respectively | The recommended dose is half the daily dose indicated for patients not receiving concomitant medications that result in moderate inhibition of CYP3A4 and potent inhibition of CYP2C19, i.e., in patients already taking: Teva-Tofacitinib 10 mg BID, reduce the dose to Teva-Tofacitinib 5 mg BID, or Teva-Tofacitinib 5 mg BID, reduce the dose to Teva-Tofacitinib 5 mg once daily. |
| Tacrolimus and Cyclosporine | CT | Coadministration of tacrolimus, a mild inhibitor of CYP3A4, increased the AUC of tofacitinib by 21% and decreased the C_{max} of tofacitinib by 9%. Coadministration of cyclosporine, a moderate inhibitor of CYP3A4, increased the AUC of tofacitinib by 73% and decreased C_{max} of tofacitinib by 17%. | There is a risk of added immunosuppression when tofacitinib is co-administered with potent immunosuppressive drugs (e.g: tacrolimus, cyclosporine, azathioprine). The combined use with these potent immunosuppressives has not been studied in patients and is not recommended. |
| Rifampin | CT | Coadministration of rifampin, a strong CYP3A4 inducer, decreased the AUC and C_{max} of tofacitinib by 84% and 74%, | Coadministration of tofacitinib with potent inducers of CYP3A4 may result in loss of or reduced clinical response /efficacy. |

| Drug | Reference | Effect | Clinical Comment |
|--|------------------|--|---|
| | | respectively | |
| Midazolam | CT | Coadministration of with midazolam, a highly sensitive CYP3A4 substrate, had no effect on midazolam PK | No dosage adjustment is required for CYP3A4 substrates such as midazolam. |
| Oral contraceptives (Ethinyl Estradiol and Levonorgestrel) | CT | Coadministration of tofacitinib with oral contraceptives had no effect on the PK of either oral contraceptive in healthy females | No dose adjustment is required for either oral contraceptives (ethinyl estradiol and levonorgestrel). |
| Metformin | CT | Coadministration of tofacitinib with metformin, a substrate of Organic Cationic Transporter and Multidrug and Toxic Compound Extrusion, had no effect on the PK of metformin | No dosage adjustment is required for metformin. |
| Legend: C = Case Study; CT = Clinical Trial; T = Theoretical | | | |

The impact of extrinsic factors on tofacitinib pharmacokinetics is summarized in Figure 1 and 2 with dosage adjustment recommendations.


Figure 1: Impact of Co-administered of drugs on Pharmacokinetics Tofacitinib



Note: Reference group is administration of tofacitinib alone; PK=Pharmacokinetics; CI=Confidence Interval

^a In RA patients the recommended dose is tofacitinib 5 mg once daily. In UC patients receiving 10 mg BID, tofacitinib dosage should be reduced to 5 mg BID, and in UC patients receiving 5 mg BID, tofacitinib dosage should be reduced to 5 mg once daily.

Figure 2: Impact of Tofacitinib on Pharmacokinetics of Co-administered Drugs



Note: Reference group is administration of concomitant medication alone; OCT= Organic Cationic Transporter; MATE= Multidrug and Toxic Compound Extrusion; PK=Pharmacokinetics; CI=Confidence Interval

Drugs that Decrease Heart Rate and/or Prolong the PR Interval

Tofacitinib resulted in a decrease in heart rate and an increase in the PR interval (see [7 WARNINGS AND PRECAUTIONS](#) and [8 ADVERSE REACTIONS](#)). Caution should be observed if Teva-Tofacitinib is used concomitantly with other drugs that lower heart rate and/or prolong the PR interval, such as antiarrhythmics, beta blockers, alpha2 adrenoceptor agonists, non-dihydropyridine calcium channel blockers, digitalis glycosides, cholinesterase inhibitors, sphingosine-1 phosphate receptor modulators, and some HIV protease inhibitors.

Combination with other therapies

Tofacitinib has not been studied and is not indicated to be used in combination with biologics such as TNF antagonists, interleukin (IL)-1R antagonists, IL-6R antagonists, IL-17 antagonists, IL-12/IL-23 antagonists, anti-CD20 monoclonal antibodies, anti-integrins, selective co-stimulation modulators, and potent immunosuppressants such as azathioprine, 6-mercaptopurine, cyclosporine, and tacrolimus because of the possibility of increased immunosuppression and increased risk of infection.

The use of tofacitinib in combination with phosphodiesterase 4 inhibitors has not been studied in tofacitinib clinical trials.

9.5 Drug-Food Interactions

Grapefruit juice affects CYP450 3A-mediated metabolism and concomitant administration with Teva-Tofacitinib should be avoided.

9.6 Drug-Herb Interactions

St John’s Wort is a CYP3A4 inducer and co-administration with Teva-Tofacitinib may result in loss of or reduced clinical response.

9.7 Drug-Laboratory Test Interactions

Interactions with laboratory tests have not been established.

10 CLINICAL PHARMACOLOGY

10.1 Mechanism of Action

Tofacitinib is a potent, selective inhibitor of the JAK family of kinases with a high degree of selectivity against other kinases in the human genome. In kinase assays, tofacitinib, inhibits JAK1, JAK2, JAK3, and to a lesser extent TyK2. In cellular settings where JAK kinases signal in pairs, tofacitinib preferentially inhibits signaling by heterodimeric receptors associated with JAK3 and/or JAK1 with functional selectivity over receptors that signal via pairs of JAK2. Inhibition of JAK1 and JAK3 by tofacitinib blocks signaling through the common gamma chain-containing receptors for several cytokines, including IL-2, -4, -7, -9, -15, and -21. These cytokines are integral to lymphocyte activation, proliferation, and function and inhibition of their signaling may thus result in modulation of multiple aspects of the immune response. In addition, inhibition of JAK1 will result in attenuation of signaling by additional pro-inflammatory cytokines, such as IL-6 and Type I interferons. At higher exposures, inhibition of erythropoietin signaling could occur via inhibition of JAK2 signaling.

10.2 Pharmacodynamics

In patients with RA, treatment with tofacitinib was associated with dose-dependent reductions of circulating CD16/56+ natural killer cells, with estimated maximum reductions occurring at approximately 8-10 weeks after initiation of therapy. These changes generally resolved within 2-6 weeks after discontinuation of treatment. Treatment with tofacitinib was associated with dose-dependent increases in B cell counts. Changes in circulating T-lymphocyte counts and T-lymphocyte subsets were small and inconsistent. The clinical significance of these changes is unknown.

Changes in total serum IgG, M, and A levels over 6-month dosing of patients with RA were small, not dose-dependent and similar to those seen on placebo.

After treatment with tofacitinib in patients with RA, rapid decreases in serum C-reactive protein (CRP) were observed and maintained throughout dosing. Changes in CRP observed with tofacitinib treatment do not reverse fully within 2 weeks after discontinuation, indicating a longer duration of pharmacodynamic activity compared to the half-life.

In a randomized, double-blind, placebo-controlled study in 200 adult RA patients treated with tofacitinib 10 mg BID or placebo, humoral responses to concomitant pneumococcal and influenza vaccines were assessed. The percentages of patients achieving a satisfactory humoral response to pneumococcal vaccines were lower for the tofacitinib group than the placebo group. This effect was more pronounced for patients receiving background methotrexate. A total of 31.6% tofacitinib-treated subjects and 61.8% placebo-treated subjects who received background methotrexate achieved a ≥ 2 -fold increase in antibody concentrations to ≥ 6 of 12 pneumococcal antigens.

In the same study, the proportion of patients achieving protective antibody levels to the influenza antigens was lower in the tofacitinib group (64.9%) compared to the placebo group (92.7%) in patients receiving background methotrexate. However, the difference in humoral response to the influenza vaccine was small with 50.9% of patients in the tofacitinib group and 58.2% in the placebo group with background methotrexate achieving a ≥ 4 -fold increase in antibody titers to ≥ 2 of 3 influenza antigens.

Similar changes in T cells, B cells and serum CRP have been observed in patients with active PsA, although reversibility was not assessed. Total serum immunoglobulins were not assessed in patients with active

PSA.

Patients with UC were not studied.

10.3 Pharmacokinetics (PK)

Tofacitinib

Following oral administration of tofacitinib, the PK profile of tofacitinib is characterized by rapid absorption (peak plasma concentrations are reached within 0.5-1 hour), rapid elimination (half-life of ~3 hours) and dose-proportional increases in systemic exposure in the therapeutic dose range. Steady state concentrations are achieved in 24-48 hours with negligible accumulation after BID administration.

A geometric mean accumulation ratio (Rac) of 1.12 following BID dosing indicates little difference between single dose and steady state concentrations as well as the predictability of steady state PK from single dose data. The dose-AUC relationship was adequately described by a linear model fit to log-both sides transformed data while the dose-C_{max} relationship were best described by a nonlinear sigmoidal, hyperbolic model fit to log-transformed C_{max} data. Although the nonlinear model provided better description of the dose-C_{max} relationship relative to a linear model, when compared to 5 mg, the mean model predicted relative changes in dose-normalized C_{max} were approximately +7% for 10 mg, +2% for 30 mg, and -10% for 50 mg doses. These small changes from linearity support the conclusion that tofacitinib C_{max} is approximately dose proportional at least up to 5 times the 10 mg dose.

Pharmacokinetics in Patients with Moderately to Severely Active UC

Population PK analysis in UC patients indicated that PK characteristics were similar to that of RA patients. There were no clinically relevant differences in tofacitinib exposure (AUC), based on age, weight, gender and race, after accounting for differences in renal function (i.e., creatinine clearance) between patients. The between-subject variability (% coefficient of variation) in AUC of tofacitinib is estimated to be approximately 23% to 25% in UC patients.

Absorption:

Tofacitinib is well-absorbed, with an absolute oral bioavailability of 74% following administration of tofacitinib. Coadministration of tofacitinib with a high-fat meal resulted in no changes in AUC while C_{max} was reduced by 32%. In clinical trials, tofacitinib was administered without regard to meal.

Distribution:

After intravenous administration, the volume of distribution is 87 L. The protein binding of tofacitinib is ~40%). Tofacitinib binds predominantly to albumin and does not appear to bind to α 1-acid glycoprotein. Tofacitinib distributes equally between red blood cells and plasma.

Metabolism:

Clearance mechanisms for tofacitinib are approximately 70% hepatic metabolism and 30% renal excretion of the parent drug. The metabolism of tofacitinib is primarily mediated by CYP3A4 with minor contribution from CYP2C19. In a human radiolabeled study, more than 65% of the total circulating radioactivity was accounted for by unchanged tofacitinib, with the remaining 35% attributed to 8 metabolites, each accounting for less than 8% of total radioactivity. The pharmacologic activity of tofacitinib is attributed to the parent molecule.

Elimination:

Approximately 94% of a radioactive dose of tofacitinib was recovered from the urine (80%) and feces

(14%), with the majority of excreted radioactivity recovered within 24 hours after dosing.

Table 18: Summary of Tofacitinib Pharmacokinetic Parameters after Repeated Oral Administration of tofacitinib 10 mg BID or Single IV Administration in Humans

| Healthy Volunteers | Oral Administration | | | IV Administration | |
|--------------------|--------------------------|--------------------|----------------------------------|---------------------|----------------------------|
| | C _{max} (ng/mL) | t _½ (h) | AUC _{0-12hrs} (ng·h/mL) | Clearance (L/h) | Volume of distribution (L) |
| RA Patients | 79.4 | 3.0 | 311 | 25 | 87 |
| PsA Patients | 116 | 3.62 | 507 | N/A (no IV data) | N/A (no IV data) |
| UC Patients | 88.9 | 3.74 | 436 | N/A (no IV data) | N/A (no IV data) |
| | 91 | 3.05 | 404 | N/A (no IV data) | N/A (no IV data) |

N/A = Not available; C_{max} = maximum plasma concentration; t_½ = terminal elimination half-life; AUC₀₋₁₂ = area under the plasma concentration-time curve from time 0 to 12 hours post dose; CL = total systemic clearance; V_{SS} = volume of distribution at steady state.

Special Populations and Conditions

Rheumatoid Arthritis and Ulcerative Colitis

Pediatrics (<18 years of age): The pharmacokinetics, safety and effectiveness of tofacitinib in pediatric patients have not been established; therefore, tofacitinib should not be used in this patient population. Pharmacokinetic of tofacitinib was characterized in an open-label, non-randomized, multi-center, Phase 1 study conducted in pediatric patients (aged from 2 to less than 18 years) with juvenile idiopathic arthritis. A total of 26 patients were enrolled in this study and treated at dosing regimens based on the children's age and body weight. The study consisted of 3 cohorts based on subject age with at least 8 subjects per cohort. Based on limited data, the PK profile of tofacitinib appears to be characterized by a rapid absorption (peak plasma concentrations were reached within 0.5-1 hour) and a rapid elimination. The average half-lives for tofacitinib were approximately 2.6h, 1.9h, and 1.8h for the Cohorts 1 (12 to <18 years), 2 (6 to <12 years) and 3 (2 to <6 years), respectively, with individual values ranging from 1.4 to 3.1h across all cohorts.

Geriatrics (>65 years of age): Population PK analysis in RA patients indicated that geriatric patients 80 years of age were estimated to have <5% higher tofacitinib AUC relative to the mean age of 55 years. Of the 3315 patients who enrolled in studies I to V, a total of 505 (15%) RA patients were 65 years of age and older, including 71 (2%) patients 75 years and older. The frequency of serious infection and other events among tofacitinib treated subjects 65 years of age and older was higher than those under the age of 65.

Of the 4362 patients enrolled in [Study RA-VI](#), 1353 patients were 65 years of age and older (891 patients were treated with tofacitinib and 462 patients were treated with TNFi), including 183 patients over 70 years of age (115 patients treated with tofacitinib and 68 patients treated with TNFi). The frequency of adverse events (including serious infections, all-cause mortality, cardiovascular events, malignancies, non-melanoma skin cancer, gastrointestinal perforations, interstitial lung disease, venous thromboembolism, and arterial thromboembolism) in patients 65 years of age and older was higher than among those under the age of 65.

There were not enough geriatric patients treated with tofacitinib (n=77) in the UC program to adequately study the effects of tofacitinib in this population. As there is a higher incidence of infections in the geriatric population in general, caution should be used when treating the geriatric (see [7 WARNINGS AND PRECAUTIONS](#)).

Sex: Based on population PK analysis, female RA patients were estimated to have 7% lower tofacitinib AUC compared to male RA patients. Female UC patients were estimated to have 15.2% higher tofacitinib AUC compared to male UC patients.

Race: In RA patients, no major differences (<5%) were estimated in tofacitinib AUC between White, Black and Asian RA patients by population PK analysis. In UC patients, population PK analysis indicated that Asian patients had 7.3% higher tofacitinib AUC compared to non-Asian patients. There was a higher incidence of adverse events in Asian patients. Therefore, tofacitinib should be used with caution in Asian patients (see [7 WARNINGS AND PRECAUTIONS](#)).

Body Weight: Population PK analysis in RA patients indicated that systemic exposure (AUC) of tofacitinib in the extremes of body weight (40 kg, 140 kg) were similar to that of a 70 kg patient. An approximately linear relationship between body weight and volume of distribution was observed, resulting in higher peak (C_{max}) and lower trough (C_{min}) concentrations in lighter patients. However, this difference is not considered to be clinically relevant. The between-subject variability (% coefficient of variation) in AUC of tofacitinib is estimated to be approximately 27%. Population PK analysis in UC patients also indicated that tofacitinib AUC did not significantly change with patient body weight.

Hepatic Impairment: tofacitinib is contraindicated in patients with severe hepatic impairment (see [2 CONTRAINDICATIONS](#)). Subjects with mild and moderate hepatic impairment had 3%, and 65% higher tofacitinib AUC, respectively, compared with healthy subjects.

No dose adjustment of tofacitinib is required in patients with mild hepatic impairment.

The recommended total daily dose in patients with moderate hepatic impairment is half the total daily dose recommended for patients with normal hepatic function. The recommended dose is Teva-Tofacitinib 5 mg BID when the indicated dose in the presence of normal hepatic function is Teva-Tofacitinib 10 mg BID; the recommended dose is Teva-Tofacitinib 5 mg once daily when the indicated dose in the presence of normal hepatic function is Teva-Tofacitinib 5 mg BID (see [4 DOSAGE AND ADMINISTRATION](#)).

Tofacitinib has not been studied in patients with severe hepatic impairment or in patients with positive hepatitis B virus or hepatitis C virus serology and should not be used in these populations.

Renal Impairment: Subjects with mild, moderate, and severe renal impairment had 37%, 43% and 123% higher tofacitinib AUC, respectively, compared with healthy subjects. In subjects with ESRD undergoing hemodialysis, the contribution of dialysis to the total clearance of tofacitinib was relatively small.

In subjects with ESRD undergoing hemodialysis, mean AUC was approximately 40% higher compared with historical healthy subject data, consistent with approximately 30% contribution of renal clearance to the total clearance of tofacitinib. Dose adjustment is recommended in ESRD patients under going hemodialysis (see [4 DOSAGE AND ADMINISTRATION](#)).

No dose adjustment of tofacitinib is required in patients with mild renal impairment.

The recommended total daily dose in patients with moderate or severe renal impairment, including patients with ESRD but not limited to those undergoing hemodialysis, is half the total daily dose recommended for patients with normal renal function. The recommended dose is Teva-Tofacitinib 5 mg BID when the indicated dose in the presence of normal renal function is Teva-Tofacitinib 10 mg BID; the


recommended dose is Teva-Tofacitinib 5 mg once daily when the indicated dose in the presence of normal renal function is Teva-Tofacitinib 5 mg BID (see [4 DOSAGE AND ADMINISTRATION](#)).

In clinical trials, tofacitinib was not evaluated in patients with baseline creatinine clearance values (estimated by the Cockcroft-Gault equation) less than 40 mL/min.

Genetic Polymorphism: Mean C_{max} and $AUC_{0-\infty}$ values of tofacitinib following administration of tofacitinib in poor metabolizers of CYP2C19 (carriers of CYP2C19*2/*2, CYP2C19*2/*3 or CYP2C19*3/*3 alleles) were approximately 15% and 17% greater, respectively, than those in normal metabolizers, indicating that CYP2C19 is a minor contributor of tofacitinib clearance.

The impact of intrinsic factors on tofacitinib following administration of tofacitinib pharmacokinetics is summarized in Figure 3 with dosage adjustment recommendations.

Figure 3: Impact of Intrinsic Factors on Tofacitinib Pharmacokinetics



PK=Pharmacokinetics; CI=Confidence Interval

Note: Reference values for weight, age, gender, and race comparisons are 70 kg, 55 years, male, and white, respectively; reference groups for renal and hepatic impairment data are subjects with normal renal and hepatic

function.

^a In RA patients the recommended dose is tofacitinib 5 mg once daily. In UC patients the recommended dose is half the total daily dose indicated for patients with normal renal and hepatic function, i.e., the recommended dose is tofacitinib 5 mg BID when the indicated dose in the presence of normal renal and hepatic function is tofacitinib 10 mg BID, and the recommended dose is tofacitinib 5 mg once daily when the indicated dose in the presence of normal renal and hepatic function is tofacitinib 5 mg BID.

^b Supplemental doses are not necessary in patients after dialysis.

Psoriatic Arthritis

Results from population PK analysis in patients with active PsA were consistent with those in patients with RA.

11 STORAGE, STABILITY AND DISPOSAL

Store between 15°C and 30°C.

12 SPECIAL HANDLING INSTRUCTIONS


This information is not available for this drug product.

PART II: SCIENTIFIC INFORMATION

13 PHARMACEUTICAL INFORMATION

Drug Substance

| | |
|---------------------|---|
| Proper name: | Tofacitinib |
| Chemical name: | (3R,4R)-4-methyl-3-(methyl-7H-pyrrolo[2,3-d]pyrimidin-4-ylamino)-β-oxo-1-piperidine propane nitrile, 2-hydroxy-1,2,3-propane tricarboxylate (1:1) . |
| Molecular formula: | C ₁₆ H ₂₀ N ₆ O•C ₆ H ₈ O ₇ |
| Molecular mass: | 504.5 g/mol (or 312.4 g/mol free base) |
| Structural formula: | |



Physicochemical properties: White to off-white powder. Solubility of tofacitinib is dependent on pH of the medium. The solubility is decreasing with elevated medium pH, where tofacitinib has the least solubility in pH 7.4 buffer and has the highest solubility in pH 1.2 buffer.

14 CLINICAL TRIALS

14.1 Clinical Trials by Indication

Rheumatoid Arthritis

Description of Clinical Studies

The efficacy and safety of tofacitinib were assessed in five randomized, double-blind, multicenter studies in patients ≥18 years with active RA diagnosed according to American College of Rheumatology (ACR) criteria. Patients had ≥6 tender and ≥6 swollen joints at randomization (≥4 swollen and ≥4 tender joints for Study II). Tofacitinib, 5 or 10 mg BID, was given as monotherapy (Study I) and in combination with nonbiologic DMARDs (Study II) in patients with an inadequate response to DMARDs (nonbiologic or biologic). Tofacitinib, 5 or 10 mg BID, was given in combination with methotrexate in patients with either an inadequate response to MTX (Studies III and Study IV) or inadequate efficacy or lack of tolerance to at least one approved TNF-inhibiting biologic agent (Study V).

The primary endpoints for Studies I and V were the proportion of patients who achieved an ACR20 response, mean change from baseline in HAQ-DI and proportion of patients who achieved DAS28-4(ESR) less than 2.6 at Month 3. The primary endpoints for Studies II, III, and IV were the proportion of patients who achieved an ACR20 response at Month 6, mean change from baseline in HAQ-DI at Month 3 and proportion of patients who achieved DAS28-4(ESR) less than 2.6 at Month 6.

Baseline demographics were generally similar among the treatment groups in each study and comparable between the studies. The mean age ranged from 50 to 56 years. Most (80 to 87%) of the patients were female. With the exception of Study A3921044 (46%), the majority (55% to 86%) of the patients in each study were white. The baseline demographics in each study are shown in Table 19.

[Study RA-VI](#) (A3921133) was a randomized, open-label (blinded endpoint), 3-arm parallel-group, multicenter, non-inferiority, safety endpoint study in patients with a diagnosis of moderate to severe active RA and on a stable dose of methotrexate. This study included 4,362 treated RA patients who were aged ≥ 50 years with at least one additional cardiovascular risk factor beyond RA. CV risk factors defined as: current cigarette smoker, diagnosis of hypertension, diabetes mellitus, family history of premature coronary heart disease, history of coronary artery disease including a history of revascularization procedure, coronary artery bypass grafting, myocardial infarction, cardiac arrest, unstable angina, acute coronary syndrome, and presence of extra-articular disease associated with RA, e.g. nodules, Sjögren's syndrome, anemia of chronic disease, pulmonary manifestations.

Patients were randomized 1:1:1 to tofacitinib 5 mg BID, tofacitinib 10 mg BID or TNFi (adalimumab 40 mg SC every 2 weeks in North America or etanercept 50 mg SC every 1 week in the rest of the world). Notably, following a recommendation by the Data and Safety Monitoring Board in February 2019, the dose of tofacitinib in the 10 mg BID arm of the study was reduced to 5 mg BID after it was determined that the frequency of pulmonary embolism was increased in the tofacitinib 10 mg BID treatment arm versus the TNF inhibitor arm. Additionally, all-cause mortality was increased in the tofacitinib 10 mg BID treatment arm versus the TNF inhibitor and tofacitinib 5 mg BID treatment arms. In the final study data, patients in the tofacitinib 10 mg BID treatment arm were analyzed in their originally randomized treatment group. The co-primary endpoints were adjudicated MACE and adjudicated malignancies (excluding NMSC) and the primary analysis for both co-primary endpoints was non-inferiority testing between the two tofacitinib doses combined relative to the TNF inhibitor control. Results showed that the prespecified non-inferiority criteria for these co-primary endpoints were not met for the primary comparison of the combined tofacitinib doses to TNFi. (see [8 ADVERSE REACTIONS](#)).

Study demographics and trial design

Table 19: Summary of patient demographics for clinical trials in RA

| Study # | Trial design | Dosage, route of administration and duration | Study subjects (n) | Age (yrs) Mean (Range) | Female (%) | Mean Disease Duration (yrs) |
|--|---|---|--------------------|------------------------|------------|-----------------------------|
| Background DMARD Studies* | | | | | | |
| A3921046 Study II Sync | MC, DB, PG, PC, R, Background DMARD 12 Months | Tofacitinib: 5 mg BID, 10 mg BID Placebo → 5 mg Placebo → 10 mg NR advance to next period at 3 months, All advance to next period at 6 months | 792 | 52.3 (18-86) | 81.4 | 8.1-10.2 |
| A3921064 Study III Standard | MC, DB, PG, PC, R, Background MTX 12 Months | Tofacitinib: 5 mg BID, 10 mg BID Placebo → 5 mg Placebo → 10 mg Adalimumab 40 mg sc QOW NR advance to next period at 3 months, All advance to next period at 6 months. | 717 | 52.9 (18-83) | 81.7 | 6.9-9.0 |
| A3921044 (1-Year Analysis) Study IV Scan | MC, DB, PG, PC, R, Background MTX 24 Months | Tofacitinib: 5 mg BID, 10 mg BID Placebo → 5 mg Placebo → 10 mg NR advance to next period at 3 months, All advance to next period at 6 months | 797 | [52.0-53.7]** (18-82) | 85.2 | 8.8-9.5 |
| A3921032 Study V Step | MC, DB, PG, PC, R, Background MTX 6 Months | Tofacitinib: 5 mg BID, 10 mg BID Placebo → Tofacitinib 5 mg BID at 3 months | 399 | 55.0 (20-84) | 84.0 | 11.2-13.0 |

| Study # | Trial design | Dosage, route of administration and duration | Study subjects (n) | Age (yrs) Mean (Range) | Female (%) | Mean Disease Duration (yrs) |
|---|-------------------------------|---|--------------------|------------------------|------------|-----------------------------|
| | | Placebo → Tofacitinib 10 mg BID at 3 months | | | | |
| Monotherapy Studies | | | | | | |
| A3921045 (Study I) Solo | MC, DB, PG, PC, R 6 Months | Tofacitinib 5 mg BID, 10 mg BID Placebo → 5 mg Tofacitinib at 3 months, Placebo → 10 mg BID Tofacitinib at 3 months | 610 | 51.8 (21-81) | 86.6 | 7.3-8.6 |
| Post-authorization safety study | | | | | | |
| A3921133 (Study RA-VI) Surveillance | MC, PG, R, OL ~5 Years | Tofacitinib 5 mg BID, 10 mg BID*** Adalimumab 40 mg sc QOW (NA) Etanercept 50 mg sc QW (ROW) | 4362 | 61.15 (50-88) | 78.2 | 10.2-10.4 |
| <p>*In addition to their randomized treatment, all patients in background DMARD studies also received methotrexate (specified in Studies 1032, 1044, and 1064, permitted in Study 1046) or other DMARDs, mostly methotrexate (Study 1046).</p> <p>** Range of mean across treatment groups.</p> <p>*** The tofacitinib 10 mg BID treatment group consists of data from patients that were switched from tofacitinib 10 mg BID to tofacitinib 5 mg BID during the trial as a result of a study modification following a recommendation by the Data and Safety Monitoring Board in February 2019</p> <p>N = number of patients randomized, MC = multicenter, DB = double blind, PG = parallel group, PC = placebo controlled, R = randomized, NR = nonresponder (patient who failed to improve at Month 3 by at least 20% from baseline in the number of swollen and tender/painful joint count), MTX = methotrexate, DMARD = disease modifying antirheumatic drug, sc = subcutaneous, QW = every week, QOW = every other week, LT = long term, OL = open label, NA = North America, ROW = Rest of world.</p> | | | | | | |

Study Results

Clinical Response:

In Studies I and V, patients treated with 5 mg BID tofacitinib had statistically superior ACR20, ACR50, and ACR70 response rates at month 3 vs. placebo-treated patients. In Studies II, III and IV, patients treated with 5 mg BID tofacitinib had statistically superior ACR20, ACR50, and ACR70 response rates at month 3 and 6 vs placebo-treated patients (Table 20). In Studies I, II and V, improvement in ACR20 response rate vs. placebo was observed within 2 weeks. In studies II, III, and IV, ACR response rates were maintained to 12 months in tofacitinib treated patients.

The percent of ACR20 responders by visit for study IV is shown in Figure 4. Similar responses were observed in Studies I, II, III and V.

The proportion of patients with DAS28-4(ESR) less than 2.6 for each study is summarized in Table 22.

Physical Function Response:

Improvement in physical functioning was measured by the HAQ-DI. Patients receiving tofacitinib 5 mg BID demonstrated significantly greater improvement from baseline in physical functioning compared to placebo at month 3 (Studies I, II, III, and V). Tofacitinib 5 mg BID treated patients exhibited significantly greater improved physical functioning compared to placebo as early as week 2 in Studies I and II. In Study III, mean HAQ-DI improvements were maintained to 12 months in tofacitinib –treated patients. At month 3, patients in the tofacitinib 5 mg BID had decreases from baseline in HAQ-DI values (Table 23) which were not less than those of adalimumab-treated patients.

Table 20: Proportion of Patients with an ACR Response

| | Percent of Patients | | | | | | | | | | |
|-------------------------------|---------------------|----------------------------------|-----------------------------|---|---------------------------|---|----------------------------------|---------------------------|---|-------------------------------------|---|
| | Monotherapy | | DMARD Inadequate Responders | | MTX Inadequate Responders | | | MTX Inadequate Responders | | TNF Inhibitor Inadequate Responders | |
| | Study I (SOLO) | | Study II (SYNC) | | Study III (Standard) | | | Study IV (SCAN) | | Study V (STEP) | |
| Response Rate | PBO N=120 | Tofacitinib 5 mg BID N=241 | PBO + DMARD N=157 | Tofacitinib 5 mg BID + DMARD N=311 | PBO + MTX N=106 | Tofacitinib 5 mg BID + MTX N=196 | ADA 40mg QW + MTX N=199 | PBO + MTX N=154 | Tofacitinib 5 mg BID + MTX N=309 | PBO N=131 | Tofacitinib 5 mg BID + MTX N=132 |
| ACR20 [†] Month 3 | 27% | 60%*** 69% | 27% 31% | 56%*** 53%*** | 26% 28% | 61%*** 52%*** | 56%*** 47%** | 27% 25% | 56%*** 51%*** | 24% | 42%* 52% |

| Percent of Patients | | | | | | | | | | | |
|---|----------------|--------|-----------------------------|--------|---------------------------|--------|--------|---------------------------|--------|-------------------------------------|--------|
| | Monotherapy | | DMARD Inadequate Responders | | MTX Inadequate Responders | | | MTX Inadequate Responders | | TNF Inhibitor Inadequate Responders | |
| | Study I (SOLO) | | Study II (SYNC) | | Study III (Standard) | | | Study IV (SCAN) | | Study V (STEP) | |
| Month 6 | NA | | | | | | | | | NA | |
| ACR50^{††} | 13% | 31%*** | 10% | 27%*** | 7% | 34%*** | 24%*** | 8% | 29%*** | 8% | 27%*** |
| Month 3 | NA | 42% | 13% | 34%*** | 12% | 37%*** | 28%** | 8% | 32%*** | NA | 37% |
| Month 6 | | | | | | | | | | | |
| ACR70^{††} | 6% | 15%* | 2% | 8%** | 2% | 12%** | 9%* | 3% | 11%** | 2% | 14%** |
| Month 3 | NA | 22% | 3% | 13%*** | 2% | 20%*** | 9%* | 1% | 15%*** | NA | 16% |
| Month 6 | | | | | | | | | | | |
| * $p < 0.05$, Tofacitinib vs. placebo + MTX/DMARD ** $p < 0.001$, Tofacitinib vs. placebo + MTX/DMARD *** $p < 0.0001$, Tofacitinib vs. placebo + MTX/DMARD † Primary endpoint, Type I error controlled †† Secondary Endpoint, Type I error not controlled | | | | | | | | | | | |

Figure 4: Percentage of ACR20 Responders by Visit for Study IV




Table 21: Proportion of Patients with DAS28-4(ESR) Less Than 2.6

| DAS28-4 (ESR) Less Than 2.6 | Monotherapy | | DMARD Inadequate Responders | | MTX Inadequate Responders | | | MTX Inadequate Responders | | TNF Inhibitor Inadequate Responders | |
|---|----------------|----------------------------------|-----------------------------|---|---------------------------|---|-------------------------------------|---------------------------|---|-------------------------------------|---|
| | Study I (SOLO) | | Study II (SYNC) | | Study III (Standard) | | | Study IV (SCAN) | | Study V (STEP) | |
| | PBO N=122 | Tofacitinib 5 mg BID N=243 | PBO + DMARD N=159 | Tofacitinib 5 mg BID + DMARD N=315 | PBO + MTX N=108 | Tofacitinib 5 mg BID + MTX N=204 | ADA 40mg QW + MTX N=204 | PBO + MTX N=160 | Tofacitinib 5 mg BID + MTX N=321 | PBO N=132 | Tofacitinib 5 mg BID + MTX N=133 |
| Proportion of responders at Month 3 (n) | 4% (5) | 5% (13) | NA | NA | NA | NA | NA | NA | NA | 2% (2) | 6% (8) |
| Proportion of Responders at Month 6 (n) | NA | NA | 3% (4) | 8%* (24) | 1% (1) | 5% (11) | 6%* (12) | 1% (2) | 6%† (19) | NA | NA |

*Statistically significant (p<0.05)
†Statistical significance could not be declared in Study IV due to Step-down procedure
BID = BID, DAS = Disease Activity Score, ESR = erythrocyte sedimentation rate, N=number of patients, n=number of patients meeting pre-specified criteria

Table 22: Mean Change from Baseline in HAQ-DI

| | Monotherapy | | DMARD Inadequate Responders | | MTX Inadequate Responders | | | MTX Inadequate Responders | | TNF Inhibitor Inadequate Responders | |
|---------------------------------|----------------|----------------------------------|-----------------------------|--|---------------------------|---|-------------------------------------|---------------------------|---|-------------------------------------|---|
| | Study I (SOLO) | | Study II (SYNC) | | Study III (Standard) | | | Study IV (SCAN) | | Study V (STEP) | |
| LS Mean Change in HAQ-DI | PBO N=109 | Tofacitinib 5 mg BID N=237 | PBO + DMARD N=147 | Tofacitinib 5 mg BID + DMARD N=292 | PBO + MTX N=98 | Tofacitinib 5 mg BID + MTX N=188 | ADA 40mg QW + MTX N=190 | PBO + MTX N=146 | Tofacitinib 5 mg BID + MTX N=294 | PBO N=118 | Tofacitinib 5 mg BID + MTX N=117 |
| Month 3* | -0.22 | -0.51*** | -0.21 | -0.47*** | -0.25 | - 0.56***- | - 0.51*** | -0.15 | -0.4† | -0.18 | -0.43** |

| | Monotherapy | DMARD Inadequate Responders | MTX Inadequate Responders | MTX Inadequate Responders | TNF Inhibitor Inadequate Responders |
|---|----------------|-----------------------------|---------------------------|---------------------------|-------------------------------------|
| | Study I (SOLO) | Study II (SYNC) | Study III (Standard) | Study IV (SCAN) | Study V (STEP) |
| | | | 0.56*** | | |
| <p>* Primary efficacy time point ** $p < 0.001$, Tofacitinib vs. placebo + MTX/DMARD *** $p < 0.0001$, Tofacitinib vs. placebo + MTX/DMARD † Statistical significance could not be declared in Study IV due to Step-down procedure BID = BID, CI = confidence interval, FAS = full analysis set, LS = least squares, N = number of patients. Results are obtained from a longitudinal linear model with change from baseline as a dependent variable and treatment, baseline, visit, region as fixed effects and patient as random effect.</p> | | | | | |

Psoriatic Arthritis

Description of Clinical Studies

The efficacy and safety of tofacitinib were assessed in 2 multicenter, randomized, double-blind, placebo-controlled trials in 816 patients 18 years of age and older with active psoriatic arthritis. All patients had active psoriatic arthritis for at least 6 months based upon the Classification Criteria for Psoriatic Arthritis (CASPAR), at least 3 tender/painful joints and at least 3 swollen joints at screening and baseline, and active plaque psoriasis at screening. Patients with different psoriatic arthritis subtypes (not mutually exclusive) were enrolled across the 2 clinical trials, including <5 joints or asymmetric involvement (21%), ≥5 joints involved (90%), distal interphalangeal (DIP) joint involvement (61%), arthritis mutilans (8%), enthesitis (80%), dactylitis (53%), total psoriatic body surface area (BSA) >3% (69%), and spondylitis (19%). Patients in these clinical trials had a diagnosis of psoriatic arthritis for a median of 5.5 years (median range 3.0-6.0 years). Of the study population randomized in the double-blind, controlled clinical studies, 54.2% were female and 94.6% were white. The mean age was 48.9 years; 77 (9.4%) patients were 65 years of age or older. All patients were required to receive treatment with a stable dose of a conventional synthetic DMARD (csDMARD; 78% received methotrexate, 13% received sulfasalazine, 7% received leflunomide, 1% received other csDMARDs) and were allowed to receive a stable low dose of oral corticosteroids (21% received equivalent to ≤10 mg/day of prednisone) and/or nonsteroidal anti-inflammatory drugs (NSAIDs; 57% received). In both clinical trials, the primary endpoints were the ACR20 response and the change in HAQ-DI at Month 3.

Study PsA-I (A3921091) was a 12-month clinical trial in 422 patients who had an inadequate response to a csDMARD (67% and 33% were inadequate responders to 1 csDMARD and ≥2 csDMARDs, respectively) and who were naïve to treatment with a TNF-inhibitor biologic DMARD (TNFi). Patients were randomized in a 2:2:2:1:1 ratio to receive tofacitinib 5 mg BID, tofacitinib 10 mg BID, adalimumab 40 mg subcutaneously once every 2 weeks, placebo to tofacitinib 5 mg BID treatment sequence, or placebo to tofacitinib 10 mg BID treatment sequence, respectively; study drug was added to background csDMARD treatment. At the Month 3 visit, all patients randomized to placebo treatment were advanced in a blinded fashion to a predetermined tofacitinib dose of 5 mg or 10 mg BID. Study PsA-I was not designed to demonstrate non-inferiority or superiority to adalimumab.

Study PsA-II (A3921125) was a 6-month clinical trial in 394 patients who had an inadequate response to at least 1 approved TNFi (66%, 19% and 15% were inadequate responders to 1 TNFi, 2 TNFi, and ≥3 TNFi, respectively). Patients were randomized in a 2:2:1:1 ratio to receive tofacitinib 5 mg BID, tofacitinib 10 mg BID, placebo to tofacitinib 5 mg BID treatment sequence, or placebo to tofacitinib 10 mg BID treatment sequence, respectively; study drug was added to background csDMARD treatment. At the Month 3 visit, placebo patients were advanced in a blinded fashion to a predetermined tofacitinib dose of 5 mg or 10 mg BID as in Study PsA-I.

Clinical Response:

Signs and symptoms

At Month 3, patients treated with tofacitinib 5 mg BID had higher ($p \leq 0.05$) response rates versus placebo for ACR20, ACR50, and ACR70 in Study PsA-I and for ACR20 and ACR50 in Study PsA-II; ACR70 response rate was also higher for tofacitinib 5 mg BID versus placebo in Study PsA-II, although the difference versus placebo was not statistically significant ($p > 0.05$) (Table 23).

Table 23: Proportion (%) of PsA Patients Who Achieved Clinical Response and Mean Change from Baseline in PsA-I and PsA-II Studies

| Treatment Group | Conventional Synthetic DMARD Inadequate Responders ^a (TNFi-Naïve) | | | TNFi Inadequate Responders ^b | |
|---------------------|---|-------------------------|--|--|-------------------------|
| | Study PsA-I | | | Study PsA-II ^c | |
| | Placebo | Tofacitinib 5 mg BID | Adalimumab 40 mg SC q2W ^f | Placebo | Tofacitinib 5 mg BID |
| N | 105 | 107 | 106 | 131 | 131 |
| ACR20 | | | | | |
| Month 3 | 33% | 50%* | 52% | 24% | 50%*** |
| Month 6 | NA | 59% | 64% | NA | 60% |
| Month 12 | NA | 68% | 60% | - | - |
| ACR50 | | | | | |
| Month 3 | 10% | 28%** | 33% | 15% | 30%* |
| Month 6 | NA | 38% | 42% | NA | 38% |
| Month 12 | NA | 45% | 41% | - | - |
| ACR70 | | | | | |
| Month 3 | 5% | 17%* | 19% | 10% | 17% |
| Month 6 | NA | 18% | 30% | NA | 21% |
| Month 12 | NA | 23% | 29% | - | - |
| ΔLEI ^d | | | | | |
| Month 3 | -0.4 | -0.8 | -1.1 | -0.5 | -1.3 |
| Month 6 | NA | -1.3 | -1.3 | NA | -1.5 |
| Month 12 | NA | -1.7 | -1.6 | - | - |
| ΔDSS ^d | | | | | |
| Month 3 | -2.0 | -3.5 | -4.0 | -1.9 | -5.2 |
| Month 6 | NA | -5.2 | -5.4 | NA | -6.0 |
| Month 12 | NA | -7.4 | -6.1 | - | - |
| PASI75 ^e | | | | | |
| Month 3 | 15% | 43%*** | 39% | 14% | 21% |
| Month 6 | NA | 46% | 55% | NA | 34% |
| Month 12 | NA | 56% | 56% | - | - |

* p<0.05; ** p<0.001; *** p<0.0001 for active treatment versus placebo at Month 3 achieved statistical significance; with the correction for type 1 error.
Abbreviations: BSA=body surface area; ΔLEI=change from baseline in Leeds Enthesitis Index; ΔDSS=change from baseline in Dactylitis Severity Score; ACR20/50/70=American College of Rheumatology ≥20%, 50%, 70% improvement; csDMARD=conventional synthetic disease modifying antirheumatic drug; N=number of randomised and treated patients; NA=Not applicable, as data for placebo treatment is not available beyond Month 3 due to placebo advanced to tofacitinib 5 mg BID or tofacitinib 10 mg BID; SC q2w=subcutaneously once every 2 weeks; TNFi=tumour necrosis factor inhibitor; PASI=Psoriasis Area and Severity index; PASI75=≥75% improvement in PASI.
a Inadequate response to at least 1 csDMARD due to lack of efficacy and/or intolerability.
b Inadequate response to a least 1 TNFi due to lack of efficacy and/or intolerability. c Study PsA-II had a duration of 6 months.
d Statistical significance cannot be claimed for these endpoints based on step-down testing procedure. Baseline score was >0 in these patients.

| | Conventional Synthetic DMARD Inadequate Responders ^a (TNFi-Naïve) | | | TNFi Inadequate Responders ^b | |
|-----------------|--|----------------------|--------------------------------------|---|----------------------|
| | Study PsA-I | | | Study PsA-II ^c | |
| Treatment Group | Placebo | Tofacitinib 5 mg BID | Adalimumab 40 mg SC q2W ^f | Placebo | Tofacitinib 5 mg BID |

e For patients with Baseline BSA ≥3% and PASI >0. f Arm is not controlled for type 1 error

As with the ACR responses, in patients treated with tofacitinib 5 mg BID in Studies PsA-I and PsA-II, each of the components of the ACR response was consistently improved from baseline at Month 3 including tender/painful and swollen joint counts, patient assessment of arthritis pain, patient and physician's global assessment of arthritis, HAQ-DI, and CRP compared to patients receiving placebo (Table 24).

Table 24: Components of ACR Response at Baseline and Month 3 in Studies PsA-I and PsA-II

| | Conventional Synthetic DMARD Inadequate Responders (TNFi-Naïve) | | | TNFi Inadequate Responders | |
|---|---|----------------------|-------------------------|----------------------------|----------------------|
| | Study PsA-I | | | Study PsA-II | |
| Treatment Group | Placebo | Tofacitinib 5 mg BID | Adalimumab 40 mg SC q2W | Placebo | Tofacitinib 5 mg BID |
| N at Baseline | 105 | 107 | 106 | 131 | 131 |
| ACR Component ^a | | | | | |
| Number of tender/painful joints (0-68) | | | | | |
| Baseline | 20.6 | 20.5 | 17.1 | 19.8 | 20.5 |
| Month 3 | 14.6 | 12.2 | 10.8 | 15.1 | 11.5 |
| Number of swollen joints (0-66) | | | | | |
| Baseline | 11.5 | 12.9 | 9.8 | 10.5 | 12.1 |
| Month 3 | 7.1 | 6.3 | 4.0 | 7.7 | 4.8 |
| Patient assessment of Arthritis pain ^b | | | | | |
| Baseline | 53.2 | 55.7 | 50.7 | 54.9 | 56.4 |
| Month 3 | 44.7 | 34.7 | 32.5 | 48.0 | 36.1 |
| Patient global assessment of arthritis ^b | | | | | |
| Baseline | 53.9 | 54.7 | 50.6 | 55.8 | 57.4 |
| Month 3 | 44.4 | 35.5 | 32.9 | 49.2 | 36.9 |
| HAQ-DI ^c | | | | | |
| Baseline | 1.11 | 1.16 | 1.10 | 1.25 | 1.26 |
| Month 3 | 0.95 | 0.81 | 0.75 | 1.09 | 0.88 |
| Physician's Global Assessment of Arthritis ^b | | | | | |
| Baseline | 53.8 | 54.6 | 50.5 | 53.7 | 53.5 |


| | Conventional Synthetic DMARD Inadequate Responders (TNFi-Naïve) | | | TNFi Inadequate Responders | |
|------------|---|------|------|----------------------------|------|
| | Study PsA-I | | | Study PsA-II | |
| Month 3 | 35.4 | 29.5 | 26.3 | 36.4 | 27.0 |
| CRP (mg/L) | | | | | |
| Baseline | 10.4 | 10.5 | 14.3 | 12.1 | 13.8 |
| Month 3 | 8.60 | 4.02 | 3.10 | 11.44 | 7.72 |

^a Data shown are mean value at baseline and at Month 3
^b Visual analog scale (VAS): 0 = best, 100 = worst
^c HAQ-DI = Health Assessment Questionnaire – Disability Index: 0 = best, 3 = worst; 20 questions; categories: dressing and grooming, arising, eating, walking, hygiene, reach, grip, and activities


The percentage of ACR20 responders by visit for Studies PsA-I and PsA-II is shown in Figure 5. In tofacitinib-treated patients in both Studies PsA-I and PsA-II, significantly higher ACR20 response rates were observed within 2 weeks compared to placebo (Figure 5).

Figure 5: Percentage of ACR20 Responders by Visit

a) Through Month 12 in Study PsA-I



b) Through Month 6 in Study PsA-II^a



In Studies PsA-I and PsA-II, the comparison of tofacitinib 5 mg BID, tofacitinib 10 mg BID, and adalimumab (Study PsA-I only) to placebo was significant (p -value ≤ 0.05) at Months 0.5, 1, 2, and 3.

BID = twice daily; SC q2w = subcutaneously once every 2 weeks.

Patients randomized to placebo treatment were advanced to either tofacitinib 5 mg or 10 mg BID in a blinded manner at Month 3; results for the tofacitinib portion of the placebo \rightarrow tofacitinib treatment sequence (i.e., post-Month 3) are not included in the figure to improve readability.

^a Study PsA-II had a duration of 6 months.

Physical Function:

Improvement in physical functioning was measured by the HAQ-DI. Patients receiving tofacitinib 5 mg BID demonstrated greater improvement ($p \leq 0.05$) from baseline in physical functioning compared to placebo at Month 3 (Table 25).

Table 25: Change from Baseline in HAQ-DI in Studies PsA-I and PsA-II

| Treatment Group | Least Squares Mean Change From Baseline in HAQ-DI | | | | |
|-----------------|--|----------------------|--------------------------------------|---|----------------------|
| | Conventional Synthetic DMARD Inadequate Responders ^a (TNFi-Naïve) | | | TNFi Inadequate Responders ^b | |
| | Study PsA-I | | | Study PsA-II | |
| | Placebo | Tofacitinib 5 mg BID | Adalimumab 40 mg SC q2W ^c | Placebo | Tofacitinib 5 mg BID |
| N | 104 | 107 | 106 | 131 | 129 |
| Month 3 | -0.18 | -0.35* | -0.38 | -0.14 | -0.39*** |
| Month 6 | NA | -0.45 | -0.43 | NA | -0.44 |
| Month 12 | NA | -0.54 | -0.45 | NA | NA |

* $p \leq 0.05$; *** $p < 0.0001$ for active treatment versus placebo at Month 3 achieved statistical significance; with the correction for type 1 error.
Abbreviations: DMARD=disease-modifying antirheumatic drug; HAQ-DI=Health Assessment Questionnaire Disability Index;
N=total number of patients in the statistical analysis; SC q2w=subcutaneously once every 2 weeks; TNFi=tumour necrosis factor inhibitor; NA=Not applicable, as data for placebo treatment is not available beyond Month 3 due to placebo advanced to tofacitinib 5 mg BID or tofacitinib 10 mg BID.

| Least Squares Mean Change From Baseline in HAQ-DI | | | | | |
|--|---------|-------------------------|--|--|-------------------------|
| Conventional Synthetic DMARD Inadequate Responders ^a (TNFi-Naïve) | | | | TNFi Inadequate Responders ^b | |
| Study PsA-I | | | | Study PsA-II | |
| Treatment Group | Placebo | Tofacitinib 5 mg BID | Adalimumab 40 mg SC q2W ^c | Placebo | Tofacitinib 5 mg BID |
| ^a Inadequate response to at least one conventional synthetic DMARD (csDMARD) due to lack of efficacy and/or intolerability. ^b Inadequate response to a least one TNF inhibitor (TNFi) due to lack of efficacy and/or intolerability. ^c Arm is not controlled for type 1 error | | | | | |

The HAQ-DI responder rate (response defined as having decrease from baseline of ≥ 0.35) at Month 3 in Studies PsA-I and PsA-II was 53% and 50%, respectively in patients receiving tofacitinib 5 mg BID, 31% and 28%, respectively in patients receiving placebo, and 53% in patients receiving adalimumab 40 mg subcutaneously once every 2 weeks (Study PsA-I only).

Ulcerative Colitis

Description of Clinical Studies

The efficacy and safety of tofacitinib for the treatment of adult patients with moderately to severely active UC (Mayo score 6 to 12 with endoscopy subscore ≥ 2 and rectal bleeding subscore ≥ 1) were assessed in 3 multicentre, double blind, randomised, placebo controlled studies: 2 identical induction studies OCTAVE Induction 1 and OCTAVE Induction 2 followed by 1 maintenance study OCTAVE Sustain. Enrolled patients had failed at least one conventional therapy, including corticosteroids, immunomodulators, and/or a TNF inhibitor. Concomitant stable doses of oral aminosalicylates and corticosteroids (prednisone daily dose up to 25 mg equivalent) were permitted with taper of corticosteroids to discontinuation mandated within 15 weeks of entering the maintenance study. tofacitinib was administered as monotherapy (i.e., without concomitant use of biologics and immunosuppressants) for UC.

Table 26: Phase 3 Clinical Trials of Tofacitinib 5 and 10 mg BID Doses in Patients with UC

| Studies | OCTAVE Induction 1 (A3921094) | OCTAVE Induction 2 (A3921095) | OCTAVE Sustain (A3921096) |
|---|--|--|---|
| Treatment groups (Randomisation ratio) | Tofacitinib 10 mg BID Placebo (4:1) | Tofacitinib 10 mg BID Placebo (4:1) | Tofacitinib 5 mg BID Tofacitinib 10 mg BID Placebo (1:1:1) |
| Number of patients enrolled | 598 | 541 | 593 |
| Study duration | 8 weeks | 8 weeks | 52 weeks |
| Primary efficacy endpoints | Remission | Remission | Remission |
| Key secondary efficacy endpoints | Improvement of endoscopic appearance of the mucosa | Improvement of endoscopic appearance of the mucosa | Improvement of endoscopic appearance of the mucosa Sustained corticosteroid-free remission among patients in |

| Studies | OCTAVE Induction 1 (A3921094) | OCTAVE Induction 2 (A3921095) | OCTAVE Sustain (A3921096) |
|---------------------------------|----------------------------------|----------------------------------|------------------------------|
| | | | remission at baseline |
| Prior TNFi failure | 51.3% | 52.1% | 44.7% |
| Prior corticosteroid failure | 74.9% | 71.3% | 75.0% |
| Prior immunosuppressant failure | 74.1% | 69.5% | 69.6% |
| Baseline corticosteroid use | 45.5% | 46.8% | 48.7% |

TNFi=tumour necrosis factor inhibitor

In addition, an open-label long-term extension study (OCTAVE Open) was also performed (see further down for more information)

Induction Efficacy Data (OCTAVE Induction 1 and OCTAVE Induction 2):

The primary endpoint of OCTAVE Induction 1 and OCTAVE Induction 2 was the proportion of patients in remission at Week 8 (i.e., a total Mayo score ≤ 2 with no individual subscore >1 , and rectal bleeding subscore of 0). The key secondary endpoint was the proportion of patients with improvement of endoscopic appearance of the mucosa at Week 8. (i.e., endoscopy subscore of 0 or 1). Central endoscopy readings were used for these endpoints.

A significantly greater proportion of patients treated with tofacitinib 10 mg BID achieved remission and improvement of endoscopic appearance of the mucosa at Week 8 compared to placebo in both studies, as shown in Table 27.

Table 27: Proportion of Patients Meeting Efficacy Endpoints at Week 8 (OCTAVE Induction 1 and OCTAVE Induction 2, Central Endoscopy Read)

| Endpoint | OCTAVE Induction 1 | | |
|---|--------------------|--------------------------|---|
| | Placebo | Tofacitinib 10 mg BID | Difference Between Tofacitinib 10 mg BID and Placebo (95% CI) |
| | N=122 | N=476 | |
| Remission ^a | 8.2% | 18.5% | 10.3 (4.3, 16.3) [‡] |
| Improvement of endoscopic appearance of the mucosa ^b | 15.6% | 31.3% | 15.7 (8.1, 23.4) [†] |
| Endpoint | OCTAVE Induction 2 | | |
| | Placebo | Tofacitinib 10 mg BID | Difference Between Tofacitinib 10 mg BID and Placebo (95% CI) |
| | N=112 | N=429 | |
| Remission ^a | 3.6% | 16.6% | 13.0 (8.1, 17.9) [†] |

| Endpoint | OCTAVE Induction 1 | | |
|---|--------------------|--------------------------|---|
| | Placebo | Tofacitinib 10 mg BID | Difference Between Tofacitinib 10 mg BID and Placebo (95% CI) |
| | N=122 | N=476 | |
| Improvement of endoscopic appearance of the mucosa ^b | 11.6% | 28.4% | 16.8 (9.5, 24.1) [†] |

† p<0.001; ‡ p<0.05.
N=number of patients in the analysis set.

a. Primary endpoint: Remission was defined as clinical remission (a Mayo score ≤2 with no individual subscore >1) and rectal bleeding subscore of 0.

b. Key secondary endpoint: improvement of endoscopic appearance of the mucosa was defined as Mayo endoscopy subscore of 0 (normal or inactive disease) or 1 (erythema, decreased vascular pattern).

In both subgroups of patients with or without prior TNF inhibitor failure, a greater proportion of patients treated with tofacitinib 10 mg BID achieved remission and improvement of endoscopic appearance of the mucosa at Week 8 as compared to placebo. This treatment difference was consistent between the 2 subgroups (Table 28).

Table 28: Proportion of Patients Meeting Primary and Key Secondary Efficacy Endpoints at Week 8 by TNF Inhibitor Therapy Subgroups (OCTAVE Induction 1 and OCTAVE Induction 2 Central Endoscopy Read)

| OCTAVE Induction 1 (A3921094) | | |
|---|---------------|-----------------------------------|
| Endpoint | Placebo N=122 | Tofacitinib 10 mg BID N=476 |
| Remission at Week 8 ^a | | |
| With prior TNF inhibitor failure | 1.6% (1/64) | 11.1% (27/243) |
| Without prior TNF inhibitor failure ^b | 15.5% (9/58) | 26.2% (61/233) |
| Improvement of endoscopic appearance of the mucosa at Week 8 ^c | | |
| With prior TNF inhibitor failure | 6.3% (4/64) | 22.6% (55/243) |
| Without prior TNF inhibitor failure ^b | 25.9% (15/58) | 40.3% (94/233) |
| OCTAVE Induction 2 (A3921095) | | |
| Endpoint | Placebo N=112 | Tofacitinib 10 mg BID N=429 |
| Remission at Week 8 ^a | | |
| With prior TNF inhibitor failure ^d | 0.0% (0/60) | 11.7% (26/222) |
| Without prior TNF inhibitor failure ^b | 7.7% (4/52) | 21.7% (45/207) |
| Improvement of endoscopic appearance of the mucosa at Week 8 ^c | | |
| With prior TNF inhibitor failure ^d | 6.7% (4/60) | 21.6% (48/222) |
| Without prior TNF inhibitor failure ^b | 17.3% (9/52) | 35.7% (74/207) |

| OCTAVE Induction 1 (A3921094) | | |
|---|---------------|-----------------------------------|
| Endpoint | Placebo N=122 | Tofacitinib 10 mg BID N=476 |
| TNF=tumour necrosis factor; N=number of patients in the analysis set. | | |
| a. Remission was defined as clinical remission (a Mayo score ≤ 2 with no individual subscore >1) and rectal bleeding subscore of 0. | | |
| b. Failed one or more conventional therapies (corticosteroid, azathioprine, 6-mercaptopurine) but did not have history of prior failure of TNF inhibitor therapy. | | |
| c. Improvement of endoscopic appearance of the mucosa was defined as Mayo endoscopy subscore of 0 (normal or inactive disease) or 1 (erythema, decreased vascular pattern). | | |
| d. Inadequate response, loss of response, or intolerance to TNF inhibitor therapy. | | |

Decreases in rectal bleeding and stool frequency subscores were observed as early as Week 2 in patients treated with tofacitinib.

Clinical response was defined as a decrease from baseline in Mayo score of ≥ 3 points and $\geq 30\%$, with an accompanying decrease in the subscore for rectal bleeding of ≥ 1 point or absolute subscore for rectal bleeding of 0 or 1. Clinical response was observed in 60% of patients treated with tofacitinib 10 mg BID compared to 33% of placebo patients in Octave Induction 1 and 55% compared to 29% in Octave Induction 2.

Maintenance (OCTAVE Sustain):

A total of 593 patients who completed 8 weeks in one of the induction studies and achieved clinical response were re-randomized into OCTAVE Sustain; 179 out of 593 (30%) patients were in remission at baseline of OCTAVE Sustain.

The primary endpoint was the proportion of patients in remission at Week 52. The 2 key secondary endpoints were the proportion of patients with improvement of endoscopic appearance of the mucosa at Week 52, and the proportion of patients with sustained corticosteroid-free remission at both Week 24 and Week 52 among patients in remission at baseline of OCTAVE Sustain.

A significantly greater proportions of patients in both the tofacitinib 5 mg BID and tofacitinib 10 mg BID treatment groups achieved the primary and two key secondary endpoints, as shown in Table 29.

Table 29: Proportion of Patients Meeting Primary and Key Secondary Efficacy Endpoints at Week 52 (Maintenance OCTAVE Sustain, Central Endoscopy Read)

| Endpoint | Placebo N=198 | Tofacitinib 5 mg BID N=198 | Difference Between Tofacitinib 5 mg BID and Placebo (95% CI) | Tofacitinib 10 mg BID N=197 | Difference Between Tofacitinib 10 mg BID and Placebo (95% CI) |
|---|------------------|-------------------------------|--|-----------------------------------|--|
| Remission ^a | 11.1% | 34.3% | 23.2 (15.3, 31.2)* | 40.6% | 29.5 (21.4, 37.6)* |
| Improvement of endoscopic appearance of the mucosa ^b | 13.1% | 37.4% | 24.2 (16.0, 32.5)* | 45.7% | 32.6 (24.2, 41.0)* |
| Sustained | | | 30.3 (17.4, | | 42.2 (27.9, 56.5)* |

| Endpoint | Placebo N=198 | Tofacitinib 5 mg BID N=198 | Difference Between Tofacitinib 5 mg BID and Placebo (95% CI) | Tofacitinib 10 mg BID N=197 | Difference Between Tofacitinib 10 mg BID and Placebo (95% CI) |
|--|--------------------|-------------------------------|--|-----------------------------------|--|
| corticosteroid-free remission at both Week 24 and Week 52 among patients in remission at baseline ^c | N = 59 5.1% | N = 65 35.4% | 43.2)* | N = 55 47.3% | |
| <p>* p<0.0001 N=number of patients in the analysis set.</p> <p>a. Remission was defined as clinical remission (a Mayo score ≤2 with no individual subscore >1) and rectal bleeding subscore of 0.</p> <p>b. Improvement of endoscopic appearance of the mucosa was defined as Mayo endoscopy subscore of 0 (normal or inactive disease) or 1 (erythema, decreased vascular pattern).</p> <p>c. Sustained corticosteroid-free remission was defined as being in remission and not taking corticosteroids for at least 4 weeks prior to the visit at both Week 24 and Week 52.</p> | | | | | |

Additionally, among the 179 patients in remission at baseline (59 in the placebo group, 65 in the tofacitinib 5 mg BID group, and 55 in the tofacitinib 10 mg BID group), the rate of patients with remission at week 52 (i.e., maintained remission) was larger with tofacitinib 5 mg BID (46%) and 10 mg BID (56%) as compared to placebo (10%).

In both subgroups of patients with or without prior TNF inhibitor failure, the proportions of patients treated with either tofacitinib 5 mg BID or tofacitinib 10 mg BID were numerically larger as compared to placebo for the primary and key secondary endpoints, however, statistical significance was not possible to determine (see Table 30).

Table 30: Proportion of Patients Meeting Primary and Key Secondary Efficacy Endpoints in Maintenance Study OCTAVE Sustain (A3921096) by TNF Inhibitor Therapy Subgroup (Central Endoscopy Read)

| Endpoint | Placebo N=198 | Tofacitinib 5 mg BID N=198 | Tofacitinib 10 mg BID N=197 |
|--|----------------|-------------------------------|--------------------------------|
| Remission at Week 52 ^a | | | |
| With prior TNF inhibitor failure ^e | 10/89 (11.2%) | 20/83 (24.1%) | 34/93 (36.6%) |
| Without prior TNF inhibitor failure ^b | 12/109 (11.0%) | 48/115 (41.7%) | 46/104 (44.2%) |
| Improvement of endoscopic appearance of the mucosa at Week 52 ^c | | | |
| With prior TNF inhibitor failure ^e | 11/89 (12.4%) | 25/83 (30.1%) | 37/93 (39.8%) |
| Without prior TNF inhibitor failure ^b | 15/109 (13.8%) | 49/115 (42.6%) | 53/104 (51.0%) |
| Sustained corticosteroid-free remission at both Week 24 and Week 52 among patients in remission at | | | |

| Endpoint | Placebo N=198 | Tofacitinib 5 mg BID N=198 | Tofacitinib 10 mg BID N=197 |
|---|---------------|-------------------------------|--------------------------------|
| baseline ^d | | | |
| With prior TNF inhibitor failure ^e | 1/21 (4.8%) | 4/18 (22.2%) | 7/18 (38.9%) |
| Without prior TNF inhibitor failure ^b | 2/38 (5.3%) | 19/47 (40.4%) | 19/37 (51.4%) |
| TNF=tumour necrosis factor; N=number of patients in the analysis set. a. Remission was defined a Mayo score ≤ 2 with no individual subscore >1 , and rectal bleeding subscore of 0 b. Patients who failed ≥ 1 conventional therapies (corticosteroid, azathioprine, 6-mercaptopurine) but did not have history of prior failure of TNF inhibitor therapy. c. Improvement of endoscopic appearance of the mucosa was defined as Mayo endoscopy subscore of 0 (normal or inactive disease) or 1 (erythema, decreased vascular pattern). d. Sustained corticosteroid-free remission was defined as being in remission and not taking corticosteroids for at least 4 weeks prior to the visit at both Week 24 and Week 52. e. Prior TNF inhibitor failure was defined in this program as inadequate response, loss of response, or intolerance to TNF inhibitor therapy. | | | |

Open-label Extension Study (OCTAVE Open):

Patients who did not achieve clinical response in one of the induction studies (Study OCTAVE Induction 1 or OCTAVE Induction 2) after 8 weeks of tofacitinib 10 mg BID, were allowed to enter an open-label extension study (OCTAVE Open). After an additional 8 weeks of tofacitinib 10 mg BID in Study OCTAVE Open, 53% (155/293) patients achieved clinical response and 14% (42/292) patients achieved remission.

14.2 Comparative Bioavailability Studies

A randomized, single-dose (1 x 10 mg), two-way crossover comparative bioavailability study of Teva-Tofacitinib tablets 10 mg (Teva Canada Limited) with XELJANZ tablets 10 mg (Pfizer Labs, Division of Pfizer Inc., USA) was conducted in healthy, adult, male and female subjects under fasting conditions. Comparative bioavailability data from 48 subjects that were included in the statistical analysis are presented in the following table:

SUMMARY TABLE OF THE COMPARATIVE BIOAVAILABILITY DATA

| Tofacitinib (1 x 10 mg) Geometric Mean Arithmetic Mean (CV %) | | | | |
|--|-----------------------|------------------------|----------------------------|-------------------------|
| Parameter | Test ¹ | Reference ² | % Ratio of Geometric Means | 90% Confidence Interval |
| AUC _T (ng·h/mL) | 352.2 368.5 (31.6) | 340.8 355.8 (29.7) | 103.3 | 100.7 - 106.0 |
| AUC _I (ng·h/mL) | 358.9 375.3 (31.3) | 347.8 362.9 (29.4) | 103.2 | 100.6 - 105.8 |
| C _{max} (ng/mL) | 107.9 111.8 (26.8) | 107.3 110.5 (24.2) | 100.5 | 95.7 - 105.5 |

| Tofacitinib (1 x 10 mg) Geometric Mean Arithmetic Mean (CV %) | | | | |
|--|--------------------|------------------------|----------------------------|-------------------------|
| Parameter | Test ¹ | Reference ² | % Ratio of Geometric Means | 90% Confidence Interval |
| T _{max} ³ (h) | 0.83 (0.33 - 2.00) | 0.83 (0.50 - 2.50) | | |
| T _½ ⁴ (h) | 2.68 (21.28) | 2.61 (21.99) | | |

¹ Teva-Tofacitinib (tofacitinib as tofacitinib citrate) tablets, 10 mg (Teva Canada Limited)

² XELJANZ (tofacitinib as tofacitinib citrate) tablets, 10 mg (Pfizer Labs, Division of Pfizer Inc., USA)

³ Expressed as the median (range) only

⁴ Expressed as the arithmetic mean (CV %) only

15 MICROBIOLOGY

No microbiological information is required for this drug product.

16 NON-CLINICAL TOXICOLOGY

Single and Repeat-Dose Toxicity

Tofacitinib caused death in rats at single oral doses of ≥ 500 mg/kg. Single intravenous doses up to 3 mg/kg did not induce local or systemic toxicity in rats. In cynomolgus monkeys emesis and decreased activity were observed at single oral doses of ≥ 200 mg/kg (divided 3 times daily [TID], ~7 hours apart).

Immune and hematopoietic organ systems were identified as main targets in repeat-dose toxicity studies. Effects on the immune system (including decreased circulating lymphocytes, lymphoid depletion of lymph nodes, spleen, thymus and bone marrow, and bacterial and viral infections) were consistent with inhibition of JAK1/3. Decreases in hemoglobin, hematocrit, erythrocyte numbers and reticulocytes were attributed to JAK2 inhibition. These effects were generally reversible during a 4-week recovery phase in the 4- and 6-week monkey and rat studies, respectively. Repeated oral doses up to 10 mg/kg once daily in rats (up to approximately 15 or 7.6 times human clinical exposure at 5 or 10 mg BID) and 1 mg/kg BID in adult cynomolgus monkeys (approximately 1 or 0.5 times human exposure at 5 or 10 mg BID) were tolerated in studies up to 6 months and 39 weeks duration, respectively. In the 39-week juvenile monkey study, the T-dependent antibody response to antigen immunization was decreased at the high dose of 5 mg/kg BID, approximately 5 or 2.5 times human exposure at 5 or 10 mg BID.

Mutagenesis

Tofacitinib was not mutagenic in the bacterial reverse mutation assay. Reproducible increases in chromosomal abnormalities were observed in a human lymphocyte *in vitro* cytogenetic assay, at high cytotoxic concentrations with metabolic activation, but no effects were observed without metabolic activation. In follow up studies, tofacitinib was not mutagenic in mammalian cells (*in vitro* CHO/HGPRT assay) and did not induce primary DNA damage in an *in vivo/in vitro* rat hepatocyte unscheduled DNA synthesis assay. Tofacitinib was also negative in the *in vivo* rat micronucleus test.

Carcinogenesis

In the 39-week repeat-dose toxicity study in adult monkeys, lymphomas were observed at the high dose of

5 mg/kg BID (approximately 6 times human exposure at 5 mg BID, or approximately 3 times the 10 mg BID dose), but not at the lower dose of 1 mg/kg BID (approximately 1 times human exposure at 5 mg BID, or approximately 0.5 times the 10 mg BID dose). No treatment-related tumors were observed in a 6-month rasH2 transgenic mouse study up to the high dose of 200 mg/kg/day, approximately 38 or 19 times human exposure at 5 or 10 mg BID.

In a 2-year rat carcinogenicity study, tofacitinib induced benign Leydig cell tumors and malignant hibernomas (tumors of brown adipose tissue) at oral doses of ≥ 30 mg/kg/day (≥ 35 times or 17 times human exposure at 5 or 10 mg BID) and benign thymomas at 100/75 mg/kg/day (approximately 187 or 94 times human exposure at 5 or 10 mg BID). No treatment-related tumors were found in rats at 10 mg/kg/day (approximately 16 or 8 times human exposure at 5 or 10 mg BID). The relevance of benign Leydig cell tumors to human risk is unknown.

Developmental and Reproductive Toxicity

Tofacitinib had no effect on fertility of male rats; however, in treated female rats tofacitinib decreased pregnancy rate, numbers of corpora lutea, implantation sites, and viable fetuses, with an increase in early resorptions at oral doses of ≥ 10 mg/kg/day (≥ 15 or 8 times human exposure at 5 or 10 mg BID). The non-observed-adverse-effect-level (NOAEL) for female fertility and early embryonic development was 1 mg/kg/day (approximately 1 or 0.6 times human exposure at 5 or 10 mg BID).

Tofacitinib was teratogenic (external, visceral and skeletal abnormalities) in rabbits and rats at oral doses of 30 and 100 mg/kg/day (approximately 13/6 and 146/73 times human exposure at 5/10 mg BID), respectively. In rabbits, teratogenic effects occurred in the absence of maternal toxicity, consisted of thoracogastroschisis, omphalocele, craniofacial malformations (microstomia, microphthalmia, and cleft lip and palate), membranous ventricular septal defects, gallbladder agenesis, short or absent tail, and skeletal malformations (fused sternbrae and vertebral and/or rib anomalies). In addition, there was an increase in postimplantation loss (early and late resorptions) and consequently, reduced number of viable fetuses. The developmental NOAEL in rabbits was 10 mg/kg/day (approximately 3 or 1.5 times human exposure at 5 or 10 mg BID). In rats, tofacitinib increased postimplantation loss (early and late resorptions), reduced fetal body weights, and increased incidences of fetal malformations at doses that induced maternal toxicity. Malformations suggestive of teratogenicity included anasarca, membranous ventricular septal defects, and skeletal abnormalities (absent cervical arch, bent limb bones, hemicentric thoracic centrum, and rib and sternal anomalies). The developmental NOAEL in rats was 30 mg/kg/day (approximately 58 or 29 times human exposure at 5 or 10 mg BID).

In the peri/postnatal development study in rats, tofacitinib decreased the number of delivered and live born pups, and reduced pup survival at oral doses of 50 mg/kg/day (approximately 73 or 36 times human exposure at 5 or 10 mg BID). There was no effect on sexual maturation, or the ability of these F1 generation rats to learn, mate and produce viable F2 generation fetuses of treatment of the dams at oral doses up to 10 mg/kg/day (up to 15 or 8 times human exposure at 5 or 10 mg BID).

Table 31: Summary of Toxicology Studies

| Study Type | Treatment Duration | Species/ Test system | Animals/ Group | Dose (mg/kg/day) ^a | Results |
|--|--------------------|----------------------|-----------------------|--|---|
| Single-Dose Toxicity | | | | | |
| Single-Dose Oral Toxicity Study in Sprague-Dawley Rats (01-2063-07) | Single Dose | Rat/Sprague-Dawley | 3M, 3F | 0, 500, 1000, 2000 (Oral gavage, 20 mL/kg, 0.5% Methylcellulose/Suspension) | <p>500 mg/kg: 1 female died on Day 1; red-stained fur (nose/muzzle); ↓ eosinophils, ↓ fibrinogen, ↑ ALT, ↑ AST, ↑ glucose, ↑ BUN.</p> <p>≥ 500 mg/kg: ↓ activity, lethargy, partially closed eyes, labored respiration, salivation; lympho-cytolysis in mesenteric lymph node and decreased numbers of lymphocytes within the minimal zone of the splenic white pulp.</p> <p>1000 mg/kg: 6/6 animals died by Day 2; necrosis of centrilobular hepatocytes.</p> <p>≥ 1000 mg/kg: lacrimation and cold to touch; stomach distension; necrosis of individual hepatocytes; lymphocytolysis within the splenic white pulp.</p> <p>2000 mg/kg: 6/6 animals died by Day 2; slow respiration and eye staining/nasal discharge.</p> |
| Single-Dose IV Toxicity Study in Rats with a 14-Day Recovery (09GR453) | Single Dose | Rat/Sprague-Dawley | 10M, 10F ^b | 0, 0.5, 1, 3 (IV, 0.5-3 mL/kg, 10mM Lactic acid in normal saline) | ≤ 3 mg/kg: None |
| Single-Day Oral Toxicity Study in Cynomolgus Monkeys (00-2063-04) | 1 Day | Monkey/Cynomolgus | 2M, 2F | 40, 200, 1000 ^c (Oral gavage, 7 mL/kg, 0.5% Methylcellulose/Suspension) | ≥ 200 mg/kg: Emesis, ↓ activity |
| Repeat-Dose Toxicity | | | | | |
| Pivotal Studies | | | | | |
| 6-Week Oral Toxicity Study with 1-Month Recovery in | 6 Weeks | Rat/Sprague-Dawley | 10-15/sex/dose | 1, 10, 100 Oral gavage, QD, 10 mL/kg | 1 mg/kg/day (LOEL): ↓ WBC count, ↓ lymphocytes, ↓ eosinophils, ↓ basophils, ↓ RBC count, ↓ HCT, ↓ HGB, lymphoid depletion |

| Study Type | Treatment Duration | Species/ Test system | Animals/ Group | Dose (mg/kg/day) ^a | Results |
|--|--------------------|----------------------|----------------|---|--|
| Sprague-Dawley Rats (01-2063-06) | | | | (0.5% Methylcellulose/ Suspension | in bone marrow. 10 mg/kg/day: Same as above, + ↓ reticulocytes, lymphoid depletion in spleen, thymus, and mesenteric lymph node. 100 mg/kg/day: Same as above, + ↑ neutrophils, ↑ AST. 100 mg/kg/day (Recovery): Recovery of reticulocytes and AST, no microscopic findings in lymphoid tissues, partial recovery of WBC count, lymphocytes, RBC parameters, and lymphoid cells in bone marrow. |
| 6-Month Oral Toxicity Study in Rats (77435) | 6 Months | Rat/Sprague-Dawley | 15/sex/dose | 1, 10, 100 (Oral gavage, QD, 10 mL/kg, 0.5% Methylcellulose/ Suspension) | 1 mg/kg/day (LOEL): ↓ WBC, ↓ lymphocytes, ↓ eosinophils, ↓ basophils, ↓ large unstained cells, ↓ RBC count, ↓ HCT, ↓ HGB, ↑ neutrophils (F), ↓ spleen weight, ↓ T lymphocytes, T-cells (CD3+), T-cell subtypes (CD4+, CD8+), B cells (CD45RA+), NK cells (CD161+). 10 mg/kg/day: Same as above, + ↓ reticulocytes; neutrophils, ↑ glucose, ↑ alkaline phosphatase; ↓ triglycerides (F), ↓ spleen weight, lymphoid atrophy (lymph nodes, spleen, thymus) (F), alveolar histiocytosis. 100 mg/kg/day: Same as above, + ↑ neutrophils, ↑ reticulocytes, ↑ globulin; ↓ triglycerides, ↑ liver weight; ↓ thymus weight, lymphoid atrophy (GALT), hepatocellular hypertrophy. |
| 1-Month Oral Toxicity Study with 1-Month Recovery in Cynomolgus Monkeys (01-2063-09) | 4 Weeks | Monkey/ Cynomolgus | 3/sex/dose | 10, 50, 100 Oral gavage, TID ^d , 5 mL/kg, 0.5% Methylcellulose/ Suspension | 10 mg/kg/day: ↓ lymphocytes, ↓ lymphocyte subsets (helper T cells, cytotoxic/suppressor T cells, and NK cells, ↓ HGB. 50 mg/kg/day: Same as above, + death, body weight loss, decreased activity, ↑ WBC, ↓ RBC count, ↓ HCT, ↓ reticulocytes, ↑ AST, ↑ ALT, ↓ Ca, ↓ neutrophil pool, slight granulocytic depletion in |

| Study Type | Treatment Duration | Species/ Test system | Animals/ Group | Dose (mg/kg/day) ^a | Results |
|--|--------------------|--|----------------|---|--|
| | | | | | bone marrow, lymphoid depletion in spleen, bacterial and viral infection secondary to immunosuppression in heart, kidney, gastrointestinal tract, buccal cavity, and skin. 100 mg/kg/day: Same as above (except no ↑ WBC count), + RBC depletion in bone marrow, and ↑ immature myeloid cells in bone marrow, lymphoid depletion in mesenteric lymph node. 50 mg/kg/day (Recovery: Complete recovery with the exceptions of partial recovery of ↑ neutrophils, ↑ ALT and ↑ AST, ↓ (CD16+, CD3-), ↓ RBC count; rebound effect in lymphocytes, (CD4+, CD3+), and (CD8+, CD3+), lymphocytes, and reticulocytes. |
| 39-Week Oral Toxicity Study in Monkeys (2003-0301) | 39 weeks | Monkey/ Cynomolgus | 4/sex/dose | 0.5, 2, 10 ^e Oral gavage, BID, 10 mL/kg, 0.5% Methylcellulose/ Suspension | 0.5 mg/kg/day (LOEL): ↓ total lymphocytes, ↓ lymphocyte subsets (T-helper, -cytotoxic/suppressor and NK cells); lymphoid hyperplasia (2/4 M). 2 mg/kg/day: Same as above +, ↓ RBC count, ↓ HCT, ↓ HGB, lymphoid hyperplasia (4/4 M) 10 mg/kg/day: Same as above, + death, ↑ reticulocytes; RBC hyperplasia in bone marrow; lymphoid hyperplasia (3/4 M, 1/4 F); lymphoma (1/4 M, 2/4 F; 2 confirmed B-cell origin), mononuclear cell infiltrates in the heart (F). |
| Genotoxicity | | | | | |
| In Vitro Studies | | | | | |
| Microbial Reverse Bacterial Mutation Assay (AMES) (01-2063-11) | In Vitro | <i>Salmonella typhimurium</i> , <i>Escherichia coli</i> | NA | 0.010-5 mg/plate Plate Incorporation for ~ 48 to 72 hours at 37°C | No genotoxic effect. No cytotoxic effect. |
| Mammalian Cell Mutation Assays (01-2063-16) | In Vitro | Chinese Hamster ovary (CHO)-K1-BH4 cells, | NA | 16-5000 µg/mL 5-hour treatment, 6-8 day incubation | - No Genotoxic effects - Substantial cytotoxicity at 950, 1000, and 1100 µg/mL with average Day 3 relative cell survivals of 43%, 29%, and 17%, respectively. |

| Study Type | Treatment Duration | Species/ Test system | Animals/ Group | Dose (mg/kg/day) ^a | Results |
|---|---|---|----------------|--|--|
| In Vitro Cytogenetics Assay (01-2063-10) | In Vitro | Human Peripheral Lymphocytes | NA | 41.8-2400 µg/mL 3 hours with activation, 3 and 24 hours without activation | Cytotoxic Effects: ~ 50% Mitotic suppression achieved in all treatments. Genotoxic Effects: tofacitinib did not significantly increase structural chromosome aberrations at 3- and 24-hour treatments without metabolic activation. At 3 hours with metabolic activation, tofacitinib increased structural chromosome aberrations at relatively cytotoxic concentrations. |
| In Vivo Studies | | | | | |
| In Vivo/In Vitro Rat Hepatocyte Unscheduled DNA Synthesis Study (01- 2063-17) | Single Dose Hepatocytes, 2-4 and 14-16 HPD | Rat/Sprague- Dawley | M | 125, 250, 250 Oral gavage, 10 mL/kg, 0.5% Methylcellulose | Toxic/Cytotoxic Effects: Hypoactivity, labored breathing and/or squinted eyes in the 500 mg/kg group Genotoxic Effects: None |
| In Vivo Cytogenetics (Rat Micronucleus) (01- 2063-12) | Once daily for 3 days | Rat/Sprague- Dawley | 6M, 6F | 62.5, 125, 250 Oral gavage, QD, 10 mL/kg, 0.5% Methylcellulose | Toxic/Cytotoxic Effects: No mortality or adverse clinical signs attributed to drug treatment was observed. A statistically significant decrease in mean percent body weight gain was evident in the male rats. The males also showed statistically significant treatment-related reduction in mean %PCE, suggestive of bone marrow toxicity. Genotoxic Effects: None. |
| Carcinogenicity | | | | | |
| <u>6-Month Oral Gavage Study in Mice</u> (8200-368) | 6 Months | Mouse/Model 001178-T (hemizygous), CB6F1/Jic- TgrasH2@Tac Mouse/Model 001178-W (homozygous wild-type), CB6F1/Jic- TgrasH2@Tac | 25/sex/dose | 25, 75, 200 Oral gavage, QD, 10 mL/kg, 0.5% (w/v) Methylcellulose/ Solution | ≥25 mg/kg/day: No evidence of treatment- related carcinogenicity. |
| <u>2-Year Oral Gavage in</u> | 103 Weeks ^f | Rat/Sprague- | 60-70/ | 10/10, 30/30, | 10 mg/kg/day: Benign angiomas of |

| Study Type | Treatment Duration | Species/ Test system | Animals/ Group | Dose (mg/kg/day) ^a | Results |
|--|--|--|---|--|---|
| Rats (6348-463) | | Dawley | sex/dose | 75/100 ^g Oral gavage, QD, 10 mL/kg, 0.5% Methylcellulose/ Solution | mesenteric lymph nodes (M). 30 mg/kg/day: Hyperplasia and benign tumors of interstitial cells of testes (M), malignant hibernomas of multiple organs (F). 75 mg/kg/day: Same as above (M). 100/75 mg/kg/day: Benign thymoma in thymus (F). |
| Investigative | | | | | |
| 14-Day Oral Investigative Study in Rats (10GR431) | 14 Days | Rat/Sprague-Dawley | 8F with BrdU pumps 5F without BrdU pumps | Oral gavage, QD, 10 mL/kg, 0.5% Methylcellulose/ Solution | Tofacitinib inhibited JAK/STAT signaling in BAT as evidenced by decreased tissue levels of phosphorylated STAT3 (pSTAT3) and pSTAT5 at doses ≥ 10 mg/kg/day. |
| Investigative Study with Rat Brown Adipocytes (11GR016) | 1 hour pre-incubation with tofacitinib then 20 minutes with oPRL and tofacitinib | Rat/Sprague-Dawley/Primary Leydig cells | In vitro | 150 mM NaCl, 0.03 mM NaHCO ₃ /Solution (oPRL), 0.1% dimethyl sulphoxide/Solution (tofacitinib) | Tofacitinib inhibited the prolactin-induced increase in STAT5A/B phosphorylation. |
| Investigative Study with Rat Primary Leydig Cells (11GR015) | 1 hour pre-incubation with tofacitinib then 15 minutes with oPRL and tofacitinib | Rat/Sprague-Dawley/ Differentiated primary brown adipocytes/ pSTAT5A/B protein | In vitro | 150 mM NaCl, 0.03 mM NaHCO ₃ /Solution (oPRL), 0.1% dimethyl sulphoxide/Solution (tofacitinib) | Tofacitinib inhibited the prolactin-induced increase in STAT5A/B phosphorylation. |
| Reproductive and Developmental Toxicity | | | | | |
| Oral Fertility and Embryonic Development Study in Male and Female Rats (05GR051) | (F) Phase 1: 14 Days pre mating, throughout cohabitation | Rat/Sprague Dawley | 20/sex/dose | 1, 10, 100 Oral Gavage, QD, 10 mL/kg | 1 mg/kg/day: No effect. 10 mg/kg/day: \uparrow Post implantation loss. 100 mg/kg/day: Same as above, + \downarrow pregnancy rate, \downarrow corpora lutea, \downarrow implantation sites, \downarrow viable fetuses, \uparrow early |

| Study Type | Treatment Duration | Species/ Test system | Animals/ Group | Dose (mg/kg/day) ^a | Results |
|--|---|--------------------------|----------------|--|--|
| | and through GD 7. (M) Phase 2: Minimum of 63 days (beginning 28 days pre-mating) | | | | resorptions, ↑ pre-implantation loss. |
| Oral Embryo-Fetal Development Study in Rats (04-2063-24) | GD 6-17 | Rat/Sprague Dawley | 20F/dose | 1, 10, 30 Oral gavage, QD, 10 mL/kg | ≥1 mg/kg/day: No effect. |
| Oral Embryo-Fetal Development Study in Rats (09GR353) | GD 6-17 | Rat/Sprague Dawley | 20F/dose | 30, 100, 300 Oral gavage, QD, 10 mL/kg | 30 mg/kg/day: No effect. 100 mg/kg/day: ↓ Viable fetuses, ↓ uterine weight, external, visceral and skeletal malformations. 300 mg/kg/day: ↓ Maternal body weight and food consumption, clinical signs of poor toleration, no viable fetuses to examine. |
| Oral Embryo-Fetal Development Study in Rabbits (05-2063-25) | GD 7-19 | Rabbit/New Zealand White | 20F/dose | 10, 30, 100 Oral gavage, QD, 2 mL/kg | 10 mg/kg/day: No effect. 30 mg/kg/day: ↓ Viable fetuses, ↓ uterine weight, external, visceral, and skeletal malformations. 100 mg/kg/day: Same as above, + ↓ fetal body weights, ↑ visceral variations. |
| Oral Developmental Peri/Postnatal Reproduction including Postnatal Behavioral/Functional Evaluation in Rats (LIA00468) | GD 6 - DL 21 (or GD 24 for rats not delivering a litter) | Rat/Sprague-Dawley | 25F/dose | Oral gavage, QD during dosage period; 10 mL/kg | 10 mg/kg/day: No effect 50 mg/kg/day: ↓ Delivered pups, ↓ liveborn pups, ↓ pup survival, ↓ pup body weight. |
| Developmental and Reproductive - Juvenile | | | | | |
| Oral Fertility Study in Juvenile Rats | PND 21-70 (M) | Rat/Sprague-Dawley | 20/sex/dose | 1, 10, 100 Oral gavage, QD, | 1 mg/kg/day: No effect. 10 mg/kg/day: ↓ BW (M), ↓ BW gain (M). |

| Study Type | Treatment Duration | Species/ Test system | Animals/ Group | Dose (mg/kg/day) ^a | Results |
|---|--------------------|----------------------|----------------|--|--|
| (10GR250) | PND 21-55 (F) | | | 10 mL/kg 0.5% (w/v) Methylcellulose/ Suspension | 100 mg/kg/day: Same as above (M&F). |
| Oral Toxicity Study in Juvenile Rats with a 2-Month Recovery (10GR307) | PND 21-49 | Rat/Sprague Dawley | 16/sex/dose | 1, 10, 100 Oral gavage, QD, 10 mL/kg 0.5% (w/v) Methylcellulose/ Suspension | 1 mg/kg/day: Females: ↓WBC, ↓ lymphocytes, eosinophils, basophils Males only: ↑ vacuolation in brown adipose tissue, ↓ T cells, ↓ helper T cells, ↓ cytotoxic T cells, ↓ B cells, ↓ NK cells. 10 mg/kg/day: Same as above, ↓ T cells, ↓ helper T cells, ↓ cytotoxic T cells, ↓ B cells, ↓ NK cells. Males: ↓ WBC, ↓ lymphocytes, eosinophils, basophils. Females: ↓ body weight and body weight gain, ↓ reticulocytes, ↓ cellularity (thymus) - females, ↓ cellularity (spleen), ↓ lymphoid cellularity-mesenteric lymph node. 100 mg/kg/day: Same as above, ↓ body weight and body weight gain (M), ↓ RBC, ↓ cellularity: inguino-femoral lymph node, mandibular lymph node. |
| 39-Week Oral Toxicity in Juvenile Monkeys with a 26-Week Recovery (Interim Report) (2501-010) | 39 Weeks | Monkey/ Cynomolgus | 4/sex/dose | 0.5, 2, 10 Oral gavage, BID, 5 mL/kg 0.5% (w/v) Methylcellulose/ Suspension | 0.5 mg/kg/day: No effect. 2 mg/kg/day: ↓ total lymphocytes (M), ↓ lymphocyte subsets (NK cells, effector CD8+ T cells, CD8+ T cells (M), ↓ thymus weight (M), ↓ spleen weight (F). 10 mg/kg/day: ↓ total lymphocytes (M + F), ↓ RBC count, ↓ HCT, ↓ HGB, ↓ lymphocyte subsets (NK cells, CD4+ and CD8+ T cells, naïve CD4+ and CD8+ T cells, central and effector memory CD8+ cells), ↓ spleen and thymus weight. |

^a Doses are expressed as mg active moiety/kg/day unless otherwise noted.

^b Five/sex were necropsied on Day 2 and 5/sex were retained for a 14-day recovery period and necropsied on Day 15.

^c 13, 67, 333 mg/kg TID; 7 hours apart.

| Study Type | Treatment Duration | Species/ Test system | Animals/ Group | Dose (mg/kg/day) ^a | Results |
|---|--------------------|----------------------|----------------|-------------------------------|---------|
| <p>^d 3.33, 16.7, 33.3mg/kg TID; 7 hours apart.</p> <p>^e 0.25, 1, 5, mg/kg BID; 12 hours apart.</p> <p>^f All surviving males in Group 4 were sacrificed on Day 654 (Week 94) of the dosing phase. All surviving males in Group 1 through Group 3 were sacrificed on Day 686 (Week 98) of the dosing phase. All surviving females were sacrificed on Day 715 (Week 103) of the dosing phase.</p> <p>^g Dose was lowered from 100 to 75 mg/kg/day starting on Day 133.</p> <p>ALT = Alanine aminotransferase; AST = Aspartate aminotransferase; BAT = Brown adipose tissue; BID = Twice daily; BrdU = 5-bromo-2'-deoxyuridine; BUN = Blood urea nitrogen; Ca = Calcium; CHO = Chinese hamster ovary; CD = Cluster of differentiation; DL = Day of lactation; F = Female; GALT = Gut associated lymphoid tissue; GGT = Gamma glutamyl transferase; GD = Gestation Day; HGB = Hemoglobin; HCT = Hematocrit; HPD = Hours postdose; IV = Intravenous; JAK = Janus kinase; LOEL = Lowest observed effect level; M = Male; NA = Not applicable; NaCl = Sodium chloride; NaHCO₃ = Sodium bicarbonate; NK = Natural killer; oPRL = Ovine prolactin; PND = Postnatal day; PCE = Polychromatic erythrocytes; pSTAT = Phosphorylated signal transducer and activator of transcription; QD = Once daily; RBC = Red blood cells; STAT = Signal transducer and activator of transcription; TID = Three times daily; WBC = White blood cells.</p> | | | | | |

17 SUPPORTING PRODUCT MONOGRAPHS

1. XELJANZ® (tofacitinib tablets 5 mg and 10 mg) , Control Number 272662, Product Monograph, Pfizer Canada ULC, Quebec, Canada, dated AUG 22, 2023.

PATIENT MEDICATION INFORMATION

READ THIS FOR SAFE AND EFFECTIVE USE OF YOUR MEDICINE

^{Pr}Teva-Tofacitinib Tofacitinib tablets

Read this carefully before you start taking Teva-Tofacitinib and each time you get a refill. This leaflet is a summary and will not tell you everything about this drug. Talk to your healthcare professional about your medical condition and treatment and ask if there is any new information about Teva-Tofacitinib.

Serious Warnings and Precautions

Serious Infections

- You should NOT take Teva-Tofacitinib if you have an active infection.
- Teva-Tofacitinib is a medicine that affects your immune system. It can lower the ability of your body to fight infections such as tuberculosis, shingles (herpes zoster) and infections caused by other bacteria, fungi, or viruses that can spread throughout the body.
- In some cases, these infections may lead to hospitalization or death.
- Most patients who developed infections were taking other medicines, such as methotrexate or corticosteroids at the same time. These medicines make it harder to fight infections.
- Your healthcare professional will closely monitor you for the signs and symptoms of infections during and after the treatment with Teva-Tofacitinib.
- Contact your healthcare professional if you have any signs or symptoms of an infection, such as:
 - fever, sweating, or chills,
 - muscle aches,
 - cough, shortness of breath,
 - blood in spit,
 - weight loss,
 - warm, red, or painful skin or sores on your body,
 - diarrhea or stomach pain,
 - burning when you urinate or urinating more often than normal,
 - feeling very tired.
- If a serious infection develops, stop taking Teva-Tofacitinib and contact your healthcare professional right away.

Cancers and immune conditions

- Lymphoma, lung cancer, and other cancers have been reported in patients treated with tofacitinib.
- Your healthcare professional will closely monitor you for signs and symptoms of cancer and other serious conditions during treatment with Teva-Tofacitinib.

Blood clots

- Blood clots in the veins of your legs or arms (deep vein thrombosis, DVT), arteries (arterial thrombosis) or lungs (pulmonary embolism, PE) can happen in some people taking Teva-

Tofacitinib. This may be life-threatening and cause death.

- Stop taking Teva-Tofacitinib and seek medical help right away if you develop any signs or symptoms of:
 - Blood clots in your leg (such as swelling, pain or tenderness); or
 - Blood clots in your lung (such as sudden unexplained chest pain or shortness of breath).

Major heart problems

- Major heart problems have been reported in Rheumatoid Arthritis patients treated with tofacitinib.
- Talk to your healthcare professional about possible heart disease risk factors before you start taking Teva-Tofacitinib.
- If you develop signs and symptoms of a heart problem, stop taking Teva-Tofacitinib and contact your healthcare professional right away. Symptoms may include:
 - new or worsening chest pain,
 - shortness of breath,
 - irregular heartbeats,
 - swelling of the legs.

What is Teva-Tofacitinib used for?

- **Rheumatoid Arthritis**
Teva-Tofacitinib (tofacitinib) is used to treat adults with rheumatoid arthritis (RA) when other treatments do not work. Teva-Tofacitinib may be taken alone or in combination with methotrexate.
- **Psoriatic Arthritis**
Teva-Tofacitinib is used to treat adults with active psoriatic arthritis (PsA) when other medicines do not work. Teva-Tofacitinib may be taken alone or in combination with methotrexate or other medicines called conventional synthetic disease modifying antirheumatic drug (csDMARD).
- **Ulcerative Colitis**
Teva-Tofacitinib is used to treat adults with moderately to severely active ulcerative colitis (UC) when other medicines do not work.

How does Teva-Tofacitinib work?

Teva-Tofacitinib is a Janus Kinase (JAK) inhibitor. JAK is a type of enzyme which helps start the immune response in your body. Teva-Tofacitinib is believed to interfere with the activity of the JAK enzyme to reduce the immune response. This helps reduce signs and symptoms of rheumatoid arthritis, psoriatic arthritis and ulcerative colitis.

What are the ingredients in Teva-Tofacitinib?

Medicinal ingredients: Tofacitinib citrate

Non-medicinal ingredients:

5 mg tablet core contains: Croscarmellose sodium, lactose monohydrate, magnesium stearate, microcrystalline cellulose. The film coat contains: HPMC/hypromellose, macrogol/PEG, titanium dioxide.

10 mg tablet core contains: Croscarmellose sodium, lactose monohydrate, magnesium stearate, microcrystalline cellulose. The film coat contains: FD&C blue #2/indigo carmine aluminum lake, HPMC/hypromellose, macrogol/PEG, titanium dioxide.

Teva-Tofacitinib comes in the following dosage forms:

5 mg and 10 mg tablets in bottles.

Do not use Teva-Tofacitinib if:

- you are allergic to tofacitinib or any other non-medicinal ingredients in Teva-Tofacitinib.
- you are pregnant or are planning to become pregnant.
- you are breast-feeding or intend to breast-feed. Talk to your healthcare professional about the best way to feed your baby while taking Teva-Tofacitinib.
- you have severe liver problems.

To help avoid side effects and ensure proper use, talk to your healthcare professional before you take Teva-Tofacitinib. Talk about any health conditions or problems you may have, including if you:

- are being treated for an infection, get a lot of infections or have infections that keep coming back;
- have diabetes, HIV/AIDS, or a weak immune system. People with these conditions have a higher chance for infections;
- have tuberculosis, or a history of tuberculosis or have been in close contact with someone with tuberculosis;
- have or have had hepatitis B or C;
- have known narrowing or blockage of your digestive tract (intestines or another part of your bowel are not as wide as normal).
- have gastrointestinal problems, including gastrointestinal perforations (tear in the stomach or intestines), diverticulitis (inflammation in parts of the large intestine), ulcers in your stomach or intestines;
- have low blood counts. Treatment with Teva-Tofacitinib can be associated with low red blood cell counts (anemia), or with low white blood cell counts (neutrophils or lymphocytes);
- have high cholesterol;
- have or have had any type of cancer;
- have liver problems;
- have kidney problems;
- have a history of interstitial lung disease (diseases that inflame or scar lung tissue);

- have muscle pain or muscle weakness;
- develop new skin lesions during or after therapy or if existing lesions change appearance;
- have received any vaccines (shots) within 1 month prior to starting Teva-Tofacitinib or are planning to get vaccinated. Certain types of vaccines (shots) should not be given when taking Teva-Tofacitinib. Before you start Teva-Tofacitinib, you should be up to date with all recommended vaccinations, including a shingles vaccine;
- have had blood clots in your legs (deep vein thrombosis), eyes (retinal venous thrombosis) or lungs (pulmonary embolism) or have been told you are at risk of blood clots;
- have problems with your blood clotting (thrombophilia);
- have chest pain, heart failure or any heart problems, or heart disease risk factors, such as if you:
 - are a current or past smoker,
 - have high blood pressure (hypertension),
 - have diabetes,
 - have a family history of premature coronary heart disease,
 - have had coronary artery disease. This is when blood vessels that supply your heart are clogged;
- have other diseases associated with rheumatoid arthritis, such as: lumps (nodules), anemia, lung problems, a type of immune problem called Sjögren’s syndrome
- are of Asian descent. You may be at increased risk of serious side effects
- have risk factors for broken bones, such as if you: are older than 65 years of age, are a woman, or take a type of medicine called corticosteroids.

Other warnings you should know about:

Blood tests and monitoring

You may need blood tests before you start Teva-Tofacitinib. These tests may be repeated while you are taking Teva-Tofacitinib. Your healthcare professional will also monitor your liver tests and blood cholesterol levels 4 to 8 weeks after you start receiving Teva-Tofacitinib and routinely thereafter. These will help your healthcare professional find out how Teva-Tofacitinib is affecting your blood and how well your liver is working

Female patients

Pregnancy and birth control

- Avoid becoming pregnant while taking Teva-Tofacitinib. It may harm your unborn baby.
- If you are of child-bearing age, use an effective method of birth control while taking Teva-Tofacitinib. Continue using birth control for 4 to 6 weeks after you stop taking Teva-Tofacitinib.

Adults aged 65 years and older

Side effects, including serious side effects, have occurred more often in patients aged 65 years and older.

Tell your healthcare professional about all the medicines you take, including any drugs, vitamins, minerals, natural supplements or alternative medicines.

The following may interact with Teva-Tofacitinib:

- other medicines used to treat rheumatoid arthritis, psoriatic arthritis or ulcerative colitis, including:
 - Biologics such as: abatacept, adalimumab, anakinra, certolizumab, etanercept, golimumab, infliximab, rituximab, secukinumab, ustekimumab, vedolizumab,
 - Other JAK inhibitors such as: baricitinib and upadacitinib.
- medicines that affect your immune system (such as azathioprine, 6-mercaptopurine, tacrolimus, sirolimus, cyclosporine)
- antiarrhythmics (medicines used to treat heart rhythm problems)
- beta-blockers (medicines used to slow the heart or lower blood pressure), and calcium channel blockers (medicines used to lower blood pressure)
- cholinesterase inhibitors (medicines used to treat Alzheimer's)
- HIV protease inhibitors
- a medicine used to treat bacterial infections like tuberculosis called rifampin, and medicines for fungal infections (such as ketoconazole, fluconazole)
- grapefruit juice
- St. John's Wort (an herbal medicine also known as hypericum perforatum). It may reduce the response to Teva-Tofacitinib.

How to take Teva-Tofacitinib:

- Always take Teva-Tofacitinib exactly as your healthcare professional tells you.
- Teva-Tofacitinib can be taken with or without food.
- Your doctor may reduce the dose if you have liver or kidney problems. You should not increase the dose.
- Teva-Tofacitinib should not be used if you have or develop a serious infection until the infection is controlled.

Your healthcare professional may prescribe Teva-Tofacitinib alone or in combination with other medication(s). If you receive treatment with another drug, your healthcare professional will tell you how to take it. Be sure to read the package leaflets for the other drugs as well as this one.

Usual dose:

Adults (18 years and older:

Rheumatoid Arthritis:

- The recommended dose is 5 mg taken by mouth twice daily.

Psoriatic Arthritis:

- The recommended dose is 5 mg, taken by mouth twice daily.

Ulcerative Colitis:

- The recommended dose is 10 mg, twice daily for the first 8 weeks. After 8 weeks, your doctor will decide to give you 5 mg or 10 mg twice daily for maintenance.
- Your doctor may decide to stop your treatment with Teva-Tofacitinib if it does not work for you within 16 weeks.

Overdose:

If you think you, or a person you are caring for, have taken too much Teva-Tofacitinib, contact a healthcare professional, hospital emergency department, or regional poison control centre immediately, even if there are no symptoms.

Missed Dose:

If you have missed your dose of Teva-Tofacitinib, take the next dose as planned at the next scheduled time. Do NOT take a double dose to make up for a forgotten dose.

What are possible side effects from using Teva-Tofacitinib?

These are not all the possible side effects you may have when taking Teva-Tofacitinib. If you experience any side effects not listed here, tell your healthcare professional.

The side effects of Teva-Tofacitinib include:

- Upper respiratory tract infection (such as a cold)
- Nasopharyngitis (nose or throat infection, runny or stuffy nose), cough
- Headache, dizziness
- Diarrhea, vomiting, nausea (feeling queasy, feeling like you may throw up)
- Indigestion (heartburn or upset stomach)
- Back pain, joint pain
- Rash
- Muscle weakness/pain

If any of the above affects you severely, tell your healthcare professional.

Teva-Tofacitinib may cause abnormal blood test results, including changes in cholesterol levels, white or red blood cell counts or creatinine levels (a protein that may increase in people with kidney problems). Your healthcare professional will decide when to perform blood tests and will interpret the results.

| Serious side effects and what to do about them | | | |
|--|---|---------------------|--|
| Symptom/ effect | Talk to your healthcare professional | | Stop taking drug and get immediate medical help |
| | Only if severe | In all cases | |
| COMMON | | | |
| Broken bones | | ✓ | |
| Cellulitis: skin infection with redness, swelling and pain | | ✓ | |
| Gastritis: stomach ache, loss of appetite | | ✓ | |
| Herpes Zoster (shingles): skin rash or blisters usually on one side of the body with itching, burning or tingling pain | | | ✓ |
| Hypertension (high blood pressure): measured high blood pressure, sometimes with headache or nosebleed | | ✓ | |
| Myocardial infarction (heart attack): pressure or squeezing pain between the shoulder blades, in the chest, jaw, left arm or upper abdomen, shortness of breath, dizziness, fatigue, light- | | | ✓ |

| Serious side effects and what to do about them | | | |
|--|--------------------------------------|--------------|---|
| Symptom/ effect | Talk to your healthcare professional | | Stop taking drug and get immediate medical help |
| | Only if severe | In all cases | |
| headedness, clammy skin, sweating, indigestion, anxiety, feeling faint and possible irregular heartbeat | | | |
| Pneumonia: infection with coughing, fever, fatigue | | ✓ | |
| Urinary tract infections: difficulty or increased need to urinate; pain or burning sensation when passing urine, pain in the pelvis or mid-back, urine that appears cloudy | | ✓ | |
| UNCOMMON | | | |
| Allergic reaction: hives, rash, swelling of the face, lips, tongue, and throat that may cause difficulty in breathing or swallowing | | | ✓ |
| Anemia/neutropenia/ lymphopenia (low blood cell counts): fatigue, loss of energy, weakness, shortness of breath | | ✓ | |
| Bronchitis: persistent cough, fatigue, shortness of breath | | ✓ | |
| Congestive heart failure: shortness of breath when you exert yourself or lie down, swelling in your legs, ankles and feet, irregular heartbeat, persistent cough | | | ✓ |
| Deep vein thrombosis (blood clot in the leg): swelling, pain or tenderness in the leg | | | ✓ |
| Retinal venous thrombosis (blood clot in the eyes): blurry vision, partial or complete loss of vision | | | ✓ |
| Flu: cough, sore throat, feverish chills | | ✓ | |
| Increased creatine kinase levels: muscle weakness and/or muscle pain | ✓ | | |
| Kidney problems: change in the amount, frequency or colour (pale or dark) of urine | | ✓ | |
| Liver problems: yellowing of the skin or eyes, dark urine, abdominal pain, nausea, throwing up, loss of appetite with itching | | | ✓ |
| Lung cancer: Worsening cough, shortness of breath, chest pain, loss of appetite, coughing up blood, fatigue, unexplained weight loss | | | ✓ |
| Lymphoma (cancer of the lymphatic system): painless swelling of lymph node, swollen tonsils, fever, chills, night sweats, feeling tired, itching, unexplained weight loss, loss of appetite, persistent coughing/ difficulty breathing or not being able to breathe, and headache | | | ✓ |
| Peripheral edema: swelling of legs and ankles or the arms and hands | | ✓ | |
| Pulmonary embolism (blood clot in the lung): sharp chest pain, coughing up blood, sudden shortness of breath | | | ✓ |
| Skin cancer: lesions during or after therapy or if existing lesions change appearance | | ✓ | |

If you have a troublesome symptom or side effect that is not listed here or becomes bad enough to interfere with your daily activities, tell your healthcare professional.

Reporting Side Effects

You can report any suspected side effects associated with the use of health products to Health Canada by:

- Visiting the Web page on Adverse Reaction Reporting (<https://www.canada.ca/en/health-canada/services/drugs-health-products/medeffect-canada/adverse-reaction-reporting.html>) for information on how to report online, by mail or by fax; or
- Calling toll-free at 1-866-234-2345.

NOTE: Contact your health professional if you need information about how to manage your side effects. The Canada Vigilance Program does not provide medical advice.

Storage:

Store between 15°C and 30°C.

Keep out of reach and sight of children.

If you want more information about Teva-Tofacitinib:

- Talk to your healthcare professional
- Find the full product monograph that is prepared for healthcare professionals and includes this Patient Medication Information by visiting the Health Canada website: (<https://www.canada.ca/en/health-canada/services/drugs-health-products/drug-products/drug-product-database.html>); or by contacting Teva Canada Limited by:

Phone: 1-800-268-4127 ext. 3;

Email: druginfo@tevacanada.com; or

Fax: 1-416-335-4472

This leaflet was prepared by:

Teva Canada Limited

30 Novopharm Court

Toronto, Ontario

M1B 2K9

Canada

www.tevacanada.com

Last Authorized: December 29, 2023.