

PRODUCT MONOGRAPH

PrCefuroxime for Injection, USP

750 mg / 10 mL vial

1.5 g / 20 mL vial

7.5 g / 100 mL Pharmacy Bulk Vial

Antibiotic

Hikma Canada Limited
5995 Avebury Road, Suite 804
Mississauga, Ontario
L5R 3P9

Date of Initial Authorization:
January 16, 2024

Control No: 280074

PRODUCT MONOGRAPH

PrCefuroxime for Injection, USP

750 mg / 10 mL vial

1.5 g / 20 mL vial

7.5 g / 100 mL Pharmacy Bulk Vial

THERAPEUTIC CLASSIFICATION

Antibiotic

ACTION AND CLINICAL PHARMACOLOGY

In vitro studies demonstrate that the bactericidal action of cefuroxime results from inhibition of bacterial cell wall synthesis by inhibiting the transpeptidase and carboxypeptidase enzymes.

INDICATIONS AND CLINICAL USES

Treatment

Cefuroxime for Injection, USP may be indicated for the treatment of patients with infections caused by susceptible strains of the designated organisms in the following diseases:

Lower Respiratory Tract Infections:

Pneumonia caused by *Streptococcus pneumoniae*, *Haemophilus influenzae* including ampicillin-resistant strains, *Klebsiella* species, *Staphylococcus aureus* including ampicillin-resistant (but not methicillin-resistant) strains, *Streptococcus pyogenes*, and *Escherichia coli*.

Urinary Tract Infections:

Caused by *Escherichia coli*, and *Klebsiella* species.

Soft Tissue Infections:

Caused by *Staphylococcus aureus*, including ampicillin-resistant (but not methicillin-resistant) strains, *Streptococcus pyogenes*, *Escherichia coli*, and *Klebsiella* species.

Meningitis:

Caused by *Staphylococcus aureus* including ampicillin-resistant (but not methicillin-resistant) strains, *Streptococcus pneumoniae*, *Haemophilus influenzae*, and *Neisseria meningitidis*.

Gonorrhea:

Caused by *Neisseria gonorrhoeae* including ampicillin-resistant strains.

Bone and Joint Infections:

Caused by *Staphylococcus aureus* (penicillinase and non-penicillinase producing strains).

To reduce the development of drug-resistant bacteria and maintain the effectiveness of Cefuroxime for Injection, USP and other antibacterial drugs, Cefuroxime for Injection, USP should be used only to treat infections that are proven or strongly suspected to be caused by susceptible bacteria. When culture and susceptibility information are available, they should be considered in selecting or modifying antibacterial therapy. In the absence of such data, local epidemiology and susceptibility patterns may contribute to the empiric selection of therapy.

Specimens for bacteriologic culture should be obtained prior to therapy in order to identify the causative organisms and to determine their susceptibility to cefuroxime. Therapy may be instituted before results of susceptibility testing are known. However, modification of the treatment may be required once these results become available.

Prevention

The preoperative prophylactic administration of Cefuroxime for Injection, USP may prevent the growth of susceptible disease-causing bacteria and thereby may decrease the incidence of certain postoperative infections:

- in patients undergoing surgical procedures (e.g., vaginal hysterectomy) that are classified as clean contaminated or potentially contaminated;
- in patients undergoing open heart surgery in whom infections at the operative site would present a serious risk.

If signs of infection occur postoperatively, culture specimens should be obtained for identification of the causative organism and appropriate antimicrobial therapy should be instituted.

CONTRAINDICATIONS

Cefuroxime for Injection, USP is contraindicated for patients who have shown Type I hypersensitivity to cefuroxime or to the cephalosporin group of antibiotics.

WARNINGS

Before therapy with Cefuroxime for Injection, USP is instituted, careful enquiry should be made to determine whether the patient has had previous hypersensitivity reactions to cefuroxime, cephalosporins, penicillins, or other drugs. Cefuroxime for Injection, USP should be

administered with caution to any patient who has shown some form of allergy, particularly to drugs. There is some clinical and laboratory evidence of partial cross-allergenicity of the cephalosporins and penicillins. If an allergic reaction to Cefuroxime for Injection, USP occurs, treatment should be discontinued and standard agents (e.g., epinephrine, antihistamines, corticosteroids) administered as required.

Pseudomembranous colitis has been reported to be associated with cefuroxime therapy (and other broad-spectrum antibiotics). Therefore, it is important to consider its diagnosis in patients administered Cefuroxime for Injection, USP who develop diarrhea. Treatment with broad-spectrum antibiotics, including cefuroxime, changes the normal flora of the colon and may permit overgrowth of *Clostridia*. Studies indicate that a toxin produced by *Clostridium difficile* is one primary cause of antibiotic-associated colitis. Mild cases of colitis may respond to drug discontinuance alone. Moderate to severe cases should be managed with fluid, electrolyte, and protein supplementation as needed. When the colitis is not relieved by discontinuance of Cefuroxime for Injection, USP administration or when it is severe, consideration should be given to the administration of vancomycin or other suitable therapy. Other possible causes of colitis should also be considered.

Severe Cutaneous Adverse Reactions

Severe cutaneous adverse reactions (SCAR) such as acute generalized exanthematous pustulosis (AGEP), drug reaction with eosinophilia and systemic symptoms (DRESS), Stevens-Johnson syndrome (SJS), and toxic epidermal necrolysis (FEN) have been reported in association with beta-lactam treatment. When SCAR is suspected, Cefuroxime for Injection, USP should be discontinued and appropriate therapy and/or measures should be taken.

Susceptibility/Resistance

Development of Drug Resistant Bacteria

Prescribing Cefuroxime for Injection, USP in the absence of a proven or strongly suspected bacterial infection is unlikely to provide benefit to the patient and risks the development of drug-resistant bacteria.

PRECAUTIONS

Cefuroxime for Injection, USP should be administered with caution to individuals with a history of gastrointestinal disease, particularly colitis.

Patients with markedly impaired renal function (i.e., creatinine clearance of 20 mL / min / 1.73 m² or less) should be placed on the special dosage schedule for cefuroxime recommended under **DOSAGE AND ADMINISTRATION**. Normal dosages in these individuals are likely to produce excessive serum concentrations of cefuroxime.

The concomitant administration of aminoglycosides and some cephalosporins has caused nephrotoxicity. Although transient elevations of BUN and serum creatinine have been seen in clinical studies, there is no evidence that cefuroxime when administered alone, is significantly nephrotoxic.

Studies suggest that the concurrent use of potent diuretics, such as furosemide and ethacrynic acid, may increase the risk of renal toxicity with cephalosporins.

Prolonged treatment with cefuroxime may result in the overgrowth of nonsusceptible organisms, including species originally sensitive to the drug. Repeated evaluation of the patient's condition is required. If superinfection occurs during therapy, appropriate measures should be taken. Should an organism become resistant during antibiotic therapy, another antibiotic should be substituted.

As with other therapeutic regimens used in the treatment of meningitis, hearing loss has been reported in a few pediatric patients treated with cefuroxime. Persistence of positive CSF cultures of *Haemophilus influenzae* at 18 - 36 hours has been noted with cefuroxime.

Pregnancy

The safety of Cefuroxime for Injection, USP in pregnancy has not been established. The use of cefuroxime in pregnant women requires that the potential benefit from the drug be weighed against the possible risk to the mother and fetus. Animal studies have shown cefuroxime to affect bone calcification in the fetus and to show maternal toxicity in the rabbit.

Nursing Mothers

Cefuroxime is excreted in human milk in low concentrations (0.5 mg / L). The clinical significance of this is unknown; therefore, caution should be exercised when Cefuroxime for Injection, USP is administered to a nursing mother.

Elderly Patients

The elimination of cefuroxime may be decreased due to impairment of renal function.

Drug-Laboratory Test Interactions

Cefuroxime may interfere with Benedict's and Fehling's tests for glycosuria. It may cause false-negative reactions in the ferricyanide test, and therefore it is recommended that either the glucose oxidase or hexokinase methods be used to determine blood / plasma glucose levels in patients receiving cefuroxime. Cefuroxime does not interfere with the assay of serum and urine creatinine by the alkaline picrate method.

ADVERSE REACTIONS

The following reactions have been observed during treatment with Cefuroxime for Injection, USP.

Hypersensitivity

Rash, and eosinophilia. Anaphylaxis, urticaria, pruritus and drug fever have also been observed with cephalosporin treatment. Like other cephalosporins, there have been rare reports of erythema multiforme, Stevens-Johnson syndrome and toxic epidermal necrolysis (exanthematic necrolysis).

Local reactions

Thrombophlebitis, stiffness at the site of injection, and inflammatory reactions at the site of injection; some degree of pain, after intramuscular injections when using water as diluent, has been observed.

Blood

Increased erythrocyte sedimentation rate and decreased hemoglobin; eosinophilia, leukopenia and neutropenia; some patients developed a positive direct Coombs' test.

Renal

Increases in BUN and serum creatinine.

Hepatic

Transient increases in serum bilirubin, transaminases and alkaline phosphatase.

Others

Drowsiness, loose stools, faint feeling, sweating, palpitations and *Candida* intertrigo.

SYMPTOMS AND TREATMENT OF OVERDOSAGE

Overdosage of cephalosporins can cause cerebral irritation leading to convulsions. Other than general supportive treatment, no specific antidote is known. Excessive serum levels of cefuroxime can be reduced by dialysis. For treatment of hypersensitivity reactions, see **WARNINGS**.

For management of a suspected drug overdose, contact your regional poison control centre.

DOSAGE AND ADMINISTRATION

Dosage

Cefuroxime for Injection, USP (sterile cefuroxime sodium) may be administered intravenously after reconstitution.

Treatment:

Dosage and route of administration should be determined by severity of infection, susceptibility

of the causative organism(s), and condition of the patient. The intravenous route is preferable for patients with severe or life-threatening infections.

The usual duration of treatment is 5 to 14 days. For β -hemolytic streptococcal infections, therapy should be continued for at least 10 days.

Adults

For most infections, the usual recommended dosage is 750 mg every 8 hours (2.25 g / day), administered intravenously. For severe or life-threatening infections, and for Gram-negative infections of the lower respiratory tract, a dosage of 1.5 g intravenous every 8 hours (4.5 g / day) is recommended.

For treatment of bacterial meningitis, a dosage of 3 g intravenous every 8 hours (9 g / day) should be used.

For bone and joint infections, a dosage of 1.5 g intravenous every 8 hours (4.5 g / day) is recommended. Surgical intervention should be performed when indicated as an adjunct to cefuroxime therapy. A course of oral antibiotics should be administered when appropriate following the completion of parenteral administration of Cefuroxime for Injection, USP.

Infants and Children (1 month to 12 years)

The usual dosage range is 30 to 100 mg / kg / day in 3 or 4 equally divided doses. A dose of 60 mg / kg / day is appropriate for most infections.

In cases of bacterial meningitis*, a dosage of 200 to 240 mg / kg / day intravenous in 3 or 4 equally divided doses should be used.

For bone and joint infections, a dosage of 70 to 150 mg / kg / day administered intravenously every 8 hours is recommended. In clinical trials, a course of oral antibiotics was administered to children following the completion of parenteral administration of Cefuroxime for Injection, USP.

Doses in excess of the maximum adult dose should not be used in infants and children.

Neonates (up to 1 month)

In the first few weeks of life, the serum half-life of cefuroxime can be 3 to 5 times that in adults. Infections in neonates should be treated with dosages in the range of 30 to 100 mg / kg / day in 2 or 3 equally divided doses.

For bacterial meningitis,* a dosage of 100 mg / kg / day intravenous in 2 or 3 equally divided doses should be used.

*Delayed sterilization of cerebral spinal fluid has been reported in a few children treated with cefuroxime for bacterial meningitis. Hearing impairment has occasionally occurred as a complication of meningitis in children treated with cefuroxime.

Prevention:

Clean contaminated or potentially contaminated surgical procedures.

The recommended dose is 1.5 g of Cefuroxime for Injection, USP administered intravenously just prior to surgery.

This may be supplemented with 750 mg at 8 and 16 hours when surgery is prolonged.

In general, prophylactic administration is normally not required after the end of surgical procedures, however, intraoperative administrations should be considered if the surgical procedure is lengthy.

In many surgical procedures, continuing prophylactic administration of any antibiotic does not appear to be related with a decreased incidence of subsequent infection, but will increase the possibility of adverse reactions and the development of bacterial resistance.

Open Heart Surgery:

The recommended dosage is 1.5 g of Cefuroxime for Injection, USP administered intravenously at the induction of anesthesia and every 12 hours thereafter for 48 hours.

Dosage in Patients with Impaired Renal Function:

For patients with markedly impaired renal function, a reduced dosage of Cefuroxime for Injection, USP must be used. For adult patients with moderate infections, dosage adjustment may be made according to the guidelines listed in **Table 1**.

Table 1 – Dosage Adjustment for Adults with Renal Insufficiency			
Creatinine Clearance		Unit Dose	Dosing Frequency
mL / min / 1.73 m²	mL / s / 1.73 m²		
> 20	> 0.33	750 mg – 1.5 g	q8h
10 – 20	0.17 – 0.33	750 mg	q12h
< 10	< 0.17	750 mg	q24h

For adults with severe infections who require doses higher than those recommended in **Table 1**, serum levels of cefuroxime should be monitored and dosage adjusted accordingly.

Studies in children with renal impairment are not sufficient to recommend specific dosages. If it is necessary to administer Cefuroxime for Injection, USP to a child with such impairment, consideration should be given to modifying the frequency of drug administration consistent with the recommendations for adults with renal impairment as indicated in **Table 1**.

When only serum creatinine levels are known, the following formulas may be used to estimate creatinine clearance. The serum creatinine must represent a steady state of renal function.

Males:

$$\text{Creatinine clearance (mL / min)} = \frac{\text{weight (kg)} \times (140 - \text{age})}{72 \times \text{serum creatinine (mg / dL)}}$$

OR

$$\text{Creatinine clearance (mL / s)} = \frac{\text{weight (kg)} \times (140 - \text{age})}{49 \times \text{serum creatinine (mcmol / L)}}$$

Females: 0.85 x male value

For patients on hemodialysis, a further 750 mg dose of Cefuroxime for Injection, USP should be administered at the end of each dialysis treatment.

Administration

Intravenous:

Cefuroxime for Injection, USP may be administered intravenously either by a bolus injection or by a short intravenous infusion over a period of approximately 30 minutes.

For continuous intravenous infusions, a solution of Cefuroxime for Injection, USP (1.5 g dissolved in 16 mL of Water for Injection) may be added to a suitable bottle containing an appropriate intravenous infusion fluid in the amount calculated to give the desired antibiotic dose.


PHARMACEUTICAL INFORMATION

Drug Substance

Proper Name: Sterile cefuroxime sodium

Chemical Name: 5-Thia-1-azabicyclo [4.2.0] oct-2-ene-2 carboxylic acid, 3-[[[(aminocarbonyl) oxy]methyl]-7-[[2-furanyl(methoxy-imino) acetyl]amino-8-oxo-, monosodium salt [6R-[6 α , 7 β (Z)]]

Structural Formula:



Molecular Formula: C₁₆H₁₅N₄NaO₈S

Molecular Weight: 446.4 g / mol

Description: Cefuroxime sodium is a white to faintly yellow crystalline powder, soluble in water (15% w/v at 25 °C), sparingly soluble in ethanol and insoluble in chloroform, toluene, ether, ethyl acetate and acetone.

Composition: Cefuroxime for Injection, USP vials contain cefuroxime sodium (expressed in terms of free acid). Freshly prepared solutions of cefuroxime are yellowish in colour, with some variations in intensity. The pH of freshly constituted solutions range from 6.0 to 8.5.

RECONSTITUTION

For Intravenous Use

Reconstitute with Sterile Water for Injection.

Reconstitution Table			
Vial size	Diluent to be added to vial	Volume to be withdrawn	Approximate Cefuroxime concentration
750 mg / 10 mL vial	8.0 mL	Total	90 mg/mL
1.5 g / 20 mL vial	16.0 mL	Total	90 mg/mL
Shake well until dissolved.			

The reconstituted solution may be further diluted with Sodium Chloride Injection 0.9% w/v, or 5% w/v Dextrose Injection. For short intravenous infusion, 1.5 g of Cefuroxime for Injection, USP is dissolved in 49 mL of Sterile Water for Injection, resulting in an approximate volume of 50 mL i.e., 30 mg / mL.

7.5 g Pharmacy Bulk Vial:

THE AVAILABILITY OF THE PHARMACY BULK VIAL IS RESTRICTED TO HOSPITALS WITH A RECOGNIZED INTRAVENOUS ADMIXTURE PROGRAM.

Cefuroxime for Injection, USP does not contain any preservatives. The Pharmacy Bulk Vial is intended for multiple dispensing for intravenous use only, employing a single puncture. Reconstitute with 77 mL Sterile Water for Injection.

Reconstitution Table			
Vial Size	Diluent to be added to vial	Volume to be withdrawn	Approximate Cefuroxime concentration
7.5 g / 100 mL	77 mL	Amount needed*	95 mg / mL
*8 mL of solution contains 750 mg of cefuroxime; 16 mL of solution contains 1.5 g of cefuroxime. Shake well until dissolved.			

Following reconstitution with Sterile Water for Injection, the solution should be dispensed for further dilution within four hours. Any unused portion of the reconstituted solution should be discarded.

As with all parenteral drug products, intravenous admixtures should be inspected visually for clarity, particulate matter, precipitate, discoloration and leakage prior to administration, whenever solution and container permit. Solutions showing haziness, particulate matter, precipitate, discoloration or leakage should not be used.

STABILITY AND STORAGE RECOMMENDATIONS

Storage

Cefuroxime for Injection, USP in the dry state should be stored between 15°C and 30°C and protected from light.

Reconstituted solution for intravenous injection should be used within 4 hours if kept between 15°C and 25°C or 48 hours if stored under refrigeration.

The further diluted solutions (1 - 30 mg / mL) for intravenous infusion should be used immediately after dilution or stored for up to 36 hours under refrigeration in the dark. Some increase in colour intensity may occur on storage.

NOTE: The pH of 2.74% w/v Sodium Bicarbonate Injection considerably affects the colour of the solution; therefore, this solution is not recommended for the dilution of Cefuroxime for Injection, USP. However, if needed, for patients receiving Sodium Bicarbonate Injection by infusion, the cefuroxime dose may be introduced into the tube of the set.

Incompatibility

Cefuroxime for Injection, USP should not be mixed in the syringe with aminoglycoside antibiotics (e.g., gentamicin sulfate, tobramycin sulfate, amikacin sulfate) due to potential interaction.

AVAILABILITY OF DOSAGE FORMS

Cefuroxime for Injection, USP (sterile cefuroxime sodium) is available as follows:

10 mL vials containing cefuroxime sodium powder equivalent to 750 mg of cefuroxime in packages of 25 vials.

20 mL vials containing cefuroxime sodium powder equivalent to 1.5 g of cefuroxime in packages of 25 vials.

100 mL Pharmacy Bulk Vials containing cefuroxime sodium powder equivalent to 7.5 g of cefuroxime, in packages of 10 vials.

MICROBIOLOGY

Cefuroxime has been shown to be active against the following organisms *in vitro*:

Gram-positive: *Streptococcus pyogenes*, *S. viridans* and *S. pneumoniae*. (Most strains of *Streptococcus faecalis* are resistant.) *Staphylococcus aureus*, both penicillin-sensitive and beta-lactamase-producing. (Some strains of methicillin-resistant *Staphylococci* have been found to be resistant to cefuroxime.) *Clostridia*.

Gram-negative: *Escherichia coli* (including beta-lactamase-producing strains), *Klebsiella*, *Enterobacter*, *Haemophilus influenzae*, *Proteus mirabilis*, *Salmonella*, *Shigella* spp., *Neisseria gonorrhoeae* and *N. meningitidis*.

The following organisms are not susceptible to cefuroxime: *Clostridium difficile*, *Pseudomonas* spp., *Campylobacter* spp., *Acinetobacter calcoaceticus*, Methicillin resistant strains of *Staphylococcus aureus*, Methicillin resistant strains of *Staphylococcus epidermidis* and *Legionella* spp. Some strains of the following genera are not susceptible to cefuroxime: *Streptococcus faecalis*, *Morganella morganii*, *Proteus vulgaris*, *Enterobacter* spp. *Citrobacter* spp., *Serratia* spp. and *Bacteroides fragilis*.

The minimum inhibitory concentrations against various organisms are shown in Tables 2, 3 and 4.

Table 2 – <i>In vitro</i> activity of cefuroxime against Gram-positive bacteria													
Organism	No. of strains	Inoculum size (CFU / mL)	Cumulative percent of strains sensitive at indicated conc. (mcg / mL)										
			< 0.005	0.01	0.03	0.06	0.12	0.25	0.5	1.0	2.0	4.0	> 4.0
<i>Staphylococcus aureus</i>													
<i>penicillin-sensitive</i>	12	10 ³								58	100		
<i>penicillin-resistant</i>	28	10 ³								14	68	100	
<i>methicillin-resistant</i>	40	10 ³								5	25	33	
<i>coagulase-negative</i>	39	10 ³				3	10	28	54	79	85		
<i>alpha- and non-hemolytic Streptococci</i>	20	10 ³	15	35	55	70	75	85	100				
<i>beta-hemolytic Streptococci</i>	40	10 ³	8	50	80	95	98	100					
<i>Streptococcus pneumoniae</i>	19	10 ³	53	100									
<i>Clostridium spp.</i>	7	10 ³				13		26			86		100

Table 3 – <i>In vitro</i> activity of cefuroxime against Gram-negative bacteria									
Organism	No. of strains	Inoculum size (CFU / mL)	Cumulative percent of strains sensitive at indicated conc. (mcg / mL)						
			< 0.125	0.25 - 0.5	1 - 4	8 - 16	32 - 62	> 125	
<i>E. coli</i>	129	10 ⁵		5	89	98	100	100	
<i>Klebsiella spp.</i>	73	10 ⁵			51	81	99	100	
<i>Acinetobacter</i>	6	10 ⁵		17	67	83	100		
<i>Enterobacter spp.</i>	138	10 ⁵			22	88	94	100	
<i>Serratia spp.</i>	8	10 ⁵				13	25	75	
<i>Proteus mirabilis</i>	27	10 ⁵			89	96	100		
<i>Proteus rettgeri</i>	4	10 ⁵	25			50	75	100	
<i>Proteus vulgaris</i>	21	10 ⁵				29	86	100	

Table 3 – <i>In vitro</i> activity of cefuroxime against Gram-negative bacteria								
Organism	No. of strains	Inoculum size (CFU / mL)	Cumulative percent of strains sensitive at indicated conc. (mcg / mL)					
			< 0.125	0.25 - 0.5	1 - 4	8 - 16	32 - 62	> 125
<i>Morganella morganii</i>	9	10 ⁵			11	44	78	100
<i>Salmonella spp.</i>	40	10 ⁵			95	98	100	
<i>Shigella spp.</i>	10	10 ⁵			90		100	
<i>B. fragilis</i>	16	10 ⁵			6	31	100	
<i>H. influenzae</i>								
<i>ampicillin-sensitive</i>	16	10 ⁵		100				
<i>ampicillin-resistant</i>	15	10 ⁵		100				

Table 4 – <i>In vitro</i> activity of cefuroxime against gonococci						
Organism	No. of strains	Inoculum size (CFU / mL)	Cumulative percent of strains sensitive at indicated conc. (mcg / mL)			
			< 0.03	0.06 - 0.25	0.5 - 2.0	> 2.0
<i>N. gonorrhoeae</i>						
<i>beta-lactamase positive</i>	110	10 ³	72	94	100	
<i>beta-lactamase negative</i>	752	10 ³	60	92	97	100

Although cefuroxime is resistant to hydrolysis by most beta-lactamases, these enzymes from certain species (*Bacteroides fragilis*, *Enterobacter* and indole-positive *Proteus* spp) have been shown to produce hydrolysis. Table 5 indicates the degree of resistance of cefuroxime to beta-lactamase inactivation.

Source of enzyme	Enzyme Class	µg of cefuroxime hydrolyzed / minute
<i>Escherichia coli</i> (R ⁺ tem)	III	< 1
<i>E. coli</i> (R ⁺ GN238)	V	4.5
<i>E. coli</i> D31	I	< 1
<i>Proteus mirabilis</i>	III	< 1
<i>Klebsiella aerogenes</i> K1	IV	54
<i>Enterobacter cloacae</i> P99	I	< 1
<i>Proteus vulgaris</i>	I	< 1
<i>Bacteroides fragilis</i> 1600	I	112
<i>Pseudomonas aeruginosa</i> 1822	I	< 1
<i>Bacillus cereus</i> 659 / H9		72
<i>Staphylococcus aureus</i> PCI*		< 1

*Activity is expressed as micrograms hydrolyzed per hour

Mice, rats and rabbits were inoculated intraperitoneally with a variety of Gram-positive and Gram-negative microbes (such as *Staphylococcus aureus*, *E. coli*, *Proteus mirabilis*, *Klebsiella*). Animals administered cefuroxime intramuscularly, showed resistance to all of these test organisms at doses from 1 to 32 mg / kg. Doses of cefuroxime ranging from 35 to 133 mg / kg / dose were needed for protection against infections from two strains of *Proteus vulgaris* and one strain of beta-lactamase-producing *E. coli*.

Susceptibility Testing

The results of susceptibility testing, by either disk-diffusion or tube-dilution techniques, should be interpreted according to the criteria shown in **Table 6**.

	Zone diameter (30 mcg cefuroxime disk)	Approximate MIC correlate
SUSCEPTIBLE (susceptible to the usual doses)	≥ 18 mm	≤ 8 mcg / mL
INTERMEDIATE (moderately susceptible)*	15 – 17 mm	16 mcg / mL
RESISTANT	≤ 14 mm	≥ 32 mcg / mL
CONTROL STRAINS		
<i>S. aureus</i> ATCC 25923	27 – 35 mm	0.5 – 2 mcg / mL
<i>E. coli</i> ATCC 25922	20 – 26 mm	2 – 8 mcg / mL

- Organisms that produce zones of 15 to 17 mm may be susceptible if the infection is confined to tissues and fluids (e.g., urine) in which high antibiotic concentrations are attained.

Only cefuroxime disks should be used, since cefuroxime has been shown by *in vitro* tests to be active against certain strains found resistant when other beta-lactamase disks are used.

PHARMACOLOGY

Animal

Subcutaneous administration of cefuroxime to mice at a dose of 4 g / kg, had no significant effect on the central nervous system, on spontaneous locomotor activity or motor coordination and no anticonvulsant, analgesic, tranquilizing or antidepressant properties. Doses of up to and including 300 mg / kg cefuroxime given intravenously to cats and dogs produced no pharmacodynamic effects on the cardiovascular or respiratory systems other than small variations in blood pressure and heart rate in the cat, which were not dose-related. However, doses of 1 and 3 g / kg produced an initial transitory tachycardia, a decrease in blood pressure followed by bradycardia and an increase in blood pressure. In neither species did cefuroxime affect the responses of the cardiovascular system to intravenously-injected neurohumoral agents. Ganglionic transmission was not affected in the cat.

Cefuroxime had no effect on isolated smooth muscle preparations at a concentration in the bathing fluid of 10^{-5} M. Only minor increases in contractile force and rate of contraction of the isolated rabbit heart (Langendorff preparation) were observed when the concentration in the perfusing fluid was increased to 10^{-2} M. A 30% solution of cefuroxime in 0.9% saline had no local anaesthetic activity nor any irritant effect to the cornea of the rabbit eye.

Cefuroxime had no significant effect on the cortical EEG of rats.

Human

Cefuroxime is poorly absorbed when given orally; following a 1 g dose, serum concentrations of less than 1.2 mcg / mL were noted and only between 1 and 1.3% of the administered dose was excreted in the urine. Cefuroxime, therefore, is used by the intravenous route.


Intravenous Administration:

Administration of cefuroxime sodium 750 mg and 1.5 g resulted in blood levels of 73 mcg / mL and 151 mcg / mL respectively (**Figure 1**), 5 minutes after the initiation of the injection.

Intravenous infusion of 750 mg cefuroxime over a 30-minute period resulted in a serum level of 51 mcg / mL at the end of the infusion. Intravenous administration of 1.5 g over a 20-minute period, produced a drug concentration of 146 mcg / mL at the end of the infusion.

Following intravenous administration, more than 95% of cefuroxime was excreted unchanged by the kidneys (**Table 7**) with excretion evenly divided between glomerular filtration and tubular secretion. The half-life of cefuroxime following intravenous injection was approximately 65 minutes.

Figure 1: Serum cefuroxime levels following intravenous injection



Route and dose (g)	Mean % urinary recovery at hours after injection							Total mean ± SD
	0- 1	1 - 2	2 - 3	3 - 4	4 - 6	6 - 12	12 - 24	
IV								
0.25	60.2	23.2	14.2	6.1	6.3	3.6	0.5	114.1 ± 6.1
0.5	41.3	23.6	13.1	6.9	5.6	4.0	0.6	95.1 ± 4.4
1.0	53.6	21.5	12.0	5.2	4.1	2.5	0.2	99.1 ± 0.2

The effect of probenecid on the pharmacokinetics of cefuroxime is shown in **Table 8**.

Pharmacokinetic variable	With probenecid	Without probenecid	Percentage change
Peak serum concentration (mcg / mL)	29.4	22.7	+ 30
Ultimate serum half-life (min)	101	76.6	+ 32
Area under curve (mcg / mL/ h)	94.4	56.8	+ 56
Apparent volume of distribution (L / 1.73m ²)	11.7	14.8	- 20
Urinary recovery: 0 - 2 h (%)	47	60.4	- 22
0 - 24 h	95.6	100.2	- 5
Renal clearance (mL / min / 1.73 m ²)	79.6	133.8	- 40
Cefuroxime / Creatinine clearance ratio	0.74	1.25	- 40

Patients With Renal Impairment:

The effect of various degrees of renal impairment on the pharmacokinetics of cefuroxime is shown in **Table 9**.

Patient No.	Mean Creatinine Clearance (mL / min)	Serum Concentration (mcg / mL)		Serum Half-Life (h)	Urinary Concentrations (mcg / mL)			
		Peak	Trough		0 - 2h	2 - 4h	4 - 6h	6 - 10h
1	21.0 (± 1.8)	101.0 – 62.4	9.2 – 8.0	4.3 (± 0.08)	150	177	145	135
2	23.0 (± 2.6)	80.3 – 72.6	9.7 – 8.0	4.2 (± 0.21)	180	225	102	85
3	12.1 – 17.8 (no mean available)	65.7 – 55.4	7.1 – 1.1	6.5 (± 0.37)	100	99	63	113
4	10.0 (± 1.4)	90.0 – 75.6	15.1 – 10.6	8.4 (± 0.41)	57	59	45	79
5	5.0 (± 2.0)	125.0 – 52.2	28.6 – 24.2	22.3 (± 2.03)	41	25	17	37

Fluid and Tissue Levels:

Cefuroxime was detected in certain fluids and tissues as indicated in **Table 10**.

Site	Dose given (mg)	Route	Concentration (mcg / mL)
Sputum	750 t.i.d. for 2 days	Intramuscular	2.0
	1500 t.i.d. for 4 days	Intramuscular	7.8
Bone	750 t.i.d. for 4 days	Intramuscular	3.9*
	1500 t.i.d. for 4 days	Intramuscular	13.5*
Skin blister	750 single	Intramuscular	9.4
Bile	750 single	Intramuscular	8.6
	1500 single	Intramuscular	22.0
Aqueous humor	1500 single	Intravenous	1.6

*mc / g

An intravenous dose of 750 mg of cefuroxime produced biliary concentrations which varied considerably between 1.3 and 26 mcg / mL. Biliary levels appear to be lowest in patients with a non-functioning gallbladder.

Following a 750 mg intramuscular dose to 6 women in labour, average concentrations of cefuroxime in amniotic fluid (18.6 mcg / mL) were comparable to those in maternal serum; average peak maternal serum concentrations of 19.2 mcg / mL were achieved after 1.2 hours, while in umbilical cord blood, the average peaks were one-third of those in the mothers.

Protein Binding:

The extent of cefuroxime bound to protein in the serum was approximately 33%.

TOXICOLOGY

Acute Toxicity

Species	No. of animals	Dose (g / kg)	Route	Deaths
Mouse	10	11	Intravenous	5
Rat	6	4	Intravenous	3
Cat	4	2	Intramuscular	2
Monkey	4	2	Intramuscular	2
Dog	4	2	Intramuscular	2

Symptoms of toxicity immediately following intravenous administration in the rat included collapse and tachypnea. During the follow-up observation period (7 days), soft feces and a slight

loss of body weight were noted in rats, while monkeys displayed diarrhea, accompanied by weight loss.

Subacute Toxicity

Rat:

When rats were treated for a month with daily subcutaneous doses of 100 mg / kg of cefuroxime, the serum potassium was increased on day 34. With doses of 200 mg / kg / day, peripheral erythrocyte values were somewhat decreased in males and with 400 mg / kg / day doses in females. Daily doses of 800 mg / kg produced moderate reactions at the injection site, forming subcutaneous lumps and occasionally ulcers. The ulcerations normally resolved within 10 days. There was also evidence of mild colitis. Rats were administered subcutaneous doses of 1.25, 2.5 and 5.0 mg / kg / day of cefuroxime for 14 days. All of the animals displayed symptoms of extreme discomfort during and immediately after injection. During autopsy, necrotic patches were noted at the injection sites of the rats administered the highest dose. All rats had watery feces, increased leukocyte count, and a dose-related reduction in hemoglobin concentration.

Rats treated with cefuroxime 50, 100, 200 and 400 mg / kg / day intravenously, for one month resulted in increased packed cell volume in all groups, increased urine output in the 200 mg / kg group and embolic reactions in many lungs, in both control and drug-treated animals. At doses of 100 and 400 mg / kg / day, a small but statistically significant reduction in spleen weights was seen.

Dog:

Administration of cefuroxime at daily intramuscular doses of 60, 180 and 540 mg / kg for 11 days produced increased kidney and liver weights. In two male dogs, this was 1.5 times the weight of the controls.

Monkey:

Cefuroxime administered intramuscularly for 29 days at doses of 150 and 450 mg / kg / day, produced a moderate reduction in erythrocytes, leukocytosis with neutrophilia, eosinophilia and soft stools.

A slight-to-moderate dose-related inflammatory reaction around the subcutaneous and intramuscular injection sites were seen in all subacute tests. Hemorrhage at the injection sites was sometimes noted, occurring more frequently at high doses.

Chronic Toxicity

Rat - Three-month, subcutaneous study:

In rats administered cefuroxime at 100, 300 and 900 mg / kg / day, a dose- and duration-related

mild-to-marked subcutaneous reaction with hemorrhage at injection sites was seen at all 3 dose levels. Slight decrease in erythrocytes with mild reticulocytosis and slight decrease of serum calcium were seen at both 300 mg and 900 mg / kg dose levels. Increased prothrombin time was noted in males dosed at 300 mg / kg / day, and in both sexes at 900 mg / kg / day.

At 900 mg / kg / day an increased total leukocyte count, decreased serum albumin and gamma-globulin, increased serum potassium (females on day 56), marginally increased blood glucose (females on day 56) and decreased alkaline phosphatase (males on day 28) were seen. There was an increase in relative weights of the liver, the kidney and the spleen in all females. Increased excretion of electrolytes and increased urinary volume in both sexes were noted at the high dose level after 77 days.

Rat - Six-month, subcutaneous study:

Rats were studied at cefuroxime doses of 50, 150 or 450 mg / kg / day. No mortality was observed. Dilatation of the lymphatics and enlargement of the spleen were seen at the higher doses. There was a statistically significant decrease in packed cell volume and hemoglobin, accompanied by reticulocytosis of a similar degree at 150 mg / kg / day. A decrease in serum alanine transaminase activity in both sexes as well as of pituitary weights in females were seen at the highest dose. A significant increase in serum cholesterol was observed in females at all dose levels while serum calcium levels were marginally decreased in males at high dose levels.

An increase in excretion of sodium and potassium was seen, however, statistically significant changes were observed in the excretion of potassium in the high dose male animals only. Other abnormalities included hyaline granular casts and erythrocytes in the urine in addition to increased urinary protein content. The incidence and severity of these changes were greatest at high dose levels.

The mean kidney weights (absolute and relative) in both sexes were increased 10 to 15% at high doses.

Dog - Six-month, toxicity study:

Dogs were administered cefuroxime for 6 months at doses of 50 mg / kg / day intramuscularly, and at doses of 150 and 450 mg / kg / day subcutaneously. Dose-related subcutaneous hemorrhage was observed at the injection site with higher doses of cefuroxime. In the group treated with the highest dose, hypochromia and increased serum iron binding capacity were seen. Serum triglyceride concentrations were increased after 20 weeks in animals given 150 and 450 mg / kg cefuroxime. Blood urea nitrogen was decreased and serum potassium was increased in the high dosage group. One dog in the 450 mg group developed Heinz body anaemia after 12 weeks, however, the causative agent was not identified.

Nephrotoxicity Studies

Mouse:

Mice were given single subcutaneous doses of cefuroxime (10 g / kg) alone or in combination with furosemide (50 mg / kg) or furosemide plus glycerol (5.4 mL / kg). Cefuroxime alone produced no nephrotoxicity; together with furosemide, proximal tubular necrosis was observed in two out of nine animals. Administration of furosemide and glycerol resulted in tubular necrosis in 5 of 8 animals but this was not affected by the addition of cefuroxime.

Rat - Single dose study:

Cefuroxime at doses up to 10 g / kg was administered either alone or together with furosemide (100 mg / kg) or furosemide plus glycerol (3.15 mL / kg). Three of 6 animals had proximal tubular necrosis of the inner cortex when given 4 g of cefuroxime alone. The frequency and severity was observed to increase with increasing doses. The frequency of tubular necrosis also increased with increasing doses. The incidence of tubular necrosis was increased when furosemide or furosemide plus glycerol were given concomitantly. Cefuroxime at 1 g / kg increased the severity of the furosemide-glycerol-included necrosis in the outer cortex. Furosemide plus glycerol therapy also decreased the dosage of cefuroxime (to 2 g / kg) needed to cause necrosis of the inner cortex.

Rat - Repeated dose study:

Rats were given cefuroxime at doses ranging from 1 to 5 g / kg / day subcutaneously for 10 days. Histological evidence of tubular necrosis were not seen at 5 g / kg, however, transient increases in urine volume, protein and enzymes, which peaked at days 2 - 3 were noted. A significant decrease in body weight of the animals was observed in the high dose group.

Combination with aminoglycosides:

Rats were administered gentamicin (35 mg / kg) for 10 days. Cefuroxime was administered either concomitantly during the 10 days or as a single dose with the ninth dose of gentamicin. Gentamicin-induced tubular necrosis was not increased by the administration of single doses of cefuroxime up to 6 g / kg / day. Protection of rats against gentamicin-induced nephrotoxicity was seen with multiple cefuroxime doses of up to 4 g / kg, but at cefuroxime doses of 6 g / kg / day, severe tubular necrosis was seen after 4 days of treatment. Similar findings were observed with amikacin and tobramycin.

Teratogenicity Studies

Mouse:

Cefuroxime was administered subcutaneously at doses of 800, 1600, 3200 and 6400 mg / kg / day from day 6 to day 15 of pregnancy. With the exception of the 3200 mg dose, there was a 15

to 21% incidence of bone immaturity at all doses tested as evidenced by a reduction in the calcification of various ossification centres of the offspring. Based on historical controls, the untreated animals had a 7% incidence of bone immaturity.

Rabbit:

Cefuroxime was administered intramuscularly at doses of 50, 100, 200 and 400 mg / kg / day from day 6 to day 18 of pregnancy. Four rabbits given 400 mg / kg, one rabbit given 200 mg / kg and one rabbit given 100 mg / kg / day, died during the experiment. The offspring had an 8, 17, 25, and 10% incidence of bone immaturity and the incidence of bone abnormalities was 8, 21, 0 and 30% at the 50, 100, 200 and 400 mg / kg dose levels, respectively.

Fertility and Reproduction Studies

Male and female mice were administered daily subcutaneous doses of 800, 1600 or 3200 mg / kg of cefuroxime prior to mating (males for 60 days and females for 14 days). The pregnant females were continued on treatment with cefuroxime until the 17th day of pregnancy. A few of their offspring were later mated to produce a second generation. Therapy had no apparent effect on gametogenesis. The fertility of the second generation was also unaffected.

Perinatal and Postnatal Studies

Mouse:

Daily subcutaneous doses of 800, 1600, or 3200 mg / kg of cefuroxime from day 16 of pregnancy until the weaning of the litters, had no effect on gestation, parturition, lactation or the health of the dams or the pups.

Rabbit:

Daily administration of 50, 100 or 200 mg / kg of cefuroxime from day 19 of pregnancy through lactation (at least 50 doses) had no effect on the litters or the development and health of the pups. Treatment related deaths were seen in 10 rabbits before parturition and one died after it had littered. Mortality was dose-related, and although believed to be the result of enteritis, a direct toxic effect could not be ruled out.

BIBLIOGRAPHY

1. Bahaeldin HK, Elbaroudy R, Omran N. The efficacy of cefuroxime in the treatment of bacterial meningitis in infants and children. *Clin Ther* 1983; 5(6):644-650.
2. Broekhuysen J, Deger F, Douchamps J, Freschi E, Mal N, Neve P, Parfait R, Siska G, Winand M. Pharmacokinetic study of cefuroxime in the elderly. *Br J Clin Pharmacol* 1981; 12:801-805.
3. Brogden RN, Heel RC, Speight TM, Avery GS. Cefuroxime: a review of its antibacterial activity, pharmacological properties and therapeutic use. *Drugs* 1979; 17:233-266.
4. Brundtzen RW, Toothaker RD, Neilson OS, Madsen PO, Welling PG, Craig WA. Pharmacokinetics of cefuroxime in normal and impaired renal function: comparison of high-pressure liquid chromatography and microbiological assays. *Antimicrob Ag Chemother* 1981; 19(3):443-449.
5. Bullen BR, Ramsden CH, Kester RC. Cefuroxime levels attained in tissues and wound exudates from severely ischaemic limbs. *Journal of Antimicrobial Chemotherapy* 1981, 7:163-169.
6. Court-Brown CM, Hughes SPF. The role of cefuroxime in total joint replacement. *Research and Clinical Forums*; 5(4):27-36.
7. Daikos GK, Kosmidis J, Stathakis C, Anyfantis A, Plakoutsis T, Papathanassiou B. Bioavailability of cefuroxime in various sites including bile, sputum and bone. *Proc Roy Soc Med* 1977; 70 (9):38-41.
8. Gold B, Rodriguez J. Cefuroxime: mechanisms of action, antimicrobial activity, pharmacokinetics, clinical applications, adverse reactions and therapeutic indications. *Pharmacother* 1983; 3(2):82-100.
9. Gower PE, Kennedy MRK, Dash CH. The effect of renal failure and dialysis on the pharmacokinetics of cefuroxime. *Proc Roy Soc Med* 1977; 70(Suppl 9):151-157.
10. Graninger W, Pichler R, Mulac KD, Bohm M, Spitzky KH. Clinical evaluation of the effect of cefuroxime in patients with pyelonephritis and osteomyelitis. *Proc Roy Soc Med* 1977; 70 (9):148-150.
11. Hedstrom SA, Lidgren L, Nilsson-Ehle I. Cefuroxime in acute septic arthritis. *Scand J. Infect Dis* 1984; 16:79-82.
12. Herold HZ, Shmueli G, Butler JK. Cefuroxime in the treatment of bone and joint infections. *Current Therapeutic Research* 1983; 33 (6):892-897.

13. Hofstetter A, Boewering R, Schillin A. Clinical evaluation of cefuroxime in patients with urinary tract infections. *Proc Roy Soc Med* 1977; 70(Suppl 9):158-161.
14. Hughes S, Robertson S, Want S, Darrell J, Kennedy M, Dash CH. Cefuroxime in bone. Cefuroxime Update: Royal Society of Medicine International Congress and Symposium Series No. 38; p.163-171.
15. Hughes SPF, Want S, Darrell JH, Dash CH, Kennedy M. Prophylactic Cefuroxime in total joint replacement. *International Orthopaedics (SICOT)* (1982)6:155-161.
16. Jandial L, Reid TMS, Jandial V. Prophylactic cefuroxime in gynaecological surgery. *Research and Clinical Forums* 1983; 5(4):45-50.
17. Jones RN, Fuchs PC, Gavan TL, Gerlach EH, Barry AL, Thornsberry C. Cefuroxime, a new parenteral cephalosporin: collaborative *in vitro* susceptibility comparison with cephalothin against 5,887 clinical bacterial isolates. *Antimicrob Ag Chemother* 1977; 12(1):47-50.
18. Jung HH, Mischokowsky T. Measurements of cefuroxime concentration in bone in dogs and man. *Arch Orthop Traumat Surg* 1979; 94:25-28.
19. La Greca G, Biasioli S, Chiaromonte S, Fabris A, Feriani M, Pisani E, Ronco C, Xerri L. Pharmacokinetics of intravenous and intraperitoneal cefuroxime during peritoneal dialysis. *Int J Clin Pharmacol Therap Toxicol* 1982; 20(2):92-94.
20. Leigh DA, Marriner J, Nisbet D, Powell HDW, Church JCT, Wise K. Bone concentrations of cefuroxime and cefamandole in the femoral head in 96 patients undergoing total hip replacement surgery. *Journal of Antimicrobial Chemotherapy* 1982, 9:303-311.
21. de Louvois J, Mulhall A, Hurley R. Cefuroxime in the treatment of neonates. *Arch Dis Childh* 1982; 57:59-62.
22. Mullinger BM, House CA. Penetration of cefuroxime and other parenteral cephalosporins into various tissues and body fluids. In Wood C, Rue Y (eds): Cefuroxime Update. Royal Society of Medicine International Congress and Symposium Series #38, pp 153-161. Academic Press, London, 1981.
23. Nelson JD, Bucholz RW, Kusmiesz H, Shelton S. Benefits and risks of sequential parenteral-oral cephalosporin therapy for suppurative bone and joint infections. *Journal of Pediatric Orthopedics* 1982; 2:255-262.
24. Nelson JD. Cefuroxime: a cephalosporin with unique applicability to pediatric practice. *Ped Inf Dis* 1983; 2(5):394-396.

25. Netland A, Muller C, Andrew E. Concentration of cefuroxime in cerebrospinal fluid in patients with bacterial meningitis. *Scand J Inf Dis* 1981; 13:273-275.
26. Neu HC, Fu KP. Cefuroxime, a beta-lactamase-resistant cephalosporin with a broad spectrum of Gram-positive and -negative activity. *Antimicrob Ag Chemother* 1978; 13(4):657-664.
27. Nyman R, Anttolainen I. Experience in the use of cefuroxime in paediatric infections. *Proc Roy Soc Med* 1977; 70(Suppl 9):191-194.
28. Peterson CD, Lake KD, Arom KV, Love KR. Antibiotic prophylaxis in openheart surgery patients: comparison of cefamandole and cefuroxime. *Drug Intelligence and Clinical Pharmacy*, 1987 September; 21:728-732.
29. Pfenninger J, Schaad UB, Lutschg J, Nussbaumer A, Zellweger U. Cefuroxime in bacterial meningitis. *Arch Dis Childh* 1982; 57:539-543.
30. Philipson A, Stiernstedt G. Pharmacokinetics of cefuroxime in pregnancy. *Am J Obstet Gynecol* 1982; 142(7):823-828.
31. Pines A, Raafat H, Mullinger BM. Cefuroxime in severe respiratory infections: a double-blind comparison of two doses. *Br J Dis Chest* 1979; 73:395-397.
32. Price JD, Fluker JL. Dose response evaluation of cefuroxime in the treatment of gonorrhoea in male and female patients. *Proc Roy Soc Med* 1977; 70(Suppl9):125-127.
33. Schaad UB, Krucko J, Pfenninger J. An extended experience with cefuroxime therapy of childhood bacterial meningitis. *Ped Inf Dis* 1984; 3(5):410-416.
34. Sirinavin S, Chiemchanya S, Visudhipan P, Lolekha S. Cefuroxime treatment of bacterial meningitis in infants and children. *Antimicrob Ag Chemother* 1984; 25(2):273-275.
35. Slama TG, Sklar SJ, Misinski J, Fess SW. Randomized comparison of cefamandole, cefazolin, and cefuroxime prophylaxis in open-heart surgery. *Antimicrobial Agents and Chemotherapy*, May 1986; 29(5):744-747.
36. Sutton A, Fleet H, Ng SH, Farthing CP, Kennedy MRK, Cox D: The role of cefuroxime and gentamicin in neonates: a comparative study. In Wood C, Rue Y (eds): *Cefuroxime Update*. Royal Society of Medicine International Congress and Symposium Series #38, pp 153-161. Academic Press, London, 1981.
37. Swedish Study Group: Cefuroxime versus ampicillin and chloramphenicol for the treatment of bacterial meningitis. *Lancet* 1982; 1:295-299.

38. Tartaglione TA, Polk RE. Review of the new second-generation cephalosporins: cefonicid, ceforanide, and cefuroxime. *Drug Intell Clin Pharm* 1985; 19(3):188-198.
39. Thomas M, Browning AR, McFarland RJ. Experience with cefuroxime in biliary disease. *Research and Clinical Forums* 1983; 5(4):85-88.
40. Walstad RA, Nilsen OG, Berg KJ. Pharmacokinetics and clinical effects of cefuroxime in patients with severe renal insufficiency. *Eur. J. Clin. Pharmacology* 24:391-398, 1983.
41. Product Monograph, Cefuroxime for Injection, USP. Fresenius Kabi Canada Ltd. Control No. 233970. April 17, 2020.

READ THIS FOR SAFE AND EFFECTIVE USE OF YOUR MEDICINE

PATIENT MEDICATION INFORMATION

PrCefuroxime for Injection, USP (cefuroxime sodium powder)

Read this carefully before you start taking **Cefuroxime for Injection, USP** and each time you get a refill. This leaflet is a summary and will not tell you everything about this drug. Talk to your healthcare professional about your medical condition and treatment and ask if there is any new information about **Cefuroxime for Injection, USP**.

What is Cefuroxime for Injection, USP used for?

Cefuroxime for Injection, USP is used for the treatment of infections caused by certain bacteria in many different parts of the body.

Cefuroxime for Injection, USP can also be used to prevent infections, before and after surgery.

Antibacterial drugs like Cefuroxime for Injection, USP treat only bacterial infections. They do not treat viral infections.

How does Cefuroxime for Injection, USP work?

Cefuroxime for Injection, USP is an antibiotic, which belongs to a class of drugs called cephalosporins. Cefuroxime for Injection, USP works by killing bacteria that cause infections in the body.

What are the ingredients in Cefuroxime for Injection, USP?

Medicinal ingredients: cefuroxime sodium

Non-medicinal ingredients: none

Cefuroxime for Injection, USP comes in the following dosage forms:

Sterile powder for injection: 750 mg / 10 mL vial, 1.5 g / 20 mL vial, 7.5 g / 100 mL vial

Do not use Cefuroxime for Injection, USP if:

You are allergic to cefuroxime or to cephalosporin antibiotics.

To help avoid side effects and ensure proper use, talk to your healthcare professional before you take Cefuroxime for Injection, USP. Talk about any health conditions or problems you may have, including if you:

- are allergic to or react badly to penicillins or other antibiotics.
- have kidney problems, as your healthcare professional may reduce your dose.
- have a history of bowel disease, particularly colitis.
- are taking other antibiotics called aminoglycosides, such as streptomycin, neomycin, gentamicin and tobramycin.

- are taking “water pills” or diuretics used to lower high blood pressure, such as furosemide and ethacrynic acid.
- are having a urine test for sugar as Cefuroxime for Injection, USP can affect the results.
- are pregnant or planning to become pregnant.
- are breastfeeding or planning to breastfeed. Cefuroxime for Injection, USP is passed into breast milk. Discuss whether you should breastfeed while taking Cefuroxime for Injection, USP with your healthcare professional.

Other warnings you should know about:

Driving and using machines: Cefuroxime for Injection, USP can cause dizziness. Do not drive or operate machinery if you are feeling dizzy.

Tell your healthcare professional about all the medicines you take, including any drugs, vitamins, minerals, natural supplements or alternative medicines.

The following may interact with Cefuroxime for Injection, USP:

- Diuretics or “water pills” used to lower high blood pressure, such as furosemide and ethacrynic acid

How to take Cefuroxime for Injection, USP:

- Cefuroxime for Injection, USP will be given to you by a healthcare professional who will inject it into a vein.
- Although you may feel better early in treatment, Cefuroxime for Injection, USP should be used exactly as directed.
- Misuse or overuse of Cefuroxime for Injection, USP could lead to the growth of bacteria that will not be killed by Cefuroxime for Injection, USP (resistance). This means that Cefuroxime for Injection, USP may not work for you in the future.
- Do not share your medicine.

Usual dose:

Your healthcare professional will decide how much, how often and for how long you will receive Cefuroxime for Injection, USP.

Overdose:

If you think you, or a person you are caring for, have taken too much Cefuroxime for Injection, USP, contact a healthcare professional, hospital emergency department, or regional poison control centre immediately, even if there are no symptoms.

Missed dose:

If you miss an appointment to receive an injection of Cefuroxime for Injection, USP contact your healthcare professional as soon as possible.

What are possible side effects from using Cefuroxime for Injection, USP?

These are not all the possible side effects you may feel when taking Cefuroxime for Injection, USP. If you experience any side effects not listed here, contact your healthcare professional.

Side effects may include:

- Drowsiness, feeling faint
- Loose stool
- Sweating
- Pain and stiffness at the injection site

Cefuroxime for Injection, USP can cause abnormal blood test results. Your healthcare professional will decide when to perform blood tests and interpret the results. Also, tell any healthcare professionals you see before having blood work that you are taking Cefuroxime for Injection, USP as it may affect your results.

Serious side effects and what to do about them			
Symptom / effect	Talk to your healthcare professional		Stop taking drug and get immediate medical help
	Only if severe	In all cases	
RARE			
Yeast infection of the skin: redness, burning and itching in the skin folds, commonly in the groin, under the breasts and in the arm pit		✓	
Pseudomembranous colitis: watery, bloody diarrhea, mucus in the stool, abdominal cramps and pain, fever			✓
VERY RARE			
Allergic Reaction: wheeziness and tightness of chest, swelling of eyelids, face or lips, or develop skin lumps or hives, or a skin rash (red spots), high fever.			✓
Skin Reaction: skin rash, which may blister, and looks like small targets (central dark spots surrounded by a paler area, with a dark ring around the edge), widespread rash with blisters and skin peeling on much of the body, particularly around the mouth, nose, eyes and genitals.			✓

Serious side effects and what to do about them			
Symptom / effect	Talk to your healthcare professional		Stop taking drug and get immediate medical help
	Only if severe	In all cases	
Heart palpitations: fast, slow or irregular heartbeat		✓	
Kidney problems: decreased urination, nausea, vomiting, swelling of extremities, fatigue		✓	
Liver problems: yellowing of the skin or eyes, dark urine, abdominal pain, nausea, vomiting, loss of appetite		✓	
Anemia: fatigue, loss of energy, weakness, shortness of breath		✓	
Low levels of white blood cells: infection, fatigue, fever, aches, pain, flu-like symptoms		✓	
Phlebitis: swelling of a vein near the injection site, with pain, tenderness, redness		✓	
UNKNOWN			
Severe Cutaneous Adverse Reaction (SCAR) (severe skin reactions that may also affect other organs: <ul style="list-style-type: none"> • Skin peeling, scaling, or blistering (with or without pus) which may also affect your eyes, mouth, nose or genitals, itching, severe rash, bumps under the skin, skin pain, skin colour changes (redness, yellowing, purplish) • Swelling and redness of eyes or face • Flu-like feeling, fever, chills, body aches, swollen glands, cough • Shortness of breath, chest pain or discomfort 			✓

If you have a troublesome symptom or side effect that is not listed here or becomes bad enough to interfere with your daily activities, talk to your healthcare professional.

Reporting Side Effects

You can report any suspected side effects associated with the use of health products to Health Canada by:

- Visiting the Web page on Adverse Reaction Reporting (<https://www.canada.ca/en/health-canada/services/drugs-health-products/medeffect-canada/adverse-reaction-reporting.html>) for information on how to report online, by mail or by fax; or
- Calling toll-free at 1-866-234-2345

NOTE: Contact your health professional if you need information about how to manage your side effects. The Canada Vigilance Program does not provide medical advice.

Storage:

- Cefuroxime for Injection, USP will be stored by your healthcare professional at room temperature (15°C and 30°C) in a dry place, protected from light.
- Keep out of reach and sight of children.

If you want more information about Cefuroxime for Injection, USP:

- Talk to your healthcare professional
- Find the full product monograph that is prepared for healthcare professionals and includes this Patient Medication Information by visiting the Health Canada website (<https://www.canada.ca/en/health-canada/services/drugs-health-products/drug-products/drug-product-database.html>); or the manufacturer's website (<https://www.hikma.com/canada>), or by calling 1-800-656-0793.

This leaflet was prepared by:

Hikma Canada Limited

Last Authorized: January 16, 2024