

PRODUCT MONOGRAPH
INCLUDING PATIENT MEDICATION INFORMATION

Pr JAMP Rivaroxaban

Rivaroxaban Tablets
Tablets 10 mg, 15 mg and 20 mg, Oral

Ph.Eur.

Anticoagulant
(ATC Classification: B01AF01)

JAMP Pharma Corporation
1310, rue Nobel
Boucherville, Québec
J4B 5H3

Date of Initial Authorization:
May 26, 2021

Date of Revision:
January 23, 2024

Submission Control Number: 280219

RECENT MAJOR LABEL CHANGES

7 WARNINGS AND PRECAUTIONS, Bleeding	03/2022
--	---------

TABLE OF CONTENTS

Section or subsections that are not applicable at the time of authorization are not listed.

RECENT MAJOR LABEL CHANGES..... 2

TABLE OF CONTENTS..... 2

PART I: HEALTH PROFESSIONAL INFORMATION 4

1 INDICATIONS..... 4

 1.1 Pediatrics4

 1.2 Geriatrics4

2 CONTRAINDICATIONS 4

4 DOSAGE AND ADMINISTRATION 5

 4.1 Dosing Considerations.....5

 4.2 Recommended Dose and Dosage Adjustment.....6

 4.4 Administration.....9

 4.5 Missed Dose10

5 OVERDOSAGE..... 10

6 DOSAGE FORMS, STRENGTHS, COMPOSITION AND PACKAGING 11

7 WARNINGS AND PRECAUTIONS 12

 7.1 Special Populations.....17

8 ADVERSE REACTIONS 18

 8.1 Adverse Reaction Overview18

 8.2 Clinical Trial Adverse Reactions.....22

 8.3 Less Common Clinical Trial Adverse Reactions26

 8.4 Abnormal Laboratory Findings: Hematologic, Clinical Chemistry and Other Quantitative DataClinical Trial Findings.....28

 8.5 Post-Market Adverse Reactions28

9 DRUG INTERACTIONS..... 29

 9.2 Drug Interactions Overview29

 9.4 Drug-Drug Interactions29

 9.5 Drug-Food Interactions.....33

 9.6 Drug-Herb Interactions33

 9.7 Drug-Laboratory Test Interactions33

10 CLINICAL PHARMACOLOGY 33

 10.1 Mechanism of Action.....33

 10.2 Pharmacodynamics33

 10.3 Pharmacokinetics36

11 STORAGE, STABILITY AND DISPOSAL 41

12 SPECIAL HANDLING INSTRUCTIONS 41

PART II: SCIENTIFIC INFORMATION 42

13 PHARMACEUTICAL INFORMATION..... 42

14 CLINICAL TRIALS 43

 14.1 Clinical Trials by Indication43

14.2 Comparative Bioavailability Studies	70
15 MICROBIOLOGY	71
16 NON-CLINICAL TOXICOLOGY	71
17 SUPPORTING PRODUCT MONOGRAPHS	72
PATIENT MEDICATION INFORMATION	73

PART I: HEALTH PROFESSIONAL INFORMATION

1 INDICATIONS

JAMP Rivaroxaban (rivaroxaban) film-coated tablets (10 mg, 15 mg, 20 mg) is indicated for the:

- prevention of venous thromboembolic events (VTE) in patients who have undergone elective total hip replacement (THR) or total knee replacement (TKR) surgery.
- treatment of venous thromboembolic events (deep vein thrombosis [DVT], pulmonary embolism [PE]) and prevention of recurrent DVT and PE.
- prevention of stroke and systemic embolism in patients with atrial fibrillation, in whom anticoagulation is appropriate.

Acute Pulmonary Embolus in hemodynamically unstable patients, or in those requiring thrombolysis or pulmonary embolectomy

For the treatment of VTE, JAMP Rivaroxaban is **not** recommended as an alternative to unfractionated heparin in patients with pulmonary embolus who are hemodynamically unstable, or who may receive thrombolysis or pulmonary embolectomy, since the safety and efficacy of rivaroxaban have not been established in these clinical situations (see [4 DOSAGE AND ADMINISTRATION](#)).

1.1 Pediatrics

Pediatrics (< 18 years of age): JAMP Rivaroxaban is not recommended for use in children below 18 years of age.

1.2 Geriatrics

Geriatrics (>65 years of age): Clinical studies have included patients with an age > 65 years (see [7 WARNINGS AND PRECAUTIONS – 7.1.4 Geriatrics](#) and [Renal Impairment](#), and [4 DOSAGE AND ADMINISTRATION – Renal Impairment](#) and [Geriatrics \(>65 years of age\)](#)).

Safety and efficacy data are available (see [14 CLINICAL TRIALS](#)).

2 CONTRAINDICATIONS

- Clinically significant active bleeding, including gastrointestinal bleeding.
- Lesions or conditions at increased risk of clinically significant bleeding, e.g, recent cerebral infarction (hemorrhagic or ischemic), active peptic ulcer disease with recent bleeding, patients with spontaneous or acquired impairment of hemostasis.
- Concomitant **systemic** treatment with strong inhibitors of **both** CYP 3A4 and P-glycoprotein (P-gp), such as cobicistat, ketoconazole, itraconazole, posaconazole, or ritonavir (see [7 WARNINGS AND PRECAUTIONS – Drug Interactions](#))
- Concomitant treatment with any other anticoagulant, including
 - unfractionated heparin (UFH), except at doses used to maintain a patent central venous or arterial catheter,
 - low molecular weight heparins (LMWH), such as enoxaparin and dalteparin,
 - heparin derivatives, such as fondaparinux, and
 - oral anticoagulants, such as warfarin, dabigatran, apixaban, edoxaban, except under circumstances of switching therapy to or from JAMP Rivaroxaban.

- Hepatic disease (including Child-Pugh Class B and C) associated with coagulopathy, and having clinically relevant bleeding risk (see [7 WARNINGS AND PRECAUTIONS – Hepatic Impairment](#))
- Pregnancy (see [7 WARNINGS AND PRECAUTIONS – 7.1.1 Pregnant Women](#))
- Nursing women (see [7 WARNINGS AND PRECAUTIONS – 7.1.2 Breast-feeding](#))
- Hypersensitivity to rivaroxaban or to any ingredient in the formulation, (see [6 DOSAGE FORMS, STRENGTHS, COMPOSITION AND PACKAGING](#)).

4 DOSAGE AND ADMINISTRATION

4.1 Dosing Considerations

As for any non-vitamin K antagonist oral anticoagulant (NOAC) drug, before initiating JAMP Rivaroxaban, ensure that the patient understands and is prepared to accept adherence to NOAC therapy, as directed.

Determine estimated creatinine clearance (eCrCl) in all patients before instituting JAMP Rivaroxaban (rivaroxaban), and monitor renal function during rivaroxaban tablets treatment, as clinically appropriate. Determination of renal function by eCrCl should occur at least once per year, and especially during circumstances when renal function may be expected to be compromised, i.e., acute myocardial infarction (AMI), acute decompensated heart failure (AHF), increased use of diuretics, dehydration, hypovolemia, etc. Clinically relevant deterioration of renal function may require dosage adjustment or discontinuation of rivaroxaban tablets (see below, [Renal Impairment](#)).

Glomerular filtration rate may be estimated by calculating eCrCl, using the Cockcroft-Gault formula:

eCrCl (mL/min)=

in males: $\frac{(140-\text{age}) (\text{years}) \times \text{weight} (\text{kg}) \times 1.23}{\text{serum creatinine} (\mu\text{mol/L})}$ or, $\frac{(140-\text{age}) (\text{yrs}) \times \text{weight} (\text{kg})}{72 \times \text{serum creatinine} (\text{mg}/100 \text{ mL})}$

in females: $\frac{(140-\text{age}) (\text{years}) \times \text{weight} (\text{kg}) \times 1.04}{\text{serum creatinine} (\mu\text{mol/L})}$ or, $\frac{(140-\text{age}) (\text{yrs}) \times \text{weight} (\text{kg}) \times 0.85}{72 \times \text{serum creatinine} (\text{mg}/100 \text{ mL})}$

Switching from Parenteral Anticoagulants to JAMP Rivaroxaban

JAMP Rivaroxaban can be started when the infusion of full-dose intravenous heparin is stopped or 0 to 2 hours before the next scheduled injection of full-dose subcutaneous low-molecular-weight heparin (LMWH) or fondaparinux. In patients receiving prophylactic heparin, LMWH or fondaparinux, JAMP Rivaroxaban can be started 6 or more hours after the last prophylactic dose.

Switching from JAMP Rivaroxaban to Parenteral Anticoagulants

Discontinue JAMP Rivaroxaban and give the first dose of parenteral anticoagulant at the time that the next JAMP Rivaroxaban dose was scheduled to be taken.

Switching from Vitamin K Antagonists (VKA) to JAMP Rivaroxaban

To switch from a VKA to JAMP Rivaroxaban, stop the VKA and determine the INR. If the INR is ≤ 2.5 , start JAMP Rivaroxaban at the usual dose. If the INR is > 2.5 , delay the start of JAMP Rivaroxaban until the INR is ≤ 2.5 (see [Considerations for INR Monitoring of VKA Activity during Concomitant JAMP Rivaroxaban Therapy](#)).

Switching from JAMP Rivaroxaban to a VKA

As with any short-acting anticoagulant, there is a potential for inadequate anticoagulation when transitioning from JAMP Rivaroxaban to a VKA. It is important to maintain an adequate level of anticoagulation when transitioning patients from one anticoagulant to another.

JAMP Rivaroxaban should be continued concurrently with the VKA until the INR is ≥ 2.0 . For the first 2 days of the conversion period, the VKA can be given in the usual starting doses without INR testing (see [Considerations for INR Monitoring of VKA Activity during Concomitant JAMP Rivaroxaban Therapy](#)).

Thereafter, while on concomitant therapy, the INR should be tested just prior to the next dose of JAMP Rivaroxaban, as appropriate. JAMP Rivaroxaban can be discontinued once the INR is > 2.0 . Once JAMP Rivaroxaban is discontinued, INR testing may be done at least 24 hours after the last dose of JAMP Rivaroxaban and should then reliably reflect the anticoagulant effect of the VKA.

Considerations for INR Monitoring of VKA Activity during Concomitant JAMP Rivaroxaban Therapy

In general, after starting VKA therapy, the initial anticoagulant effect is not readily apparent for at least 2 days, while the full therapeutic effect is achieved in 5-7 days. Consequently, INR monitoring in the first 2 days after starting a VKA is rarely necessary. Likewise, the INR may remain increased for a number of days after stopping VKA therapy.

Although JAMP Rivaroxaban therapy will lead to an elevated INR, depending on the timing of the measurement (see [10.2 Pharmacodynamics](#)), the INR is not a valid measure to assess the anticoagulant activity of JAMP Rivaroxaban. The INR is only calibrated and validated for VKA and should not be used for any other anticoagulant, including JAMP Rivaroxaban.

When switching patients from JAMP Rivaroxaban to a VKA, the INR should only be used to assess the anticoagulant effect of the VKA, and not that of JAMP Rivaroxaban. Therefore, while patients are concurrently receiving JAMP Rivaroxaban and VKA therapy, if the INR is to be tested, it should not be before 24 hours after the previous dose of JAMP Rivaroxaban, and should be just prior to the next dose of JAMP Rivaroxaban, since at this time the remaining JAMP Rivaroxaban concentration in the circulation is too low to have a clinically important effect on the INR. If INR testing is done earlier than just prior to the next dose of JAMP Rivaroxaban, the reported INR will not reflect the anticoagulation effect of the VKA only, because JAMP Rivaroxaban use may also affect the INR, leading to aberrant readings (see [10.2 Pharmacodynamics](#)).

4.2 Recommended Dose and Dosage Adjustment

Prevention of VTE after THR or TKR

The recommended dose is one 10 mg tablet once daily. JAMP Rivaroxaban 10 mg may be taken with or without food. The initial dose should be taken within 6 to 10 hours after surgery, provided that hemostasis has been established. If hemostasis is not established, treatment should be delayed.

The duration of administration depends on the type of surgery:

- After elective THR surgery, patients should be administered JAMP Rivaroxaban for 35 days.
- After elective TKR surgery, patients should be administered JAMP Rivaroxaban for 14 days.

Treatment of VTE and Prevention of recurrent DVT and PE

JAMP Rivaroxaban is NOT recommended as an alternative to unfractionated heparin in patients with acute pulmonary embolus who are hemodynamically unstable, or who may receive thrombolysis or pulmonary embolectomy, since the safety and efficacy of rivaroxaban tablets have not been established in these

clinical situations (see [1 INDICATIONS](#)).

The recommended dose for the initial treatment of acute DVT or PE is 15 mg twice daily (one tablet in the morning and one in the evening) for the first 3 weeks followed by 20 mg once daily for the continued treatment and prevention of recurrent DVT and PE.

Short duration of therapy (at least 3 months) should be considered in patients with DVT or PE provoked by major transient risk factors (e.g. recent major surgery or trauma). The duration of therapy should be individualized after careful assessment of the treatment benefit against the risk for bleeding.

Following completion of at least 6 months treatment for DVT or PE, the recommended dose for prevention of recurrent DVT and PE is 20 mg or 10mg once daily based on an individual assessment of the risk of recurrent DVT and PE against the risk for bleeding. For example, in patients in whom the risk of recurrent DVT or PE is considered high, such as those with complicated comorbidities who are at high risk of VTE recurrence, a dose of 20mg should be considered.

Longer duration of therapy should be considered in patients with DVT or PE provoked by permanent risk factors, unprovoked DVT or PE, or a history of recurrent DVT or PE.

The recommended maximum daily dose is 30 mg during the first 3 weeks of treatment and 20 mg thereafter.

JAMP Rivaroxaban 15 mg and 20 mg tablets should be taken with food. JAMP Rivaroxaban 10 mg tablets may be taken with or without food.

Prevention of Stroke and Systemic Embolism in Patients with Atrial Fibrillation

The recommended dose is one 20 mg tablet of JAMP Rivaroxaban taken once daily with food (see [10.3 Pharmacokinetics, Absorption](#)).

For patients with moderate renal impairment (CrCl 30 – 49 mL/min), the recommended dose is 15 mg once daily with food (see [Renal Impairment](#) below).

The recommended maximum daily dose is 20 mg.

Acute myocardial infarction (AMI): Consideration should be given to discontinuing JAMP Rivaroxaban in the setting of acute myocardial infarction should the treatment of myocardial infarction involve invasive procedures, such as percutaneous coronary revascularization, or coronary artery bypass surgery. Similar consideration should be given if thrombolytic therapy is to be initiated, because bleeding risk may increase. Patients with acute myocardial infarction should be treated according to current clinical guidelines. In this setting, JAMP Rivaroxaban may be resumed, when deemed clinically appropriate, for the prevention of stroke and systemic embolism upon completion of these revascularization procedures.

Concomitant use of ASA or clopidogrel with JAMP Rivaroxaban in patients with atrial fibrillation increases the risk of bleeding. Concomitant use of ASA or other antiplatelet agents based on medical need to prevent myocardial infarction should be undertaken with caution. Close clinical surveillance is recommended.

Other situations requiring thrombolytic therapy: JAMP Rivaroxaban should be discontinued in situations such as acute ischemic stroke where current clinical practice calls for administering thrombolytic therapy.

JAMP Rivaroxaban treatment may be subsequently resumed as soon as is deemed clinically appropriate. Measurement of a PT time, in seconds, using the Neoplastin reagent, may inform therapeutic decision-making (see [7 WARNINGS AND PRECAUTIONS – Monitoring and Laboratory Tests](#)).

Concomitant use of JAMP Rivaroxaban 10 mg, 15 mg and 20 mg with antiplatelet agents: The concomitant use of JAMP Rivaroxaban with antiplatelet agents increases the risk of bleeding (see [7 WARNINGS AND PRECAUTIONS – Bleeding](#)). If concomitant antiplatelet therapy is contemplated with JAMP Rivaroxaban 10 mg, 15 mg, and 20 mg, a careful assessment of the potential risks should be made against potential benefits, weighing risk of increased bleeding against expected benefit.

Patients with nonvalvular atrial fibrillation who undergo PCI with stent placement: Patients with nonvalvular atrial fibrillation who undergo PCI with stent placement should receive a reduced dose of 15 mg JAMP Rivaroxaban once daily (or 10 mg JAMP Rivaroxaban once daily for patients with moderate renal impairment [CrCl 30 – 49 mL/min]) in combination with a P2Y₁₂ inhibitor (e.g, clopidogrel). This treatment regimen is recommended for a maximum of 12 months after PCI with stent placement (see [10.2 Pharmacodynamics, Patients with nonvalvular atrial fibrillation who undergo PCI with stent placement](#)). After completion of the antiplatelet therapy, rivaroxaban dosage should be changed to the standard dose for patients with atrial fibrillation.

Cardioversion: Patients can be maintained on JAMP Rivaroxaban while being cardioverted (see [10.2 Pharmacodynamics, Patients undergoing cardioversion](#)).

Hepatic Impairment

JAMP Rivaroxaban is contraindicated in patients with hepatic disease (including Child-Pugh Class B and C) associated with coagulopathy and having clinically relevant bleeding risk. Patients with severe hepatic impairment or chronic hepatic disease were excluded from pivotal clinical trials.

The limited clinical data for patients with moderate hepatic impairment indicate a significant increase in the pharmacological activity. JAMP Rivaroxaban should be used with caution in these patients (see [2 CONTRAINDICATIONS, 7 WARNINGS AND PRECAUTIONS - Hepatic Impairment](#), and [10.3 Pharmacokinetics – Hepatic Insufficiency](#)).

The limited data available for patients with mild hepatic impairment without coagulopathy indicate that there is no difference in pharmacodynamic response or pharmacokinetics as compared to healthy subjects.

Renal Impairment

Table 1 – Dosage and Administration for Adult Patients According to Renal Function

Creatinine Clearance (CRCL)	Normal >80 mL/min	Mild 50-80 mL/min	Moderate 30-49 mL/min	Severe* 15 - < 30 mL/min	< 15 mL/min
Indication					
Prevention of VTE After THR or TKR	10 mg od			10 mg od	JAMP Rivaroxaban is not recommended
Treatment of VTE and Prevention of Recurrent DVT and PE	15 mg bid for 3 weeks, followed by 20 mg od			15 mg bid for 3 weeks, followed by 20 mg od	
Prevention of recurrent DVT and PE following completion of at least 6 months treatment	10 mg od or 20 mg od			10 mg od or 20 mg od	

Prevention of Stroke and Systemic Embolism in Patients with Atrial Fibrillation	20 mg od	15 mg od	15 mg od	
--	----------	----------	----------	--

od=once daily, bid=twice daily

*must be used with caution

JAMP Rivaroxaban should be used with caution in patients receiving other drugs which increase rivaroxaban plasma concentrations. Physicians should consider the benefit/risk of anticoagulant therapy before administering JAMP Rivaroxaban to patients with moderate renal impairment with a creatinine clearance close to the severe renal impairment category (CrCl < 30 mL/min) or with a potential to have deterioration of renal function during therapy. Renal function should be followed carefully in these patients (see [7 WARNINGS AND PRECAUTIONS – Renal Impairment](#) and [9.4 Drug-Drug Interactions](#)).

In patients with severe renal impairment (CrCl 15 - < 30 mL/min), rivaroxaban plasma levels may be significantly elevated compared to healthy volunteers (1.6-fold on average) which may lead to an increased bleeding risk. Due to limited clinical data, rivaroxaban tablets must be used with caution in these patients. No clinical data are available for patients with CrCl < 15 mL/min. Use is not recommended in patients with CrCl < 15 mL/min. Patients who develop acute renal failure while on JAMP Rivaroxaban should discontinue such treatment.

Gender, Race, or Body Weight

No dose adjustment is required for gender or race. (see [10.3 Pharmacokinetics – Sex, Ethnic Origin, and Body Weight](#))

Geriatrics (>65 years of age)

No dose adjustment is generally required for the elderly. Increasing age may be associated with declining renal function (see [7 WARNINGS AND PRECAUTIONS – Renal Impairment](#), and [4 DOSAGE AND ADMINISTRATION – Renal Impairment](#)).

Pediatrics (<18 years of age)

JAMP Rivaroxaban is not recommended for use in children below 18 years of age.

4.4 Administration

Tablet:

Swallow the tablet with liquid.

JAMP Rivaroxaban 15 mg and 20 mg tablets should be taken with food (see [10.3 Pharmacokinetics, Absorption](#)).

Administration of Crushed Tablets:

For patients who are unable to swallow whole tablets, JAMP Rivaroxaban tablets may be crushed and mixed with applesauce immediately prior to use and administered orally. After the administration of a crushed JAMP Rivaroxaban 15 mg or 20 mg tablet, the dose should be immediately followed by food.

A crushed JAMP Rivaroxaban tablet may be also administered via nasogastric (NG) tube. After confirming gastric placement of the NG tube, the crushed tablet should be suspended in 50 mL of water and administered via the NG tube after which it should be flushed with water. Because rivaroxaban absorption is dependent on the site of drug release in the GI tract, avoid administration of JAMP

Rivaroxaban distal to the stomach as this can result in reduced absorption and therefore reduced drug exposure. After the administration of a crushed JAMP Rivaroxaban 15 mg or 20 mg tablet, the dose should then be immediately followed by enteral feeding (see [10.3 Pharmacokinetics, Absorption](#)).

An *in vitro* compatibility study indicated that there is no adsorption of rivaroxaban from a water suspension of a crushed JAMP Rivaroxaban tablet to PVC or silicone nasogastric (NG) tubing.

4.5 Missed Dose

It is essential to adhere to the dosage schedule provided.

- JAMP Rivaroxaban 10 mg, 15 mg, or 20 mg tablets taken **once** a day:

If a dose is missed, the patient should take JAMP Rivaroxaban immediately and continue on the following day with the once daily intake as before. A double dose should not be taken to make up for a missed tablet.

- JAMP Rivaroxaban 15 mg taken **twice** a day:

If a dose is missed during the 15 mg twice daily treatment phase the patient should take the next dose immediately to ensure the intake of 30 mg total dose per day. In this case two 15 mg tablets may be taken at once. The following day the patient should continue with the regular 15 mg twice daily intake schedule as recommended.

5 OVERDOSAGE

For management of a suspected drug overdose, contact your regional poison control centre.

Overdose following administration of JAMP Rivaroxaban may lead to hemorrhagic complications due to its pharmacodynamic properties.

In adults, rare cases of overdose up to 600 mg have been reported without bleeding complications or other adverse reactions. No further increase in average plasma exposure is expected due to limited absorption at supratherapeutic doses of 50 mg or above in adults, however, no data are available at supratherapeutic doses in children, because of a solubility ceiling effect.

A specific reversal agent for JAMP Rivaroxaban is not available. The use of activated charcoal to reduce absorption in case of JAMP Rivaroxaban overdose may be considered. Administration of activated charcoal up to 8 hours after overdose may reduce the absorption of JAMP Rivaroxaban.

Due to the high plasma protein binding, JAMP Rivaroxaban is not expected to be removed by dialysis (see [10.3 Pharmacokinetics, Distribution](#)).

Management of Bleeding

In the event of hemorrhagic complications in a patient receiving JAMP Rivaroxaban, treatment should be temporarily discontinued, and the source of bleeding investigated. JAMP Rivaroxaban has a half-life of approximately 5 to 13 hours in adults. Consideration should be given to the resumption of antithrombotic therapy when clinically appropriate to adequately control risk of underlying thrombosis.

Management of bleeding should be individualised according to the severity and location of the hemorrhage. Appropriate symptomatic treatment should be used as needed, such as mechanical compression (e.g. for severe epistaxis), surgical hemostasis with bleeding control procedures, fluid replacement and hemodynamic support, blood products (packed red cells or fresh frozen plasma, depending on associated anemia or coagulopathy) or platelets.

If bleeding cannot be controlled by the above measures, consider administration of one of the following procoagulants:

- activated prothrombin complex concentrate (APCC), e.g, FEIBA
- prothrombin complex concentrate (PCC)
- recombinant Factor-VIIa (rFVIIa)

However, there is currently only very limited experience with the use of these products in adults receiving rivaroxaban tablets.

In a randomized, double-blind, placebo-controlled study, a non-activated prothrombin complex concentrate (PCC) given to 6 healthy male subjects who had previously received rivaroxaban tablets, completely reversed its anticoagulant effect within 15 minutes, based on coagulation tests. Although this study may have important clinical implications, this effect of PCC has not yet been confirmed in patients with active bleeding who have been previously treated with rivaroxaban tablets.

Protamine sulfate and vitamin K are not expected to affect the anticoagulant activity of rivaroxaban tablets. There is limited experience with tranexamic acid and no experience with aminocaproic acid and aprotinin in adults receiving rivaroxaban tablets. There is no experience on the use of these agents in children receiving rivaroxaban tablets. There is neither scientific rationale for benefit or experience with the systemic hemostatic desmopressin in individuals receiving rivaroxaban tablets.

The prothrombin time (PT), measured in seconds, is influenced by rivaroxaban tablets in a dose-dependent way with a close correlation to plasma concentrations if the Neoplastin® reagent is used. In patients who are bleeding, measuring the PT (Neoplastin® reagent) may be useful to assist in determining an excess of anticoagulant activity. INR should **NOT** be used to assess the anticoagulant effect of JAMP Rivaroxaban (see [7 WARNINGS AND PRECAUTIONS – Monitoring and Laboratory Tests](#) and [10.2 Pharmacodynamics](#)).

6 DOSAGE FORMS, STRENGTHS, COMPOSITION AND PACKAGING

Table 2 – Dosage Forms, Strengths, Composition and Packaging

Route of Administration	Dosage Form / Strength/Composition	Non-medicinal Ingredients
Oral	Film-coated tablet, 10 mg, 15 mg and 20 mg	Cellulose microcrystalline, croscarmellose sodium, hypromellose, lactose monohydrate, magnesium stearate, sodium lauryl sulphate. The tablet film coating consists of hypromellose, iron oxide red, polyethylene glycol and titanium dioxide.

10 mg Tablets:

Round, biconvex, pink film-coated tablets, engraved by '10' on one side and plain on the other side. JAMP Rivaroxaban 10 mg tablets are supplied in bottle of 100 tablets.

15 mg Tablets:

Round, biconvex, red film-coated tablets, engraved by '15' on one side and plain on the other side. JAMP Rivaroxaban tablets 15 mg are supplied in bottle of 100 tablets.

20 mg Tablets:

Round, biconvex, dark red film-coated tablets, engraved by '20' on one side and plain on the other side. JAMP Rivaroxaban tablets 20 mg are supplied in bottle of 100 tablets.

7 WARNINGS AND PRECAUTIONS**General****PREMATURE DISCONTINUATION OF ANY ORAL ANTICOAGULANT, INCLUDING JAMP Rivaroxaban, INCREASES THE RISK OF THROMBOTIC EVENTS.**

To reduce this risk, consider coverage with another anticoagulant if JAMP Rivaroxaban is discontinued for a reason other than pathological bleeding or completion of a course of therapy.

Bleeding

JAMP Rivaroxaban, like other anticoagulants, should be used with caution in patients with an increased bleeding risk. Bleeding can occur at any site during therapy with JAMP Rivaroxaban. The possibility of a hemorrhage should be considered in evaluating the condition of any anticoagulated patient. Any unexplained fall in hemoglobin or blood pressure should lead to a search for a bleeding site. Patients at high risk of bleeding should not be prescribed JAMP Rivaroxaban (see [2 CONTRAINDICATIONS](#)).

Should severe bleeding occur, treatment with JAMP Rivaroxaban must be discontinued and the source of bleeding investigated promptly.

Close clinical surveillance (looking for signs of bleeding or anemia) is recommended throughout the treatment period, especially in the presence of multiple risk factors for bleeding (see [Table 3](#) below).

Table 3 – Factors Which Increase Hemorrhagic Risk

Factors increasing rivaroxaban plasma levels	Severe renal impairment (CrCl < 30 mL/min)
	Concomitant systemic treatment with strong inhibitors of both CYP 3A4 and P-gp
Pharmacodynamic interactions	NSAID
	Platelet aggregation inhibitors, including ASA, clopidogrel, prasugrel, ticagrelor
	Selective serotonin reuptake inhibitors (SSRI), and serotonin norepinephrine reuptake inhibitors (SNRIs)
Diseases / procedures with special hemorrhagic risks	Congenital or acquired coagulation disorders
	Thrombocytopenia or functional platelet defects
	Uncontrolled severe arterial hypertension
	Active ulcerative gastrointestinal disease
	Recent gastrointestinal bleeding
	Vascular retinopathy, such as hypertensive or diabetic

	Recent intracranial hemorrhage
	Intraspinal or intracerebral vascular abnormalities
	Recent brain, spinal or ophthalmological surgery
	Bronchiectasis or history of pulmonary bleeding
Others	Age > 75 years

Concomitant use of drugs affecting hemostasis increases the risk of bleeding. Care should be taken if patients are treated concomitantly with drugs affecting hemostasis such as non-steroidal anti-inflammatory drugs (NSAIDs), acetylsalicylic acid (ASA), platelet aggregation inhibitors or selective serotonin reuptake inhibitors (SSRI), and serotonin norepinephrine reuptake inhibitors (SNRIs) (see also [9 DRUG INTERACTIONS](#)).

In patients with atrial fibrillation and having a condition that warrants single or dual antiplatelet therapy, a careful assessment of the potential benefits against the potential risks should be made before combining this therapy with JAMP Rivaroxaban.

Concomitant ASA use (almost exclusively at a dose of 100 mg or less) with either rivaroxaban tablets or warfarin during the ROCKET-AF trial was identified as an independent risk factor for major bleeding (see also [9 DRUG INTERACTIONS](#)).

The antiplatelet agents, prasugrel and ticagrelor, have not been studied with rivaroxaban tablets, and are not recommended as concomitant therapy.

The use of thrombolytics should generally be avoided during acute myocardial infarction (AMI) or acute stroke in patients treated with rivaroxaban, due to expected increased risk of major bleeding (see [4 DOSAGE AND ADMINISTRATION](#), [Other situations requiring thrombolytic therapy](#)).

Patients with malignant disease may simultaneously be at higher risk of bleeding and thrombosis than patients without malignant disease. In patients with malignant disease who undergo antithrombotic treatment with rivaroxaban, risk of bleeding may increase, in particular in the gastrointestinal, genitourinary tract or respiratory system. The individual benefit of antithrombotic treatment should be weighed against risk for bleeding in patients with active cancer dependent on tumor location, antineoplastic therapy and stage of disease.

Cardiovascular

Patients with valvular disease

JAMP Rivaroxaban is not indicated and is not recommended for thromboprophylaxis in patients having recently undergone transcatheter aortic valve replacement (TAVR). Results from a randomized controlled clinical study (GALILEO) showed that the rivaroxaban tablets regimen failed to demonstrate clinical benefit compared with an antiplatelet strategy. In the intention-to-treat analysis, all-cause mortality, thromboembolic and bleeding events occurred more frequently in patients randomized to the rivaroxaban tablets regimen. A causal relationship between rivaroxaban tablets and all-cause mortality could not be established.

Safety and efficacy of rivaroxaban tablets have not been studied in patients with other prosthetic heart valves or other valve procedures, or those with hemodynamically significant rheumatic heart disease, especially mitral stenosis. There are no data to support that rivaroxaban tablets provides adequate anticoagulation in patients with prosthetic heart valves, with or without atrial fibrillation. Therefore, the use of JAMP Rivaroxaban is not recommended in this setting.

Of note, in the pivotal Phase III ROCKET AF trial that evaluated rivaroxaban tablets in the prevention of stroke in atrial fibrillation, 14% of patients had other valvular disease including aortic stenosis, aortic regurgitation, and/or mitral regurgitation. Patients with a history of mitral valve repair were also not excluded from the study. Mitral valve repair rates are not known in ROCKET AF, since information on mitral valve repair status was not specifically collected in this study.

Patients with antiphospholipid syndrome

JAMP Rivaroxaban is not recommended for patients with a history of thrombosis who are diagnosed with antiphospholipid syndrome. In particular for patients who are triple positive (for lupus anticoagulant, anticardiolipin antibodies, and anti-beta 2-glycoprotein I antibodies), treatment with rivaroxaban is associated with an increased rate of recurrent thrombotic events compared with vitamin K antagonists.

Patients with nonvalvular atrial fibrillation who undergo PCI (Percutaneous Coronary Intervention) with stent placement

Clinical data are available from an open label interventional study with the primary objective to assess safety in patients with nonvalvular atrial fibrillation who undergo PCI with stent placement. Data on efficacy in this population are limited (see [4 DOSAGE AND ADMINISTRATION – Patients with nonvalvular atrial fibrillation who undergo PCI with stent placement](#); [10.2 Pharmacodynamics, Patients with nonvalvular atrial fibrillation who undergo PCI with stent placement](#)).

Drug Interactions

Drug interaction studies have only been performed in adults. (see [9 DRUG INTERACTIONS](#)).

Interaction with strong inhibitors of both CYP 3A4 and P-gp

The use of JAMP Rivaroxaban is contraindicated in patients receiving concomitant systemic treatment with strong inhibitors of both CYP 3A4 and P-gp, such as cobicistat, ketoconazole, itraconazole, posaconazole, or ritonavir. These drugs may increase JAMP Rivaroxaban plasma concentrations to a clinically relevant degree, i.e, 2.6-fold on average, which increases bleeding risk.

Dronedarone should not be used concomitantly with rivaroxaban since it may increase exposure of rivaroxaban through P-gp and CYP3A4 inhibition, and thereby the risk of bleeding.

Interaction with moderate CYP 3A4 inhibitors

The azole anti-mycotic, fluconazole, a moderate CYP 3A4 inhibitor, or erythromycin, have no clinically relevant effect on rivaroxaban exposure (1.4-fold and 1.3-fold increase, respectively) and may be co-administered with JAMP Rivaroxaban in patients with normal renal function (see [9 DRUG INTERACTIONS](#)).

The use of rivaroxaban tablets in subjects with mild and moderate renal impairment concomitantly treated with combined P-gp and moderate CYP 3A4 inhibitors such as erythromycin increased exposure to rivaroxaban by 1.8- and 2.0-fold, respectively, compared to subjects with normal renal function without medication. If such use must be undertaken, caution is required.

Interaction with strong CYP 3A4 inducers

The concomitant use of JAMP Rivaroxaban with strong inducers of CYP 3A4, such as rifampicin, and the anticonvulsants, phenytoin, carbamazepine, phenobarbital, reduces rivaroxaban exposure (see [9.4 Drug-Drug Interactions](#)). Combined use of JAMP Rivaroxaban with strong inducers should generally be avoided, since efficacy of rivaroxaban tablets may be compromised (see [9.4 Drug-Drug Interactions](#)).

Hepatic Impairment

Patients with significant hepatic disease (e.g, acute clinical hepatitis, chronic active hepatitis, liver cirrhosis) were excluded from clinical trials. Therefore, JAMP Rivaroxaban is contraindicated in patients with hepatic disease (including Child-Pugh Class B and C) associated with coagulopathy and having clinically relevant bleeding risk.

The limited data available for patients with mild hepatic impairment without coagulopathy indicate that there is no difference in pharmacodynamic response or pharmacokinetics as compared to healthy subjects.

No clinical data are available in children with hepatic impairment.

Surgery / Procedural Interventions

As with any anticoagulant, patients on JAMP Rivaroxaban who undergo surgery or invasive procedures are at increased risk for bleeding. In these circumstances, temporary discontinuation of JAMP Rivaroxaban may be required.

If a patient concomitantly receiving platelet aggregation inhibitors is to undergo elective surgery and anti-platelet effect is not desired, platelet aggregation inhibitors should be discontinued as directed by the manufacturer's prescribing information.

Limited clinical data are available for patients undergoing fracture-related surgery of the lower limbs. These patients were from a subgroup which was not pre-specified for enrollment in an international, non-interventional (no exclusion criteria), open label cohort study designed to compare the incidence of symptomatic thromboembolic events in patients undergoing elective hip or knee surgery while not randomly assigned to treatment with JAMP Rivaroxaban or any local standard-of-care pharmacological therapy.

Pre-Operative Phase

If an invasive procedure or surgical intervention is required, JAMP Rivaroxaban 10 mg, 15 mg and 20 mg should be stopped at least 24 hours before the intervention, if possible, due to increased risk of bleeding, and based on clinical judgment of the physician. If a patient is to undergo elective surgery and anti-platelet effect is not desired, platelet aggregation inhibitors should be discontinued as per current treatment guidelines. If the procedure cannot be delayed, the increased risk of bleeding should be assessed against the urgency of the intervention. Although there are limited data, in patients at higher risk of bleeding or in major surgery where complete hemostasis may be required, consider stopping JAMP Rivaroxaban two to four days before surgery, depending on clinical circumstances.

Peri-Operative Spinal/Epidural Anesthesia, Lumbar Puncture

When neuraxial (epidural/spinal) anesthesia or spinal puncture is performed, patients treated with antithrombotics for prevention of thromboembolic complications are at risk for developing an epidural or spinal hematoma that may result in long-term neurological injury or permanent paralysis.

The risk of these events is even further increased by the use of indwelling epidural catheters or the

concomitant use of drugs affecting hemostasis. Accordingly, the use of JAMP Rivaroxaban, at doses greater than 10 mg, is not recommended in patients undergoing anesthesia with post-operative indwelling epidural catheters. The risk may also be increased by traumatic or repeated epidural or spinal puncture. If traumatic puncture occurs, the administration of JAMP Rivaroxaban should be delayed for 24 hours.

Patients who have undergone epidural puncture and who are receiving JAMP Rivaroxaban 10 mg should be frequently monitored for signs and symptoms of neurological impairment (e.g, numbness or weakness of the legs, bowel or bladder dysfunction). If neurological deficits are noted, urgent diagnosis and treatment is necessary.

The physician should consider the potential benefit versus the risk before neuraxial intervention in patients anticoagulated or to be anticoagulated for thromboprophylaxis and use JAMP Rivaroxaban 10 mg only when the benefits clearly outweigh the possible risks. An epidural catheter should not be withdrawn earlier than 18 hours after the last administration of JAMP Rivaroxaban. JAMP Rivaroxaban should be administered not earlier than 6 hours after the removal of the catheter in adults.

There is no clinical experience with the use of rivaroxaban tablets 15 mg and 20 mg, in this situation.

To reduce the potential risk of bleeding associated with the concurrent use of rivaroxaban and neuraxial (epidural/spinal) anesthesia or lumbar puncture, consider the pharmacokinetic profile of rivaroxaban. Placement or removal of an epidural catheter or lumbar puncture is best performed when the anticoagulant effect of rivaroxaban is estimated to be low. However, the exact timing to reach a sufficiently low anticoagulant effect in each patient is not known and should be weighed against the urgency of a diagnostic procedure.

Post-Procedural Period

JAMP Rivaroxaban should be restarted following an invasive procedure or surgical intervention as soon as adequate hemostasis has been established and the clinical situation allows, in order to avoid unnecessary increased risk of thrombosis.

Renal Impairment

Following oral dosing with JAMP Rivaroxaban, there is a direct relationship between pharmacodynamic effects and the degree of renal impairment (see [10.3 Pharmacokinetics – Renal Insufficiency](#)).

Determine estimated creatinine clearance (eCrCl) in all patients before instituting JAMP Rivaroxaban (see [4 DOSAGE AND ADMINISTRATION](#)).

JAMP Rivaroxaban should be used with caution in patients with moderate renal impairment (CrCl 30-49 mL/min), especially in those concomitantly receiving other drugs which increase rivaroxaban plasma concentrations (see [4 DOSAGE AND ADMINISTRATION – Renal Impairment](#), and [9.4 Drug-Drug Interactions](#)).

Physicians should consider the benefit/ risk of anticoagulant therapy before administering JAMP Rivaroxaban to patients with moderate renal impairment having a creatinine clearance close to the severe renal impairment category (CrCl < 30 mL/min), or in those with a potential to have deterioration of renal function to severe impairment during therapy.

In patients with severe renal impairment (CrCl 15 - < 30 mL/min), rivaroxaban plasma levels may be significantly elevated compared to healthy volunteers (1.6-fold on average) which may lead to an increased bleeding risk. Due to limited clinical data, rivaroxaban tablets must be used with caution in

these patients. No clinical data are available for patients with CrCl < 15 mL/min. Use is not recommended in patients with CrCl < 15ml/min. Patients who develop acute renal failure while on JAMP Rivaroxaban should discontinue such treatment.

Due to the high plasma protein binding, i.e, about 95%, rivaroxaban is not expected to be removed by dialysis.

Lactose Sensitivity

JAMP Rivaroxaban contains lactose. Patients with rare hereditary problems of lactose or galactose intolerance (e.g, the Lapp lactase deficiency or glucose-galactose malabsorption) should not take JAMP Rivaroxaban.

Monitoring and Laboratory Tests

The prothrombin time (PT), measured in seconds, is influenced by JAMP Rivaroxaban in a dose-dependent way with a close correlation to plasma concentration if the Neoplastin® reagent is used. In patients who are bleeding, measuring the PT using the Neoplastin® reagent may be useful to assist in determining an excess of anticoagulant activity (see [4 DOSAGE AND ADMINISTRATION – Considerations for INR Monitoring of VKA Activity during Concomitant JAMP Rivaroxaban Therapy](#)).

Although JAMP Rivaroxaban therapy will lead to an elevated INR, depending on the timing of the measurement (see [10.2 Pharmacodynamics](#)), the INR is not a valid measure to assess the anticoagulant activity of JAMP Rivaroxaban. The INR is only calibrated and validated for VKA and should not be used for any other anticoagulant, including JAMP Rivaroxaban.

At recommended doses, JAMP Rivaroxaban affects the measurement of the aPTT and Heptest®. These tests are not recommended for the assessment of the pharmacodynamic effects of JAMP Rivaroxaban (see [10.2 Pharmacodynamics](#)).

Converting patients from warfarin to JAMP Rivaroxaban, or from JAMP Rivaroxaban to warfarin, increases prothrombintime by the Neoplastin® reagent in seconds (or INR values) more than additively (e.g, individual INR values up to 12 may be observed) during concomitant therapy, whereas effects on aPTT and endogenous thrombin potential are additive (see [10.2 Pharmacodynamics](#)).

Anti-Factor-Xa activity is influenced by JAMP Rivaroxaban in a dose-dependent fashion. If it is desired to test the pharmacodynamic effects of JAMP Rivaroxaban during the switching period, tests of anti-Factor-Xa activity can be used as they are not affected by warfarin. Use of these tests to assess the pharmacodynamic effects of JAMP Rivaroxaban requires calibration and should not be done unless JAMP Rivaroxaban -specific calibrators and controls are available (see [10.2 Pharmacodynamics](#)).

Although there is no need for routine monitoring of anticoagulation effect of JAMP Rivaroxaban during clinical practice, in certain infrequent situations such as overdosage, acute bleeding, urgent surgery, in cases of suspected non-compliance, or in other unusual circumstances, assessment of the anticoagulant effect of rivaroxaban may be appropriate. Accordingly, measuring PT using the Neoplastin reagent, or Factor-Xa assay using rivaroxaban-specific calibrators and controls, may be useful to inform clinical decisions in these circumstances.

7.1 Special Populations

7.1.1 Pregnant Women

No data are available on the use of rivaroxaban tablets in pregnant women.

Based on animal data, use of rivaroxaban tablets is contraindicated throughout pregnancy (see [2 CONTRAINDICATIONS](#), and [16 NON-CLINICAL TOXICOLOGY – Reproductive and Developmental Toxicology](#)).

If JAMP Rivaroxaban is to be used in women of childbearing potential, pregnancy should be avoided.

7.1.2 Breast-feeding

No data are available on the use of rivaroxaban tablets in nursing mothers. In rats, rivaroxaban tablets is secreted into breast milk. Therefore, JAMP Rivaroxaban should only be administered after breastfeeding is discontinued (see [2 CONTRAINDICATIONS](#), and [16 NON-CLINICAL TOXICOLOGY – Reproductive and Developmental Toxicology](#)).

7.1.3 Pediatrics

Pediatric Use (< 18 years of age): JAMP Rivaroxaban is not recommended for use in children below 18 years of age.

7.1.4 Geriatrics

Geriatrics (> 65 years of age): Increasing age is associated with declining renal function. Both of these factors have been observed to result in increased systemic exposure to rivaroxaban, and consequently increased bleeding (see [7 WARNINGS AND PRECAUTIONS – Renal Impairment](#), and [4 DOSAGE AND ADMINISTRATION – Renal Impairment](#)).

Increasing age may increase hemorrhagic risk.

Use with caution in elderly patients, especially those taking concomitant medications that increase systemic exposure of JAMP Rivaroxaban (see [7 WARNINGS AND PRECAUTIONS, Drug Interactions](#), and [9 DRUG INTERACTIONS](#)).

8 ADVERSE REACTIONS

8.1 Adverse Reaction Overview

Prevention of VTE after THR or TKR

The safety of rivaroxaban tablets 10 mg has been evaluated in three randomized, double-blind, active-control Phase III studies (RECORD 1, RECORD 2, and RECORD 3). In the Phase III studies, 4657 patients undergoing total hip replacement or total knee replacement surgery were randomized to rivaroxaban tablets, with 4571 patients actually receiving rivaroxaban tablets.

In RECORD 1 and 2, a total of 2209 and 1228 THR patients, respectively, were randomized to rivaroxaban tablets 10 mg od. In RECORD 1, the treatment period for both groups were 35±4 days postoperatively. In RECORD 2, patients randomized to rivaroxaban tablets were treated for 35 ±4 days postoperatively, and patients randomized to enoxaparin received placebo after day 12±2 until day 35±4 postoperatively. In RECORD 3, a total of 1220 TKR patients were randomized to rivaroxaban tablets 10 mg od, and both groups received study drug until day 12±2 postoperatively.

Treatment of VTE and Prevention of Recurrent DVT and PE

The safety of rivaroxaban tablets has been evaluated in four Phase III trials with 6790 patients treated up to 21 months. Patients were exposed to 15 mg rivaroxaban tablets twice daily for 3 weeks followed by:

- 20 mg once daily (EINSTEIN DVT, EINSTEIN PE) or
- 20 mg once daily after at least 6 months of treatment for DVT or PE (EINSTEIN Extension), or

- 20 mg or 10 mg rivaroxaban tablets once daily after at least 6 months of treatment for DVT or PE (EINSTEIN CHOICE).

The mean treatment duration was 194 days in EINSTEIN DVT, 183 days in EINSTEIN PE, 188 days in EINSTEIN Extension and 290 days in EINSTEIN CHOICE.

The incidence of adverse events resulting in permanent discontinuation of study drug was 5.0% for rivaroxaban tablets and 4.4% for enoxaparin/VKA (pooled data from EINSTEIN DVT and EINSTEIN PE), 6.5% for rivaroxaban tablets and 3.4% for placebo (EINSTEIN Extension) and 4.5% for rivaroxaban tablets 10 mg, 4.5% for rivaroxaban tablets 20 mg and 4.2% for ASA (EINSTEIN CHOICE).

Prevention of Stroke and Systemic Embolism in Patients with Atrial Fibrillation (SPAF)

In the pivotal double-blind ROCKET AF study, a total of 14,264 patients with atrial fibrillation at risk for stroke and systemic embolism were randomly assigned to treatment with either rivaroxaban (7,131) or warfarin (7,133) in 45 countries. Patients received rivaroxaban tablets 20 mg orally once daily (15 mg orally once daily in patients with moderate renal impairment [CrCl: 30-49 mL/min]) or dose-adjusted warfarin titrated to a target INR of 2.0 to 3.0. The safety population included patients who were randomized and took at least 1 dose of study medication. In total, 14,236 patients were included in the safety population, with 7,111 and 7,125 patients in rivaroxaban and warfarin groups, respectively. The mediantime on treatment was 19 months and overall treatment duration was up to 41 months.

The incidence of adverse events resulting in permanent discontinuation of study drug was 15.8% in the rivaroxaban group and 15.2% in the warfarin group.

Bleeding

Due to the pharmacological mode of action, JAMP Rivaroxaban is associated with an increased risk of occult or overt bleeding from any tissue and organ (see [7 WARNINGS AND PRECAUTIONS – Bleeding](#), and [Drug Interactions](#)). The risk of bleeding may be increased in certain patient groups, e.g, patients with uncontrolled severe arterial hypertension and/or on concomitant medication affecting hemostasis (see [Table 3](#)). The signs, symptoms, and severity (including fatal outcome) will vary according to the location and degree or extent of the bleeding and/or anemia. Hemorrhagic complications may present as weakness, paleness, dizziness, headache or unexplained swelling, dyspnea, and unexplained shock. In some cases, as a consequence of anemia, symptoms of cardiac ischemia like chest pain or angina pectoris have been observed. Known complications secondary to severe bleeding such as compartment syndrome and renal failure due to hypoperfusion have been reported for rivaroxaban tablets. Therefore, the possibility of a hemorrhage should be considered in evaluating the medical condition in any anticoagulated patient.

Major or severe bleeding may occur and, regardless of location, may lead to disabling, life-threatening or even fatal outcomes.

Since the adverse event profiles of the patient populations treated with rivaroxaban tablets for different indications are not interchangeable, a summary description of major and total bleeding is provided by indication, in [Table 4](#) for VTE prevention in patients undergoing elective THR or TKR surgery, in [Table 5](#) for Treatment of VTE and prevention of recurrent DVT and PE, in [Table 6](#) for stroke prevention in atrial fibrillation.

Table 4 - RECORD 1, 2, and 3 (VTE Prevention After THR or TKR) – Treatment-Emergent Bleeding Events (Safety Population with Central Adjudication) in Patients Randomized to rivaroxaban tablets (First Dose 6 to 8 Hours Postoperatively) or Enoxaparin (First Dose 12 Hours Preoperatively)

		Major Bleeding ^a n (%)	Major Bleeding Including Surgical Site Bleeding Events Associated With Hemoglobin Drops or Transfusions n (%)	Any Bleeding (Major or Nonmajor) n (%)
RECORD 1 (THR)	Rivaroxaban tablets (N=2209) 10 mg od po for 35±4 days	6 (0.3)	40 (1.8)	133 (6.0)
	Enoxaparin (N=2224) 40 mg od SC for 36±4 days	2 (0.1)	33 (1.5)	131 (5.9)
	<i>P</i> -Value	0.18	0.41	0.90
RECORD 2 (THR)	Rivaroxaban tablets (N=1228) 10 mg od po for 35±4 days	1 (0.1)	23 (1.9)	81 (6.6)
	Enoxaparin (N=1229) 40 mg od SC for 12±2 days	1 (0.1)	19 (1.6)	68 (5.5)
	<i>P</i> -Value	1.00	0.54	0.273
RECORD 3 (TKR)	Rivaroxaban tablets (N=1220) 10 mg od po for 12±2 days	7 (0.6)	21 (1.7)	60 (4.9)
	Enoxaparin (N=1239) 40 mg od SC for 13±2 days	6 (0.5)	17 (1.4)	60 (4.8)
	<i>P</i> -Value	0.79	0.52	1.00
Pooled Analysis (RECORD 1, 2, 3)	Rivaroxaban tablets (N=4657)	14 (0.3)	84 (1.8)	274 (5.9)
	Enoxaparin (N=4692) 40 mg od SC	9 (0.2)	69 (1.5)	259 (5.5)
	<i>P</i> -Value	0.31	0.22	0.48

^a Major bleeding events included: (1) fatal, (2) bleeding into a critical organ (e.g, retroperitoneal, intracranial, intraocular or intraspinal bleeding/hemorrhagic puncture), (3) bleeding requiring reoperation, (4) clinically overt extra-surgical site bleeding associated with ≥2 g/dL fall in hemoglobin or leading to infusion of ≥2 units of whole blood or packed cells.

See [Table 19](#) and [Table 21](#) for additional details.

od = once daily, po = oral, SC = subcutaneous

Table 5 - Treatment-Emergent Bleeding Events and Results – Safety Population with Central Adjudication - Pooled Analysis, EINSTEIN DVT, EINSTEIN PE, EINSTEIN Extension and EINSTEIN CHOICE (Treatment of VTE and Prevention of Recurrent DVT and PE)

Bleeding event	Pooled EINSTEIN DVT and EINSTEIN PE			EINSTEIN Extension		EINSTEIN CHOICE		
	Rivaroxaban tablets N=4130	Enox/VKA N=4116	HR (95% CI) P-value for superiority	20 mg od N=598	Placebo N=590	Rivaroxaban tablets 10 mg N=1127	Rivaroxaban tablets 20 mg N=1107	ASA 100 mg N=1131
	n (%)	n (%)		n (%)	n (%)	n (%)	n (%)	n (%)
Major and Clinically Relevant Non-major Bleeding ^a	388 (9.4)	412 (10.0)	0.93 (0.81-1.06) P=0.27	36 (6.0)	7 (1.2)	27 (2.4)	36 (3.3)	23(2.0)
Major Bleeding ^b	40 (1.0)	72 (1.7)	0.54 (0.37-0.80) P=0.0018*	4 (0.7)	0	5 (0.4)	6 (0.5)	3 (0.3)
Fatal Bleeding	3 (<0.1)	8 (0.2)	-	0	0	0	1 (<0.1)	1 (<0.1)
Intracranial	2 (<0.1)	4 (<0.1)	-	0	0	0	0	1 (<0.1)
Non-Fatal Critical Organ Bleeding	10 (0.2)	29 (0.7)	-	0	0	2 (0.2)	4 (0.4)	1 (<0.1)
Intracranial	3 (<0.1)	10 (0.2)	-	0	0	1 (<0.1)	3 (0.3)	1 (<0.1)
Non-Fatal Non-Critical Organ Bleeding (Fall in Hb ≥ 2 g/dL and/or Transfusions ≥ 2 Units	27 (0.7)	37 (0.9)	-	4	0	3 (0.3)	1 (<0.1)	1 (<0.1)
Gastrointestinal	12 (0.3)	20 (0.5)	-	3	0	2 (0.2)	1 (<0.1)	1 (<0.1)
Clinically Relevant Non-Major Bleeding	357 (8.6)	357 (8.7)	0.99 (0.85-1.14) P=0.84	32 (5.4)	7 (1.2)	22 (2.0)	30 (2.7)	20 (1.8)

^a Primary safety outcome for Pooled EINSTEIN DVT and EINSTEIN PE.

^b Primary safety outcome for EINSTEIN Extension and EINSTEIN CHOICE. Major bleeding event was defined as overt bleeding associated with a fall in hemoglobin of 2 g/dL or more; or leading to a transfusion of 2 or more units of packed red blood cells or whole blood; or that occurred in a critical site: intracranial, intraocular, pericardial, intra-articular, intramuscular with compartment syndrome, retroperitoneal; or contributing to death. In EINSTEIN Extension, some patients had more than one event.

See [Table 4](#) for definition of other footnotes.

Clinically relevant non-major bleeding pooled from both EINSTEIN DVT and EINSTEIN PE from a mucosal site occurred in 7.2 % of patients in the rivaroxaban tablets group and 6.0 % of subjects in the enoxaparin/VKA group. Major bleeding from a mucosal site was observed in 0.6 % of the rivaroxaban tablets group and 0.7 % of the enoxaparin/VKA group.

Table 6 – ROCKET AF (Prevention of Stroke and Systemic Embolism in Patients with Atrial Fibrillation (SPAF))–Time to the First Occurrence of Bleeding Events While on Treatment (up to Last Dose Plus 2 Days) - Safety Analysis

	Rivaroxaban tablets	Warfarin	HR (95% CI); P-value
	n (%/year)	n (%/year)	
Major and Non-major Clinically Relevant Bleeding	1475 (14.91)	1449 (14.52)	1.03 (0.96,1.11); 0.442
Major Bleeding ^a	395 (3.60)	386 (3.45)	1.04 (0.90,1.20); 0.576
Hemoglobin Drop	305 (2.77)	254 (2.26)	1.22 (1.03,1.44); 0.019*
Transfusion (> 2 units)	183 (1.65)	149 (1.32)	1.25 (1.01,1.55); 0.044*
Critical Organ Bleed	91 (0.82)	133 (1.18)	0.69 (0.53,0.91); 0.007*
Intracranial Hemorrhage	55 (0.49)	84 (0.74)	0.67 (0.47, 0.94); 0.019*
Fatal Bleed	27 (0.24)	55 (0.48)	0.50 (0.31,0.79); 0.003*
Non-major Clinically Relevant Bleeding	1185 (11.80)	1151 (11.37)	1.04 (0.96,1.13); 0.345

^a See [Table 4](#) and [Table 5](#) for definition of other footnotes.

* Statistically significant at nominal 0.05 (two-sided).

See [Table 28](#), [Table 32](#), and [Table 34](#) for additional details.

Mucosal major bleeding was more common in the rivaroxaban tablets group (2.4%/year) as compared to the warfarin group (1.6%/year; HR 1.52 (1.25, 1.83) $P < 0.001$). Most of the mucosal major bleeding was from a gastrointestinal site.

Intracranial hemorrhage and upper gastrointestinal hemorrhage resulting in death were observed in 24/55 (43.6%) and 1/204 (0.5%) rivaroxaban tablets patients who experienced these adverse events, respectively, compared to 42/84 (50.0%) and 3/125 (2.4%) warfarin patients who experienced these same events, respectively.

8.2 Clinical Trial Adverse Reactions

Clinical trials are conducted under very specific conditions. The adverse reaction rates observed in the clinical trials; therefore, may not reflect the rates observed in practice and should not be compared to the rates in the clinical trials of another drug. Adverse reaction information from clinical trials may be useful in identifying and approximating rates of adverse drug reactions in real-world use.

The most common treatment-emergent adverse events in the three Phase III studies for VTE prevention in elective THR and TKR surgery are presented below in [Table 7](#).

Table 7 – Treatment-Emergent Adverse Drug Reactions Occurring in >1% of Any Treatment Group –Pooled Data of RECORD 1, 2, 3 (VTE Prevention After THR or TKR) – (Patients Valid for Safety Analysis^a)

	Rivaroxaban tablets (N=4571)		Enoxaparin (N=4601)	
	n	(%)	n	(%)
Blood and lymphatic system disorders				
Thrombocytosis (including platelet count increased)	77	(1.68)	86	(1.87)
Gastrointestinal disorders				
Nausea	402	(8.79)	402	(8.74)
Diarrhea	101	(2.21)	134	(2.91)
Abdominal and gastrointestinal pain (including upper abdominal pain, stomach discomfort)	88	(1.93)	88	(1.91)
Dyspepsia (including epigastric discomfort)	40	(0.88)	49	(1.06)
Vomiting	371	(8.12)	392	(8.52)
Constipation	293	(6.41)	319	(6.93)
General Disorders and Administration Site Conditions				
Fever	420	(9.19)	427	(9.28)
Decreased general strength and energy (including asthenia, fatigue)	56	(1.23)	45	(0.98)
Edema peripheral	190	(4.16)	160	(3.48)
Injury, poisoning, and post-procedural complications				
Anemia (including laboratory parameter)	263	(5.75)	292	(6.35)
Post procedural hemorrhage	200	(4.38)	192	(4.17)
Wound secretion	125	(2.73)	92	(2.00)
Investigations				
Increase in LDH	37	(0.81)	49	(1.06)
Increase in transaminases	123	(2.69)	190	(4.13)
Increase in Gamma-glutamyltransferase	74	(1.62)	121	(2.63)
Increase in alkaline phosphatase	35	(0.77)	56	(1.22)
Musculoskeletal, Connective Tissue, and Bone Disorders				
Pain in extremity	74	(1.62)	55	(1.20)
Nervous System Disorders				
Dizziness	149	(3.26)	142	(3.09)
Headache	105	(2.30)	96	(2.09)
Syncope (including loss of consciousness)	71	(1.55)	37	(0.80)
Skin and subcutaneous tissue disorders				
Pruritus (including uncommon cases of generalized pruritus)	97	(2.12)	73	(1.59)
Rash	56	(1.23)	57	(1.24)
Vascular disorders				
Hypotension (including blood pressure decreased)	146	(3.19)	147	(3.19)
Hematoma	47	(1.03)	53	(1.15)

Note: Incidence = number of events/number at risk, where: number of events = number of patients reporting the event; number at risk = number of patients in reference population

Only treatment emergent adverse events which occurred up to 2 days after the last dose of study medication are included.

^a Started after administration of oral study medication (rivaroxaban tablets or matching placebo tablet).

The most common treatment-emergent adverse events reported by patients valid for safety analysis in the 3 Phase III studies for treatment of VTE and prevention of recurrent DVT and PE are presented in [Table 8](#).

Table 8 - Treatment-Emergent Adverse Reactions occurring in >1% of Any Treatment Group – pooled EINSTEIN DVT (11702 DVT) and EINSTEIN PE (11702 PE); EINSTEIN Extension (11899); EINSTEIN CHOICE (16416)^b (Treatment of VTE and Prevention of Recurrent DVT and PE) - Safety Analysis

	Pooled EINSTEIN DVT and EINSTEIN PE		EINSTEIN Extension		EINSTEIN CHOICE		
	Rivaroxaban tablets (N=4130) n (%)	ENOXAPARIN/VKA (N=4116) n (%)	Rivaroxaban tablets (N=598) n (%)	Placebo (N=590) n (%)	Rivaroxaban tablets 10 mg (N=1127) n (%)	Rivaroxaban tablets 20 mg (N=1107) n (%)	ASA 100 mg (N=1131) n (%)
Blood and lymphatic system disorders							
Anemia	84 (2.03)	62 (1.51)	4 (0.67)	2 (0.34)	1 (<0.1)	3 (0.3)	0
Cardiac disorder							
Tachycardia	55 (1.33)	43 (1.04)	2 (0.33)	0	0	1 (<0.1)	0
Eye disorders							
Conjunctival hemorrhage	39 (0.94)	47 (1.14)	6 (1.00)	0	2 (0.2)	6 (0.5)	4 (0.4)
Gastrointestinal disorders							
Gingival bleeding	93 (2.25)	104 (2.53)	11 (1.84)	2 (0.34)	14 (1.2)	28 (2.5)	12 (1.1)
Rectal hemorrhage	90 (2.18)	56 (1.36)	4 (0.67)	4 (0.68)	9 (0.8)	6 (0.2)	7 (0.6)
Abdominal pain	69 (1.67)	53 (1.29)	2 (0.33)	7 (1.19)	1 (<0.1)	3 (0.3)	2 (0.2)
Abdominal pain upper	71 (1.72)	50 (1.21)	10 (1.67)	1 (0.17)	2 (0.2)	2 (0.2)	5 (0.4)
Constipation	187 (4.53)	174 (4.23)	6 (1.00)	5 (0.85)	2 (0.2)	0	7 (0.6)
Diarrhea	179 (4.33)	164 (3.98)	7 (1.17)	8 (1.36)	4 (0.4)	4 (0.4)	1 (<0.1)
Dyspepsia	60 (1.45)	54 (1.31)	8 (1.34)	4 (0.68)	1 (<0.1)	3 (0.3)	4 (0.4)
Nausea	153 (3.70)	160 (3.89)	7 (1.17)	6 (1.02)	3 (0.3)	3 (0.3)	2 (0.2)
Vomiting	69 (1.67)	96 (2.33)	3 (0.50)	6 (1.02)	0	4 (0.4)	2 (0.2)
General disorders and administration site conditions							
Pyrexia	111 (2.69)	108 (2.62)	5 (0.84)	7 (1.19)	1 (<0.1)	2 (0.2)	0
Edema peripheral	128 (3.10)	135 (3.28)	13 (2.17)	17 (2.88)	0	0	1 (<0.1)
Asthenia	61 (1.48)	60 (1.46)	4 (0.67)	6 (1.02)	1 (<0.1)	1 (<0.1)	1 (<0.1)
Fatigue	90 (2.18)	68 (1.65%)	6 (1.00)	3 (0.51)	1 (<0.1)	1 (<0.1)	3 (0.3)
Injury, poisoning and post-procedural complications							
Wound hemorrhage	59 (1.43)	65 (1.58)	11 (1.84)	7 (1.19)	11 (1.0)	11(1.0)	8 (0.7)
Contusion	145 (3.51)	197 (4.79)	19 (3.18)	16 (2.71)	0	2 (0.2)	0
Subcutaneous hematoma	44 (1.07)	61 (1.48)	0	2 (0.34)	33 (2.9)	24 (2.2)	33 (2.9)
Investigations							
Alanine aminotransferase increased ^c	72 (1.74)	129 (3.13)	2 (0.33)	4 (0.68)	-	-	-
Aspartate aminotransferase increased ^c	32 (0.77)	44 (1.07)	4 (0.67)	3 (0.51)	-	-	-
Musculoskeletal, connective tissue and bone disorders							
Pain in extremity	230 (5.57)	221 (5.37)	29 (4.85)	35 (5.93)	4 (0.4)	2 (0.2)	1 (<0.1)

	Pooled EINSTEIN DVT and EINSTEIN PE		EINSTEIN Extension		EINSTEIN CHOICE		
	Rivaroxaban tablets (N=4130)n (%)	ENOXAPARIN/VKA (N=4116) n (%)	Rivaroxaban tablets (N=598) n (%)	Placebo (N=590) n (%)	Rivaroxaban tablets 10 mg (N=1127) n (%)	Rivaroxaban tablets 20 mg (N=1107) n (%)	ASA 100 mg (N=1131) n (%)
Nervous system disorders							
Headache	284 (6.88)	242 (5.88)	18 (3.01)	15 (2.54)	3 (0.3)	4 (0.4)	3 (0.3)
Dizziness	102 (2.47)	108 (2.62)	6 (1.00)	8 (1.36)	5 (0.4)	4 (0.4)	3 (0.3)
Renal and urinary disorders							
Hematuria	111 (2.69)	113 (2.75)	13 (2.17)	2 (0.34)	0	3 (0.3)	0
Reproductive system and breast disorders							
Menorrhagia ^a	122 (2.95)	64 (1.55)	5 (0.84)	2 (0.34)	10 (0.9)	15 (1.4)	2 (0.2)
Vaginal hemorrhage	54 (1.31)	23 (0.56)	1 (0.17)	5 (0.85)	4 (0.4)	5 (0.5)	2 (0.2)
Respiratory, thoracic and mediastinal disorders							
Epistaxis	307 (7.43)	271 (6.58)	24 (4.01)	11 (1.86)	41 (3.6)	41 (3.7)	29 (2.6)
Hemoptysis	100 (2.42)	98 (2.38)	1 (0.17)	1 (0.17)	0	6 (0.5)	1 (<0.1)
Skin and subcutaneous tissue disorders							
Pruritus	83 (2.01)	58 (1.41)	2 (0.33)	2 (0.34)	8 (0.7)	3 (0.3)	3 (0.3)
Rash	97 (2.35)	89 (2.16)	5 (0.84)	7 (1.19)	5 (.4)	3 (0.3)	4 (0.4)
Vascular disorders							
Hematoma	91 (2.20)	150 (3.64)	7 (1.17)	8 (1.36)	0	1 (<0.1)	1 (<0.1)

NB: - Percentages calculated with the number of subjects in each group as denominator

- Incidence is based on number of subjects, not number of events

- Treatment-Emergent (pooled EINSTEIN DVT and EINSTEIN PE) = events that start after randomization and up to 2 days after the last dose of study medication

- Treatment-Emergent (EINSTEIN Extension) = events that start on or after the first dose of study medication and up to 2 days after the last dose of study medication

^a Observed as very common for rivaroxaban in women <55 years in pooled 11702 DVT and 11702 PE studies

^b According to the protocol, a targeted AE reporting was applied in this study, i.e. all serious adverse events (SAEs), all AEs of special interest, independent if serious or not, all non-serious AEs leading to a permanent study medication discontinuation, and all pregnancies (and their outcomes) in a patient or of the patient's partner needed to be captured on the eCRF and were reported to PV within 24 hours. Investigators could collect AEs on the eCRF, if deemed important.

^c As laboratory measurements related to AST/ALT in Einstein CHOICE were not scheduled but performed as needed, the information is not available.

The most common identified treatment-emergent adverse drug reactions in the pivotal Phase III study, ROCKET AF, for prevention of stroke and systemic embolism in patients with atrial fibrillation are presented in [Table 9](#).

Table 9 – Treatment-Emergent Adverse Reactions Occurring in >1% of Any Treatment Group – ROCKET AF (Prevention of Stroke and Systemic Embolism in Patients with Atrial Fibrillation (SPAF)) - Safety Analysis

	Rivaroxaban Tablets (N=7111)		Warfarin (N=7125)	
	n	(%)	n	(%)
Blood and lymphatic system disorders				
Anemia	219	(3.08)	143	(2.01)
Eye disorders				
Conjunctival hemorrhage	104	(1.46)	151	(2.12)
Gastrointestinal disorders				
Diarrhea	379	(5.33)	397	(5.57)
Gingival bleeding	263	(3.70)	155	(2.18)
Nausea	194	(2.73)	153	(2.15)
Rectal hemorrhage	149	(2.10)	102	(1.43)
Abdominal pain upper	127	(1.79)	120	(1.68)
Vomiting	114	(1.60)	111	(1.56)
Dyspepsia	111	(1.56)	91	(1.28)
Abdominal pain	107	(1.50)	118	(1.66)
Gastrointestinal hemorrhage	100	(1.41)	70	(0.98)
General Disorders and Administration Site Conditions				
Edema peripheral	435	(6.12)	444	(6.23)
Fatigue	223	(3.14)	221	(3.10)
Asthenia	125	(1.76)	106	(1.49)
Pyrexia	72	(1.01)	87	(1.22)
Injury, poisoning and post-procedural complications				
Contusion	196	(2.76)	291	(4.08)
Investigations				
Alanine aminotransferase increased	144	(2.03)	112	(1.57)
Musculoskeletal, Connective Tissue, and Bone Disorders				
Pain in extremity	191	(2.69)	208	(2.92)
Nervous System Disorders				
Dizziness	433	(6.09)	449	(6.30)
Headache	324	(4.56)	363	(5.09)
Syncope	130	(1.83)	108	(1.52)
Renal and urinary disorders				
Hematuria	296	(4.16)	242	(3.40)
Respiratory tract disorders				
Epistaxis	721	(10.14)	609	(8.55)
Hemoptysis	99	(1.39)	100	(1.40)
Skin and subcutaneous tissue disorders				
Ecchymosis	159	(2.24)	234	(3.28)
Pruritus	120	(1.69)	118	(1.66)
Rash	112	(1.58)	129	(1.81)
Vascular disorders				
Hematoma	216	(3.04)	330	(4.63)
Hypotension	141	(1.98)	130	(1.82)

NB: Incidence is based on number of subjects, not number of events

Treatment-Emergent = events that start on or after the first dose of study medication and up to 2 days after the last dose of study medication

8.3 Less Common Clinical Trial Adverse Reactions

Incidence is $\geq 0.1\%$ to $< 1\%$ unless specified.

VTE Prevention in Elective THR and TKR Surgery

Cardiac Disorders: tachycardia

Gastrointestinal Disorders: dry mouth, gastrointestinal tract hemorrhage (including gingival bleeding, rectal hemorrhage, hematemesis)

General Disorders and Administration Site Conditions: feeling unwell (including malaise), localized edema

Hepatobiliary Disorders: hepatic impairment ($\geq 0.01\%$ to $< 0.1\%$)

Immune System Disorders: hypersensitivity, anaphylaxis, allergic edema and angioedema, dermatitis allergic

Investigations: bilirubin conjugated increased (with or without concomitant increase of ALT) ($\geq 0.01\%$ to $< 0.1\%$), blood bilirubin increased, increased amylase, increased lipase

Renal and Urinary Disorders: renal impairment (including serum creatinine increased, blood urea increased)

Respiratory Tract Disorders: epistaxis

Skin and Subcutaneous Tissue Disorders: contusion, urticaria (including rare cases of generalized urticaria)

Vascular Disorders: urogenital tract hemorrhage

Treatment of VTE and Prevention of Recurrent DVT and PE

Incidence is $\geq 0.1\%$ to $< 1\%$ (pooled EINSTEIN DVT, EINSTEIN PE and EINSTEIN Extension) unless specified. Patients rolled over from EINSTEIN DVT or EINSTEIN PE into EINSTEIN Extension are considered as one patient (N=4556).

Cardiac Disorder: tachycardia

Gastrointestinal Disorders: gastrointestinal hemorrhage, hematochezia, hemorrhoidal hemorrhage, melena, mouth hemorrhage, abdominal discomfort, abdominal pain lower, dry mouth

General Disorders and Administration Site Conditions: asthenia, feeling abnormal, malaise

Hepatobiliary Disorders: hepatic impairment

Immune System Disorders: hypersensitivity

Injury, Poisoning and Post-procedural Complications: post-procedural hemorrhage, traumatic hematoma, traumatic hemorrhage, subcutaneous hematoma

Investigations: hemoglobin decreased, aspartate aminotransferase increased, liver function test abnormal, hepatic enzyme increased, transaminases increased, blood bilirubin increased, bilirubin conjugated increased (with or without concomitant increase of ALT), gamma-glutamyl transferase increased, blood alkaline phosphatase increased

Nervous System Disorders: syncope, cerebral and intra cranial hemorrhage ($\geq 0.01\%$ to $< 0.1\%$)

Reproductive System and Breast Disorders: menometrorrhagia, metrorrhagia

Skin and Subcutaneous Tissue Disorders: urticaria, ecchymosis, skin hemorrhage, dermatitis allergic

(≥0.01% to <0.1%)

Vascular Disorders: hypotension

In other clinical studies with rivaroxaban tablets, occurrences of vascular pseudoaneurysm formation following percutaneous intervention have been observed. Very rare cases of adrenal hemorrhage have been reported.

Prevention of Stroke and Systemic Embolism in Patients with Atrial Fibrillation (SPAF)

Cardiac Disorders: tachycardia

Eye Disorders: eye hemorrhage, vitreous hemorrhage

Gastrointestinal Disorders: melena, upper gastrointestinal hemorrhage, hemorrhoidal hemorrhage, hematochezia, mouth hemorrhage, lower gastrointestinal hemorrhage, anal hemorrhage, gastric ulcer hemorrhage, gastritis hemorrhagic, gastric hemorrhage, hematemesis, abdominal discomfort, abdominal pain lower, dry mouth

General Disorders and Administration Site Conditions: malaise

Hepatobiliary Disorders: hepatic impairment, hyperbilirubinemia, jaundice (≥ 0.01% to < 0.1%)

Immune System Disorders: hypersensitivity, anaphylaxis (≥ 0.01% to < 0.1%), allergic edema and angioedema

Injury, Poisoning, and Post-procedural Complications: post-procedural hemorrhage, wound hemorrhage, traumatic hematoma, incision site hemorrhage, subdural hematoma, subcutaneous hematoma, periorbital hematoma

Investigations: hemoglobin decreased, hematocrit decreased, blood bilirubin increased, liver function test abnormal, aspartate aminotransferase increased, hepatic enzyme increased, blood urine present, creatinine renal clearance decreased, blood creatinine increased, blood urea increased, blood alkaline phosphatase increased, lipase increased, bilirubin conjugated increased (with or without concomitant increase of ALT) (≥ 0.01% to < 0.1%)

Musculoskeletal, Connective Tissue, and Bone Disorders: hemarthrosis, muscle hemorrhage (≥ 0.01% to < 0.1%)

Nervous System Disorders: loss of consciousness, hemorrhagic stroke, hemorrhage intracranial

Renal and Urinary Disorders: renal impairment

Reproductive System Disorders: vaginal hemorrhage, metrorrhagia

Skin and Subcutaneous Tissue Disorders: dermatitis allergic, rash pruritic, rash erythematous, rash generalized, pruritus generalized, urticaria, skin hemorrhage

Vascular Disorders: hemorrhage, bleeding varicose vein

8.4 Abnormal Laboratory Findings: Hematologic, Clinical Chemistry and Other Quantitative Data Clinical Trial Findings

In Phase III clinical trials, in VTE prevention, Treatment of VTE and prevention of recurrent DVT and PE, and SPAF the incidence of increases in transaminases in the rivaroxaban tablets and comparator arms were similar, see [Table 7](#), [Table 8](#), and [Table 9](#) above.

8.5 Post-Market Adverse Reactions

The following adverse reactions have been identified during post-approval use of rivaroxaban tablets. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

Blood and the Lymphatic System Disorders: agranulocytosis, thrombocytopenia

Hepatobiliary Disorders: cholestasis, hepatitis (including hepatocellular injury)

Immune System Disorders: anaphylaxis, allergic edema and angioedema (with or without urticaria)

Respiratory, Thoracic and Mediastinal Disorders: eosinophilic pneumonia

Skin and Subcutaneous Tissue Disorders: Stevens-Johnson syndrome, drug reaction with eosinophilia and systemic symptoms (DRESS)

9 DRUG INTERACTIONS

9.2 Drug Interactions Overview

Drug interaction studies have only been performed in adults. (see [7 WARNINGS AND PRECAUTIONS – Drug Interactions](#)).

Rivaroxaban neither inhibits nor induces CYP 3A4 or any other major CYP isoenzymes.

Concomitant use of drugs affecting hemostasis increases the risk of bleeding. Care should be taken if patients are treated concomitantly with drugs affecting hemostasis such as nonsteroidal anti-inflammatory drugs (NSAIDs), acetylsalicylic acid, and platelet aggregation inhibitors. Due to the increased bleeding risk, generally avoid concomitant use with other anticoagulants (see [7 WARNINGS AND PRECAUTIONS – Bleeding](#)).

9.4 Drug-Drug Interactions

The use of JAMP Rivaroxaban is contraindicated in patients receiving concomitant systemic treatment with strong inhibitors of both CYP 3A4 and P-gp such as cobicistat, ketoconazole, itraconazole, posaconazole, or ritonavir). These drugs may increase rivaroxaban tablets plasma concentrations to a clinically relevant degree, i.e, 2.6-fold on average, which may lead to bleeding. Theazole anti-mycotic, fluconazole, a moderate CYP 3A4 inhibitor, has less effect on rivaroxaban exposure and may be co-administered with caution (see [2 CONTRAINDICATIONS](#), and [7 WARNINGS AND PRECAUTIONS – Drug Interactions](#)).

In the ROCKET AF clinical trial in patients with atrial fibrillation, no apparent increase in major bleeding was observed in patients in whom amiodarone, a moderate CYP 3A4 inhibitor, was co-administered with rivaroxaban.

Drugs strongly inhibiting only one of the rivaroxaban tablets elimination pathways, either CYP 3A4 or P-gp, are expected to increase rivaroxaban tablets plasma concentrations to a lesser extent. The expected increase is considered less clinically relevant, see [Table 10](#).

Table 10 – Established or Potential Drug-Drug Interactions

Concomitant Drug Class: Drug Name	Reference	Effect	Clinical Comment
Azole antimycotic: ketoconazole	CT	Co-administration of rivaroxaban tablets with the azole-antimycotic ketoconazole(400 mg od) a strong CYP 3A4 and P-gp inhibitor, led to a 2.6-fold increase in mean rivaroxaban tablets steady state AUC and a 1.7-fold increase in mean rivaroxaban tablets Cmax, with significant increases in its pharmacodynamic effects.	The use of JAMP Rivaroxaban is contraindicated in patients receiving systemic treatment with ketoconazole (see 2 CONTRAINDICATIONS and 7 WARNINGS AND PRECAUTIONS – Drug Interactions and Renal Impairment).
Cobicistat	C	Cobicistat is a strong CYP 3A4 and P-gp inhibitor. Coadministration with cobicistat may result in increased plasma concentration of rivaroxaban, leading to an increased bleeding risk.	The use of JAMP Rivaroxaban is contraindicated in patients receiving systemic treatment with cobicistat (see 2 CONTRAINDICATIONS and 7 WARNINGS AND PRECAUTIONS – Drug Interactions).
Dronedarone	C	In a retrospective cohort study using the Truven Health MarketScan database in the US, a significantly increased risk of ICD-diagnoses of bleeding leading to hospitalization or emergency department visit was observed, driven by gastrointestinal bleeding, in NVAf patients with the concomitant use of rivaroxaban and dronedarone compared to those taking rivaroxaban alone.	Dronedarone should not be used concomitantly with JAMP Rivaroxaban since it may increase exposure of JAMP Rivaroxaban through P-gp and CYP3A4 inhibition, and thereby the risk of bleeding.
fluconazole	CT	Administration of the moderate CYP 3A4 inhibitor fluconazole (400 mg once daily) led to a 1.4-fold increase in mean rivaroxaban tablets AUC and a 1.3-fold increase in mean Cmax.	No dose adjustment is required.
Protease inhibitor: ritonavir	CT	Co-administration of rivaroxaban tablets with the HIV protease inhibitor ritonavir (600 mg bid), a strong CYP 3A4 and P-gp inhibitor, led to a 2.5-fold increase in mean rivaroxaban tablets AUC and a 1.6-fold increase in mean rivaroxaban tablet Cmax, with significant increases in its pharmacodynamic effects.	The use of JAMP Rivaroxaban is contraindicated in patients receiving systemic treatment with ritonavir (see 2 CONTRAINDICATIONS and 7 WARNINGS AND PRECAUTIONS – Drug Interactions and Renal Impairment).

Concomitant Drug Class: Drug Name	Reference	Effect	Clinical Comment
Anti-infectives: erythromycin	CT	Erythromycin (500 mg tid), which inhibits CYP 3A4 and P-gp moderately, led to a 1.3-fold increase in mean rivaroxaban tablets AUC and C _{max} .	No dose adjustment is required. For patients with renal impairment see 7 WARNINGS AND PRECAUTIONS – Drug Interactions and 10.3 Pharmacokinetics – Special Populations and Conditions – Renal Insufficiency .
clarithromycin	CT	Clarithromycin (500 mg bid), considered a strong CYP 3A4 inhibitor and moderate P-gp inhibitor, led to a 1.5-fold increase in mean rivaroxaban, and a 1.4-fold increase in C _{max} .	The use of JAMP Rivaroxaban in combination with clarithromycin may increase the risk of bleeding particularly in patients with underlying disease conditions, and elderly. Caution is required.
rifampicin	CT	Co-administration of rivaroxaban tablets with the strong CYP 3A4 and P-gp inducer rifampicin led to an approximate 50% decrease in mean rivaroxaban tablets AUC, with parallel decreases in its pharmacodynamic effects.	Strong CYP 3A4 inducers should generally be avoided in combination with JAMP Rivaroxaban, as such use can be expected to result in inadequate anticoagulation.
Anticonvulsants: phenytoin carbamazepine phenobarbital	T	The concomitant use of rivaroxaban tablets with strong CYP 3A4 inducers (e.g, phenytoin, carbamazepine, or phenobarbital) may also lead to a decreased rivaroxaban tablets plasma concentration.	Strong CYP 3A4 inducers should generally be avoided in combination with JAMP Rivaroxaban, as such use can be expected to result in inadequate anticoagulation.
Nonsteroidal Anti-inflammatory Drugs (NSAID): naproxen	CT	Co-administration with naproxen did not affect rivaroxaban tablets bioavailability and pharmacokinetics. No clinically relevant prolongation of bleeding time was observed when 500 mg naproxen was pre-administered 24 hours before concomitant administration of single doses of rivaroxaban tablets 15 mg and naproxen 500 mg in healthy subjects.	Concomitant use with JAMP Rivaroxaban may increase the risk of bleeding. Promptly evaluate any signs or symptoms of blood loss (see 7 WARNINGS AND PRECAUTIONS – Drug Interactions).
acetylsalicylic acid (ASA)	CT	No clinically significant pharmacokinetic or pharmacodynamic interactions were observed when 500 mg ASA was pre-administered 24 hours before concomitant administration of single doses of rivaroxaban tablets 15 mg and ASA 100 mg in healthy subjects.	Concomitant use with JAMP Rivaroxaban increases the risk of bleeding. Promptly evaluate any signs or symptoms of blood loss (see 7 WARNINGS AND PRECAUTIONS – Drug Interactions). For patients in the ROCKET AF trial, concomitant ASA use (almost exclusively at 100 mg or less) was identified as an independent risk factor for major bleeding with both

Concomitant Drug Class: Drug Name	Reference	Effect	Clinical Comment
			rivaroxaban tablets and warfarin.
Antiplatelet drugs: clopidogrel	CT	In two drug interaction studies of 11 and 13 healthy subjects, clopidogrel 300 mg was pre-administered 24 hours before concomitant administration of single doses of rivaroxaban tablets 15 mg and clopidogrel 75 mg in healthy subjects. Clopidogrel with or without rivaroxaban tablets led to an approximately 2-fold increase in the median bleeding time (normal range 2 - 8 minutes). In these studies, between 30% and 40% of subjects who received both rivaroxaban tablets and clopidogrel had maximum bleeding times of up to 45 minutes. Rivaroxaban tablets alone did not lead to a change in bleeding time at 4 hours or 2 days after administration. There was no change in the pharmacokinetics of either drug.	Concomitant use with JAMP Rivaroxaban increases the risk of bleeding. Promptly evaluate any signs or symptoms of blood loss (see 7 WARNINGS AND PRECAUTIONS – Drug Interactions).
Antithrombotic: enoxaparin	CT	After combined administration of enoxaparin (40 mg single dose) with rivaroxaban tablets (10 mg single dose), an additive effect on anti-Factor-Xa activity was observed, without any additional effects on clotting tests (PT, aPTT). Enoxaparin did not affect the bioavailability and pharmacokinetics of rivaroxaban tablets.	Co-administration of rivaroxaban tablets at doses ≥ 10 mg with other anticoagulants or antithrombotic therapy has not been adequately studied in clinical trials. Due to the increased bleeding risk, generally avoid concomitant use with other anticoagulants (see 7 WARNINGS AND PRECAUTIONS – Drug Interactions).
Selective serotonin reuptake inhibitors (SSRI), and serotonin norepinephrine reuptake inhibitors (SNRIs)	T, CT	When concomitantly used in the rivaroxaban tablets clinical program, numerically higher rates of major or non-major clinically relevant bleeding were observed	As with other anticoagulants, patients on rivaroxaban tablets are at increased risk of bleeding in case of concomitant use with SSRIs or SNRIs due to their reported effect on platelets.

Legend: C = Case Study; CT=Clinical Trial; T=Theoretical

No pharmacokinetic interaction was observed between warfarin and rivaroxaban tablets.

There were no mutual pharmacokinetic interactions observed between rivaroxaban tablets and midazolam (substrate of CYP 3A4), digoxin (substrate of P-gp), or atorvastatin (substrate of CYP 3A4 and P-gp).

Co-administration of the proton pump inhibitor, omeprazole, the H2-receptor antagonist, ranitidine, the antacid, aluminum hydroxide / magnesium hydroxide, or naproxen, clopidogrel, or enoxaparin did not

affect rivaroxaban tablets bioavailability or pharmacokinetics.

9.5 Drug-Food Interactions

JAMP Rivaroxaban 10 mg may be taken with or without food. JAMP Rivaroxaban 15 mg and 20 mg should be taken with food. (see [10.3 Pharmacokinetics](#)).

Grapefruit juice is a moderate CYP 3A4 inhibitor. Therefore, an increase in rivaroxaban tablets exposure following grapefruit juice consumption is not expected to be clinically relevant.

9.6 Drug-Herb Interactions

The concomitant use of rivaroxaban tablets with strong CYP 3A4 inducers (e.g. St. John's Wort) may lead to a decreased rivaroxaban tablets plasma concentration. Strong CYP 3A4 inducers should generally be avoided in combination with JAMP Rivaroxaban, as such use can be expected to result in inadequate anticoagulation.

9.7 Drug-Laboratory Test Interactions

Although various clotting parameter tests (PT, aPTT, Heptest®) are affected by the mode of action of rivaroxaban tablets, none of these clotting tests have been demonstrated to reliably assess the anticoagulant activity of rivaroxaban following rivaroxaban tablets administration under usual conditions (see [7 WARNINGS AND PRECAUTIONS– Monitoring and Laboratory Tests](#), and [10.2 Pharmacodynamics](#)).

The prothrombin time (PT), measured in seconds, is influenced by rivaroxaban tablets in a dose-dependent way with a close correlation to plasma concentrations if the Neoplastin® reagent is used. In patients who are bleeding, measuring the PT (Neoplastin® reagent) in seconds, but not INR, may be useful to assist in determining an excess of anticoagulant activity (see [7 WARNINGS AND PRECAUTIONS – Monitoring and Laboratory Tests](#)).

10 CLINICAL PHARMACOLOGY

10.1 Mechanism of Action

Rivaroxaban is a highly selective, direct, antithrombin independent Factor-Xa inhibitor with high oral bioavailability.

Activation of Factor-X to Factor-Xa (FXa) via the intrinsic and extrinsic pathway plays a central role in the cascade of blood coagulation. FXa directly converts prothrombin to thrombin through the prothrombinase complex and, ultimately, this reaction leads to fibrin clot formation and activation of platelets by thrombin. One molecule of FXa is able to generate more than 1000 molecules of thrombin due to the amplification nature of the coagulation cascade. In addition, the reaction rate of prothrombinase-bound FXa increases 300,000-fold compared to that of free FXa and causes an explosive burst of thrombin generation. Selective inhibitors of FXa can terminate the amplified burst of thrombin generation, thereby diminishing thrombin-mediated activation of coagulation.

10.2 Pharmacodynamics

There is a clear correlation between plasma rivaroxaban concentration and the degree of anticoagulant effect. The maximal effect (E_{max}) of rivaroxaban on pharmacodynamic parameters occurs at the same time as C_{max} .

- A dose-dependent inhibition of Factor-Xa (FXa) activity was observed over the complete dose range closely following the pharmacokinetic profiles which provides the 'proof of mechanism' in humans. Inhibition of FXa activity versus rivaroxaban plasma concentration follows a maximum effect (E_{max})

model. There is a close correlation between FXa inhibition and plasma concentrations with an r value of 0.97.

FXa assay tests require calibration and should not be used unless rivaroxaban-specific calibrators and controls are available.

- Prothrombin time (PT), measured in seconds, is influenced by rivaroxaban in a dose-dependent way with a close correlation to plasma concentrations ($r = 0.98$) if the Neoplastin® reagent is used. Other reagents would provide different results.
- Although JAMP Rivaroxaban therapy will lead to an elevated INR, depending on the timing of the measurement, the INR is not a valid measure to assess the anticoagulant activity of JAMP Rivaroxaban. The INR is only calibrated and validated for VKA and should not be used for any other anticoagulant (see [7 WARNINGS AND PRECAUTIONS – Monitoring and Laboratory Tests](#)).

In patients who are bleeding, measuring the PT (Neoplastin® reagent) may be useful to assist in determining an excess of anticoagulant activity (see [7 WARNINGS AND PRECAUTIONS – Monitoring and Laboratory Tests](#)).

Figure 1 and Figure 2 below show the relative measured effects of rivaroxaban 20 mg once daily for the PT test using the Neoplastin® reagent (Figure 1) and that expressed by the INR (Figure 2).

Figure 1: PT Prolongation (Neoplastin® reagent): Relative prolongation expressed as median of ratio to baseline with warfarin / rivaroxaban treatment and rivaroxaban alone, following last day of warfarin (Day -1) and 4 days of 20 mg rivaroxaban od, PD set, n=84





Figure 2: INR prolongation: Relative prolongation expressed as median of ratio to baseline with warfarin/ rivaroxaban treatment and rivaroxaban alone, following last day of warfarin (Day -1) and 4 days of 20 mg rivaroxaban od, PK/PD set, n=84



The usual expected effect of rivaroxaban tablets on PT when the Neoplastin® reagent is used is shown in [Table 11](#) below.

Table 11 – PT (Neoplastin® reagent) by Indication, Following rivaroxaban tablets Administration

Indication	Rivaroxaban Tablets Dosage	Plasma concentration C _{max} (mcg/L)	Plasma concentration C _{trough} (mcg/L)	Range of (5/95 percentile) PT (Neoplastin®) C _{max}	Range of (5/95 percentile) PT (Neoplastin®) C _{trough}
Prevention of VTE After THR or TKR	10 mg od	101 (7 – 273) ^a	14 (4 – 51) ^c	13 to 25 seconds ^a	12-17 seconds ^c
Treatment of VTE and Prevention of Recurrent DVT and PE	15 mg bid	---	---	17 to 32 seconds ^a	14–24 seconds ^c
	20 mg od	215 (22– 535) ^a	32 (6–239) ^d	15 to 30 seconds ^a	13–20 seconds ^d
Prevention of Stroke in Patients with Atrial Fibrillation	15 mg od	229 (178 – 313) ^b	57 (18 – 136) ^e	10 to 50 seconds ^b	12–26 seconds ^e
	20 mg od	249 (184 – 343) ^b	44 (12 – 137) ^e	14 to 40 seconds ^b	11–26 seconds ^e

^a 2 to 4 hours after drug administration (t_{max})

^b 1 to 4 hours after drug administration (t_{max})

^c 8 to 16 hours after drug administration (t_{min})

^d 18 to 30 hours after drug administration (t_{min})

^e 16 to 32 hours after drug administration (t_{min})

- The activated partial thromboplastin time (aPTT) is prolonged dose-dependently; however, the slope is rather flat and does not allow a sufficient discrimination at the relevant plasma concentrations. Therefore, aPTT is not considered to be adequate for following the pharmacodynamic effects. The r value for aPTT is 0.99.
- Heptest® is prolonged dose-dependently and correlates closely with plasma concentrations, following a curvilinear model. Despite the r value of 0.99 for the relation to plasma concentrations, the Heptest® is not considered optimal to assess the pharmacodynamic effects due to the curvilinear relationship.

QT Prolongation

No QTc prolonging effects were observed in healthy men and women older than 50 years. The treatment difference in QTcF 3 hours post-dose in comparison to placebo as well as QTcF, QTcI and QT analyses at the time of t_{max} and for post-dose changes in mean and maximum QTcF did not show any dose-related QTcF prolongation at both the 45 mg and the 15 mg dose of rivaroxaban. All changes in LS-means, including their 95% CI, were below 5 milliseconds.

Patients undergoing cardioversion

A prospective, randomized, open-label, multicenter, exploratory study with blinded endpoint evaluation (X-VerT) was conducted in 1504 patients with non-valvular atrial fibrillation scheduled for cardioversion to compare rivaroxaban with dose-adjusted VKA (randomized 2:1). The rate of stroke occurring within 42 days of cardioversion was low and similar across treatment groups, i.e., rivaroxaban (0.20%) and VKA (0.41%). The rate of major bleeding was also low and similar across treatment groups, i.e., rivaroxaban (0.61%) and VKA (0.80%).

Patients with nonvalvular atrial fibrillation who undergo PCI with stent placement

In a randomized, open label, multicentre study (PIONEER AF-PCI) in patients with nonvalvular atrial fibrillation who underwent PCI with stent placement for primary atherosclerotic disease, the 12-month safety of two antithrombotic regimens was compared. One group of 696 patients received rivaroxaban 15 mg o.d. (10 mg o.d. in patients with CrCl 30-49 mL/min) in combination with a P2Y12 inhibitor (e.g, clopidogrel), while a second group of 697 patients received dose-adjusted VKA plus DAPT. Patients with a history of stroke or TIA were excluded from the trial.

The primary safety endpoint, clinically significant bleeding events [a composite of TIMI major bleeding, TIMI minor bleeding and Bleeding Requiring Medical Attention (BRMA)] occurred in 109 patients (15.7%) on the rivaroxaban regimen and in 167 patients (24.0%) on the VKA regimen (HR 0.59; 95% CI 0.47-0.76; $p < 0.001$). This difference in bleeding risk was primarily a result of significantly fewer BRMA events in patients on the rivaroxaban regimen. While a consistent treatment effect for all 3 components of the composite was observed, the low number of TIMI major and TIMI minor bleeding events during the trial prevented the demonstration of a significant difference between the two regimens for these endpoints. The secondary endpoint, a composite of CV death, MI or stroke, occurred in 41 patients (5.9%) on rivaroxaban and in 36 patients (5.2%) on VKA; stent thrombosis occurred in 5 patients on rivaroxaban and in 4 patients on VKA. The study was not designed to compare efficacy between the treatment arms, preventing any conclusions regarding efficacy.

10.3 Pharmacokinetics

Rivaroxaban pharmacokinetics are linear with no relevant accumulation beyond steady state after multiple doses. Variability in pharmacokinetics is moderate with interindividual variability (coefficient of variation) ranging from 30% to 40%.

Absorption

Rivaroxaban is a low solubility, high permeability compound. Rivaroxaban is readily absorbed after oral administration as tablet (C_{max} after 2 to 4 hours).

Oral bioavailability of rivaroxaban is high (80-100%) due to almost complete absorption with/without food (at doses up to 15 mg) and lack of relevant presystemic first-pass extraction of this low-clearance drug.

The following information is based on the data obtained in adults.

The absolute bioavailability of rivaroxaban is approximately 100% for doses up to 10 mg. Rivaroxaban is rapidly absorbed with maximum concentrations (C_{max}) appearing 2 to 4 hours after tablet intake.

Intake with food does not affect rivaroxaban AUC or C_{max} for doses up to 10 mg. Rivaroxaban 10mg tablets can be taken with or without food. Due to reduced extent of absorption an oral bioavailability of 66% was determined for the 20 mg tablet under fasting conditions. When rivaroxaban 20 mg tablets are taken together with food, increases in mean AUC by 39% and mean C_{max} by 76% were observed when compared to tablet intake under fasting conditions, indicating almost complete absorption and high oral bioavailability when this dose was taken with food.

The bioavailability of rivaroxaban 10 mg, 15 mg and 20 mg tablets under fed conditions, and 10 mg tablets under fasted conditions, demonstrated dose-proportionality. JAMP Rivaroxaban 15 mg and 20 mg tablets should be taken with food. (see [4 DOSAGE AND ADMINISTRATION](#)).

Rivaroxaban pharmacokinetic parameters behave in a linear fashion; no evidence of undue accumulation beyond steady-state was seen after multiple doses.

Interindividual variability (CV%) of rivaroxaban pharmacokinetics ranges from 30% to 40%. This may be increased on the day of surgery and on the following day when interindividual variability is 70%.

Table 12 – Summary of PK Parameters After Oral Administration of 10 mg of Rivaroxaban in Humans

	C_{max} [mcg/L]	$t_{1/2}$ [h]	AUC [mcg*h/L]	Clearance, Urinary Excretion	Volume of Distribution
Healthy (Young) Subjects	approx. 114 ^a	5-9	approx. 817	CL _{sys} = approx. 10 L/h CL _R = 3 – 4 L/h Ae _{ur} = 30% - 40%	V _{ss} = approx. 50 L
Patients	approx. 125	7-11	approx. 1170	N/A (no IV data) ^b Ae _{ur} = 22%	N/A (no IV data)

^a = 2 – 4 hours after drug administration (t_{max})

^b = not available

AUC = area under the plasma-concentration time curve; Ae_{ur} = amount of drug excreted unchanged into urine; CL_{sys} = systemic clearance (after intravenous administration); CL_R = renal clearance; C_{max} = maximum plasma concentration; $t_{1/2}$ = terminal elimination half-life; t_{max} = time to reach C_{max}; V_{ss} = volume of distribution at steady state

Absorption of rivaroxaban is dependent on the site of drug release in the GI tract. A 29% and 56% decrease in AUC and C_{max} compared to orally ingested tablet was reported when rivaroxaban granulate is released in the proximal small intestine. Exposure is further reduced when drug is released in the distal small intestine, or ascending colon. Avoid administration of rivaroxaban distal to the stomach as this can result in reduced absorption and related drug exposure.

In an open-label, randomized, 3-period, 3-treatment crossover comparative bioavailability study conducted in 44 healthy male and female subjects, the bioavailability (AUC_T and C_{max}) of rivaroxaban following a single 20 mg dose as a crushed 20 mg tablet mixed in applesauce and administered orally, or as a crushed 20 mg tablet suspended in water and administered via NG tube was comparable to a whole 20 mg tablet administered orally. Each rivaroxaban treatment was taken with a standardized liquid meal. Given, the predictable, dose-proportional pharmacokinetic profile of rivaroxaban, the bioavailability results from this study are likely applicable to lower rivaroxaban doses.

Distribution

Plasma protein binding for rivaroxaban in humans is high at approximately 92% to 95% in vitro, with serum albumin being the main binding component. No concentration dependency and no gender difference in fraction unbound were detected. Mean rivaroxaban protein-bound fractions determined *ex vivo* in healthy subjects ranged from 90% to 95%. The volume of distribution is moderate with V_{ss} being approximately 50 L.

Due to its high plasma protein binding, rivaroxaban is not expected to be removed by dialysis.

The binding of rivaroxaban to plasma proteins is fully reversible. In accordance with other species, rivaroxaban is mainly located in plasma; the human plasma-to-blood partition coefficient is 1.40.

Metabolism

Rivaroxaban is eliminated by metabolic degradation (approximately 2/3 of administered dose) as well as by direct renal excretion of unchanged active compound (approximately 1/3 of administered dose). In all investigated species, the oxidative degradation of the morpholinone moiety (catalyzed via CYP 3A4/CYP 3A5 and CYP 2J2 and leading via cleavage of the ring and further oxidation to metabolite M-1) was the

major site of biotransformation of rivaroxaban. Unchanged rivaroxaban is the most important compound in human plasma with no major or active circulating metabolites being present. No metabolic conversion of rivaroxaban to its enantiomer was observed in humans.

Taking excretion data and metabolite profiles derived from the mass balance study in man into consideration, present data from the CYP reaction phenotyping study suggests that contribution of CYP 3A4/CYP 3A5 accounts for approximately 18% and CYP 2J2 for approximately 14% of total rivaroxaban elimination, respectively. Besides this oxidative biotransformation, hydrolysis of the amide bonds (approximately 14%) and active, transporter-mediated renal excretion of unchanged drug (approximately 30%) play important roles as elimination pathways.

Elimination

Rivaroxaban and its metabolites have a dual route of elimination, via both renal (66% in total) and biliary/fecal routes; 36% of the administered dose is excreted unchanged via the kidneys via glomerular filtration and active secretion.

The clearance and excretion of rivaroxaban is as follows:

- 1/3 of the active drug is cleared as unchanged drug by the kidneys
- 1/3 of the active drug is metabolized to inactive metabolites and then excreted by the kidneys
- 1/3 of the active drug is metabolized to inactive metabolites and then excreted by the fecal route.

Rivaroxaban has been identified in vitro to be a substrate both of the active transporter P glycoprotein (P-gp) and of the multidrug transport protein BCRP ('breast cancer resistance protein').

With an average systemic plasma clearance of approximately 10 L/h, rivaroxaban is a low-clearance drug lacking relevant first-pass extraction. Mean terminal elimination half-lives of rivaroxaban are in the range of 5 h to 9 h after steady-state tablet dosing regimens in young subjects. Mean terminal elimination half-lives between 11 h to 13 h were observed in the elderly.

Special Populations and Conditions

Geriatrics (> 65 Years of Age)

Clinical studies have been conducted in older ages, with results of prolonged terminal half-lives (11 to 13 hours in elderly versus 5 to 9 hours in young subjects) accompanied by increases of rivaroxaban tablets exposure (approximately 50%) compared to young healthy subjects. This difference may be due to reduced renal function in the elderly (see [2 CONTRAINDICATIONS](#), [7 WARNINGS AND PRECAUTIONS – Renal Impairment](#), and [4 DOSAGE AND ADMINISTRATION – Renal Impairment](#)).

Results from a set of Phase I studies indicate for the target population of elderly higher mean AUC values by 52% in males and by 39% in females when compared to young subjects of the same gender, accompanied by an increase in C_{max} by 35% in both genders and by terminal half-lives between 11 and 13 h. Investigating subjects older than 75 years confirmed the expectation, leading to approximately 41% higher AUC values in comparison to young subjects (90% CI [1.20 – 1.66]), mainly due to reduced (apparent) total body clearance and renal clearance. No relevant age effects could be observed for C_{max} (C_{max} ratio 1.08; 90% CI [0.94-1.25]) or t_{max} .

Pediatrics (< 18 Years of Age)

PT (Neoplastin® reagent), aPTT, and anti-factor Xa activity by a calibrated quantitative test) display a

close correlation to rivaroxaban plasma concentrations in children. The correlation between anti-factor Xa to plasma concentrations is linear with a slope close to 1. The lower limit of quantifications must be considered when the anti-factor Xa test is used to quantify plasma concentrations of rivaroxaban in children.

Sex

There were no relevant differences in pharmacokinetics and pharmacodynamics between male and female subjects, especially when taking into account body weight differences. An exploratory analysis did not reveal relevant differences in rivaroxaban exposure between male and female children.

Body Weight

In adults, extremes in body weight (<50 kg vs >120 kg) had only a small influence on rivaroxaban plasma concentrations (less than 25%).

Ethnic Origin

In adults, no clinically relevant interethnic differences among Caucasian, African-American, Hispanic, Japanese or Chinese patients were observed regarding pharmacokinetics and pharmacodynamics.

Hepatic Insufficiency

A Phase I study in adults investigated the influence of impaired hepatic function in cirrhotic patients (Child-Pugh Class A or B, number of patients 8 per group) on the pharmacodynamics and pharmacokinetics of a single dose of rivaroxaban.

In patients with mild hepatic impairment (Child-Pugh Class A), there was no difference as compared to healthy volunteers with respect to either pharmacodynamics (inhibition of Factor-Xa activity [1.08-fold for AUC and 0.98-fold for E_{max}]), prolongation of prothrombin time (1.02-fold for AUC and 1.06-fold for E_{max}), or pharmacokinetics (both total and unbound AUC [1.15 for total and 0.91-fold increase for unbound] and C_{max} [0.97 for total and 0.78-fold for unbound]).

Child-Pugh Class B patients had lower baseline Factor-Xa activity levels (0.64 U/mL) compared to healthy subjects and Child-Pugh Class A patients (0.85 U/mL, for both patient populations). Inhibition of Factor-Xa activity was more pronounced in Child-Pugh Class B patients compared to both healthy subjects and Child-Pugh Class A patients. The increase of inhibition was 2.6-fold AUC_(0-tn) and 1.2-fold maximal effect (E_{max}). The group difference was statistically significant, both for AUC_(0-tn) ($P < 0.01$) as well as for E_{max} ($P < 0.05$) of inhibition of Factor-Xa activity. In line with these results, a relevant difference in prolongation of PT was observed between healthy subjects and Child-Pugh Class B patients. The increase of prolongation was 2.1-fold (AUC_(0-tn)) and 1.4-fold (E_{max}). A statistically significant group-difference was observed for AUC_(0-tn) ($P < 0.0004$) as well as E_{max} ($P < 0.0001$).

Pharmacokinetic parameters also indicated a significant increase in Child-Pugh Class B patients as compared to healthy volunteers both on AUC pharmacokinetics (both total and unbound AUC [2.27-fold for total and 2.57-fold increase for unbound]) and C_{max} (1.27-fold for total and 1.38-fold for unbound).

A PK/PD analysis showed that the slope of the prothrombin time/plasma concentration correlation is increased by more than 2-fold for Child-Pugh Class B patients as compared to healthy volunteers. Since the global clotting test PT assesses the extrinsic pathway that is comprised of the coagulation Factor-VII, Factor-X, Factor-V, Factor-II, and Factor-I which are synthesized in the liver, impaired liver function can also result in prolongations of PT in the absence of anticoagulant therapy.

The PK/PD changes observed in Child-Pugh Class B patients are markers for the severity of the underlying hepatic disease which is expected to lead to a subsequent increased bleeding risk in this patient group.

JAMP Rivaroxaban is contraindicated in patients with hepatic disease (including Child-Pugh Class B and C) associated with coagulopathy, and having clinically relevant bleeding risk (see [2 CONTRAINDICATIONS](#), and [7 WARNINGS AND PRECAUTIONS – Hepatic Impairment](#)).

The safety and pharmacokinetics of single-dose rivaroxaban tablets (10 mg) were evaluated in a study in healthy subjects (n=16) and subjects with varying degrees of hepatic impairment (see [Table 13](#)). No patients with severe hepatic impairment (Child-Pugh C) were studied. Compared to healthy subjects with normal liver function, significant increases in rivaroxaban exposure were observed in subjects with moderate hepatic impairment (Child-Pugh B). Increases in pharmacodynamic effects were also observed.

Table 13 - Percent Increase of Rivaroxaban PK and PD Parameters from Normal in Subjects with Hepatic Insufficiency from a Dedicated Hepatic Impairment Study

Parameter		Hepatic Impairment Class (Child-Pugh Class)	
		Mild (Child Pugh A)	Moderate (Child-Pugh B)
		N=8	N=8
Exposure (% increase relative to normal)	AUC	15	127
	C _{max}	0	27
FXa Inhibition (% increase relative to normal)	AUC	8	159
	E _{max}	0	24
PT Prolongation (% increase relative to normal)	AUC	6	114
	E _{max}	2	41

PT = Prothrombin time; FXa =Coagulation factor Xa; AUC =Area under the concentration or effect curve; C_{max} =maximum concentration; E_{max} = maximum effect; and CrCl = creatinine clearance

Renal Insufficiency

As active rivaroxaban is partially cleared via the kidneys (30% to 40% of the dose), there is a direct but moderate correlation of systemic exposure to rivaroxaban with degree of renal impairment.

In a Phase I study in adults, following oral single dosing with rivaroxaban 10 mg in subjects with mild (CrCl 50 – 79 mL/min), moderate (CrCl 30 – 49 mL/min), or severe (CrCl 15 – 29 mL/min) renal impairment, rivaroxaban plasma concentrations (AUC) were increased 1.4-, 1.5-, and 1.6-fold, respectively compared to healthy subjects with normal renal function (CrCl ≥ 80 mL/min).

The overall inhibition of Factor-Xa activity (AUC_(0-48h) of effect versus time) was increased in these groups by a factor of 1.5, 1.9, and 2.0, respectively. The relative prolongation of prothrombin time (PT) was also affected by renal impairment and showed even more pronounced effects. AUC_(0-48h) of effect versus time was increased by a factor of 1.3, 2.2, and 2.4, respectively.

In Phase II, rivaroxaban plasma concentrations (AUC) were increased 1.2- and 1.5-fold in subjects with mild and moderate renal impairment respectively compared to healthy subjects with normal renal function and the peak inhibition of Factor-Xa activity (AUC_(0-48h) of effect versus time) was increased in these groups by a factor of 1.0 and 1.3 respectively. In a pooled analysis of Phase III THR or TKR subjects with mild and moderate renal impairment, the peak PT was increased by 1.0-, and 1.1-fold compared to subjects with normal renal function.

In Phase II (VTE treatment), rivaroxaban plasma concentrations (AUC) were 1.3- and 1.5-fold in subjects with mild and moderate renal impairment, respectively, compared to subjects with normal renal function. In Phase III subjects (VTE treatment) with mild renal impairment, the peak PT was increased by

1.1-fold, and 1.2-fold for moderate renal impairment compared to subjects with normal renal function.

In patients with atrial fibrillation evaluated in Phase III, the peak PT was increased by 1.2-fold for both mild and moderate renal impairment compared to subjects with normal renal function.

There was no evidence of substantial drug accumulation in patients with mild or moderate renal impairment.

The safety and pharmacokinetics of single-dose rivaroxaban tablets (10 mg) were evaluated in a study in healthy subjects [CrCl \geq 80 mL/min (n=8)] and in subjects with varying degrees of renal impairment (see [Table 14](#)). Compared to healthy subjects with normal creatinine clearance, rivaroxaban exposure increased in subjects with renal impairment. Increases in pharmacodynamic effects were also observed.

Table 14 - Percent Increase of Rivaroxaban PK and PD Parameters from Normal in Subjects with Renal Insufficiency from a Dedicated Renal Impairment Study

Parameter		CrCl (mL/min)		
		50 to 79	30 to 49	15 to 29
		N=8	N=8	N=8
Exposure (% increase relative to normal)	AUC	44	52	64
	C _{max}	28	12	26
FXa Inhibition (% increase relative to normal)	AUC	50	86	100
	E _{max}	9	10	12
PT Prolongation (% increase relative to normal)	AUC	33	116	144
	E _{max}	4	17	20

PT = Prothrombin time; FXa = Coagulation factor Xa; AUC = Area under the concentration or effect curve; C_{max} = maximum concentration; E_{max} = maximum effect; and CrCl = creatinine clearance

In subjects with mild renal impairment, the combined P-gp and moderate CYP 3A4 inhibitor erythromycin (500 mg three times a day) led to a 1.8-fold increase in mean rivaroxaban AUC and 1.6-fold increase in C_{max} when compared to subjects with normal renal function without co-medication. In subjects with moderate renal impairment, erythromycin led to a 2.0-fold increase in mean rivaroxaban AUC and 1.6-fold increase in C_{max} when compared to subjects with normal renal function without co-medication (see [7 WARNINGS AND PRECAUTIONS – Drug Interactions](#)). Subjects with either mild or moderate renal impairment had a 1.2- and 1.4-fold increase in Factor Xa inhibition, respectively, and a prolongation of prothrombin time of 1.7- and 1.75-fold in subjects with mild and moderate renal impairment, respectively.

11 STORAGE, STABILITY AND DISPOSAL

Store JAMP Rivaroxaban tablets at 15°C to 30°C.

Store in a safe place out of the reach of children.

12 SPECIAL HANDLING INSTRUCTIONS

No special handling instructions are required for this product.

PART II: SCIENTIFIC INFORMATION

13 PHARMACEUTICAL INFORMATION

Drug Substance


Proper Name: Rivaroxaban

Chemical Name: 5-Chloro-N-((5S)-2-oxo-3-[4-(3-oxo-4-morpholinyl)phenyl]-1,3-oxazolidin-5-yl)methyl)-2-thiophene-carboxamide

Molecular Formula and Molecular C₁₉ H₁₈ Cl N₃ O₅ S

Mass: 435.89 g / mol

Structural Formula:



Physicochemical Properties: Rivaroxaban is a pure (S)-enantiomer. It is an odorless, nonhygroscopic, white to yellowish powder. Rivaroxaban is practically insoluble in water, freely soluble in dimethyl sulfoxide, practically insoluble in anhydrous ethanol and in heptane.

14 CLINICAL TRIALS

14.1 Clinical Trials by Indication

Prevention of VTE after THR or TKR

Table 15 – Summary of the Pivotal Studies for the Prevention of Venous Thromboembolic Events in Patients Undergoing Elective Total Hip Replacement (THR) or Total Knee Replacement (TKR) Surgery

Study	Study Design	Treatment Regimen	Patient Populations
RECORD 1 ^a	THR patients prospectively randomized to rivaroxaban tablets or enoxaparin; noninferiority, double-blind, double-dummy design; multinational study.	<p>Rivaroxaban tablets 10 mg od oral for 35±4 days (first dose administered 6 to 8 h postoperatively)</p> <p>Enoxaparin 40 mg od SC for 36±4 days (first dose administered 12 h preoperatively)</p>	<p>Randomized 4541 (2266 rivaroxaban tablets, 2275 enoxaparin) Safety Population 4433 (2209 rivaroxaban tablets, 2224 enoxaparin) mITT 3153 (1595 rivaroxaban tablets, 1558 enoxaparin) mITT (for Major VTE) 3364 (1686 rivaroxaban tablets, 1678 enoxaparin) Per Protocol 3029 (1537 rivaroxaban tablets, 1492 enoxaparin)</p>
RECORD 3 ^a	TKR patients prospectively randomized to rivaroxaban tablets or enoxaparin; noninferiority, double-blind, double-dummy design; multinational study.	<p>Rivaroxaban tablets 10 mg od oral for 12±2 days (first dose administered 6 to 8 h postoperatively)</p> <p>Enoxaparin 40 mg od SC for 13±2 days (first dose administered 12 h preoperatively)</p>	<p>Randomized 2531 (1254 rivaroxaban tablets, 1277 enoxaparin) Safety Population 2459 (1220 rivaroxaban tablets, 1239 enoxaparin) mITT 1702 (824 rivaroxaban tablets, 878 enoxaparin) mITT (for Major VTE) 1833 (908 rivaroxaban tablets, 925 enoxaparin) Per Protocol 1631 (793 rivaroxaban tablets, 838 enoxaparin)</p>

^a The mean age of patients in RECORD 1 and 3 was 63.2±11.4, and 67.6±9 years, respectively.

Safety population = The safety population comprised those patients who received at least 1 dose of study drug. mITT = A subject was considered valid for the modified intent-to-treat (MITT) analysis if the subject was (1) valid for safety analysis; (2) had undergone the appropriate surgery; and (3) had an adequate assessment of thromboembolism.

mITT (for Major VTE) = A subject was valid for MITT analysis of major VTE, if the subject was (1) valid for safety analysis; (2) had undergone the appropriate surgery; and (3) had an adequate assessment of thromboembolism for major VTE.

Per Protocol = the per-protocol (PP) population was to include patients who were (1) valid for the MITT analysis; (2) had an adequate assessment of thromboembolism that, in case of a positive finding, was done not later than 36 h after stop of active study drug, in case of no finding, was done not later than 72 h after the end of active study drug; and (3) had no major protocol deviations.

Major VTE = composite of proximal DVT, nonfatal PE, or VTE-related death

od = once daily

SC = subcutaneous

The pivotal studies were designed to demonstrate the efficacy of rivaroxaban tablets (rivaroxaban) for the prevention of venous thromboembolic events, i.e., proximal and distal deep vein thrombosis (DVT) and pulmonary embolism (PE) in patients undergoing elective total hip replacement (THR) or total knee replacement (TKR) surgery. A once daily dose of 10 mg was selected for all Phase III studies in the prevention of VTE in patients undergoing THR or TKR surgery, based on clinical data generated in Phase II studies. Over 9,500 patients (7,050 in THR surgery; 2,531 in TKR surgery) were studied in these controlled randomized double-blind studies (RECORD 1, 2, and 3).

Pivotal Studies

The RECORD 1 and 3 studies were multicenter, multinational, prospective, double-blind, double-dummy studies in patients randomized to rivaroxaban tablets or to enoxaparin, see [Table 15](#). A non-inferiority was adopted with the pre-specification that, if non-inferiority was shown, a second analysis would be undertaken to determine if the efficacy of rivaroxaban tablets were superior to that of enoxaparin. RECORD 1 was conducted in patients undergoing elective THR surgery while RECORD 3 was conducted in patients undergoing elective TKR surgery. In both studies, rivaroxaban tablets 10 mg once daily started not earlier than 6 hours postoperatively was compared with an enoxaparin dosage regimen of 40 mg once daily started 12 hours preoperatively, as recommended in many countries worldwide. The dose of enoxaparin sodium approved for use in thromboprophylaxis in conjunction with elective THR or TKR surgery in Canada is subcutaneous 30 mg twice daily with the first dose to be administered 12 to 24 hours postoperatively. The primary endpoint was Total VTE a composite of any DVT (distal or proximal), nonfatal PE, or death from any cause. The main secondary endpoint was Major VTE, a composite endpoint comprising proximal DVT, nonfatal pulmonary embolism (PE), and VTE-related death. Other pre-specified secondary efficacy endpoints included the incidence of DVT (any thrombosis, including proximal and distal) and the incidence of symptomatic VTE.

Men and women of 18 years or older scheduled for elective surgery could be enrolled provided that they had no active or high risk of bleeding or other conditions contraindicating treatment with low-molecular-weight heparin, no significant liver disease, were not pregnant or breastfeeding women, or were not using HIV-protease inhibitors.

In RECORD 1 and 3, demographic and surgical characteristics were similar between the two groups except for a significantly larger number of females in RECORD 3 (rivaroxaban tablets 70% and enoxaparin 66%, $P = 0.03$). The reasons for exclusion of patients from various analyses in both studies were also similar.

The results of the non-inferiority analysis of Total VTE for RECORD 1 and 3 are presented in [Table 16](#). For the primary efficacy analysis, the difference between the incidences in the rivaroxaban tablets group and the enoxaparin group were estimated, after stratification according to country using the Mantel-Haenszel weighting, and the corresponding asymptotic two-sided 95% confidence interval was determined. Tests for non-inferiority and superiority were both based on the 95% confidence interval. Non-inferiority was shown if the lower limit of the CI was above the pre-specified non-inferiority margin; -3.5% in RECORD 1 and -4% in RECORD 3.

Table 16 – RECORD 1 (THR) and RECORD 3 (TKR): Non-inferiority Analysis of Total VTE^a, the Primary Composite Efficacy Endpoint, and its Components –Per Protocol (PP)^b Population Through the Double-Blind Treatment Period

	RECORD 1 (THR)		RECORD 3 (TKR)	
	Rivaroxaban Tablets 10 mg od N=1537 n (%)	Enoxaparin 40 mg od N=1492 n (%)	Rivaroxaban Tablets 10 mg od N=793 n (%)	Enoxaparin 40 mg od N=838 n (%)
Total VTE^a (primary composite endpoint)	13 (0.9%)	50 (3.4%)	74 (9.3%)	152 (18.1%)
	Absolute Risk Reduction ^c 2.5% (1.5% to 3.6%; <i>P</i> <0.001)		Absolute Risk Reduction ^c 8.7% (5.4% to 12.0%; <i>P</i> <0.001)	
DVT (proximal and/or distal)	11 (0.7)	47 (3.2)	74 (9.3)	147 (17.5)
Nonfatal PE	2 (0.1)	1 (<0.1)	0	3 (0.4)
Death from all causes	1 (<0.1)	2 (0.1)	0	2 (0.2)

^a Total VTE = DVT (proximal and/or distal), nonfatal PE, or death from all causes

^b PP = the per-protocol (PP) population was to include patients who were (1) valid for the MITT analysis; (2) had an adequate assessment of thromboembolism that, in case of a positive finding, was done not later than 36 h after stop of active study drug, in case of no finding, was done not later than 72 h after the end of active study drug; and (3) had no major protocol deviations

^c Mantel-Haenszel Weighted Reduction to Enoxaparin (Non-inferiority was shown if the lower limit of the CI was above the pre-specified non-inferiority margin; -3.5% in RECORD 1 and -4% in RECORD 3)

In both pivotal studies, the per-protocol analysis for the primary endpoint showed that rivaroxaban tablets 10 mg od/day (first dose 6 to 8 hours postoperatively) was not inferior to enoxaparin 40 mg/day (first dose 12 to 24 hours preoperatively).

Since non-inferiority was shown, a pre-specified superiority analysis was undertaken to determine if the efficacy of rivaroxaban tablets was superior to that of enoxaparin in the modified intent-to-treat population (mITT). The superiority analysis of Total VTE and data for the main secondary endpoint (Major VTE) and other secondary endpoints for RECORD 1 and 3 are presented in [Table 17](#) and [Table 18](#), respectively.

Table 17 – RECORD 1 (THR): Superiority Analysis for Total VTE (Primary Composite Endpoint)^a, Major VTE (Main Secondary Endpoint)^b and Their Components, and Other Selected Efficacy Endpoints – Modified ITT^c (MITT) Population Through the Double-Blind Treatment Period

Parameter	Rivaroxaban tablets 10 mg		Enoxaparin 40 mg		Absolute Risk Reduction ^d % (95% CI)	P-Value	Relative Risk Reduction % (95% CI)	P-Value
	n/N	% (95% CI)	n/N	% (95% CI)				
Total VTE	18/1595	1.1% (0.7% to 1.8%)	58/1558	3.7% (2.8% to 4.8%)	2.6% (1.5% to 3.7%)	<0.001	70% (49%-82%)	<i>P</i> <0.001
Major VTE	4/1686	0.2% (0.1% to 0.6%)	33/1678	2.0% (1.4% to 2.8%)	1.7% (1.0% to 2.5%)	<0.001	88% (66%-96%)	<i>P</i> <0.001
Death from all causes	4/1595	0.3% (0.1% to 0.6%)	4/1558	0.3% (0.1% to 0.7%)	0.0% (-0.4% to 0.4%)	1.00	--	--
Nonfatal PE	4/1595	0.3% (0.1% to 0.6%)	1/1558	0.1% (<0.1% to 0.4%)	-0.2% (-0.6% to 0.1%)	0.37	--	--
DVT (proximal and/or distal)	12/1595	0.8% (0.4% to 1.3%)	53/1558	3.4% (2.6% to 4.4%)	2.7% (1.7% to 3.7%)	<0.001	--	--
Proximal DVT	1/1595	0.1% (<0.1% to 0.4%)	31/1558	2.0% (1.4% to 2.8%)	1.9% (1.2% to 2.7%)	<0.001	--	--
Distal DVT only	11/1595	0.7% (0.3% to 1.2%)	22/1558	1.4% (0.9% to 2.1%)	0.7% (0.0% to 1.5%)	0.04	--	--
VTE-related death	0/1595	0%	1/1558	<0.1%	--	--	--	--
Symptomatic VTE^e	6/2193	0.3% (0.1% to 0.6%)	11/2206	0.5% (0.3% to 0.9%)	0.2% (-0.1% to 0.6%)	0.22	--	--

^{a.} Total VTE = composite of DVT (proximal and/or distal), nonfatal PE, or death from all causes

^{b.} Major VTE = composite of proximal DVT, nonfatal PE, or VTE-related death

^{c.} MITT = subject valid for safety analysis, has undergone appropriate surgery, has adequate assessment of thromboembolism

^{d.} Mantel-Haenszel Weighted Reduction to Enoxaparin given for all endpoints except nonfatal PE and death from all causes, for which unweighted (exact) estimates were given. Superiority was shown if the lower limit of the CI was above zero.

^{e.} Safety population for Symptomatic VTE (patients valid for safety analysis who underwent the appropriate surgery). The safety population was used because assessment of symptomatic events is possible in the greater population, regardless of the availability of an adequate venographic assessment.

Table 18 – RECORD 3 (TKR): Superiority Analysis for Total VTE (Primary Composite Endpoint)^a, Major VTE (Main Secondary Endpoint)^b and Their Components, and Other Selected Efficacy Endpoints – Modified ITT (MITT)^c Population Through the Double-Blind Treatment Period

Parameter	Rivaroxaban tablets 10 mg		Enoxaparin 40 mg		Absolute Risk Reduction ^d % (95% CI)	P-Value	Relative Risk Reduction % (95% CI)	P-Value
	n/N	% (95% CI)	n/N	% (95% CI)				
Total VTE	79/824	9.6% (7.7% to 11.8%)	166/878	18.9% (16.4% to 21.7%)	9.2% (5.9% to 12.4%)	<0.001	49% (35%-61%)	<0.001
Major VTE	9/908	1.0% (0.5% to 1.9%)	24/925	2.6% (1.7% to 3.8%)	1.6% (0.4% to 2.8%)	0.01	62% (18%-82%)	0.016
Death from all causes	0/824	0% (0.0% to 0.5%)	2/878	0.2% (0.0% to 0.8%)	0.2% (-0.2% to 0.8%)	0.23	--	--
Nonfatal PE	0/824	0% (0.0% to 0.3%)	4/878	0.5% (0.1% to 1.2%)	0.5% (0.0% to 1.2%)	0.06	--	--
DVT (proximal and/or distal)	79/824	9.6% (7.7% to 11.8%)	160/878	18.2% (15.7% to 20.9%)	8.4% (5.2% to 11.7%)	<0.001	--	--
Proximal DVT	9/824	1.1% (0.5% to 2.1%)	20/878	2.3% (1.4% to 3.5%)	1.1% (-0.1% to 2.3%)	0.07	--	--
Distal DVT only	70/824	8.5% (6.7% to 10.6%)	140/878	15.9% (13.6% to 18.5%)	7.3% (4.3% to 10.4%)	<0.001	--	--
VTE-related death	0/824	0%	0/878	0%	--	--	--	--
Symptomatic VTE^e	8/1201	0.7% (0.3% to 1.3%)	24/1217	2.0% (1.3% to 2.9%)	1.3% (0.4% to 2.2%)	0.005	--	--

a. Total VTE = composite of DVT (proximal and/or distal), nonfatal PE, or death from all causes.

b. Major VTE = composite of proximal DVT, nonfatal PE, or VTE-related death

c. MITT = subject valid for safety analysis, has undergone appropriate surgery, has adequate assessment of thromboembolism

d. Mantel-Haenszel Weighted Reduction to Enoxaparin given for all endpoints except nonfatal PE and death from all causes, for which unweighted (exact) estimates were given. Superiority was shown if the lower limit of the CI was above zero.

e. Safety population for Symptomatic VTE (patients valid for safety analysis who underwent the appropriate surgery). The safety population was used because assessment of symptomatic events is possible in the greater population, regardless of the availability of an adequate venographic assessment.

The efficacy results of the pre-specified analysis using a modified intent-to-treat population indicate that rivaroxaban tablets 10 mg administered postoperatively once daily is superior in preventing DVT to enoxaparin 40 mg once daily (first dose 12 hours preoperatively). The Canadian approved dosage regimen for enoxaparin is 30 mg every 12 hours (first dose is to be administered 12 to 24 hours postoperatively).

There are no definitive head-to-head studies to compare the safety and efficacy of the Canadian approved enoxaparin dosage regimen to the enoxaparin dosage regimen used in the RECORD 1 and 3 studies.

In the safety population of 3429 patients treated with rivaroxaban tablets and 3463 patients treated with enoxaparin in the pivotal studies (RECORD 1 and 3), the results observed for bleeding events have been summarized in [Table 19](#). In RECORD 1, serious drug-related treatment-emergent adverse events were reported in 26 (1.2%) for rivaroxaban tablets and 23 (1.0%) for enoxaparin. In RECORD 3, serious drug-related treatment-emergent adverse events were reported in 26 (2.1%) for rivaroxaban tablets and 19 (1.5%) for enoxaparin.

Table 19 – RECORD 1 and 3: Detailed Overview of Treatment-Emergent Bleeding Events (Safety Population)^a

	RECORD 1 (THR)			RECORD 3 (TKR)		
	Rivaroxaban Tablets 10 mg od N=2209	Enoxaparin 40 mg od N=2224	P-Value	Rivaroxaban tablets 10 mg od N=1220	Enoxaparin 40 mg od N=1239	P-Value
Any Bleeding n (%) (95% CI)	133 (6.0%) (5.1% to 7.1%)	131 (5.9%) (5.0% to 7.0%)	0.90	60 (4.9%) (3.8%-6.3%)	60 (4.8%) (3.7%-6.2%)	1.0
Major Bleeding^b n (%) (95% CI)	6 (0.3%) (0.1%-0.6%)	2 (0.1%) (<0.1%-0.3%)	0.18	7 (0.6%) (0.2%-1.2%)	6 (0.5%) (0.2%-1.1%)	0.79
Fatal Bleeding^c	1 (<0.1%) ^b	0 (0.0%)	--	0 (0.0%)	0 (0.0%)	--
Bleeding into a critical organ n (%)	1 (<0.1%)	0 (0.0%)	--	1 (0.1%)	2 (0.2%)	--
Bleeding leading to reoperation n (%)	2 (0.1%)	1 (<0.1%)	--	5 (0.4%)	4 (0.3%)	--
Clinically overt extra-surgical site bleeding leading to a fall in hemoglobin n (%)	2 (0.1%)	1 (<0.1%)	--	1 (0.1%)	0 (0.0%)	--
Clinically overt extra-surgical site bleeding leading to transfusion of ≥2 units of blood n (%)	2 (0.1%)	1 (<0.1%)	--	1 (0.1%)	0 (0.0%)	--
Nonmajor Bleeding^d n (%)	128 (5.8%)	129 (5.8%)	--	53 (4.3%)	54 (4.4%)	--

	RECORD 1 (THR)			RECORD 3 (TKR)		
	Rivaroxaban Tablets 10 mg od N=2209	Enoxaparin 40 mg od N=2224	P-Value	Rivaroxaban tablets 10 mg od N=1220	Enoxaparin 40 mg od N=1239	P-Value
Clinically relevant nonmajor bleedingn (%)	65 (2.9%)	54 (2.4%)	--	33 (2.7%)	28 (2.3%)	--
Hemorrhagic wound complications^e n (%)	34 (1.5%)	38 (1.7%)	--	25 (2.0%)	24 (1.9%)	--

- a Patients may have had more than one type of event, and an event could fall into more than one category; adjudicated treatment-emergent bleeding events included those beginning after the initiation of the study drug and up to 2 days after last dose of the study drug.
- b Major bleeding events included: (1) fatal, (2) bleeding into a critical organ (e.g, retroperitoneal, intracranial, intraocular, or intraspinal bleeding/hemorrhagic puncture), (3) bleeding requiring reoperation, (4) clinically overt extra-surgical site bleeding associated with ≥ 2 g/dL fall in hemoglobin or leading to infusion of ≥ 2 units of whole blood or packed cells.
- c The event occurred before the administration of the first dose of rivaroxaban.
- d Nonmajor bleeding events were bleeding events that did not fulfill the criteria of major bleeding.
- e Composite of excessive wound hematoma and reported surgical-site bleeding.

Phase III Supportive Study

RECORD 2 was a randomized, double-blind, double-dummy, prospective study conducted in 2509 randomized patients (safety population = 2457; mITT = 1733) undergoing THR. The aim of RECORD 2 was to assess extended thromboprophylaxis with rivaroxaban tablets for 35 ± 4 days. RECORD 2 was similar in study design, inclusion/exclusion criteria and endpoints to RECORD 1, except that enoxaparin 40 mg once daily (first dose given preoperatively) was given for a shorter duration (12 ± 2 days) than rivaroxaban tablets 10 mg od (35 ± 4 days). Comparative efficacy claims to enoxaparin may not be drawn from this study, due to the differences in the treatment duration of rivaroxaban tablets and enoxaparin.

Table 20 – RECORD 2 (THR): Superiority Analysis for Total VTE (Primary Composite Endpoint)^a, Major VTE (Main Secondary Endpoint)^b and Their Components, and Other Selected Efficacy Endpoints – Modified ITT^c (MITT) Population Through the Double-Blind Treatment Period

Parameter	Rivaroxaban tablets 10 mg od for 35±4 days		Enoxaparin 40 mg for 12±2 days		Absolute Risk Reduction ^d	P-Value	Relative Risk Reduction	P-Value
	n/N	% (95% CI)	n/N	% (95% CI)	% (95% CI)		% (95% CI)	
Total VTE	17/864	2.0% (1.2% to 3.1%)	81/869	9.3% (7.5% to 11.5%)	7.3% (5.2% to 9.4%)	<0.0001	79% (65% to 87%)	< 0.001
Major VTE	6/961	0.6% (0.2% to 1.4%)	49/962	5.1% (3.8% to 6.7%)	4.5% (3.0% to 6.0%)	<0.0001	88% (71% to 95%)	< 0.001
Death from all causes	2/864	0.2% (<0.1% to 0.8%)	6/869	0.7% (0.3% to 1.5%)	0.5% (-0.2% to 1.3%)	0.29	--	--
Nonfatal PE	1/864	0.1% (<0.1% to 0.6%)	4/869	0.5% (0.1% to 1.2%)	0.3% (-0.2% to 1.1%)	0.37	--	--
DVT (proximal and/or distal)	14/864	1.6% (0.9% to 2.7%)	71/869	8.2% (6.4% to 10.2%)	6.5% (4.5% to 8.5%)	<0.0001	--	--
Proximal DVT	5/864	0.6% (0.2% to 1.3%)	44/869	5.1% (3.7% to 6.7%)	4.5% (2.9% to 6.0%)	<0.0001	--	--
Distal DVT only	9/864	1.0% (0.5% to 2.0%)	27/869	3.1% (2.1% to 4.5%)	2.0% (0.7% to 3.3%)	0.0025	--	--
VTE-related death	0/864	0%	1/869	0.1%	--	--	--	--
Symptomatic VTE^e	3/1212	0.2% (<0.1% to 0.7%)	15/1207	1.2% (0.7% to 2.0%)	1.0% (0.3% to 1.8%)	0.0040	--	--

a. Total VTE = composite of DVT (proximal and/or distal), nonfatal PE, or death from all causes.

b. Major VTE = composite of proximal DVT, nonfatal PE, or VTE-related death

c. MITT = subject valid for safety analysis, has undergone appropriate surgery, has adequate assessment of thromboembolism

d. Mantel-Haenszel Weighted Reduction to Enoxaparin given for all endpoints except nonfatal PE and death from all causes, for which unweighted (exact) estimates were given. Superiority was shown if the lower limit of the CI was above zero.

e. Safety population for Symptomatic VTE (patients valid for safety analysis who underwent the appropriate surgery). The safety population was used because assessment of symptomatic events is possible in the greater population regardless of the availability of an adequate venographic assessment.

Table 21 – RECORD 2 (THR): Detailed Overview of Treatment-Emergent Bleeding Events (Safety Population)^a

	Rivaroxaban tablets 10 mg od for 35±4 days N=1228	Enoxaparin 40 mg od for 12±2 days N=1229	P-Value
Any Bleeding n (%) (95% CI)	81 (6.6%) (5.3% to 8.1%)	68 (5.5%) (4.3% to 7.0%)	0.27
Major Bleeding^b n (%) (95% CI)	1 (0.1%) (0.0–0.5)	1 (0.1%) (0.0–0.5)	1.00
Fatal bleeding	0 (0.0%)	0 (0.0%)	--
Bleeding into a critical organ n (%)	0 (0.0%)	1 (0.1%)	--
Bleeding leading to reoperation n (%)	0 (0.0%)	0 (0.0%)	--
Clinically overt extra-surgical site bleeding leading to a fall in hemoglobin n (%)	1 (0.1%)	0 (0.0%)	--
Clinically overt extra-surgical site bleeding leading to transfusion of ≥2 units of blood n (%)	1 (0.1%)	0 (0.0%)	--
Nonmajor Bleeding^c n (%)	80 (6.5%)	67 (5.5%)	--
Clinically relevant nonmajor bleeding n (%)	40 (3.3%)	33 (2.7%)	--
Hemorrhagic wound complications^d n (%)	20 (1.6%)	21 (1.7%)	--

a Patients may have had more than one type of event, and an event could fall into more than one category; adjudicated treatment-emergent bleeding events included those beginning after the initiation of the study drug and up to 2 days after last dose of the study drug.

b Major bleeding events included: (1) fatal, (2) bleeding into a critical organ (e.g, retroperitoneal, intracranial, intraocular, or intraspinal bleeding/hemorrhagic puncture), (3) bleeding requiring reoperation, (4) clinically overt extra-surgical site bleeding associated with ≥2 g/dL fall in hemoglobin or leading to infusion of ≥2 units of whole blood or packed cells.

c Nonmajor bleeding events were bleeding events that did not fulfill the criteria of major bleeding.

d Composite of excessive wound hematoma and reported surgical-site bleeding.

The results from this study demonstrate that extended duration prophylaxis with 10 mg rivaroxaban tablets od for 35 days provided clinically meaningful decreases in Total VTE, Major VTE, and symptomatic VTE in THR patients without an increased risk of bleeding.

Treatment of VTE and prevention of recurrent DVT and PE

Table 22 - Summary of the Pivotal Studies for the Treatment of VTE and Prevention of Recurrent DVT and PE

Study	Study Design	Treatment Regimen	Patient Population
EINSTEIN DVT	multicenter, randomized, open-label, event-driven non-inferiority study for efficacy	Rivaroxaban tablets 15 mg bid for 3 weeks followed by 20 mg od 3, 6 or 12 months ^a	Randomized 3449 (1731 rivaroxaban tablets, 1718 Enox/VKA) Safety Population 3429 (1718 rivaroxaban tablets, 1711 Enox/VKA) Per Protocol 3096 (1525 rivaroxaban tablets, 1571 Enox/VKA)
EINSTEIN PE		Standard Therapy Enoxaparin bid bridging to therapeutic VKA 3, 6 or 12 months ^a	Randomized 4833, (2420 rivaroxaban tablets, 2413 Enox/VKA) Safety Population 4817 (2412 rivaroxaban tablets, 2405 Enox/VKA) Per Protocol 4462 (2224 rivaroxaban tablets, 2238 Enox/VKA)
EINSTEIN Extension	multicenter, randomized, double-blind, placebo-controlled, event-driven, superiority study for efficacy in subjects with symptomatic proximal DVT or PE	Rivaroxaban tablets 20 mg once daily or placebo for 6 or 12 months ^a	Randomized 1197 (602 rivaroxaban tablets, 594 placebo)
EINSTEIN CHOICE	multicenter, randomized, double-blind, double-dummy, active-comparator (ASA), event-driven, superiority study for efficacy in subjects with symptomatic DVT and/or PE	Rivaroxaban tablets 10 mg, or 20 mg or ASA 100 mg once daily ^b	Randomized 3396 (1121 rivaroxaban tablets 20 mg, 1136 rivaroxaban tablets 10 mg, 1139 ASA 100 mg)

a Treatment duration as determined by investigator

b Individual (actual) treatment duration depends on the individual randomization date: either 12 months, 9 to <12 months or 6 months

Safety population = The safety population comprised those subjects who received at least one dose of study medication.

bid = twice daily; od = once daily; VKA = vitamin K antagonist; enox = enoxaparin; ASA= acetylsalicylic acid

The EINSTEIN clinical development program consisted of four Phase III studies. The EINSTEIN DVT and EINSTEIN PE studies evaluated the treatment of VTE and prevention of recurrent DVT and PE. The EINSTEIN Extension study evaluated the benefit of continued treatment in subjects for whom clinical uncertainty regarding the absolute risk-benefit of extended duration existed.

Patients with VTE who were treated either with rivaroxaban or enoxaparin/VKA for 6 or 12 months in EINSTEIN DVT or EINSTEIN PE, or who were treated for 6 to 14 months with VKA and in whom there was equipoise to continue anticoagulant treatment were eligible for enrollment into EINSTEIN Extension.

Subjects considered to have been adequately treated with 6 to 12 months of therapy or those who required more prolonged anticoagulation therapy were not included.

In EINSTEIN CHOICE, patients with confirmed symptomatic VTE who completed 6-12 months of anticoagulant treatment and in whom there was equipoise to continue anticoagulant treatment were eligible for the study. Patients with an indication for continued therapeutic-dosed anticoagulation were excluded.

Duration of administration in EINSTEIN DVT was up to 12 months (i.e., 3, 6 or 12 months) as determined by the investigator, prior to randomization, based on local risk assessment and guidelines. Nearly half of the subjects were treated for 6 to 9 months.

In EINSTEIN DVT, and EINSTEIN PE rivaroxaban tablets were compared to the standard dual-drug regimen of enoxaparin administered for at least 5 days in combination with VKA until the PT/INR was in therapeutic range (≥ 2.0). VKA alone was then continued, dose-adjusted to maintain the PT/INR values within the therapeutic range of 2.0 to 3.0.

Table 23 – Co-morbid Diseases and Characteristics of Patients in EINSTEIN DVT, EINSTEIN PE and EINSTEIN Extension – ITT Population

	EINSTEIN DVT	EINSTEIN PE	EINSTEIN Extension	EINSTEIN CHOICE
Males (%)	57%	53%	58%	55%
Age, mean (years)	56	58	58	59
Creatinine Clearance (mL/min)				
< 50	7%	8%	7%	5%
50 to < 80	23%	25%	21%	25%
≥ 80	68%	66%	62%	70%
Risk Factors				
Patients with idiopathic DVT/PE	48%	49%	59%	41%
Recent surgery or trauma	19%	17%	4.1%	13%
Immobilization	15%	16%	14%	11%
Previous VTE	19%	19%	16%	18%
Mean TTR, Enox/VKA arm	58% ^a	63% ^b	n/a	n/a
North American subjects	64%	63%	n/a	n/a
Pre-randomization anticoagulation ^c	73%	92%	n/a	n/a
Actual Treatment Duration in rivaroxaban tablets arm				
≥ 3 months	92%	92%	91%	n/a
≥ 6 months	68%	73%	62%	n/a
≥ 12 months	3%	4%	2%	n/a

a. unadjusted Mean TTR. Adjusted Mean TTR is 60%.

b. Adjusted mean TTR.

c. Pre-randomization anticoagulation was limited to 24 hours in the majority of cases.

n/a=not applicable

Table 24 - Efficacy outcomes in EINSTEIN DVT, EINSTEIN PE and EINSTEIN Extension – ITT population

	EINSTEIN DVT			EINSTEIN PE			EINSTEIN Extension		
	Rivaroxaban tablets N=1731	Enox/VKA N=1718	HR (95% CI) P-value	Rivaroxaban tablets N=2419	Enox/VKA N=2413	HR (95% CI) P-value	Rivaroxaban tablets N=602	Placebo N=594	HR (95% CI) P-value
Symptomatic Recurrent VTE ^b	36 (2.1%)	51 (3.0%)	0.68 (0.44-1.04) <i>P</i> <0.001 ^a	50 (2.1%)	44 (1.8%)	1.12 (0.754-1.68) <i>P</i> =0.0026 ^a	8 (1.3%)	42 (7.1%)	0.18 (0.09-0.39) <i>P</i> <0.001
Type of Symptomatic Recurrent VTE									
Fatal PE	1 (<0.1%)	0	-	3 (0.1%)	1 (<0.1%)	-	0	1 (0.2%)	-
Death where PE could not be ruled out	3 (0.2%)	6 (0.3%)	-	8 (0.3%)	6 (0.2%)	-	1 (0.2%)	0	-
Recurrent PE only	20 (1.2%)	18 (1.0%)	-	23 (1.0%)	20 (0.8%)	-	2 (0.3%)	13 (2.2%)	-
Recurrent DVT plus PE	1 (<0.1%)	0	-	0	2 (<0.1%)	-	n.a.	n.a.	-
Recurrent DVT only	14 (0.8%)	28 (1.6%)	-	18 (0.7%)	17 (0.7%)	-	5 (0.8%)	31 (5.2%)	-
Symptomatic recurrent VTE and all-cause mortality	69 (4.0%)	87 (5.1%)	0.72 (0.53-0.99) <i>P</i> =0.044 ^c	97 (4.0%)	82 (3.4%)	1.16 (0.86-1.55) <i>P</i> =0.3333 ^c	8 (1.3%)	43 (7.2%)	0.18 (0.085-0.38) <i>P</i> <0.0001 ^c
Net Clinical Benefit	51 (2.9%)	73 (4.2%)	0.67 (0.47-0.95) <i>P</i> =0.027 ^c	83 (3.4%)	96 (4.0%)	0.85 (0.63-1.14) <i>P</i> =0.2752 ^c	12 (2.0%)	42 (7.1%)	0.28 (0.15-0.53) <i>P</i> <0.0001
All On-Treatment Vascular Events	12 (0.7%)	14 (0.8%)	0.79 (0.36–1.71) <i>P</i> =0.55 ^c	35 (1.5%)	37 (1.5%)	0.94 (0.59-1.49) <i>P</i> =0.7780 ^c	3 (0.5)	44 (0.7%)	0.74 (0.17-3.3) <i>P</i> =0.69
All-cause Mortality	38 (2.2%)	49 (2.9%)	0.67 (0.44–1.02) <i>P</i> =0.06 ^c	58 (2.4%)	50 (2.1%)	1.13 (0.77-1.65) <i>P</i> =0.5260	1 (0.2%)	2 (0.3%)	-

- a. P-value for non-inferiority (one-sided)
- b. Some patients had more than one event
- c. P-value for superiority (two-sided)
- n.a.=not assessed

Table 25 - Efficacy outcomes in EINSTEIN CHOICE


	Rivaroxaban tablets 10 mg N=1127	Rivaroxaban tablets 20 mg N=1107	ASA 100 mg N=1131	Rivaroxaban 20 mg vs. ASA 100 mg HR^a (95% CI) P-value	Rivaroxaban 10 mg vs. ASA 100 mg HR^a (95% CI) P-value
Symptomatic Recurrent VTE ^b	13 (1.2%)	17 (1.5%)	50 (4.4%)	0.34 (0.20-0.59) <i>P</i> = 0.0001 ^c	0.26 (0.14-0.47) <i>P</i> <0.001 ^c
Symptomatic recurrent VTE and all-cause mortality	15 (1.3%)	23 (2.1%)	55 (4.9%)	0.42 (0.26-0.68) <i>P</i> =0.0005	0.27 (0.15-0.47) <i>P</i> <0.0001
Net Clinical Benefit	17 (1.5%)	23 (2.1%)	53 (4.7%)	0.44 (0.27-0.71) <i>P</i> = 0.0009 ^c	0.32 (0.18-0.55) <i>P</i> = <0.0001 ^c

- a. P-value for non-inferiority (one-sided)
 - b. Some patients had more than one event
 - c. P-value for superiority (two-sided)
- FAS =Full Analysis SET

EINSTEIN DVT

EINSTEIN DVT met its principal objective demonstrating that rivaroxaban tablets was non-inferior to enoxaparin/VKA for the primary outcome of symptomatic recurrent VTE (HR of 0.68 [95% CI = 0.44 - 1.04], $P < 0.001$) (Table 24 and Figure 3). The results of per-protocol analyses were similar to those of the intention-to-treat analysis. The pre-specified test for superiority was not statistically significant ($P = 0.0764$). The incidence rates for the principal safety outcome (major or clinically relevant non-major bleeding events), as well as the secondary safety outcome (major bleeding events), were similar for both groups (HR of 0.97 [95% CI = 0.76-1.22], $P = 0.77$ and HR of 0.65 [95% CI = 0.33-1.30], $P = 0.21$, respectively). The pre-defined secondary outcome of net clinical benefit, (the composite of the primary efficacy outcome and major bleeding events), was reported with a HR of 0.67 ([95% CI = 0.47-0.95], nominal $P = 0.03$) in favour of rivaroxaban tablets. The relative efficacy and safety findings were consistent regardless of pre-treatment (none, LMWH, unfractionated heparin or fondaparinux) as well as among the 3, 6 and 12-month durations. In terms of other secondary outcomes, vascular events during study treatment occurred in 12 patients (0.7%) in the rivaroxaban tablets arm and 14 patients (0.8%) in the enoxaparin/VKA group (HR of 0.79 [95% CI = 0.36-1.71], $P = 0.55$), and total mortality accounted for 38 (2.2%) vs. 49 (2.9%) patients in the rivaroxaban tablets vs. enoxaparin/VKA arms, respectively, within intended treatment duration ($P = 0.06$).

Figure 3: Kaplan-Meier Cumulative Event Rates for the Primary Efficacy Outcome in EINSTEIN-DVT –Intention-to-Treat Population




EINSTEIN PE

EINSTEIN PE met its principal objective demonstrating that rivaroxaban tablets was non-inferior to enoxaparin/VKA for the primary efficacy outcome of symptomatic recurrent VTE (HR of 1.12 [95% CI: 0.75-1.68], $P=0.0026$) (Table 24 and Figure 4). The results of per-protocol analyses were similar to those of the intention-to-treat analysis. The pre-specified test for superiority was not statistically significant ($P=0.5737$). The incidence rate of the principal safety outcome (major or clinically relevant non-major bleeding events) was similar for both groups (HR of 0.90 [95% CI: 0.76 to 1.07] $P=0.2305$). For major bleeding events, the incidence rate was nominally lower in favour of rivaroxaban tablets treatment group (HR of 0.49 [95% CI: 0.31 – 0.79]; $P=0.003$). The pre-defined secondary outcome of net clinical benefit (the composite of the primary efficacy outcome and major bleeding events) was reported with a HR of 0.85 ([95% CI: 0.63-1.14]; $P=0.27$) in favour of rivaroxaban tablets. The relative efficacy and safety findings were consistent regardless of pre-treatment (none, LMWH, unfractionated heparin or fondaparinux) as well as among the 3, 6 and 12 month durations. In terms of other secondary outcomes, vascular events during study treatment occurred in 41 patients (1.7%) in the rivaroxaban tablets arm and 39 patients (1.6%) in the enoxaparin/VKA group (HR of 1.04 [95% CI = 0.67- 1.61], $P = 0.86$), and total mortality accounted for 58 (2.4%) vs. 50 (2.1%) patients in the rivaroxaban tablets vs. enoxaparin/VKA arms, respectively, within intended treatment duration ($P = 0.53$).

Figure 4: Kaplan-Meier analysis: cumulative rate of primary efficacy outcome in study 11702 PE - ITT Population


Graphic: Kaplan-Meier Cumulative Rate of the Primary efficacy outcome
 Timepoint: Event or censoring up to the Intended treatment duration
 Population: Intent to treat



EINSTEIN Extension

In the EINSTEIN Extension study, rivaroxaban tablets was superior to placebo for the primary efficacy outcome with a HR of 0.18 [95% CI = 0.09-0.39], $P < 0.001$ (i.e, a relative risk reduction of 82%) (Table 24 and Figure 5). For the principal safety outcome (major bleeding events) there was no significant difference between patients treated with rivaroxaban tablets compared to placebo ($P = 0.11$). The pre-defined secondary outcome of net clinical benefit, defined as the composite of the primary efficacy outcome and major bleeding events, was reported with a HR of 0.28 ([95% CI = 0.15-0.53], $P < 0.001$) in favour of rivaroxaban tablets. In terms of other secondary outcomes, vascular events occurred in 3 patients in the rivaroxaban tablets arm and 4 patients in the placebo group (HR of 0.74 [95% CI = 0.17-3.3], $P = 0.69$), and total mortality accounted for 1 (0.2%) vs. 2 (0.3%) of patients in the rivaroxaban tablets vs. placebo arms, respectively.

Figure 5: Kaplan-Meier Cumulative Event Rates for the Primary Efficacy Outcome in EINSTEIN Extension



EINSTEIN CHOICE

In EINSTEIN CHOICE 3,396 patients with confirmed symptomatic DVT and/or PE who completed 6-12 months of therapeutic-dose anticoagulation and who did not have an indication for continued anticoagulation in therapeutic doses, were studied for the prevention of fatal PE or non-fatal symptomatic recurrent DVT/PE. Patients with an indication for continued therapeutic-dosed anticoagulation were excluded from the study. The treatment duration was up to 12 months depending on the individual randomization date (median: 351 days). Rivaroxaban tablets 20 mg once daily and rivaroxaban tablets 10 mg once daily was compared with 100 mg acetylsalicylic acid once daily.

The primary efficacy outcome was symptomatic recurrent VTE defined as the composite of recurrent DVT or fatal or non-fatal PE. The secondary efficacy outcome was the composite of the primary efficacy outcome, MI, ischemic stroke, or non-CNS systemic embolism.


In the EINSTEIN CHOICE study, the primary efficacy objective for superiority was met for both rivaroxaban tablets 20 mg and 10 mg versus acetylsalicylic acid 100mg. The secondary efficacy outcome was significantly reduced when comparing rivaroxaban tablets 20 mg or 10 mg vs. 100 mg acetylsalicylic acid. The principal

safety outcome (major bleeding events) was similar for patients treated with rivaroxaban tablets 20 mg and 10 mg once daily compared to 100 mg acetylsalicylic acid. The secondary safety outcome (non-major bleeding associated with treatment cessation of more than 14 days) was similar when comparing rivaroxaban tablets 20 mg or 10 mg vs. 100 mg acetylsalicylic acid. Outcomes were consistent across the patients with provoked and unprovoked VTE (see [Table 25](#)).

In a prespecified net clinical benefit analysis (NCB) (primary efficacy outcome plus major bleeding events) of EINSTEIN CHOICE, a HR of 0.44 (95% CI 0.27 - 0.71, p = 0.0009) for rivaroxaban tablets 20 mg once daily vs 100 mg acetylsalicylic acid once daily and a HR of 0.32 (95% CI 0.18 - 0.55, p <0.0001) for rivaroxaban tablets 10mg once daily vs 100 mg acetylsalicylic acid once daily were reported.

Figure 6: Kaplan-Meier analysis: cumulative event rates of the primary efficacy outcome until the end of individual intended treatment duration (FAS)

Kaplan-Meier plot of cumulative rate of the Primary Efficacy Outcome up to the end of individual intended treatment duration (full analysis set)



Bayer: /by-sasp/patdb/projects/597939/16416/stat/prod/pgms/af_14_2_2_adtte_efficacy_km_cum.sas erjli 19DEC2016 13:59

PHASE IV STUDY

Phase IV clinical studies (XALIA) was done to evaluate the effects of rivaroxaban use under real-world (clinical practice) conditions.

XALIA

In addition to the Phase III EINSTEIN program, a prospective, non-interventional, open-label cohort study (XALIA) investigated the long-term safety of rivaroxaban tablets under real-world conditions (central outcome adjudication including recurrent VTE, major bleeding and death. In 2619 rivaroxaban

tablets-treated patients, rates of major bleeding, recurrent VTE and all-cause mortality for rivaroxaban tablets were 0.7%, 1.4% and 0.5%, respectively.

These results are consistent with the established safety profile of rivaroxaban tablets in this population.

Prevention of Stroke and Systemic Embolism in Patients with Atrial Fibrillation

Table 26 – Summary of the ROCKET AF Trial, a Phase III Clinical Trial in Atrial Fibrillation

Study	Study Design	Treatment Regimen	Populations
ROCKET AF	double-blind, double-dummy prospective randomized parallel-group multinational study	<p>Rivaroxaban tablets 20 mg od (15 mg od for patients with moderate renal impairment [CrCl 30 – 49 mL/min])</p> <p>Warfarin dose adjusted to an INR of 2.5 (range 2.0 to 3.0)</p>	<p>Randomized 14,264 (7131 rivaroxaban tablets, 7133 warfarin)</p> <p>Safety Population 14,236 (7111 rivaroxaban tablets, 7125 warfarin)</p> <p>Per Protocol 14,054 (7008 rivaroxaban tablets, 7046 warfarin)</p>

Randomized = The randomized/ intent-to-treat population represent all uniquely randomized patients.
 Safety population = The safety population comprised those patients who received at least 1 dose of study drug.
 Per Protocol = The per-protocol population was all intent to-treat patients excluding those who have specific pre-defined major protocol deviations that occur by the time of enrollment into the study or during the trial.
 od = once daily

Evidence for the effectiveness of rivaroxaban tablets are derived from the ROCKET AF trial, a prospective, randomized, double-blind, double-dummy, parallel-group, multicenter, pivotal clinical study comparing the efficacy and safety of once daily oral rivaroxaban tablets with dose-adjusted warfarin in patients with atrial fibrillation at risk of stroke or systemic embolism. In addition to documented atrial fibrillation, patients had prior stroke, TIA or systemic embolism, or 2 or more of the following risk factors without prior stroke:

- clinical heart failure and/or left ventricular ejection fraction $\leq 35\%$
- hypertension
- age ≥ 75 years
- diabetes mellitus

Patients with prosthetic heart valves, or those with hemodynamically significant rheumatic heart disease, especially mitral stenosis, were excluded from the ROCKET AF study, and thus were not evaluated. These trial results do not apply to patients these conditions, whether in the presence or absence of atrial fibrillation (see [7 WARNINGS AND PRECAUTIONS – Cardiovascular, Patients with valvular disease](#)).

The primary objective of this study was to demonstrate that rivaroxaban tablets, a direct Factor-Xa inhibitor, was non-inferior to warfarin in reducing the occurrence of the composite endpoint of stroke and systemic embolism. If non-inferiority was shown, a pre-specified step-wise multiple testing procedure was undertaken to determine whether rivaroxaban tablets was superior to warfarin in primary and secondary endpoints.

The study design, treatment regimen and patient populations are summarized in [Table 27](#) and [Table 28](#). A total of 14,264 patients were randomized with a mean age of 71 years (range 25 to 97 years) and a mean CHADS₂ score of 3.5. Patients were randomized to 20 mg once daily rivaroxaban tablets (15 mg in patients

with moderate renal impairment at screening) or to dose-adjusted warfarin, titrated to an INR of 2.0 to 3.0. ROCKET AF had a mean treatment duration of 572 days of rivaroxaban tablets given as a fixed dose without routine coagulation monitoring.

ROCKET AF studied patients with significant co-morbidities, e.g, 55% secondary prevention population (prior stroke / TIA / systemic embolism), see [Table 27](#). For patients randomized to warfarin, the time-in-therapeutic range (TTR) of 2.0 to 3.0 was a mean of 55% (cf. 64% in North American patients).

Table 27 – Co-morbid Diseases and Characteristics of Patients in ROCKET AF Trial – ITT Population

Heart failure and/or left ventricular ejection fraction \leq 35%	62%
Hypertension	91%
Age \geq 75 years	44%
Female	40%
Diabetes	40%
Prior Stroke / TIA / Systemic Embolism	55%
Stroke ^a	34%
TIA ^a	22%
Systemic Embolism ^a	4%
Valvular Disease (not meeting exclusion criteria) ^b	14%
Mean CHADS ₂	3.5
Prior VKA Use	62%
Prior MI	17%

a Some patients may have had more than one event, so sum of individual components does not add up to 55%.

b Patients with prosthetic heart valves, or those with hemodynamically significant rheumatic heart disease, especially mitral stenosis were excluded from ROCKET AF. Patient with other valvular disease including aortic stenosis, aortic regurgitation, and/or mitral regurgitation did not meet the exclusion criteria.

ITT Population= 14, 264 patients

At baseline, 36.5% of patients were on chronic ASA, 2.4% on anticoagulants other than VKAs, 8.7% on Class III antiarrhythmics, 54.5% on angiotensin converting enzyme (ACE) inhibitors, 22.7% on angiotensinreceptor blockers, 60.0% on diuretics, 24.0% on oral antidiabetics, and 65.5% on beta blockers.

ROCKET AF demonstrated that in patients with atrial fibrillation, rivaroxaban tablets is non-inferior to warfarin in the primary efficacy endpoint, a composite of prevention of stroke and systemic embolism in the per protocol population, on-treatment analysis (rivaroxaban: 1.71%/year, warfarin 2.16%/year, HR 0.79, 95% CI 0.66-0.96, $P < 0.001$). As non-inferiority was met, rivaroxaban tablets was tested, as per the pre-specified analysis, for superiority in primary and secondary endpoints. Rivaroxaban tablets demonstrated superiority over warfarin for stroke and systemic embolism in the safety population, on-treatment analysis (HR 0.79, 95% CI 0.65 to 0.95, $P = 0.015$), see [Table 28](#) and [Figure 7](#) below.

Table 28 – ROCKET AF – Time to the First Occurrence of Total Stroke and Systemic Embolism, While on Treatment (up to Last Dose Plus 2 Days) – Safety Population

Parameter	Rivaroxaban tablets (N=7061)		Warfarin (N=7082)		Rivaroxaban tablets vs Warfarin	
	n	%/year	n	%/year	Hazard Ratio (95% CI)	P-value for superiority
Total stroke and systemic embolism (Primary Efficacy Outcome)	189	1.70	243	2.15	0.79 (0.65,0.95)	0.015*
Total Stroke	184	1.65	221	1.96	0.85 (0.70,1.03)	0.092
Hemorrhagic Stroke	29	0.26	50	0.44	0.59 (0.37,0.93)	0.024*
Ischemic Stroke	149	1.34	161	1.42	0.94 (0.75,1.17)	0.581
Unknown Stroke Type	7	0.06	11	0.10	0.65 (0.25,1.67)	0.366
Systemic Embolism	5	0.04	22	0.19	0.23 (0.09,0.61)	0.003*
Other Endpoints						
All Cause Death	208	1.87	250	2.21	0.85 (0.70,1.02)	0.073
Vascular Death	170	1.53	193	1.71	0.89 (0.73,1.10)	0.289
Myocardial Infarction	101	0.91	126	1.12	0.81 (0.63,1.06)	0.121

Safety population on-treatment analysis = Events (Adjudicated by CEC) While on Treatment (up to Last Dose Plus 2 Days) – Safety Population


Hazard ratio (95% CI) and P-value from Cox proportional hazard model with treatment group as a covariate.

p-value (two-sided) for superiority of rivaroxaban tablets versus warfarin in hazard ratio

* Statistically significant

While the pre-specified primary analysis for superiority used the on-treatment data set for the safety population, an intention-to treat (ITT) analysis was also conducted. In this analysis, the primary endpoint occurred in 269 patients in the rivaroxaban group (2.1% per year) and in 306 patients in the warfarin group (2.4% per year) (hazard ratio, 0.88; 95% CI, 0.74 to 1.03; $P < 0.001$ for non-inferiority; $P = 0.12$ for superiority).

Figure 7: Kaplan-Meier curve of time to first total stroke or systemic embolism in the ROCKET AF trial safety population, on-treatment analysis, includes the 15 mg and 20 mg doses of rivaroxaban



The analysis of the principal safety endpoint demonstrates rivaroxaban tablets has a similar rate to warfarin for the composite of major and non-major clinically relevant bleeding, see [Table 29](#) below.

Table 29 – ROCKET AF – Time to the First Occurrence of Bleeding Events While on Treatment (up to Last Dose Plus 2 Days) – Safety Population

Parameter	Rivaroxaban tablets (N=7111)		Warfarin (N=7125)		Rivaroxaban tablets vs Warfarin	
	n	%/year	N	%/year	Hazard Ratio (95% CI)	P-value
Major and Non-major clinically relevant bleeding event (Principal Safety Endpoint)	1475	14.91	1449	14.52	1.03 (0.96,1.11)	0.442
Major Bleeding	395	3.60	386	3.45	1.04 (0.90,1.20)	0.576
Hemoglobin Drop (2g/dL)	305	2.77	254	2.26	1.22 (1.03,1.44)	0.019*
Transfusion (> 2 units)	183	1.65	149	1.32	1.25 (1.01,1.55)	0.044*
Critical Organ Bleed	91	0.82	133	1.18	0.69 (0.53,0.91)	0.007*
Intracranial Hemorrhage	55	0.49	84	0.74	0.67 (0.47, 0.94)	0.019*
Fatal Bleed	27	0.24	55	0.48	0.50 (0.31,0.79)	0.003*

Non-major Clinically Relevant Bleeding	1185	11.80	1151	11.37	1.04 (0.96,1.13)	0.345
--	------	-------	------	-------	------------------	-------

All analyses are based on the time to the first event.

Hemoglobin drop = a fall in hemoglobin of 2 g/dL or more.

Transfusion = a transfusion of 2 or more units of packed red blood cells or whole blood.

Critical organ bleeding are cases where CEC bleeding site=intracranial, intraspinal, intraocular, pericardial, intra-articular, intramuscular with compartment syndrome or retroperitoneal.

Hazard ratio (95% CI) and P-value from Cox proportional hazard model with treatment group as a covariate.

P-value (two-sided) for superiority of rivaroxaban tablets versus Warfarin in hazard ratio.

* Statistically significant

The incidences of increased liver function tests were low and comparable between the two groups, see [Table 30](#).

Table 30 – ROCKET AF – Incidence of Pre-specified Post-baseline Liver Function Abnormalities – Safety Population

Parameter	Rivaroxaban tablets (N=7111)		Warfarin (N=7125)		Rivaroxaban tablets vs Warfarin
	n/J	%	n/J	%	Hazard Ratio (95% CI)
ALT> 3xULN	203/6979	2.91	203/7008	2.90	1.01 (0.83,1.23)
ALT > 3xULN and TBL > 2xULN	31/6980	0.44	33/7012	0.47	0.95 (0.58,1.55)

ULN = Upper Limit of Normal Range, n = Number of patients with events, N= Number of patients valid for safety population, J = Number of patients with non-missing lab values, TBL: Total Bilirubin

Hazard Ratio (95% CI): time to event analysis using a Cox model with the treatment as the covariate.

The event rates for efficacy and safety outcomes stratified by age groups are presented in [Table 31](#) and [Table 32](#). The event rates for efficacy and safety outcomes stratified by renal function are presented in [Table 33](#) and [Table 34](#).

Table 31 – Efficacy Outcomes by Age Groups in the ROCKET AF Trial, While on Treatment (up to LastDose Plus 2 Days) – Safety Population

	Rivaroxaban tablets		Warfarin		Rivaroxaban tablets vs Warfarin	
	n/J	Event rate (%/yr)	n/J	Event rate (%/yr)	Hazard Ratio (95% CI)	P-value
Total Stroke and Systemic Embolism (Primary Efficacy Outcome)						
All Patients	189/7061	1.70	243/7082	2.15	0.79 (0.65,0.95)	0.015*
< 65 years	43/1642	1.59	42/1636	1.53	1.04 (0.68,1.58)	-
65 to 75 years	77/2767	1.74	98/2768	2.18	0.79 (0.59,1.07)	-
> 75 years	69/2652	1.73	103/2678	2.54	0.68 (0.50,0.92)	-
> 80 years	40/1305	2.17	46/1281	2.39	0.91 (0.60,1.40)	-
≥ 85 years	7/ 321	1.75	9/ 328	1.91	0.92 (0.34,2.47)	-
Total Stroke						
All Patients	184/7061	1.65	221/7082	1.96	0.85 (0.70,1.03)	0.092
< 65 years	42/1642	1.55	36/1636	1.31	1.18 (0.76,1.84)	-

	Rivaroxaban tablets		Warfarin		Rivaroxaban tablets vs Warfarin	
	n/J	Event rate (%/yr)	n/J	Event rate (%/yr)	Hazard Ratio (95% CI)	P-value
65 to 75 years	75/2767	1.69	90/2768	2.00	0.84 (0.62,1.14)	-
> 75 years	67/2652	1.68	95/2678	2.34	0.72 (0.52,0.98)	-
> 80 years	38/1305	2.06	42/1281	2.18	0.95 (0.61,1.48)	-
Ischemic Stroke						
All Patients	149/7061	1.34	161/7082	1.42	0.94 (0.75,1.17)	0.581
< 65 years	30/1642	1.11	23/1636	0.84	1.32(0.77,2.28)	-
65 to 75 years	68/2767	1.53	66/2768	1.47	1.04 (0.74,1.46)	-
> 75 years	51/2652	1.28	72/2678	1.77	0.72 (0.50,1.03)	-
> 80 years	26/1305	1.41	33/1281	1.71	0.83 (0.50,1.39)	-
Hemorrhagic Stroke						
All Patients	29/7061	0.26	50/7082	0.44	0.59 (0.37,0.93)	0.024*
< 65 years	9/1642	0.33	12/1636	0.44	0.76 (0.32,1.80)	-
65 to 75 years	4/2767	0.09	19/2768	0.42	0.21 (0.07,0.62)	-
> 75 years	16/2652	0.40	19/2678	0.47	0.86 (0.44,1.67)	-
> 80 years	12/1305	0.65	9/1281	0.47	1.40 (0.59,3.31)	-
Vascular Death						
All Patients	170/7061	1.53	193/7082	1.71	0.89 (0.73,1.10)	0.289
< 65 years	35/1642	1.29	44/1636	1.60	0.81 (0.52,1.26)	-
65 to 75 years	66/2767	1.49	70/2768	1.56	0.95 (0.68,1.33)	-
> 75 years	69/2652	1.73	79/2678	1.94	0.89 (0.64,1.23)	-
> 80 years	34/1305	1.84	35/1281	1.81	1.01 (0.63,1.62)	-
≥ 85 years	15/ 321	3.75	12/ 328	2.54	1.44 (0.67,3.08)	-

Safety population on-treatment analysis = Events (Adjudicated by CEC) While on Treatment (up to Last Dose Plus 2 Days) – Safety Population

n=number of patients with events, J=number of patients in each subgroup.

Hazard ratio (95% CI) and p-value from Cox proportional hazard model with treatment group as a covariate.

P-value (two-sided) for superiority of rivaroxaban tablets versus warfarin in hazard ratio

* Statistically significant

Table 32 – Bleeding Endpoints by Age Groups in the ROCKET AF trial, While on Treatment (up to Last Dose Plus 2 Days) – Safety Population

	Rivaroxaban tablets		Warfarin		Rivaroxaban tablets vs. Warfarin	
	n/J	Event rate (%/yr)	n/J	Event rate (%/yr)	Hazard Ratio (95% CI)	P-value
Major and Non-major Clinically Relevant Bleeding Event (Principal Safety Endpoint)						
All Patients	1475/7111	14.91	1449/7125	14.52	1.03 (0.96,1.11)	0.442

	Rivaroxaban tablets		Warfarin		Rivaroxaban tablets vs. Warfarin	
	n/J	Event rate (%/yr)	n/J	Event rate (%/yr)	Hazard Ratio (95% CI)	P-value
< 65 years	241/1646	9.73	260/1642	10.41	0.93 (0.78,1.11)	-
65 to 75 years	541/2777	13.59	556/2781	13.95	0.98 (0.87,1.10)	-
> 75 years	693/2688	20.18	633/2702	18.09	1.12(1.00,1.25)	-
> 80 years	362/1320	22.79	313/1298	18.84	1.20 (1.04,1.40)	-
≥85 years	89/ 326	25.46	90/ 335	22.29	1.13 (0.84,1.52)	-
Major Bleeding						
All Patients	395/7111	3.60	386/7125	3.45	1.04 (0.90,1.20)	0.576
< 65 years	59/1646	2.21	59/1642	2.16	1.02 (0.71,1.46)	-
65 to 75 years	133/2777	3.04	148/2781	3.34	0.91 (0.72,1.15)	-
> 75 years	203/2688	5.16	179/2702	4.47	1.15 (0.94,1.41)	-
> 80 years	118/1320	6.50	86/1298	4.50	1.44 (1.09,1.90)	-
≥85 years	28/ 326	7.05	32/ 335	6.91	1.01 (0.61,1.67)	-
Intracranial Hemorrhage						
All Patients	55/7111	0.49	84/7125	0.74	0.67 (0.47,0.93)	0.019*
< 65 years	13/1646	0.48	17/1642	0.62	0.78 (0.38,1.60)	-
65 to 75 years	13/2777	0.29	34/2781	0.75	0.39 (0.20,0.73)	-
> 75 years	29/2688	0.72	33/2702	0.81	0.89 (0.54,1.47)	-
> 80 years	22/1320	1.18	15/1298	0.77	1.54 (0.80,2.96)	-
Fatal Bleeding						
All Patients	27/7111	0.24	55/7125	0.48	0.50 (0.31,0.79)	0.003*
< 65 years	7/1646	0.26	11/1642	0.40	0.65 (0.25,1.66)	-
65 to 75 years	7/2777	0.16	19/2781	0.42	0.37 (0.16,0.89)	-
> 75 years	13/2688	0.32	25/2702	0.61	0.53 (0.27,1.03)	-
> 80 years	10/1320	0.54	12/1298	0.62	0.87 (0.38,2.02)	-
Non-major Clinically Relevant Bleeding						
All Patients	1185/7111	11.80	1151/7125	11.37	1.04 (0.96,1.13)	0.345
< 65 years	191/1646	7.62	210/1642	8.32	0.91 (0.75,1.11)	-
65 to 75 years	444/2777	11.00	445/2781	11.02	1.00 (0.88,1.14)	-
> 75 years	550/2688	15.74	496/2702	13.93	1.13 (1.00,1.28)	-
> 80 years	276/1320	17.06	249/1298	14.74	1.15 (0.97,1.37)	-

Safety population on-treatment analysis = Events (Adjudicated by CEC) While on Treatment (up to Last Dose Plus 2 Days) – Safety Population

n=number of patients with events, J=number of patients in each subgroup.

Hazard ratio (95% CI) and P-value from Cox proportional hazard model with treatment group as a covariate.

p-value (two-sided) for superiority of rivaroxaban tablets versus warfarin in hazard ratio

* Statistically significant

Table 33 – Efficacy Outcomes Stratified by Renal Function at Study Entry in the ROCKET AF Trial, While on Treatment (up to Last Dose Plus 2 Days) – Safety Population

	Rivaroxaban tablets		Warfarin		Rivaroxaban tablets vs Warfarin	
	n/J ^a	Event rate (%/yr)	n/J ^a	Event rate (%/yr)	Hazard Ratio (95% CI)	P-value
Total Stroke and Systemic Embolism (Primary Efficacy Outcome)						
All Patients	189/7061	1.70	243/7082	2.15	0.79 (0.65,0.95)	0.015*
30 – 49 mL/min	50/1481	2.36	60/1452	2.80	0.84 (0.58,1.22)	-
50 – 80 mL/min	91/3290	1.74	128/3396	2.39	0.73 (0.56,0.96)	-
> 80 mL/min	47/2278	1.25	54/2221	1.43	0.87 (0.59,1.28)	-
Total Stroke						
All Patients	184/7061	1.65	221/7082	1.96	0.85 (0.70,1.03)	0.092
30 – 49 mL/min	49/1481	2.31	52/1452	2.42	0.95 (0.64,1.40)	-
50 – 80 mL/min	88/3290	1.68	120/3396	2.24	0.75 (0.57,0.99)	-
> 80 mL/min	46/2278	1.22	48/2221	1.27	0.95 (0.64,1.43)	-
Ischemic Stroke						
All Patients	149/7061	1.34	161/7082	1.42	0.94 (0.75,1.17)	0.581
30 – 49 mL/min	43/1481	2.03	39/1452	1.82	1.11(0.72,1.72)	-
50 – 80 mL/min	69/3290	1.32	89/3396	1.66	0.80 (0.58,1.09)	-
> 80 mL/min	36/2278	0.95	32/2221	0.85	1.12 (0.70,1.80)	-
Hemorrhagic Stroke						
All Patients	29/7061	0.26	50/7082	0.44	0.59 (0.37,0.93)	0.024*
30 – 49 mL/min	6/1481	0.28	11/1452	0.51	0.55 (0.20,1.48)	-
50 – 80 mL/min	15/3290	0.29	25/3396	0.47	0.62 (0.33,1.17)	-
> 80 mL/min	8/2278	0.21	14/2221	0.37	0.57 (0.24,1.35)	-
Vascular Death						
All Patients	170/7061	1.53	193/7082	1.71	0.89 (0.73,1.10)	0.289
30 – 49 mL/min	55/1481	2.59	54/1452	2.52	1.02 (0.70,1.49)	-
50 – 80 mL/min	75/3290	1.43	91/3396	1.69	0.85 (0.62,1.15)	-
> 80 mL/min	40/2278	1.06	47/2221	1.24	0.85 (0.56,1.29)	-

Safety population on-treatment analysis = Events (Adjudicated by CEC) While on Treatment (up to Last Dose Plus 2Days) – Safety Population

n=number of patients with events, J=number of patients in each subgroup.

^a Patients with CrCl< 30 mL/min or missing baseline CrCl are excluded from the rows of CrCl subgroups (30-49mL/min, 50-80 mL/min, >80 mL/min). The patients are, however, included in the “All Patients” rows. Hazard ratio (95% CI) and p-value from Cox proportional hazard model with treatment group as a covariate. P-value (two-sided) for superiority of rivaroxaban tablets versus warfarin in hazard ratio

* Statistically significant

Table 34 – Bleeding Endpoints Stratified by Renal Function at Study Entry in the ROCKET AF Trial, While on Treatment (up to Last Dose Plus 2 Days) – Safety Population

	Rivaroxaban tablets		Warfarin		Rivaroxaban tablets vs Warfarin	
	n/J ^a	Event rate (%/yr)	n/J ^a	Event rate (%/yr)	Hazard Ratio (95% CI)	p-value
Major and Non-Major Clinically Relevant Bleeding Event (Principal Safety Endpoint)						
All Patients	1475/7111	14.91	1449/7125	14.52	1.03 (0.96,1.11)	0.442
30 – 49 mL/min	336/1498	17.87	341/1472	18.28	0.98 (0.84,1.14)	-
50 – 80 mL/min	725/3313	15.74	719/3410	15.30	1.04 (0.93,1.15)	-
> 80 mL/min	412/2288	12.15	388/2230	11.42	1.06(0.92,1.21)	-
Major Bleeding						
All Patients	395/7111	3.60	386/7125	3.45	1.04 (0.90,1.20)	0.576
30 – 49 mL/min	99/1498	4.72	100/1472	4.72	1.00 (0.76,1.32)	-
50 – 80 mL/min	183/3313	3.54	197/3410	3.72	0.95 (0.78,1.17)	-
> 80 mL/min	112/2288	3.02	89/2230	2.38	1.26 (0.95,1.67)	-
Intracranial Hemorrhage						
All Patients	55/7111	0.49	84/7125	0.74	0.67 (0.47,0.93)	0.019*
30 – 49 mL/min	15/1498	0.70	19/1472	0.88	0.80 (0.41,1.57)	-
50 – 80 mL/min	27/3313	0.51	43/3410	0.80	0.64 (0.40,1.04)	-
> 80 mL/min	13/2288	0.34	22/2230	0.58	0.59 (0.30,1.17)	-
Fatal Bleeding						
All Patients	27/7111	0.24	55/7125	0.48	0.50 (0.31,0.79)	0.003*
30 – 49 mL/min	6/1498	0.28	16/1472	0.74	0.38 (0.15,0.97)	-
50 – 80 mL/min	14/3313	0.27	24/3410	0.45	0.60 (0.31,1.16)	-
> 80 mL/min	7/2288	0.19	15/2230	0.40	0.46 (0.19,1.14)	-
Non-major Clinically Relevant Bleeding						
All Patients	1185/7111	11.80	1151/7125	11.37	1.04 (0.96,1.13)	0.345
30 – 49 mL/min	261/1498	13.67	259/1472	13.61	1.01 (0.85,1.19)	-
50 – 80 mL/min	596/3313	12.77	570/3410	11.94	1.08 (0.96,1.21)	-
> 80 mL/min	327/2288	9.48	321/2230	9.36	1.01 (0.86,1.18)	-

Safety population on-treatment analysis = Events (Adjudicated by CEC) While on Treatment (up to Last Dose Plus 2 Days) – Safety Population

n=number of patients with events, J=number of patients in each subgroup

^a Patients with CrCl < 30 mL/min or missing baseline CrCl are excluded from the rows of CrCl subgroups (30-49 mL/min, 50-80 mL/min, >80 mL/min). The patients are, however, included in the “All Patients” rows. Hazard ratio (95% CI) and p-value from Cox proportional hazard model with treatment group as a covariate. P-value (two-sided) for superiority of rivaroxaban tablets versus warfarin in hazard ratio

* Statistically significant

PHASE IV STUDY

Phase IV clinical studies (XANTUS) was done to evaluate the effects of rivaroxaban use under real-world (clinical practice) conditions.

XANTUS

In addition to the Phase III ROCKET AF study, a prospective, single-arm, post-authorization, non-interventional, open-label cohort study (XANTUS) with central outcome adjudication including thromboembolic events and major bleeding has been conducted. 6,704 patients with non-valvular atrial fibrillation were enrolled for prevention of stroke and non-central nervous system (CNS) systemic embolism under real-world conditions. The mean CHADS2 score of the population was 1.9. Major bleeding incidence was 2.1 per 100 patient years. Fatal hemorrhage incidence was 0.2 per 100 patient years and intracranial hemorrhage incidence was 0.4 per 100 patient years. Stroke or non-CNS systemic embolism incidence was 0.8 per 100 patient years. These results are consistent with the established safety profile of rivaroxaban tablets in this population.

14.2 Comparative Bioavailability Studies

A two-treatment, two-sequence, single-dose (1 x 10 mg) crossover comparative bioavailability study of JAMP Rivaroxaban, 10 mg tablets (JAMP Pharma Corporation, Canada), and ^{Pr}XARELTO® (Rivaroxaban), 10 mg tablets (Bayer Inc., Canada), was conducted in healthy adult Asian males under fasting conditions. A summary of the rivaroxaban bioavailability data from 29 subjects is presented in the following table:

SUMMARY TABLE OF THE COMPARATIVE BIOAVAILABILITY DATA

Rivaroxaban (1 × 10 mg) Geometric Mean Arithmetic Mean (CV %)				
Parameter	Test ¹	Reference ²	% Ratio of Geometric Means	90% Confidence Interval
AUC _T (ng•h/mL)	1400.60 1630.73 (31.52)	1502.73 1695.24 (29.90)	93.0	87.2 - 99.1
AUC _I (ng•h/mL)	1483.54 1685.12 (30.44)	1552.80 1738.30 (29.30)	95.3	90.8 - 100.2
C _{max} (ng/mL)	177.41 200.59 (31.94)	202.53 224.37 (31.23)	87.2	80.5 - 94.3
T _{max} ³ (h)	2.00 (1.00 - 4.52)	2.67 (1.00 - 4.52)		
T _½ ⁴ (h)	7.08 (44.00)	5.47 (28.32)		

¹ JAMP Rivaroxaban (Rivaroxaban) tablets 10 mg (JAMP Pharma Corporation, Canada)

² ^{Pr}XARELTO® (Rivaroxaban) tablets 10 mg (Bayer Inc., Canada)

³ Expressed as the median (range) only

⁴ Expressed as the Arithmetic mean (CV%) only

A two-treatment, two-sequence, single-dose (1 x 20 mg) crossover comparative bioavailability study of JAMP Rivaroxaban, 20 mg tablets (JAMP Pharma Corporation, Canada), and PrXARELTO® (Rivaroxaban), 20 mg tablets (Bayer Inc., Canada), was conducted in healthy adult Asian males under high fat, high calorie fed conditions. A summary of the bioavailability data from 21 subjects is presented in the following table:

SUMMARY TABLE OF THE COMPARATIVE BIOAVAILABILITY DATA

Rivaroxaban (1 x 20 mg) Geometric Mean Arithmetic Mean (CV %)				
Parameter	Test ¹	Reference ²	% Ratio of Geometric Means	90% Confidence Interval
AUC _T (ng•h/mL)	4225.22 4345.45 (24.81)	4236.13 4347.15 (25.51)	100.1	96.4-103.9
AUC _I (ng•h/mL)	4291.04 4416.84 (25.24)	4284.82 4398.63 (25.69)	100.5	97.0-104.1
C _{max} (ng/mL)	411.86 420.63 (20.72)	428.19 438.92 (22.67)	96.7	89.4-104.6
T _{max} ³ (h)	4.33 (2.00-8.02)	4.00 (2.00-8.00)		
T _½ ⁴ (h)	7.00 (42.94)	6.33 (22.36)		

¹ JAMP Rivaroxaban (Rivaroxaban) tablets, 20 mg (JAMP Pharma Corporation, Canada)

² PrXARELTO® (Rivaroxaban) tablets, 20 mg (Bayer Inc., Canada)

³ Expressed as the median (range) only

⁴ Expressed as the Arithmetic mean (CV%) only

15 MICROBIOLOGY

No microbiological information is required for this drug product.

16 NON-CLINICAL TOXICOLOGY

General Toxicology (repeat-dose studies)

Rivaroxaban was tested in repeat-dose studies up to 6 months in rats and up to 12 months in dogs. Based on the pharmacological mode of action, a NOEL could not be established due to effects on clotting time. All adverse findings, except for a slight body weight gain reduction in rats and dogs, could be related to an exaggerated pharmacological mode of action of the compound. In dogs, at very high exposures, severe spontaneous bleedings were observed. The NOAELs after chronic exposure are 12.5 mg/kg in rats and 5 mg/kg in dogs.

Carcinogenicity

In 2-year carcinogenicity studies, rivaroxaban was tested in mice, up to 60 mg/kg/day (reaching systemic exposure similar to humans) and in rats (up to 3.6-fold higher than in humans) without demonstration of carcinogenic potential.

Genotoxicity

No genotoxicity was observed in a test for gene mutation in bacteria (Ames-Test), in an in vitro test for chromosomal aberrations, or in the in vivo micronucleus test.

Reproductive and Developmental Toxicology

Rivaroxaban was tested in developmental toxicity studies at exposure levels of up to 38-fold (rat) and up to 89-fold (rabbit) above the therapeutic exposure in humans. The toxicological profile is mainly characterized by maternal toxicity due to exaggerated pharmacodynamic effects.

Up to the highest dose tested, no primary teratogenic potential was identified.

[¹⁴C]Rivaroxaban -related radioactivity penetrated the placental barrier in rats. In none of the fetal organs and tissues did the exposure in terms of maximum concentrations or AUC exceed the maternal blood exposure. The average exposure in the fetuses based on AUC₍₀₋₂₄₎ reached about 20% of the exposure in maternal blood. The AUC in the mammary glands was approximately equivalent to the AUC in the blood, which indicates secretion of radioactivity into milk (see [2 CONTRAINDICATIONS](#)).

Rivaroxaban did not show an effect on male or female fertility up to 200 mg/kg.

[¹⁴C]Rivaroxaban was administered orally to lactating Wistar rats (day 8 to 10 post partum) as a single oral dose of 3 mg/kg body weight.

[¹⁴C]Rivaroxaban -related radioactivity was secreted into the milk of lactating rats only to a low extent in relation to the administered dose: The estimated amount of radioactivity excreted with milk was 2.12% of the maternal dose within 32 hours after administration (see [2 CONTRAINDICATIONS](#)).

Juvenile Toxicity

Rivaroxaban was tested in male and female juvenile rats up to 3-month treatment duration. Treatment started at postnatal day 4 at doses of 6, 20 and 60 mg/kg/day. Rivaroxaban was generally well tolerated except for signs indicating exaggerated pharmacodynamics (increased coagulation time). No evidence of target organ-specific toxicity or toxicity developing organs was seen.

17 SUPPORTING PRODUCT MONOGRAPHS

1. Xarelto[®] (Tablets, 2.5 mg, 10 mg, 15 mg, 20 mg), submission control 268349, Product Monograph, (Bayer Inc.) April 17, 2023.

PATIENT MEDICATION INFORMATION
READ THIS FOR SAFE AND EFFECTIVE USE OF YOUR MEDICINE

PrJAMP Rivaroxaban

Rivaroxaban Tablets

Read this carefully before you start taking **JAMP Rivaroxaban** and each time you get a refill. This leaflet is a summary and will not tell you everything about this drug. Talk to your healthcare professional about your medical condition and treatment and ask if there is any new information about **JAMP Rivaroxaban**.

What is JAMP Rivaroxaban used for?

JAMP Rivaroxaban 10 mg, 15 mg and 20 mg tablets are used in adults to:

- prevent blood clots after major hip or knee surgery
- treat and prevent blood clots in the veins of the legs, arms or lungs
- prevent blood clots in the brain (stroke) and other blood vessels in the body of patients who have atrial fibrillation (irregular heart rhythm)

How does JAMP Rivaroxaban work?

JAMP Rivaroxaban belongs to a group of medicines called anticoagulants (blood thinners). It helps prevent bloodclots from forming by directly blocking the activity of a clotting factor called Factor-Xa.

What are the ingredients in JAMP Rivaroxaban?

Medicinal ingredient: Rivaroxaban

Non-medicinal ingredients: Cellulose microcrystalline, croscarmellose sodium, hypromellose, iron oxide red, lactose monohydrate, magnesium stearate, polyethylene glycol, sodium lauryl sulfate and titanium dioxide.

JAMP Rivaroxaban comes in the following dosage forms:

- Tablets: 10 mg, 15 mg and 20 mg

Do not use JAMP Rivaroxaban if you:

- Are allergic to rivaroxaban or to any other ingredients in JAMP Rivaroxaban. JAMP Rivaroxaban contains lactose.
- Are actively bleeding, especially if you are excessively bleeding.
- Have any injuries or conditions that may increase your risk of bleeding. This includes:
 - active or recent bleeding in the brain
 - active or recent bleeding in the stomach or gut
 - have difficulties to stop bleeding (bleeding disorder)
- Take medicines to treat HIV / AIDS, such as cobicistat and ritonavir.
- Take medicines to treat fungal infections, such as ketoconazole, itraconazole and posaconazole.
- Take dronedarone, a medicine used to control an abnormal heart rhythm called atrialfibrillation.
- Take any other medicines used to prevent or treat blood clots. This includes:
 - warfarin, dabigatran, apixaban, edoxaban, heparin
 - low molecular weight heparin (LMWH), such as enoxaparin and dalteparin
 - heparin derivatives, such as fondaparinux

- Have severe liver disease, which can lead to an increased risk of bleeding.
- Are pregnant, think you might be pregnant or plan on becoming pregnant.
- Are breastfeeding.

To help avoid side effects and ensure proper use, talk to your healthcare professional before you take JAMP Rivaroxaban. Talk about any health conditions or problems you may have, including if you:

- Have an increased risk of bleeding. This includes if you:
 - have severe kidney problems or reduced kidney function
 - are taking Non-Steroidal Anti-Inflammatory Drugs (NSAIDs), which are used to relieve pain and reduce inflammation, such as ibuprofen, naproxen and celecoxib
 - are taking acetylsalicylic acid (ASA, or ASPIRIN) or any other medicines used to prevent stroke or heart disease, such as clopidogrel, prasugrel and ticagrelor
 - are taking medicines to treat depression or other psychological conditions
 - have low levels of platelets in the blood or your platelets do not work properly
 - have very high blood pressure, not controlled by medical treatment
 - have or have had an ulcer in the stomach or bowel
 - have a problem with the blood vessels in the back of the eyes (retinopathy)
 - have problems with the blood vessels in the brain or spinal column
 - had surgery on the brain, spinal column or eyes
 - have bronchiectasis, which is a chronic disease where the bronchial tubes of the lungs become damaged, widened, and thickened
 - have a history of bleeding into the lungs
 - are older than 75 years of age
 - have been diagnosed with cancer, had a relapse of cancer or have been treated for cancer in the last 6 months
 - have liver problems
 - are having surgery for any reason, including an operation that involves a catheter or injection into the spinal column (e.g., for epidural or spinal anesthesia or pain reduction)
- Have damaged heart valves or heart valve disease.
- Have a prosthetic heart valve.
- Have antiphospholipid syndrome, also known as Hughes syndrome.
This is a disorder of the immune system that causes an increased risk of blood clots.
- Have a stent, which is a small metal coil inserted into a blocked blood vessel to restore blood flow by keeping it open.
- Have a history of stroke, with or without bleeding in the brain.
- Are taking rifampicin, a medicine used to treat bacterial infections, including tuberculosis.
- Are taking medicines used to prevent epilepsy or seizures, such as phenytoin, carbamazepine and phenobarbital.
- Are lactose intolerant. JAMP Rivaroxaban contains lactose.

Other warnings you should know about:

Do not stop taking JAMP Rivaroxaban without first talking to your healthcare professional. If you stop taking JAMP Rivaroxaban, blood clots may form and cause a stroke, heart attack, or other serious complications. This can lead to severe disability or even death.

Risk of Bleeding: As with other blood thinners, taking JAMP Rivaroxaban may increase the risk of bleeding from any part of the body, including internal organs. It may be serious or even life-threatening. Tell your healthcare professional **right away** if you have any unusual bleeding or bruising. For examples of bleeding, see the **Serious side effects and what to do about them** table below.

Spinal or epidural blood clots (hematoma): People who take JAMP Rivaroxaban and have medicine injected into their spinal or epidural area, or have a spinal puncture, have a risk of forming a spinal or epidural blood clot. This can cause long-term or permanent loss of the ability to move (paralysis). The risk of developing a spinal or epidural blood clot is higher if:

- A thin tube called an epidural catheter is placed in your back to give certain medicines.
- You take NSAIDs or a medicine to prevent blood from clotting.
- You have a history of difficult or repeated epidural or spinal punctures.
- You have a history of problems with the spinal column or have had surgery on the spinal column.

If you take JAMP Rivaroxaban and receive spinal anesthesia or have a spinal puncture, your healthcare professional will watch you closely for symptoms of spinal or epidural blood clots.

Tell your healthcare professional **right away** if you develop the following symptoms after an epidural or a spinal procedure:

- Back pain
- Tingling
- Numbness
- Muscle weakness (especially in your legs and feet)
- Or loss of control of the bowels or bladder (incontinence)

Surgery:

- Tell your healthcare professional if you are going to have any surgeries, including dental procedures, while taking JAMP Rivaroxaban. There is an increased risk of bleeding if you take JAMP Rivaroxaban during these procedures. It can be life-threatening.
- Your healthcare professional may ask you to stop taking JAMP Rivaroxaban before the surgery, and to resume taking JAMP Rivaroxaban after the procedure. It is important that you take JAMP Rivaroxaban before and after the procedure exactly at the times the healthcare professional has instructed.

Children and adolescents:

- JAMP Rivaroxaban should **not** be used in children and adolescents under 18 years of age for any conditions.

Pregnancy:

- JAMP Rivaroxaban should **not** be used during pregnancy. Taking it during pregnancy may harm an unborn baby.
- Your healthcare professional may ask you, to use a highly effective birth control method while taking JAMP Rivaroxaban.

- If you discover that you are pregnant while taking JAMP Rivaroxaban, contact your healthcare professional **as soon as possible**.

Breastfeeding: JAMP Rivaroxaban should **not** be used while breastfeeding. It may pass into breast milk and harm a breastfed baby. JAMP Rivaroxaban will only be given to you once you have stopped breastfeeding.

Check-ups and testing: Your healthcare professional may:

- evaluate your risk of bleeding before starting JAMP Rivaroxaban and regularly during treatment. This includes performing blood tests to check the amount of red blood cells in the blood, and monitoring blood pressure.
- do blood tests before you start taking JAMP Rivaroxaban and during treatment to check if your kidneys are working properly.

Depending on the test results, your healthcare professional may discontinue your treatment with JAMP Rivaroxaban.

Tell your healthcare professional about all the medicines you take, including any drugs, vitamins, minerals, natural supplements or alternative medicines.

The following may interact with JAMP Rivaroxaban:

- Non-Steroidal Anti-Inflammatory Drugs (NSAIDs), which are used to relieve pain and reduce inflammation, such as ibuprofen, naproxen and celecoxib.
- Acetylsalicylic acid (ASA, or ASPIRIN), used to relieve fever and pain.
- Medicines used to prevent stroke or heart disease, such as clopidogrel, prasugrel and ticagrelor.
- Medicines used to prevent or treat blood clots. This includes:
 - warfarin, dabigatran, apixaban, edoxaban, heparin
 - low molecular weight heparin (LMWH), such as enoxaparin and dalteparin
 - heparin derivatives, such as fondaparinux
- Medicines used to treat HIV / AIDS, such as cobicistat and ritonavir.
- Medicines used to treat fungal infections, such as ketoconazole, itraconazole and posaconazole.
- Dronedaron, a medicine used to control an abnormal heart rhythm called atrial fibrillation.
- Medicines used to treat bacterial infections, such as clarithromycin and rifampicin.
- Medicines used to control epilepsy or seizures, such as phenytoin, carbamazepine and phenobarbital.
- Medicines used to treat depression or other psychological conditions, such as Selective Serotonin Reuptake Inhibitors (SSRIs) and Serotonin Norepinephrine Reuptake Inhibitors (SNRIs).
- St. John's Wort (also known as *Hypericum perforatum*), a herbal remedy used to treat depression.

How to take JAMP Rivaroxaban:

- Take:
 - JAMP Rivaroxaban tablets exactly as your healthcare professional tells you
 - JAMP Rivaroxaban tablets at the same time every day to help you remember it
 - JAMP Rivaroxaban 10 mg tablets **with or without** food
 - JAMP Rivaroxaban 15 mg and 20 mg tablets **with** food

- Swallow tablet whole, preferably with water. Do **not** split tablets.
- JAMP Rivaroxaban tablets may be crushed and mixed with applesauce. Take the applesauce mixture right away after you have mixed it. A crushed 10 mg tablet can be taken with or without food. Eat food right after taking a crushed 15 mg or 20 mg tablet. The healthcare professional may give you the crushed JAMP Rivaroxaban tablet also via a tube.
- Do **not** stop taking or giving JAMP Rivaroxaban without talking to the healthcare professional first.

Usual dose:

- **Prevention of blood clots after major hip or knee surgery**

Usual adult dose: Take one 10 mg tablet once a day with or without food.

Take the first tablet 6 to 10 hours after your operation. Then take a tablet every day until your healthcare professional tells you to stop.

If you have had a major hip operation, you will usually take JAMP Rivaroxaban for 35 days.

If you have had a major knee operation, you will usually take JAMP Rivaroxaban for 14 days.

- **Treatment and prevention of blood clots in the veins of the legs or arms, or blood vessels of the lungs**

Adults:

- **Day 1 to 21:** Take one 15 mg tablet TWICE a day (in the morning and evening) with food.
- **Day 22 and onwards:** Take one 20 mg tablet ONCE a day with food.

After at least 6 months treatment, your healthcare professional may decide to continue treatment with either one 20 mg tablet once a day or one 10 mg tablet once a day.

The 10 mg tablet may be taken with or without food.

This is a long-term treatment, and you should continue to take JAMP Rivaroxaban until your healthcare professional says otherwise.

- **Prevention of blood clots in your brain (stroke) and in other blood vessels of your body if you have atrial fibrillation**

Usual adult dose: Take one 20 mg tablet once a day with food.

If your kidneys are not working properly, your healthcare professional may prescribe 15 mg once a day with food.

This is a long-term treatment, and you should continue to take JAMP Rivaroxaban until your healthcare professional says otherwise.

The recommended maximum daily dose is 20 mg.

Overdose:

Taking too much JAMP Rivaroxaban may cause excessive bleeding.

If you think you, or a person you are caring for, have taken too much JAMP Rivaroxaban, contact a healthcare professional, hospital emergency department, or regional poison control centre immediately, even if there are no symptoms.

Missed Dose:

- If you take JAMP Rivaroxaban 15 mg tablets **twice** a day and miss a dose, take the missed dose as soon as you remember. If you forget to take a dose, you can take two 15 mg tablets at the same time to get a total of two tablets (30 mg) in one day. Do not take more than two 15 mg tablets in one day. The next day, take JAMP Rivaroxaban as usual.
- If you take JAMP Rivaroxaban 10 mg, 15 mg or 20 mg tablets **once** a day and miss a dose, take the missed dose as soon as you remember on the same day. If this is not possible, you should skip the missed dose and take the next dose on the following day as usual. Do not take a double dose to make up for the one that was missed.

What are possible side effects from using JAMP Rivaroxaban?

These are not all the possible side effects you may have when taking JAMP Rivaroxaban. If you experience any side effects not listed here, tell your healthcare professional.

As JAMP Rivaroxaban acts on the blood clotting system, most side effects are related to signs of bruising or bleeding. In some cases, bleeding may not be obvious, such as unexplained swelling.

Patients treated with JAMP Rivaroxaban may also experience the following side effects: Nausea, vomiting, stomach ache, constipation, diarrhea, indigestion, and decreased general strength and energy.

JAMP Rivaroxaban can cause abnormal blood test results. The healthcare professional may perform blood tests before you take JAMP Rivaroxaban and regularly during treatment. They will tell you if the test results are abnormal and if you need treatment to correct these side effects.

Serious side effects and what to do about them			
Symptom / effect	Talk to your healthcare professional		Stop taking drug and get immediate medical help
	Only if severe	In all cases	
COMMON			
Anemia (decreased number of red blood cells): pale complexion, weakness, tiredness, dizziness, headache, breathlessness, unusually fast heartbeat, or chest pain		✓	
Bleeding from hemorrhoids	✓		
Bleeding from the surgical wound, an injury or other medical procedure		✓	
Bleeding gums for longer than 5 minutes when you brush your teeth		✓	

Serious side effects and what to do about them			
Symptom / effect	Talk to your healthcare professional		Stop taking drug and get immediate medical help
	Only if severe	In all cases	
Bleeding in the eye	✓		
Bleeding under the skin	✓		
Blood in your urine, (red / pink tinge to urine)		✓	
Fever		✓	
Gastrointestinal (GI) bleeding (bleeding anywhere along the GI tract between mouth and anus): blood in vomit, black tarry stool, bright red blood in your stool or coming from rectum, rapid pulse, low blood pressure, low urine flow, confusion, weakness, dizziness		✓	
Genital bleeding in postmenopausal women		✓	
Hypotension (low blood pressure): dizziness, fainting, lightheadedness, blurred vision, nausea, vomiting, fatigue (may occur when you go from lying or sitting to standing up)		✓	
Increased or more frequent menstrual bleeding	✓		
Itchy skin or rash		✓	
Localized swelling		✓	
Nosebleed lasting more than 5 minutes		✓	
Pain or swelling in your limbs		✓	
Tachycardia (abnormally fast heartbeat)		✓	
Unexpected bruising		✓	
UNCOMMON			
Decreased urine output	✓		
Hemarthrosis (bleeding into a joint): joint pain, swelling		✓	
Hemoptysis : coughing up blood		✓	

Serious side effects and what to do about them			
Symptom / effect	Talk to your healthcare professional		Stop taking drug and get immediate medical help
	Only if severe	In all cases	
Intracerebral hemorrhage (bleeding in the brain): sudden, severe headache; confusion; nausea and vomiting; seizures; loss of consciousness			✓
Oozing from the surgical wound		✓	
Seizure (fit) reported in adults: loss of consciousness with uncontrollable shaking			✓
Syncope (fainting): a temporary loss of consciousness due to a sudden drop in blood pressure		✓	
RARE			
Allergic Reaction: difficulty swallowing or breathing, wheezing; drop in blood pressure; feeling sick to your stomach and throwing up; hives or rash; swelling of the face, lips, tongue or throat			✓
Liver Disorder: yellowing of the skin or eyes, dark urine, pale stools, abdominal pain, nausea, vomiting, loss of appetite		✓	
UNKNOWN FREQUENCY			
Agranulocytosis (decrease in white blood cells): frequent infection with fever, chills, sore throat		✓	
Compartment Syndrome: increased pressure within legs or arms after a bleed, with pain, swelling, numbness or paralysis		✓	
Eosinophilic Pneumonia (accumulation of a type of white blood cell that causes inflammation in the lung): shortness of breath, cough, wheezing, chest pain, fever, and unintended weight loss		✓	

Serious side effects and what to do about them			
Symptom / effect	Talk to your healthcare professional		Stop taking drug and get immediate medical help
	Only if severe	In all cases	
Stevens-Johnson Syndrome (SJS) (severe skin rash): redness, blistering and/or peeling of the skin and/or inside of the lips, eyes, mouth, nasal passages or genitals, accompanied by fever, chills, headache, cough, body aches or swollen glands			✓

If you have a troublesome symptom or side effect that is not listed here or becomes bad enough to interfere with your daily activities, tell your healthcare professional.

Reporting Side Effects

You can report any suspected side effects associated with the use of health products to Health Canada by:

- Visiting the Web page on Adverse Reaction Reporting (<https://www.canada.ca/en/health-canada/services/drugs-health-products/medeffect-canada/adverse-reaction-reporting.html>) for information on how to report online, by mail or by fax; or
- Calling toll-free at 1-866-234-2345.

NOTE: Contact your health professional if you need information about how to manage your side effects. The Canada Vigilance Program does not provide medical advice.

Storage:

- Keep JAMP Rivaroxaban tablets at room temperature, 15-30°C.
- Keep out of reach and sight of children.
- Do not use JAMP Rivaroxaban after the expiry date which is stated on the bottle after EXP. The expiry date refers to the last day of that month.
- Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help to protect the environment.

If you want more information about JAMP Rivaroxaban:

- Talk to your healthcare professional.
- Find the full Product Monograph that is prepared for healthcare professionals and includes this Patient Medication Information by visiting the Health Canada website: (<https://www.canada.ca/en/health-canada/services/drugs-health-products/drug-products/drug-product-database.html>); or by calling 1-866-399-9091.

This leaflet was prepared by:

JAMP Pharma Corporation
1310, rue Nobel
Boucherville, Quebec,
J4B 5H3, Canada

Last Revised: January 23, 2024