

PRODUCT MONOGRAPH
INCLUDING PATIENT MEDICATION INFORMATION

^{Pr} **GLICLAZIDE MR**

Gliclazide

Modified-release tablets 30 mg

Modified-release breakable tablets 60 mg

Hypoglycemic sulfonylurea

Oral antidiabetic agent

Sanis Health Inc.
1 President's Choice Circle
Brampton, Ontario
L6Y 5S5

Date of Initial Authorization:
January 27, 2022

Date of Revision:
MAR 27, 2024

Submission Control Number: 280770

RECENT MAJOR LABEL CHANGES

Non Applicable

TABLE OF CONTENTS

Sections or subsections that are not applicable at the time of authorization are not listed.

RECENT MAJOR LABEL CHANGES	2
TABLE OF CONTENTS	2
PART I: HEALTH PROFESSIONAL INFORMATION	4
1 INDICATIONS	4
1.1 Pediatrics (< 18 years of age)	4
1.2 Geriatrics (≥ 65 years of age)	4
2 CONTRAINDICATIONS	4
4 DOSAGE AND ADMINISTRATION	5
4.1 Dosing Considerations	5
4.2 Recommended Dose and Dosage Adjustment.....	5
4.4 Administration	5
4.5 Missed Dose.....	6
5 OVERDOSAGE	7
6 DOSAGE FORMS, STRENGTHS, COMPOSITION AND PACKAGING	8
7 WARNINGS AND PRECAUTIONS	8
7.1 Special Populations	13
7.1.1 Pregnant Women.....	13
7.1.2 Breast-feeding.....	13
7.1.3 Pediatrics (< 18 years of age)	13
7.1.4 Geriatrics (≥ 65 years of age)	13
8 ADVERSE REACTIONS	14
8.1 Adverse Reaction Overview	14
8.2 Clinical Trial Adverse Reactions	14

8.3	Less Common Clinical Trial Adverse Reactions	17
8.4	Abnormal Laboratory Findings: Hematologic, Clinical Chemistry and Other Quantitative Data	18
8.5	Post-Market Adverse Reactions.....	18
9	DRUG INTERACTIONS	19
9.1	Serious Drug Interactions.....	19
9.2	Drug Interactions Overview	19
9.3	Drug-Behavioural Interactions	20
9.4	Drug-Drug Interactions	20
9.5	Drug-Food Interactions	23
9.6	Drug-Herb Interactions	23
9.7	Drug-Laboratory Test Interactions.....	23
10	CLINICAL PHARMACOLOGY.....	24
10.1	Mechanism of Action	24
10.2	Pharmacodynamics	25
10.3	Pharmacokinetics.....	26
11	STORAGE, STABILITY AND DISPOSAL	27
12	SPECIAL HANDLING INSTRUCTIONS	27
PART II: SCIENTIFIC INFORMATION		28
13	PHARMACEUTICAL INFORMATION	28
14	CLINICAL TRIALS	28
14.1	Clinical Trials	29
14.3	Comparative Bioavailability Studies	30
15	MICROBIOLOGY.....	31
16	NON-CLINICAL TOXICOLOGY	31
PATIENT MEDICATION INFORMATION		41

PART I: HEALTH PROFESSIONAL INFORMATION

1 INDICATIONS

GLICLAZIDE MR (gliclazide modified-release tablets 30 and 60 mg) is indicated for:

- Control of hyperglycemia in gliclazide responsive diabetes mellitus of stable, mild, non-ketosis prone, maturity onset or adult type which cannot be controlled by proper dietary management and exercise, or when insulin therapy is not appropriate.

1.1 Pediatrics (< 18 years of age)

Safety and effectiveness of GLICLAZIDE MR in pediatrics have not been established; therefore, Health Canada has not authorized an indication for pediatric use.

1.2 Geriatrics (≥ 65 years of age)

No significant differences in efficacy and tolerance were observed between patients over 65 years of age and younger patients, however greater sensitivity of some older individuals cannot be ruled out (see [4 DOSAGE AND ADMINISTRATION](#) and [7.1.4 Geriatrics](#)).

2 CONTRAINDICATIONS

GLICLAZIDE MR is contraindicated in patients:

- With known hypersensitivity or allergy to gliclazide, other sulfonylureas, sulfonamides, or to any ingredient in the formulation, including any non-medical ingredient, or component of the container. For a complete listing, see [6 DOSAGE FORMS, STRENGTHS, COMPOSITION AND PACKAGING](#).
- With unstable and/or insulin-dependent (Type 1) diabetes mellitus, particularly juvenile diabetes, diabetic ketoacidosis, diabetic pre-coma and coma.
- During stress conditions such as serious infection, trauma or surgery.
- In the presence of severe hepatic impairment (see [7 WARNINGS AND PRECAUTIONS, Hepatic/Biliary/Pancreatic](#)).
- In the presence of severe renal impairment (see [7 WARNINGS AND PRECAUTIONS, Renal](#)).
- Treated with miconazole via systemic route or oromucosal gel (see [9.1 Serious Drug Interactions, 9.4 Drug-Drug Interactions](#)).
- During pregnancy and lactation (see [7.1.1 Pregnant Women](#) and [7.1.2 Breast-](#)

[feeding](#)).

4 DOSAGE AND ADMINISTRATION

4.1 Dosing Considerations

Determination of the proper dosage for GLICLAZIDE MR for each patient should be made on the basis of frequent determinations of blood glucose during dose titration and throughout maintenance.

The daily dose of GLICLAZIDE MR may vary from 30 to 120 mg once daily (i.e., one half tablet to 2 tablets of GLICLAZIDE MR 60 mg, or 1 to 4 tablets of GLICLAZIDE MR 30 mg).

4.2 Recommended Dose and Dosage Adjustment

The recommended starting dose of GLICLAZIDE MR is 30 mg daily, i.e. one half tablet of GLICLAZIDE MR 60 mg or one tablet of GLICLAZIDE MR 30 mg, even in elderly patients (over 65 years old).

A single daily dose provides effective blood glucose control. The single daily dose may be between 30 mg and 120 mg. The daily dose should not exceed 120 mg.

Dose adjustment should be carried out in steps of 30 mg, according to the blood glucose response. Each step should last for at least two weeks.

4.4 Administration

It is recommended that the medication be taken at breakfast time. The 30 mg tablets cannot be split in half and should be swallowed whole. The 60 mg tablets can be halved. Both the 30 mg and 60 mg tablets must not be chewed or crushed.

- Previously untreated patients should commence with a dose of 30 mg and will benefit from dose adjustment until the appropriate dose is reached.
- One GLICLAZIDE MR 60 mg tablet is equivalent to two GLICLAZIDE MR 30 mg tablets. The breakability of the GLICLAZIDE MR modified-release 60 mg tablet allows the use of a dose of 30 mg as a half tablet and of 90 mg as one and a half tablets.
- Half a tablet of GLICLAZIDE MR 60 mg or one tablet of GLICLAZIDE MR 30 mg corresponds to one tablet of GLICLAZIDE 80 mg.
- GLICLAZIDE MR can replace an antidiabetic treatment without any transitional period. If a patient is switched from a hypoglycemic sulfonylurea with a prolonged half-life

(i.e. chlorpropamide) he/she should be carefully monitored (for 1 to 2 weeks) in order to avoid hypoglycemia due to possible residual effects of the previous therapy.

Pediatrics (< 18 years of age)

Safety and effectiveness of GLICLAZIDE MR in pediatrics have not been established; therefore, Health Canada has not authorized an indication for pediatric use.

Geriatrics (≥ 65 years of age)

No significant differences in efficacy and tolerance were observed between patients over 65 years of age and younger patients, however greater sensitivity of some older individuals cannot be ruled out. Patients over 65 years of age should be started with GLICLAZIDE MR 30 mg with dosage adjustments being made cautiously.

Hepatic or Renal Impairment

Patients with renal or hepatic impairment should be started with GLICLAZIDE MR 30 mg with dosage adjustments being made cautiously (see 7 [WARNINGS AND PRECAUTIONS, Endocrine and Metabolism](#)).

Patients receiving Insulin

Maturity onset diabetics with no ketoacidosis or history of metabolic decompensation and whose insulin requirements are less than 40 units per day may be considered for GLICLAZIDE MR therapy after cessation of insulin.

If a change from insulin to GLICLAZIDE MR is contemplated in such a patient, discontinue insulin for a period of 2 or 3 days to determine whether any therapy other than dietary regulation and exercise is needed. During this insulin-free interval, test the patient's urine at least 3 times daily for glucose and ketone bodies and monitor the results carefully. The appearance of significant ketonuria accompanied by glucosuria within 12 to 24 hours after the withdrawal of insulin, strongly suggests that the patient is ketosis prone, and precludes the change from insulin to sulfonylurea therapy.

4.5 Missed Dose

If a dose is forgotten, the patient should be advised to skip the missed dose and take his or her usual dose at the regular time the next day. The dose taken on the next day should not be increased to account for the missed dose.

5 OVERDOSAGE

Symptoms

Overdosage with sulfonylureas may result in hypoglycemia but it should be noted that the dosage which causes such hypoglycemia varies widely and may be within the accepted therapeutic range in sensitive individuals.

The manifestations of hypoglycemia include sweating, flushing or pallor, numbness, chilliness, hunger, trembling, headache, dizziness, increased pulse rate, palpitations, increased blood pressure and apprehensiveness in mild cases. In more severe cases, coma appears.

However, symptoms of hypoglycemia are not necessarily as typical as those described above and sulfonylureas may cause insidious development of symptoms mimicking cerebrovascular insufficiency.

Treatment of Overdosage

Discontinue medication and treat hypoglycemia by giving dextrose promptly and in sufficient quantity.

Some sulfonylurea-induced hypoglycemias may be refractory to treatment and susceptible to relapse especially in elderly or malnourished patients. Continuous dextrose infusions for hours or days have been necessary.

Strict monitoring should be continued until the doctor is sure that the patient is out of danger.

Severe hypoglycaemic reactions, with coma, convulsions or other neurological disorders are possible and must be treated as a medical emergency, requiring immediate hospitalization.

Dialysis is of no benefit to patients due to the strong binding of gliclazide to proteins.

For management of a suspected drug overdose contact your regional Poison Control Center.

6 DOSAGE FORMS, STRENGTHS, COMPOSITION AND PACKAGING

Table 1 – Dosage Forms, Strengths, Composition and Packaging

Route of Administration	Dosage Form / Strength/Composition	Non-medicinal Ingredients
oral	30 mg modified release tablet	anhydrous colloidal silica, calcium hydrogen phosphate dihydrate, hypromellose, magnesium stearate, maltodextrin
oral	60 mg modified release tablet	anhydrous colloidal silica, hypromellose, lactose monohydrate, magnesium stearate, maltodextrin

Dosage forms:

GLICLAZIDE MR 30 mg are white oblong tablets, engraved with "30" on both faces. Each tablet contains 30 mg of gliclazide.

GLICLAZIDE MR 60 mg are white oblong breakable tablets, scored on both faces, engraved with "60" on both faces. Each tablet contains 60 mg of gliclazide.

Packaging:

GLICLAZIDE MR 30 mg is available in bottles of 100 tablets.

GLICLAZIDE MR 60 mg is supplied in bottles of 100 tablets.

7 WARNINGS AND PRECAUTIONS

General

Use of GLICLAZIDE MR must be considered as treatment in addition to proper dietary regimen and not as substitute for diet.

Careful selection of patients is important. It is imperative that there be rigid attention to diet, careful adjustment of dosage and instruction of the patient on hypoglycemic reactions, their recognition, remedies and control as well as regular, thorough medical follow-up.

Since the effects of oral hypoglycemic agents on the vascular changes and other long-term sequelae of diabetes mellitus are not fully known, patients receiving such drugs must be closely observed for both short- and long-term complications. Periodic assessment of cardiovascular, ophthalmic, renal and hepatic status is advisable.

GLICLAZIDE MR use is not recommended with medications containing alcohol, phenylbutazone (systemic route) and danazol and precautions are required when used with chlorpromazine, glucocorticoids, ritodrine, salbutamol, terbutaline and anticoagulant therapy (see [9.4 Drug-Drug Interactions](#)).

Carcinogenesis and Mutagenesis

See [16 NON-CLINICAL TOXICOLOGY](#).

Driving and Operating Machinery

Treatment with GLICLAZIDE MR can have effects on ability to drive and use machines. Patients should be made aware of the symptoms of hypoglycaemia and should be careful if driving or operating machinery, especially at the beginning of treatment (see [9.3 Drug-behavioral interactions](#)).

Endocrine and Metabolism

Dysglycaemia

Fluoroquinolones should be used with caution in patients receiving GLICLAZIDE MR. Hypoglycaemia and hyperglycaemia have been reported in diabetic patients receiving concomitant treatment with fluoroquinolones, especially in elderly patients. Careful monitoring of blood glucose is recommended in all patients taking GLICLAZIDE MR and a fluoroquinolone concomitantly.

Hypoglycemic reactions

As with other sulfonylurea drugs, manifestations of hypoglycemia including dizziness, lack of energy, drowsiness, headache and sweating have been observed and weakness, nervousness, shakiness and paresthesia have also been reported. All sulfonylurea drugs can induce severe hypoglycemia. Particularly susceptible are elderly subjects, patients with impaired hepatic or renal function, those who are debilitated or malnourished and patients with primary or secondary adrenal insufficiency. Some cases may be severe and prolonged.

Hospitalisation may be necessary and glucose administration may need to be continued for several days. Hypoglycemia may be difficult to recognize in elderly patients and in patients receiving beta-blockers.

Possible other symptoms of hypoglycaemia are intense hunger, nausea, vomiting, lassitude, sleep disorders, agitation, aggression, poor concentration, reduced awareness and slowed reactions, depression, confusion, visual and speech disorders, aphasia, tremor, paresis, sensory disorders, feeling of powerlessness, loss of self-control, delirium, convulsions, shallow respiration, bradycardia, drowsiness and loss of consciousness, possibly resulting in coma and lethal outcome. In addition, signs of adrenergic counter-regulation may be observed: clammy skin, anxiety, tachycardia, hypertension, palpitations, angina pectoris and cardiac arrhythmia.

This treatment should only be prescribed if the patient is likely to have a regular food intake (including breakfast). It is important to have a regular carbohydrate intake due to the increased risk of hypoglycemia if a meal is taken late, if an inadequate amount of food is consumed or if the food is low in carbohydrate. Hypoglycemia is more likely to occur during periods of low-calorie diet, following prolonged or strenuous exercise, following alcohol intake or during the administration of a combination of hypoglycemic agents.

Usually, hypoglycemic symptoms disappear after intake of carbohydrates (sugar). However, artificial sweeteners have no effect. Experience with other sulphonylureas shows that hypoglycemia can recur even when measures prove effective initially.

If a hypoglycemic episode is severe or prolonged, and even if it is temporarily controlled by intake of sugar, immediate medical treatment or even hospitalisation are required.

Other factors which increase the risk of hypoglycemia are: overdose of GLICLAZIDE MR, certain endocrine disorders (thyroid disorders, hypopituitarism and adrenal insufficiency) as well as withdrawal of prolonged and/or high dose corticosteroid therapy, severe vascular disease (severe coronary heart disease, severe carotid impairment, diffuse vascular disease) and concomitant administration of certain medicines (see [9.4 Drug-Drug Interactions](#)).

Patients with Porphyria

Cases of acute porphyria (which can cause severe abdominal pain, gastrointestinal symptoms, unspecified neurologic symptoms along with chronic, blistering lesions on sun-exposed skin) have been reported with the use of sulfonylurea drugs. Therefore, caution should be taken in the administration of GLICLAZIDE MR, as it may precipitate attacks of acute porphyria in patients with porphyria.

Poor Blood Glucose Control

The efficacy of gliclazide, in reducing glucose to the desired level decreases over a long period of time in many patients: this may be due to progression in the severity of the diabetes, or to a reduced response to treatment. This phenomenon is known as secondary failure and should be distinguished from primary failure, when the drug is ineffective when prescribed as first-line treatment. Adequate dose adjustment and compliance with dietary measures should be considered before classifying the patient as secondary failure. If a loss of adequate blood glucose-lowering response to GLICLAZIDE MR is detected, the drug should be discontinued.

In patients stabilized on gliclazide therapy, loss of blood sugar control may occur in cases of acute intercurrent disease such as a fever and serious infection, in stressful situations such as trauma or surgery, or if used concomitantly with herbs such as St. John's Wort (*Hypericum perforatum*) preparations or any treatment that may interact with gliclazide metabolism (see [9.6 Drug-Herb Interactions](#)). Under these conditions, discontinuation of GLICLAZIDE MR and administration of insulin should be considered.

Hematologic

Treatment of patients with glucose-6-phosphate dehydrogenase (G6PD)-deficiency with sulfonylurea agents can lead to haemolytic anaemia. Since GLICLAZIDE MR belongs to the class of sulfonylurea agents, caution should be used in patients with G6PD-deficiency and a non- sulfonylurea alternative should be considered.

Hepatic/Biliary/Pancreatic

The metabolism and excretion of sulfonylureas including GLICLAZIDE MR may be slowed in patients with impaired hepatic function. Isolated cases of impairment of liver function with cholestasis and jaundice, and hepatitis which can regress after withdrawal of the drug or may lead to life-threatening liver failure have been observed. Discontinue treatment if cholestatic jaundice appears. Therefore, GLICLAZIDE MR is contraindicated in patients with severe hepatic impairment (see [2 CONTRAINDICATIONS](#) and [7 WARNINGS AND PRECAUTIONS, Monitoring and Laboratory Tests](#)).

Monitoring and Laboratory Tests

Measurement of glycated haemoglobin levels (or fasting venous plasma glucose) is recommended in assessing blood glucose control. Blood glucose self-monitoring is also recommended.

Blood glucose control in a patient receiving GLICLAZIDE MR treatment may be affected by fever, infection, surgical intervention or when used concomitantly with St. John's Wort (*Hypericum perforatum*) preparations. Close monitoring is required in these patients. In some cases, it may be necessary to administer insulin.

Hepatic function should be assessed before initiating therapy and the liver function should be assessed periodically in patients with mild to moderately impaired hepatic function.

In patients with mild to moderately impaired renal function, renal function should be assessed periodically. Blood glucose and glycated hemoglobin levels should be regularly monitored in all patients.

Elderly patients (malnourished, with impaired hepatic, renal, or adrenal function) will require periodic monitoring and special care.

Peri-Operative Considerations

In patients stabilized on gliclazide therapy, loss of blood sugar control may occur in cases of acute intercurrent disease or in stressful situations such as trauma or surgery. Under these conditions, discontinuation of GLICLAZIDE MR (gliclazide) and administration of insulin should be considered (see [7 WARNINGS AND PRECAUTIONS, Endocrine and Metabolism, Poor blood glucose control](#)).

Renal

The metabolism and excretion of sulfonylureas including GLICLAZIDE MR may be slowed in patients with impaired renal function. If hypoglycemia should occur in such patients, it may be prolonged and appropriate management should be instituted. Therefore, GLICLAZIDE MR is contraindicated in patients with severe renal impairment (see [2 CONTRAINDICATIONS and 7 WARNINGS AND PRECAUTIONS - Monitoring and Laboratory Tests](#)).

Sensitivity/Resistance

Due to the presence of lactose in GLICLAZIDE MR 60 mg, patients with hereditary problems of galactose intolerance, glucose-galactose malabsorption or the Lapp lactose deficiency should not take GLICLAZIDE MR 60 mg. GLICLAZIDE MR 30 mg does not contain lactose.

Skin

Serious skin and hypersensitivity reactions including rash, pruritus, urticaria, angioedema, erythema, maculopapular rashes, bullous reactions (such as Stevens-Johnson syndrome, toxic epidermal necrolysis and bullous pemphigoid) and drug rash with eosinophilia and systemic symptoms (DRESS) have been reported.

Bullous pemphigoid:

Cases of bullous pemphigoid requiring hospitalisation have been reported with the use of gliclazide (see [8.5 Post-Market Adverse Reactions](#)). In reported cases, patients typically recovered with topical or systemic immunosuppressive treatment and discontinuation of the gliclazide. Tell patients to report development of blisters or erosions while receiving GLICLAZIDE MR. If bullous pemphigoid is suspected, GLICLAZIDE MR should be discontinued and referral to a dermatologist should be considered for diagnosis and appropriate treatment.

7.1 Special Populations

7.1.1 Pregnant Women

Gliclazide is contraindicated in pregnancy. It is recommended that insulin be used during pregnancy in diabetic women (see [2 CONTRAINDICATIONS](#)).

Uncontrolled diabetes (gestational or not) is associated with a higher incidence of congenital abnormalities and perinatal mortality. Blood glucose control should be optimal around the time of conception to reduce the risk of congenital malformations.

7.1.2 Breast-feeding

GLICLAZIDE MR (gliclazide) is contraindicated in breast-feeding mothers. Some sulfonylurea drugs are excreted in human milk although it is not known whether gliclazide is one of them. Because the potential for hypoglycemia in nursing infants may exist, the product is contraindicated in breast-feeding mothers (see [2 CONTRAINDICATIONS](#)).

7.1.3 Pediatrics (< 18 years of age)

Safety and effectiveness of GLICLAZIDE MR in pediatrics (<18 years of age) have not been established; therefore, Health Canada has not authorized an indication for pediatric use.

7.1.4 Geriatrics (≥ 65 years of age)

Efficacy and tolerance of GLICLAZIDE MR, prescribed using the same therapeutic regimen in subjects over 65 years, has been confirmed in clinical trials, however greater sensitivity of some older individuals cannot be ruled out.

Severe hypoglycemia can be induced by all sulfonylurea drugs. Elderly subjects are particularly susceptible.

8 ADVERSE REACTIONS

8.1 Adverse Reaction Overview

GLICLAZIDE MR 30 mg (gliclazide) has been evaluated for safety in controlled clinical trials in 955 patients, of which 728 were treated in long-term studies for up to 10 months, in comparison with gliclazide 80 mg tablets.

The most frequent adverse drug reactions are hypoglycaemia and gastrointestinal disturbances (including abdominal pain, nausea, vomiting, dyspepsia, diarrhea, constipation).

Serious adverse drug reactions that resulted in hospitalization during clinical trials were malaise, acute renal failure, and thrombophlebitis.

8.2 Clinical Trial Adverse Reactions

Clinical trials are conducted under very specific conditions. The adverse reaction rates observed in the clinical trials; therefore, may not reflect the rates observed in practice and should not be compared to the rates in the clinical trials of another drug. Adverse reaction information from clinical trials may be useful in identifying and approximating rates of adverse drug reactions in real-world use.

Hypoglycemia (see [8 WARNINGS AND PRECAUTIONS, Endocrine and Metabolism](#))

Severe hypoglycemia which mimics acute CNS disorders may occur. Hepatic and/or renal impairment, malnutrition, debility, advanced age, alcoholism, adrenal or pituitary insufficiency may be predisposing factors.

In long-term studies, the percentage of patients experiencing hypoglycemic episodes was similar between patients treated with GLICLAZIDE MR 30 mg (11.6%) and those treated with

gliclazide 80 mg tablets (11.1%). However, the number of hypoglycemic episodes for 100 patient-months was lower in the GLICLAZIDE MR 30 mg group (3.5) than in the gliclazide 80 mg tablets group (4.8).

Analysis in elderly patients (over 65 years old) showed that this population experienced, overall, less hypoglycemia than the whole population with a prevalence of hypoglycemic episodes lower in the GLICLAZIDE MR 30 mg group (2.6 hypoglycemic episodes for 100 patient-months) than in the gliclazide 80 mg tablets group (4.1).

Other adverse events

Adverse events reported during controlled clinical trials with GLICLAZIDE MR 30 mg were those expected in the population of interest, a population whose underlying disease is recognized atheromatous risk factor.

Adverse events that have been reported in at least 1.0% of diabetic patients in long-term controlled studies, whatever their relationship to treatment, are listed by body system in Table 2. The most frequent adverse events were unspecific of the disease as respiratory infections or back pain.

Table 2 – Adverse events reported in ≥1% of type 2 diabetic patients in long-term controlled studies with GLICLAZIDE MR 30 mg vs. Gliclazide 80 mg

	GLICLAZIDE MR 30 mg (n= 728) %	Gliclazide 80 mg tablets (n=734) %
Cardiovascular disorders		
Hypertension	3.2	3.7
Angina Pectoris	2.1	2.2
Oedema legs	1.2	1.4
Gastrointestinal disorders		
Diarrhea	2.5	2.0
Constipation	1.6	1.2
Gastritis	1.2	0.5
Gastroenteritis	1.1	1.5
Nausea	1.1	0.7
Abdominal pain	1.1	1.4
General disorders		
Asthenia	2.2	2.6
Headache	3.8	4.6
Inflicted injury	4.3	4.5
Infections and infestations		
Infection viral	7.7	5.6
Otitis media	1.1	0.8

Metabolism and nutrition disorders		
Hypoglycemia	11.6	11.1
Hyperglycemia	1.9	2.2
Lipid metabolism disorder	1.4	0.5
Hyperlipaemia	1.0	0.8
Musculoskeletal and connective tissue disorders		
Back pain	5.2	4.1
Arthralgia	3.0	3.5
Arthrosis	2.2	2.2
Arthritis	1.4	2.3
Tendinitis	1.1	1.0
Myalgia	2.3	1.5
Nervous System Disorders		
Dizziness	2.2	2.3
Neuralgia	1.2	0.7
Ophthalmologic disorders		
Conjunctivitis	1.0	0.8
Psychiatric disorders		
Depression	1.9	1.2
Insomnia	1.1	2.0
Renal and urinary disorders		
Urinary tract infections	2.6	3.0
Respiratory, thoracic and mediastinal disorders		
Rhinitis	4.4	4.6
Bronchitis	4.4	4.6
Pharyngitis	4.3	3.5
Upper respiratory infection	3.3	3.7
Coughing	2.1	2.0
Pneumonia	1.5	1.4
Sinusitis	1.5	1.1
Skin and subcutaneous tissue disorders		
Dermatitis	1.6	1.2
Rash	1.0	1.2
Skin disorder	1.9	2.0
Pruritus	1.0	0.4

Analysis of adverse events in sub-populations led to similar pattern as in the whole population and showed that sex, age and renal insufficiency had no significant influence on the safety profile of 30 mg.

8.3 Less Common Clinical Trial Adverse Reactions

Adverse events other than those already specifically mentioned in this product monograph and that have been reported with GLICLAZIDE MR 30 mg during long-term studies in more than one patient and/or that have been previously reported with gliclazide 80 mg tablets or with other sulfonylurea drugs include the following (drug relationship has not been proved for all cases):

Cardiovascular disorders: arteritis, cardiac failure, cerebrovascular disorder, coronary artery disorder, epistaxis, hypotension, myocardial infarction, palpitation, tachycardia, thrombophlebitis, vein disorder.

Ear/Nose/Throat disorders: hearing decreased, tinnitus.

Endocrine disorders: hypothyroidism. A decrease in the uptake of radioactive iodine by the thyroid gland has been reported with other sulfonylurea drugs. This has not been shown with gliclazide 80 mg tablets during a study involving 15 patients.

Gastrointestinal disorders: abdominal pain, anal fissure, appetite increased, colitis, duodenal ulcer, epigastric fullness, faecal incontinence, flatulence, gastric irritation, gastroesophageal reflux, GI neoplasm benign, hemorrhoids, melena, dry mouth, oesophagitis, saliva increased, tooth ache, tooth disorder, vomiting. These reactions are generally dose-related and may disappear when the dose is reduced.

General disorders: allergy, carpal tunnel syndrome, chest pain, fever, infection, fungal infection, leg pain, malaise, pain, weight increase.

Hepatobiliary Disorders: increased liver enzymes, hepatitis, hepatomegaly.

Metabolic and nutritional disorders: gout, glycosuria, hypercholesterolemia, hypertriglyceridemia, thirst. Cases of hepatic porphyria and disulfiram-like reactions have been described with sulfonylurea drugs. Clinical experience to date has shown that gliclazide 80 mg tablets has a low incidence of disulfiram type reactions.

Musculo-skeletal and connective tissue disorders: arthropathy, bursitis, hernia congenital, skeletal pain, spine malformation.

Nervous System disorders: anxiety, confusion, depression, insomnia, nervousness, neuropathy.

Ophthalmologic disorders: cataract, conjunctival haemorrhage, diplopia, glaucoma, abnormal lacrimation, retinal disorder, abnormal vision, vitreous disorder, xerophthalmia.

Renal and urinary disorders: albuminuria, cystitis, nocturia, polyuria, renal calculus, renal cyst.

Reproductive system and breast disorders: balanoposthitis, benign female breast neoplasm, impotence, mastitis, menstrual disorder, prostatic disorder, vaginitis.

Respiratory: asthma, dyspnea, tracheitis.

Skin and subcutaneous tissue disorders: fungal dermatitis, eczema, erythema, hyperkeratosis, maculopapular or morbiliform rash, nail disorder, onychomycosis, pruritus, dry skin, skin ulceration, urticaria.

These reactions may persist during treatment, which must be then interrupted. Cases of porphyria tarda and of photosensitivity have also been described with sulfonylurea drugs.

8.4 Abnormal Laboratory Findings: Hematologic, Clinical Chemistry and Other Quantitative Data

Clinical Trial Findings

The pattern of laboratory tests abnormalities previously observed with gliclazide 80 mg tablets was similar to that for other sulfonylureas. Occasional mild to moderate elevations of hepatic enzymes, LDH and creatinine and decrease in natremia have been observed. These abnormalities frequently encountered with treated or untreated diabetic patients are rarely associated with clinical symptoms and generally not considered to be drug related. As with all hypoglycemic sulfonylurea drugs, a few rare cases of leukopenia, agranulocytosis, thrombocytopenia and anemia have been reported with gliclazide 80 mg tablets. No laboratory test abnormalities other than those already reported with gliclazide 80 mg tablets have been observed during controlled clinical trials performed on GLICLAZIDE MR 30 mg.

8.5 Post-Market Adverse Reactions

In post-marketing experience with GLICLAZIDE MR, gastrointestinal disturbance, including abdominal pain, nausea, vomiting, dyspepsia, diarrhea and constipation have been reported. Skin and subcutaneous tissue disorders, rash, pruritus, urticaria, angioedema, erythema,

maculopapular rashes, bullous reactions (such as Stevens-Johnson syndrome, toxic epidermal necrolysis and bullous pemphigoid) and drug rash with eosinophilia and systemic symptoms (DRESS) have been reported.

The most serious adverse drug reactions reported with gliclazide are hypoglycaemic coma, pancytopenia, thrombocytopenia, hepatitis, cholestatic jaundice, pyrexia, pancreatitis acute and skin reactions (pruritus and rash).

The following adverse events have also been observed with gliclazide: cases of erythrocytopenia, agranulocytosis, hemolytic anemia, allergic vasculitis, hyponatremia, and elevated liver enzyme levels (AST, ALT, alkaline phosphatase); isolated cases of impairment of liver function with cholestasis and jaundice which can regress after withdrawal of the drug or may lead to life-threatening liver failure. Discontinue treatment if cholestatic jaundice appears.

9 DRUG INTERACTIONS

9.1 Serious Drug Interactions

Serious Drug Interactions

The concomitant use of miconazole and gliclazide is contraindicated (see 2 [CONTRAINDICATIONS, 9.4 Drug-Drug Interactions](#)).

9.2 Drug Interactions Overview

As a result of drug interaction, hypoglycemia may be potentiated when a sulfonylurea is used concurrently with agents such as: long-acting sulfonamides, tuberculostatics, NSAIDs, fibrates, monoamine oxidase inhibitors, salicylates, probenecid, beta-blockers, azole antifungal agents (oral and parenteral preparations), H₂ receptor antagonists, angiotensin converting enzyme inhibitors and clarithromycin. In addition, while not approved for use with other antidiabetic agents, hypoglycaemia is potentiated when gliclazide is used in combination with other antidiabetic agents.

Certain drugs tend to induce hyperglycemia and may lead to loss of blood sugar control. These include diuretics (thiazides, furosemide), corticosteroids, oral contraceptives (estrogen plus progestogen), chlorpromazine, ritodrine, salbutamol, terbutaline and nicotinic acid in pharmacologic doses.

Barbiturates should be used with caution in patients receiving an oral hypoglycemic agent since they may reduce the hypoglycemic effect.

Sulfonylureas may potentiate the action of anticoagulants. Adjustment of the anticoagulant dose may be necessary.

9.3 Drug-Behavioural Interactions

This treatment should only be prescribed if the patient is likely to have a regular food intake (including breakfast). It is important to have a regular carbohydrate intake due to the increased risk of hypoglycemia if a meal is taken late, if an inadequate amount of food is consumed or if the food is low in carbohydrate. Hypoglycemia is more likely to occur during periods of low-calorie diet and following prolonged or strenuous exercise.

Intolerance to alcohol (disulfiram-like reaction: flushing, sensation of warmth, giddiness, nausea and occasionally tachycardia) may occur in patients treated with sulfonylurea. Alcohol increases the hypoglycaemic reaction (by inhibiting compensatory reactions) that can lead to the onset of hypoglycaemic coma. Avoid alcohol or medicines containing alcohol.

Treatment with GLICLAZIDE MR can have effects on ability to drive and use machines. Patients should be made aware of the symptoms of hypoglycaemia and should be careful if driving or operating machinery, especially at the beginning of treatment.

9.4 Drug-Drug Interactions

Table 3 – Established or Potential Drug-Drug Interactions

Gliclazide	Reference	Effect	Clinical Comment
Miconazole (systemic route, oromucosal gel)	C	Increases the risk of hypoglycaemia	<u>Contra-indicated combination.</u> Increases the hypoglycaemic effect with possible onset of hypoglycaemic symptoms, or even coma.
Phenylbutazone (systemic route)	C	Increases the risk of hypoglycaemia	<u>Combination is not recommended.</u> Increases the hypoglycaemic effect of sulphonylureas (displaces their binding to plasma proteins and/or reduces their elimination). It is preferable to use a different anti-inflammatory agent, or else to warn the patient and emphasise the importance of self-

Table 3 – Established or Potential Drug-Drug Interactions

			monitoring. Where necessary, adjust the dose during and after treatment with the anti-inflammatory agent.
Other antidiabetic agents (insulins, acarbose, biguanides)	C	Increases the risk of hypoglycaemia	<u>Combinations requiring precautions for use.</u> Potentiation of the blood glucose lowering effect and thus, in some instances, hypoglycaemia may occur.
Beta-blockers	C	Increases the risk of hypoglycaemia	<u>Combinations requiring precautions for use.</u> Potentiation of the blood glucose lowering effect and thus, in some instances, hypoglycaemia may occur.
Fluconazole	C	Increases the risk of hypoglycaemia	<u>Combinations requiring precautions for use.</u> Potentiation of the blood glucose lowering effect and thus, in some instances, hypoglycaemia may occur.
Angiotensin converting enzyme inhibitors	C	Increases the risk of hypoglycaemia	<u>Combinations requiring precautions for use.</u> Potentiation of the blood glucose lowering effect and thus, in some instances, hypoglycaemia may occur.
H2-receptor antagonists	C	Increases the risk of hypoglycaemia	<u>Combinations requiring precautions for use.</u> Potentiation of the blood glucose lowering effect and thus, in some instances, hypoglycaemia may occur.
MAOIs	C	Increases the risk of hypoglycaemia	<u>Combinations requiring precautions for use.</u> Potentiation of the blood glucose lowering effect and thus, in some instances, hypoglycaemia may occur.
Sulfonamides	C	Increases the risk of hypoglycaemia	<u>Combinations requiring precautions for use.</u> Potentiation of the blood glucose lowering effect and thus, in some instances, hypoglycaemia may occur.

Nonsteroidal anti-inflammatory agents	C	Increases the risk of hypoglycaemia	<u>Combinations requiring precautions for use.</u> Potentiation of the blood glucose lowering effect and thus, in some instances, hypoglycaemia may occur.
Clarithromycin	T	Increases the risk of hypoglycaemia	<u>Combinations requiring precautions for use.</u> May potentiate the hypoglycemic action of sulfonylurea agents.
Danazol	C	Causes an increase in blood glucose levels	<u>Combination is not recommended because of diabetogenic effect of danazol.</u> If the use of this active substance cannot be avoided, warn the patient and emphasise the importance of urine and blood glucose monitoring. It may be necessary to adjust the dose of the antidiabetic agent during and after treatment with danazol.
Chlorpromazine (neuroleptic agent)	C	Causes an increase in blood glucose levels	<u>Combination requiring precautions during use.</u> High doses (>100 mg per day of chlorpromazine) increase blood glucose levels (reduced insulin release). Warn the patient and emphasise the importance of blood glucose monitoring. It may be necessary to adjust the dose of the antidiabetic active substance during and after treatment with the neuroleptic agent.
Glucocorticoids (systemic and local route: intra- articular, cutaneous and rectal preparations) and tetracosactide	C	Causes an increase in blood glucose levels	<u>Combination requiring precautions during use.</u> Increase in blood glucose levels with possible ketosis (reduced tolerance to carbohydrates due to glucocorticoids). Warn the patient and emphasise the importance of blood glucose monitoring, particularly at the start of treatment. It may be necessary to adjust the dose of the antidiabetic

			active substance during and after treatment with glucocorticoids.
Ritodrine, salbutamol, terbutaline (I.V.)	C	Causes an increase in blood glucose levels	<u>Combination requiring precautions during use.</u> Increased blood glucose levels due to beta-2 agonist effects. Emphasise the importance of monitoring blood glucose levels. If necessary, switch to insulin.
Anticoagulant therapy (Warfarin and other)	C	Potential of anticoagulation	<u>Combination which must be taken into account.</u> Sulfonylureas may lead to potentiation of anticoagulation during concurrent treatment. Adjustment of the anticoagulant may be necessary.
Drugs containing alcohol	C	Increases the risk of hypoglycaemia	Intolerance to alcohol (disulfiram-like reaction: flushing, sensation of warmth, giddiness, nausea and occasionally tachycardia) may occur in patients treated with sulfonylurea.
Fluoroquinolones	T	Increases the risk of dysglycaemia	<u>Combinations requiring precautions for use.</u> Warn the patient and emphasise the importance of monitoring blood glucose

Legend: C = Case Study; CT = Clinical Trial; T = Theoretical

9.5 Drug-Food Interactions

Interactions with food have not been established.

9.6 Drug-Herb Interactions

St. John's Wort

Pharmacodynamic interactions between gliclazide and the herbal remedy St. John's Wort may occur and may lead to hyperglycemia or loss of blood glucose control.

9.7 Drug-Laboratory Test Interactions

Interactions with laboratory tests have not been established.

10 CLINICAL PHARMACOLOGY

10.1 Mechanism of Action

GLICLAZIDE MR (gliclazide) is a hypoglycemic agent of the sulfonylurea group. The hypoglycemic action of GLICLAZIDE MR is related to an improvement in insulin secretion from the functioning beta cells of the pancreas. It potentiates the insulin release, improves the dynamics of insulin. Increase in postprandial insulin and C-peptide secretion persists after two years of treatment. Gliclazide has extra-pancreatic actions. These metabolic actions are accompanied by hemovascular effects. However, the mechanism of action regarding these effects is still poorly understood. The clinical significance of these effects has not been established.

Effects on insulin release. In type 2 diabetics, gliclazide restores the first peak of insulin secretion in response to glucose and increases the second phase of insulin secretion. A significant increase in insulin response is seen in response to stimulation induced by a meal or glucose.

Extra-pancreatic effects. It has been demonstrated that gliclazide increases peripheral insulin sensitivity:

- In muscle: the action of insulin on glucose uptake, measured during an euglycemic hyperinsulinemic clamp is significantly increased (+35%), due to an improvement in peripheral sensitivity to insulin. This leads to an improvement in diabetes control. Gliclazide acts mainly by potentiating insulin action on muscle glycogen synthetase. Moreover, results of studies on the muscle are consistent with a post-transcriptional action of gliclazide on GLUT4 glucose carriers;
- In the liver: studies on glucose turnover show that gliclazide decreases hepatic glucose production, leading to an improvement in fasting blood glucose levels.

Hemovascular effects. Gliclazide decreases microthrombosis by two mechanisms which may be involved in complications of diabetes:

- A partial inhibition of platelet aggregation and adhesion, with a decrease in the markers of platelet activation (beta thromboglobulin, thromboxane B2);
- A restoration of the vascular endothelium fibrinolytic activity with an increase in t-PA activity.

Antioxidant effects. A controlled clinical study in diabetics has confirmed the antioxidant effects of gliclazide that were already demonstrated in clinical pharmacology: reduction in plasma levels of lipid peroxides, increase in the activity of erythrocyte superoxide dismutase.

10.2 Pharmacodynamics

Hypoglycemic activity

The main mechanism of action of gliclazide consists of an increase in the insulin secretory potential of pancreatic beta-cells in a situation of hyperglycemia. This effect of gliclazide on insulin secretion is maintained during long-term treatment in type 2 diabetic patients. It was observed that the administration of gliclazide was followed by:

- a consistent and significant decrease in fasting blood glucose;
- a more than 1% decrease in mean glycated hemoglobin;
- an inhibition by 12 to 27% of the rise in blood glucose after a standard meal or an oral glucose load.

A slight and transitory increase in mean fasting plasma insulin levels was occasionally observed with gliclazide treatment.

Regarding the biphasic nature of insulin secretion, the first peak that is severely blunted in type 2 diabetes is improved during gliclazide treatment.

In addition to the effect of gliclazide on the secretion of insulin, extrapancreatic effects have also been evidenced. Gliclazide improves peripheral sensitivity to insulin and increases glucose utilization rate:

- with euglycemic hyperinsulinemic clamps in obese and non-obese type 2 diabetic patients, it has been shown that gliclazide, after 3 months of treatment, increases the disappearance rate and metabolic clearance of glucose at the highest insulin infusion rates (100 and 300 mU/kg/h);
- in comparison to diet treatment, gliclazide also enhances insulin-stimulated glucose metabolism after 8 weeks of treatment by potentiating insulin action on skeletal muscle glycogen synthetase.

Studies on glucose turnover have also shown that basal hepatic glucose production, measured by tracer methodology, was markedly reduced (28-50%) after 3 months of treatment.

Hemovascular activity

Gliclazide possesses anti-platelet properties which are independent of its antidiabetic action, and improves the fibrinolytic potential in diabetic patients:

-numerous studies have shown inhibitory effects of gliclazide on platelet aggregation and hyperadhesiveness. A statistically significant 22% decrease in collagen-induced platelet aggregation has been observed after 3 and 6 months of treatment with gliclazide in 15 patients previously well controlled under glibenclamide. A concentration-dependent inhibition of PAF-induced platelet aggregation has also been reported with gliclazide in vitro in platelet-rich plasma from healthy subjects and type 2 diabetic patients. Finally, a consistent decrease in markers of platelet activation (e.g. beta thromboglobuline and thromboxane B2 levels) has been observed with gliclazide whether glycemic control improved or not;

- change to gliclazide of patients treated since several years by chlorpropamide is followed by normalisation of the t-PA activity, sustained over 24-48 months. This has been confirmed by 2 studies in type 1 and glibenclamide-treated type 2 diabetics: in both, the addition of gliclazide to insulin or the switch to this sulfonylurea were followed by significant increase in t-PA and in the activity of the intrinsic fibrinolytic system.

Antioxidant activity

Gliclazide is a strong free radical scavenging agent, an effect demonstrated both in vitro and in patient. In 17 type 2 diabetic patients switched to gliclazide and seen at regular intervals during a 36-week period, peroxidized lipids and oxidized damaged IgG decreased significantly. These effects of gliclazide on the oxidative stress have been confirmed in a double-blind study in diabetic patients. Highly significant and sustained decrease in peroxidized lipid levels and increase in erythrocyte superoxide dismutase activity were obtained with gliclazide, but not with glibenclamide.

10.3 Pharmacokinetics

Absorption

Gliclazide is slowly and completely absorbed from the gastro-intestinal tract (mean absolute bioavailability of 97%). After administration, plasma concentrations rise gradually and the maximum concentration is usually reached after about 6 hours, with a plateau maintained for another 4 to 6 hours. Intra- individual variability is low. Food intake does not affect the rate and extent of absorption. The relationship between the dose administered and the area under the concentration curve as a function of time is linear. Linear kinetics were observed with GLICLAZIDE MR 30 mg in the dose range up to 120 mg.

Distribution

The volume of distribution is relatively small, which can partially be explained by high protein binding (about 95%). A single daily dose of GLICLAZIDE MR 30 mg maintains effective gliclazide plasma concentrations over 24 hours.

Metabolism

Although more than 90% of unchanged gliclazide is found in plasma following oral administration, this is extensively metabolized with little of the unchanged compound (<1%) found in urine. Six principal metabolites have been identified in urine, essentially oxidized and hydroxylated derivatives, and two glucuronoconjugates. No active metabolites have been detected in plasma.

Elimination

Gliclazide metabolites and conjugates are primarily (60-70%) eliminated via the urine, with about 10 to 20% elimination via feces.

The mean elimination half-life is 16 h (range 12-20 h).

Special Populations and Conditions

- **Pediatrics (< 18 years of age)**
Safety and effectiveness of GLICLAZIDE MR in pediatrics have not been established; therefore, Health Canada has not authorized an indication for pediatric use.
- **Geriatrics (≥ 65 years of age)** No clinically significant modifications in the pharmacokinetic parameters have been observed in elderly patients.
- **Sex** No significant relationship was found between any of the pharmacokinetic parameters and the covariates gender, body weight and creatinine clearance.

11 STORAGE, STABILITY AND DISPOSAL

Store at room temperature (15°C-30°C).

Keep out of reach of children and pets.

Unused medication should not be disposed of down the drain or in household garbage.

12 SPECIAL HANDLING INSTRUCTIONS

No special requirements.

PART II: SCIENTIFIC INFORMATION

13 PHARMACEUTICAL INFORMATION


Drug Substance

Proper name: gliclazide

Chemical name: 1-(hexahydrocyclopenta[c]pyrrol-2(1H)-yl)-3-[(4-methylphenyl)sulphonyl]urea (European Pharmacopoeia)

Molecular formula and molecular mass: C₁₅H₂₁N₃O₃S, 323.4

Structural formula:



Physicochemical properties:

Appearance: white or almost white powder.
Solubility: practically insoluble in water
freely soluble in methylene chloride
sparingly soluble in acetone
slightly soluble in ethanol 96%.

Acid function pKa: 5.6

Overall distribution coefficient of gliclazide between water and octanol at pH 7.4 (Log D pH 7.4) is 0.4

Melting point: Approximately 168°C

14 CLINICAL TRIALS

14.1 Clinical Trials

Diabetes mellitus

Two pivotal controlled clinical studies involving a total of 888 type 2 diabetic patients have been conducted during the initial development of the modified-release (MR) formulation of gliclazide.

The first study was a phase II, multicenter, comparative, randomized, double-blind trial designed to evaluate the dose/efficacy relationship of the MR formulation administered once daily and to determine its minimum effective dose. Placebo and five gliclazide MR doses (15, 30, 60, 90 and 135 mg) were assessed over 8 weeks in 224 patients (35 to 39 patients per group). The lowest tested dose (15 mg once daily) slightly decreased fasting plasma glucose (FPG) but the effect of this dose on glycated hemoglobin (HbA1c) was not clinically significant. The first gliclazide MR dose demonstrating clinically relevant efficacy on both parameters was 30 mg once daily. For doses above 30 mg, the efficacy of the gliclazide MR formulation was confirmed with a good clinical and biological acceptability. This study thus demonstrated that 30 mg of gliclazide MR administered once daily is the minimum effective dose for initiating treatment in type 2 diabetic patients.

The second study was a large phase III, multinational, comparative, randomized, double-blind trial aimed at demonstrating the therapeutic equivalence of gliclazide MR 30 mg compared to the gliclazide 80 mg immediate release formulation. A total of 664 patients were randomized in two parallel groups, one assigned to gliclazide 80 mg (336 patients) and one to gliclazide MR 30 mg (328 patients). After a 4-month dose escalating period allowing patient-tailored titration, patients entered a maintenance period of 6 months. Gliclazide 80 mg was administered at 80, 160, 240 or 320 mg/day, with doses above 80 mg given twice daily; gliclazide MR 30 mg was always administered once daily at breakfast time at 30, 60, 90 or 120 mg/day. The study demonstrated that after 10 months of treatment, gliclazide MR 30 mg is at least as effective as gliclazide 80 mg in controlling HbA1c and FPG levels of type 2 diabetic patients. The therapeutic equivalence was actually achieved with lower daily doses of the MR formulation, 30 mg of gliclazide MR producing a similar effect as 80 mg of gliclazide immediate release formulation. The general safety of both formulations was good with no difference in type and incidence of adverse events. With regard to hypoglycemia, the number of patients experiencing hypoglycemic episodes was almost the same in both groups. However, the number of hypoglycemic episodes was lower in the gliclazide MR group than in the gliclazide 80 mg group.

14.3 Comparative Bioavailability Studies

Two pivotal comparative bioavailability studies demonstrate comparable bioavailability between GLICLAZIDE MR 60 and GLICLAZIDE MR 30 mg tablets at equivalent doses. The therapeutic equivalence between GLICLAZIDE MR 30 mg and GLICLAZIDE 80 mg, the immediate release formulation has also been demonstrated. The efficacy and safety data of GLICLAZIDE MR 60 mg thus relies on GLICLAZIDE MR 30 mg and GLICLAZIDE 80 mg data.

Two randomized crossover comparative bioavailability studies were conducted with gliclazide MR 60 mg tablet and gliclazide MR 30 mg tablet administered as a single oral 60 mg dose (1 X 60 mg or 2 X 30 mg), to healthy adult male volunteers under non-standard fasting conditions or following the administration of a high-fat meal. Data from twenty-three (high-fat meal) and twenty-four (fasting conditions) subjects who completed each study is presented in the tables below.

**Table 4 - SUMMARY TABLE OF THE COMPARATIVE BIOAVAILABILITY DATA
(FASTING CONDITIONS)
Gliclazide
(1 x 60 mg or 2 x 30 mg)
From measured data Geometric Mean
Arithmetic Mean (CV%)**

Parameter (unit)	Gliclazide MR 60 mg*	Gliclazide MR 30 mg&	% Ratio of Geometric Least Square Means	90% Confidence Interval
AUC _T (ng.h/ml)	35762 36904 (27%)	36316 38336 (37%)	98%	93% - 105%
AUC _I (ng.h/ml) [£]	37048 38629 (32%)	36406 38077 (33%)	102%	98% - 106%
C _{max} (ng/ml)	1367 1394 (20%)	1571 1616 (24%)	87%	81% - 94%
T _{max} (h) #	8.5 (3.0 - 24)	5.0 (4.0 - 16)		
T _{1/2} (h) ###	14.5 (35%)	14.6 (30%)		

*: Test, administered as 1 x 60 mg
 &: Reference, administered as 2 x 30 mg
 £: n = 23
 #: expressed as median (range)
 ###: expressed as arithmetic mean (CV%)

**Table 5 - SUMMARY TABLE OF THE COMPARATIVE BIOAVAILABILITY DATA
(FED CONDITIONS)**

Gliclazide

(1 x 60 mg or 2 x 30 mg)

From measured data Geometric Mean
Arithmetic Mean (CV%)

Parameter (unit)	Gliclazide MR 60 mg*	Gliclazide MR 30 mg&	% Ratio of Geometric Least Square Means	90% Confidence Interval
AUC _T (ng.h/ml)	39019 39865 (21%)	38206 39273 (23%)	100%	97% - 103%
AUC _I (ng.h/ml)	40596 41634 (22%)	39588 40849 (25%)	100%	97% - 103%
C _{max} (ng/ml)	2101 2117 (12%)	2141 2178 (19%)	96%	91% - 102%
T _{max} (h) #	7.0 (5.0 - 10)	7.0 (4.0 - 16)		
T _{1/2} (h) ###	14.3 (20%)	13.9 (19%)		

*: Test, administered as 1 x 60 mg

&: Reference, administered as 2 x 30 mg

#: expressed as median (range)

###: expressed as arithmetic mean (CV%)

Bioequivalence was demonstrated between one 60 mg gliclazide modified release tablet and two ½ 60 mg gliclazide modified release tablets following a single oral administration to healthy adult male volunteers, indicating that division of the 60 mg gliclazide modified release tablet did not significantly affect the rate or extent of absorption of gliclazide.

15 MICROBIOLOGY

No microbiological information is required for this drug product.

16 NON-CLINICAL TOXICOLOGY

General Toxicology

Acute toxicity

Table 6: Gliclazide acute toxicity data in different species

Species	Mean weight	Number of animals per lot	LD ₅₀ (mg/kg)	
Mouse CD-SPF	25 g	10 M 10 F	> 3 000	
Mouse ICR-HAN	20 g	10 M 10 F	> 4 000	
Rat SD-SPF	250 g	10 M	3733 < 5200	2679
		10 F	3407 < 5467	2123
Rat CFY	110 g	6 M 6 F	> 4 000	
Tricolor Guinea Pig	240 g	4 M	48 hours: 1732 < 1999 1501	10 days: 1599 < 2016 1269
		4 F	2244 < 2509 1944	2068 < 2553 1675
Beagle Dog	7 kg	3 M 3 F	> 3000	

The LD 50 is greater than 3000 mg/kg in the mouse, rat and dog (i.e., 300 times the therapeutic dose) and than 2000 mg/kg in the guinea-pig (i.e. 200 times the therapeutic dose).

Symptomatology is essentially linked to the hypoglycemic effect of the drug.

Sub-chronic toxicity

- Maximum tolerated dose:

In the dog, this dose is between 150 and 200 mg/kg by daily administration.

-Four-week oral toxicity study in the Beagle dog:

Groups of 4 Beagle dogs (2 males, 2 females) were treated for 30 days with 0, 15, 30, 45 or 90 mg/kg/day. At the dose of 90 mg/kg, 2 animals died as a result of prolonged hypoglycemic coma following 2 weeks of treatment. All others showed normal behavior, with the exception of an increase in the weight of the liver. No evidence was found of any change in biochemical (apart from the fall in blood glucose), hematological and histopathological parameters.

-Two-month oral toxicity study in the guinea-pig:

Groups of 10 guinea-pigs (5 males, 5 females), were treated 6 days out of 7 for 2 months with 0, 25, 50 or 100 mg/kg/day. Only male animals in the 50 mg/kg group showed delayed weight gain. All others had normal biochemical, hematological and histopathological results.

Chronic toxicity

-Six-month study in the Sprague-Dawley rat:

Groups of 20 rats (10 males, 10 females) weighing 300 g, were treated for 6 days out of 7 for 6 months with 0, 25, 100 or 200 mg/kg/day. Seven deaths occurred as a result of technical problems. All other animals showed normal behavior and haematological results. From a biochemical standpoint, blood urea decreased significantly in the male rats as did blood glucose in the males of the 100 mg/kg/day group. Histological examination showed an increase in the weight of the liver and kidneys in male animals, not accompanied by any histological lesion.

In a six-month rat study carried out in Japan with higher doses (0, 50, 100, 200, 400 and 800 mg/kg/day) females exhibited greater systemic toxicity when compared with males, suggesting that females may be more sensitive to the product: slight increases in liver enzymes together with slight decreases in erythrocytes counts, hematocrit values and hemoglobin concentrations at doses of 200 mg/kg and higher.

-Six-month study in the Beagle dog:

Groups of 6 dogs (3 males, 3 females) were treated daily for 6 months with 0, 15 or 30 mg/kg of gliclazide or 50 mg/kg of tolbutamide.

From a clinical standpoint:

- 3 deaths (one at 15 mg/kg, two at 30 mg/kg) in the gliclazide group as a result of hypoglycemic coma;
- 1 convulsion, 4 cases of severe gastro-intestinal disturbances in the tolbutamide group;
- weight changes and food consumption were similar with both drugs.

From a laboratory standpoint:

- 40% fall in blood glucose in animals treated with gliclazide;
- signs of hepatotoxicity in the tolbutamide group.

From a histological standpoint:

- increase in weight of the liver in the 3 deaths of the gliclazide group;
- increase in the weight of the liver and lesions of toxic hepatitis in 5 animals out of 6 of the tolbutamide group.

•

-Twelve-month oral toxicity study in the Beagle dog:

Groups of 8 dogs (4 males, 4 females) were treated for 12 months with 0, 12 or 24 mg/kg/day of gliclazide.

Four animals in each group were sacrificed after 90 days.

- there were no deaths;
- no evidence of any modification in behavior and body weight;
- significant fall in blood glucose;
- fluctuation in certain parameters (liver enzymes, lipid profile, creatinine);
- at autopsy: swelling of the renal and hepatic parenchyma and at the highest dose a slight increase in the weight of the thyroid and slight decrease in the weight of the pituitary gland.

•

-Twelve-month oral toxicity study in the rhesus monkey:

Groups of 8 rhesus monkeys (4 males, 4 females) were treated daily for 12 months with 0, 20, 60 or 180 mg/kg of gliclazide.

- no evidence was found of any modification in weight gain nor food consumption;
- significant fall in blood glucose;
- irregular rise in some liver enzymes in some animals;
- no abnormality by histopathological examination.

Carcinogenicity

Specific carcinogenicity studies have not been performed; the following safety data are now available:

- gliclazide belongs to the chemical class of the phenylsulfonyleurea which did not demonstrate any mutagenic or carcinogenic potential. Its metabolic pathway is consistent with the general metabolic pathway of the class;
- gliclazide was not associated with any mutagenic action in the numerous studies performed;
- long term toxicity studies did not reveal any evidence of carcinogenicity;
- gliclazide has been studied in several thousands of patients during clinical trials and has been marketed for numerous years all over the world and in particular in Europe and Japan without any suspicion of carcinogenicity.

Mutagenicity

Gliclazide

The mutagenic potential of gliclazide has been sought using six mutagenesis tests, i.e.:

- 2 gene mutation tests (Ames test);
- 1 *in vitro* chromosomal aberration test (human lymphocyte test);
- 2 *in vivo* chromosomal tests (micronucleus test);
- 1 unscheduled DNA synthesis test.

Gene mutation tests

Ames test

1st test

In this test, gliclazide was used in the presence of 5 strains of *Salmonella typhimurium* (TA 1535/1537/1538/98/100) at the doses of 0, 0.005, 0.01, 0.05, 0.1, 0.5, 1, 3, 5 and 8 mg/petri dish, with and without metabolic activators. Positive controls were used for each strain with and without metabolic activators.

The qualitative test showed no mutagenic effect. The quantitative test at doses of 0.005 mg to 8 mg/dish showed no significant increase in the number of revertants.

Thus no mutagenic effect was seen under the experimental conditions adopted.

2nd test

This test used 7 strains of *Salmonella typhimurium* (TA 97/98/100/102/1535/1537/1538) at the doses of 0, 0.05, 0.1, 0.5, 1, 3, 5 and 8 mg of gliclazide per petri dish, in the presence and absence of metabolic activator. Positive controls were used for each strain, with and without metabolic activators.

No mutagenic effect was seen in the qualitative test. No mutagenic activity was detected in the quantitative test under the experimental conditions described.

In vitro chromosomal aberration test

Possible clastogenic potential action of gliclazide on activated lymphocytes in culture was studied by the human lymphocyte test with and without metabolic activators. Maximum tolerated doses determined in the preliminary toxicity test were 0.033 mg/ml with metabolic activators and 0.1 mg/ml without metabolic activator.

Gliclazide was used at the following concentrations:

- 0, 0.003, 0.01 and 0.033 mg/ml with metabolic activators;
- 0, 0.01, 0.033 and 0.1 mg/ml without metabolic activator.

Cyclophosphamide (0.02 mg/ml) and bleomycin (0.250 mg/ml) were used as positive controls with and without metabolic activators. Gliclazide was not found to have any clastogenic activity under the experimental conditions described.

In vivo chromosomal aberration

Micronucleus test

1st test

The test used three groups of 10 OF1 mice: 1 negative control, 1 gliclazide high dose (2 g/kg x 2), 1 gliclazide low dose (1 g/kg x 2) and one group of 5 positive control mice given cyclophosphamide (50 mg/kg x 2). No evidence was found of any significant variation in the number of erythrocyte micronuclei. Gliclazide was not associated with any mutagenic action detectable by the micronucleus test.

2nd test

The test used SPF Swiss mice as follows:

- 24 mice for the preliminary toxicology test which determined the maximum administrable dose as 3 g/kg;
- 108 mice in the phase 1 genetic toxicology test with study of effect/time relationship at the maximum administrable dose (MAD) (sacrifice of animals at times 24, 48 and 72 hours);
- 60 mice in the phase 2 genetic toxicology test with study of the dose/effect relationship at the time defined in phase 1 (t = 24 h) and using the following doses: 0, 750 (MAD/4), 1500 (MAD/2) and 3000 mg/kg (MAD).

Cyclophosphamide 50 mg/kg was used as positive control. Gliclazide was found to be free of any clastogenic activity under the experimental conditions adopted in this trial involving oral administration in the Swiss mouse.

Unscheduled DNA synthesis

The potential of gliclazide to induce unscheduled DNA synthesis in the liver of orally dosed male Wistar rats was investigated using an *in vivo/in vitro* procedure. Doses of 0, 632.5 and 2000 mg/kg of gliclazide were administered by gavage. Two samples were planned and collected approximately 12-14 h or 2-4 h after dosing. Primary cultures of hepatocytes were prepared from 3 animals per dose. *In vitro*, the aim was to determine the net grain count. Plasma levels of gliclazide were measured 2 hours after dosing with 2000 mg/kg. Under the conditions of this study, gliclazide did not induce unscheduled DNA synthesis in rats properly exposed to the drug.

Paratoluenesulfonamide (PTS)

PTS is a gliclazide degradation impurity which may occur in the dosage form. The mutagenic potential of PTS is well documented in the literature since this compound is also a degradation product of saccharin. The following *in vitro* and *in vivo* tests support the qualification of this impurity:

In vitro tests

Ames test

Strains of *Salmonella typhimurium* (TA 1530/1535/1538/98/100) were tested for doses $\leq 4.10^{-2}$ M. No mutagenic effect was observed. The same result was reported for the strains TA 1535/1537/1538/98/100 at doses up to 18000 $\mu\text{g}/\text{plate}$, with and without metabolic activation. In a ZLM medium (with lower content of glucose and citrate) with a metabolic activator, PTS induced a slight increase over the revertant frequency in the strain TA 98 at doses $\geq 9600 \mu\text{g}/\text{plate}$.

SCE test on CHO-K1 cells

Concentrations of 0, 14, 200 and 400 $\mu\text{g}/\text{ml}$ did not show any significant difference after a 24-hour treatment in comparison with the DMSO at a concentration of 50 $\mu\text{g}/\text{ml}$.

Test on human embryo cells

The RSa cells (ouabain-resistant) were exposed to PTS concentrations $\leq 1800 \mu\text{g}/\text{ml}$. In comparison with a UV exposure, used as a positive control, no induction of mutation to ouabain-resistance was observed after a 24-hour treatment.

In vivo tests

Drosophila test

No mutagenic effect was reported with PTS administered by abdominal injection at a dose of 5 mM. In one study, an induction of recessive lethal sex-linked mutation was observed at a concentration of 2.5 mM.

Micronucleus test

No significant increase in the micronuclei rate was reported after intraperitoneal or oral administration (2 x 855 mg/kg) in male and female mice.

Reproductive and Developmental Toxicology

Teratogenicity

Teratogenicity studies have been carried out in three species: mouse, rat and rabbit.

- In the CD/SPF mouse (group of 30 females), administration of gliclazide at doses of 0, 50, 250 and 500 mg/kg/day starting from mating and throughout gestation did not modify fertilization and abortion rates and had no apparent teratogenic effect.
- In the CFY-SPF rat (groups of 20 females), administration of gliclazide at doses of 0, 50, 100 and 200 mg/kg/day from the 6th to the 15th day of gestation did not show any embryotoxic effect.
- In the SD/SPF rat (groups of 60 females), administration of gliclazide at the doses of 0, 15, 30, 60, 120, 240 and 480 mg/kg/day starting from mating and throughout gestation had no effect on fertilization, gestation, mean number of foetuses or incidence of foetal abnormalities. The number of offspring surviving at 48 hours was decreased in the 15, 60, 120 and 480 mg/kg groups. No other abnormality was seen.
- In the common rabbit (group of 15 females), administration of gliclazide at doses of 0, 10, 25 and 50 mg/kg/day from the 6th to the 18th day of gestation had no effect on the number of foetal resorptions, percentage of abortion nor the mean number of foetuses per litter.
- In the New Zealand rabbit (group of 6 females), administration of gliclazide at doses of 0, 50, 75, 100 and 200 mg/kg/day for 13 days followed by an observation period of 8 days, was associated with maternotoxicity and embryotoxicity in the form of gastro-intestinal and renal lesions accompanied by anorexia and weight loss. However, there was no evidence of any teratogenic effect.

Fertility and reproduction

In the SD rat, groups of 40 females and of 20 males were treated for 8 and 70 days respectively before mating and until weaning in the females, and until 15 days after littering in the males, with gliclazide at doses of 0, 10, 50 and 200 mg/kg/day.

There was no evidence of any change in fertilization nor abortion rates. Foetal resorption, placental haemorrhage and foetal atrophy rates were unaffected. The genital tract of treated parents showed no abnormality imputable to treatment. No embryotoxic effect was seen on foetuses of females sacrificed before littering. In females in which gestation was allowed to run

to term, a significant decrease in the viability of offspring was seen at 48 hours. No abnormality was seen during the study of fertility and reproduction in first generation offspring born of treated animals.

PATIENT MEDICATION INFORMATION

READ THIS FOR SAFE AND EFFECTIVE USE OF YOUR MEDICINE

Pr GLICLAZIDE MR

Gliclazide Modified release tablets

Gliclazide Modified release breakable tablets

Read this carefully before you start taking **GLICLAZIDE MR** and each time you get a refill. This leaflet is a summary and will not tell you everything about this drug. Talk to your healthcare professional about your medical condition and treatment and ask if there is any new information about **GLICLAZIDE MR**.

What is GLICLAZIDE MR used for?

GLICLAZIDE MR is used along with diet and exercise to lower blood sugar levels in adults with type 2 diabetes.

How does GLICLAZIDE MR work?

GLICLAZIDE MR belongs to a group of medicines called hypoglycemic (antidiabetic) drugs. They are part of a subgroup of medicines called sulfonylureas. It helps improve the release of insulin in the body.

What are the ingredients in GLICLAZIDE MR?

Medicinal ingredients: Gliclazide.

Non-medicinal ingredients:

- GLICLAZIDE MR 30 mg tablets contain: anhydrous colloidal silica, calcium hydrogen phosphate dihydrate, hypromellose, magnesium stearate, maltodextrin.
- GLICLAZIDE MR 60 mg tablets contain: anhydrous colloidal silica, hypromellose, lactose monohydrate, magnesium stearate, maltodextrin.

GLICLAZIDE MR comes in the following dosage forms:

- 30 mg modified-release tablets
- 60 mg modified-release breakable tablets

Do not use GLICLAZIDE MR if you:

- are allergic to gliclazide, other sulfonylureas, sulfonamides, or to any of the ingredients of these medicines.
- have unstable and/or insulin dependent type 1 diabetes (particularly Juvenile diabetes)
- have diabetic ketoacidosis (DKA)
- have diabetic pre-coma and coma
- have a serious infection, trauma or surgery
- have severe liver problems
- have severe kidney problems
- are taking medicine containing miconazole (used to treat fungal or yeast infections)
- are pregnant
- are breast-feeding

To help avoid side effects and ensure proper use, talk to your healthcare professional before you take GLICLAZIDE MR. Talk about any health conditions or problems you may have, including if you:

- have or have had liver problems
- have or have had kidney problems
- are pregnant or planning to get pregnant
- are breast-feeding
- have a blood disease called G6PD-deficiency anemia (this is a condition where red blood cells break down faster than they are made)
- have porphyria (a condition that results in chemicals building up in your body that results in symptoms affecting your skin, nervous system and other areas)
- are lactose intolerant (GLICLAZIDE MR 60 mg contains lactose)

Other warnings you should know about:

- **Effect on Blood Sugar Levels**
 - **Low blood sugar levels (hypoglycemia):**
 - GLICLAZIDE MR may cause low blood sugar levels (a condition called hypoglycemia). You should talk to your healthcare professional about symptoms of low blood sugar levels and what to do if you have these

symptoms. You should also test your blood sugar levels as instructed by your healthcare professional.

- See the **Serious Side Effects Table** for symptoms of low blood sugar levels (hypoglycemia).
 - **If you experience these symptoms:**
 - **Immediately eat or drink something containing sugar. Good sources of sugar are:**
 - **orange juice**
 - **corn syrup**
 - **honey**
 - **sugar cubes or table sugar (dissolved in water)**
 - **Talk to your healthcare professional as soon as you can.**
- **High blood sugar levels (hyperglycemia):**
 - Your blood sugar may get too high (a condition called hyperglycemia) if you experience a fever, infection, surgery, or trauma (stressful conditions).
 - In such cases, contact your healthcare professional as your medication may need to be adjusted.
 - Symptoms of high blood sugar levels include the following: thirst, frequent urination, dry mouth, dry itchy skin, skin infections and reduced performance.
- **Serious Skin Reactions:** (for example DRESS, Stevens-Johnson Syndrome, Toxic Epidermal Necrolysis, bullous pemphigoid, hypersensitivity Syndrome) that can be any combination of red itchy rash with blisters and peeling of the skin and/or of the lips, eyes, mouth, nasal passages or genitals.
 - You may also have a fever, chills, headache, cough, body aches or joint pain.
 - You may have less or dark urine, yellow skin or eyes.
 - If you have these symptoms, you should stop taking GLICLAZIDE MR immediately and talk to your healthcare professional.
- **Children and adolescents (under 18 years of age):**

GLICLAZIDE MR is not recommended for use in patients under 18 years of age.

- **Driving and Operating Machinery:**

Your alertness and reactions may be impaired due to low blood sugar levels, especially when starting to take GLICLAZIDE MR. This may affect your ability to drive or to operate machines.

Tell your healthcare professional about all the medicines you take, including any drugs, vitamins, minerals, natural supplements or alternative medicines.

Serious Drug Interactions

GLICLAZIDE MR (gliclazide) should not be taken if you are also taking miconazole (a medicine used to treat bacterial or yeast infections).

The following may interact with GLICLAZIDE MR:

- medicines that are used to treat diabetes such as other antidiabetic agents
- medicines used to treat bacterial infections (antibiotics) such as sulfonamides/sulfa drugs, clarithromycin and fluoroquinolones.
 - If GLICLAZIDE MR is taken at the same time as fluoroquinolones, low and high blood sugar can occur.
 - Talk to your healthcare professional about monitoring your blood sugar levels if taking these medicines together. This is especially important if you are elderly.
- medicines used to treat tuberculosis (anti-tuberculosis)
- medicines used to treat fungal infections (anti-fungal) such as fluconazole.
- medicines used to treat inflammation (non-steroidal anti-inflammatory drugs (NSAIDs) such as phenylbutazone or corticosteroids.
- salicylates (used to treat pain, fever and inflammation) such as acetylsalicylic acid.
- medicines used to treat high blood pressure and certain heart conditions like:
 - angiotensin converting enzyme (ACE) inhibitors,
 - beta blockers
 - diuretics such as thiazides and furosemide
- medicines used to thin blood (anticoagulant therapy) such as warfarin
- medicines used to treat high levels of fats in the blood such as fibrates and nicotinic acid
- H2-receptor antagonists (used to treat acid reflux/heartburn)
- monoamine oxidase inhibitors (used to treat depression)
- chlorpromazine (used to treat certain psychiatric conditions)
- probenecid (used to treat high levels of uric acid in the blood and gout)

- medicines used to treat asthma such as salbutamol and terbutaline
- ritodrine (used to stop premature labor)
- medicines used to treat seizures (barbiturates) such as sedatives and anti-seizure medications.
- oral contraceptives used for birth control such as estrogen plus progestogen.
- danazol (used to treat breast cysts and endometriosis, a condition where tissue similar to the lining of the uterus grows outside the uterus)
- St. John's wort, a herbal product used to treat depression
 - Using St. John's Wort with GLICLAZIDE MR may lead to high blood sugar levels and loss of blood sugar control.
- alcohol
 - Avoid drinking alcohol and taking medicines containing alcohol while you are taking GLICLAZIDE MR. This can lead to a drop in blood sugar levels.

Medicines like diuretics, corticosteroids, oral contraceptives, chlorpromazine, ritodrine, salbutamol, terbutaline, danazol and nicotinic acid may lead to high blood sugar levels. See **Other Warnings** for more information on high blood sugar levels (hyperglycemia).

How to take GLICLAZIDE MR:

- Take GLICLAZIDE MR exactly as your healthcare professional has told you. Check with your healthcare professional if you are not sure.
- Take GLICLAZIDE MR once daily at breakfast.
- The 30 mg tablets cannot be split in half. Swallow the tablet whole with water.
- The 60 mg tablets can be halved.
- Do not crush or chew the tablets of either strength.

Usual dose:

30 mg per day. This is one tablet of GLICLAZIDE MR 30 mg or half tablet of GLICLAZIDE MR 60 mg.

The daily dose should not exceed 120 mg.

Talk to your healthcare professional about checking your blood sugar levels. You should test your blood sugar levels as instructed by your healthcare professional to make sure that your blood sugar levels are being controlled.

Your healthcare professional will check your blood sugar levels during regular visits, especially when you start taking this medicine.

Overdose:

If you think you, or a person you are caring for, have taken too much GLICLAZIDE MR, contact a healthcare professional, hospital emergency department, or regional poison control centre immediately, even if there are no symptoms.

Missed Dose:

If you miss a dose of this medicine, skip the missed dose. Take your next dose at the regular time the next day. Do not double dose.

What are possible side effects from using GLICLAZIDE MR?

These are not all the possible side effects you may have when taking GLICLAZIDE MR. If you experience any side effects not listed here, tell your healthcare professional.

- low blood sugar (hypoglycemia)
- high blood sugar (hyperglycemia)
- viral infection
- infection of the nose, sinuses, or throat (upper respiratory infection)
- runny nose
- sore throat
- cough
- back, muscle and joint pain
- headache
- high blood pressure (hypertension)
- chest pain (angina)
- leg swelling
- diarrhea
- constipation
- abdominal pain
- nausea
- dizziness
- skin rash
- itching
- depression

- urinary tract infection
- pink eye (conjunctivitis)

Serious side effects and what to do about them			
Symptom / effect	Talk to your healthcare professional		Stop taking drug and get immediate medical help
	Only if severe	In all cases	
COMMON			
Low blood sugar level (hypoglycemia): anxious feeling, chest pain or pressure, chills, clammy skin, cold sweats, confusion, cool pale skin, depression, difficulty in concentration, dizziness, drowsiness, excessive hunger, fast or irregular heartbeat, headache, high blood pressure, nausea, nervousness, shakiness, shortness of breath, unsteady walk, unusual tiredness or weakness		√	
UNCOMMON			
Chest pain or pressure, and/or shortness of breath			√
Oedema , swelling of the legs or unexpected weight gain		√	
Skin rash , redness, itching or hives			√
Unexplained fever , chills or sore throat			√
Yellowing of skin or eyes, dark- coloured urine or light- coloured bowel movements (e.g. jaundice)			√

Serious side effects and what to do about them			
Symptom / effect	Talk to your healthcare professional		Stop taking drug and get immediate medical help
	Only if severe	In all cases	
VERY RARE			
Allergic inflammation of blood vessels (vasculitis)			√
Blood abnormalities with symptoms of sore throat, fever, mouth sore, unusual bleeding or bruising, low level of red blood cells (anemia)			√
Low sodium level in blood combined with symptoms of tiredness, weakness and confusion (hyponatraemia)			√
Rapid swelling of tissues such as eyelids, face, lips, mouth, tongue or throat that may result in breathing difficulty (angioedema)			√
Serious Skin Reactions (DRESS, Stevens-Johnson Syndrome, Toxic Epidermal Necrolysis, bullous pemphigoid, hypersensitivity Syndrome): any combination of red itchy rash with blisters and peeling of the skin and/or of the lips, eyes, mouth, nasal passages or genitals. It often goes with fever, chills, headache, cough, body aches or joint pain. You may have less or dark urine, yellow skin or eyes			√

If you have a troublesome symptom or side effect that is not listed here or becomes bad enough

to interfere with your daily activities, tell your healthcare professional.

Reporting Side Effects

You can report any suspected side effects associated with the use of health products to Health Canada by:

- Visiting the Web page on Adverse Reaction Reporting (<https://www.canada.ca/en/health-canada/services/drugs-health-products/medeffect-canada/adverse-reaction-reporting.html>) for information on how to report online, by mail or by fax; or
- Calling toll-free at 1-866-234-2345.

NOTE: Contact your health professional if you need information about how to manage your side effects. The Canada Vigilance Program does not provide medical advice.

Storage:

- Keep out of reach and sight of children and pets.
- GLICLAZIDE MR should be stored at room temperature (15°C to 30°C).
- Medicines should not be disposed of down the drain or in household garbage. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

If you want more information about GLICLAZIDE MR:

- Talk to your healthcare professional
- Find the full product monograph that is prepared for healthcare professionals and includes this Patient Medication Information by visiting the Health Canada website: (<https://health-products.canada.ca/dpd-bdpp/index-eng.jsp>); the manufacturer's website (<https://www.sanis.com/products/>), or by calling 1-866-236-4076.

This leaflet was prepared by Sanis Health Inc.

Last Revised: MAR 27, 2024