

PRODUCT MONOGRAPH  
INCLUDING PATIENT MEDICATION INFORMATION

Pr **SANDOZ® FESOTERODINE FUMARATE**  
fesoterodine fumarate extended-release tablets

Extended-release Tablets 4 mg and 8 mg Oral

Anticholinergic - Antispasmodic Agent

Sandoz Canada Inc.  
110 de Lauzon Street  
Boucherville, Quebec  
J4B 1E6

Date of Initial Authorization:  
OCT 25, 2021

Date of Revision  
APR 26, 2024

Submission Control No: 281797

**RECENT MAJOR LABEL CHANGES**

Not applicable

**TABLE OF CONTENTS**

**Sections or subsections that are not applicable at the time of authorization are not listed.**

**TABLE OF CONTENTS ..... 2**

**PART I: HEALTH PROFESSIONAL INFORMATION..... 4**

**1 INDICATIONS..... 4**

1.1 Pediatrics.....4

1.2 Geriatrics.....4

**2 CONTRAINDICATIONS ..... 4**

**4 DOSAGE AND ADMINISTRATION ..... 5**

4.1 Dosing Considerations .....5

4.2 Recommended Dose and Dosage Adjustment .....5

4.3 Reconstitution .....5

4.4 Administration.....5

4.5 Missed Dose .....6

**5 OVERDOSAGE..... 6**

**6 DOSAGE FORMS, STRENGTHS, COMPOSITION AND PACKAGING ..... 6**

**7 WARNINGS AND PRECAUTIONS ..... 7**

7.1 Special Populations.....10

7.1.1 Pregnant Women .....10

7.1.2 Breast-feeding .....10

7.1.3 Pediatrics.....10

7.1.4 Geriatrics.....10

**8 ADVERSE REACTIONS ..... 11**

8.1 Adverse Reaction Overview.....11

8.2 Clinical Trial Adverse Reactions .....11

8.2.1 Clinical Trial Adverse Reactions – Pediatrics.....13

8.3 Less Common Clinical Trial Adverse Reactions .....14

8.3.1 Less Common Clinical Trial Adverse Reactions – Pediatrics.....14

8.4 Abnormal Laboratory Findings: Hematologic, Clinical Chemistry and Other

	Quantitative Data .....	14
	8.5 Post-Market Adverse Reactions .....	14
<b>9</b>	<b>DRUG INTERACTIONS .....</b>	<b>14</b>
	9.2 Drug Interactions Overview .....	14
	9.3 Drug-Behavioural Interactions.....	15
	9.4 Drug-Drug Interactions .....	15
	9.5 Drug-Food Interactions.....	16
	9.6 Drug-Herb Interactions.....	17
	9.7 Drug-Laboratory Test Interactions.....	17
<b>10</b>	<b>CLINICAL PHARMACOLOGY .....</b>	<b>17</b>
	10.1 Mechanism of Action.....	17
	10.2 Pharmacodynamics .....	17
	10.3 Pharmacokinetics .....	18
<b>11</b>	<b>STORAGE, STABILITY AND DISPOSAL .....</b>	<b>21</b>
<b>12</b>	<b>SPECIAL HANDLING INSTRUCTIONS.....</b>	<b>21</b>
	<b>PART II: SCIENTIFIC INFORMATION .....</b>	<b>22</b>
<b>13</b>	<b>PHARMACEUTICAL INFORMATION .....</b>	<b>22</b>
<b>14</b>	<b>CLINICAL TRIALS .....</b>	<b>23</b>
	14.1 Trial Design and Study Demographics .....	23
	14.2 Study Results .....	23
	14.3 Comparative Bioavailability Studies.....	28
	14.4 Immunogenicity.....	30
<b>15</b>	<b>MICROBIOLOGY .....</b>	<b>30</b>
<b>16</b>	<b>NON-CLINICAL TOXICOLOGY .....</b>	<b>30</b>
<b>17</b>	<b>SUPPORTING PRODUCT MONOGRAPHS.....</b>	<b>33</b>
	<b>PATIENT MEDICATION INFORMATION.....</b>	<b>34</b>

## **PART I: HEALTH PROFESSIONAL INFORMATION**

### **1 INDICATIONS**

Sandoz Fesoterodine Fumarate (fesoterodine fumarate extended-release tablet) is indicated for:

- The treatment of patients with overactive bladder with symptoms of urinary frequency, urgency, or urge incontinence, or any combination of these symptoms.

#### **1.1 Pediatrics**

Pediatrics (< 18 years of age): Based on the data submitted and reviewed by Health Canada, the safety and efficacy of fesoterodine fumarate extended-release in pediatric patients has not been established; therefore, Health Canada has not authorized an indication for pediatric use. See [7 WARNINGS AND PRECAUTIONS, 7.1.3 Pediatrics](#); [8 ADVERSE REACTIONS, 8.2 Clinical Trial Adverse Drug Reactions, 8.2.1 Pediatrics](#); [10 CLINICAL PHARMACOLOGY, 10.3 Pharmacokinetics, Special populations, Pediatrics](#).

#### **1.2 Geriatrics**

Geriatrics ( $\geq$  65 years of age): Based on clinical studies, no apparent overall differences were observed in safety between older (patients  $\geq$  65 years) and younger patients (patients < 65 years) on fesoterodine extended-release tablets. Therefore, dosage adjustment for geriatric patients may not be required. See [7 WARNINGS AND PRECAUTIONS, 7.1.4 Geriatrics](#); [8 ADVERSE REACTIONS, 8.2 Clinical Trial Adverse Drug Reactions](#); [10 CLINICAL PHARMACOLOGY 10.3 Pharmacokinetics, Special Populations, Geriatrics](#).

### **2 CONTRAINDICATIONS**

Sandoz Fesoterodine Fumarate (fesoterodine fumarate extended-release tablet) is contraindicated in patients with:

- Urinary retention
- Gastric retention
- Uncontrolled narrow-angle glaucoma
- Hypersensitivity to tolterodine L-tartrate tablets, tolterodine L-tartrate extended-release capsules, soya, peanuts, lactose.
- Hypersensitivity to this drug, or to any ingredient in the formulation, including any non-medicinal ingredient, or component of the container. For a complete listing, see [6 DOSAGE FORMS, STRENGTHS, COMPOSITION AND PACKAGING](#).

## 4 DOSAGE AND ADMINISTRATION

### 4.1 Dosing Considerations

Dosing of Sandoz Fesoterodine Fumarate (fesoterodine fumarate) may be affected by the following:

- Individual response and tolerability
- Impaired hepatic function and renal impairment
- Potent CYP3A4 inhibitors

(See [7 WARNINGS AND PRECAUTIONS](#) and [4 DOSAGE AND ADMINISTRATION, 4.2 Recommended Dose and Dosage Adjustment](#).)

### 4.2 Recommended Dose and Dosage Adjustment

The recommended starting dose of Sandoz Fesoterodine Fumarate is 4 mg once daily. Based upon individual response and tolerability, the dose may be increased to 8 mg once daily.

The daily dose of Sandoz Fesoterodine Fumarate should not exceed 4 mg in the following populations:

- Patients with severe renal impairment ( $CL_{CR} < 30$  mL/min)
- Patients taking potent CYP3A4 inhibitors, such as ketoconazole, itraconazole, miconazole, and clarithromycin.

Sandoz Fesoterodine Fumarate is not recommended for use in patients with severe hepatic impairment (Child-Pugh C). Dosage adjustment may not be necessary for elderly patients ( $\geq 65$  years of age).

See [7 WARNINGS AND PRECAUTIONS, Hepatic/Biliary/Pancreatic](#); [7 WARNINGS AND PRECAUTIONS, Renal](#); See [7 WARNINGS AND PRECAUTIONS, 7.1.4 Geriatrics](#); [10 CLINICAL PHARMACOLOGY 10.3 Pharmacokinetics, Special Populations, Geriatrics](#).

Health Canada has not authorized an indication for pediatric use. See [1 INDICATIONS, 1.1 Pediatrics](#); [7 WARNINGS AND PRECAUTIONS, 7.1.3 Pediatrics](#); [10 CLINICAL PHARMACOLOGY, 10.3 Pharmacokinetics, Special populations, Pediatrics](#)

### 4.3 Reconstitution

Not applicable.

### 4.4 Administration

Sandoz Fesoterodine Fumarate tablets should be taken with liquid and swallowed whole. Sandoz Fesoterodine Fumarate can be administered with or without food, and should not be chewed, divided, or crushed. Sandoz Fesoterodine Fumarate may be taken during the day or at night (see [10 CLINICAL PHARMACOLOGY, 10.3 Pharmacokinetics Daytime versus Nighttime](#)).

#### 4.5 Missed Dose

If a dose of Sandoz Fesoterodine Fumarate tablets is missed, then it should be taken as soon as the patient remembers unless it is almost time for the next dose, in which case the patient should not take the missed dose. Patients should not take 2 doses at the same time to make up for a missed dose.

#### 5 OVERDOSAGE

Overdosage with fesoterodine could result in severe antimuscarinic effects and should be treated accordingly.

Treatment of overdosage with fesoterodine should consist of gastric lavage and activated charcoal. Treatments for symptoms are recommended as follows. For severe central anticholinergic effects (hallucinations, severe excitation), an anticholinesterase agent, such as physostigmine, may be used. If excitation and convulsions occur, administer an anticonvulsant, such as diazepam. Patients with respiratory insufficiency should be given respiratory assistance. If respiratory arrest occurs, patients should be given artificial respiration. Patients with tachycardia may be treated with a beta-blocker, and those with urinary retention may be catheterized. Patients with troublesome mydriasis may be placed in a dark room or treated with pilocarpine eye drops, or both. ECG should be monitored.

For management of a suspected drug overdose, contact your regional poison control centre.

#### 6 DOSAGE FORMS, STRENGTHS, COMPOSITION AND PACKAGING

Table 1 – Dosage forms, strengths, composition and packaging

Route of Administration	Dosage Form / Strength	All Nonmedicinal Ingredients
Oral	Extended-release tablets 4 mg, 8 mg	Fumaric acid, glycerol dibehenate, hypromellose (methocel K4M and K100M premium CR), hydroxypropyl cellulose, lactose monohydrate, microcrystalline cellulose, pullulan, talc. The film coating contains the following inactive ingredients: Lecithin (soya), macrogol/PEG, FD&C blue #2/ indigo carmine aluminum lake, polyvinyl alcohol-part. hydrolyzed, talc, titanium dioxide.

## Description

Sandoz Fesoterodine Fumarate (fesoterodine fumarate extended-release tablets) is available as:

- 4 mg tablets (light blue oval, biconvex, film-coated tablets debossed with I11 on one side and plain on other side)
- 8 mg tablets (blue, oval, biconvex, film-coated tablets debossed with I17 on one side and plain on other side.)

Sandoz Fesoterodine Fumarate (fesoterodine fumarate extended-release tablets) is supplied as follows:

- Bottles of 30 and 100 tablets
- Blister Strips boxed as 30 tablets

## 7 WARNINGS AND PRECAUTIONS

### Carcinogenesis and Mutagenesis

See [16 NON-CLINICAL TOXICOLOGY, Carcinogenicity](#)

### Cardiovascular

Fesoterodine, like other antimuscarinic drugs, is associated with increased heart rate that correlates with increasing dose. See [10 CLINICAL PHARMACOLOGY, 10.2 Pharmacodynamics, Cardiac Electrophysiology and Hemodynamics](#). Although there are no clinical trial or post-marketing data to confirm the potential for fesoterodine to aggravate certain pre-existing cardiac conditions, this product is in the class anticholinergic medications which are known to have cardiac effects. Prescribers should therefore use caution when prescribing Sandoz Fesoterodine Fumarate to patients with ischemic heart disease, congestive heart failure, cardiac arrhythmias, or tachycardia.

### Driving and Operating Machinery

Patients should be advised not to engage in potentially hazardous activities, such as driving or operating a vehicle or potentially dangerous machinery, until they know how Sandoz Fesoterodine Fumarate may affect them (see [8 ADVERSE REACTIONS, 8.5 Post-Market Adverse Reactions](#)).

### Endocrine and Metabolism

CYP3A4: Caution should be exercised when prescribing or up-titrating fesoterodine from 4 mg to 8 mg in patients in whom an increased exposure to the active metabolite is expected, such as with concomitant administration of CYP3A4 inhibitors.

In the presence of a potent CYP3A4 inhibitor (e.g. ketoconazole, itraconazole, miconazole and clarithromycin), doses of Sandoz Fesoterodine Fumarate greater than 4 mg are not recommended.

In the presence of moderate CYP3A4 inhibitors (e.g. fluconazole), no dosing adjustments are recommended.

While the effect of weak CYP3A4 inhibitors (e.g. cimetidine) was not examined in a clinical study, some pharmacokinetic interaction is expected, though less than what was observed with moderate CYP3A4 inhibitors. See [4 DOSAGE AND ADMINISTRATION](#); [9 DRUG INTERACTIONS, 9.4 Drug-Drug Interactions](#); [10 CLINICAL PHARMACOLOGY, 10.3 Pharmacokinetics, Metabolism](#)).

CYP2D6: A subset of individuals are poor metabolizers for CYP2D6 (see [10 CLINICAL PHARMACOLOGY, 10.3 Pharmacokinetics, Metabolism](#)).

Compared with CYP2D6 extensive metabolizers not taking ketoconazole (a potent CYP3A4 inhibitor), further increases in the exposure to the active metabolite of fesoterodine were observed in subjects who were CYP2D6 poor metabolizers taking ketoconazole. See [9 DRUG INTERACTIONS, 9.4 Drug-Drug Interactions](#).

### **Gastrointestinal**

Patients at Risk of Gastric Retention: Sandoz Fesoterodine Fumarate (fesoterodine fumarate extended-release tablets), like other antimuscarinic drugs, should be administered with caution to patients with decreased gastrointestinal motility, including patients with severe constipation and to patients with gastrointestinal obstruction disorders (e.g. pyloric stenosis) because of the risk of gastric retention (see [2 CONTRAINDICATIONS](#)).

### **Genitourinary**

Patients at Risk of Urinary Retention: Sandoz Fesoterodine Fumarate, like other antimuscarinic drugs, should be administered with caution to patients with clinically significant bladder outlet obstruction because of the risk of urinary retention. See [2 CONTRAINDICATIONS](#); [9 DRUG INTERACTIONS, 9.2 Drug Interactions Overview](#).

### **Hepatic/Biliary/Pancreatic**

Sandoz Fesoterodine Fumarate should be administered with caution to patients with impaired hepatic function. In patients with mild to moderate hepatic impairment, no dosage adjustment is required. Fesoterodine is not recommended for use in patients with severe hepatic impairment. See [4 DOSAGE AND ADMINISTRATION](#), [10 CLINICAL PHARMACOLOGY, 10.3 Pharmacokinetics, Special Populations and Conditions, Hepatic Insufficiency](#).



## Immune

Angioedema of the face, lips, tongue, and/or larynx has been reported with fesoterodine. In some cases angioedema occurred after the first dose. Angioedema associated with upper airway swelling may be life-threatening. If involvement of the tongue, hypopharynx, or larynx occurs, fesoterodine should be promptly discontinued and appropriate therapy and/or measures to ensure a patent airway should be promptly provided. See [8 ADVERSE REACTIONS, 8.5 Post-Market Adverse Reactions](#).

Lactose: Sandoz Fesoterodine Fumarate tablets contain lactose. Patients with rare hereditary problems of galactose intolerance, the Lapp lactase deficiency or glucose-galactose malabsorption should not take this medicinal product. See [2 CONTRAINDICATIONS](#).

## Neurologic

Sandoz Fesoterodine Fumarate, like other antimuscarinic drugs, should be administered with caution to patients with myasthenia gravis.

Sandoz Fesoterodine Fumarate is associated with anticholinergic central nervous system (CNS) effects. A variety of CNS anticholinergic effects have been reported, including headache, dizziness, and somnolence (see [8.5 Post-Market Adverse Reactions](#)). Patients should be monitored for signs of anticholinergic CNS effects, particularly after beginning treatment or increasing the dose. Advise patients not to drive or operate heavy machinery until they know how Sandoz Fesoterodine Fumarate affects them. If a patient experiences anticholinergic CNS effects, dose reduction or drug discontinuation should be considered.

## Ophthalmologic

Sandoz Fesoterodine Fumarate, like other antimuscarinic drugs, should be used with caution in patients being treated for narrow-angle glaucoma (see [2 CONTRAINDICATIONS](#)).

## Renal

Sandoz Fesoterodine Fumarate should be administered with caution to patients with impaired renal function. In patients with mild to moderate renal impairment, no dosage adjustment is required. Doses of fesoterodine greater than 4 mg are not recommended in patients with severe renal impairment ( $CL_{CR} < 30$  mL/min). See [4 DOSAGE AND ADMINISTRATION; 10 CLINICAL PHARMACOLOGY, 10.3 Pharmacokinetics, Special Populations and Conditions, Renal Insufficiency](#).

## Reproductive Health: Female and Male Potential

- **Fertility:** No clinical trials have been conducted to assess the effect of fesoterodine on human fertility.  
Findings in mice at maternally toxic doses at exposures approximately 5 to 19 times (based

on lowest and highest total systemic exposure) those at the Maximum Recommended Human Dose (MRHD) show an effect on female fertility, however, the clinical implications of these animal findings are not known. Fesoterodine had no effect on male reproductive function or fertility in mice at doses up to 45 mg/kg/day. See [16 Non-Clinical Toxicology, Reproductive and Developmental Toxicology](#).

- **Teratogenic Risk**

No dose-related teratogenicity was observed in reproduction studies performed in mice and rabbits. See [16 Non-Clinical Toxicology, Reproductive and Developmental Toxicology](#).

## **Special Populations**

### **7.1.1 Pregnant Women**

There are no adequate data from the use of fesoterodine in pregnant women. The potential risk for humans is unknown. Therefore, fesoterodine should be used during pregnancy only if the potential benefit to the mother outweighs the potential risk to the foetus. Women of childbearing potential should be considered for treatment only if using adequate contraception. In animal reproduction studies, oral administration of fesoterodine to pregnant mice and rabbits during organogenesis resulted in fetotoxicity at maternal exposures that were 7 and 6 times the MRHD, respectively, based on the lowest unbound systemic exposure. See [16 Non-Clinical Toxicology, Reproductive and Developmental Toxicology](#).

### **7.1.2 Breast-feeding**

It is not known whether fesoterodine is excreted into human milk; therefore, breastfeeding is not recommended during treatment with fesoterodine

### **7.1.3 Pediatrics**

Pediatrics (< 18 years of age): The safety and efficacy of fesoterodine in pediatric patients have not been established. See [8 ADVERSE REACTIONS, 8.2 Clinical Trial Adverse Drug Reactions, 8.2.1 Pediatrics](#); [10 CLINICAL PHARMACOLOGY, 10.3 Pharmacokinetics, Special populations, Pediatrics](#).

### **7.1.4 Geriatrics**

Geriatrics (≥ 65 years of age): No overall differences in safety or effectiveness were observed between patients younger than 65 years of age and those 65 years of age or older in the clinical studies. However, patients in these studies were highly selected and relatively healthy. The pharmacokinetics of fesoterodine are not significantly influenced by age. Dose adjustment may not be required for the elderly. See [10 CLINICAL PHARMACOLOGY 10.3 Pharmacokinetics, Special Populations, Geriatrics](#).

## 8 ADVERSE REACTIONS

### 8.1 Adverse Reaction Overview

Due to the pharmacological properties of fesoterodine, treatment may cause mild to moderate antimuscarinic effects like dry mouth, constipation, dry eyes, and dyspepsia.

### 8.2 Clinical Trial Adverse Reactions

Clinical trials are conducted under very specific conditions. The adverse reaction rates observed in the clinical trials; therefore, may not reflect the rates observed in practice and should not be compared to the rates in the clinical trials of another drug. Adverse reaction information from clinical trials may be useful in identifying and approximating rates of adverse drug reactions in real-world use. The safety of fesoterodine fumarate extended-release tablet was primarily evaluated in Phase 2 and 3 controlled trials in a total of 2859 patients with overactive bladder of which 2288 were treated with fesoterodine. Of this total, 782 received fesoterodine 4 mg/day, and 785 received fesoterodine 8 mg/day in Phase 2 or 3 studies with treatment periods of 8 or 12 weeks. Approximately 80% of these patients had >10 weeks exposure to fesoterodine.

A total of 1964 patients participated in two 12-week, Phase 3 efficacy and safety studies and subsequent open-label extension studies. In these 2 studies combined, 554 patients received fesoterodine 4 mg/day and 566 patients received fesoterodine 8 mg/day.

In Phase 2 and 3 placebo-controlled trials combined, the incidences of serious adverse events in patients receiving placebo, fesoterodine 4 mg, and fesoterodine 8 mg were 1.9%, 3.5%, and 2.9%, respectively. All serious adverse events were judged to be not related or unlikely to be related to study medication by the investigator, except for four patients receiving fesoterodine who reported one serious adverse event each: angina, chest pain, gastroenteritis, and QT prolongation on ECG.

The most commonly reported adverse event in patients treated with fesoterodine was dry mouth. The incidence of dry mouth was higher in those taking 8 mg/day (35%) and in those taking 4 mg/day (19%), as compared to placebo (7%). Dry mouth led to discontinuation in 0.4%, 0.4%, and 0.8% of patients receiving placebo, fesoterodine 4 mg, and fesoterodine 8 mg, respectively. For those patients who reported dry mouth, most had their first occurrence of the event within the first month of treatment.

The second most commonly reported adverse event was constipation. The incidence of constipation was 2% in those taking placebo, 4% in those taking fesoterodine 4 mg/day, and 6% in those taking fesoterodine 8 mg.

Table 2 lists adverse events, regardless of causality, that were reported in the combined Phase 3, randomized, placebo-controlled trials at an incidence greater than placebo and in 1% or more of patients treated with fesoterodine 4 or 8 mg once daily for up to 12 weeks.

**Table 2 - Adverse events with an incidence exceeding the placebo rate and reported by  $\geq 1\%$  of patients from double-blind, placebo-controlled Phase 3 trials of 12 weeks treatment duration**

<b>System organ class/Preferred term</b>	<b>Placebo N=554 %</b>	<b>Fesoterodine 4mg/day N=554 %</b>	<b>Fesoterodine 8mg/day N=566 %</b>
Gastrointestinal disorders			
Dry mouth	7.0	18.8	34.6
Constipation	2.0	4.2	6.0
Dyspepsia	0.5	1.6	2.3
Nausea	1.3	0.7	1.9
Abdominal pain upper	0.5	1.1	0.5
Infections			
Urinary tract infection	3.1	3.2	4.2
Upper respiratory tract infection	2.2	2.5	1.8
Eye Disorders			
Dry eyes	0	1.4	3.7
Renal and urinary disorders			
Dysuria	0.7	1.3	1.6
Urinary retention	0.2	1.1	1.4
Respiratory disorders			
Cough	0.5	1.6	0.9
Dry Throat	0.4	0.9	2.3
General disorders			
Edema peripheral	0.7	0.7	1.2
Musculoskeletal disorders			
Back pain	0.4	2.0	0.9
Psychiatric disorders			
Insomnia	0.5	1.3	0.4
Investigations			
ALT increased	0.9	0.5	1.2
GGT increased	0.4	0.4	1.2
Skin disorders			
Rash	0.5	0.7	1.1

ALT=alanine aminotransferase, GGT=gamma glutamyltransferase

Patients also received fesoterodine for up to three years in open-label extension phases of one Phase 2 and two Phase 3 controlled trials. In all open-label trials combined, 857, 701, 529, and 105 patients received fesoterodine for at least 6 months, 1 year, 2 years, and 3 years respectively.

The adverse events observed during long-term, open-label studies were similar to those observed in the 12-week, placebo-controlled studies, and included dry mouth, constipation, dry eyes, dyspepsia and abdominal pain. Similar to the controlled studies, most adverse events of dry mouth and constipation were mild to moderate in intensity. Serious adverse events, judged to be at least possibly related to study medication by the investigator, and reported more than once during the open-label treatment period of up to 3 years included urinary retention (3 cases), diverticulitis (3 cases), constipation (2 cases), irritable bowel syndrome (2 cases), and electrocardiogram QT corrected interval prolongation (2 cases).

The safety of fesoterodine was further established in two additional 12-week, active- and placebo-controlled, double-blind, randomized studies comparing fesoterodine with tolterodine ER 4 mg and placebo. In these studies combined, 1527 patients received fesoterodine 8 mg, 1552 patients received tolterodine ER 4 mg, and 755 patients received placebo. The most common treatment-emergent adverse events (dry mouth, constipation, and headache) reported with fesoterodine during these 2 studies were similar to those observed in the 12-week, placebo-controlled studies.

Fesoterodine was associated with an increase in heart rate that correlated with increasing dose, a well-characterized effect described for antimuscarinic drugs. In the placebo-controlled phase 3 studies in patients with overactive bladder, the mean increases in heart rate compared to placebo were approximately 3-4 beats/minute in the 4 mg/day group and 3-5 beats/minute in the 8 mg/day group. See [10 CLINICAL PHARMACOLOGY, 10.2 Pharmacodynamics, Cardiac Electrophysiology and Hemodynamics](#).

Geriatrics ( $\geq 65$  years of age): Of 1567 patients who received fesoterodine 4mg/day or 8mg/day in the Phase 2 and 3, placebo-controlled, efficacy and safety studies, 515 (33%) were 65 years of age or older, and 140 (9%) were 75 years of age or older. No overall differences in safety or efficacy were observed between patients younger than 65 years of age and those 65 years of age or older in these studies; however, the incidence of antimuscarinic adverse events, including dry mouth, constipation, dyspepsia, increase in residual urine, dizziness (at 8mg only) and urinary tract infection, was higher in patients 75 years of age and older as compared to younger patients.

### **8.2.1 Clinical Trial Adverse Reactions – Pediatrics**

Pediatrics (<18 years of age): An open-label pediatric Phase 2 study with fesoterodine in overactive bladder (N=10) or neurogenic detrusor overactivity (NDO: N=11) patients aged 9 to 17 with body weight >25 kg was conducted. Patients received 4 mg once daily (N = 21) for 4 weeks, followed by dose escalation to 8 mg once daily (N = 20) for a further 4 weeks. One patient with NDO receiving fesoterodine 8 mg once daily experienced a treatment-related serious adverse event of constipation which required hospitalization and temporary discontinuation of fesoterodine. The safety and efficacy of fesoterodine in pediatric populations have not been established.

### **8.3 Less Common Clinical Trial Adverse Reactions**

This information is not available for this drug product.

#### **8.3.1 Less Common Clinical Trial Adverse Reactions – Pediatrics**

Less common clinical trial adverse reactions in the pediatric population have not been identified. Fesoterodine fumarate is not authorized for use in pediatric patients. See [1 INDICATIONS, 1.1 Pediatrics](#).

### **8.4 Abnormal Laboratory Findings: Hematologic, Clinical Chemistry and Other Quantitative Data**

In clinical trials comparing fesoterodine to placebo, cases of markedly elevated liver enzymes (ALT increased, GGT increased) were reported at a frequency no different than placebo. The relation to fesoterodine treatment is unclear.

### **8.5 Post-Market Adverse Reactions**

The following events have been reported in association with fesoterodine use in worldwide post-marketing experience:

- Eye disorders: Blurred vision;
- Cardiac disorders: Palpitations;
- Central nervous system disorders: Dizziness, headache; somnolence, hypoesthesia
- Skin and subcutaneous tissue disorders: Angioedema including angioedema with airway obstruction, face edema, hypersensitivity reactions, urticaria, pruritus, rash, pharyngeal oedema, pharyngeal swelling;
- Renal and urinary disorders: Urinary retention.

Because these spontaneously reported events are from the worldwide post-marketing experience, the frequency of the events and the role of fesoterodine in their causation cannot be reliably determined.

## **9 DRUG INTERACTIONS**

### **9.2 Drug Interactions Overview**

Coadministration of fesoterodine fumarate extended-release tablet with other medicinal products with anticholinergic properties may result in more pronounced therapeutic and/or adverse effects. Fesoterodine is rapidly metabolized to active metabolite, 5-hydroxymethyl tolterodine (5-HMT), by nonspecific esterases; this active metabolite of fesoterodine is further metabolized, principally via CYP2D6 and CYP3A4. At therapeutic concentrations, 5-HMT does not inhibit CYP isoenzymes 1A2, 2B6, 2C8, 2C9, 2C19, 2D6, 2E1, or 3A4 and does not induce CYP isoenzymes 1A2, 2B6, 2C9, 2C19, or 3A4.

Use with Other Concomitant Therapies: Alpha-blockers for lower urinary tract symptoms (LUTS) in

men: fesoterodine fumarate extended-release tablet efficacy was not established in a study of men 40 years and older with overactive bladder symptoms taking an alpha-blocker for lower urinary tract symptoms (LUTS). No excess incidence of acute urinary retention was demonstrated. However, urinary treatment-emergent events such as urinary retention and dysuria were reported more often by men in the fesoterodine add-on group relative to the placebo add-on group (urinary retention: 2.3% versus 0.4% and dysuria: 3.2% versus 0.6%). Caution should be used when administering fesoterodine fumarate extended-release tablet to men with possible bladder outlet obstruction. See [7 WARNINGS AND PRECAUTIONS, Genitourinary](#).

### 9.3 Drug-Behavioural Interaction

Interactions with individual behaviour have not been established.

### 9.4 Drug-Drug Interactions

The drugs listed in Table 3 are based on either drug interaction case reports or studies, or potential interactions due to the expected magnitude and seriousness of the interaction (i.e., those identified as contraindicated).

**Table 3 - Established or Potential Drug-Drug Interactions**

Proper/Common name	Source of Evidence	Effect	Clinical Comment
Ketoconazole (potent CYP3A4 inhibitors)	CT	The effect of ketoconazole 200 mg twice daily for 5 days increased $C_{max}$ and AUC of the active metabolite of fesoterodine by 2.0- and 2.3-fold, respectively after oral administration of fesoterodine 8 mg to CYP2D6 extensive metabolizers. In CYP2D6 poor metabolizers, the effect of ketoconazole 200 mg twice daily for 5 days increased $C_{max}$ and AUC of the active metabolite of fesoterodine by 2.1- and 2.5-fold, respectively. Furthermore, in subjects who were CYP2D6 poor metabolizers and taking ketoconazole versus subjects who were CYP2D6 extensive metabolizers and not taking ketoconazole, the $C_{max}$ and AUC increased by 4.5 and 5.7 fold, respectively. The effect of ketoconazole 200 mg once a day for 5 days increased $C_{max}$ and AUC of the active metabolite of fesoterodine by 2.2-fold in CYP2D6 extensive metabolizers and 1.5- and 1.9- fold, respectively, in CYP2D6 poor metabolizers. Furthermore, in subjects who were CYP2D6 poor metabolizers and taking ketoconazole versus subjects who were CYP2D6 extensive metabolizers and not taking ketoconazole, the $C_{max}$ and AUC increased by 3.4 and 4.2 fold, respectively.	Dose of fesoterodine greater than 4 mg are not recommended in patients taking potent CYP3A4 inhibitors such as ketoconazole, itraconazole, miconazole and clarithromycin

**Table 3 - Established or Potential Drug-Drug Interactions**

Proper/Common name	Source of Evidence	Effect	Clinical Comment
Fluconazole (moderate CYP3A4 inhibitors)	CT	Co-administration of fesoterodine 8 mg with fluconazole 200 mg twice daily increased $C_{max}$ and $AUC_{inf}$ of the active metabolite of fesoterodine by approximately 19% (11% - 28%) and 27% (18% - 36%), respectively.	The increase in the active metabolite of fesoterodine is not considered clinically relevant. No dosage adjustment is recommended when fesoterodine is co-administered with a moderate CYP3A4 inhibitor.
Cimetidine (Weak CYP3A4 inhibitors)	T	The effect of weak CYP3A4 inhibitors was not examined; it is not expected to be in excess of the effect of moderate inhibitors.	
Rifampicin (CYP3A4 inducers)	CT	Following induction of CYP3A4 by rifampicin 600 mg once a day, $C_{max}$ and AUC of the active metabolite of fesoterodine decreased by approximately 70% and 75%, respectively, after oral administration of fesoterodine 8 mg. The terminal half-life of the active metabolite was not changed.	Induction of CYP3A4 may lead to reduced plasma levels of the active metabolite of fesoterodine. No dosing adjustments are recommended in the presence of CYP3A4 inducers such as rifampicin or carbamazepine. However, concomitant use of CYP3A4 inducers is not recommended.
CYP2D6 inhibitors	T	In poor metabolizers for CYP2D6, $C_{max}$ and AUC of the active metabolite were increased 1.7- and 2-fold, respectively.	The interaction with CYP2D6 inhibitors was not tested clinically. No dosing adjustments are recommended in the presence of CYP2D6 inhibitors.
Warfarin	CT	A clinical study has shown in healthy volunteers that fesoterodine 8 mg once daily has no significant effect on the PK or the anticoagulant activity of a single 25 mg dose of warfarin. Standard therapeutic monitoring for warfarin should be continued.	
Oral contraceptives	CT	In the presence of fesoterodine, there were no clinically significant changes in the plasma concentrations of combined oral contraceptives containing 0.03 mg ethinyl estradiol and 0.15mg levonorgestrel	

Legend: C = Case Study; CT = Clinical Trial; T = Theoretical

## 9.5 Drug-Food Interactions

Fesoterodine tablets can be taken with or without food. There is no clinically relevant effect of



food on the pharmacokinetics of fesoterodine. Concomitant food intake increased the active metabolite of fesoterodine AUC by 19% and  $C_{max}$  by 18%. See [4 DOSAGE AND ADMINISTRATION, 4.4 Administration](#); [10 CLINICAL PHARMACOLOGY, 10.3 Pharmacokinetics, Effect of Food](#).

### 9.6 Drug-Herb Interactions

Interactions with herbal products have not been established.

### 9.7 Drug-Laboratory Test Interactions

Interactions with laboratory tests have not been established.

## 10 CLINICAL PHARMACOLOGY

### 10.1 Mechanism of Action

Fesoterodine is a unique competitive muscarinic receptor antagonist. After oral administration, fesoterodine is rapidly and extensively hydrolyzed by nonspecific esterases to its active metabolite, 5-hydroxymethyl tolterodine (5-HMT), which is responsible for the antimuscarinic activity of fesoterodine. The conversion of fesoterodine fumarate extended-release tablet to its active metabolite is not dependent on cytochrome P450 enzymes.

Muscarinic receptors play a role in contractions of urinary bladder smooth muscle and stimulation of salivary secretion. Inhibition of these receptors in the bladder is presumed to be the mechanism by which fesoterodine produces its effects.

### 10.2 Pharmacodynamics

In a urodynamic study involving patients with involuntary detrusor contractions, the effects after the administration of fesoterodine on the volume at first detrusor contraction and bladder capacity were assessed. Administration of fesoterodine increased the volume at first detrusor contraction and bladder capacity in a dose-dependent manner. These findings are consistent with an antimuscarinic effect on the bladder.

### Cardiac Electrophysiology and Hemodynamics

The effect of fesoterodine 4 mg (therapeutic dose) and 28 mg (supratherapeutic dose) on the ECG parameters was evaluated in a double-blind, randomized, placebo- and positive-controlled (moxifloxacin 400 mg once a day) parallel group trial with once daily treatment over a period of 3 days in 261 male and female subjects aged 44 to 65 years. Electrocardiographic parameters were measured over a 24-hour period at pre-dose, after the first administration, and after the third administration of study medication. Fesoterodine 28 mg was chosen because this dose, when administered to CYP2D6 extensive metabolizers, results in an exposure to the active metabolite that is similar to the exposure in a CYP2D6 poor metabolizer receiving fesoterodine 8 mg together with CYP3A4 blockade. The study demonstrated that

fesoterodine at doses of 4 and 28 mg/day did not prolong the QTc interval, the QRS duration, or the PR interval in a treatment related manner.

Fesoterodine was associated with an increase in heart rate that correlated with increasing dose, a well-characterized effect described for antimuscarinic drugs. On day 3 of the study described above, when compared to placebo, the mean increases in heart rate averaged over 24 h, were 3 beats/minute for 4 mg/day fesoterodine and 11 beats/minute for 28 mg/day fesoterodine (see [ADVERSE REACTIONS](#)).

Routine safety monitoring in this study included blood pressure assessment. On day 3 of treatment, blood pressure measurements were performed at 4-5 h post-dosing. Mean changes from baseline in systolic blood pressure were -1.9 mmHg (90% CI: -4.0, 0.1) with fesoterodine 4 mg/day, 0.3 mmHg (90% CI: -2.4, 2.9) with fesoterodine 28 mg/day, and -3.8 mmHg (90% CI: -6.1, -1.5) with placebo. Mean changes from baseline in diastolic blood pressure were 1.4 mmHg (90% CI: -0.2, 3.1) with fesoterodine 4 mg/day, 3.7 mmHg (90% CI: -1.9, 5.4) with fesoterodine 28 mg/day, and -2.9 mmHg (90% CI -4.7, -1.1) with placebo. In phase 3 controlled clinical trials, systolic and diastolic blood pressure was assessed at steady state at each clinic visit. No difference from placebo was observed with fesoterodine at either 4mg/day or 8mg/day.

### **Cognitive Testing in Healthy Elderly**

A phase I, 4-treatment, cross-over, double-blind, placebo- and positive-controlled study in elderly healthy volunteers (n=20, mean age 72 years) evaluated the effect of fesoterodine 4 mg and 8 mg, placebo and alprazolam 1 mg (positive control) on a computer-based battery of cognitive tests and memory tests (CogState Tests: comprised of a detection task, an identification task, a one-card learning task, a continuous paired associate learning task, and the Groton maze learning task) and the Rey Auditory Verbal Learning Tests (RAVLT). There were no statistically significant differences between fesoterodine 4 mg and placebo (p=0.1198) or between fesoterodine 8 mg and placebo (p=0.2459) for the primary endpoint (CogState detection task). The validity of the study assessment was confirmed by the results of the positive control. Similar results were obtained for all other pharmacodynamics endpoints, including the battery of Rey Auditory Verbal Learning Tests, which were similarly statistically non-significant between fesoterodine and placebo.

### **10.3 Pharmacokinetics**

A summary of pharmacokinetic parameters for the active metabolite (5-HMT) after a single dose of fesoterodine 4 mg and 8 mg in extensive and poor metabolizers of CYP2D6 from subjects in a fasted state is provided in Table 4.

**Table 4 - Summary of geometric mean [CV] pharmacokinetic parameters for the active metabolite (5-HMT) after a single dose of Fesoterodine 4 mg and 8 mg in extensive and poor CYP2D6 metabolizers from subjects in a fasted state**

Parameter	Fesoterodine 4 mg		Fesoterodine 8 mg	
	EM (n=16)	PM (n=8)	EM (n=16)	PM (n=8)
C <sub>max</sub> (ng/mL)	1.89 [43%]	3.45 [54%]	3.98 [28%]	6.90 [39%]
AUC <sub>0-tz</sub> (ng*h/mL)	21.2 [38%]	40.5 [31%]	45.3 [32%]	88.7 [36%]
t <sub>max</sub> (h) <sup>a</sup>	5 [2-6]	5 [5-6]	5 [3-6]	5 [5-6]
t <sub>1/2</sub> (h)	7.31	7.31	8.59	7.66

EM = extensive CYP2D6 metabolizer, PM = poor CYP2D6 metabolizer, CV=coefficient of variation

C<sub>max</sub> = maximum plasma concentration, AUC<sub>0-tz</sub> = area under the concentration time curve from zero up to the last measurable plasma concentration, t<sub>max</sub> = time to reach C<sub>max</sub>, t<sub>1/2</sub>= terminal half-life

<sup>a</sup> Data presented as median (range)

### Absorption

After oral administration, fesoterodine is well absorbed. Due to rapid and extensive hydrolysis by nonspecific esterases to its active metabolite 5-hydroxymethyl tolterodine (5-HMT), fesoterodine cannot be detected in plasma. Bioavailability of the active metabolite 5-HMT is 52%. After single or multiple-dose oral administration of fesoterodine in doses from 4 mg to 28 mg, plasma concentrations of the active metabolite are proportional to the dose. Maximum plasma levels are reached after approximately 5 hours. No accumulation occurs after multiple-dose administration.

### Distribution

Plasma protein binding of the active metabolite 5-HMT is low (approximately 50%) and is bound to albumin and alpha-1-acid glycoprotein. The mean steady-state volume of distribution following intravenous infusion of the active metabolite is 169 L.

### Metabolism

After oral administration, fesoterodine is rapidly and extensively hydrolyzed to its active metabolite 5-HMT. The active metabolite is further metabolized in the liver to its carboxy, carboxy-N-desisopropyl, and N-desisopropyl metabolites via two major pathways involving CYP2D6 and CYP3A4. None of these metabolites contribute significantly to the antimuscarinic activity of fesoterodine.

Variability in CYP2D6 Metabolism: A subset of individuals (approximately 7% of Caucasians and approximately 2% of African Americans) are poor metabolizers for CYP2D6. C<sub>max</sub> and AUC of the active metabolite are increased 1.7- and 2-fold, respectively, in CYP2D6 poor metabolizers as compared to extensive metabolizers.

## Elimination

Hepatic metabolism and renal excretion contribute significantly to the elimination of the active metabolite. After oral administration of fesoterodine, approximately 70% of the administered dose was recovered in urine as the active metabolite (16%), carboxy metabolite (34%), carboxy-N-desisopropyl metabolite (18%), or N-desisopropyl metabolite (1%), and a smaller amount (7%) was recovered in feces.

The apparent terminal half-life following oral administration is approximately 7 hours.

## Daytime versus Nighttime

Sandoz Fesoterodine Fumarate may be taken during the day or at night. In a randomized, open-label, 2-period, 2-treatment crossover, single-dose study of fesoterodine 8 mg tablets in healthy subjects, the relative bioavailability of the active metabolite of fesoterodine, as measured by AUC<sub>inf</sub> ratio, was estimated to be about 93%, and the 90% CI was contained entirely within the bioequivalence limits of 80% to 125%. The C<sub>max</sub> ratio was estimated to be about 78%.

When compared to daytime dosing, the modest lowering of C<sub>max</sub> at nighttime is unlikely to be of clinical relevance for antimuscarinic efficacy.

**Effect of Food:** There is no clinically relevant effect of food on the pharmacokinetics of fesoterodine. See [4 DOSAGE AND ADMINISTRATION, 4.4 Administration](#). In a study of the effects of food on the pharmacokinetics of fesoterodine in 16 healthy male volunteers, concomitant food intake increased the active metabolite of fesoterodine AUC by approximately 19% and C<sub>max</sub> by 18%.

## Special Populations and Conditions

- **Pediatrics:** An open-label pediatric Phase 2 pharmacokinetic study with fesoterodine was conducted in 21 patients. Efficacy and safety of fesoterodine in the pediatric population have not been established. Therefore, fesoterodine should not be used in pediatric patients. See [8 ADVERSE REACTIONS, 8.2 Clinical Trial Adverse Drug Reactions, 8.2.1 Pediatrics](#).
- **Geriatrics:** The pharmacokinetics of fesoterodine was not significantly influenced by age (see [4 DOSAGE AND ADMINISTRATION](#)).
- **Sex:** The pharmacokinetics of fesoterodine was not significantly influenced by sex. Following a single 8 mg oral dose of fesoterodine, the mean (+/-SD) AUC and C<sub>max</sub> for the active metabolite of fesoterodine in 12 elderly men (mean age 67 years) were 51.8 +/- 26.1 h\*ng/mL and 3.8 +/- 1.7 ng/mL, respectively. In the same study, the mean (+/-SD) AUC and C<sub>max</sub> in 12 elderly women (mean age 68 years) were 56.0 +/- 28.8 h\*ng/mL and 4.6 +/- 2.3 ng/mL, respectively.
- **Ethnic Origin:** The pharmacokinetics of fesoterodine was not significantly influenced by ethnic origin. The effects of Caucasian or Black race on the pharmacokinetics of fesoterodine were examined in a study of 12 Caucasian and 12 Black African young male volunteers. Each subject received a single oral dose of 8 mg fesoterodine. The mean (+/- SD) AUC and C<sub>max</sub> for the active metabolite of fesoterodine in Caucasian males were 73.0 +/- 27.8 h\*ng/mL and 6.1 +/-

2.7 ng/mL, respectively. The mean( $\pm$  SD) AUC and C<sub>max</sub> in Black males were 65.8  $\pm$  23.2 h\*ng/mL and 5.5  $\pm$  1.9 ng/mL, respectively. In single- and multiple-dose studies in Japanese and Korean young male volunteers, following administration of 4 and 8 mg fesoterodine, the AUC and C<sub>max</sub> of the active metabolite of fesoterodine increased in proportion with dose, and were similar to those in Western studies.

- **Hepatic Insufficiency:** In patients with moderate (Child-Pugh B) hepatic impairment, C<sub>max</sub> and AUC of the active metabolite were increased 1.4- and 2.1-fold, respectively, as compared to healthy subjects. No dose adjustment is recommended in patients with mild or moderate hepatic impairment. Subjects with severe hepatic impairment (Child-Pugh C) have not been studied; therefore Sandoz Fesoterodine Fumarate is not recommended for use in these patients. See [4 DOSAGE AND ADMINISTRATION](#); [7 WARNINGS AND PRECAUTIONS, Hepatic/Biliary/Pancreatic](#).
- **Renal Insufficiency:** In patients with mild or moderate renal impairment (CLCR ranging from 30- 80 mL/min), C<sub>max</sub> and AUC of the active metabolite 5-HMT were increased up to 1.5- and 1.8- fold respectively, as compared to healthy subjects. In patients with severe renal impairment (CLCR < 30 mL/min), C<sub>max</sub> and AUC were increased 2.0- and 2.3-fold, respectively. In patients with mild or moderate renal impairment, no dose adjustment is required. Doses of Sandoz Fesoterodine Fumarate greater than 4 mg are not recommended in patients with severe renal impairment. See [4 DOSAGE AND ADMINISTRATION](#); [7 WARNINGS AND PRECAUTIONS, Renal](#).

## 11 STORAGE, STABILITY AND DISPOSAL

Store at controlled room temperature at 15 – 25°C. Protect from moisture.

Any unused medicinal product should be disposed of in accordance with local requirements.

## 12 SPECIAL HANDLING INSTRUCTIONS

There are no special handling instructions for Sandoz Fesoterodine Fumarate.

## PART II: SCIENTIFIC INFORMATION

### 13 PHARMACEUTICAL INFORMATION

#### Drug Substance

Proper name: Fesoterodine fumarate

Chemical name: Isobutyric acid 2-((R)-3-diisopropylammonium-1-phenylpropyl)-4-(hydroxymethyl)phenyl ester hydrogen fumarate (IUPAC)


CAS Name: Propanoic acid, 2-methyl, 2-[(1R)-3-[bis(1-methylethyl)amino]-1-phenylpropyl]-4-(hydroxymethyl)phenyl ester, (2E)-2-butenedioate (1:1) (salt)

CAS Number: 286930-03-8

Molecular formula: C<sub>30</sub>H<sub>41</sub>NO<sub>7</sub>

Molecular mass: 527.66 daltons (salt); 411.59 daltons (base)

Structural formula:



Physicochemical properties:

- Appearance: Fesoterodine fumarate is a white to off-white powder.
- Melting Point: The typical melting point is 105°C.
- pKa: Fesoterodine: pKa = 10.31 ± 0.01, at 23.4°C. Dibasic fumaric acid: pKa,1=2.94, pKa,2=4.46 at 20 °C
- pH: The pH of aqueous solutions varies between 4.0 and 3.5 in a concentration range between 0.0005 mol/L and 0.5 mol/L.

Solubility:

Solvents	Fesoterodine Fumarate [mg/mL]	Solubility Term
0.9 % NaCl solution	551	Freely soluble

Water	542	Freely soluble
Toluene	0.14	Very slightly soluble
Heptane	0.03	Practically insoluble
Acetone	205	Freely soluble
Methanol	574	Freely soluble

## 14 CLINICAL TRIALS

### 14.1 Trial Design and Study Demographics

The efficacy of fixed doses of fesoterodine 4 mg and 8 mg taken orally once daily was evaluated in two Phase 3 randomized, double-blind, placebo-controlled, 12-week studies. The co-primary endpoints were the change from Baseline to Week 12 in the average number of micturitions per 24 hours and a change from Baseline to Week 12 in the average number of UUI episodes per 24 hours (US analysis) or treatment response derived from the Treatment Benefit Scale (European analysis). Secondary endpoints included change in mean voided volume, daytime micturitions, urgency episodes per 24 hours, number of continent days per week, and change in severity of urgency episodes. A summary of the Patient Demographics for Study 1 and 2 is provided in Table 5.

**Table 5 - Summary of Patient Demographics for Study 1 and Study 2**

Study #	Study design	Dosage, route of administration and duration	Study Subjects (n)	Mean age (Range)	Sex	Ethnic origin
Study 1 (Europe, Australia, New Zealand, South Africa)	Randomized, double-blind, double-dummy, placebo- and active-controlled, parallel-arm	Fesoterodine 4 mg Fesoterodine 8 mg Placebo Tolterodine ER	N=272 N=288 N=285 N=290	57 years (19-86)	81 % F 19 % M	97 % White
Study 2 (USA)	Randomized, double-blind, placebo-controlled, parallel-arm	Fesoterodine 4 mg Fesoterodine 8 mg Placebo	N= 283 N= 279 N= 274	59 years (21-91)	76 % F 24 % M	82 % White 9 % Black 8% Other

F= Female; M=Male

### 14.2 Study Results

Fesoterodine-treated patients had statistically significant mean reductions in the number of micturitions per 24 hours and in the number of urge incontinence episodes per 24 hours at the

end of treatment compared with placebo-treated patients. Likewise, the response rate (% of patients reporting that their condition has been “greatly improved” or “improved” using a 4-point Treatment Benefit Scale) was significantly greater with fesoterodine compared with placebo. Furthermore, fesoterodine improved the mean change in the voided volume per micturition, and the mean change in the number of continent days per week. See Table 6).

**Table 6 - Mean changes from baseline to end of treatment for primary and selected secondary endpoints**

Parameter	Study 1				Study 2		
	Placebo	Feso 4 mg	Feso 8 mg	Tolterodine ER 4 mg	Placebo	Feso 4 mg	Feso 8 mg
<b>Number of micturitions per 24 hours #</b>							
	N=279	N=265	N=276	N=283	N=266	N=267	N=267
Baseline	12.0	11.6	11.9	11.5	12.2	12.9	12.0
Change from Baseline	-1.02	-1.74	-1.94	-1.69	-1.02	-1.86	-1.94
p-value vs Pbo	-	<0.001	<0.001	0.001	-	0.032	<0.001
<b>Responder rate# (treatment response)</b>							
	N=279	N=265	N=276	N=283	N=266	N=267	N=267
Responder rate	53.4%	74.7%	79.0%	72.4%	45.1%	63.7%	74.2%
p-value vs Pbo	-	<0.001	<0.001	<0.001	-	<0.001	<0.001
<b>Number of urge incontinence episodes per 24 hours #</b>							
	N=211	N=199	N=223	N=223	N=205	N=228	N=218
Baseline	3.7	3.8	3.7	3.8	3.7	3.9	3.9
Change from Baseline	-1.20	-2.06	-2.27	-1.83	-1.00	-1.77	-2.42
p-value vs Pbo	-	0.001	<0.001	0.008	-	0.003	<0.001
<b>Number of continent days per week</b>							
	N=211	N=199	N=223	N=223	N=205	N=228	N=218
Baseline	0.8	0.8	0.6	0.6	0.6	0.7	0.7
Change from Baseline	2.1	2.8	3.4	2.5	1.4	2.4	2.8
p-value vs Pbo	-	0.007	<0.001	0.139	-	<0.001	<0.001
<b>Voided volume per micturition (ml)</b>							
	N=279	N=265	N=276	N=283	N=266	N=267	N=267
Baseline	150	160	154	154	159	152	156
Change from Baseline	10	27	33	24	8	17	33
p-value vs Pbo	-	<0.001	<0.001	0.002	-	0.150	<0.001

The values are the mean change from baseline and the p-values represent the difference in LS mean vs Placebo.

# primary end points; Feso=fesoterodine


vs=versus; Pbo= placebo

In addition, sustained efficacy was shown during a 3-year open-label extension of one phase 2 and 2 phase 3 studies. Long-term treatment with fesoterodine resulted in maintained or continued improvement in all efficacy and health-related quality of life measures.

Figures 1-4: The following figures show change from baseline over time in number of micturitions and urge urinary incontinence episodes per 24 hours in the two Phase 3 studies (Study 1 and Study 2, respectively).




Figure 1 - Change in number of micturitions per 24 h (Study 1)




p-values vs Placebo: \* p< 0.001 for fesoterodine 4 mg and 8 mg; †p= 0.001 for tolterodine ER 4 mg

Figure 2 - Change in urge incontinence episodes per 24 h (Study 1)




p-values vs Placebo: \*  $p < 0.001$  for fesoterodine 4 mg and 8 mg; † $p = 0.008$  for tolterodine ER 4 mg

Figure 3 - Number of micturitions per 24 h (Study 2)



P-values vs Placebo: \*  $p < 0.001$  for fesoterodine 8 mg;  $†p = 0.032$  for fesoterodine 4 mg

Figure 4 - Change in urge incontinence episodes per 24 h (Study 2)



P-values vs Placebo: \*  $p < 0.001$  for fesoterodine 4 mg and 8 mg

### 14.3 Comparative Bioavailability Studies

#### CLINICAL TRIALS

**Fed Bioequivalence Study:** A randomized, two-way crossover, single-dose ( $1 \times 8$  mg), oral bioequivalence study was conducted to compare Sandoz Fesoterodine Fumarate (fesoterodine fumarate) 8 mg

extended-release tablets (Sandoz Canada Inc.) and Toviaz® (fesoterodine fumarate) 8 mg extended-release tablets (Pfizer Canada Inc.) in 22 healthy, adult male subjects under high-calorie, high-fat fed conditions. Plasma concentrations of the active metabolite of fesoterodine, 5-hydroxy methyl tolterodine, were measured and used to calculate the following pharmacokinetic parameters.

**SUMMARY TABLE OF THE COMPARATIVE BIOAVAILABILITY DATA**

5-Hydroxy Methyl Tolterodine (1 × 8 mg Fesoterodine Fumarate) Geometric Mean Arithmetic Mean (CV %)				
Parameter	Test <sup>a</sup>	Reference <sup>b</sup>	% Ratio of Geometric Means	90% Confidence Interval
AUC <sub>T</sub> (pg·h/mL)	77697.73 80022.26 (24.58)	74025.76 76972.48 (29.24)	104.49	98.84-110.47
AUC <sub>I</sub> (pg·h/mL)	78846.91 80981.59- (23.67)	74887.48 77760.76 (28.78)	104.87	99.18-110.89
C <sub>max</sub> (pg/mL)	6174.36 6400.89 (31.29)	6389.99 6541.12 (21.57)	96.51	85.27-109.24
T <sub>max</sub> <sup>c</sup> (h)	4.51 (1.00-7.00)	4.50 (2.00-6.00)		
T <sub>½</sub> <sup>d</sup> (h)	4.67 (20.01)	4.94 (20.47)		

<sup>a</sup> Sandoz Fesoterodine Fumarate (fesoterodine fumarate extended-release tablets) 8 mg (Sandoz Canada Inc.)

<sup>b</sup> TOVIAZ® (fesoterodine fumarate extended-release tablets) 8 mg (Pfizer Canada Inc.)

<sup>c</sup> Expressed as the median (range) only

<sup>d</sup> Expressed as the arithmetic mean (CV%) only

**Fasting Bioequivalence Study:** A randomized, two-way crossover, single-dose (1 × 8 mg), oral bioequivalence study was conducted to compare Sandoz Fesoterodine Fumarate (fesoterodine fumarate) 8 mg extended-release tablets (Sandoz Canada Inc.) and Toviaz® (fesoterodine fumarate) 8 mg extended-release tablets (Pfizer Canada Inc.) in 41 healthy, adult male subjects under fasting conditions. Plasma concentrations of the active metabolite of fesoterodine, 5-hydroxy methyl tolterodine, were measured and used to calculate the following pharmacokinetic parameters.

## SUMMARY TABLE OF THE COMPARATIVE BIOAVAILABILITY DATA

5-Hydroxy Methyl Tolterodine (1 x 8 mg Fesoterodine Fumarate) Geometric Mean Arithmetic Mean (CV %)				
Parameter	Test <sup>a</sup>	Reference <sup>b</sup>	% Ratio of Geometric Means	90 % Confidence Interval
AUCT (pg·h/mL)	82740.05 86125.94 (29.58)	77683.58 80952.35 (30.00)	106.30	103.29-109.40
AUC <sub>i</sub> (pg·h/mL)	83890.44 87244.06(29.26)	78578.16 81837.13(29.84)	106.57	103.60-109.62
C <sub>max</sub> (pg/mL)	6403.41 6672.97 (30.13)	6302.21 6555.48(29.41)	101.22	95.79-106.97
T <sub>max</sub> <sup>c</sup> (h)	6.50 (1.50-10.00)	5.00 (4.00-7.50)		
T <sub>½</sub> <sup>d</sup> (h)	5.60 (22.32)	5.44 (24.43)		

<sup>a</sup> Sandoz Fesoterodine Fumarate (fesoterodine fumarate extended-release tablets) 8 mg (Sandoz Canada Inc.)

<sup>b</sup> TOVIAZ<sup>®</sup> (fesoterodine fumarate extended-release tablets) 8 mg (Pfizer Canada Inc.)

<sup>c</sup> Expressed as the median (range) only

<sup>d</sup> Expressed as the arithmetic mean (CV%) only

### 14.4 Immunogenicity

This information is not available for this drug product.

## 15 MICROBIOLOGY

No microbiological information is required for this drug product.

## 16 NON-CLINICAL TOXICOLOGY

General Toxicology:

Toxicological studies have been performed in mice, rats, rabbits and dogs. The selection of the species is justified by *in vitro* and *in vivo* metabolism studies. The local tolerance was tested in

guinea-pigs and rabbits. In all studies, except the acute oral and intravenous toxicity studies in mice and rats and some dose-range finding studies, negative control groups have been employed. Adequate positive control substances have proven the integrity of the genotoxicity test-battery, the skin sensitization test and the immuno-toxicology study.

### **Single-Dose Toxicity:**

The NOEL and LD<sub>50</sub> for both mice and rats were 100 and ≥ 316 mg/kg following oral administration, and 10 and 31.6 mg/kg after intravenous administration of fesoterodine.

### **Repeat-Dose Toxicity:**

- Rodent

In rodents, the signs of toxicity were different in mice and rats after oral administration of fesoterodine. The NOEL was 5 mg/kg in both species after 13 weeks and in mice after 26 weeks of treatment.

- Dogs

No mortality occurred in dogs after oral treatment with 0, 0.5, 2.5 or 10 mg/kg fesoterodine for 13-weeks or 0, 0.5, 2.5 or 12.5 mg/kg fesoterodine for 9-months. No overt toxicity was noted and no fesoterodine-specific target organs could be identified. Mainly antimuscarinic effects were observed in the form of reduced lacrimal secretion leading to conjunctivitis in the high dose groups, a tightly filled gall bladder due to sphincter closure (after 9 months) and an increased heart rate starting at 2.5 mg/kg (dose dependent in females). No changes in the electrical complexes of the ECG were seen. In addition, the body weights were reduced in the males starting at 2.5 mg/kg and in the females in the high dose group in the 9-month study. Clinical hematology and biochemistry revealed increased platelet counts and urea concentration in blood in the high-dosed animals. All noted test-substance related effects were reversible after a 4-week recovery period. The NOEL was 0.5 mg/kg in these studies, and the NOAEL was 2.5 mg/kg.

### **Carcinogenicity:**

No evidence of drug-related carcinogenicity was found in 24-month studies with oral administration to mice and rats. The unbound AUC of 5-HMT at the highest tolerated dose in mice corresponded to 17 to 31 times (females) and 6 to 15 times (males) the unbound human 5-HMT AUC value (46.2 ng\*h/mL) in fed human poor CYP2D6 metabolizers reached with fesoterodine 8 mg, which is the Maximum Recommended Human Dose (MRHD). In rats, the unbound 5-HMT AUC at the highest tolerated dose corresponded to 4 to 13 times (females) and 6 to 24 times (males), the unbound human 5-HMT AUC (46.2 ng\*h/mL) in fed human poor CYP2D6 metabolizers at the MRHD.

**Genotoxicity:**

Fesoterodine was not mutagenic or genotoxic in vitro (Ames tests, chromosome aberration tests) or in vivo (mouse micronucleus test).

**Reproductive and Developmental Toxicology:**

Fesoterodine had no effect on male reproductive function or fertility in mice at doses up to 45 mg/kg/day. The maternal No-Observed-Effect Level (NOEL) and the NOEL for effects on reproduction and early embryonic development were both 15 mg/kg/day. At a dose of 45 mg/kg/day, the resulting exposures are approximately 5 to 19 times (based on lowest and highest total systemic exposure) those at the MRHD, a lower number of corpora lutea, implantation sites and viable fetuses was observed in female mice administered fesoterodine for 2 weeks prior to mating and continuing through Day 7 of gestation. Reproduction studies have shown minor embryotoxicity (increased number of resorptions, pre-implantation and post-implantation losses). At the NOEL, based on human unbound 5-HMT AUC in the fed state (46.2 ng\*h/mL), the unbound systemic exposure was 1 to 2.4 times higher in mice than in humans at the MRHD, whereas based on peak plasma concentrations, the unbound exposure in mice was 8 to 15 times higher. The Lowest-Observed-Effect Level (LOEL) for maternal toxicity was 45 mg/kg/day.

No dose-related teratogenicity was observed in reproduction studies performed in mice and rabbits. In mice treated at 7-30 times the human MRHD (45mg/kg/day, oral; based on unbound 5-HMT AUC in comparison to fed human poor metabolizers at 8 mg fesoterodine), decreased live fetuses and reduced F1 fetal body weight were observed. One fetus with cleft palate was observed at each dose (15, 45, and 75 mg/kg/day), at an incidence within the background historical range. In rabbits treated at 6 to 18 times the human MRHD (27 mg/kg/day, oral; based on unbound AUC comparison to human poor metabolizers at 8 mg fesoterodine), incompletely ossified sternbrae (retardation of bone development) were observed in fetuses. Oral administration of 30 mg/kg/day fesoterodine to mice in a pre- and post-natal development study resulted in decreased body weight of the dams and delayed ear opening of the pups. No effects were noted on mating and reproduction of the F1 dams or on the F2 offspring.

**Special Toxicology:**

This information is not available for this drug product.

**Juvenile Toxicity:**

This information is not available for this drug product.



## 17 SUPPORTING PRODUCT MONOGRAPHS

1. Pfizer Canada Inc. Product Monograph: <sup>Pf</sup>TOVIAZ<sup>®</sup> Extended-Release Tablets 4 mg and 8 mg, submission control 274307, Product Monograph Pfizer Canada Inc. Oct. 12, 2023.

## **PATIENT MEDICATION INFORMATION**

### **READ THIS FOR SAFE AND EFFECTIVE USE OF YOUR MEDICINE**

#### **Sandoz Fesoterodine Fumarate**

#### **(fesoterodine fumarate extended-release tablets)**

Read this carefully before you start taking **Sandoz Fesoterodine Fumarate** and each time you get a refill. This leaflet is a summary and will not tell you everything about this drug. Talk to your healthcare professional about your medical condition and treatment and ask if there is any new information about **Sandoz Fesoterodine Fumarate**.

#### **What is Sandoz Fesoterodine Fumarate used for?**

Sandoz Fesoterodine Fumarate is used to treat the symptoms of an overactive bladder in adults. Symptoms of an overactive bladder can include any of the following or combination of the following:

- having to urinate often
- having a strong need to urinate right away
- leaking or wetting accidents due to a strong need to urinate

#### **How does Sandoz Fesoterodine Fumarate work?**

Sandoz Fesoterodine Fumarate works by helping to relax the muscles of the bladder. This helps to relieve many of the symptoms of an overactive bladder.

#### **What are the ingredients in Sandoz Fesoterodine Fumarate?**

Medicinal ingredients: Fesoterodine fumarate

Non-medicinal ingredients: Fumaric acid, glycerol dibehenate, hypromellose (methocel K4M and K100M premium CR), hydroxypropyl cellulose, lactose monohydrate, microcrystalline cellulose, pullulan, talc. The film coating contains the following inactive ingredients: Lecithin (soya), macrogol/PEG, FD&C blue #2/ indigo carmine aluminum lake, polyvinyl alcohol-part. hydrolyzed, talc, titanium dioxide.

#### **Sandoz Fesoterodine Fumarate comes in the following dosage forms:**

- Sandoz Fesoterodine Fumarate 4 mg extended-release tablets (light blue, oval, biconvex, film-coated tablets debossed with I11 on one side and plain on other side)
- Sandoz Fesoterodine Fumarate 8 mg extended-release tablets (blue oval, biconvex, film-coated tablets debossed with I17 on one side and plain on other side)

#### **Do not use Sandoz Fesoterodine Fumarate if:**

- You are not able to completely empty your bladder (urinary retention).

- You have delayed or slow emptying of the contents in your stomach (gastric retention).
- You have an eye problem called uncontrolled-narrow angle glaucoma which leads to an increase of the pressure inside the eyeball.
- You are allergic to Sandoz Fesoterodine Fumarate or any of its ingredients. For a complete list, see [What are the ingredients in Sandoz Fesoterodine Fumarate?](#)
- You are allergic to lactose, peanuts or soya.
- You are allergic to medicines that contain tolterodine (Detrol® or Detrol® LA).

**To help avoid side effects and ensure proper use, talk to your healthcare professional before you take Sandoz Fesoterodine Fumarate. Talk about any health conditions or problems you may have, including if you:**

- Have or have had problems with your heart, such as:
  - abnormal increases in heart rate or rhythm (arrhythmia), or
  - ischemic heart disease, or
  - heart failure
- Take medicines, such as ketoconazole, itraconazole, miconazole, clarithromycin and cimetidine.
- Have stomach problems affecting passage and digestion of food.
- Have severe constipation.
- Have gastrointestinal obstruction disorders (e.g. pyloric stenosis).
- Have problems emptying your bladder or if you have weak urine stream.
- Have liver problems.
- Have a rare hereditary problem of galactose intolerance, the Lapp lactase deficiency or glucose- galactose malabsorption. Sandoz Fesoterodine Fumarate contains lactose.
- Have myasthenia gravis (a chronic autoimmune neuromuscular disease which causes muscle weakness).
- Are receiving treatment for an eye problem called narrow-angle glaucoma.
- Have kidney problems.
- Are pregnant, or trying to become pregnant.
- Are breastfeeding your child.
- Have a family history of soya intolerance. Sandoz Fesoterodine Fumarate contains soya lecithin.

**Other warnings you should know about:**

**Angioedema and Allergic Reactions:** Sandoz Fesoterodine Fumarate can cause:

- angioedema (swelling of face or tongue, difficulty breathing) and
- severe allergic reactions (hives, trouble breathing, abdominal cramps, rapid heartbeat and feeling faint).

If you experience any of these symptoms, stop taking Sandoz Fesoterodine Fumarate and see your healthcare professional **right away**.

**Driving and Operating Machinery:** Until you know how Sandoz Fesoterodine Fumarate affects you, do not drive or operate a vehicle or potentially dangerous machinery, especially when you first start treatment, or when your dose is changed. Taking Sandoz Fesoterodine Fumarate can cause

side effects such as:

- blurred vision,
- dizziness, and
- drowsiness.

**Tell your healthcare professional about all the medicines you take, including any drugs, vitamins, minerals, natural supplements or alternative medicines.**

The following **may interact with Sandoz Fesoterodine Fumarate:**

- Other drugs similar to Sandoz Fesoterodine Fumarate (antimuscarinic/anticholinergic)
- Drugs to treat fungal infections (such as, fluconazole, ketoconazole, miconazole or itraconazole)
- Drugs to treat ulcers or gastroesophageal reflux disease (GERD) (such as cimetidine)
- Drugs to treat bacterial infections (such as erythromycin, clarithromycin)
- Drugs to treat some types of cancer (such as vinblastine)
- Drugs to treat depression
- Antipsychotics (drugs to stabilize thinking and behavior)

**How to take Sandoz Fesoterodine Fumarate:**

- Take Sandoz Fesoterodine Fumarate as instructed by your healthcare professional. Do not increase, decrease or stop taking Sandoz Fesoterodine Fumarate without first talking to your healthcare professional.
- Take Sandoz Fesoterodine Fumarate with liquid and swallow the tablet **whole**. Sandoz Fesoterodine Fumarate should not be chewed, divided, or crushed.

REMEMBER: This medication is for YOU. Never give it to others. It may harm them even if their symptoms are the same as yours.

**Usual Adult dose:**

- The usual starting dose is 4 mg once daily.
- The maximum daily dose for Sandoz Fesoterodine Fumarate is 8 mg.
- Your healthcare professional may give you the lower 4 mg dose of Sandoz Fesoterodine Fumarate if you have certain medical conditions.
- Based on how you respond to Sandoz Fesoterodine Fumarate upon your response and tolerability, your healthcare professional may increase your dose to 8 mg once daily.

**Overdose:**

Some of the signs of an overdose could be:

- Dry mouth
- Constipation
- Dry eyes
- Seeing or believing things that are not there (hallucinations)
- Feeling excited

- Convulsions
- Trouble breathing
- Fast heart rate
- Dilated pupils
- Problems emptying your bladder

If you think you, or a person you are caring for, have taken too much Sandoz Fesoterodine Fumarate, contact a healthcare professional, hospital emergency department, or regional poison control centre immediately, even if there are no symptoms.

### **Missed Dose:**

If you miss taking your dose, take it as soon as you remember. But if it is almost time for you to take the next dose, skip the missed dose and just take it at your usual time. Do not take two doses at the same time to make up for the dose that you missed.

### **What are possible side effects from using Sandoz Fesoterodine Fumarate?**

These are not all the possible side effects you may have when taking Sandoz Fesoterodine Fumarate. If you experience any side effects not listed here, tell your healthcare professional.

A very common side effect is dry mouth. If you experience dry mouth after taking Sandoz Fesoterodine Fumarate, there are a few ways that might help relieve the symptoms:

- Carry a bottle of water and sip a little bit throughout the day or suck on ice chips to provide moisture
- Chew sugarless gum or suck on sugarless hard candy to stimulate saliva production
- Avoid eating salty or spicy food
- Avoid drinking carbonated, caffeinated and alcoholic beverages
- Avoid using mouth rinses that contain alcohol, as they may dry out the mouth
- Use a humidifier at night
- Ask your healthcare professional to recommend an over-the-counter saliva substitute or oral lubricant

Common side effects are:

- Constipation
- Upset stomach
- Nausea
- Urinary tract infection
- Upper respiratory tract infection
- Dry eyes
- Painful urination
- Difficulty emptying your bladder
- Dry throat
- Swelling of your lower legs and hands
- Skin rash

- Blurred vision
- Dizziness
- Headache

Uncommon side effects are:

- Cough
- Back pain
- Abdominal pain
- Trouble sleeping
- Drowsiness

Other side effects that have been reported after the drug was marketed include:

- blurred vision
- dizziness, headache
- heart palpitations
- hypersensitivity reactions
- hives, and itching
- throat numbness, swelling and soreness
- reduced sense of touch, temperature, and pain (numbness).

Check with your healthcare professional or pharmacist right away if you have any bothersome or unusual effects while taking Sandoz Fesoterodine Fumarate.

Serious side effects and what to do about them			
Symptom / effect	Talk to your healthcare professional		Stop taking drug and get immediate medical help
	Only if severe	In all cases	
<b>UNCOMMON</b>			
<b>Severe allergic reactions:</b> sudden wheeziness and chest pain or tightness; or swelling of eyelids, face, lips, tongue or throat			✓
<b>Urinary retention</b> (inability to pass urine or to empty your bladder): pain			✓
<b>Angioedema</b> (swelling of tissue under the skin): difficulty breathing; swollen face, hands and feet, genitals, tongue, throat; Swelling of the digestive tract causing diarrhea, nausea or vomiting			✓

If you have a troublesome symptom or side effect that is not listed here or becomes bad enough to interfere with your daily activities, tell your healthcare professional.

### **Reporting Side Effects**

You can report any suspected side effects associated with the use of health products to Health Canada by:

- Visiting the Web page on Adverse Reaction Reporting (<https://www.canada.ca/en/health-canada/services/drugs-health-products/medeffect-canada/adverse-reaction-reporting.html>) for information on how to report online, by mail or by fax; or
- Calling toll-free at 1-866-234-2345.

*NOTE: Contact your healthcare professional if you need information about how to manage your side effects. The Canada Vigilance Program does not provide medical advice.*

### **Storage:**

Store at controlled room temperature at 15 – 25°C. Protect from moisture.

You should not use your medication after the expiration date printed on the carton and label.

Keep out of reach and sight of children.

### **If you want more information about Sandoz Fesoterodine Fumarate:**

- Talk to your healthcare professional
- Find the full product monograph that is prepared for healthcare professionals and includes this Patient Medication Information by visiting the Health Canada website: <https://www.canada.ca/en/health-canada/services/drugs-health-products/drug-products/drug-product-database.html>; the manufacturer's website [www.sandoz.ca], or by calling 1-800-361-3062.

This leaflet was prepared by Sandoz Canada Inc.

Last Revised:

APR 26, 2024