

PRODUCT MONOGRAPH  
INCLUDING PATIENT MEDICATION INFORMATION

**PrLUMAKRAS®**  
sotorasib tablets  
tablets, 120 mg, 240 mg, oral  
Antineoplastic Agent  
ATC Code: L01XX73

“LUMAKRAS, indicated for:

- the treatment of adult patients with Kirsten rat sarcoma viral oncogene homolog (*KRAS*) *G12C*-mutated locally advanced (not amenable to curative therapy) or metastatic non-small cell lung cancer (NSCLC) who have received at least one prior systemic therapy

has been issued market authorization with conditions, pending the results of trials to verify its clinical benefit. Patients should be advised of the nature of the authorization. For further information for LUMAKRAS please refer to Health Canada’s Notice of Compliance with conditions - drug products web site: <https://www.canada.ca/en/health-canada/services/drugs-health-products/drug-products/notice-compliance/conditions.html>”

Amgen Canada Inc.  
6775 Financial Drive, Suite 300  
Mississauga, Ontario  
L5N 0A4

Date of Initial Authorization:  
SEP 09, 2021

Date of Revision:  
OCT 28, 2024

Submission Control Number: 276210

© 2021-2024 Amgen Canada Inc., All rights reserved.

**This product has been authorized under the Notice of Compliance with Conditions (NOC/c) for one or all of its indicated uses.**

### **What is a Notice of Compliance with Conditions (NOC/c)?**

An NOC/c is a form of market approval granted to a product on the basis of promising evidence of clinical effectiveness following review of the submission by Health Canada.

Products authorized under Health Canada's NOC/c policy are intended for the treatment, prevention or diagnosis of a serious, life-threatening or severely debilitating illness. They have demonstrated promising benefit, are of high quality and possess an acceptable safety profile based on a benefit/risk assessment. In addition, they either respond to a serious unmet medical need in Canada or have demonstrated a significant improvement in the benefit/risk profile over existing therapies. Health Canada has provided access to this product on the condition that sponsors carry out additional clinical trials to verify the anticipated benefit within an agreed upon time frame.

### **What will be different about this Product Monograph?**

The following Product Monograph will contain boxed text at the beginning of each major section clearly stating the nature of the market authorization. Sections for which NOC/c status holds particular significance will be identified in the left margin by the symbol (NOC/c). These sections may include, but are not limited to, the following:

- Indications;
- Clinical Pharmacology;
- Warnings and Precautions;
- Adverse Reactions;
- Dosage and Administration; and
- Clinical Trials.

### **Adverse Reaction Reporting and Re-Issuance of the Product Monograph**

Health care providers are encouraged to report Adverse Reactions associated with normal use of these and all drug products to Health Canada's Canada Vigilance Program at 1-866-234-2345. The Product Monograph will be re-issued in the event of serious safety concerns previously unidentified or at such time as the sponsor provides the additional data in support of the product's clinical benefit. Once the latter has occurred, and in accordance with the NOC/c policy, the conditions associated with market authorization will be removed.

**RECENT MAJOR LABEL CHANGES**

4.2 Recommended Dose and Dosage Adjustment	10 / 2024
4.4 Administration	10 / 2023
7 WARNINGS AND PRECAUTIONS	8 / 2024

**TABLE OF CONTENTS**

Sections or subsections that are not applicable at the time of authorization are not listed.

<b>RECENT MAJOR LABEL CHANGES</b> .....	<b>3</b>
<b>TABLE OF CONTENTS</b> .....	<b>3</b>
<b>PART I: HEALTH PROFESSIONAL INFORMATION</b> .....	<b>5</b>
<b>1 INDICATIONS</b> .....	<b>5</b>
1.1 Pediatrics .....	5
1.2 Geriatrics .....	5
<b>2 CONTRAINDICATIONS</b> .....	<b>5</b>
<b>3 SERIOUS WARNINGS AND PRECAUTIONS BOX</b> .....	<b>5</b>
<b>4 DOSAGE AND ADMINISTRATION</b> .....	<b>5</b>
4.1 Dosing Considerations .....	5
4.2 Recommended Dose and Dosage Adjustment .....	6
4.3 Reconstitution .....	7
4.4 Administration .....	7
4.5 Missed Dose .....	8
<b>5 OVERDOSAGE</b> .....	<b>8</b>
<b>6 DOSAGE FORMS, STRENGTHS, COMPOSITION AND PACKAGING</b> .....	<b>8</b>
<b>7 WARNINGS AND PRECAUTIONS</b> .....	<b>9</b>
7.1 Special Populations .....	10
7.1.1 Pregnant Women .....	10
7.1.2 Breast-feeding .....	10
7.1.3 Pediatrics .....	10
7.1.4 Geriatrics .....	10
<b>8 ADVERSE REACTIONS</b> .....	<b>10</b>
8.1 Adverse Reaction Overview .....	10
8.2 Clinical Trial Adverse Reactions .....	11
8.2.1 Clinical Trial Adverse Reactions – Pediatrics .....	13

8.3	Less Common Clinical Trial Adverse Reactions .....	13
8.3.1	Less Common Clinical Trial Adverse Reactions – Pediatrics .....	13
8.4	Abnormal Laboratory Findings: Hematologic, Clinical Chemistry and Other Quantitative Data .....	13
<b>9</b>	<b>DRUG INTERACTIONS.....</b>	<b>14</b>
9.1	Serious Drug Interactions .....	14
9.2	Drug Interactions Overview .....	14
9.3	Drug-Behavioural Interactions .....	14
9.4	Drug-Drug Interactions .....	14
9.5	Drug-Food Interactions.....	15
9.6	Drug-Herb Interactions .....	15
9.7	Drug-Laboratory Test Interactions.....	15
<b>10</b>	<b>CLINICAL PHARMACOLOGY .....</b>	<b>15</b>
10.1	Mechanism of Action .....	15
10.2	Pharmacodynamics .....	16
10.3	Pharmacokinetics .....	16
<b>11</b>	<b>STORAGE, STABILITY AND DISPOSAL.....</b>	<b>18</b>
<b>12</b>	<b>SPECIAL HANDLING INSTRUCTIONS.....</b>	<b>18</b>
<b>PART II: SCIENTIFIC INFORMATION.....</b>		<b>19</b>
<b>13</b>	<b>PHARMACEUTICAL INFORMATION.....</b>	<b>19</b>
<b>14</b>	<b>CLINICAL TRIALS .....</b>	<b>20</b>
14.1	Trial Design and Study Demographics .....	20
14.2	Study Results .....	21
<b>15</b>	<b>MICROBIOLOGY.....</b>	<b>21</b>
<b>16</b>	<b>NON-CLINICAL TOXICOLOGY .....</b>	<b>21</b>
<b>PATIENT MEDICATION INFORMATION .....</b>		<b>23</b>

“LUMAKRAS, indicated for:

- the treatment of adult patients with Kirsten rat sarcoma viral oncogene homolog (*KRAS*) *G12C*-mutated locally advanced (not amenable to curative therapy) or metastatic non-small cell lung cancer (NSCLC) who have received at least one prior systemic therapy

has been issued market authorization with conditions, pending the results of trials to verify its clinical benefit. Patients should be advised of the nature of the authorization. For further information for LUMAKRAS please refer to Health Canada’s Notice of Compliance with conditions - drug products web site: <https://www.canada.ca/en/health-canada/services/drugs-health-products/drug-products/notice-compliance/conditions.html>”

## PART I: HEALTH PROFESSIONAL INFORMATION

### 1 INDICATIONS

**NOC/c** LUMAKRAS® (sotorasib) is indicated for:

- the treatment of adult patients with Kirsten rat sarcoma viral oncogene homolog (*KRAS*) *G12C*-mutated locally advanced (not amenable to curative therapy) or metastatic non-small cell lung cancer (NSCLC) who have received at least one prior systemic therapy.

This indication is issued market authorization with conditions based on objective response rate (ORR) and duration of response (DOR) (see 14 CLINICAL TRIALS). Continued approval for this indication may be contingent upon verification and description of clinical benefit in a confirmatory trial(s).

#### 1.1 Pediatrics

**Pediatrics (< 18 years of age):** No data are available to Health Canada; therefore, Health Canada has not authorized an indication for pediatric use.

#### 1.2 Geriatrics

**Geriatrics (≥ 65 years of age):** In clinical studies, no overall differences in safety or efficacy were observed between geriatric patients (≥ 65 years old) and younger patients.

### **NOC/c** 2 CONTRAINDICATIONS

LUMAKRAS is contraindicated in patients who are hypersensitive to this drug or to any ingredient in the formulation, including any non-medicinal ingredient, or component of the container. For a complete listing, see 6 DOSAGE FORMS, STRENGTHS, COMPOSITION AND PACKAGING.

### 3 SERIOUS WARNINGS AND PRECAUTIONS BOX

At the time of authorization, no serious warning or precaution had been identified.

### **NOC/c** 4 DOSAGE AND ADMINISTRATION

#### 4.1 Dosing Considerations

Confirm presence of *KRAS G12C* mutation using a validated test prior to initiation of LUMAKRAS.

## 4.2 Recommended Dose and Dosage Adjustment

The recommended dosage of LUMAKRAS is 960 mg (eight 120 mg tablets or four 240 mg tablets) orally once daily until disease progression or unacceptable toxicity.

### Dose Modifications

LUMAKRAS dose reduction levels are summarized in Table 1. Dosage modifications for adverse reactions are provided in Table 2.

If adverse reactions occur, a maximum of two dose reductions are permitted. Discontinue LUMAKRAS if patients are unable to tolerate the minimum dose of 240 mg once daily.

**Table 1. Recommended Dose Reduction Levels for LUMAKRAS**

Dose Reduction Level	Dose
First dose reduction	480 mg (4 x 120 mg tablets or 2 x 240 mg tablets) once daily
Second dose reduction	240 mg (2 x 120 mg tablets or 1 x 240 mg tablet) once daily

**Table 2. Recommended Dose Modifications for LUMAKRAS**

Adverse Reaction	Severity <sup>a</sup>	Dosage Modification
Hepatotoxicity (see 7 WARNINGS AND PRECAUTIONS)	Grade 2 AST or ALT with symptoms or Grade 3 or 4 AST or ALT	<ul style="list-style-type: none"> <li>Withhold LUMAKRAS until recovery to ≤ Grade 1 or baseline.</li> <li>Resume LUMAKRAS at the next lower dose level.</li> </ul>
	AST or ALT > 3 x ULN with total bilirubin > 2 x ULN in the absence of alternative causes	<ul style="list-style-type: none"> <li>Permanently discontinue LUMAKRAS.</li> </ul>
Interstitial Lung Disease (ILD)/ pneumonitis (see 7 WARNINGS AND PRECAUTIONS)	Any Grade	<ul style="list-style-type: none"> <li>Stop LUMAKRAS if ILD/pneumonitis is suspected</li> <li>Permanently discontinue LUMAKRAS if ILD/pneumonitis is confirmed</li> </ul>
Nausea or vomiting despite appropriate supportive care (including anti-emetic therapy)	Grade 3 to 4	<ul style="list-style-type: none"> <li>Withhold LUMAKRAS until recovery to ≤ Grade 1 or baseline</li> <li>Resume LUMAKRAS at the next lower dose level.</li> </ul>
Diarrhea despite appropriate supportive care (including anti-diarrheal therapy)	Grade 3 to 4	<ul style="list-style-type: none"> <li>Withhold LUMAKRAS until recovery to ≤ Grade 1 or baseline.</li> <li>Resume LUMAKRAS at the next lower dose level.</li> </ul>
Other adverse reactions	Grade 3 to 4	<ul style="list-style-type: none"> <li>Withhold LUMAKRAS until recovery to ≤ Grade 1 or baseline.</li> <li>Resume LUMAKRAS at the next lower dose level.</li> </ul>

ALT = alanine aminotransferase; AST = aspartate aminotransferase; ULN = upper limit of normal

<sup>a</sup> Grading defined by National Cancer Institute Common Terminology Criteria for Adverse Events (NCI CTCAE) version 5.0

## Hepatic Impairment

No dose adjustment is recommended for patients with mild (Child-Pugh A) hepatic impairment. There are limited data on daily LUMAKRAS dosing in subjects with moderate or severe hepatic impairment (Child-Pugh B or C). Use LUMAKRAS in patients with moderate or severe hepatic impairment only if the benefits outweigh the risks (see 10.3 Pharmacokinetics).

Monitor LUMAKRAS related toxicities in patients with hepatic impairment more frequently since these patients may be at increased risk for adverse reactions including hepatotoxicity.

## Renal Impairment

Based on population pharmacokinetic analysis, no dose adjustment is recommended for patients with mild and moderate renal impairment (estimated glomerular filtration rate (eGFR) as determined by Modification of Diet in Renal Disease formula (MDRD):  $\geq 45$  mL/min/1.73m<sup>2</sup>). However, since data were limited in patients with moderate renal impairment, caution should be exercised when treating these patients (see 10.3 Pharmacokinetics, Special populations).

LUMAKRAS has not been studied in patients with severe renal impairment (eGFR as determined by MDRD:  $< 30$  mL/min/1.73m<sup>2</sup>).

### 4.3 Reconstitution

Not applicable.

### 4.4 Administration

Take LUMAKRAS at the same time each day with or without food. Swallow tablets whole. Do not chew, crush, or split tablets.

#### Administration to Patients Who Have Difficulty Swallowing Solids

Disperse tablets in 120 mL (4 ounces) of non-carbonated, room temperature water without crushing. Do not use other liquids. Stir until tablets are dispersed into small pieces (the tablets will not completely dissolve) and drink immediately. The appearance of the mixture may range from pale to bright yellow. Rinse the container with an additional 120 mL (4 ounces) of water and drink immediately. If the mixture is not consumed immediately, stir the mixture again to ensure that tablets are dispersed. Consume within two hours of preparation.

If administration through a nasogastric (NG) tube (8 Fr and above) or percutaneous endoscopic gastrostomy (PEG) tube (24 Fr and above) is required, follow the process described above for the initial dispersion. Draw up the mixture into a catheter tip disposable syringe and immediately inject the mixture through the feeding tube. Rinse the container with extra 120 mL of water and flush it down the feeding tube. The resulting dispersed suspension and rinse should be administered as per the NG or PEG tube manufacturer's instructions with appropriate water flushes. Silicone and polyurethane feeding tubes can be used. Administer within two hours of preparation, stored at room temperature.

#### Coadministration of LUMAKRAS with Acid-Reducing Agents

Avoid coadministration of proton pump inhibitors (PPIs) and H<sub>2</sub> receptor antagonists with LUMAKRAS. If treatment with an acid-reducing agent cannot be avoided, take LUMAKRAS 4 hours before or 10 hours after administration of a local antacid (see 9.4 Drug-Drug Interactions and 10.3 Pharmacokinetics).

#### 4.5 Missed Dose

If a dose of LUMAKRAS is missed by greater than 6 hours, resume treatment as prescribed the next day.

If vomiting occurs after taking LUMAKRAS, do not take an additional dose. Resume treatment as prescribed the next day.

#### 5 OVERDOSAGE

There is no clinical experience with overdose with LUMAKRAS. In the event of an overdose, treat the patient symptomatically and institute supportive measures as required.

For management of a suspected drug overdose, contact your regional poison control centre.

#### 6 DOSAGE FORMS, STRENGTHS, COMPOSITION AND PACKAGING

**Table 3. Dosage Forms, Strengths, Composition and Packaging**

Route of Administration	Dosage Form / Strength/Composition	Non-medicinal Ingredients
oral	sotorasib tablet, 120 mg	croscarmellose sodium, iron oxide yellow, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, talc, titanium dioxide
oral	sotorasib tablet, 240 mg	croscarmellose sodium, iron oxide yellow, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, talc, titanium dioxide

#### Product Characteristics:

##### 120 mg Tablet

Yellow, oblong-shaped, immediate release, film-coated tablet, debossed with “AMG” on one side and “120” on the opposite side.

- HDPE bottles with child-resistant polypropylene closures and aluminum foil induction seal liners containing 120 film-coated tablets.
- HDPE bottle with child-resistant polypropylene closures and aluminum foil induction seal liner containing 240 film-coated tablets.
- PVC/Aclar blisters with aluminum foil backing in each carton. Each blister card contains 8 film-coated tablets. Carton contains 30 blister cards for a total of 240 film-coated tablets.
- PVC/PVDC blisters with aluminum foil backing in each carton. Each blister card contains 8 film-coated tablets. Carton contains 30 blister cards for a total of 240 film-coated tablets.

##### 240 mg Tablet

Yellow, oval-shaped, immediate release, film-coated tablet, debossed with “AMG” on one side and “240” on the opposite side.



- HDPE bottles with child-resistant polypropylene closures and aluminum foil induction seal liners containing 30 film-coated tablets.
- PVC/Aclar blisters with aluminum foil backing in each carton. Each blister card contains 10 film-coated tablets. Carton contains 3 blister cards for a total of 30 film-coated tablets.

## NOC/c 7 WARNINGS AND PRECAUTIONS

### Hepatic/Biliary/Pancreatic

#### Hepatotoxicity

LUMAKRAS can cause hepatotoxicity, which may lead to drug-induced liver injury and hepatitis. Among 405 patients who received 960 mg once daily (QD) LUMAKRAS in CodeBreak 100, 18.5% reported events associated with hepatotoxicity. Drug-induced liver injury and hepatitis occurred in 1.5% (all grades) and 1.0% (Grade 3) of the population.

Serious cases of increased aspartate aminotransferase (AST) / increased alanine aminotransferase (ALT) can occur (see 8.1 Adverse Reaction Overview). A total of 17% of patients who received LUMAKRAS had increased AST/ALT, 5.2% were Grade 3 and 0.5% were Grade 4. Increased AST/ALT leading to dose interruption and/or reduction occurred in 6.7% of patients. LUMAKRAS was discontinued due to AST/ALT increases in 1.7% of patients. The median time to first onset for increased AST/ALT was 44 days (range: 2 days to 722).

Monitor liver function tests (ALT, AST and total bilirubin) prior to the start of LUMAKRAS, every 3 weeks for the first 3 months, then once a month or as clinically indicated, with more frequent testing in patients who develop transaminase and/or bilirubin elevations. Withhold, reduce dose or permanently discontinue LUMAKRAS based on severity of adverse reaction (see 4.2 Recommended Dose and Dosage Adjustment).

### Respiratory

#### Interstitial Lung Disease (ILD)/Pneumonitis

Severe or life-threatening ILD/pneumonitis can occur in patients treated with LUMAKRAS with prior exposure to immunotherapy or radiotherapy. Among 405 patients who received 960 mg QD LUMAKRAS in CodeBreak 100, ILD/pneumonitis occurred in 1.2% of patients (see 8.1 Adverse Reaction Overview), most cases were Grade 3 or 4 and serious. The median time to first onset for ILD/pneumonitis was 126 days (range: 15 to 704 days). LUMAKRAS was discontinued due to ILD/pneumonitis in 0.7% of patients. ILD/pneumonitis leading to dose interruption and/or reduction occurred in 0.2% of the patients. Monitor patients for new or worsening pulmonary symptoms indicative of ILD/pneumonitis (e.g. dyspnea, cough, fever). Immediately withhold LUMAKRAS in patients with suspected ILD/pneumonitis and permanently discontinue LUMAKRAS if no other potential causes of ILD/pneumonitis are identified (see 4.2 Recommended Dose and Dosage Adjustment).

### Reproductive Health: Female and Male Potential

Refer to Section 7.1 Special Populations, 7.1.1 Pregnant Women

#### • Fertility

There are no clinical studies to evaluate the effect of LUMAKRAS on fertility. (see 16 NON-CLINICAL TOXICOLOGY, Impairment of Fertility).

## 7.1 Special Populations

### 7.1.1 Pregnant Women

There are no clinical studies with LUMAKRAS use in pregnant women.

In rat and rabbit embryo-fetal development studies, oral sotorasib did not affect embryo survival at exposures up to 4 times the human exposure at the 960 mg clinical dose. However, there were slight decreases in fetal body weights, and the number of ossified metacarpals in fetuses at approximately 2.3 times the human exposure, based on AUC, at the clinical dose of 960 mg, which was associated with maternal toxicity (see 16 NON-CLINICAL TOXICOLOGY, Reproductive and Developmental Toxicology).

Inform the patient of the potential hazards to the fetus if LUMAKRAS is used during pregnancy, or if the patient becomes pregnant while taking LUMAKRAS.

### 7.1.2 Breast-feeding

There are no clinical studies on the presence of LUMAKRAS or its metabolites in human milk, the effects on the breastfed child or on milk production. Because of the potential risk for LUMAKRAS to cause adverse effects in breastfed children, advise women not to breastfeed during treatment with LUMAKRAS and for 1 week after the final dose.

### 7.1.3 Pediatrics

**Pediatrics (< 18 years of age):** Safety and effectiveness of LUMAKRAS in pediatric patients have not been established. No data are available to Health Canada; therefore, Health Canada has not authorized an indication for pediatric use.

### 7.1.4 Geriatrics

Of the 405 patients in CodeBreak 100 who received the recommended dose of LUMAKRAS at 960 mg once daily, 47.2% were 65 years or older. In clinical studies, no overall differences in safety or efficacy were observed in comparison with younger patients. No dose adjustment is required for geriatric patients.

## NOC/c 8 ADVERSE REACTIONS

### 8.1 Adverse Reaction Overview

The data described in the 7 WARNINGS AND PRECAUTIONS section reflect the safety of LUMAKRAS in 405 patients with any tumour type who received 960 mg orally once daily as monotherapy in a multicenter, open-label, single-arm trial (CodeBreak 100), 37.0% were exposed for 6 months or longer and 18.8% were exposed for greater than one year.

#### Non-Small Cell Lung Cancer

The data described below reflect the safety of LUMAKRAS in 214 patients with *KRAS G12C*-mutated locally advanced or metastatic NSCLC who received 960 mg orally once daily in a multicenter, open-label, single-arm trial (CodeBreak 100). This information excludes any event with disease progression-related preferred terms. The median duration of exposure to LUMAKRAS was 5.6 months (range: 0.2 to 43.2 months) with 48.6% exposed for at least 6 months, 36.0% exposed for at least 9 months, and 28% exposed for at least 1 year. The patient and disease characteristics were: median age of 66 years (range 37 to 86) with 57% ≥ 65 years and 12% ≥ 75 years; 55% female; 79% White, 15% Asian, 3% Black; 29% Eastern

Cooperative Oncology Group Performance Status (ECOG PS) 0, 71% ECOG PS 1, 1% ECOG PS  $\geq$  2.

Serious adverse events occurred in 56.1% of patients treated with LUMAKRAS. Serious adverse reactions in  $\geq$  2% of patients were pneumonia (9.3%), musculoskeletal pain (5.1%), hepatotoxicity (3.3%), and dyspnea (3.3%).

Fatal adverse events, excluding disease progression, occurred in 8.4% of the population. Fatal adverse reactions due to pneumonia occurred in 0.9% of patients who received LUMAKRAS.

Permanent discontinuation due to an adverse event occurred in 10.3% of patients who received LUMAKRAS. Adverse reactions resulting in permanent discontinuation of LUMAKRAS included hepatotoxicity (4.2%), pneumonitis (1.4%), increased blood ALP (0.9%), pneumonia (0.5%), dyspnea (0.5%), diarrhea (0.5%) and vomiting (0.5%).

The most common adverse reactions ( $\geq$  20%) were diarrhea, musculoskeletal pain, fatigue, nausea, cough, arthralgia, hepatotoxicity, and dyspnea. The most common Grade 3-4 laboratory abnormalities ( $\geq$ 5%) were ALT increase (11%) and AST increase (9%).

Dosage interruptions and/or reduction due to an adverse event occurred in 38.3% of patients who received LUMAKRAS. The most frequent adverse reactions that resulted in dosage interruption and/or reduction ( $\geq$ 2%) were hepatotoxicity (10.7%), diarrhea (9.8%), pneumonia (3.7%), blood ALP increased (3.3%), nausea (3.3%), musculoskeletal pain (2.8%) and fatigue (2.8%).

Table 4 summarizes the common adverse reactions observed in CodeBreak 100.

## 8.2 Clinical Trial Adverse Reactions

Clinical trials are conducted under very specific conditions. The adverse reaction rates observed in the clinical trials; therefore, may not reflect the rates observed in practice and should not be compared to the rates in the clinical trials of another drug. Adverse reaction information from clinical trials may be useful in identifying and approximating rates of adverse drug reactions in real-world use.

The safety of LUMAKRAS was evaluated in 214 patients with *KRAS G12C*-mutated locally advanced or metastatic NSCLC who received 960 mg orally once daily as monotherapy.

Adverse reactions reported in LUMAKRAS clinical studies are displayed below.

**Table 4. Adverse Reactions Occurring in  $\geq$  10% (Any grades) or  $\geq$  2% (Grade 3 or 4) of NSCLC Patients Who Received LUMAKRAS in CodeBreak 100, Excluding Preferred Terms Related to Disease-progression**

	LUMAKRAS N = 214	
	All Grades (%)	Grade 3 or 4 (%)
<b>Gastrointestinal disorders</b>		
Diarrhea	46	5.6
Nausea	30	0.9
Vomiting	19	1.4

**Table 4. Adverse Reactions Occurring in  $\geq 10\%$  (Any grades) or  $\geq 2\%$  (Grade 3 or 4) of NSCLC Patients Who Received LUMAKRAS in CodeBreak 100, Excluding Preferred Terms Related to Disease-progression**

	LUMAKRAS N = 214	
	All Grades (%)	Grade 3 or 4 (%)
Constipation	20	0.5
Abdominal Pain <sup>a</sup>	18	1.4
<b>Hepatobiliary disorders</b>		
Hepatotoxicity <sup>b</sup>	24	11.2
<b>Respiratory</b>		
Cough <sup>c</sup>	28	1.9
Dyspnea <sup>d</sup>	22	4.2
Pulmonary Embolism	3	2.8
<b>Musculoskeletal Disorders</b>		
Musculoskeletal Pain <sup>e</sup>	39	8.9
Arthralgia	24	2.8
<b>General disorders and administration site conditions</b>		
Fatigue <sup>f</sup>	31	3.3
Edema <sup>g</sup>	16	0.5
<b>Metabolism and nutrition disorders</b>		
Decreased appetite	15	0.9
<b>Infections</b>		
Pneumonia <sup>h</sup>	16	8.9
<b>Skin and Subcutaneous Tissue Disorders</b>		
Rash <sup>i</sup>	15	0

<sup>a</sup> Abdominal pain includes abdominal pain, abdominal pain upper, abdominal pain lower

<sup>b</sup> Hepatotoxicity includes alanine aminotransferase increased, aspartate aminotransferase increased, blood bilirubin increased, drug-induced liver injury, hepatitis, transaminases abnormal, transaminases increased, hypertransaminases.

<sup>c</sup> Cough includes cough, productive cough, and upper-airway cough

<sup>d</sup> Dyspnea includes dyspnea and dyspnea exertional

<sup>e</sup> Musculoskeletal pain includes back pain, bone pain, musculoskeletal chest pain, neck pain, musculoskeletal pain, myalgia, bone pain, musculoskeletal stiffness, spinal pain, non-cardiac chest pain and pain in extremity

<sup>f</sup> Fatigue includes fatigue and asthenia

<sup>g</sup> Edema includes edema peripheral, edema, edema generalized, localized edema

<sup>h</sup> Pneumonia includes pneumonia, pneumonia bacterial, pneumonia influenza, pneumonia staphylococcal, pneumonia aspiration

<sup>i</sup> Rash includes dermatitis, dermatitis acneiform, rash, rash maculo-papular, rash pustular

### 8.2.1 Clinical Trial Adverse Reactions – Pediatrics

Trials were not conducted in pediatric populations.

### 8.3 Less Common Clinical Trial Adverse Reactions

Respiratory: Pneumonitis (1.9%)

#### 8.3.1 Less Common Clinical Trial Adverse Reactions – Pediatrics

Trials were not conducted in pediatric populations.

### 8.4 Abnormal Laboratory Findings: Hematologic, Clinical Chemistry and Other Quantitative Data

**Table 5. Laboratory Abnormalities Worsening from Baseline in  $\geq 20\%$  of Patients in CodeBreak 100<sup>a</sup>**

Laboratory Abnormality	LUMAKRAS NSCLC, 960 mg QD N = 214	
	Any Worsening Change from Baseline n (%)	Worsening $\geq 3$ Grades from Baseline n (%)
Lymphocytes - Decrease	115 (53.7)	9 (4.2)
Hemoglobin - Decrease	96 (44.9)	1 (0.5)
Calcium (Corrected) - Decrease	90 (42.1)	2 (0.9)
Alanine Amino Transferase - Increase	84 (39.3)	23 (10.7)
Aspartate Amino Transferase - Increase	84 (39.3)	19 (8.9)
Alkaline Phosphatase - Increase	79 (36.9)	5 (2.3)
Urine Protein - Increase	72 (33.6)	10 (4.7)
Sodium - Decrease <sup>b</sup>	71 (33.2)	2 (0.9)
Albumin - Decrease	53 (24.8)	1 (0.5)
Activated Partial Thromboplastin Time - Increase	51 (23.8)	3 (1.4)
Creatinine - Increase	48 (22.4)	0 (0.0)

QD= once daily

<sup>a</sup> Grading defined by NCI CTCAE version 5.0

<sup>b</sup> Including 19 subjects reported hyponatremia in CodeBreak 100. The remaining 3 out of 22 subjects reported Grade 1 hyponatremia but without providing laboratory details.

### Post-Market Findings

At the time of authorization, no post-market findings were identified.

## 9 DRUG INTERACTIONS

### 9.1 Serious Drug Interactions

At the time of authorization, no serious drug interactions were identified.

### 9.2 Drug Interactions Overview

Drug interactions were observed when sotorasib was co-administered with an acid-reducing agent, a strong CYP3A4 inducer, a P-glycoprotein (P-gp) substrate, CYP3A4 substrates and Breast Cancer Resistance Protein (BCRP) substrates (see 10.3 Pharmacokinetics). The related findings and effects are discussed further below in Table 6.

### 9.3 Drug-Behavioural Interactions

At the time of authorization, drug-behavioural interactions have not been established.

### 9.4 Drug-Drug Interactions

The drugs listed in this table are based on either drug interaction case reports or studies, or potential interactions due to the expected magnitude and seriousness of the interaction (ie, those identified as contraindicated).

**Table 6. Established or Potential Drug-Drug Interactions**

	Source of Evidence	Effect	Clinical comment
Acid-Reducing Agents Effect on LUMAKRAS	CT	Coadministration of LUMAKRAS with PPI (omeprazole) or an H <sub>2</sub> receptor antagonist (famotidine) led to a ↓ in sotorasib concentrations.	Avoid coadministration of PPIs and H <sub>2</sub> receptor antagonists with LUMAKRAS since it may reduce the efficacy of sotorasib. If treatment with an acid-reducing agent cannot be avoided, take LUMAKRAS 4 hours before or 10 hours after administration of a local antacid (see 10.3 Pharmacokinetics).
Strong CYP3A4 Inducers Effect on LUMAKRAS	CT	Coadministration of LUMAKRAS with a strong CYP3A4 inducer (rifampin) led to a ↓ in sotorasib concentrations.	Avoid coadministration of strong CYP3A4 inducers with LUMAKRAS since it may reduce the efficacy of sotorasib (see 10.3 Pharmacokinetics).
LUMAKRAS Effect on CYP3A4 Substrates	CT	LUMAKRAS is a moderate CYP3A4 inducer. LUMAKRAS ↓ CYP3A4 substrates.	Avoid coadministration of LUMAKRAS and CYP3A4 sensitive substrates, for which minimal concentration changes may lead to therapeutic failures of the substrate. If coadministration cannot be avoided, increase the CYP3A4 substrate dosage in accordance with approved product labelling.

**Table 6. Established or Potential Drug-Drug Interactions**

	Source of Evidence	Effect	Clinical comment
LUMAKRAS Effect on P-glycoprotein (P-gp) substrates	CT	Coadministration of LUMAKRAS with a P-gp substrate (digoxin) led to an ↑ in digoxin concentrations.	Avoid coadministration of LUMAKRAS and P-gp substrates, for which minimal concentration changes may lead to increased adverse reactions. If coadministration cannot be avoided, decrease the P-gp substrate dosage in accordance with approved product labeling (see 10.3 Pharmacokinetics).
LUMAKRAS Effect on BCRP Substrates	CT	Coadministration of LUMAKRAS with a BCRP substrate led to an ↑ in the plasma concentrations of the BCRP substrate which may ↑ the effects of these substrates (see 10.3 Pharmacokinetics).	LUMAKRAS is a weak BCRP inhibitor. When coadministered with LUMAKRAS, monitor for adverse reactions of the BCRP substrate and decrease the BCRP substrate dosage in accordance with its product labeling.

CT = Clinical Trial

### 9.5 Drug-Food Interactions

Refer to 10.3 Pharmacokinetics, Absorption, *Effect of Food* for more information.

### 9.6 Drug-Herb Interactions

At the time of authorization, interactions with herbal products have not been established.

### 9.7 Drug-Laboratory Test Interactions

At the time of authorization, interactions with laboratory tests have not been established.

## NOC/c 10 CLINICAL PHARMACOLOGY

### 10.1 Mechanism of Action

Sotorasib is an oral, potent and highly selective inhibitor of KRAS<sup>G12C</sup>, a tumour-restricted, mutant-oncogenic form of the RAS GTPase, KRAS. Sotorasib forms an irreversible, covalent bond with the unique cysteine of KRAS<sup>G12C</sup>, locking the protein in an inactive state that prevents downstream signaling without affecting wild-type KRAS. Sotorasib blocked KRAS signaling, inhibited cell growth, promoted apoptosis, and was associated with anti-tumour inflammatory responses and immunity only in *KRAS p.G12C* tumour models. Sotorasib inhibited KRAS<sup>G12C</sup> *in vitro* and *in vivo* with minimal detectable off-target activity. In mice with xenografts of human or mouse tumour cells, sotorasib treatment led to tumour regressions at clinically relevant concentrations.

## 10.2 Pharmacodynamics

### Cardiac electrophysiology

The effect of sotorasib on the QT interval was assessed in 156 patients administered LUMAKRAS 960 mg once daily in clinical studies. LUMAKRAS did not prolong the QT interval to any clinically relevant extent.

### 10.3 Pharmacokinetics

The pharmacokinetics of sotorasib have been characterized in patients with *KRAS G12C*-mutated solid tumours, including NSCLC. Sotorasib exhibited non-linear, time-dependent, pharmacokinetics over the dose range of 180 mg to 960 mg once daily with similar systemic exposure (ie  $AUC_{0-24h}$  and  $C_{max}$ ) across doses at steady state, likely due to low solubility. Sotorasib systemic exposure ( $AUC_{0-24h}$  and  $C_{max}$ ) was comparable between film-coated tablets and film-coated tablets predispersed in water administered under fasted conditions. Steady state of sotorasib concentrations was reached within 22 days. No accumulation with multiple dosing was observed. The pharmacokinetics of sotorasib in NSCLC subjects is described in Table 7.

**Table 7. Summary of LUMAKRAS Pharmacokinetic Parameters in NSCLC Patients**

	$C_{max}$ (ng/mL)	$T_{max}$ (h)	$t_{1/2}$ (h)	$AUC_{inf}$ (hour•ng/mL)	$AUC_{0-24h}$ (hour•ng/mL)	CL/F (L/hour)	Vd (L)	AR
<b>960 mg QD, N = 25 Day 1</b>	5870 (8320, 59%)	1.0 (0.23 – 6.3)	5.56 (1.69) <sup>a</sup>	75500 (86700, 51%) <sup>a</sup>	73000 (81500, 44%) <sup>d</sup>	12.7 (15.2, 75%) <sup>a</sup>	98.2 (112, 61%) <sup>a</sup>	NC
<b>960 mg QD N = 21 Day 8</b>	4970 (7180, 55%)	1.0 (0.25 – 10)	5.19 (1.99) <sup>b</sup>	NC	36600 (43900, 58%) <sup>c</sup>	26.2 (32.5, 76%) <sup>c</sup>	211 (367, 135%) <sup>c</sup>	0.56 (0.65, 59%)

AR= accumulation ratio, QD= once daily,  $C_{max}$  = maximum observed drug concentration;  $T_{max}$  = time to reach  $C_{max}$ ;  $t_{1/2}$  = terminal half-life;  $AUC_{inf}$  = area under the drug concentration-time curve from time 0 to infinity;  $AUC_{0-24h}$  = area under the concentration-time curve from time 0 to 24 hour postdose; CL/F = apparent drug clearance observed at terminal phase; Vd = volume of distribution observed at terminal phase; NC = not calculated

Data presented as Geometric Mean (Mean, CV%) for all PK Parameters except for  $t_{max}$ , and  $t_{1/2}$  which is presented as Median (Range) and Mean (SD), respectively.

Values are reported to 3 significant figures except for  $t_{max}$  and CV% which are presented as 2 significant figures and the nearest integer, respectively. Day 8 observed PK was used to approximate steady state PK parameters

<sup>a</sup>N = 15 <sup>b</sup>N = 16, <sup>c</sup>N = 18, <sup>d</sup>N = 19

Pharmacokinetics of sotorasib have been characterized in patients with *KRAS G12C*-mutated solid tumours, including NSCLC, and healthy subjects.

### Absorption:

Following an oral, single-dose or multiple dose administration, the sotorasib median time to achieve peak concentration is 1 hour.

### Effect of Food

Following administration of a single 360 mg dose (3 X 120 mg) of sotorasib to healthy subjects with a high-fat, high-calorie meal, there was no effect on  $C_{max}$ , AUC increased by 38%, and



$T_{max}$  was delayed from 0.5 hours to 1.75 hours compared to administration under fasted conditions.

**Distribution:**

The mean volume of distribution at steady state of sotorasib was 211 L. *In vitro*, plasma protein binding of sotorasib is 89%.

**Metabolism:**

The main metabolic pathways of sotorasib are non-enzymatic glutathione conjugation and oxidative metabolism with CYP3As.

**Elimination:**

The mean plasma terminal elimination half-life (+/-SD) is 5 (2) hours. The mean oral clearance (CV%) (CL/F) was 12.7L/hour (75%) following a single oral dose of 960 mg and increased to 26.2L/hour (76%) at Day 8 following multiple dose administration.

**Excretion:**

After a single dose of radiolabeled sotorasib, approximately 74% of the dose was recovered in the feces (53% unchanged) and 6% (1% unchanged) recovered in the urine.

**Special Populations and Conditions**

No clinically meaningful differences in the pharmacokinetics of sotorasib were observed based on age (31 to 86 years), sex, race (75% Caucasian, 13% Asian and 8% Black), body weight, line of therapy, ECOG PS (0 and 1), or mild renal impairment (estimated glomerular filtration rate (eGFR) as determined by Modification of Diet in Renal Disease (MDRD) formula:  $\geq 60$  mL/min/1.73m<sup>2</sup>).

Based on limited data in the population pharmacokinetics analysis (n=37 out of 500 subjects, 7.4% of the total dataset), no clinically meaningful differences in the pharmacokinetics of sotorasib were observed in patients with moderate renal impairment (eGFR as determined by MDRD formula: 45 - 59 mL/min/1.73m<sup>2</sup>).

The effect of severe renal impairment on sotorasib pharmacokinetics has not been studied.

**Hepatic Impairment**

According to a population pharmacokinetic analysis, mild hepatic impairment (AST or ALT < 2.5 x ULN or total bilirubin < 1.5 x ULN) did not notably affect the pharmacokinetics of sotorasib.

In a dedicated clinical study, following administration of a single 960 mg dose,  $AUC_{inf}$  of total sotorasib decreased by 25.4% in subjects with moderate impairment (Child-Pugh B) and increased by 3.6% in subjects with severe impairment (Child-Pugh C), compared with subjects with normal liver function. The unbound  $AUC_{inf}$  of sotorasib increased by 1.8-fold in subjects with moderate hepatic impairment and by 6.3-fold in subjects with severe hepatic impairment.

## **Drug Interaction Studies**

### ***Clinical Studies***

#### ***Effect of Other Drugs on Sotorasib***

**Acid-Reducing Agents:** Coadministration of repeat doses of omeprazole (PPI) with a single dose of 960 mg LUMAKRAS decreased sotorasib  $C_{max}$  by 65% and AUC by 57% under fed conditions, and decreased sotorasib  $C_{max}$  by 57% and AUC by 42% under fasted conditions.

Coadministration of a single dose of famotidine ( $H_2$  receptor antagonist) given 10 hours prior to and 2 hours after a single dose of 960 mg LUMAKRAS decreased sotorasib  $C_{max}$  by 35% and AUC by 38% (see 9.4 Drug-Drug Interactions).

**Strong CYP3A4 Inducers:** Coadministration of LUMAKRAS with repeat doses of rifampin (a strong CYP3A4 inducer) decreased sotorasib  $C_{max}$  by 35% and AUC by 51% (see 9.4 Drug-Drug Interactions).

**Other Drugs:** No clinically meaningful effect on the exposure of sotorasib was observed following coadministration of LUMAKRAS with itraconazole (a combined strong CYP3A4 and P-gp inhibitor), and a single dose of rifampin (an OATP1B1/1B3 inhibitor), or metformin (a MATE1/MATE2-K substrate).

#### ***Effect of Sotorasib on Other Drugs***

**CYP3A substrates:** Sotorasib is a CYP3A4 inducer. Coadministration of LUMAKRAS with midazolam (a sensitive CYP3A substrate) decreased midazolam  $C_{max}$  by 48% and AUC by 53% (see 9.4 Drug-Drug Interactions).

**P-gp substrates:** Sotorasib is a P-gp inhibitor. Co-administration of LUMAKRAS with digoxin (a P-gp substrate) increased digoxin  $C_{max}$  by 91% and AUC by 21%.

**MATE1/MATE2-K substrates:** No clinically meaningful effect on the exposure of metformin (a MATE1/MATE2-K substrate) was observed following coadministration of LUMAKRAS.

**BCRP substrates:** Sotorasib is a BCRP inhibitor. Coadministration of LUMAKRAS with rosuvastatin (a BCRP substrate) increased rosuvastatin  $C_{max}$  by 70% and AUC by 34% (see 9.4 Drug-Drug Interactions).

### ***In Vitro Studies***

**Cytochrome P450 (CYP) Enzymes:** Sotorasib may induce CYP2C8, CYP2C9, CYP2C19, and CYP2B6. *In vitro* data indicated sotorasib does not inhibit CYP1A2, CYP2B6, CYP2C8, CYP2C9, CYP2C19, or CYP2D6.

**Transporter Systems:** *In vitro* data indicated that sotorasib is a substrate of P-gp.

## **11 STORAGE, STABILITY AND DISPOSAL**

No special storage conditions required.

Store at 15°C to 30°C.

## **12 SPECIAL HANDLING INSTRUCTIONS**

Dispose of any unused medicinal product or waste material in accordance with local requirements.


**PART II: SCIENTIFIC INFORMATION****13 PHARMACEUTICAL INFORMATION****Drug Substance**

Proper /Common name: sotorasib

Chemical name: 6-fluoro-7-(2-fluoro-6-hydroxyphenyl)-(1*M*)-1-[4-methyl-2-(propan-2-yl)pyridine 3-yl]-4-[(2*S*)-2-methyl-4-(prop-2-enoyl)piperazin-1-yl]-pyrido[2,3-*d*]pyrimidin-2(1*H*)-one

Molecular formula and molecular mass: C<sub>30</sub>H<sub>30</sub>F<sub>2</sub>N<sub>6</sub>O<sub>3</sub> 560.6 g/mol

Structural formula:



Physicochemical properties: A white to off-white to yellow to light brown powder with a pKa of 8.06 and 4.56 (determined electrophoretically) and with a pH 5.6 at 0.06 mg/mL (deionized water). It is slightly hygroscopic, gaining < 0.3 wt%, water between 0% to 90% Relative Humidity (RH). The log of the distribution coefficient (LogD – octanol/water) at pH 7.4 is 2.44. The solubility of sotorasib in the aqueous media decreases over the range of pH 1.2 to 6.8 from 1.3 mg/mL to 0.03 mg/mL.

## 14 CLINICAL TRIALS

### NOC/c 14.1 Trial Design and Study Demographics

**Table 8. Summary of patient demographics for clinical trials in Locally Advanced or Metastatic NSCLC**

Study #	Study design	Dosage, route of administration and duration	Study subjects (n)	Median Age (Range)	Gender
20170543 (CodeBreak 100)	Single-arm, open label, multicenter trial	960 mg, oral, once daily until disease progression or unacceptable toxicity	126	64 years (37 to 80 years)	Male and Female

#### **Advanced KRAS G12C-mutated NSCLC in previously treated patients (CodeBreak 100 Study)**

The efficacy of LUMAKRAS was demonstrated in a single-arm, open-label, multicenter trial (CodeBreak 100) that enrolled patients with locally advanced or metastatic *KRAS G12C*-mutated NSCLC who had disease progression after receiving prior therapy. Key eligibility criteria included progression on an immune checkpoint inhibitor and/or platinum-based chemotherapy, an Eastern Cooperative Oncology Group Performance Status (ECOG PS) of 0 or 1, and at least one measurable lesion as defined by Response Evaluation Criteria in Solid Tumours (RECIST v1.1), adequate hematological, hepatic, renal and cardiac function parameters. Subjects with active brain metastases were excluded.

All patients were required to have prospectively identified *KRAS G12C*-mutated NSCLC in tumour tissue samples using the QIAGEN theascreen *KRAS* RGQ PCR Kit performed in a central laboratory. A total of 124 patients had at least one measurable lesion at baseline as assessed by Blinded Independent Central Review (BICR) according to RECIST v1.1 and were treated with LUMAKRAS 960 mg once daily until disease progression or unacceptable toxicity. The median duration of treatment was 5.5 months (range 0 to 34) with 48% of patients treated for  $\geq 6$  months and 25% of patients treated for  $\geq 12$  months. The major efficacy outcome measure was objective response rate (ORR) and duration of response (DOR) as evaluated by a BICR according to RECIST v1.1).

The baseline demographic and disease characteristics of the study population were: median age 64 years (range 37 to 80) with 47%  $\geq 65$  years and 8%  $\geq 75$  years, 50% Female, 82% White, 15% Asian, 2% Black; 70% ECOG PS 1, 96% had stage IV disease, 99% with nonsquamous histology, 81% former smokers, 12% current smokers, 5% never smokers. All patients received at least 1 prior line of systemic therapy for metastatic NSCLC; 43% received only 1 prior line of therapy, 35% received 2 prior lines of therapy, 22% received 3 prior lines of therapy; 91% received prior anti-PD-1/PD-L1 immunotherapy, 90% received prior platinum-based chemotherapy, 81% received both platinum-based chemotherapy and anti-PD-1/PD-L1. The sites of known extra-thoracic metastasis included 48% bone, 21% brain, and 21% liver.

## 14.2 Study Results

### Advanced KRAS G12C-mutated NSCLC in previously treated patients (CodeBreak 100 Study)

Efficacy results are summarized in Table 9. The ORR was 37% (95% CI: 29, 46). The patients with objective responses had DOR ranging from 1.3 to 32.6 months, and 4.3% were still on study with ongoing response after a median duration of follow-up of 29.2 months. The median time to response (TTR) was 1.4 months (range 1.2 to 10.1), with 70% of responses occurring within the first 7 weeks. Consistent efficacy results were seen in patients with *KRAS G12C* mutation identified in either tissue or plasma specimens.

**Table 9. Efficacy Results in CodeBreak 100 for Patients with *KRAS G12C*-mutated NSCLC**

Efficacy Parameter	LUMAKRAS N = 124
<b>ORR</b> (95% CI) <sup>a</sup>	37.1 (28.6, 46.2)
Complete response, %	4.0
Partial response, %	33.1
<b>DOR</b> <sup>a</sup>	
Median <sup>b</sup> , months (range)	11.1 (1.3, 32.6)
95% CI	6.9, 15.0
Patients with duration ≥ 6 months <sup>b</sup> , %	71.3

CI = confidence interval; DOR = duration of response; ORR = objective response rate

<sup>a</sup> Response-related efficacy outcome as per BICR, RECIST 1.1

<sup>b</sup> Estimate using Kaplan-Meier method

Data cut-off period Sept 2022

## 15 MICROBIOLOGY

No microbiological information is required for this drug product.

## 16 NON-CLINICAL TOXICOLOGY

### General Toxicology

In the 3-month rat toxicology study, renal toxicity including minimal to marked histologic tubular degeneration/necrosis and increased kidney weight, urea nitrogen, creatinine, and urinary biomarkers of renal tubular injury, were present at sotorasib doses ≥ 60 mg/kg (approximately 0.4 times the human exposure at the clinical dose of 960 mg based on AUC). Increases in cysteine S-conjugate β-lyase pathway metabolism in the rat kidney compared to human may make rats more susceptible to renal toxicity due to local formation of a putative sulfur-containing metabolite.

In the 3-month dog toxicology study, there were sotorasib-induced findings in the liver (centrilobular hepatocellular hypertrophy), pituitary gland (hypertrophy of basophils), and thyroid gland (marked follicular cell atrophy, moderate to marked colloid depletion, and mild to moderate follicular cell hypertrophy) at doses  $\geq$  200 mg/kg (approximately 0.35 times the human exposure at the clinical dose of 960 mg based on AUC). These findings may be due to an adaptive response to hepatocellular enzyme induction and subsequent reduced thyroid hormone levels (secondary hypothyroidism), although thyroid levels were not measured in dogs.

**Carcinogenicity:**

Carcinogenicity studies have not been performed with sotorasib.

**Genotoxicity:**

Sotorasib was not mutagenic in an *in vitro* bacterial reverse mutation (Ames) assay. Sotorasib was not genotoxic in the *in vivo* rat micronucleus and comet assays.

**Reproductive and Developmental Toxicology:**

In a rat embryo-fetal developmental study, once daily oral administration of sotorasib to pregnant rats during the period of organogenesis resulted in maternal toxicity at 540 mg/kg (equivalent to approximately 4 times the human exposure, based on AUC, at the clinical dose of 960 mg). Sotorasib did not affect embryo-fetal survival or development at doses up to 540 mg/kg.

In a rabbit embryo-fetal developmental study, once daily administration of sotorasib during the period of organogenesis resulted in lower fetal body weights, and a reduction in the number of ossified metacarpals, in fetuses at the 100 mg/kg dose level (equivalent to approximately 2.3 times the human exposure, based on AUC, at the clinical dose of 960 mg), which was associated with maternal toxicity including effects such as decreased body weight gain and decreased food consumption during the dosing phase.

**Impairment of Fertility:**

Fertility/early embryonic development studies were not conducted with sotorasib. There were no adverse effects on female or male reproductive organs in general toxicology studies conducted in dogs and rats.

## PATIENT MEDICATION INFORMATION

### READ THIS FOR SAFE AND EFFECTIVE USE OF YOUR MEDICINE

Pr **LUMAKRAS**<sup>®</sup>

**sotorasib tablets**

Read this carefully before you start taking **LUMAKRAS** and each time you get a refill. This leaflet is a summary and will not tell you everything about this drug. Talk to your healthcare professional about your medical condition and treatment and ask if there is any new information about **LUMAKRAS**.

#### What is **LUMAKRAS** used for?

Please see the boxed text below.

“For the following indication LUMAKRAS has been approved with conditions (NOC/c). This means it has passed Health Canada’s review and can be bought and sold in Canada, but the manufacturer has agreed to complete more studies to make sure the drug works the way it should. For more information, talk to your healthcare professional.”

- LUMAKRAS is used to treat adults with non-small cell lung cancer (NSCLC) with an abnormal gene called “*KRAS G12C*”. This cancer:
  - cannot be removed by surgery or other treatment, or has spread to other parts of the body, and
  - has been treated with at least one type of cancer treatment before.
- LUMAKRAS is not approved for use in children and adolescents under 18 years of age. Your healthcare professional will test your cancer for abnormal *KRAS G12C* and make sure that LUMAKRAS is right for you.

#### What is a Notice of Compliance with Conditions (NOC/c)?

A Notice of Compliance with Conditions (NOC/c) is a type of approval to sell a drug in Canada. Health Canada only gives a NOC/c to a drug that treats, prevents, or helps identify a serious or life-threatening illness. The drug must show promising proof that it works well, is of high quality, and is reasonably safe. Also, the drug must either respond to a serious medical need in Canada or be much safer than existing treatments.

Drug makers must agree in writing to clearly state on the label that the drug was given an NOC/c, to complete more testing to make sure the drug works the way it should, to actively monitor the drug’s performance after it has been sold, and to report their findings to Health Canada.

#### How does **LUMAKRAS** work?

LUMAKRAS helps block the abnormal *KRAS G12C* protein in your lung cancer. This may slow down or stop the growth and spread of the lung cancer.

If you have any questions about how LUMAKRAS works or why this medicine has been prescribed for you, ask your healthcare professional.

**What are the ingredients in LUMAKRAS?**

Medicinal ingredients: sotorasib

Non-medicinal ingredients:

croscarmellose sodium, iron oxide yellow, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, talc, titanium dioxide

**LUMAKRAS comes in the following dosage forms:**

Tablet, 120 mg, 240 mg

**Do not use LUMAKRAS if:**

- You are allergic to sotorasib or any of the other ingredients in LUMAKRAS or the packaging.

**To help avoid side effects and ensure proper use, talk to your healthcare professional before you take LUMAKRAS. Talk about any health conditions or problems you may have, including if you:**

- Have or had liver problems.
- Have or had lung problems.
- Have an intolerance to lactose.

**Other warnings you should know about:**

LUMAKRAS can cause the following side effects, including:

- **Liver problems: Hepatotoxicity** (liver damage), including **liver injury** and **hepatitis** (liver inflammation) can happen in patients taking LUMAKRAS. Increased levels of liver enzyme levels can happen as well. You will have regular blood tests done before starting your treatment and then every 3 weeks during the first 3 months, then once a month after. You might need more tests depending on your health. These blood tests will tell your healthcare professional how your liver is working.
- **Lung or breathing problems: Interstitial lung disease (ILD) / pneumonitis** (inflamed or scarred lungs) can happen while taking LUMAKRAS. They might cause death. Tell your healthcare professional if you have new or worsening shortness of breath, cough, or fever. If these symptoms become severe, get emergency medical help right away.

See the “Serious side effects and what to do about them” table, below, for more information on these and other serious side effects.

**Pregnancy, contraception and breastfeeding:****Female patients**

- Tell your healthcare professional if you are pregnant, able to get pregnant or think you are pregnant. There are specific risks you should discuss with your healthcare professional.
- Tell your healthcare professional right away if you become pregnant or think you may be pregnant during your treatment with LUMAKRAS.
- Do not breastfeed while you are taking LUMAKRAS and for at least 1 week after your last dose.



**Tell your healthcare professional about all the medicines you take, including any drugs, vitamins, minerals, natural supplements or alternative medicines.**

**The following medicines may interact with LUMAKRAS:**

- Medicines used to reduce stomach acid and to treat stomach ulcers, indigestion and heartburn such as:
  - dexlansoprazole, esomeprazole, lansoprazole, omeprazole, pantoprazole sodium, or rabeprazole (medicines known as 'proton pump inhibitors')
  - ranitidine, famotidine, cimetidine (medicines known as 'H<sub>2</sub> receptor antagonists')
- Medicines used to treat tuberculosis, such as rifampin
- Medicines used to treat epilepsy, such as carbamazepine, phenytoin, or phenobarbital
- St. John's Wort (a herbal medicine used to treat depression)
- Enzalutamide (used to treat prostate cancer)
- Medicines used to treat severe pain, such as alfentanil or fentanyl
- Medicines used to prevent organ rejection in transplants, such as cyclosporin, sirolimus, everolimus, or tacrolimus
- Medicines used to lower cholesterol levels, such as simvastatin, atorvastatin, lovastatin, or rosuvastatin
- Midazolam (used to treat acute seizures or as a sedative before or during surgery or medical procedures)
- Medicines used to treat heart rhythm problems, such as dronedarone, amiodarone or digoxin
- Medicines to treat blood clots (anticoagulants), such as rivaroxaban or apixaban

**Please note this list of medicines does not cover all possible medications. Please consult with your healthcare professional.**

**How to take LUMAKRAS:**

- Take LUMAKRAS exactly as your healthcare professional tells you to take it. Do not change your dose or stop taking LUMAKRAS unless your healthcare professional tells you to.
- LUMAKRAS can be taken with or without food.
- If you need to take an antacid medicine (to reduce stomach acid), take LUMAKRAS either 4 hours before or 10 hours after the antacid.
- Swallow tablets whole. Do not chew, crush, or split tablets.

If you cannot swallow LUMAKRAS tablets whole:

- Place your daily dose of LUMAKRAS in a glass of 120 mL (4 ounces) non-carbonated, room temperature water without crushing the tablets. Do not use any other liquids.
- Stir until the tablets are in small pieces (the tablets will not completely dissolve). The water mixture may have a pale to bright yellow look.
- Drink the LUMAKRAS and water mixture right away.
- Add another 120 mL (4 ounces) of water and drink right away to make sure that you have taken the full dose of LUMAKRAS.
- If you do not drink the mixture right away, stir the mixture again. Drink within two hours of making the mixture.
- If you have a feeding tube your healthcare professional will provide guidance on how to take your medicine.

**Usual dose:**

**Recommended dose:** 960 mg (eight 120 mg tablets or four 240 mg tablets) once per day. Take by mouth at the same time each day.

Your healthcare professional may lower your dose, stop your treatment for a period of time or recommend that you stop treatment completely. This may happen if you:

- experience serious side effects, or
- your disease gets worse.

**Overdose:**

Contact your healthcare professional right away if you take more tablets than recommended.

If you think you, or a person you are caring for, have taken too much LUMAKRAS, contact a healthcare professional, hospital emergency department, or regional poison control centre immediately, even if there are no symptoms.

**Missed Dose:**

If you miss a dose of LUMAKRAS, do not take the dose if 6 hours have passed from your regular scheduled time. Take your next dose at your regular scheduled time.

If you vomit at any time after taking LUMAKRAS, do not take another dose. Take the next dose at your usual time.

**What are possible side effects from using LUMAKRAS?**

These are not all the possible side effects you may have when taking LUMAKRAS. If you experience any side effects not listed here, tell your healthcare professional.

- Diarrhea
- Nausea, vomiting
- Feeling tired (fatigue)
- Cough

- Stomach pain
- Constipation
- Decreased appetite
- Joint pain (arthralgia)
- Rash

LUMAKRAS can cause abnormal blood test results. Your healthcare professional will do blood tests during your treatment. These will tell your healthcare professional how LUMAKRAS is affecting your blood and liver.

<b>Serious side effects and what to do about them</b>			
<b>Symptom / effect</b>	<b>Talk to your healthcare professional</b>		<b>Stop taking drug and get immediate medical help</b>
	<b>Only if severe</b>	<b>In all cases</b>	
<b>VERY COMMON</b>			
<b>Diarrhea</b>	X		
<b>Edema:</b> unusual swelling of the arms, hands, legs, feet and ankles, face or airway passages	X		
<b>Hepatotoxicity</b> (swelling of your liver), <b>Hepatitis</b> (liver inflammation), <b>Liver Injury:</b> jaundice (yellowing of the skin or whites of eyes), urine turns dark, light-coloured stool, loss of appetite for several days or longer, nausea, stomach pain, fever, fatigue, weakness, vomiting		X	
<b>Nausea</b>	X		
<b>Musculoskeletal pain</b> (pain that affects the muscles and tendons along with bones): muscle pain, limb pain, joint pain and bone pain	X		
<b>Vomiting</b>	X		
<b>Lung or breathing problems (pneumonia, pneumonitis, interstitial lung disease):</b> new or worsening lung problems, trouble breathing, shortness of breath, chest pain, cough, fluid build up in the lungs, or fever		X	

Serious side effects and what to do about them			
Symptom / effect	Talk to your healthcare professional		Stop taking drug and get immediate medical help
	Only if severe	In all cases	
<b>COMMON</b>			
<b>Pulmonary embolism</b> (blood clot in the lung): chest pain that may increase with deep breathing, cough, coughing up bloody sputum, shortness of breath		X	

If you have a troublesome symptom or side effect that is not listed here or becomes bad enough to interfere with your daily activities, tell your healthcare professional.

### Reporting Side Effects

You can report any suspected side effects associated with the use of health products to Health Canada by:

- Visiting the Web page on Adverse Reaction Reporting (<https://www.canada.ca/en/health-canada/services/drugs-health-products/medeffect-canada.html>) for information on how to report online, by mail or by fax; or
- Calling toll-free at 1-866-234-2345.

*NOTE: Contact your health professional if you need information about how to manage your side effects. The Canada Vigilance Program does not provide medical advice.*

### Storage:

This medicine does not require any special storage conditions.

Store at 15°C to 30°C.

Keep LUMAKRAS out of the sight and reach of children.

Ask your pharmacist how to throw away drugs you no longer use.

### If you want more information about LUMAKRAS:

- Talk to your healthcare professional
- Find the full product monograph that is prepared for healthcare professionals and includes this Patient Medication Information by visiting the Health Canada website: (<https://www.canada.ca/en/health-canada/services/drugs-health-products/drug-products/drug-product-database.html>); the manufacturer's website [www.amgen.ca](http://www.amgen.ca), or by calling 1-866-50-AMGEN (1-866-502-6436).

This leaflet was prepared by Amgen Canada Inc.

Last Revised: OCT 28, 2024