

PRODUCT MONOGRAPH  
INCLUDING PATIENT MEDICATION INFORMATION

<sup>Pr</sup>**ACCEL-ONDANSETRON ODT**

ondansetron

Orally disintegrating tablets, 4 mg and 8 mg, oral

Mfr. Std.

Antiemetic

5-HT<sub>3</sub> receptor antagonist

ATC code A04AA01

ACCEL PHARMA INC.  
119 Labrosse Avenue  
Pointe-Claire, QC, H9R1A3, Canada

Date of Initial Authorization:  
February 08, 2023

Date of Revision:  
October 30, 2024

Submission Control Number: 286958

**RECENT MAJOR LABEL CHANGES**

Section	Date
<a href="#">1 INDICATIONS</a>	10/2024
<a href="#">4 DOSAGE AND ADMINISTRATION, 4.1 Dosing Considerations</a>	10/2024
<a href="#">4 DOSAGE AND ADMINISTRATION, 4.2 Recommended Dose and Dosage Adjustment</a>	10/2024
<a href="#">4 DOSAGE AND ADMINISTRATION, 4.4 Administration</a>	10/2024
<a href="#">7 WARNINGS AND PRECAUTIONS, Neurologic, Serotonin toxicity / Neuroleptic Malignant Syndrome</a>	10/2024

**TABLE OF CONTENTS**

<b>RECENT MAJOR LABEL CHANGES</b> .....	<b>2</b>
<b>TABLE OF CONTENTS</b> .....	<b>2</b>
<b>PART I: HEALTH PROFESSIONAL INFORMATION</b> .....	<b>4</b>
<b>1 INDICATIONS</b> .....	<b>4</b>
1.1 Pediatrics .....	4
1.2 Geriatrics .....	4
<b>2 CONTRAINDICATIONS</b> .....	<b>5</b>
<b>4 DOSAGE AND ADMINISTRATION</b> .....	<b>5</b>
4.1 Dosing Considerations .....	5
4.2 Recommended Dose and Dosage Adjustment .....	5
4.4 Administration .....	8
4.5 Missed Dose .....	9
<b>5 OVERDOSAGE</b> .....	<b>9</b>
<b>6 DOSAGE FORMS, STRENGTHS, COMPOSITION AND PACKAGING</b> .....	<b>10</b>
<b>7 WARNINGS AND PRECAUTIONS</b> .....	<b>10</b>
7.1 Special Populations .....	13
7.1.1 Pregnant Women .....	13
7.1.2 Breast-feeding .....	14
7.1.3 Pediatrics .....	14
7.1.4 Geriatrics .....	14
<b>8 ADVERSE REACTIONS</b> .....	<b>14</b>

8.1	Adverse Reaction Overview.....	14
8.2	Clinical Trial Adverse Reactions .....	14
8.5	Post-Market Adverse Reactions .....	15
<b>9</b>	<b>DRUG INTERACTIONS .....</b>	<b>17</b>
9.1	Serious Drug Interactions .....	17
9.2	Drug Interactions Overview.....	17
9.3	Drug-Behavioural Interactions.....	17
9.4	Drug-Drug Interactions .....	18
9.5	Drug-Food Interactions.....	20
9.6	Drug-Herb Interactions.....	20
9.7	Drug-Laboratory Test Interactions.....	20
<b>10</b>	<b>CLINICAL PHARMACOLOGY .....</b>	<b>20</b>
10.1	Mechanism of Action.....	20
10.2	Pharmacodynamics .....	20
10.3	Pharmacokinetics .....	22
<b>11</b>	<b>STORAGE, STABILITY AND DISPOSAL .....</b>	<b>24</b>
<b>12</b>	<b>SPECIAL HANDLING INSTRUCTIONS.....</b>	<b>24</b>
	<b>PART II: SCIENTIFIC INFORMATION .....</b>	<b>25</b>
<b>13</b>	<b>PHARMACEUTICAL INFORMATION.....</b>	<b>25</b>
<b>14</b>	<b>CLINICAL TRIALS .....</b>	<b>25</b>
14.1	Clinical Trials by Indication .....	25
14.2	Comparative Bioavailability Studies.....	27
<b>15</b>	<b>MICROBIOLOGY .....</b>	<b>28</b>
<b>16</b>	<b>NON-CLINICAL TOXICOLOGY .....</b>	<b>28</b>
<b>17</b>	<b>SUPPORTING PRODUCT MONOGRAPHS.....</b>	<b>31</b>
	<b>PATIENT MEDICATION INFORMATION.....</b>	<b>32</b>

## PART I: HEALTH PROFESSIONAL INFORMATION

### 1 INDICATIONS

Accel-Ondansetron ODT (ondansetron) is indicated for:

- the prevention of nausea and vomiting associated with mildly and moderately emetogenic chemotherapy, and radiotherapy.
- the maintenance of antiemesis following intravenous doses of ondansetron used for the prevention of nausea and vomiting associated with highly emetogenic chemotherapy, including cisplatin.
- the prevention of post-operative nausea and vomiting.

#### 1.1 Pediatrics

##### **Prevention of Nausea and Vomiting Associated with Mildly and Moderately Emetogenic Chemotherapy**

- **Pediatrics (4-12 years of age):** Based on the data submitted and reviewed by Health Canada, the safety and efficacy of ACCEL-ONDANSETRON ODT in pediatric patients 4-12 years of age has been established. Therefore, Health Canada has authorized an indication for pediatric use. (see [4.2 Recommended Dose and Dosage Adjustment](#)).
- **Pediatrics (<4 years of age):** No data are available to Health Canada; therefore, Health Canada has not authorized an indication for the use of ACCEL-ONDANSETRON ODT in children less than 4 years of age.

##### **Prevention of Nausea and Vomiting Associated with Radiotherapy**

No data are available to Health Canada; therefore, Health Canada has not authorized an indication for this use in any pediatric population.

##### **Prevention of Post-Operative Nausea and Vomiting**

No data are available to Health Canada; therefore, Health Canada has not authorized an indication for this use in any pediatric population.

#### 1.2 Geriatrics

##### **Prevention of Nausea and Vomiting Associated with Mildly and Moderately Emetogenic Chemotherapy, and Radiotherapy**

Efficacy and tolerance of ondansetron were similar to that observed in younger adults (see [4.2 Recommended Dose and Dosage Adjustment, Prevention of Less Emetogenic Chemotherapy Induced Nausea and Vomiting, Geriatrics](#); [7 WARNINGS AND PRECAUTIONS, Cardiovascular](#); [7.1.4 Geriatrics](#); [10.3 Pharmacokinetics, Geriatrics](#)).

## Prevention of Post-Operative Nausea and Vomiting

Clinical experience in the use of ondansetron in the prevention and treatment of post-operative nausea and vomiting in elderly patients is limited. Therefore, Health Canada has not authorized an indication for this use in the geriatric population.

## 2 CONTRAINDICATIONS

- ACCEL-ONDANSETRON ODT (ondansetron) is contraindicated in patients who are hypersensitive to this drug or to any ingredient in the formulation, including any non-medicinal ingredient, or component of the container. For a complete listing, see [6 DOSAGE FORMS, STRENGTHS, COMPOSITION AND PACKAGING](#).
- The concomitant use of apomorphine with ondansetron is contraindicated based on reports of profound hypotension and loss of consciousness when apomorphine was administered with ondansetron (see [9.1 Serious Drug Interactions](#); [9.4 Drug-Drug Interactions](#)).

## 4 DOSAGE AND ADMINISTRATION

### 4.1 Dosing Considerations

- ACCEL-ONDANSETRON ODT (ondansetron) is an orally disintegrating tablet. When injectable ondansetron is used, the product monograph for ondansetron hydrochloride dihydrate injection should be consulted.
- In patients with moderate or severe hepatic impairment, the total daily dose should not exceed 8 mg (see [7 WARNINGS AND PRECAUTIONS, Hepatic/Biliary/Pancreatic](#)).
- Exposure-response modelling predicted a greater effect on QTcF in patients  $\geq 75$  years of age compared to young adults (see [10.3 Pharmacokinetics, Geriatrics](#)).
- Dosing considerations that reduce cardiac risks:
  - Carefully follow the dosing guidelines
  - Use the minimum effective dose.
  - Use oral ondansetron formulations, if possible (lower  $C_{max}$ ).

### 4.2 Recommended Dose and Dosage Adjustment

#### Prevention of Highly Emetogenic Chemotherapy Induced Nausea and Vomiting

##### Adults

ACCEL-ONDANSETRON ODT is not indicated for this use.

Oral formulations of ondansetron, such as ACCEL-ONDANSETRON ODT, are not intended for the prevention of nausea and vomiting associated with highly emetogenic chemotherapy, such as cisplatin. An intravenous formulation should be given as an initial dose prior to chemotherapy and within the first 24 hours. **Note: ACCEL-**

**ONDANSETRON ODT is only available as oral disintegrating tablets. A product monograph for ondansetron hydrochloride injection should be consulted.**

- **Pediatrics (<18 years)**

Ondansetron is not indicated for this use in children.

### **Maintenance of Antiemesis Following Intravenous Doses of Ondansetron Used for the Prevention of Nausea and Vomiting Associated with Highly Emetogenic Chemotherapy**

- **Adults**

For the maintenance of anti-emesis established by intravenous ondansetron:

- 8 mg orally every 8 hours, for up to 5 days; following the first 24 hours.

- **Pediatrics (<18 years of age)**

ACCEL-ONDANSETRON ODT is not indicated for this use in children under 18 years of age.

- **Geriatrics**

Efficacy and tolerance in patients 65 years of age and older were similar to that seen in younger adults. No dosage adjustment is required in this population (see [7.1.4 Geriatrics](#); [10.3 Pharmacokinetics, Geriatrics](#)).

- **Hepatic Insufficiency**

No dosage adjustment is needed in patients with mild hepatic impairment.

In patients with moderate or severe hepatic impairment (Child-Pugh score  $\geq 7$ ), do not exceed a total daily dose of 8 mg (Child-Pugh score of 10 or greater) (see [7 WARNINGS AND PRECAUTIONS, Hepatic/Biliary/Pancreatic](#); [10.3 Pharmacokinetics](#)). This may be given as a single oral dose.

- **Renal Insufficiency**

No alteration of daily dosage, frequency of dosing, or route of administration is required in patients with impaired renal function (see [10.3 Pharmacokinetics, Renal Insufficiency](#)).

### **Prevention of Nausea and Vomiting Associated with Mildly and Moderately Emetogenic Chemotherapy Adults**

8 mg orally, 1-2 hours prior to chemotherapy, followed by 8 mg orally, twice daily for up to 5 days

- **Pediatrics (4- <18 years of age)**

Following chemotherapy\*: 4 mg orally, every 8 hours, for up to 5 days.

\*For prevention of chemotherapy induced nausea and vomiting in children 4-17 years of age, intravenous ondansetron should be administered 30 minutes before chemotherapy. **Note: ACCEL-ONDANSETRON ODT is only available as an orally disintegrating tablet. A product monograph for ondansetron hydrochloride injection should be consulted.**

- **Pediatrics (<4 years of age)**

ACCEL-ONDANSETRON ODT is not indicated for this use in children under 4 years of age.

- **Geriatrics**

Efficacy and tolerance in patients 65 years of age and older were similar to that seen in younger adults indicating no need to alter dosage schedules in this population (see [10.3 Pharmacokinetics, Geriatrics](#)).

- **Hepatic Insufficiency**

No dosage adjustment is needed in patients with mild hepatic impairment.

In patients with moderate or severe hepatic impairment (Child-Pugh score  $\geq 7$ ), do not exceed a total daily dose of 8 mg (Child-Pugh score of 10 or greater) (see [7 WARNINGS AND PRECAUTIONS, Hepatic/Biliary/Pancreatic; 10.3 Pharmacokinetics](#)). This may be given as a single oral dose.

- **Renal Insufficiency**

No alteration of daily dosage, frequency of dosing, or route of administration is required in patients with impaired renal function (see [10.3 Pharmacokinetics, Renal Insufficiency](#)).

## Prevention of Radiotherapy Induced Nausea and Vomiting

- **Adults**

8 mg orally, given 1-2 hours before radiotherapy followed by 8 mg orally, given every 8 hours, for up to 5 days after a course of treatment

- **Pediatrics (<18 years of age)**

ACCEL-ONDANSETRON ODT is not indicated for this use in the pediatric population.

- **Geriatrics**

Efficacy and tolerance in patients 65 years of age and older were similar to that seen in younger adults, indicating no need to alter dosage schedules in this population (see [Adults](#), and [4.1 Dosing Considerations; 10.3 Pharmacokinetics, Geriatrics](#)).

- **Hepatic Insufficiency**

No dosage adjustment is needed in patients with mild hepatic impairment.

In patients with moderate or severe hepatic impairment (Child-Pugh score  $\geq 7$ ), do not exceed a total daily dose of 8 mg (Child-Pugh score of 10 or greater) (see [7 WARNINGS AND PRECAUTIONS, Hepatic/Biliary/Pancreatic; 10.3 Pharmacokinetics](#)). This may be given as a single oral dose.

- **Renal Insufficiency**

No alteration of daily dosage, frequency of dosing, or route of administration is required in patients with impaired renal function (see [10.3 Pharmacokinetics, Renal Insufficiency](#)).

#### Prevention of Post-Operative Nausea and Vomiting

- **Adults**

16 mg orally, administered 1 hour prior to induction of anaesthesia.

- **Pediatrics (<18 years of age)**

ACCEL-ONDANSETRON ODT is not indicated for this use in the pediatric population.

- **Geriatrics**

ACCEL-ONDANSETRON ODT is not indicated for this use in patients 65 years of age and older.

- **Hepatic Insufficiency**

No dosage adjustment is needed in patients with mild hepatic impairment.

In patients with moderate or severe hepatic impairment (Child-Pugh score  $\geq 7$ ), do not exceed a total daily dose of 8 mg (Child-Pugh score of 10 or greater) (see [7 WARNINGS AND PRECAUTIONS, Hepatic/Biliary/Pancreatic; 10.3 Pharmacokinetics](#)). This may be given as a single oral dose.

- **Renal Insufficiency**

No adjustment of daily dosage, frequency of dosing, or route of administration is required in patients with impaired renal function (see [10.3 Pharmacokinetics, Renal Insufficiency](#)).

#### 4.4 Administration

Patient instructions:

- **Do not** try to push ACCEL-ONDANSETRON ODT out through the foil backing on the blister pocket.
- Just prior to use, with **dry** fingers, carefully peel back the foil backing where indicated by the arrow. Gently push the tablet out of the blister pocket.
- Immediately place the ACCEL-ONDANSETRON ODT tablet on top of the tongue. It will disperse within seconds, then swallow as normal.



#### 4.5 Missed Dose

If a dose is missed, it should be taken as soon as possible. However, if it is almost time for the next dose, the missed dose should be skipped and the patient returned to their regular dosing schedule. **Do not double doses.**

If a dose is missed, and the patient feels nauseated or vomits, the missed dose should be taken as soon as possible.

If a dose of ACCEL-ONDANSETRON ODT is vomited, do not repeat the dose. ACCEL-ONDANSETRON ODT is rapidly absorbed.

### 5 OVERDOSAGE

At present there is little information concerning overdose with ondansetron. Individual doses of 84 mg and 145 mg and total daily doses as large as 252 mg have been administered with only mild side effects. There is no specific antidote for ondansetron, therefore, in cases of suspected overdose, symptomatic and supportive therapy should be given as appropriate.

The use of Ipecac to treat overdose with ondansetron is not recommended as patients are unlikely to respond due to the antiemetic action of ondansetron itself.

“Sudden blindness” (amaurosis) of 2 to 3 minutes duration plus severe constipation occurred in one patient that was administered 72 mg of ondansetron intravenously as a single dose. Hypotension (and faintness) occurred in another patient that took 48 mg of oral ondansetron. Following infusion of 32 mg over only a 4-minute period, a vasovagal episode with transient second-degree heart block was observed. Neuromuscular abnormalities, autonomic instability, somnolence, and a brief generalized tonic-clonic seizure (which resolved after a dose of benzodiazepine) were observed in a 12 months’ old infant who ingested seven or eight 8-mg ondansetron tablets (approximately forty times the recommended 0.1-0.15 mg/kg dose for a paediatric patient). In all instances, the events resolved completely.

Ondansetron prolongs QT interval in a dose-dependent fashion (see [10.2 Pharmacodynamics](#)). ECG monitoring is recommended in cases of overdose.

Cases consistent with serotonin syndrome have been reported in young children following oral overdose (see [7 WARNINGS AND PRECAUTIONS, Serotonin Toxicity/Neuroleptic Malignant Syndrome](#)).

For management of a suspected drug overdose, contact your regional poison control centre.
---

## 6 DOSAGE FORMS, STRENGTHS, COMPOSITION AND PACKAGING

**Table 1 – Dosage Forms, Strengths, Composition and Packaging**

Route of Administration	Dosage Form / Strength/Composition	Non-medicinal Ingredients
Oral	Orally disintegrating tablets /4 mg and 8 mg ondansetron	Aspartame (see <a href="#">7 WARNINGS AND PRECAUTIONS, General</a> ), Basic Butylated Methacrylate Copolymer, Crospovidone (Type A), Flavor Strawberry, Magnesium Stearate, Mannitol, Mannitol granular, Silica colloidal anhydrous

**4 mg Orally Disintegrating Tablets:** Round, white to off-white, flat, bevelled edged tablets, plain on both the surfaces, having strawberry odour. Each tablet contains 4 mg ondansetron.

ACCEL-ONDANSETRON ODT is available in a blister pack of 10 tablets packaged in a carton.

**8 mg Orally Disintegrating Tablets:** Round, white to off-white, flat, bevelled edged tablets, plain on one surface and “8” engraved on opposite surface, having strawberry odour. Each tablet contains 8 mg ondansetron.

ACCEL-ONDANSETRON ODT is available in a blister pack of 10 tablets packaged in a carton.

## 7 WARNINGS AND PRECAUTIONS

### General

ACCEL-ONDANSETRON ODT is not effective in preventing motion-induced nausea and vomiting.

### Cardiovascular

- **QTc Interval Prolongation**

Ondansetron prolongs the QT interval (see [10.2 Pharmacodynamics, Electrocardiography](#)).

The magnitude of QTc prolongation will depend on the peak serum ondansetron concentration (C<sub>max</sub>), which is substantially determined by the route of administration, the dose and the infusion rate of intravenous ondansetron. In addition, post-marketing cases of torsade de pointes have been reported in patients using ondansetron. Torsade de pointes is a polymorphic ventricular tachyarrhythmia. Generally, the risk of torsade de pointes increases with the magnitude of QTc prolongation produced by the drug. Torsade de pointes may be asymptomatic or experienced by the patient as dizziness, palpitations, syncope, or

seizures. If sustained, torsade de pointes can progress to ventricular fibrillation and sudden cardiac death.

Avoid ondansetron in patients with congenital long QT syndrome. Ondansetron should be administered with caution to patients who have or may develop prolongation of QTc, including congestive heart failure, bradyarrhythmias or patients taking other medicinal products that lead to either QT prolongation or electrolyte abnormalities (see [9.4 Drug-Drug Interactions](#)).

Hypokalaemia, hypocalcaemia and hypomagnesaemia should be corrected prior to ondansetron administration.

Additional risk factors for torsade de pointes in the general population include, but are not limited to, the following:

- female gender;
- age 65 years or older;
- baseline prolongation of the QT/QTc interval;
- presence of genetic variants affecting cardiac ion channels or regulatory proteins;
- family history of sudden cardiac death at <50 years;
- cardiac disease (e.g., myocardial ischemia or infarction, left ventricular hypertrophy, cardiomyopathy, conduction system disease);
- history of arrhythmias (especially ventricular arrhythmias, atrial fibrillation, or recent conversion from atrial fibrillation);
- bradycardia (<50 beats per minute);
- acute neurological events (e.g., intracranial or subarachnoid haemorrhage, stroke, intracranial trauma);
- nutritional deficits (e.g., eating disorders, extreme diets);
- diabetes mellitus;
- autonomic neuropathy.

- **Myocardial Ischemia and Coronary Artery Spasm**

Ondansetron can cause coronary artery vasospasm and myocardial ischemia, which may lead to myocardial infarction. In some cases, the symptoms appeared immediately after IV infusion, or shortly after oral administration, including after low doses in patients without significant, known, pre-existing cardiovascular disease or other risk factors. Caution is advised during and after ondansetron administration and close monitoring is recommended in patients with known or suspected ischemic or vasospastic coronary artery disease or other significant underlying cardiovascular disease.

## Driving and Operating Machinery

In psychomotor testing ondansetron does not impair performance or cause sedation.

## Endocrine and Metabolism

- **Phenylketonuria**

ACCEL-ONDANSETRON ODT contains aspartame. Exercise caution when using ACCEL-ONDANSETRON ODT in patients with phenylketonuria.

## Gastrointestinal

As ondansetron is known to increase large bowel transit time, patients with signs of subacute intestinal obstruction should be monitored following administration of ACCEL-ONDANSETRON ODT.

## Hepatic/Biliary/Pancreatic

Abnormal liver function test results have been reported, as well as liver failure in clinical trial cancer patients. See [8.2 Clinical Trial Adverse Reactions, Hepatic/Biliary/Pancreatic](#); and [8.5 Post-Market Adverse Reactions](#).

## Immune

Cross-reactive hypersensitivity has been reported between different 5-HT<sub>3</sub> antagonists. Patients who have experienced hypersensitivity reactions to one 5-HT<sub>3</sub> antagonist have experienced more severe reactions upon being challenged with another drug of the same class. The use of a different 5-HT<sub>3</sub> receptor antagonist is not recommended as a replacement in cases in which a patient has experienced even a mild hypersensitivity type reaction to another 5-HT<sub>3</sub> antagonist.

## Neurologic

- **Serotonin Toxicity /Neuroleptic Malignant Syndrome**

On rare occasions, serotonin toxicity, also known as serotonin syndrome, has been reported with ondansetron, particularly during combined use with other serotonergic drugs (see [9.4 Drug-Drug Interactions](#)).

Serotonin toxicity is characterized by neuromuscular excitation, autonomic stimulation (e.g. tachycardia, flushing) and altered mental state (e.g. anxiety, agitation, hypomania). In accordance with the Hunter criteria, serotonin toxicity diagnosis is likely when, in the presence of at least one serotonergic agent, one of the following is observed:

- Spontaneous clonus
- Inducible clonus or ocular clonus with agitation or diaphoresis
- Tremor and hyperreflexia
- Hypertonia and body temperature > 38°C and ocular clonus or inducible clonus

Neuroleptic malignant syndrome has also been rarely reported with ondansetron, particularly during combined use with neuroleptic/antipsychotic drugs. The clinical manifestations of neuroleptic malignant syndrome often overlap with those of serotonin toxicity, including hyperthermia, hypertonia, altered mental status, and autonomic instability. In contrast to serotonin toxicity, patients with neuroleptic malignant syndrome may present with “lead pipe” muscle rigidity as well as hyporeflexia.

Ondansetron should be used with caution in patients receiving other serotonergic drugs or antipsychotics/neuroleptics. If concomitant treatment with ondansetron, and other serotonergic drugs and/or antipsychotics/neuroleptics is clinically warranted, careful observation of the patient is advised, particularly during treatment initiation and dose increases (see [9.4 Drug-Drug Interactions](#)). Serotonin toxicity and neuroleptic malignant syndrome may result in potentially life-threatening conditions. If serotonin toxicity or neuroleptic malignant syndrome is suspected, discontinuation of ACCEL-ONDANSETRON ODT, should be considered.

## Reproductive Health: Female and Male Potential

Pregnancy status should be verified for females of reproductive potential prior to starting treatment with ACCEL-ONDANSETRON ODT.

Females of reproductive potential should be advised of the possible harm ACCEL-ONDANSETRON ODT can cause to the developing foetus (see [7.1.1 Pregnant Women](#)). Sexually active females of reproductive potential should use effective contraception (methods that result in less than 1% pregnancy rates) when using ACCEL-ONDANSETRON ODT and for two days after stopping treatment with ACCEL-ONDANSETRON ODT.

### 7.1 Special Populations

#### 7.1.1 Pregnant Women

The use of ondansetron during pregnancy is not recommended.

Ondansetron use during early pregnancy has been associated with a small increase in orofacial malformations. Despite some limitations in methodology, several human epidemiological studies have noted an increase in orofacial clefts in infants of women administered ondansetron during the first trimester of pregnancy. Regarding cardiac malformations, the epidemiological studies showed conflicting results.

Ondansetron is not teratogenic in animals (see [16 NON-CLINICAL TOXICOLOGY, Reproductive and Developmental Toxicology](#)).

### 7.1.2 Breast-feeding

Ondansetron is excreted in the milk of lactating rats. It is not known if it is excreted in human milk. However, breast-feeding is not recommended during treatment with ondansetron.

### 7.1.3 Pediatrics

Insufficient information is available to inform dosage recommendations for children 3 years of age or younger. ACCEL-ONDANSETRON ODT is not indicated for use in children less than 4 years of age (see [1.1 Pediatrics \(<18 years of age\)](#)).

### 7.1.4 Geriatrics

Early Phase I studies in healthy elderly volunteers showed a slight age-related decrease in clearance, and an increase in half-life of ondansetron. A greater effect on QTcF is predicted in patients  $\geq 75$  years of age compared to young adults, based on exposure-response modelling. See [10.3 Pharmacokinetics, Geriatrics](#).

## 8 ADVERSE REACTIONS

### 8.1 Adverse Reaction Overview

The most frequent adverse events reported in controlled clinical trials were headache (11%) and constipation (4%). Other adverse events include sensations of flushing or warmth (< 1%).

### 8.2 Clinical Trial Adverse Reactions

Clinical trials are conducted under very specific conditions. The adverse reaction rates observed in the clinical trials may not reflect the rates observed in practice and should not be compared to the rates in the clinical trials of another drug. Adverse drug reaction information from clinical trials is useful for identifying drug-related adverse events and for approximating rates.

Ondansetron has been administered to over 2500 patients worldwide in controlled clinical trials and has been well tolerated.

### Cardiovascular

There have been rare reports of tachycardia, angina (chest pain), bradycardia, hypotension, syncope and electrocardiographic alterations.

**Central Nervous System**

There have been rare reports of seizures. Movement disorders and dyskinesia have been reported in two large clinical trials of ondansetron at a rate of 0.1 – 0.3%.

**Dermatological**

Rash has occurred in approximately 1% of patients receiving ondansetron.

**Eye Disorder**

There have been reports of transient visual disturbances, including blurred vision and, in very rare cases, transient blindness, during or shortly after ondansetron treatment. These were observed generally within the recommended dosing range and predominantly during intravenous administration. The majority of blindness cases resolved within 20 minutes.

**Hepatic/Biliary/Pancreatic**

There were transient increases of SGOT and SGPT of over twice the upper limit of normal in approximately 5% of patients. These increases did not appear to be related to dose or duration of therapy. There have been reports of liver failure and death in patients with cancer receiving concurrent medications including potentially hepatotoxic cytotoxic chemotherapy and antibiotics. The etiology of the liver failure is unclear.

**Hypersensitivity**

Rare cases of immediate hypersensitivity reactions sometimes severe, including anaphylaxis, bronchospasm, urticaria and angioedema have been reported.

**Metabolic**

There have been rare reports of hypokalaemia.

**Other**

There have been reports of abdominal pain, weakness and xerostomia.

**8.5 Post-Market Adverse Reactions**

Over 250 million patient treatment days of ondansetron have been supplied since the launch of the product worldwide. The following events have been spontaneously reported during

post-approval use of ondansetron, although the link to ondansetron cannot always be clearly established.

The adverse event profiles in children and adolescents were comparable to that seen in adults.

### **Cardiovascular Disorders**

There have been rare reports (< 0.01%) of myocardial infarction, myocardial ischemia, angina, chest pain with or without ST segment depression, arrhythmias (including ventricular or supraventricular tachycardia, premature ventricular contractions, and atrial fibrillation), electrocardiographic alterations (including second degree heart block), palpitations and syncope.

Rarely and predominantly with intravenous ondansetron, transient ECG changes including QTc interval prolongation, Torsade de Pointes, ventricular fibrillation, coronary artery spasm, myocardial ischemia, cardiac arrest, and sudden death have been reported (see [7 WARNINGS AND PRECAUTIONS, Cardiovascular](#)).

### **Eye Disorder**

There have been very rare cases of transient blindness following ondansetron treatment, generally within the recommended dosing range and predominantly during intravenous administration.

The majority of blindness cases reported resolved within 20 minutes. Although most patients had received chemotherapeutic agents, including cisplatin a few cases of transient blindness occurred following ondansetron administration for the treatment of post-operative nausea or vomiting and in the absence of cisplatin treatment. Some cases of transient blindness were reported as cortical in origin.

### **Hepatic/ Biliary / Pancreatic**

Occasional asymptomatic increases in liver function tests have been reported.

### **Immune Disorders**

Rare cases of hypersensitivity reactions, sometimes severe (e.g., laryngeal oedema, stridor, laryngospasm and cardiopulmonary arrest) have also been reported.

### **Nervous System Disorders**

Transient episodes of dizziness (< 0.1%) have been reported predominantly during or upon completion of IV infusion of ondansetron.



Uncommon reports (< 1%) suggestive of extrapyramidal reactions including oculogyric crisis/dystonic reactions (e.g., oro-facial dyskinesia, opisthotonos, tremor, etc.), movement disorders and dyskinesia have been reported without definitive evidence of persistent clinical sequelae.

Serotonin syndrome and neuroleptic malignant syndrome-like events have been reported with 5-HT<sub>3</sub> receptor antagonist antiemetics, including ondansetron, when given in combination with other serotonergic and/or neuroleptic drugs (see [7 WARNINGS AND PRECAUTIONS, Neurologic](#)).

### **Respiratory, Thoracic and Mediastinal Disorders**

There have also been rare reports of hiccups.

### **Skin and Subcutaneous Tissue Disorders**

Very rare reports have been received for bullous skin and mucosal reactions, including fatal cases. These reports include toxic skin eruptions, such as Stevens-Johnson syndrome and toxic epidermal necrolysis, and have occurred in patients taking other medications that can be associated with bullous skin and mucosal reactions.

## **9 DRUG INTERACTIONS**

### **9.1 Serious Drug Interactions**

Serious Drug Interactions (see [9.4 Drug-Drug Interactions](#))

- Apomorphine (see [2 CONTRAINDICATIONS](#))
- QTc-Prolonging drugs
- Serotonergic agents

### **9.2 Drug Interactions Overview**

Ondansetron is extensively metabolised by multiple hepatic cytochrome P450 enzymes (predominantly CYP3A4, also CYP2D6 and CYP1A2), and clearance is reduced in hepatic insufficiency (see [10.3 Pharmacokinetics, Hepatic Insufficiency](#)).

CYP 3A4 inducers can increase ondansetron clearance (see [9.4. Drug-Drug Interactions, CYP 3A4 Inducers](#)).

Ondansetron does not, itself, appear to induce or inhibit the cytochrome P450 drug-metabolizing enzyme system of the liver.

### **9.3 Drug-Behavioural Interactions**

Potential interactions, in terms of individual behavioural risks, have not been established.

## 9.4 Drug-Drug Interactions

### CYP 3A4 Inducers

Patients treated with inducers of CYP3A4 (i.e. phenytoin, carbamazepine, and rifampicin) demonstrated an increase in clearance of oral ondansetron and a decrease in ondansetron blood concentrations.

In a pharmacokinetic study of 16 epileptic patients maintained chronically on carbamazepine or phenytoin (CYP 3A4 inducers), reduction in AUC,  $C_{max}$  and  $T_{1/2}$  of ondansetron was observed. This resulted in a significant increase in clearance. However, no dosage adjustment can be recommended, due to inter-subject variability in the available data.

### Cytochrome P450 Inhibitors

No effect on ondansetron clearance secondary to enzyme inhibition or reduced activity (e.g. CYP2D6 genetic deficiency) has been identified, to date.

### QTc-Prolonging Drugs

The concomitant use of ondansetron with another QTc-prolonging drug should be carefully considered to determine that the therapeutic benefit outweighs the potential risk (see [7 WARNINGS AND PRECAUTIONS, Cardiovascular](#); [9.1 Serious Drug Interactions](#)). Drugs that have been associated with QTc interval prolongation and/or torsade de pointes include, but are not limited to, the examples in the following list. Chemical/pharmacological classes are listed if some, although not necessarily all, class members have been implicated in QTc prolongation and/or torsade de pointes:

- Class IA antiarrhythmics (e.g. quinidine, procainamide, disopyramide)
- Class III antiarrhythmics (e.g. amiodarone, sotalol, ibutilide, dronedarone)
- Class 1C antiarrhythmics (e.g. flecainide, propafenone)
- antiemetics (e.g. dolasetron, palonosetron, granisetron, droperidol, chlorpromazine, prochlorperazine)
- tyrosine kinase inhibitors (e.g. vandetanib, sunitinib, nilotinib, lapatinib)
- antipsychotics (e.g. chlorpromazine, pimozide, haloperidol, ziprasidone)
- antidepressants (e.g. citalopram, fluoxetine, venlafaxine, tricyclic/tetracyclic, amitriptyline, imipramine, maprotiline)
- opioids (e.g. methadone;
- domperidone
- macrolide antibiotics and analogues (e.g. erythromycin, clarithromycin, telithromycin, tacrolimus)
- quinolone antibiotics (e.g. moxifloxacin, levofloxacin, ciprofloxacin)
- antimalarials (e.g. quinine, chloroquine)
- azole antifungals (e.g. ketoconazole, fluconazole, voriconazole)

- histone deacetylase inhibitors (e.g. vorinostat)
- beta-2 adrenoceptor agonists (e.g. salmeterol, formoterol)

### Drugs that Cause Electrolyte Abnormalities

The use of ondansetron with drugs that can disrupt electrolyte levels should be avoided. Such drugs include, but are not limited to, the following:

- loop, thiazide, other related diuretics
- laxatives and enemas
- amphotericin B
- high dose corticosteroids

### Tramadol

Data from small studies indicate that ondansetron may reduce the analgesic effect of tramadol.

### Apomorphine

Based on reports of profound hypotension and loss of consciousness when ondansetron was administered with apomorphine hydrochloride, concomitant use with apomorphine is contraindicated (see [2 CONTRAINDICATIONS](#); [9.1 Serious Drug Interactions](#)).

### Serotonergic Drugs

As with other serotonergic agents, serotonin syndrome/toxicity, a potentially life-threatening condition, may occur with 5-HT<sub>3</sub> receptor antagonist antiemetic treatment when given in combination with other agents that may affect the serotonergic neurotransmitter system, including triptans, selective serotonin reuptake inhibitors (SSRIs), serotonin and norepinephrine reuptake inhibitors (SNRIs), tricyclic antidepressants (TCAs), other 5-HT<sub>3</sub> receptor antagonists, lithium, sibutramine, fentanyl and its analogues, dextromethorphan, tramadol, tapentadol, meperidine, methadone, pertazocine, St. John's wort (*Hypericum perforatum*) and MAOIs, including phenelzine, moclobemide, tranylcypromine, linezolid (an antibiotic which is a reversible non-selective MAOI) and methylene blue (see [7 WARNINGS AND PRECAUTIONS, Neurologic](#); [9.1 Serious Drug Interactions](#)).

### Antipsychotics/Neuroleptics

Neuroleptic malignant syndrome has been rarely reported with ondansetron, particularly during combined use with neuroleptic/antipsychotic drugs (e.g. haloperidol, olanzapine, quetiapine, risperidone) (see [7 WARNINGS AND PRECAUTIONS, Neurologic](#)).

The above list of potential drug interactions is not comprehensive. Current information sources should be consulted for newly approved drugs that prolong the QTc interval, affect the serotonergic system, increase CYP 3A4 enzyme activity, or cause electrolyte disturbances, as well as for older drugs for which these effects have recently been established.

### **9.5 Drug-Food Interactions**

Interactions with food have not been established.

### **9.6 Drug-Herb Interactions**

Interactions with herbal products have not been established.

Exercise caution when using ondansetron with herbal products that may alter serotonin levels or act as CYP3A4 inducers/inhibitors, such as St. John's wort, ginseng, L-tryptophan, valerian, and Ginkgo biloba.

### **9.7 Drug-Laboratory Test Interactions**

Interactions with laboratory tests have not been established.

## **10 CLINICAL PHARMACOLOGY**

### **10.1 Mechanism of Action**

ACCEL-ONDANSETRON ODT (ondansetron) is a selective antagonist of the serotonin receptor subtype, 5-HT<sub>3</sub>. Its precise mode of action in the control of chemotherapy induced nausea and vomiting is not known.

Cytotoxic chemotherapy and radiotherapy are associated with the release of serotonin (5-HT) from enterochromaffin cells of the small intestine, presumably initiating a vomiting reflex through stimulation of 5-HT<sub>3</sub> receptors located on vagal afferents. Ondansetron may block the initiation of this reflex. Activation of vagal afferents may also cause a central release of serotonin from the chemoreceptor trigger zone of the area postrema, located on the floor of the fourth ventricle. Thus, the antiemetic effect of ondansetron is probably due to the selective antagonism of 5-HT<sub>3</sub> receptors on neurons located in either the peripheral or central nervous systems, or both.

The mechanisms of ondansetron's antiemetic action in post-operative nausea and vomiting are not known.

### **10.2 Pharmacodynamics**

Serotonin receptors of the 5-HT<sub>3</sub> type are present both peripherally and on vagal nerve terminals. Ondansetron probably acts by preventing activation of these receptors or receptors located in other regions of the central nervous system. Both the peripheral and central nervous systems appear to be involved since both abdominal vagotomy and microinjection of ondansetron and other 5-HT<sub>3</sub> antagonists directly into the area postrema eliminate cisplatin-

induced emesis, while 5-HT<sub>1</sub>-like (methiothepin maleate) and 5-HT<sub>2</sub> (ketanserin) antagonists have no effect.

Ondansetron is highly selective for 5-HT<sub>3</sub> receptors and shows negligible binding to other receptors such as 5-HT<sub>1</sub>-like, 5-HT<sub>2</sub>,  $\alpha$ <sub>1</sub> and  $\alpha$ <sub>2</sub> adrenoceptors,  $\beta$ <sub>1</sub> and  $\beta$ <sub>2</sub> adrenoceptors, D<sub>1</sub> and D<sub>2</sub> muscarinic, nicotinic, GABA<sub>A</sub>, H<sub>1</sub> and H<sub>2</sub> receptors.

The pharmacological specificity of ondansetron may explain the observed lack of extrapyramidal side effects often seen following similar therapy with metoclopramide, which preferentially binds to dopamine receptors of the D<sub>2</sub> subtype.

*In vivo* pharmacodynamic studies have investigated the effects of ondansetron on gastric emptying, small bowel transit time and oesophageal motility.

Both oral (16 mg tid) and intravenous (5-10 mg) doses of ondansetron failed to produce a significant effect on gastric emptying in both healthy volunteers and in patients suffering from delayed gastric emptying. However, in one study intravenous doses of 8 mg did increase gastric emptying in over half the volunteers tested.

Intravenous infusion of either 1 mg or 5 mg ondansetron tended to increase small bowel transit times and single intravenous doses of 10 mg ondansetron have been reported to decrease sphincter pressure in the lower oesophagus in some subjects.


### **Electrocardiography**

A study in cloned human cardiac ion channels has shown ondansetron has the potential to affect cardiac repolarisation via blockade of hERG potassium channels at clinically relevant concentrations. Dose-dependent QT prolongation has been observed in a thorough QT study in human volunteers

The effect of ondansetron on the QT<sub>c</sub> interval was evaluated in a double blind, randomized, placebo and positive (moxifloxacin) controlled, crossover study in 58 healthy adult men and women. Ondansetron was tested at single doses of 8 mg and 32 mg infused intravenously over 15 minutes. At the highest tested dose of 32 mg, prolongation of the Fridericia-corrected QT<sub>c</sub> interval ( $QT/RR^{0.33}=QTcF$ ) was observed from 15 min to 4 h after the start of the 15 min infusion, with a maximum mean (upper limit of 90% CI) difference in QT<sub>c</sub>F from placebo after baseline-correction of 19.6 (21.5) msec at 20 min. At the lower tested dose of 8 mg, QT<sub>c</sub> prolongation was observed from 15 min to 1 h after the start of the 15 minute infusion, with a maximum mean (upper limit of 90% CI) difference in QT<sub>c</sub>F from placebo after baseline-correction of 5.8 (7.8) msec at 15 min. The magnitude of QT<sub>c</sub> prolongation with ondansetron is expected to be greater if the infusion rate is faster than 15-minutes. The 32 mg intravenous dose of ondansetron must not be administered.

No treatment-related effects on the QRS duration or the PR interval were observed at either the 8 or 32 mg dose.

### LS Mean Difference (90% CI) in QTcF Interval Between Treatment and Placebo Over Time



An ECG assessment study has not been performed for orally administered ondansetron. On the basis of pharmacokinetic-pharmacodynamic modelling, an 8 mg oral dose of ondansetron is predicted to cause a mean QTcF increase of 0.7 ms (90% CI -2.1, 3.3) at steady-state, assuming a mean maximal plasma concentration of 24.7 ng/mL (95% CI 21.1, 29.0).

The magnitude of QTc prolongation at the recommended 5 mg/m<sup>2</sup> dose in pediatrics has not been studied, but pharmacokinetic-pharmacodynamic modelling predicts a mean increase of 6.6 ms (90% CI 2.8, 10.7) at maximal plasma concentrations.

### 10.3 Pharmacokinetics

#### Absorption

Pharmacokinetic studies in human volunteers showed peak plasma levels of 20-30 ng/mL at around 1½ hours after an 8 mg oral dose of ondansetron. Repeat dosing of an 8 mg tablet every 8 hours for 6 days increased the peak plasma value to 40 ng/mL.

#### Distribution

The absolute bioavailability of ondansetron in humans was approximately 60% and the plasma protein binding was approximately 73%.

#### Metabolism

*In vitro* metabolism studies have shown that ondansetron is a substrate for human hepatic cytochrome P<sub>450</sub> enzymes, including CYP1A2, CYP2D6 and CYP3A4. In terms of overall

ondansetron turnover, CYP3A4 played the predominant role. Because of the multiplicity of metabolic enzymes capable of metabolising ondansetron, it is likely that inhibition or loss of one enzyme (e.g. CYP2D6 enzyme deficiency) will be compensated by others and may result in little change in overall rates of ondansetron clearance. CYP 3A4 inducers can increase clearance (see [9.4 Drug-Drug Interactions, CYP 3A4 inducers](#)).

## Elimination

Following extensive metabolism of an orally or intravenously administered dose, ondansetron is excreted in the urine and faeces. In humans, less than 10% of the dose is excreted unchanged in the urine. The major urinary metabolites are glucuronide conjugates (45%), sulphate conjugates (20%) and hydroxylation products (10%).

The half-life of ondansetron after either an 8 mg oral dose or intravenous dose was approximately 3-4 hours and may be extended to 6-8 hours in the elderly.

## Special Populations and Conditions

- **Geriatrics**

Early Phase I studies in healthy elderly volunteers showed a slight age-related decrease in clearance, and an increase in half-life of ondansetron. However, wide inter-subject variability resulted in considerable overlap in pharmacokinetic parameters between young (< 65 years of age) and elderly subjects (≥ 65 years of age) and there were no overall differences in safety or efficacy observed between young and elderly cancer patients enrolled in CINV clinical trials (see [4.2 Recommended Dose and Dosage Adjustment, Geriatrics](#)).

There is limited data involving patients ≥75 years of age. Based on more recent ondansetron plasma concentrations and exposure-response modelling, a greater effect on QTcF is predicted in patients ≥75 years of age compared to young adults. (See [4.2 Recommended Dose and Dosage Adjustment, Geriatrics](#))

- **Genetic Polymorphism**

**CYP 2D6:** The elimination half-life and plasma levels of a single 8 mg intravenous dose of ondansetron did not differ between subjects classified as poor and extensive metabolisers of sparteine and debrisoquine (CYP 2D6 substrates). No alteration of daily dosage or frequency of ondansetron dosing is recommended for patients known to be CYP 2D6 poor metabolisers.

- **Hepatic Insufficiency**

Ondansetron is extensively metabolized by the liver. The clearance of an 8 mg intravenous dose of ondansetron was significantly reduced and the serum half-life significantly prolonged in subjects with severe impairment of hepatic function. In

patients with moderate or severe impairment of hepatic function, reductions in dosage are therefore recommended (see [4.2 Recommended Dose and Dosage Adjustment](#)).

There is no experience in patients who are clinically jaundiced.

- **Renal Insufficiency**

Renal impairment is not expected to significantly influence the total clearance of ondansetron, as renal clearance represents only 5% of the overall clearance. No dosage adjustment is required in patients with impaired renal function (see [4.2 Recommended Dose and Dosage Adjustment](#)).

## **11 STORAGE, STABILITY AND DISPOSAL**

ACCEL-ONDANSETRON ODT (ondansetron) orally disintegrating tablets should be stored at 15 - 30°C. Keep in a safe place out of reach and sight of children. There is no specific instruction for disposal.

## **12 SPECIAL HANDLING INSTRUCTIONS**

No special handling instructions are required for this drug product.



**PART II: SCIENTIFIC INFORMATION****13 PHARMACEUTICAL INFORMATION****Drug Substance**


Proper name: ondansetron

Chemical name: 1,2,3,9-tetrahydro-9-methyl-3-[(2-methyl-1H-imidazol-1-yl) methyl]-4H carbazol-4-one.

Molecular Formula: C<sub>18</sub>H<sub>19</sub>N<sub>3</sub>O

Molecular mass: 293.4 g/mol

Structural formula:



Description and Solubility:

Ondansetron is a white to off-white powder. Literature indicates that several crystalline polymorphs of ondansetron (base) exist. Ondansetron is very soluble in acid solutions, slightly soluble in acetone, isopropyl alcohol, methanol and practically insoluble in water.

**14 CLINICAL TRIALS****14.1 Clinical Trials by Indication****Clinical trial data**

The clinical trial data on which the original indication was authorized is not available.

Clinical trial results showing the number and percentage of patients exhibiting a complete response to ondansetron (0 emetic episodes) are shown in the tables below for both post-operative and chemotherapy induced emesis.

### Prevention of Chemotherapy Induced Emesis

**Table 2 - Prevention of Chemotherapy Induced Emesis - Response Over 24 Hours<sup>1</sup>**

<b>Dose</b>	<b>Ondansetron 3 doses of 0.15 mg/kg</b>	<b>Placebo 3 doses of placebo</b>
# of patients	14	14
Treatment Response:		
0 emetic episodes	2 (14%)	0 (0%)
1 – 2 emetic episodes	8 (57%)	0 (0%)
<sup>1</sup> Results are from an initial study using a different dosing regimen.		

### Prevention of Post-Operative Emesis

**Table 3 - Prevention of Post-Operative Emesis – Response Over 24 Hours<sup>1</sup>**

<b>Dose</b>	<b>Oral Prevention</b>		
	<b>Ondansetron 16 mg od</b>	<b>Placebo</b>	<b><i>p value</i></b>
# of patients	253	250	
Treatment Response: 0 emetic episodes	126 (50%)	79 (32%)	< 0.001
<sup>1</sup> The majority of patients included in the prevention and treatment of post-operative nausea and vomiting studies using ondansetron have been adult women receiving balanced anaesthesia for gynaecological surgery.			

## Prevention of Radiotherapy Induced Emesis

**Table 4 - Prevention of Radiotherapy Induced Emesis - Response Over 24 Hours<sup>1</sup>**

Oral Treatment			
Dose	Ondansetron 8 mg PO tid <sup>2</sup>	Metoclopramide 10 mg PO tid <sup>2</sup>	p value
# of patients	38	44	
Treatment Response: 0 emetic episodes	37 (97%)	20 (45%)	< 0.001
<p><sup>1</sup> results from a study of adult male and female patients receiving single high dose radiotherapy (800 to 1,000 cGy) over an anterior or posterior field size of <math>\geq 80</math> cm<sup>2</sup> to the abdomen.</p> <p><sup>2</sup> Patients received the first dose of ondansetron 8 mg tablets or metoclopramide (10 mg) 1-2 hours before radiotherapy. If radiotherapy was given in the morning, 2 additional doses of study treatment were given (1 tablet late afternoon and 1 tablet before bedtime). If radiotherapy was given in the afternoon, patients took only 1 further tablet that day before bedtime. Patients continued oral medication on a 3 times a day basis for 3-5 days.</p>			

### 14.2 Comparative Bioavailability Studies

A double blind, balanced, randomized, single-dose, two-treatment, two-sequence, two-period, crossover, oral comparative bioavailability study comparing ACCEL-ONDANSETRON ODT 8 mg Orally Disintegrating Tablets (Accel Pharma Inc.) and ZOFTRAN<sup>®</sup> ODT 8 mg tablets (Novartis Pharmaceuticals Canada Inc.) was conducted in healthy, adult, male, human subjects under fasting conditions. Comparative bioavailability data from 24 subjects that were included in the statistical analysis are presented in the following table:

**Comparative bioavailability data from 24 subjects**

Ondansetron (1 x 8 mg) Geometric Mean Arithmetic Mean (CV %)				
<b>Parameter</b>	<b>Test<sup>1</sup></b>	<b>Reference<sup>2</sup></b>	<b>% Ratio of Geometric Means</b>	<b>Confidence Interval (90%)</b>
AUC <sub>T</sub> (ng·h/mL)	316.40 339.76 (39.49)	299.28 323.01 (40.59)	105.7	98.8 - 113.2
AUC <sub>I</sub> (ng·h/mL)	339.24 371.13 (45.21)	324.10 358.76 (49.28)	104.7	96.7 – 113.3
C <sub>max</sub> (ng/mL)	44.63 46.59 (30.09)	42.98 44.70 (26.58)	103.8	98.4 - 109.6
T <sub>max</sub> <sup>3</sup> (h)	2.00 (1.00 - 4.00)	1.88 (0.75 - 4.00)		
T <sub>½</sub> <sup>4</sup> (h)	5.62 (35.49)	5.90 (33.91)		
<sup>1</sup> ACCEL-ONDANSETRON ODT (ondansetron) Orally Disintegrating Tablets, 8 mg (Accel Pharma Inc.) <sup>2</sup> ZOFTRAN® ODT (ondansetron) Orally Disintegrating Tablets, 8 mg (Novartis Pharmaceuticals Canada Inc.) <sup>3</sup> Expressed as the median (range) only <sup>4</sup> Expressed as the arithmetic mean (CV%) only				

**15 MICROBIOLOGY**

No microbiological information is required for this product.

**16 NON-CLINICAL TOXICOLOGY****General Toxicology**

- Acute Toxicity**

Single doses of ondansetron up to the LD<sub>50</sub> in mice and in rats were generally well tolerated. Reactions, including tremor and convulsive behaviour, occurred only at near lethal levels.

**Table 5 - Acute Toxicity**

Species	LD <sub>50</sub> (mg/kg)
	Oral
Mice	10-30
Rats	100-150

All deaths resulted from the acute effects of treatment, the observed clinical signs being consistent with the central nervous system effects associated with behavioural depression. These effects were not associated with any apparent histopathological changes in the brain. No target organ toxicity was identified.

**Long-term Toxicity****Table 6 - Subacute Toxicity Studies**

Species	Route	Dose (mg/kg/day)	Duration of Study	Results
Rats	Oral	160	7 weeks	Well-tolerated
	IV	12	5 weeks	Well-tolerated
Dogs	Oral	7.5-25	5 weeks	Transient post-dosing clinical reactions associated with behavioural depression (at highest dose levels)
	IV	2-8	5 weeks	

Maximum daily dose levels in rats were found to be higher when doses were gradually increased. Identical doses were rapidly lethal to rats not previously exposed to ondansetron. Post-dosing reactions, in both rats and dogs, included ataxia, exophthalmia, mydriasis, tremor and respiratory changes. Increases in liver enzymes (SGPT and SGOT) were noted at high dose levels. Dogs dosed at 6.75 mg/kg/day intravenously exhibited vein irritancy in the form of constriction and thickening, creating resistance to needle penetration. The changes were noted after seven days treatment but were reversed by decreasing the dose concentration.

**Table 7 - Chronic Toxicity**

Species	Duration	Max. no-effect Dose (mg/kg/day)	Effects
Rat	18 months	1	Usually transient and restricted to highest dose

Species	Duration	Max. no-effect Dose (mg/kg/day)	Effects
Dog	12 months	12	

### Genotoxicity

No evidence of mutagenicity was observed in microbial mutagen tests using mutant strains of *Salmonella typhimurium*, *Escherichia coli* or *Saccharomyces cerevisiae*, with or without a rat-liver post-mitochondrial metabolizing system.

There was also no evidence of damage to genetic material noted in in vitro V-79 mammalian cell mutation studies, in vitro chromosome aberration tests using human peripheral lymphocytes, or *in vivo* chromosome aberration assays in mouse bone marrow.

### Carcinogenicity

**Table 8 - Carcinogenicity Studies**

Species	Route	Dose (mg/kg/day)	Duration of Study	Results
Mice	Oral	1-40 (max. oral dose 30)	2 years	No treatment related increases in tumour incidence.
Rats	Oral	1-25 (max. oral dose 10)	2 years	Proportion of benign/malignant tumours also remained Consistent with the pathological background of the Animals studied.

There was no evidence of a tumourigenic effect of ondansetron in any tissue.

### Reproductive and Developmental Toxicology

Ondansetron was not teratogenic in rats and rabbits at dosages up to the maximum non-convulsive level, (rat: 15 mg/kg/day, rabbit: 30 mg/kg/day; the maternal dose was approximately 6 and 24 times the maximum recommended human oral dose of 24 mg/day, respectively, based on body surface area). No adverse effects on pregnancy or foetal and post-natal development were detected in rats and no foetal abnormalities were observed in rabbits after oral administration of ondansetron.

A slight maternal toxicity was observed at the highest dose level in intravenous organogenesis (4.0 mg/kg/day) studies in the rabbit. Effects included maternal body weight loss and

increased incidence of early foetal death. In a rat fertility study, there was a dose-related decrease in the proportion of surviving pups of the F2 generation; however, the significance of this is unclear.

Administration of ondansetron to pregnant rats and rabbits indicated there was foetal exposure to low levels of ondansetron and its metabolites. Ondansetron is retained in the foetal eye presumably bound to melanin. In rats, the transfer of ondansetron and its metabolites into breast milk was extensive. The concentration of unchanged ondansetron in breast milk was higher than in corresponding plasma samples.

Daily administration of ondansetron at dosages up to 15 mg/kg/day to pregnant rats (a maternal dose of approximately 6 times the maximum recommended human oral dose of 24 mg/day, based on body surface area) from day 17 of pregnancy to litter day 22 had no effects on pregnancy of the parental generation or on post-natal development and mating of the F1 generation. Foetal development of the F2 generation was comparable to controls; however, the number of implantations and viable foetuses was reduced in the highest dosage group when compared with controls.

## **17 SUPPORTING PRODUCT MONOGRAPHS**

1. Zofran ODT Orally Disintegrating Tablets, 4 mg and 8 mg, submission control 278769, Product Monograph, Sandoz Canada Inc. (February 27, 2024).

## **PATIENT MEDICATION INFORMATION**

### **READ THIS FOR SAFE AND EFFECTIVE USE OF YOUR MEDICINE**

#### **Pr ACCEL-ONDANSETRON ODT**

#### **ondansetron orally disintegrating tablets, Mfr. Std.**

Read this carefully before you start taking **ACCEL-ONDANSETRON ODT** and each time you get a refill. This leaflet is a summary and will not tell you everything about this drug. Talk to your healthcare professional about your medical condition and treatment and ask if there is any new information about **ACCEL-ONDANSETRON ODT**.

#### **What is ACCEL-ONDANSETRON ODT used for?**

##### **Children (4 to 17 years of age):**

ACCEL-ONDANSETRON ODT (ondansetron) is used to treat nausea and vomiting during certain types of chemotherapy.

##### **Adults:**

ACCEL-ONDANSETRON ODT is used:

- to prevent nausea and vomiting during certain types of chemotherapy and radiotherapy, and
- to prevent nausea and vomiting after surgery.

##### **Patients (65 years of age and older):**

ACCEL-ONDANSETRON ODT is used to prevent nausea and vomiting during chemotherapy and radiotherapy.

#### **How does ACCEL-ONDANSETRON ODT work?**

ACCEL-ONDANSETRON ODT is a medication known as an antiemetic.

Treatments such as cancer chemotherapy and radiotherapy are associated with the release of a natural substance (serotonin). The release of serotonin can make you feel sick and vomit. The way that ACCEL-ONDANSETRON ODT works is not known, but it is thought to help stop the effects of serotonin to reduce the effects of nausea and vomiting.

#### **What are the ingredients in ACCEL-ONDANSETRON ODT?**

- Medicinal ingredient: ondansetron



- Non-medicinal ingredients: aspartame, basic butylated methacrylate copolymer, crospovidone (Type A), flavor strawberry, magnesium stearate, mannitol, mannitol granular, silica colloidal anhydrous.

**ACCEL-ONDANSETRON ODT comes in the following dosage forms:**

Orally disintegrating tablets: 4 mg and 8 mg

**Do not use ACCEL-ONDANSETRON ODT if:**

- you are allergic to ondansetron or to any of the other ingredients in ACCEL-ONDANSETRON ODT.
- you are taking a medicine called apomorphine (used to treat Parkinson's disease).

**To help avoid side effects and ensure proper use, talk to your healthcare professional before you take ACCEL-ONDANSETRON ODT. Talk about any health conditions or problems you may have, including if you:**

- have had an allergic reaction to medicines that are similar to ACCEL-ONDANSETRON ODT such as medicines containing granisetron or palonosetron.
- are pregnant or planning to become pregnant. ACCEL-ONDANSETRON ODT is not recommended for use during pregnancy.
- are breast feeding or planning to breastfeed. ACCEL-ONDANSETRON ODT can pass into your breast milk and affect your baby.
- have liver problems.
- have signs of intestinal obstruction or blockage.
- have or have had heart or blood vessel problems, including if you are at a higher risk for these problems. Risk factors include, but are not limited to, if you:
  - o have family members who have or have had heart or blood vessel problems,
  - o smoke,
  - o have high blood pressure,
  - o have high cholesterol levels,
  - o have diabetes, or
  - o are overweight.
- have a condition called phenylketonuria, as ACCEL-ONDANSETRON ODT contains aspartame.
- are taking medications that affect the serotonin in your body (e.g., serotonergic and neuroleptic medications). If you are unsure, ask your healthcare professional.
- have QT/QTc prolongation (a heart rhythm condition) or a family history of QT/QTc prolongation.
- are taking medications that may lead to QT/QTc prolongation or electrolyte imbalances. If you are unsure, ask your healthcare professional.

- have low blood levels of potassium, magnesium, or calcium.

**Other warnings you should know about:**

**Serotonin toxicity (also known as Serotonin syndrome):** ACCEL-ONDANSETRON ODT can cause serotonin toxicity, a rare but potentially life-threatening condition. It can cause serious changes in how your brain, muscles and digestive system work. You may develop serotonin toxicity if you take ACCEL-ONDANSETRON ODT with certain anti-depressants or migraine medications.

Serotonin syndrome symptoms include:

- fever, sweating, shivering, diarrhea, nausea, vomiting;
- muscle shakes, jerks, twitches or stiffness, overactive reflexes, loss of coordination;
- fast heartbeat, changes in blood pressure;
- confusion, agitation, restlessness, hallucinations, mood changes, unconsciousness, and coma.

**Myocardial ischemia (lack of blood flow to the heart):** Treatment with ACCEL-ONDANSETRON ODT can cause myocardial ischemia which can lead to a heart attack. This may happen shortly after ACCEL-ONDANSETRON ODT administration. Some symptoms of myocardial ischemia can include sudden chest pain, pressure or discomfort, feeling faint, feeling anxious, shortness of breath, irregular heartbeat, nausea, and sudden heavy sweating. Your healthcare professional will monitor your health during and after administration of ACCEL-ONDANSETRON ODT. However, if you notice any symptoms of myocardial ischemia, tell your healthcare professional right away. They may reduce or stop your treatment, and may recommend another therapy.

**QT/QTc prolongation:** ACCEL-ONDANSETRON ODT can affect the electrical activity of your heart known as QT/QTc prolongation. This effect can be measured with an electrocardiogram (ECG). In rare cases, QT/QTc prolongation can cause changes to the rhythm of your heart (e.g., fast, slow or irregular heartbeats). This can lead to dizziness, palpitations (sensation of rapid, pounding, or irregular heart beat), fainting, or death. You are at a higher risk if you have a heart disease, are taking certain interacting medicines, are a female, or are over the age of 65 years. It is important to follow the instructions of your healthcare professional with regard to dosing or any special tests. If you experience any symptoms of a possible heart rhythm problem, you should seek immediate medical attention.

**Severe allergic reactions:** ACCEL-ONDANSETRON ODT can cause allergic reactions in certain individuals. Symptoms of a severe allergic reaction can include wheezing, sudden chest pain, tightness of the chest, heart throbbing, swelling of eyelids, face or lips, or develop a skin rash, skin lumps or hives. If you notice any signs of a severe allergic reaction, **contact your healthcare professional immediately. Do not take any more medicine unless your healthcare professional tells you to do so.**

**Pregnancy:**

- If you are pregnant, there are specific risks for your unborn baby that you must discuss with your healthcare professional.
- If you are able to get pregnant, you may be asked to take a pregnancy test before starting your treatment with ACCEL-ONDANSETRON ODT.
- You should use effective birth control while you are taking ACCEL-ONDANSETRON ODT, and for at least 2 days after stopping ACCEL-ONDANSETRON ODT. Ask your healthcare professional about options of effective birth control.
- If you become pregnant while taking ACCEL-ONDANSETRON ODT, tell your healthcare professional right away.

**Tell your healthcare professional about all the medicines you take, including any drugs, vitamins, minerals, natural supplements or alternative medicines.**

**Serious Drug Interactions**

- **Do not** take ACCEL-ONDANSETRON ODT if you are taking apomorphine (a medicine used to treat Parkinson's Disease). This can cause serious side effects such as extremely high blood pressure and loss of consciousness.
- QTc- Prolonging medicines (see examples below)
- Serotonergic medicines (see examples below)

**The following may interact with ACCEL-ONDANSETRON ODT:**

- medicines called CYP3A4 inducers (e.g., phenytoin, carbamazepine, and rifampicin);
- medicines that can affect electrolyte levels (e.g., diuretics, laxatives, enemas, amphotericin B, and high doses of corticosteroids);
- herbal products (e.g. St. John's Wort (*Hypericum perforatum*), methylene ginseng, L-tryptophan, valerian and Ginkgo biloba)

## QTc-Prolonging medicines:

- medicines used to treat heart rhythm disorders (e.g., quinidine, procainamide, disopyramide, amiodarone, sotalol, ibutilide, dronedarone, flecainide, and propafenone);
- medicines used to treat vomiting and nausea called antiemetics (e.g., dolasetron, palonosetron, granisetron, droperidol, chlorpromazine, prochlorperazine, and domperidone);
- medicines called tyrosine kinase inhibitors (e.g., vandetanib, sunitinib, nilotinib, and lapatinib);

- medicines used to manage psychosis or schizophrenia called antipsychotics (e.g., chlorpromazine, pimozide, haloperidol, olanzapine, quetiapine, risperidone and ziprasidone);
- medicines used to treat depression called antidepressants (e.g., citalopram, fluoxetine, venlafaxine, tricyclic/tetracyclic antidepressants, amitriptyline, imipramine, and maprotiline);
- medicines used to treat pain called opioids (e.g., methadone and tramadol);
- medicines used to treat bacterial infections called antibiotics (e.g., erythromycin, clarithromycin, telithromycin, tacrolimus, moxifloxacin, levofloxacin, and ciprofloxacin);
- medicines used to treat malaria called antimalarials (e.g., quinine and chloroquine);
- medicines used to treat fungal infections called azole antifungals (e.g., ketoconazole, fluconazole, and voriconazole);
- medicines used to treat cancer (e.g., vorinostat);
- medicines called beta-2 adrenoceptor agonists (e.g., salmeterol and formoterol);

#### Serotonergic medicines:

- medicines used to treat migraines (e.g. triptans)
- medicines used to treat depression (e.g. Selective Serotonin-Reuptake Inhibitors (SSRIs), Serotonin Noradrenalin Reuptake Inhibitors (SNRIs))
- medicines used to treat mood disorders (e.g.lithium)
- medicines used for weight loss (e.g. sibutramine)
- medicines used to treat pain (e.g. fentanyl and its analogues, tramadol, tapentadol, methadone, meperidine, pentazocine)
- medicines used to relieve cough cause by colds (e.g. dextromethorphan)
- monoamine oxidase inhibitors (MAOIs) including phenelzine, moclobemide, linezolid, and methylene blue).

If you are unsure about any medications you are taking, ask your healthcare professional.

#### How to take ACCEL-ONDANSETRON ODT:

- Take ACCEL-ONDANSETRON ODT exactly as your healthcare professional has told you. Talk to your healthcare professional if you are not sure.
- **Do not** take more doses or take them more often than your healthcare professional prescribes. If, however, you vomit within one hour of taking your medicine, you should take the same amount of medicine again. If vomiting persists, consult your healthcare professional.
- Do not try to push or force ACCEL-ONDANSETRON ODT out through the foil package without first peeling the foil back. To take ACCEL-ONDANSETRON ODT out of the foil package, tear along the perforations (dotted lines) to separate one tablet in its blister unit from the rest of the package. Just before you need to take it, with **dry** fingers,

carefully peel back the foil at the place indicated by the arrow, and gently push the ACCEL-ONDANSETRON ODT tablet out of the blister pocket.

- To take the ACCEL-ONDANSETRON ODT, promptly place it on top of your tongue, let it dissolve, and then swallow. It will dissolve very quickly.

**Usual dose:****Adults and Patients (65 years of age and older):**

Your healthcare professional will determine the dose that is right for you and how long you should take it. Your dose will depend on why you are prescribed ACCEL-ONDANSETRON ODT, your age, current health, and if you take certain other medications. Your healthcare professional may monitor your health throughout your treatment and may interrupt, reduce or stop your dose.

**Children (4 to 17 years of age):**

After chemotherapy, take 4 mg every 8 hours, for up to 5 days.

**Overdose:**

If you think you, or a person you are caring for, have taken too much ACCEL-ONDANSETRON ODT, contact a healthcare professional, hospital emergency department, or regional poison control centre immediately, even if there are no symptoms.

**Missed Dose:**

- If you miss a dose and do not feel sick, take the next dose when it is due. **Do not double the dose to make up for the one you missed.**
- If you forget to take your medicine and you feel sick or you vomit, take a dose as soon as possible.
- If you take a tablet and then vomit, **do not** take another one. ACCEL-ONDANSETRON ODT dissolves very quickly and is absorbed rapidly.

**What are possible side effects from using ACCEL-ONDANSETRON ODT?**

These are not all the possible side effects you may have when taking ACCEL-ONDANSETRON ODT. If you experience any side effects not listed here, tell your healthcare professional.

Some side effects may include:

- feeling of flushing or warmth
- hiccups
- headache
- feeling tired

- constipation
- diarrhea

There is no need to stop taking your medicine, but you should tell your healthcare professional about these symptoms at your next visit.

If you feel unwell or have any symptoms that you do not understand, you should contact your healthcare professional immediately.

<b>Serious side effects and what to do about them</b>			
<b>Symptom / effect</b>	<b>Talk to your healthcare professional</b>		<b>Stop taking drug and get immediate medical help</b>
	<b>Only if severe</b>	<b>In all cases</b>	
<b>UNCOMMON</b>			
<b>Heart problems</b> (disorders affecting your heart muscle, valves or rhythm): chest pain, chest discomfort, high blood pressure, irregular heart rhythm, shortness of breath, or fainting.			√
<b>Movement disorders</b> (including dyskinesia): loss of coordination or balance, speech or limb movements, muscle spasms, difficulty walking, tremor, upward rolling of the eyes, or abnormal muscular stiffness.			√
<b>Seizures:</b> loss of consciousness with uncontrollable shaking, <b>visual disturbances (e.g., blurred vision).</b>			√
<b>RARE</b>			
Eye problems such as blurred vision		√	
<b>Hypokalemia</b> (low level of potassium in the blood): muscle weakness, muscle spasms, cramping, constipation, feeling of skipped heart beats or palpitations, fatigue, tingling, or numbness			√
<b>Hypotension</b> (low blood pressure): dizziness, fainting, light-headedness, blurred vision, nausea, or vomiting.			√
<b>Immediate severe allergic reaction:</b> swelling of the mouth, throat, difficulty in breathing, rash, hives, or increased heart rate.			√

<b>Serious side effects and what to do about them</b>			
<b>Symptom / effect</b>	<b>Talk to your healthcare professional</b>		<b>Stop taking drug and get immediate medical help</b>
	<b>Only if severe</b>	<b>In all cases</b>	
<b>Liver problems:</b> yellowing of your skin and eyes (jaundice), unusual dark urine and pale stools, pain or swelling in the right upper abdomen, unusual tiredness, nausea, or vomiting.			√
<b>Myocardial ischemia</b> (lack of blood flow to the heart which can lead to heart attack): sudden chest pain, pressure or discomfort, feeling faint, feeling anxious, shortness of breath, irregular heartbeat, nausea, or sudden heavy sweating.			√
<b>Prolongation of QT interval</b> (a heart rhythm condition): irregular heartbeat, palpitations, dizziness, fainting, loss of consciousness, or seizures.			√
<b>Serotonin syndrome:</b> a reaction which may cause feelings of agitation or restlessness, flushing, muscle twitching, involuntary eye movements, heavy sweating, high body temperature (> 38°C), or rigid muscles.			√
<b>Neuroleptic Malignant Syndrome:</b> pronounced muscle stiffness or inflexibility with high fever, rapid or irregular heartbeat, sweating, state of confusion or reduced consciousness			√
<b>VERY RARE</b>			
Eye problems such as temporary blindness		√	
<b>Stevens-Johnson Syndrome (SJS) and Toxic Epidermal Necrolysis (TEN)</b> (severe skin reactions): redness, blistering or peeling of the skin and/or inside of the lips, eyes, mouth, nasal passages or genitals, fever, chills, headache, cough, body aches, or swollen glands.			√

If you have a troublesome symptom or side effect that is not listed here or becomes bad enough to interfere with your daily activities, tell your healthcare professional.

**Reporting Side Effects**

You can report any suspected side effects associated with the use of health products to Health Canada by:

- Visiting the Web page on Adverse Reaction Reporting (<https://www.canada.ca/en/health-canada/services/drugs-health-products/medeffect-canada/adverse-reaction-reporting.html>) for information on how to report online, by mail or by fax; or
- Calling toll-free at 1-866-234-2345.

*NOTE: Contact your healthcare professional if you need information about how to manage your side effects. The Canada Vigilance Program does not provide medical advice.*

**Storage:**

- ACCEL-ONDANSETRON ODT should be stored at room temperature between 15 - 30°C.
- Keep your medicine in a safe place out of reach and sight of children. Your medicine may harm them.

**If you want more information about ACCEL-ONDANSETRON ODT:**

- Talk to your healthcare professional.
- Find the full Product Monograph that is prepared for healthcare professionals and includes this Patient Medication Information by visiting the Health Canada website (<https://www.canada.ca/en/health-canada/services/drugs-health-products/drug-products/drug-product-database.html>); the manufacturer's website <http://www.accelpharma.com>, or by calling 1-877-822-2235.

This leaflet was prepared by Accel Pharma Inc.

Last Revised: October 30, 2024