

PRODUCT MONOGRAPH
INCLUDING PATIENT MEDICATION INFORMATION

^{Pr} **CLARITHROMYCIN**

Clarithromycin Tablets

Film-coated tablets, 250 mg and 500 mg, Oral

Manufacturer's Standard

Antibiotic

NOTE: WHEN USED IN COMBINATION WITH ACID ANTISECRETORY DRUGS AND OTHER ANTIMICROBIALS FOR THE ERADICATION OF HELICOBACTER PYLORI, THE PRODUCT MONOGRAPH FOR THOSE AGENTS SHOULD BE CONSULTED.

Sivem Pharmaceuticals ULC
4705 Dobrin Street
Saint-Laurent, Quebec, Canada
H4R 2P7
www.sivem.ca

Date of Initial Authorization:
JUL 21, 2015

Date of Revision:
JUL 10, 2025

Submission Control Number: 296811

RECENT MAJOR LABEL CHANGES

| | |
|---|---------|
| 2 CONTRAINDICATIONS | 07/2025 |
| 7 WARNINGS AND PRECAUTIONS, General | 07/2025 |

TABLE OF CONTENTS

Sections or subsections that are not applicable at the time of authorization are not listed.

RECENT MAJOR LABEL CHANGES 2

TABLE OF CONTENTS 2

PART I: HEALTH PROFESSIONAL INFORMATION 4

1 INDICATIONS 4

 1.1 Pediatrics..... 5

 1.2 Geriatrics..... 5

2 CONTRAINDICATIONS 5

3 SERIOUS WARNINGS AND PRECAUTIONS BOX 7

4 DOSAGE AND ADMINISTRATION 7

 4.1 Dosing Considerations 7

 4.2 Recommended Dose and Dosage Adjustment 7

 4.4 Administration 9

 4.5 Missed Dose 9

5 OVERDOSAGE 9

6 DOSAGE FORMS, STRENGTHS, COMPOSITION AND PACKAGING 10

7 WARNINGS AND PRECAUTIONS..... 10

 7.1.1 Pregnant Women..... 15

 7.1.2 Breast-feeding..... 16

 7.1.3 Pediatrics..... 16

 7.1.4 Geriatrics..... 17

8 ADVERSE REACTIONS 17

 8.1 Adverse Reaction Overview 17

 8.2 Clinical Trial Adverse Reactions 17

| | |
|---|-----------|
| 8.4 Abnormal Laboratory Findings: Hematologic, Clinical Chemistry and Other | 23 |
| Quantitative Data..... | 23 |
| 8.5 Post-Market Adverse Reactions..... | 25 |
| 9 DRUG INTERACTIONS | 27 |
| 9.1 Serious Drug Interactions..... | 27 |
| 9.2 Drug Interactions Overview | 27 |
| 9.3 Drug-Behavioural Interactions..... | 29 |
| 9.4 Drug-Drug Interactions | 29 |
| 9.5 Drug-Food Interactions..... | 47 |
| 9.6 Drug-Herb Interactions | 47 |
| 9.7 Drug-Laboratory Test Interactions..... | 47 |
| 10 CLINICAL PHARMACOLOGY | 47 |
| 10.1 Mechanism of Action | 47 |
| 10.2 Pharmacodynamics..... | 47 |
| 10.3 Pharmacokinetics..... | 48 |
| 11 STORAGE, STABILITY AND DISPOSAL | 56 |
| 12 SPECIAL HANDLING INSTRUCTIONS..... | 56 |
| PART II: SCIENTIFIC INFORMATION | 57 |
| 13 PHARMACEUTICAL INFORMATION | 57 |
| 14 CLINICAL TRIALS | 58 |
| 14.1 Trial Design and Study Demographics | 58 |
| 14.2 Study Results..... | 58 |
| 14.3 Comparative Bioavailability Studies | 66 |
| 15 MICROBIOLOGY | 68 |
| 16 NON-CLINICAL TOXICOLOGY..... | 77 |
| 17 SUPPORTING PRODUCT MONOGRAPHS | 82 |
| PATIENT MEDICATION INFORMATION | 83 |

PART I: HEALTH PROFESSIONAL INFORMATION

1 INDICATIONS

CLARITHROMYCIN (clarithromycin tablets) may be indicated in the treatment of mild to moderate infections caused by susceptible strains of the designated microorganisms in the diseases listed below:

- Upper Respiratory Tract
 - Pharyngitis/tonsillitis, caused by *Streptococcus pyogenes* (Group A beta-hemolytic streptococci).
 - Acute maxillary sinusitis caused by *Streptococcus pneumoniae* (*S. pneumoniae*), *Haemophilus influenzae* (*H. influenzae*), and *Moraxella* (*Branhamella*) *catarrhalis* [*M. (Branhamella) catarrhalis*].
- Lower Respiratory Tract
 - Acute bacterial exacerbation of chronic bronchitis caused by *S. pneumoniae*, *H. influenzae* (including beta-lactamase producing strains), *M. (Branhamella) catarrhalis* (including beta-lactamase producing strains).
 - Pneumonia caused by *S. pneumoniae* and *Mycoplasma pneumoniae* (*M. pneumoniae*). See [7 WARNINGS AND PRECAUTIONS, Sensitivity/Resistance](#).
- Uncomplicated Skin and Skin Structure Infections
 - Uncomplicated Skin and Skin Structure Infections caused by *Streptococcus pyogenes* (*S. pyogenes*), *Staphylococcus aureus* (*S. aureus*). See [7 WARNINGS AND PRECAUTIONS, Sensitivity/Resistance](#).
- Mycobacterial Infections
 - CLARITHROMYCIN is indicated for the prevention of disseminated *Mycobacterium avium* complex (MAC) disease in patients with advanced HIV infection, and for the treatment of disseminated mycobacterial infections due to *Mycobacterium avium* (*M. avium*) and *Mycobacterium intracellulare* (*M. intracellulare*). See [14 CLINICAL TRIALS, Mycobacterial Infections](#).
- Eradication of *Helicobacter pylori*
 - CLARITHROMYCIN in the presence of acid suppression (with omeprazole) with another antibiotic (amoxicillin) is indicated for the eradication of *Helicobacter pylori* (*H. pylori*) that may result in decreased recurrence of duodenal ulcer in patients with active duodenal ulcers and who are *H. pylori* positive. See [14 CLINICAL TRIALS, Eradication of *Helicobacter pylori*, Triple Therapy: Clarithromycin /omeprazole/amoxicillin](#).

(For additional information on the use of CLARITHROMYCIN in triple therapy for the treatment of *H. pylori* infection and active duodenal ulcer recurrence, refer to the Hp-PAC® Product Monograph.)

To reduce the development of drug-resistant bacteria and maintain the effectiveness of CLARITHROMYCIN and other antibacterial drugs, CLARITHROMYCIN should be used only to treat infections that are proven or strongly suspected to be caused by susceptible bacteria. When culture and susceptibility information are available, they should be considered in selecting or modifying antibacterial therapy. In the absence of such data, local epidemiology and susceptibility patterns may contribute to the empiric selection of therapy.

1.1 Pediatrics

Pediatrics (< 12 years of age): Based on the data submitted and reviewed by Health Canada, the safety and efficacy of CLARITHROMYCIN in pediatric patients has not been established; therefore, Health Canada has not authorized an indication for pediatric use.

1.2 Geriatrics

Geriatrics (> 65 years of age): Dosage adjustment should be considered in elderly patients with severe renal impairment. See [7 WARNINGS AND PRECAUTIONS, 7.1 Special Populations, 7.1.4 Geriatrics](#).

2 CONTRAINDICATIONS

- CLARITHROMYCIN is contraindicated in:
 - Patients with a known hypersensitivity to clarithromycin, erythromycin, other macrolide antibacterial agents or to any ingredient in the formulation, including any non-medicinal ingredient, or component of the container. See [6 DOSAGE FORMS, STRENGTHS, COMPOSITION AND PACKAGING](#).
 - Patients with a history of cholestatic jaundice/hepatic dysfunction associated with prior use of clarithromycin.
 - Patients who suffer from severe hepatic failure in combination with renal impairment. See [7 WARNINGS AND PRECAUTIONS, Hepatic/Biliary/Pancreatic](#), [7 WARNINGS AND PRECAUTIONS, Renal](#), [4 DOSAGE AND ADMINISTRATION, Dosing Considerations](#) and [4 DOSAGE AND ADMINISTRATION, 4.2 Recommended Dose and Dosage Adjustment](#).
 - Patients with history of QT prolongation (congenital or documented acquired QT prolongation) or ventricular cardiac arrhythmia, including torsades de pointes. See [7 WARNINGS AND PRECAUTIONS](#) and [9 DRUG INTERACTIONS, 9.4 Drug-Drug Interactions](#).
 - Patients with electrolyte disturbances (hypokalaemia or hypomagnesaemia, due to the risk of prolongation of the QT interval and torsades de pointes).

- Concomitant therapy with astemizole, cisapride, domperidone, pimozone, terfenadine.

There have been post-marketing reports of drug interactions when clarithromycin and/or erythromycin are co-administered with astemizole, cisapride, pimozone or terfenadine resulting in cardiac arrhythmias (QT prolongation, ventricular tachycardia, ventricular fibrillation, and torsades de pointes) most likely due to inhibition of hepatic metabolism of these drugs by erythromycin and clarithromycin. Fatalities have been reported. See [9 DRUG INTERACTIONS, 9.4 Drug-Drug Interactions, Table 11](#).

- Concomitant therapy with saquinavir due to potentially life-threatening cardiac arrhythmia.

Concomitant therapy with HMG-CoA reductase inhibitors (statins) that are extensively metabolized by CYP3A4 (lovastatin or simvastatin) due to an increased risk of myopathy, including rhabdomyolysis. See [9 DRUG INTERACTIONS, 9.4 Drug-Drug Interactions, Table 11](#).

Concomitant therapy with ergot alkaloids (e.g., ergotamine or dihydroergotamine) as this may result in ergot toxicity. See [9 DRUG INTERACTIONS, 9.4 Drug-Drug Interactions, Table 11](#).

Concomitant administration with **oral** midazolam. See [9 DRUG INTERACTIONS, 9.4 Drug-Drug Interactions, Table 11](#).

- Concomitant administration with lomitapide. See [9 DRUG INTERACTIONS](#).

Concomitant therapy with colchicine due to the risk of life threatening and fatal colchicine toxicity. This risk may be further increased with concomitant medications metabolized by P-glycoprotein or strong CYP3A inhibitors. See [9 DRUG INTERACTIONS, 9.4 Drug-Drug Interactions, Table 11](#).

- Concomitant therapy with ivabradine due to potential increase in ivabradine levels and risk of excessive bradycardia. See [9 DRUG INTERACTIONS, 9.4 Drug-Drug Interactions, Table 11](#).

- Concomitant therapy with ticagrelor or ranolazine*.

* Not marketed in Canada.

3 SERIOUS WARNINGS AND PRECAUTIONS BOX

Serious Warnings and Precautions

- Clarithromycin should not be used in **pregnancy** except where no alternative therapy is appropriate, particularly during the first 3 months of pregnancy. If pregnancy occurs while taking the drug, the patient should be apprised of the potential hazard to the fetus. See [7 WARNINGS AND PRECAUTIONS, 7.1 Special Populations, 7.1.1 Pregnant Women](#).
- The concomitant administration of clarithromycin and drugs metabolized by CYP3A and/or transported by P-gp may result in significant safety concerns. See [7 WARNINGS AND PRECAUTIONS](#) and [9 DRUG INTERACTIONS, 9.2 Drug Interactions Overview](#).

4 DOSAGE AND ADMINISTRATION

4.1 Dosing Considerations

- CLARITHROMYCIN may be given with or without meals.
- In patients with a combination of hepatic (mild to moderate) and renal impairments or in the presence of severe renal impairment, decreased dosage of clarithromycin or prolonged dosing intervals might be appropriate. See [4 DOSAGE AND ADMINISTRATION, 4.2 Recommended Dose and Dosage Adjustment](#).
- Clarithromycin is contraindicated in patients with severe hepatic failure in combination with renal impairment. See [2 CONTRAINDICATIONS](#).
- In patients with renal impairment and a creatinine clearance < 30 mL/min, the dosage of CLARITHROMYCIN should be reduced by one-half, i.e., up to 250 mg once daily, or 250 mg twice daily in more severe infections. Dosage should not be continued beyond 14 days in these patients.

4.2 Recommended Dose and Dosage Adjustment

- Adults with Respiratory Tract or Skin Infections
 - The adult dosage of CLARITHROMYCIN is 250 mg to 500 mg every 12 hours (Table 1) for 7 to 14 days. For infections caused by less susceptible organisms, the upper dosage should be used.

Table 1 - Adult Dosage Guidelines

| Infection | Dosage (b.i.d.) | Duration |
|---|---------------------------------------|-------------------------|
| Upper Respiratory Tract Pharyngitis/tonsillitis Acute maxillary sinusitis | 250-500 mg 250 mg 500 mg | 10 days 7 to 14 days |
| Lower Respiratory Tract Acute exacerbation of chronic bronchitis and pneumonia | 250-500 mg 250-500 mg | 7 to 14 days |
| Uncomplicated Skin and Skin Structure Infections | 250 mg | 7 to 14 days |

Legend: b.i.d. = twice daily

- In the treatment of Group A streptococcus infections, therapy should be continued for 10 days. The usual drug of choice in the treatment of streptococcal infections and the prophylaxis of rheumatic fever is penicillin administered by either the intramuscular or the oral route.
- Clarithromycin is generally effective in the eradication of *S. pyogenes* from the nasopharynx; however, data establishing the efficacy of clarithromycin in the subsequent prevention of rheumatic fever are not presently available.
- Renal Impairment
 - In patients with severe renal impairment (creatinine clearance < 30 mL/min., the dosage of CLARITHROMYCIN should be reduced by one-half, i.e. 250 mg once daily, or 250 mg twice daily in more severe infections. Dosage should not be continued beyond 14 days in these patients. The safety and efficacy of 500 mg clarithromycin in patients with severe renal impairment has not been established.
- Hepatic Impairment
 - In patients with a combination of hepatic (mild to moderate) and renal impairments, decreased dosage of clarithromycin or prolonged dosing intervals may be appropriate. Clarithromycin may be administered without dosage adjustment in the presence of hepatic impairment if there is normal renal function.
 - Clarithromycin is contraindicated in patients with severe hepatic failure in combination with renal impairment. See [2 CONTRAINDICATIONS](#).
- Eradication of *Helicobacter pylori*
 - Triple Therapy: CLARITHROMYCIN/omeprazole/amoxicillin: The recommended dose is clarithromycin 500 mg twice daily in conjunction with omeprazole 20 mg daily and amoxicillin 1000 mg twice daily for 10 days. See [14 CLINICAL TRIALS, Eradication of *Helicobacter pylori*, Triple Therapy: Clarithromycin/omeprazole/amoxicillin](#).

For more information on omeprazole or amoxicillin, refer to their respective Product Monographs, under [4 DOSAGE AND ADMINISTRATION](#).

(For additional information on the use of CLARITHROMYCIN in triple therapy for the treatment of *H. pylori* infection and active duodenal ulcer recurrence, refer to the Hp-PAC® Product Monograph.)

- Adults with Mycobacterial Infections
 - Prophylaxis: The recommended dose of CLARITHROMYCIN for the prevention of disseminated *M. avium* disease is 500 mg twice daily.
 - Treatment: Clarithromycin is recommended as the primary agent for the treatment of disseminated infection due to MAC. Clarithromycin should be used in combination with other antimycobacterial drugs which have shown *in vitro* activity against MAC, including ethambutol and rifampin. Although no controlled clinical trial information is available for combination therapy with clarithromycin, the U.S. Public Health Service Task Force has provided recommendations for the treatment of MAC.

The recommended dose for mycobacterial infections in adults is 500 mg twice daily.

Treatment of disseminated MAC infections in AIDS patients should continue for life if clinical and mycobacterial improvement are observed.

4.4 Administration

CLARITHROMYCIN may be taken with or without food.

4.5 Missed Dose

If a dose of clarithromycin is missed, the patient should take the dose as soon as possible and then return to their normal scheduled dose. However, if a dose is skipped, the patient should not double the next dose.

5 OVERDOSAGE

Activated charcoal may be administered to aid in the removal of unabsorbed drug. General supportive measures are recommended.

Reports indicate that the ingestion of large amounts of clarithromycin can be expected to produce gastrointestinal symptoms. Adverse reactions accompanying overdose should be treated by the prompt elimination of unabsorbed drug and supportive measures.

Clarithromycin is protein bound (70%). No data are available on the elimination of clarithromycin by hemodialysis or peritoneal dialysis.

For management of a suspected drug overdose, contact your regional poison control centre.

6 DOSAGE FORMS, STRENGTHS, COMPOSITION AND PACKAGING

Table 2 – Dosage Forms, Strengths, Composition and Packaging

| Route of Administration | Dosage Form/ Strength / Composition | Non-medicinal Ingredients |
|-------------------------|-------------------------------------|--|
| Oral | Film-coated tablets, 250 mg | colloidal silica hydrated, croscarmellose sodium, magnesium stearate, microcrystalline cellulose, opadry yellow (D&C Yellow #10 aluminum lake, hydroxypropyl methylcellulose, lactose monohydrate, polyethylene glycol 4000, titanium dioxide) and powdered cellulose. |
| Oral | Film-coated tablets, 500 mg | colloidal silica hydrated, croscarmellose sodium, magnesium stearate, microcrystalline cellulose, opadry yellow (D&C Yellow #10 aluminum lake, hydroxypropyl methylcellulose, lactose monohydrate, polyethylene glycol 4000, titanium dioxide) and powdered cellulose. |

CLARITHROMYCIN 250 mg tablets are supplied as yellow, oblong, convex, film-coated tablet, scored on both sides, with 'CL' and '250' on a side contains 250 mg of clarithromycin for oral administration, and are available in HDPE bottles of 100 tablets.

CLARITHROMYCIN 500 mg tablets are supplied as yellow, oblong, convex, film-coated tablet, scored on both sides, with 'CL' and '500' on a side contains 500 mg of clarithromycin for oral administration, and are available in HDPE bottles of 100 tablets.

7 WARNINGS AND PRECAUTIONS

Please see [3 SERIOUS WARNINGS AND PRECAUTIONS BOX](#).

General

Clarithromycin should be administered with caution to any patient who has demonstrated some form of drug allergy, particularly to structurally related-drugs. If an allergic reaction to clarithromycin occurs, administration of the drug should be discontinued. Serious hypersensitivity reactions may require epinephrine, antihistamines, or corticosteroids. See [7 WARNINGS AND PRECAUTIONS, Immune, Hypersensitivity Reactions](#).

Long-term use may, as with other antibiotics, result in colonization with increased numbers of non-susceptible bacteria and fungi. If superinfections occur, appropriate therapy should be instituted.

Patients Infected with Human Immunodeficiency Virus

Several studies of Human Immunodeficiency Virus (HIV)-positive patients receiving clarithromycin for treatment of MAC infection have shown poorer survival in those patients randomized to receive doses higher than 500 mg twice daily. The explanation for the poorer survival associated with doses higher than 500 mg twice daily has not been determined. Treatment or prophylaxis of MAC infection with clarithromycin should not exceed the approved dose of 500 mg twice daily.

Myasthenia Gravis

Exacerbation of symptoms of myasthenia gravis and new onset of symptoms of myasthenic syndrome has been reported in patients receiving clarithromycin therapy.

Use of Clarithromycin with Other Drugs

Use of clarithromycin with other drugs may lead to drug-drug interactions.

Atypical Antipsychotics (quetiapine)

Due to inhibition of CYP3A by clarithromycin, co-administration of clarithromycin with quetiapine results in increased quetiapine concentrations. Serious and life-threatening quetiapine-related adverse reactions, including malignant neuroleptic syndrome, have been reported. Clarithromycin should not be used in combination with quetiapine unless clinically necessary. See [9 DRUG INTERACTIONS](#). Monitoring and dose reductions may be required.

Oral Hypoglycemic Agents/Insulin

The concomitant use of clarithromycin and oral hypoglycaemic agents (such as sulphonylurias) and/or insulin can result in significant hypoglycaemia. Careful monitoring of glucose is recommended. See [9 DRUG INTERACTIONS, 9.4 Drug-Drug Interactions, Table 11](#).

Oral Anticoagulants

There is a risk of serious hemorrhage and significant elevations in International Normalized Ratio (INR) and prothrombin time when clarithromycin is co-administered with warfarin. INR and prothrombin times should be frequently monitored while patients are receiving clarithromycin and oral anticoagulants concurrently. [9 DRUG INTERACTIONS, 9.4 Drug-Drug Interactions, Table 11](#).

Caution should be exercised when clarithromycin is co-administered with direct acting oral anticoagulants such as dabigatran, rivaroxaban, apixaban and edoxaban, particularly to patients at high risk of bleeding. See [9 DRUG INTERACTIONS, 9.4 Drug-Drug Interactions](#).

HMG-CoA Reductase Inhibitors

Concomitant use of clarithromycin with lovastatin or simvastatin is contraindicated. See [2 CONTRAINDICATIONS](#). Caution should be exercised when prescribing clarithromycin with other

statins. Rhabdomyolysis has been reported in patients taking clarithromycin and statins. Patients should be monitored for signs and symptoms of myopathy. In situations where the concomitant use of clarithromycin with statins cannot be avoided, it is recommended to prescribe the lowest registered dose of the statin. Use of a statin that is not dependent on CYP3A metabolism (e.g., fluvastatin) can be considered. [9 DRUG INTERACTIONS, 9.4 Drug-Drug Interactions, Table 11.](#)

Triazolobenzodiazepines and Related Benzodiazepines

Caution is advised regarding the concomitant administration of clarithromycin with triazolobenzodiazepines (such as triazolam and alprazolam), or with other benzodiazepines (such as intravenous midazolam) due to the serious risk of central nervous system (CNS) effects (e.g., somnolence and confusion). [9 DRUG INTERACTIONS, 9.4 Drug-Drug Interactions, Table 11.](#)

Concomitant administration with oral midazolam is contraindicated. See [2 CONTRAINDICATIONS](#)

Calcium Channel Blockers

Caution is advised regarding the concomitant administration of clarithromycin and calcium channel blockers metabolized by CYP3A4 (e.g., verapamil, amlodipine, diltiazem) due to the risk of hypotension. See [9 DRUG INTERACTIONS, 9.4 Drug-Drug Interactions, Table 11.](#)

Hypotension, bradyarrhythmias, and lactic acidosis have been observed in patients receiving concurrent verapamil, belonging to the calcium channel blockers drug class. See [9 DRUG INTERACTIONS, 9.4 Drug-Drug Interactions, Table 11.](#)

Other Drugs

For other established or potential drug-drug interactions and their mechanisms, see [2 CONTRAINDICATIONS](#) and [9 DRUG INTERACTIONS, 9.4 Drug-Drug Interactions.](#)

Carcinogenesis and Mutagenesis

See [16 NON-CLINICAL TOXICOLOGY.](#)

Cardiovascular

Cardiovascular Events

Prolonged cardiac repolarisation and QT interval, imparting a risk of developing cardiac arrhythmia and torsades de pointes, have been seen in treatment with macrolides, including clarithromycin. See [8 ADVERSE REACTIONS.](#) Fatalities have been reported. Elderly patients may be more susceptible to drug-associated effects on the QT interval.

As the following situations may lead to an increased risk for ventricular arrhythmias (including torsades de pointes), clarithromycin should be used with caution in patients with coronary artery disease, cardiac insufficiency, conduction disturbances, clinically significant bradycardia (e.g., < 50 bpm), or when concomitantly taking with other medicinal products associated with QT prolongation, due to the risk for QT prolongation and torsades de pointes. See [9 DRUG](#)

[INTERACTIONS](#). Clarithromycin must not be given to patients with electrolyte disturbances such as hypomagnesaemia or hypokalemia. See [2 CONTRAINDICATIONS](#).

Clarithromycin is contraindicated in patients with congenital or documented acquired QT prolongation or history of ventricular arrhythmia, including torsades de pointes. Clarithromycin is also contraindicated in patients with hypokalaemia due to the risk of QT prolongation and torsades de pointes. Concomitant administration of clarithromycin with astemizole, cisapride, domperidone, pimozone, terfenadine and saquinavir is also contraindicated. See [2 CONTRAINDICATIONS](#).

Epidemiological studies investigating the risk of adverse cardiovascular outcomes with macrolides have shown variable results. Studies have identified risks of arrhythmia, myocardial infarction and cardiovascular mortality associated with macrolides including clarithromycin. Consideration of these findings should be balanced with treatment benefits when prescribing clarithromycin.

Driving and Operating Machinery

There are no data on the effect of clarithromycin on the ability to drive or use machines. The potential for dizziness, vertigo, confusion and disorientation, which may occur with the medication, should be taken into account before patients drive or use machines.

Gastrointestinal

Clostridioides difficile-Associated Disease

Clostridioides difficile-associated disease (CDAD) has been reported with use of many antibacterial agents, including clarithromycin. CDAD may range in severity from mild diarrhea to fatal colitis. It is important to consider this diagnosis in patients who present with diarrhea, or symptoms of colitis, pseudomembranous colitis, toxic megacolon, or perforation of colon subsequent to the administration of any antibacterial agent. CDAD has been reported to occur over 2 months after the administration of antibacterial agents.

Treatment with antibacterial agents may alter the normal flora of the colon and may permit overgrowth of *Clostridioides difficile*. *Clostridioides difficile* produces toxins A and B, which contribute to the development of CDAD. CDAD may cause significant morbidity and mortality. CDAD can be refractory to antimicrobial therapy.

If the diagnosis of CDAD is suspected or confirmed, appropriate therapeutic measures should be initiated. Mild cases of CDAD usually respond to discontinuation of antibacterial agents not directed against *Clostridioides difficile*. In moderate to severe cases, consideration should be given to management with fluids and electrolytes, protein supplementation, and treatment with an antibacterial drug clinically effective against *Clostridioides difficile*. Surgical evaluation should be instituted as clinically indicated, as surgical intervention may be required in certain severe cases. See [8 ADVERSE REACTIONS](#).

Hepatic/Biliary/Pancreatic

Caution is advised in patients with impaired hepatic function.

Clarithromycin is principally excreted by the liver and kidney. In patients with a combination of hepatic (mild to moderate) and renal impairments, decreased dosage of clarithromycin or prolonged dosing intervals might be appropriate. See [4 DOSAGE AND ADMINISTRATION, Recommended Dose and Dosage Adjustment](#).

Clarithromycin is contraindicated in patients with severe hepatic failure in combination with renal impairment. See [2 CONTRAINDICATIONS](#).

Hepatic dysfunction, including increased liver enzymes, and hepatocellular and/or cholestatic hepatitis, with or without jaundice, has been reported with clarithromycin. This hepatic dysfunction may be severe and is usually reversible. In some instances, hepatic failure with fatal outcomes has been reported and generally has been associated with serious underlying diseases and/or concomitant medications. Discontinue clarithromycin immediately if signs and symptoms of hepatitis occur, such as anorexia, jaundice, dark urine, pruritus, or tender abdomen.

Immune

Hypersensitivity Reactions

In the event of severe acute hypersensitivity reactions, such as anaphylaxis, severe cutaneous adverse reactions (SCAR) (e.g., acute generalized exanthematous pustulosis (AGEP), Stevens-Johnson syndrome (SJS), toxic epidermal necrolysis (TEN) and drug rash with eosinophilia and systemic symptoms (DRESS)), clarithromycin therapy should be discontinued immediately and appropriate treatment should be urgently initiated.

Renal

Caution should be exercised when administering clarithromycin to patients with moderate to severe renal impairment.

Clarithromycin is principally excreted by the liver and kidney. In patients with a combination of hepatic (mild to moderate) and renal impairments or in the presence of severe renal impairment, decreased dosage of clarithromycin or prolonged dosing intervals might be appropriate. See [4 DOSAGE AND ADMINISTRATION, 4.2 Recommended Dose and Dosage Adjustment](#).

Clarithromycin is contraindicated in patients with severe hepatic failure in combination with renal impairment. See [2 CONTRAINDICATIONS](#).

For the eradication of *H. pylori*, amoxicillin and clarithromycin should not be administered to patients with renal impairment since the appropriate dosage in this patient population has not yet been established.

Reproductive Health: Female and Male Potential

- **Fertility**

See [16 NON-CLINICAL TOXICOLOGY; Reproductive and Developmental Toxicology](#). Please See [3 SERIOUS WARNINGS AND PRECAUTIONS BOX](#). See [7 WARNINGS AND PRECAUTIONS, 7.1.1 Pregnant Women](#).

- **Teratogenic Risk**

Please See [3 SERIOUS WARNINGS AND PRECAUTIONS BOX](#). See [7 WARNINGS AND PRECAUTIONS, 7.1.1 Pregnant Women](#).

Sensitivity/Resistance

The development of resistance (11 out of 19 breakthrough isolates in 1 study) has been seen in HIV positive patients receiving clarithromycin for prophylaxis and treatment of MAC infection.

In view of the emerging resistance of *Streptococcus pneumoniae*, *Staphylococcus aureus* and *Streptococcus pyogenes* to macrolides, it is important that susceptibility testing be performed when prescribing clarithromycin for community-acquired pneumonia and uncomplicated skin and skin structure infections.

To avoid failure of the eradication treatment with a potential for developing antimicrobial resistance and a risk of failure with subsequent therapy, patients should be instructed to follow closely the prescribed regimen.

Development of Drug-Resistant Bacteria

Prescribing CLARITHROMYCIN in the absence of a proven or strongly suspected bacterial infection is unlikely to provide benefit to the patient and risks the development of drug-resistant bacteria.

Antibiotic Resistance in Relation to *Helicobacter pylori* Eradication

Use of any antimicrobial therapy, such as clarithromycin, to treat *H. pylori* infection may select for drug-resistant organisms.

Triple Therapy with Omeprazole

Among the 113 triple therapy recipients with pretreatment *H. pylori* isolates susceptible to clarithromycin, 2/102 patients (2%) developed resistance after treatment with omeprazole, clarithromycin, and amoxicillin. Among patients who received triple therapy, 6/108 (5.6%) patients had pretreatment *H. pylori* isolates resistant to clarithromycin. Of these 6 patients, 3 (50%) had *H. pylori* eradicated at follow-up, and 3 (50%) remained positive after treatment. In 5/113 (4.4%) patients, no susceptibility data for clarithromycin pretreatment were available. Development of clarithromycin resistance should be considered as a possible risk especially when less efficient treatment regimens are used.

Special Populations

7.1.1 Pregnant Women

There are no adequate and well-controlled studies in pregnant women. The benefits against risk, particularly during the first 3 months of pregnancy should be carefully weighed by a physician. Based on variable results obtained from animal studies and experience in humans, the possibility of adverse effects on embryofoetal development cannot be excluded. Some observational studies evaluating exposure to clarithromycin during the first and second trimester have reported an increased risk of miscarriage compared to no antibiotic use or other antibiotic use during the same period. The available epidemiological studies on the risk of major congenital malformations with use of macrolides including clarithromycin during pregnancy provide conflicting results. Therefore, use during pregnancy is not advised without carefully weighing the benefits against risks. See [3 SERIOUS WARNINGS AND PRECAUTIONS BOX](#).

Four teratogenicity studies in rats (3 with oral doses and 1 with intravenous doses up to 160 mg/kg/day administered during the period of major organogenesis) and 2 in rabbits (at oral doses up to 125 mg/kg/day or intravenous doses of 30 mg/kg/day administered during gestation days 6 to 18) failed to demonstrate any teratogenicity from clarithromycin. Two additional oral studies in a different rat strain at similar doses and similar conditions demonstrated a low incidence of cardiovascular anomalies at doses of 150 mg/kg/day administered during gestation days 6 to 15. Plasma levels after 150 mg/kg/day were 2 times the human serum levels.

Four studies in mice revealed a variable incidence of cleft palate following oral doses of 1000 mg/kg/day during gestation days 6 to 15. Cleft palate was also seen at 500 mg/kg/day. The 1000 mg/kg/day exposure resulted in plasma levels 17 times the human serum levels. In monkeys, an oral dose of 70 mg/kg/day produced fetal growth retardation at plasma levels that were 2 times the human serum levels.

Embryonic loss has been seen in monkeys and rabbits. See [16 NON-CLINICAL TOXICOLOGY, Reproductive and Developmental Toxicology](#).

7.1.2 Breast-feeding

The safety of clarithromycin for use during breast feeding of infants has not been established. Clarithromycin is excreted in human breast milk in small amounts. It has been estimated that an exclusively breastfed infant would receive about 1.7% of the maternal weight-adjusted dose of clarithromycin.

Preweaned rats, exposed indirectly via consumption of milk from dams treated with 150 mg/kg/day for 3 weeks, were not adversely affected, despite data indicating higher drug levels in milk than in plasma.

7.1.3 Pediatrics

Pediatrics (<6 months to 12 years of age): Use of CLARITHROMYCIN in children under 12 years

of age has not been studied.

The safety of clarithromycin has not been studied in MAC patients under the age of 20 months.

Neonatal and juvenile animals tolerated clarithromycin in a manner similar to adult animals. Young animals were slightly more intolerant to acute overdosage and to subtle reductions in erythrocytes, platelets and leukocytes, but were less sensitive to toxicity in the liver, kidney, thymus and genitalia.

Increased valproate and phenobarbital concentrations and extreme sedation were noted in a 3 year old patient coincident with clarithromycin therapy. Cause and effect relationship cannot be established. However, monitoring of valproate and phenobarbital concentrations may be considered.

7.1.4 Geriatrics

Geriatrics (> 65 years of age): Dosage adjustment should be considered in elderly patients with severe renal impairment. In a steady-state study in which healthy elderly subjects (age 65 to 81 years old) were given 500 mg every 12 hours, the maximum concentrations of clarithromycin and 14-OH clarithromycin were increased. The AUC was also increased. These changes in pharmacokinetics parallel known age-related decreases in renal function. In clinical trials, elderly patients did not have an increased incidence of adverse events when compared to younger patients.

8 ADVERSE REACTIONS

8.1 Adverse Reaction Overview

The majority of side effects observed in clinical trials involving 3563 patients treated with clarithromycin were of a mild and transient nature. Fewer than 3% of adult patients without mycobacterial infections discontinued therapy because of drug-related side effects. The most common drug-related adverse reactions in adults taking clarithromycin were nausea, diarrhea, abdominal pain, dyspepsia, headache, dysgeusia (taste perversion) and vomiting.

8.2 Clinical Trial Adverse Reactions

Clinical trials are conducted under very specific conditions. The adverse reaction rates observed in the clinical trials; therefore, may not reflect the rates observed in practice and should not be compared to the rates in the clinical trials of another drug. Adverse reaction information from clinical trials may be useful in identifying and approximating rates of adverse drug reactions in real-world use.

Patients with Respiratory Tract or Skin Infections

Table 3 provides a listing of adverse reactions from clinical trials or post-marketing surveillance as well as adverse events reported during post-marketing surveillance. Adverse events reported

during post-marketing surveillance may include patients treated for various infections and are not limited to patients with respiratory tract or skin infections.

Table 3 - Adverse Events/Adverse Drug Reactions in Patients with Respiratory Tract or Skin Infections or Other Infections Treated with Clarithromycin Tablets

| System Organ Class | Adverse Reaction/Adverse Event |
|--|--|
| Blood and lymphatic system disorders | Eosinophilia Anemia Leukopenia Thrombocythemia Thrombocytopenia |
| Cardiac disorders* | Electrocardiogram QT prolonged Ventricular tachycardia Torsades de pointes |
| Ear and labyrinth disorders | Vertigo Tinnitus Ear disorder Deafness**** |
| Eye disorders | Visual disturbance Conjunctivitis |
| Gastrointestinal disorders | Constipation Flatulence Dry mouth Glossitis Stomatitis Gastrointestinal disorder Tongue discolouration Tooth discolouration Pancreatitis |
| General disorders and administration site conditions | Asthenia Pain Chest pain |
| Hepatobiliary disorders | Hepatomegaly Hepatic function abnormal Hepatitis Hepatitis cholestatic Jaundice (cholestatic and hepatocellular) Hepatic failure *** |
| Immune system disorders | Anaphylactic reaction Myasthenia gravis |

| System Organ Class | Adverse Reaction/Adverse Event |
|---|---|
| Infections and infestations | Infection Colitis pseudomembranous Candidiasis Rhinitis Pharyngitis Vaginal candidiasis Vaginal infection |
| Investigations | Increased liver enzymes |
| Metabolism and nutrition disorders | Anorexia Hypoglycemia** |
| Musculoskeletal and connective tissue disorders | Back pain Myalgia |
| Nervous system disorders | Dizziness Somnolence Convulsion Parosmia Dysgeusia Ageusia |
| Psychiatric disorders | Nervousness Anxiety Insomnia Nightmare Depression Confusional state Disorientation Depersonalisation Hallucination Psychotic disorder |
| Renal and urinary disorders | Hematuria Nephritis interstitial |
| Reproductive system and breast disorders | Dysmenorrhea |
| Respiratory, thoracic and mediastinal disorders | Cough Dyspnea Asthma |
| Skin and subcutaneous tissue disorders | Severe cutaneous adverse reactions (SCAR) (e.g., Acute generalized exanthematous pustulosis (AGEP), Stevens-Johnson syndrome (SJS), Toxic epidermal necrosis (TEN), Drug rash with eosinophilia and systemic symptoms |

| System Organ Class | Adverse Reaction/Adverse Event |
|--------------------|--|
| | (DRESS)) Pruritus Rash Hyperhidrosis Urticaria |

*As with other macrolides, QT prolongation, ventricular tachycardia, and torsades de pointes have been reported with clarithromycin.

**There have been reports of hypoglycemia, some of which have occurred in patients on concomitant oral hypoglycemic agents or insulin.

***Hepatic dysfunction may be severe and is usually reversible. Hepatic failure with fatal outcome has been reported and generally has been associated with serious underlying diseases and/or concomitant medications.

**** There have been reports of hearing loss with clarithromycin which is usually reversible upon withdrawal of therapy.

In studies of adults with pneumonia comparing clarithromycin to erythromycin base or erythromycin stearate, there were significantly fewer adverse events involving the digestive system in patients treated with clarithromycin.

Patients with Mycobacterial Infections

In patients with acquired immune deficiency syndrome (AIDS) and other immunocompromised patients treated with the higher doses of clarithromycin over long periods of time for prevention or treatment of mycobacterial infections, it was often difficult to distinguish adverse events possibly associated with clarithromycin administration from underlying signs of HIV disease or intercurrent illness.

Prophylaxis

Discontinuation due to adverse events was required in 18% of AIDS patients receiving clarithromycin 500 mg twice daily, compared to 17% of patients receiving placebo in a randomized, double-blind study. Primary reasons for discontinuation in the clarithromycin-treated patients include headache, nausea, vomiting, depression and taste perversion. The most frequently reported adverse events with an incidence of 2% or greater, excluding those due to the patient's concurrent condition, are listed in Table 4. Among these events, taste perversion was the only event that had significantly higher incidence in the clarithromycin-treated compared to the placebo-treated group.

Table 4 - Percentage of Adverse Events* in Immunocompromised Adult Patients Receiving Prophylaxis Against *M. avium* Complex

| System Organ Class† | Adverse Reaction | Clarithromycin (N=339) % | Placebo (N=339) % |
|---------------------|------------------|-----------------------------|----------------------|
| | | | |

| | | | |
|--|----------------|-------|------|
| Gastrointestinal disorders | Abdominal pain | 5.0% | 3.5% |
| | Nausea | 11.2% | 7.1% |
| | Diarrhea | 7.7% | 4.1% |
| | Vomiting | 5.9% | 3.2% |
| | Dyspepsia | 3.8% | 2.7% |
| | Flatulence | 2.4% | 0.9% |
| Nervous system disorders | Dysgeusia | 8.0% | 0.3% |
| | Headache | 2.7% | 0.9% |
| Skin and subcutaneous tissue disorders | Rash | 3.2% | 3.5% |

* Includes those events possibly or probably related to study drug and excludes concurrent conditions.

‡ ≥ 2% Adverse Event Incidence Rates for either treatment group.

Treatment of Patients with Mycobacterial Infections

Excluding those patients who discontinued therapy due to complications of their underlying non-mycobacterial diseases (including death), approximately 14% of the patients discontinued therapy due to drug-related adverse events.

In adult patients, the most frequently reported adverse events with an incidence of 3% or greater, excluding those due to the patient's concurrent condition, are listed in Table 5 by the total daily dose the patient was receiving at the time of the event. A total of 867 patients were treated with clarithromycin for mycobacterial infections. Of these, 43% reported one or more adverse events. Most of these events were described as mild to moderate in severity, although 14% were described as severe.

Incidence of adverse events was higher in patients taking 4000 mg total daily doses compared to lower doses (Table 5).

Table 5 - Percentage of Adverse Events* in Immunocompromised Adult Patients Treated with Clarithromycin for Mycobacterial Infections Presented by Total Daily Dose at Time of the Event

| System Organ Class | Adverse Reaction | 1000 mg (N=463) | 2000 mg (N=516) | 4000 mg (N=87) |
|-----------------------------|--------------------|-----------------|-----------------|----------------|
| Ear and labyrinth disorders | Hearing impaired** | 3% | 2% | 5% |
| Gastrointestinal disorders | Nausea | 11% | 16% | 40% |
| | Vomiting | 7% | 9% | 24% |
| | Abdominal Pain | 5% | 7% | 20% |
| | Diarrhea | 4% | 6% | 17% |
| | Flatulence | 1% | 2% | 7% |
| | Constipation | 1% | < 1% | 5% |

| System Organ Class | Adverse Reaction | 1000 mg (N=463) | 2000 mg (N=516) | 4000 mg (N=87) |
|---|--------------------------------------|--------------------|--------------------|-------------------|
| | Dry Mouth | < 1% | 0% | 5% |
| Investigations | Aspartate aminotransferase increased | 2% | 2% | 11% |
| | Alanine aminotransferase increased | 1% | 1% | 9% |
| Nervous system disorders | Dysgeusia | 6% | 7% | 29% |
| | Headache | 2% | 2% | 7% |
| Psychiatric disorders | Insomnia | < 1% | < 1% | 6% |
| Respiratory, thoracic and mediastinal disorders | Dyspnea | < 1% | < 1% | 7% |
| Skin and subcutaneous tissue disorders | Rash | 4% | 3% | 2% |

* Related adverse events considered to be definitely, probably, possibly or remotely related to study events.

** Sum of patients with deafness, ear disorder, partial transitory deafness, and/or tinnitus.

N= Number of adverse events.

Patients with *Helicobacter pylori* Infection

Triple Therapy: clarithromycin/omeprazole/amoxicillin

A summary of drug-related adverse event incidence rates is presented in Table 6.

Table 6 - Summary of Drug-Related Adverse Event Incidence Rates by System Organ Class

| System Organ Class | Patients With Drug-Related Adverse Events (% of Patients Treated)* | |
|--|---|---|
| | Omeprazole + Clarithromycin + Amoxicillin (N=137) | Omeprazole + Clarithromycin (N=130) |
| Cardiac disorders | 0 (0%) | 1 (1%) |
| Ear and labyrinth disorders | 1 (1%) | 2 (2%) |
| Eye disorders | 0 (0%) | 1 (1%) |
| Gastrointestinal disorders | 24 (18%) | 21 (16%) |
| General disorders and administration site conditions | 5 (4%) | 0 (0%) |
| Hepatobiliary disorders | 2 (1%) | 0 (0%) |
| Infections and infestations | 1 (1%) | 1 (1%) |
| Investigations | 9 (7%) | 0 (0%) |
| Nervous system disorders | 15 (11%) | 30 (23%) |
| Psychiatric disorders | 1 (1%) | 1 (1%) |

| System Organ Class | Patients With Drug-Related Adverse Events (% of Patients Treated)* | |
|---|---|---|
| | Omeprazole + Clarithromycin + Amoxicillin (N=137) | Omeprazole + Clarithromycin (N=130) |
| Reproductive system and breast disorders | 1 (1%) | 0 (0%) |
| Respiratory, thoracic and mediastinal disorders | 1(1%) | 0 (0%) |
| Skin and subcutaneous tissue disorders | 3 (2%) | 1 (1%) |

* Patients with more than 1 event within a system organ class are counted only once in the total for that system organ class

Note: there is a statistical difference (Fisher's exact two-sided, p-value = 0.009) between omeprazole + clarithromycin + amoxicillin (11%) versus omeprazole + clarithromycin (23%) in regard to nervous system disorders.

8.3 Less Common Clinical Trial Adverse Events

The following adverse drug reactions are applicable to all indications approved for this formulation.

Blood and Lymphatic System Disorders: eosinophilia and neutropenia

Gastrointestinal Disorders: abdominal distension

General Disorders and

Administration Site Conditions: chest pain, chills, fatigue, influenza and malaise

Hepatobiliary Disorders: cholestasis, gamma-glutamyltransferase increased and hepatitis

Investigations: blood alkaline phosphatase increased and blood lactate dehydrogenase increased

Other adverse reactions have been observed in different patient populations and during post-marketing surveillance. See [8 ADVERSE REACTIONS, Clinical Trial Adverse Drug Reactions, Table 3.](#)

8.4 Abnormal Laboratory Findings: Hematologic, Clinical Chemistry and Other Quantitative Data

Clinical Trial Findings

Patients with Respiratory Tract or Skin Infections

Changes in laboratory values with possible clinical significance reported during clinical studies or during post-marketing surveillance are displayed in Table 7.

Table 7 -Abnormal Hematologic and Clinical Chemistry Findings in Patients with Respiratory Tract or Skin Infections Treated with Clarithromycin Tablets

| System Organ Class | Laboratory Values | Frequency |
|--------------------|---|----------------------------|
| Investigations | Alanine aminotransferase increased Aspartate aminotransferase increased Gamma-glutamyltransferase increased Blood alkaline phosphatase increased Blood lactate dehydrogenase increased Blood bilirubin increased Blood creatinine increased White blood cell count decreased | Uncommon (Less than 1%) |
| | Prothrombin time prolonged Blood urea increased | 1% 4% |

Patients with Mycobacterial Infections

In immunocompromised patients receiving prophylaxis against *M. avium*, those laboratory values outside the extreme high or low limit for the specified test were analyzed (Table 8).

Table 8 - Percentage of Patients* Exceeding Extreme Laboratory Value in Immunocompromised Patients Receiving Prophylaxis Against *M. avium* Complex

| System Organ Class | Laboratory Values | Clarithromycin 500 mg BID | | Placebo | |
|--------------------|--|------------------------------|----|---------|----|
| | | | | | |
| Investigations | Hemoglobin decreased < 8 g/dL | 4/118 | 3% | 5/103 | 5% |
| | Platelet count decreased < 50 x 10 ⁹ /L | 11/249 | 4% | 12/250 | 5% |
| | White blood cell count decreased < 1 x 10 ⁹ /L | 2/103 | 4% | 0/95 | 0% |
| | Aspartate aminotransferase increased > 5 x ULN | 7/196 | 4% | 5/208 | 2% |
| | Alanine aminotransferase increased > 5 x ULN | 6/217 | 3% | 4/232 | 2% |

| System Organ Class | Laboratory Values | Clarithromycin 500 mg BID | | Placebo | |
|--------------------|---|------------------------------|----|---------|----|
| | | | | | |
| | Blood alkaline phosphatase increased > 5 x ULN | 5/220 | 2% | 5/218 | 2% |

* Includes only patients with baseline values within the normal range or borderline high (hematology variables) and within the normal range or borderline low (chemistry variables).

Legend: BID = twice daily; ULN = Upper Limit of Normal

Treatment of Patients with Mycobacterial Infections

In immunocompromised patients treated with clarithromycin for mycobacterial infections, evaluations of laboratory values were made by analysing those values outside the seriously abnormal level (i.e., the extreme high or low limit) for the specified test (Table 9).

Table 9 - Percentage of Immunocompromised Adult Patients Treated with Clarithromycin for Mycobacterial Infections who had On-Treatment Laboratory Values that Were Outside the Seriously Abnormal Level Presented by Total Daily Dose

| System Organ Class | Laboratory Values | Seriously Abnormal Level | 1000 mg | 2000 mg | 4000 mg |
|--------------------|--------------------------------------|---------------------------|---------|---------|---------|
| Investigations | Aspartate aminotransferase increased | > 5 x ULN | 3% | 2% | 4% |
| | Alanine aminotransferase increased | > 5 x ULN | 2% | 2% | 7% |
| | Platelet count decreased | < 50 x 10 ⁹ /L | 2% | 2% | 4% |
| | White blood cell count decreased | < 1 x 10 ⁹ /L | 0% | 2% | 0% |
| | Blood urea increased | > 50 mg/dL | < 1% | < 1% | 4% |

Legend: ULN = Upper Limit of Normal

8.5 Post-Market Adverse Reactions

The following list of adverse events is a compilation of adverse reactions from Post-marketing Surveillance and Post-marketing Clinical Studies for all clarithromycin formulations.

Table 10 - Post-Market Adverse Drug Reactions

| System Organ Class | Adverse Event |
|--------------------------------------|---|
| Blood and lymphatic system disorders | Agranulocytosis, leukopenia, thrombocytopenia |

| System Organ Class | Adverse Event |
|--|---|
| Cardiac disorders ¹ | Atrial fibrillation, cardiac arrest, electrocardiogram QT prolonged, extrasystoles, palpitations, Torsades de pointes, ventricular fibrillation, ventricular tachycardia |
| Ear and labyrinth disorders | Deafness, hearing impaired, hearing loss ² , tinnitus, vertigo |
| Gastrointestinal disorders | Abdominal pain, constipation, dry mouth, dyspepsia, eructation, esophagitis, flatulence, gastritis, glossitis, pancreatitis, stomatitis, tongue discolouration, tooth discolouration, vomiting |
| General disorders and administration site conditions | Asthenia |
| Hepatobiliary disorders | Hepatic failure ³ , hepatitis, hepatitis cholestatic, jaundice (cholestatic and hepatocellular) |
| Immune system disorders | Angioedema, anaphylactic reaction, anaphylactoid reaction, anaphylaxis, hypersensitivity, myasthenia gravis |
| Infections and infestations | Candidiasis, cellulitis, pseudomembranous colitis, vaginal infection |
| Investigations | Albumin globulin ratio abnormal, alanine aminotransferase increased, aspartate aminotransferase increased, blood creatinine increased, blood urea increased, international normalized ratio (INR) increased ⁴ , liver enzymes increased, liver function test abnormal, prothrombin time prolonged ⁴ , urine color abnormal ⁵ |
| Metabolism and nutrition disorders | Anorexia, decreased appetite |
| Musculoskeletal and connective tissue disorders | Musculoskeletal stiffness, myalgia, myopathy, rhabdomyolysis ⁷ |
| Nervous system disorders | Ageusia, alteration of sense of smell, anosmia, convulsions, dizziness, dysgeusia, dyskinesia, headache, loss of consciousness, paraesthesia, parosmia, tremor, somnolence |
| Psychiatric disorders | Abnormal dreams, anxiety, confusion, depersonalization, depression, disorientation, hallucination, insomnia, mania, psychosis |
| Renal and urinary disorders | Interstitial nephritis, renal failure |
| Respiratory, thoracic and mediastinal disorders | Asthma, pulmonary embolism |
| Skin and subcutaneous tissue disorders | Severe cutaneous adverse reactions (SCAR) (e.g., acute generalized exanthematous |

| System Organ Class | Adverse Event |
|--------------------|---|
| | pustulosis (AGEP), Stevens-Johnson syndrome (SJS), toxic epidermal necrolysis (TEN), drug rash with eosinophilia and systemic symptoms (DRESS)), acne, dermatitis bullous, Henoch-Schonlein purpura, hyperhidrosis, pruritus, rash, urticaria |
| Vascular disorders | Hemorrhage ⁴ , vasodilation |

- 1 As with other macrolides, QT prolongation, ventricular tachycardia, and torsades de pointes have been reported with clarithromycin.
- 2 There have been reports of hearing loss with clarithromycin which is usually reversible upon withdrawal of therapy.
- 3 Hepatic dysfunction may be severe and is usually reversible. Hepatic failure with fatal outcome has been reported and generally has been associated with serious underlying diseases and/or concomitant medications.
- 4 When clarithromycin is co-administered with warfarin.
- 5 Symptom of hepatic failure.
- 6 In some of the reports of rhabdomyolysis, clarithromycin was administered concomitantly with other drugs known to be associated with rhabdomyolysis (such as statins, fibrates, colchicine or allopurinol).

Colchicine

There have been post-marketing reports of colchicine toxicity with concomitant use of clarithromycin and colchicine, especially in the elderly, some of which occurred in patients with renal insufficiency. Deaths have been reported in some patients. See [2 CONTRAINDICATIONS](#).

9 DRUG INTERACTIONS

9.1 Serious Drug Interactions

| Serious Drug Interactions |
|---|
| <ul style="list-style-type: none"> • Concomitant administration of clarithromycin with astemizole, cisapride, domperidone, colchicine, pimozide, ivabradine, terfenadine, lovastatin, simvastatin, ergot alkaloids (e.g., ergotamine, dihydroergotamine) is contraindicated. See 2 CONTRAINDICATIONS and 9 DRUG INTERACTIONS, 9.4 Drug-Drug Interactions. • Clarithromycin is an inhibitor of the cytochrome P450 3A isoform subfamily (CYP3A) and the P-glycoprotein transporter (P-gp). The concomitant administration of clarithromycin and drugs metabolized by CYP3A and/or transported by P-gp may lead to an increase in the plasma concentrations of the co-administered drug which could result in clinically significant safety concerns. |

9.2 Drug Interactions Overview

Many categories of drugs are metabolized by CYP3A and/or transported by P-gp located in the liver and in the intestine. Some drugs may inhibit or induce the activities of CYP3A and/or P-gp. Administration of such inhibitors or inducers may impact upon the metabolism. In some cases serum concentrations may be increased and in others decreased. Care must therefore be exercised when co-administering such drugs.

Effects of Clarithromycin on Other Drugs

Clarithromycin is an inhibitor of CYP3A and P-gp. This inhibition may lead to increased or prolonged serum levels of those drugs also metabolized by CYP3A or transported by P-gp when co-administered with clarithromycin. For such drugs the monitoring of their serum concentrations may be necessary.

Clarithromycin should be used with caution in patients receiving treatment with other drugs known to be CYP3A and/or P-gp substrates, especially if the CYP3A/P-gp substrate has a narrow safety margin (e.g., carbamazepine) and/or the substrate is extensively metabolized by CYP3A or transported by P-gp. Dosage adjustments may be considered, and when possible, serum concentrations of these drugs should be monitored closely in patients concurrently receiving clarithromycin.

Clarithromycin should be used with caution in patients receiving treatment with hydroxychloroquine and chloroquine as these medicines are known to prolong the QT interval due to the potential to induce cardiac arrhythmia and serious adverse cardiovascular events.

The following drugs or drug classes are known or suspected to be metabolized by the same CYP3A isozyme: alprazolam, astemizole, carbamazepine, cilostazol, cisapride, corticosteroids (eg. Methylprednisolone), cyclosporine, disopyramide, domperidone, ergot alkaloids, ibrutinib, ivabradine, lomitapide, lovastatin, midazolam, omeprazole, oral anticoagulants (e.g. warfarin, rivaroxaban, apixaban), atypical antipsychotics (e.g. quetiapine), pimozide, quinidine, rifabutin, sildenafil, simvastatin, tacrolimus, terfenadine, triazolam and vinblastine, but this list is not comprehensive. See [9 DRUG INTERACTIONS, 9.4 Drug-Drug Interactions](#).

Direct acting oral anticoagulants (DOACs): The DOACs dabigatran and edoxaban are substrates for the efflux transporter P-gp. Rivaroxaban and apixaban are metabolised via CYP3A4 and are also substrates for P-gp. Caution should be exercised when clarithromycin is co-administered with these agents particularly to patients at high risk of bleeding. See [7 WARNINGS AND PRECAUTIONS, General](#).

With certain drugs, co-administration of clarithromycin is contraindicated or should be avoided (see [Table 11](#)).

Effects of Other Drugs on Clarithromycin

Clarithromycin is a substrate of CYP3A. Co-administration of strong inducers of the cytochrome

P450 metabolism system may accelerate the metabolism of clarithromycin and thus lower exposure to clarithromycin while increasing exposure to its metabolite 14-OH-clarithromycin which could impair the intended therapeutic effect. Furthermore, it might be necessary to monitor the plasma levels of the CYP3A inducer, which could be increased owing to the inhibition of CYP3A by clarithromycin (see also the relevant product information for the CYP3A4 inducer administered). Co-administration of potent CYP3A inhibitors may lead to increased exposure to clarithromycin and decreased exposure to its metabolite 14-OH-clarithromycin. Clarithromycin dosage adjustment or consideration of alternative treatments may be required.

Bi-Directional Drug Interactions

Bi-directional drug interactions are complex and may occur if both of the interacting drugs are substrates and inhibitors/inducers of CYP3A.

Additional Mechanisms

Interactions with clarithromycin have been reported with drugs metabolized by cytochrome P450 isoforms other than CYP3A system. Additional mechanisms, such as effects upon absorption, may also be responsible for interaction between drugs, including zidovudine and clarithromycin.

9.3 Drug-Behavioural Interactions

This information is not available for this drug product.

9.4 Drug-Drug Interactions

Some of the drug-drug interactions which have been reported between clarithromycin-macrolides and other drugs or drug categories are listed in [Table 11](#).

The drugs listed in this table are based on either drug interaction case reports or studies, or potential interactions due to the expected magnitude and seriousness of the interaction (i.e., those identified as contraindicated).

Table 11 - Established or Potential Drug-Drug Interactions with Clarithromycin

| Concomitant Medication | Source of Evidence | Effect | Clinical Comments |
|-------------------------------|---------------------------|--|--|
| Astemizole*/ Terfenadine | CT | terfenadine-acid metabolite concentrations increase ↑ QT interval | Macrolides have been reported to alter the metabolism of terfenadine resulting in increased serum levels of terfenadine which has occasionally been associated with cardiac arrhythmias such as QT prolongation, ventricular tachycardia, ventricular fibrillation and torsades de |

| Concomitant Medication | Source of Evidence | Effect | Clinical Comments |
|------------------------|--------------------|--|---|
| | | | <p>pointes (see 2 CONTRAINDICATIONS).</p> <p>In a study involving 14 healthy volunteers, the concomitant administration of clarithromycin film-coated tablets and terfenadine resulted in a 2 to 3-fold increase in the serum levels of the acid metabolite of terfenadine, MDL 16, 455, and in prolongation of the QT interval. Similar effects have been observed with concomitant administration of astemizole and other macrolides.</p> |
| Atazanavir | CT | <p>↑ clarithromycin levels</p> <p>↑ atazanavir AUC</p> | <p>Both clarithromycin and atazanavir are substrates and inhibitors of CYP3A, and there is evidence of a bi-directional drug interaction. Co-administration of clarithromycin (500 mg twice daily) with atazanavir (400 mg once daily) resulted in a 2-fold increase in exposure to clarithromycin and a 70% decrease in exposure to 14-OH-clarithromycin, with a 28% increase in the AUC of atazanavir.</p> <p>Because of the large therapeutic window for clarithromycin, no dosage reduction should be necessary in patients with normal renal function. For patients with moderate renal function (creatinine clearance 30 to 60 mL/min), the dose of clarithromycin should be decreased by 50%. For patients</p> |

| Concomitant Medication | Source of Evidence | Effect | Clinical Comments |
|---|--------------------|---|--|
| | | | with creatinine clearance < 30 mL/min, the dose of clarithromycin should be decreased by 75% using an appropriate clarithromycin formulation. Doses of clarithromycin greater than 1000 mg per day should not be co-administered with protease inhibitors. |
| <u>Atypical Antipsychotics</u> (e.g. quetiapine) | | Potential ↑ in concentrations of quetiapine and other atypical antipsychotics | <p>Clarithromycin should not be used in combination with quetiapine unless clinically necessary. Due to CYP3A inhibition by clarithromycin, concentrations of quetiapine are expected to increase, which can result in serious and/or life-threatening adverse reactions, including malignant neuroleptic syndrome.</p> <p>For other atypical antipsychotic drugs (aripiprazole and risperidone) metabolized by CYP3A4, it is also recommended that concomitant administration with clarithromycin be avoided due to potential pharmacokinetic interactions.</p> |
| <u>Calcium Channel Blockers</u> (e.g., verapamil, amlodipine, diltiazem) | C | Potential ↑ in verapamil concentrations | Caution is advised regarding the concomitant administration of clarithromycin and calcium channel blockers metabolized by CYP3A4 (e.g., verapamil, amlodipine, diltiazem) due to the risk of hypotension. Plasma concentrations of clarithromycin as well as calcium channel blockers may increase due to the interaction. Hypotension, |

| Concomitant Medication | Source of Evidence | Effect | Clinical Comments |
|-------------------------|--------------------|---|---|
| | | | bradyarrhythmias, and lactic acidosis have been observed in patients receiving concurrent verapamil, belonging to the calcium channel blockers drug class. |
| Carbamazepine | C | ↑ levels of carbamazepine | Clarithromycin administration in patients receiving carbamazepine has been reported to cause increased levels of carbamazepine. Blood level monitoring of carbamazepine should be considered. |
| Cisapride*/ Pimozide | C | ↑ levels of cisapride ↑ levels of pimozide | Elevated cisapride levels have been reported in patients receiving clarithromycin and cisapride concomitantly. This may result in QT prolongation and cardiac arrhythmias including ventricular tachycardia, ventricular fibrillation and torsade de pointes. Similar effects have been observed in patients taking clarithromycin and pimozide concomitantly (see 2 CONTRAINDICATIONS). |
| Colchicine | C | Potential colchicine toxicity | Colchicine is a substrate for both CYP3A and the efflux transporter, P-gp. Clarithromycin and other macrolides are known to inhibit CYP3A and P-gp. When clarithromycin and colchicine are administered together, inhibition of P-gp and/or CYP3A by clarithromycin may lead to increased exposure to colchicine. This risk may be further increased with concomitant medications |

| Concomitant Medication | Source of Evidence | Effect | Clinical Comments |
|------------------------|--------------------|--|--|
| | | | metabolized by P-glycoprotein or strong CYP3A inhibitors. Concomitant use of clarithromycin and colchicine is contraindicated. See 2 CONTRAINDICATIONS . |
| Cyclosporine | C | ↑ levels of cyclosporine | There have been reports of elevated cyclosporine serum concentrations when clarithromycin and cyclosporine are used concurrently. Cyclosporine levels should be monitored and the dosage should be adjusted as necessary. Patients should also be monitored for increased cyclosporine toxicity. |
| Didanosine | CT | No change in didanosine pharmacokinetics in HIV-infected patients (n=12) | Simultaneous administration of clarithromycin film-coated tablets and didanosine to 12 HIV-infected adult patients resulted in no statistically significant change in didanosine pharmacokinetics. |
| Digoxin | C | ↑ levels of digoxin | Digoxin is thought to be a substrate for the efflux transporter, P-gp. Clarithromycin is known to inhibit P-gp. When clarithromycin and digoxin are administered together, inhibition of P-gp by clarithromycin may lead to increased exposure to digoxin. Elevated digoxin serum concentrations have been reported in patients receiving clarithromycin film-coated tablets and digoxin concomitantly. |

| Concomitant Medication | Source of Evidence | Effect | Clinical Comments |
|----------------------------|--------------------|---|---|
| | | | <p>In post-marketing surveillance some patients have shown clinical signs consistent with digoxin toxicity, including potentially fatal arrhythmias. Serum digoxin levels should be carefully monitored while patients are receiving digoxin and clarithromycin simultaneously.</p> |
| Disopyramide/ Quinidine | C | <p>↑ levels of disopyramide, resulting in ventricular fibrillation & QT prolongation (rarely reported)</p> <p>Torsades de pointes</p> | <p>Increased disopyramide plasma levels, resulting in ventricular fibrillation and QT prolongation, coincident with the co-administration of disopyramide and clarithromycin have rarely been reported.</p> <p>There have been post-marketed reports of torsades de pointes occurring with concurrent use of clarithromycin and quinidine or disopyramide. Electrocardiograms should be monitored for QTc prolongation during co-administration of clarithromycin with these drugs. Serum levels of these medications should be monitored during clarithromycin therapy.</p> <p>There have been post marketing reports of hypoglycemia with the concomitant administration of clarithromycin and disopyramide. Therefore blood glucose levels should be monitored during concomitant administration of clarithromycin and disopyramide.</p> |

| Concomitant Medication | Source of Evidence | Effect | Clinical Comments |
|--|--------------------|---|---|
| Domperidone | C, P | ↑ levels of domperidone, resulting in QT prolongation and cardiac arrhythmias | Elevated domperidone levels have been reported in patients receiving a potent CYP3A4 inhibitor and domperidone concomitantly. This may result in QT prolongation and cardiac arrhythmias including ventricular tachycardia, ventricular fibrillation and torsades de pointes. Hence, co-administration of domperidone with QT-prolonging medicines and/or potent CYP3A4 inhibitors such as clarithromycin is contraindicated. See 2 CONTRAINDICATIONS . |
| <u>Ergot alkaloids</u> Ergotamine/ Dihydroergotamine | C | Potential ischemic reactions Potential ergot toxicity | Post-marketing reports indicate that co-administration of clarithromycin with ergotamine or dihydroergotamine has been associated with acute ergot toxicity characterized by severe peripheral vasospasm, dysesthesia, and ischemia of the extremities and other tissues including the central nervous system. Concomitant administration of clarithromycin and ergot alkaloids is contraindicated (see 2 CONTRAINDICATIONS). |
| Etravirine | CT | ↓ clarithromycin ↑ 14-OH-clarithromycin | Clarithromycin exposure was decreased by etravirine; however, concentrations of the active metabolite, 14-OH-clarithromycin, were increased. Because 14-OH-clarithromycin has reduced activity against <i>Mycobacterium avium</i> complex (MAC), overall activity against this pathogen may be altered; |

| Concomitant Medication | Source of Evidence | Effect | Clinical Comments |
|--|--------------------|---|--|
| | | | therefore alternatives to clarithromycin should be considered for the treatment of MAC. |
| Fluconazole | CT | ↑ clarithromycin C _{min} & AUC | <p>Concomitant administration of fluconazole 200 mg daily and clarithromycin 500 mg twice daily to 21 healthy volunteers led to increases in the mean steady-state clarithromycin C_{min} and AUC of 33% and 18%, respectively.</p> <p>Steady-state concentrations of 14-OH clarithromycin were not significantly affected by concomitant administration of fluconazole. No clarithromycin dose adjustment is necessary.</p> |
| <p><u>HMG-CoA Reductase Inhibitors</u></p> <p>Lovastatin/ Simvastatin</p> <p>Atorvastatin Rosuvastatin</p> | <p>C</p> <p>C</p> | Rhabdomyolysis (rarely reported) | <p>Concomitant use of clarithromycin with lovastatin or simvastatin is contraindicated (See 2 CONTRAINDICATIONS) as these statins are extensively metabolized by CYP3A4 and concomitant treatment with clarithromycin increases their plasma concentration, which increases the risk of myopathy, including rhabdomyolysis. Reports of rhabdomyolysis have been received for patients taking clarithromycin concomitantly with these statins. If treatment with clarithromycin cannot be avoided, therapy with lovastatin or simvastatin must be suspended during the course of treatment. See 7 WARNINGS AND PRECAUTIONS, HMG-CoA Reductase Inhibitors.</p> |

| Concomitant Medication | Source of Evidence | Effect | Clinical Comments |
|------------------------|--------------------|--|---|
| | | | <p>Rare reports of rhabdomyolysis have also been reported in patients taking atorvastatin or rosuvastatin concomitantly with clarithromycin. Concurrent use of atorvastatin and clarithromycin may result in increased atorvastatin exposure.</p> <p>Caution should be exercised when prescribing clarithromycin with statins. In situations where the concomitant use of clarithromycin with statins cannot be avoided, it is recommended to prescribe the lowest registered dose of the statin. Use of a statin that is not dependent on CYP3A metabolism (e.g., fluvastatin) can be considered. Patients should be monitored for signs and symptoms of myopathy.</p> |
| Itraconazole | CT, P | ↑ levels of clarithromycin ↑ levels of itraconazole | Both clarithromycin and itraconazole are substrates and inhibitors of CYP3A, leading to a bi-directional drug interaction. Clarithromycin may increase the plasma levels of itraconazole, while itraconazole may increase the plasma levels of clarithromycin. Patients taking itraconazole and clarithromycin concomitantly should be monitored closely for signs or symptoms of increased or prolonged pharmacologic effect. |
| Ivabradine | T | Potential increase in serum concentration | May significantly increase blood levels and effects of ivabradine; may lead to excessive |

| Concomitant Medication | Source of Evidence | Effect | Clinical Comments |
|---|--------------------|---|--|
| | | | bradycardia. See 2 CONTRAINDICATIONS |
| Lansoprazole / Omeprazole | CT | Mild change of lansoprazole and 14-OH clarithromycin concentrations ↑ omeprazole C _{max} & AUC ₀₋₂₄ ↑ levels of clarithromycin | One study demonstrated that concomitant administration of clarithromycin and lansoprazole resulted in mild changes of serum concentrations of lansoprazole and 14-OH clarithromycin. However, no dosage adjustment is considered necessary based on these data. Clarithromycin 500 mg three times daily was given in combination with omeprazole 40 mg once daily to healthy subjects. The steady-state plasma concentrations of omeprazole were increased (i.e., C _{max} , AUC ₀₋₂₄ , and t _{1/2} increased by 30%, 89%, and 34%, respectively), by concomitant administration of clarithromycin. The mean 24-hour gastric pH value was 5.2 when omeprazole was administered alone and 5.7 when co-administered with clarithromycin. To a lesser extent, omeprazole administration increases the serum concentrations of clarithromycin. Omeprazole administration also increases tissue and mucus concentrations of clarithromycin. |
| <u>Oral</u> <u>Anticoagulants</u> Warfarin / Acenocoumarol | C | ↑ anticoagulant effect | There have been reports of increased anticoagulant effect when clarithromycin and oral anticoagulants are used concurrently. Anticoagulant |

| Concomitant Medication | Source of Evidence | Effect | Clinical Comments |
|---|---------------------|---------------------|--|
| | | | <p>parameters should be closely monitored. Adjustment of the anticoagulant dose may be necessary.</p> <p>Clarithromycin has also been reported to increase the anticoagulant effect of acenocoumarol.</p> <p>There is a risk of serious hemorrhage and significant elevations in International Normalized Ratio (INR) and prothrombin time when clarithromycin is co-administered with warfarin. INR and prothrombin times should be frequently monitored while patients are receiving clarithromycin and oral anticoagulants concurrently. See 7 WARNINGS AND PRECAUTIONS, Use with Other Drugs, Oral Anticoagulants.</p> |
| <p><u>Oral Hypoglycemic Agents</u> (e.g. insulin)</p> | <p>C P</p> | <p>Hypoglycemia</p> | <p>The concomitant use of clarithromycin and oral hypoglycaemic agents (such as sulphonylurias) and/or insulin can result in significant hypoglycaemia. With certain hypoglycaemic drugs such as nateglinide, pioglitazone, repaglinide and rosiglitazone, inhibition of CYP3A enzyme by clarithromycin may be involved and could cause hypoglycaemia when used concomitantly. Careful monitoring of glucose is recommended.</p> |
| <p><u>Phospho-</u></p> | <p>P</p> | <p>↑</p> | <p>Sildenafil, tadalafil, and</p> |

| Concomitant Medication | Source of Evidence | Effect | Clinical Comments |
|---|--------------------|---|---|
| <p><u>diesterase inhibitors</u> (e.g., sildenafil, tadalafil, vardenafil)</p> | | <p>phosphodiesterase inhibitor exposure</p> | <p>vardenafil are metabolized, at least in part, by CYP3A, and CYP3A may be inhibited by concomitantly administered clarithromycin. Coadministration of clarithromycin with sildenafil, tadalafil or vardenafil would likely result in increased phosphodiesterase inhibitor exposure. Reduction of sildenafil, tadalafil and vardenafil dosages should be considered when these drugs are co-administered with clarithromycin.</p> |
| <p>Rifabutin</p> | <p>C</p> | <p>↓ clarithromycin ↑ rifabutin</p> | <p>Clarithromycin has been reported to increase serum and tissue concentration of rifabutin and thus may increase the risk of toxicity. Clarithromycin levels decrease when co-administered with rifabutin.</p> <p>Concomitant administration of clarithromycin and rifabutin in the treatment of <i>Mycobacterial Avium</i> complex infections resulted in rifabutin-associated uveitis.</p> <p>A case control study in AIDS patients showed that concomitant administration of rifabutin and clarithromycin resulted in an approximately 50% reduction in serum clarithromycin concentration, approximately 77% increase in the area under the plasma concentration-time curve of rifabutin, and a 236% increase in</p> |

| Concomitant Medication | Source of Evidence | Effect | Clinical Comments |
|-------------------------|--------------------|---|--|
| | | | <p>the area under the plasma concentration-time curve of rifabutin's active metabolite. The increase in rifabutin and/or its metabolite contributed to the development of uveitis (the incidence of uveitis was 14% in patients weighing > 65 kg, 45% in patients between 55 and 65 kg, and 64% in patients < 55 kg).</p> |
| Ritonavir/ Indinavir | CT | <p>↑ clarithromycin C_{max}, C_{min}, & AUC</p> <p>↑ indinavir AUC ↑ clarithromycin AUC</p> | <p>A pharmacokinetic study demonstrated that the concomitant administration of ritonavir 200 mg every 8 hours and clarithromycin 500 mg every 12 hours resulted in a marked inhibition of the metabolism of clarithromycin. The clarithromycin C_{max} increased by 31%, C_{min} increased 182% and AUC increased by 77% with concomitant administration of ritonavir. An essentially complete inhibition of the formation of 14-[R]-hydroxy-clarithromycin was noted. Because of the large therapeutic window for clarithromycin, no dosage reduction should be necessary in patients with normal renal function. However, for patients with renal impairment, the following dosage adjustments should be considered: For patients with creatinine clearance 30 to 60 mL/min the dose of clarithromycin should be reduced by 50%. For patients with creatinine clearance <</p> |

| Concomitant Medication | Source of Evidence | Effect | Clinical Comments |
|------------------------|--------------------|---|---|
| | | | <p>30 mL/min the dose of clarithromycin should be decreased by 75%. Doses of clarithromycin greater than 1 g/day should not be co-administered with ritonavir.</p> <p>Similar dose adjustments should be considered in patients with reduced renal function when ritonavir is used as a pharmacokinetic enhancer with other HIV protease inhibitors including atazanavir and saquinavir.</p> <p>One study demonstrated that the concomitant administration of clarithromycin and indinavir resulted in a metabolic interaction; the clarithromycin AUC increased by 53% and the indinavir AUC was increased by 20%, but the individual variation was large. No dose adjustment is necessary with normal renal function.</p> |
| Saquinavir | CT | <p>↑ saquinavir AUC and C_{max}</p> <p>↑ clarithromycin AUC</p> | <p>Both clarithromycin and saquinavir are substrates and inhibitors of CYP3A, and there is evidence of a bi-directional drug interaction.</p> <p>Concomitant administration of clarithromycin (500 mg twice daily) and saquinavir (soft gelatin capsules, 1200 mg three times daily) for 7 days to 12 healthy volunteers resulted in steady-state AUC and C_{max} values of saquinavir which were</p> |

| Concomitant Medication | Source of Evidence | Effect | Clinical Comments |
|------------------------|--------------------|--|--|
| | | | <p>177% (108-269%) and 187% (105-300%) higher than those seen with saquinavir alone. Clarithromycin AUC and C_{max} values were approximately 40% higher than those seen with clarithromycin alone.</p> <p>[Clarithromycin AUC ↑ 45% (17-81%) and C_{max} ↑ 39% (10-76%); 14-OH clarithromycin metabolite AUC ↓ 24% (5-40%) and C_{max} ↓ 34% (14-50%)].</p> <p>QTc prolongation has been reported in patients taking saquinavir along with ritonavir and also in patients taking clarithromycin. Concurrent administration of saquinavir and clarithromycin is contraindicated (see 2 CONTRAINDICATIONS).</p> |
| Tacrolimus | P | Potential ↑ in tacrolimus concentrations | Concomitant administration of tacrolimus and clarithromycin may result in increased plasma levels of tacrolimus and increased risk of toxicity. |
| Theophylline | P | Potential ↑ in theophylline concentrations | <p>Clarithromycin use in patients who are receiving theophylline may be associated with an increase of serum theophylline concentrations.</p> <p>Monitoring of serum theophylline concentrations should be considered for patients receiving high doses of theophylline or with baseline concentrations in the upper therapeutic range.</p> |
| Tolterodine | P | ↑ serum tolterodine | The primary route of metabolism for tolterodine is via |

| Concomitant Medication | Source of Evidence | Effect | Clinical Comments |
|---|--------------------|-----------------|---|
| | | concentrations | the 2D6 isoform of cytochrome P450 (CYP2D6). However, in a subset of the population devoid of CYP2D6, the identified pathway of metabolism is via CYP3A. In this population subset, inhibition of CYP3A results in significantly higher serum concentrations of tolterodine. A reduction of tolterodine dosage may be necessary in the presence of CYP3A inhibitors, such as clarithromycin in the CYP2D6 poor metabolizer population. |
| <p><u>Triazolobenzodiazepines</u> (e.g., triazolam, alprazolam)</p> <p><u>Other related benzodiazepines</u> (e.g., midazolam)</p> | CT, C, P | ↑ midazolam AUC | <p>When midazolam was co-administered with clarithromycin tablets (500 mg twice daily), midazolam AUC was increased 2.7-fold after intravenous administration of midazolam and 7-fold after oral administration. Concomitant administration of oral midazolam and clarithromycin is contraindicated. (See 2 CONTRAINDICATIONS). If intravenous midazolam is co-administered with clarithromycin, the patient must be closely monitored to allow dose adjustment of midazolam. A drug-drug interaction study between oromucosal midazolam and clarithromycin has not been conducted.</p> <p>The same precautions should also apply to other benzodiazepines that are metabolized by CYP3A, including</p> |

| Concomitant Medication | Source of Evidence | Effect | Clinical Comments |
|---|--------------------|---|---|
| | | | <p>triazolam and alprazolam. For benzodiazepines which are not dependent on CYP3A for their elimination (temazepam, nitrazepam, lorazepam), a clinically important interaction with clarithromycin is unlikely.</p> <p>There have been post-marketing reports of drug interactions and central nervous system (CNS) effects (e.g., somnolence and confusion) with the concomitant use of clarithromycin and triazolam. Monitoring the patient for increased CNS pharmacological effects is suggested.</p> |
| Zidovudine | C | Potential ↓ in zidovudine concentrations | <p>Simultaneous oral administration of clarithromycin film-coated tablets and zidovudine to HIV-infected adult patients may result in decreased steady-state zidovudine concentrations. Clarithromycin appears to interfere with the absorption of simultaneously administered oral zidovudine, and therefore this interaction can be largely avoided by staggering the doses of clarithromycin and zidovudine. Similar interaction studies have not been conducted with clarithromycin extended-release (ER) and zidovudine.</p> |
| <u>Other drugs metabolized by CYP3A</u> (e.g., alfentanil, bromocriptine, | C, P | Potential increase in serum concentration | <p>Interactions with erythromycin and/or clarithromycin have been reported with a number of other drugs metabolized by CYP3A, such as alfentanil,</p> |

| Concomitant Medication | Source of Evidence | Effect | Clinical Comments |
|--|--------------------|---|--|
| cilostazol, methylprednisolone, vinblastine) | | | <p>bromocriptine, cilostazol, ibrutinib, methylprednisolone or vinblastine.</p> <p>Serum concentrations of drugs metabolized by CYP3A should be monitored closely in patients concurrently receiving erythromycin or clarithromycin.</p> |
| <u>Other drugs metabolized by cytochrome P450 isoforms other than CYP3A</u> (e.g., hexobarbital, phenytoin, and valproate) | C, P | Potential change in serum concentration | <p>Interactions with erythromycin and/or clarithromycin have been reported with drugs metabolized by other cytochrome P450 isoforms (i.e., not CYP3A), such as hexobarbital, phenytoin, and valproate.</p> <p>Serum concentrations of these drugs should be monitored closely in patients concurrently receiving erythromycin or clarithromycin.</p> |
| <u>Other drug inducers of the cytochrome P450 system</u> (e.g, efavirenz, nevirapine, rifampin, rifabutin, rifampicin, phenobarbital, rifapentine) | CT, P | ↓ levels of clarithromycin | <p>Strong inducers of the cytochrome P450 metabolism system such as efavirenz, nevirapine, rifampin, rifabutin, rifampicin, phenobarbital and rifapentine* may accelerate the metabolism of clarithromycin and thus lower the plasma levels of clarithromycin, while increasing those of 14-OH-clarithromycin, a metabolite that is also microbiologically active.</p> <p>Since the microbiological activities of clarithromycin and 14-OH-clarithromycin are different for different bacteria, the intended therapeutic effect could be impaired during</p> |

| Concomitant Medication | Source of Evidence | Effect | Clinical Comments |
|------------------------|--------------------|--------|---|
| | | | concomitant administration of clarithromycin and enzyme inducers. |

Legend: C = Case Study; CT = Clinical Trial; P = Potential Interactions with other drugs have not been established.

*not marketed in Canada

Combination Therapy with Omeprazole and/or Amoxicillin

For more information on drug interactions for omeprazole and amoxicillin, refer to their respective Product Monographs, under DRUG INTERACTIONS.

9.5 Drug-Food Interactions

CLARITHROMYCIN may be given with or without meals.

Interactions with food have not been established.

9.6 Drug-Herb Interactions

St. John's Wort (*Hypericum perforatum*) is an inducer of CYP3A and may induce the metabolism of clarithromycin. This may result in sub-therapeutic levels of clarithromycin leading to reduced efficacy.

9.7 Drug-Laboratory Test Interactions

Interactions with laboratory tests have not been established.

10 CLINICAL PHARMACOLOGY

10.1 Mechanism of Action

General

Clarithromycin exerts its antibacterial action by binding to the 50S ribosomal subunit of susceptible bacteria and suppressing protein synthesis.

10.2 Pharmacodynamics

Eradication of *Helicobacter pylori*

H. pylori is now established as a major etiological factor in duodenal ulcer disease. The presence of *H. pylori* may damage the mucosal integrity due to the production of enzymes (catalase, lipases, phospholipases, proteases, and urease), adhesins and toxins; the generated inflammatory response contributes to mucosal damage.

The concomitant administration of an antimicrobial(s) such as clarithromycin and an antisecretory agent, improves the eradication of *H. pylori* as compared to individual drug administration. The higher pH resulting from antisecretory treatment optimizes the environment for the pharmacologic action of the antimicrobial agent(s) against *H. pylori*.

10.3 Pharmacokinetics

A summary of clarithromycin pharmacokinetic parameters following the administration of clarithromycin film-coated tablets is provided in Table 12. See [10 CLINICAL PHARMACOLOGY, DETAILED PHARMACOLOGY](#).

Table 12 - Clarithromycin Pharmacokinetic Parameters Following the Administration of Clarithromycin Film-coated Tablets

| Single dose* | C _{max} (mg/L) | t _{max} (hr) | t _½ (hr) | AUC _{0-t} (mg•hr/L) |
|-----------------------------|----------------------------|--------------------------|------------------------|---------------------------------|
| 250 mg Mean | 1 | 1.5 | 2.7 | 5.47 |
| 500 mg Mean | 1.77 | 2.2 | --- | 11.66 |
| Multiple Doses** | | | | |
| 250 mg BID Mean | 1 | --- | 3 to 4 | 6.34 |
| 500 mg BID Mean | 3.38 | 2.1 | 5 to 7 | 44.19 |

* Single doses (from Tables 14 & 15)

** Multiple doses (from Table 14)

Legend: BID = twice daily

Absorption:

The absolute bioavailability of 250 mg and 500 mg clarithromycin tablets is approximately 50%. Food slightly delays the onset of clarithromycin absorption but does not affect the extent of bioavailability. Therefore, CLARITHROMYCIN tablets may be given without regard to meals.

In fasting healthy human subjects, peak serum concentrations are attained within 2 hours after oral dosing. Steady-state peak serum clarithromycin concentrations, which are attained within 2 to 3 days, are approximately 1 mg/L with a 250 mg dose twice daily and 2 to 3 mg/L with a 500 mg dose twice daily. The elimination half-life of clarithromycin is about 3 to 4 hours with

250 mg twice daily dosing but increases to about 5 to 7 hours with 500 mg administered twice daily.

Clarithromycin displays non-linear pharmacokinetics at clinically relevant doses, producing greater than proportional increases in AUC with increasing dose. The degree of non-linearity is reduced on chronic clarithromycin administration (i.e. at steady state). The nonlinearity of the pharmacokinetics of the principle metabolite, 14-OH clarithromycin, is slight at the recommended doses of 250 mg and 500 mg administered twice daily. With 250 mg twice daily, 14-OH clarithromycin attains a peak steady-state concentration of about 0.6 mg/L and has an elimination half-life of 5 to 6 hours. With a 500 mg twice daily dose, the peak steady-state of 14-OH concentrations of clarithromycin are slightly higher (up to 1 mg/L) and its elimination half-life is about 7 hours. With either dose, the steady-state concentration of this metabolite is generally attained within 2 to 3 days.

Adult Patients with HIV

Steady-state concentrations of clarithromycin and 14-OH clarithromycin observed following administration of 500 mg doses of clarithromycin twice a day to adult patients with HIV-infection were similar to those observed in healthy volunteers. However, at the higher clarithromycin doses which may be required to treat mycobacterial infections, clarithromycin concentrations can be much higher than those observed at 500 mg clarithromycin doses. In adult HIV-infected patients taking 2000 mg/day in two divided doses, steady-state clarithromycin C_{max} values ranged from 5 to 10 mg/L. C_{max} values as high as 27 mg/L have been observed in HIV-infected adult patients taking 4000 mg/day in two divided doses of clarithromycin tablets.

Elimination half-lives appeared to be lengthened at these higher doses as well. The higher clarithromycin concentrations and longer elimination half-lives observed at these doses are consistent with the known nonlinearity in clarithromycin pharmacokinetics.

Clarithromycin and Omeprazole

Clarithromycin 500 mg three times daily and omeprazole 40 mg once daily were studied in fasting healthy adult subjects. When clarithromycin was given alone as 500 mg every 8 hours, the mean steady-state C_{max} value was approximately 3.8 mcg/mL and the mean C_{min} value was approximately 1.8 mcg/mL. The mean AUC₀₋₈ for clarithromycin was 22.9 mcg·hr/mL. The T_{max} and half-life were 2.1 hours and 5.3 hours, respectively, when clarithromycin was dosed at 500 mg three times daily. When clarithromycin was administered with omeprazole, increases in omeprazole half-life and AUC₀₋₂₄ were observed. For all subjects combined, the mean omeprazole AUC₀₋₂₄ was 89% greater and the harmonic mean for omeprazole t_½ was 34% greater when omeprazole was administered with clarithromycin than when omeprazole was administered alone. When clarithromycin was administered with omeprazole, the steady-state C_{max}, C_{min}, and AUC₀₋₈ of clarithromycin were increased by 10%, 27%, and 15%, respectively over values achieved when clarithromycin was administered with placebo.

Distribution:

Clarithromycin distributes readily into body tissues and fluids, and provides tissue concentrations that are higher than serum concentrations. Examples from tissue and serum concentrations are presented in Table 13.

Table 13 - Representative Clarithromycin Tissue and Serum Concentrations Following the Administration of 250 mg BID of Clarithromycin Film-coated Tablets

| Tissue Type | Concentrations | |
|-------------|----------------|--------------|
| | Tissue (mcg/g) | Serum (mg/L) |
| Tonsil | 1.6 | 0.8 |
| Lung | 8.8 | 1.7 |
| Leukocytes* | 9.2 | 1.0 |

**in vitro* data.

Legend: BID = twice daily

Metabolism:

Clarithromycin is principally excreted by the liver and kidney. The major metabolite found in urine is 14-OH-clarithromycin.

Elimination:

At 250 mg twice daily, approximately 20% of an orally administered dose of clarithromycin film-coated tablet is excreted in the urine as the unchanged parent drug. The urinary excretion of unchanged clarithromycin is somewhat greater (approximately 30%) with 500 mg twice daily dosing. The renal clearance of clarithromycin is, however, relatively independent of the dose size and approximates the normal glomerular filtration rate. The major metabolite found in urine is 14-OH-clarithromycin which accounts for an additional 10 to 15% of the dose with twice daily dosing at either 250 mg or 500 mg. Most of the remainder of the dose is eliminated in the feces, primarily *via* the bile. About 5 to 10% of the parent drug is recovered from the feces. Fecal metabolites are largely products of N-demethylation, 14-hydroxylation or both.

Special Populations and Conditions

- **Pediatrics**

See [10 CLINICAL PHARMACOLOGY, 10.3 Pharmacokinetics, Absorption.](#)

- **Geriatrics**

Dosage adjustment should be considered in elderly with severe renal impairment. In a steady-state study in which healthy elderly subjects (age 65 to 81 years old) were given 500 mg of clarithromycin every 12 hours, the maximum concentrations of clarithromycin and 14-OH clarithromycin were increased. The AUC was also increased. These changes in pharmacokinetics parallel known age-related decreases in renal function. In clinical trials, elderly patients did not have an increased incidence of adverse events when compared to younger patients.

- **Hepatic Insufficiency**
The steady-state concentrations of clarithromycin in subjects with impaired hepatic function did not differ from those in normal subjects; however, the 14-OH clarithromycin concentrations were lower in the hepatically impaired subjects. The decreased formation of 14-OH clarithromycin was at least partially offset by an increase in renal clearance of clarithromycin in subjects with impaired hepatic function when compared to healthy subjects. See [7 WARNINGS AND PRECAUTIONS, Hepatic/Biliary/Pancreatic](#) and [4 DOSAGE AND ADMINISTRATION, 4.2 Recommended Dose and Dosage Adjustment](#).
- **Renal Insufficiency**
The elimination of clarithromycin was impaired in patients with impaired renal function. The daily dose of clarithromycin should be limited to 500 mg in patients with severe renal impairment (creatinine clearance < 30 mL/min). See [7 WARNINGS AND PRECAUTIONS, Renal](#) and [4 DOSAGE AND ADMINISTRATION, 4.2 Recommended Dose and Dosage Adjustment](#).

DETAILED PHARMACOLOGY

General

Helicobacter pylori

The presence of *H. pylori* may damage the mucosal integrity and defenses so that exposure to acid/pepsin, even in normal concentrations, produces ulceration.

H. pylori displays potent urease activity which may produce an alkaline environment around the organism. Excess ammonia produced by urea hydrolysis is toxic to mucosal cells and may lead to parietal cell failure and/or to a disturbance of the normal negative feedback of acid to the antral G-cells which secrete gastrin. In addition, *H. pylori* produces catalases, lipases, phospholipases, proteases, adhesins and toxins. These enzymes may further degrade the mucous layer and damage the epithelial cell membrane. Also, the presence of *H. pylori* stimulates an active inflammatory response which contributes to mucosal damage.

Gustavson *et al.* (1995) showed that concentrations of 39.3, 23.1 and 25.2 mcg/g clarithromycin were achieved in the gastric mucosa 2, 4, and 6 hours respectively after administering 500 mg clarithromycin three times daily and that corresponding concentrations of the 14-OH metabolite were 3.2, 1.1, and 4.1 mcg/g respectively. Similar results were obtained whether or not clarithromycin was given alone or together with 40 mg omeprazole once daily (Logan *et al.*, 1995). Although the activity of the 14-OH metabolite is about half of the parent drug and its concentrations are lower, it may still contribute antibacterial activity.

Pharmacokinetics

Pharmacokinetics for clarithromycin and 14-OH-clarithromycin metabolite following the oral administration of a single dose or multiple doses of clarithromycin are outlined below

Pharmacokinetics for clarithromycin and 14-OH-clarithromycin metabolite was first studied following the oral administration of a single dose of 250 mg or 500 mg or multiple doses of clarithromycin 250 mg tablet.

Single Dose

Plasma levels were determined in 20 subjects following oral administration of a single dose of 250 mg or 500 mg of clarithromycin under fasting conditions. C_{max} occurred at 1.00 and 1.77 (mg/L) and T_{max} were 1.5 and 2.2 hours, respectively for the 250 mg and 500 mg. (Table 14, Figures 1 and 2).

Table 14 - Mean (\pm SD) Pharmacokinetic Parameters for Clarithromycin Administered as a Single Dose in the Absence of Food

| Variable | Clarithromycin Dose | |
|-----------------------------------|------------------------------|-------------------------------|
| | 250 mg | 500 mg |
| Number of male evaluable patients | 20 | 20 |
| C_{max} (mg/L) | 1.00 \pm 0.34 | 1.77 \pm 0.65 |
| $C_{max}/100 \text{ mg}^1$ | 0.40 | 0.35 |
| T_{max} (hr) | 1.5 \pm 0.8 | 2.2 \pm 0.7 |
| AUC (mg.hr/L) | 5.47 \pm 1.93 ² | 11.66 \pm 3.67 ³ |
| AUC/100 mg ¹ | 2.19 | 2.33 |

$$^1C_{max}/100 \text{ mg} = C_{max} \times \frac{100 \text{ mg}}{\text{dose}}; \text{ AUC}/100 \text{ mg} = \text{AUC} \times \frac{100 \text{ mg}}{\text{dose}}$$

$$^2\text{AUC}_{0-12 \text{ hr}}$$

$$^3\text{AUC}_{0-14 \text{ hr}}$$




Figure 1: Plasma Clarithromycin Concentration (mg/mL) vs Time Following Oral Administration of a Single Dose of Clarithromycin 250 mg.




Figure 2: Plasma Clarithromycin Concentration (mg/L) vs Time Following Oral Administration of a Single Dose of Clarithromycin 500 mg.

Multiple Dose

Representative estimated pharmacokinetic parameters for clarithromycin and 14-OH-clarithromycin metabolite after a single oral 250 mg dose and after the 5th dose of clarithromycin administered orally at 250 mg twice daily are listed in Table 15.

Table 15 - Representative Estimated Single and Multiple-Dose Pharmacokinetic Parameters for Clarithromycin and 14-OH Clarithromycin

| Variables | Single Dose (250 mg) | | Multiple Dose after 5 th Dose (250 mg BID) | |
|-------------------------------|----------------------|-------------|---|-------------|
| | Clari. | 14-OH | Clari. | 14-OH |
| C _{max} (mg/L) | 0.74 ± 0.24 | 0.61 ± 0.17 | 1.00 ± 0.29 | 0.63 ± 0.19 |
| t _½ (hr) | 2.7 | 4.2 | 3.5 | 4.7 |
| AUC ₀₋₁₂ (mg·hr/L) | 4.27 ± 1.52 | 4.91 ± 1.12 | 6.34 ± 1.82 | 4.72 ± 1.29 |

Legend: Clari. = clarithromycin; 14-OH = 14-OH-clarithromycin; BID = twice daily

The pharmacokinetics of clarithromycin and its 14-OH metabolite indicate that the steady-state concentration is achieved by the 5th dose using 250 mg of clarithromycin twice daily.

The mean plasma concentration-time along the predicted curves for clarithromycin and 14-OH-clarithromycin metabolite are shown in Figure 3.




Figure 3: Mean Plasma Concentrations of Clarithromycin and 14-OH Clarithromycin vs Time Following Seven 250 mg B.I.D. Oral Doses of Clarithromycin.

At 250 mg twice daily, approximately 20% of an orally administered dose is excreted in the urine as the unchanged parent drug. The urinary excretion of unchanged clarithromycin is

somewhat greater (approximately 30%) with 500 mg twice daily dosing. The renal clearance of clarithromycin is, however, relatively independent of the dose size and approximates the normal glomerular filtration rate. The major metabolite found in urine is 14-OH-clarithromycin which accounts for an additional 10 to 15% of the dose with twice daily dosing at either 250 mg or 500 mg.

Most of the remainder of the dose is eliminated in the feces, primarily via the bile. About 5 to 10% of the parent drug is recovered from the feces. Fecal metabolites are largely products of N-demethylation, 14-hydroxylation or both.

The steady-state concentrations of clarithromycin in subjects with impaired hepatic function did not differ from those in normal subjects; however, the 14-OH clarithromycin concentrations were lower in the hepatically impaired subjects. The decreased formation of 14-OH clarithromycin was at least partially offset by an increase in renal clearance of clarithromycin in the subjects with impaired hepatic function when compared to healthy subjects.

The pharmacokinetics of clarithromycin were also altered in subjects with impaired renal function (see [7 WARNINGS AND PRECAUTIONS, Renal](#) and [4 DOSAGE AND ADMINISTRATION, 4.2 Recommended Dose and Dosage Adjustment](#)).

Clarithromycin and Omeprazole

A pharmacokinetic study was conducted with clarithromycin 500 mg three times daily and omeprazole 40 mg once daily. When clarithromycin was given alone at 500 mg every 8 hours, the mean steady-state C_{max} value was approximately 31% higher and the mean C_{min} value was approximately 119% higher than when clarithromycin is compared with a previous study at 500 mg every 12 hours. The mean AUC_{0-24} for clarithromycin was 65% greater when 500 mg clarithromycin was given every 8 hours rather than every 12 hours. Neither T_{max} nor half-life values appeared substantially different between the every 8-hour and every 12-hour regimens.

When clarithromycin was administered with omeprazole, increases in omeprazole half-life and AUC_{0-24} were observed. For all subjects combined, the mean omeprazole AUC_{0-24} was 89% greater and the harmonic mean for omeprazole $t_{1/2}$ was 34% greater when omeprazole was administered with clarithromycin than when omeprazole was administered alone. When clarithromycin was administered with omeprazole, the steady-state C_{max} , C_{min} , and AUC_{0-8} of clarithromycin were increased by 10%, 27%, and 15%, respectively over values achieved when clarithromycin was administered with placebo.

At steady state, clarithromycin gastric mucus concentrations 6 hours post-dosing were approximately 25-fold higher in the clarithromycin/omeprazole group compared with the clarithromycin alone group. Six hours post-dosing, mean clarithromycin gastric tissue concentrations were approximately 2-fold higher when clarithromycin was given with omeprazole than when clarithromycin was given with placebo.

Clarithromycin distributes readily into body tissues and fluids, and provides tissue concentrations that are higher than serum concentrations. Examples from tissue and serum concentrations are presented in [Table 16](#).

Table 16 - Representative Clarithromycin Tissue and Serum Concentrations

| Tissue Type | Concentrations (after 250 mg BID) | |
|-------------|--------------------------------------|----------------|
| | Tissue (mcg/g) | Serum (mcg/mL) |
| Tonsil | 1.6 | 0.8 |
| Lung | 8.8 | 1.7 |
| Leukocytes* | 9.2 | 1.0 |

* *in vitro* data

Legend: BID = twice daily

11 STORAGE, STABILITY AND DISPOSAL

Store film-coated tablets at controlled room temperature between 15°C and 30°C in a tightly closed container.

12 SPECIAL HANDLING INSTRUCTIONS

Not applicable.

PART II: SCIENTIFIC INFORMATION

13 PHARMACEUTICAL INFORMATION


Drug Substance

Proper Name: Clarithromycin

Chemical Name: (3R*, 4S*, 5S*, 6R*, 7R*, 9R*, 11R*, 12R*, 13S*, 14R*)-4[(2,6-dideoxy-3-C-methyl-3-O-methyl- α -L-ribo-hexopyranosyl)oxy]-14-ethyl-12, 13-dihydroxy-7-methoxy-3,5,7,9,11, 13-hexamethyl-6-[[[3,4,6-trideoxy-3-(dime-thylamino)-beta-D-xylo-hexopyranosyl]oxy]oxacyclotetradecane-2-10-dione.

Molecular Formula and molecular mass: $C_{38}H_{69}NO_{13}$, 747.95 g/mol

Structural Formula:



Physicochemical Properties: Clarithromycin is a white to off-white crystalline powder. It is slightly soluble in methanol, ethanol and acetonitrile, and practically insoluble in water. The pKa of clarithromycin is 8.48; the pH of a 0.2% (Methanol: Water, 5:95) slurry is 8.8. The partition coefficient of clarithromycin is influenced by the pH of the water phase and polarity of the organic phase. For octanol (dipole moment = 0.25): water, the partition coefficient varies from 5.63 to 46.0 for pH water increases from 2 to 8. The melting point of clarithromycin is approximately 225°C.

14 CLINICAL TRIALS

14.1 Trial Design and Study Demographics

See [14 CLINICAL TRIALS, 14.2 Study Results](#) - for trial design and study demographics by product and indication.

14.2 Study Results

Mycobacterial Infections – Prophylaxis

Table 17 - Summary of Demographics and Trial Design Prophylaxis Against *M. avium* Complex

| Study # | Trial Design | Dosage, Route of Administration and Duration | Study Subjects Immunocompromised Patients with CD ₄ Counts <100 cells/mcL | Mean Age (Range) |
|---------|--------------|---|--|------------------|
| 561 | Double blind | clarithromycin 500 mg BID (~ 10.6 months) Placebo BID (8.2 months) | 341 341 | Adult |

Legend: BID = twice daily

More patients in the placebo arm than the clarithromycin arm discontinued prematurely from the study (75.6% and 67.4%, respectively). However, if premature discontinuations due to *Mycobacterium avium* complex (MAC) or death are excluded, approximately equal percentages of patients on each arm (54.8% on clarithromycin and 52.5% on placebo) discontinued study drug early for other reasons.

Table 18 - Summary of Efficacy Results in Immunocompromised Adult Patients Receiving Prophylaxis Against *M. avium* Complex

| | Clarithromycin | Placebo | Hazard Ratio (95% CI) | p-value | Risk Reduction |
|--|-----------------|-----------------|-----------------------|---------|----------------|
| MAC Bacteremia | | | | | |
| # patients developed MAC | 19/333 (5.7%) | 53/334 (15.9%) | 0.307 (0.177, 0.533) | <0.001* | -69.3% |
| Survival | | | | | |
| # patients died | 106/341 (31.1%) | 136/341 (39.9%) | 0.710 (0.533, 0.934) | 0.014* | 28.2% |
| Emergence of MAC Signs/Symptoms | | | | | |

| | Clarithromycin | Placebo | Hazard Ratio (95% CI) | p- value | Risk Reduction |
|---|------------------------------|----------------------------------|----------------------------------|---------------------|---------------------------|
| | # meeting criterion/total | # meeting criterion/tot al | | | |
| Wt. Loss>10% | 5/333 (2%) | 23/322 (7%) | 0.179 (0.067, 0.481) | 0.001* | 82.1% |
| Moderate/severe pyrexia | 2/332 (<1%) | 10/329 (3%) | 0.191 (0.041, 0.883) | 0.034* | 80.9% |
| Moderate/severe night sweats | 1/325 (<1%) | 7/327 (2%) | 0.130 (0.016, 1.081) | 0.059 | 87.0% |
| Mod./severe night sweats or pyrexia | 2/325 (<1%) | 13/326 (4%) | 0.140 (0.031, 0.632) | 0.011* | 86.0% |
| Moderate/severe anemia | 0/319 (0%) | | 0 | | |
| Grade 3 or 4 LFT | 3/325 (<1%) | | 0.739 (0.118, 4.649) | 0.747 | |
| Quality of Life Subscores (time to first decrease of ≥10 points) | | | | | |
| | # meeting criterion/total | # meeting criterion/tot al | | | |
| Overall health | 180/317 (57%) | 184/318 (58%) | 0.809 (0.645, 1.015) | 0.068 | |
| Physical function | 210/299 (70%) | 236/306 (77%) | 0.781 (0.637, 0.956) | 0.017* | -21.9% |
| Role function | 111/189 (59%) | 131/211 (62%) | 0.922 (0.690, 1.233) | 0.585 | |
| Social function | 187/327 (57%) | 197/331 (60%) | 0.823 (0.662, 1.024) | 0.08 | |
| Cognitive function | 174/336 (52%) | 170/339 (50%) | 0.990 (0.790, 1.240) | 0.929 | |
| Pain | 201/331 (61%) | 217/336 (65%) | 0.902 (0.731, 1.113) | 0.355 | |
| Mental health | 179/336 (53%) | 184/338 (54%) | 0.842 (0.672, 1.055) | 0.134 | |
| Energy/fatigue | 208/328 (63%) | 217/335 (65%) | 0.784 (0.636, 0.966) | 0.022* | -21.6% |
| Health distress | 170/335 (51%) | 191/335 (57%) | 0.807 (0.647, 1.007) | 0.057 | |
| Quality of life | 199/330 (60%) | 199/333 (60%) | 0.902 (0.727, 1.120) | 0.352 | |

| | Clarithromycin | Placebo | Hazard Ratio (95% CI) | p-value | Risk Reduction |
|-------------------------|----------------|---------------|-----------------------|---------|----------------|
| Hospitalization | | | | | |
| # patients hospitalized | 166/339 (49%) | 189/330 (57%) | 0.764 (0.610, 0.955) | 0.018* | -23.6% |

On an intent-to-treat basis, the 1-year cumulative incidence of MAC bacteremia was 5.0% for patients randomized to clarithromycin and 19.4% for patients randomized to placebo (Table 19). While only 19 of the 341 patients randomized to clarithromycin developed MAC, 11 of these cases were resistant to clarithromycin. The patients with resistant MAC bacteremia had a median baseline CD₄ count of 10 cells/mm³ (range 2 to 25 cells/mm³). Information regarding the clinical course and response to treatment of the patients with resistant MAC bacteremia is limited. The 8 patients who received clarithromycin and developed susceptible MAC bacteremia had a median baseline CD₄ count of 25 cells/mm³ (range 10 to 80 cells/mm³). Comparatively, 53 of the 341 placebo patients developed MAC; none of these isolates were resistant to clarithromycin. The median baseline CD₄ count was 15 cells/mm³ for placebo patients that developed MAC.




Figure 4: Survival of All Randomized Immunocompromized Adult Patients Receiving Clarithromycin in Prophylaxis Against *M. avium* complex or Placebo

Table 19 - Cumulative Incidence of MAC Bacteremia and Mortality in Immunocompromised Adult Patients Receiving Prophylaxis Against *M. avium* Complex

| | Cumulative Incidence of MAC Bacteremia‡ | | Cumulative Mortality | |
|----------|---|---------|----------------------|---------|
| | Clarithromycin | Placebo | Clarithromycin | Placebo |
| 6 month | 1.0% | 9.5% | 6.4% | 9.3% |
| 12 month | 5.0% | 19.4% | 20.8% | 29.7% |
| 18 month | 10.1% | 26.8% | 36.8% | 46.8% |

‡ from Kaplan-Meier estimates

Since the analysis at 18 months includes patients no longer receiving prophylaxis the survival benefit of clarithromycin may be underestimated.

Treatment of Mycobacterial Infections

Three studies summarized in Table 20 were designed to evaluate the following end points:

- Change in MAC bacteremia or blood cultures negative for *M. avium*
- Change in clinical signs and symptoms of MAC infection including one or more of the following: fever, night sweats, weight loss, diarrhea, splenomegaly, and hepatomegaly.

Table 20 - Summary of Demographics and Trial Design Efficacy of Clarithromycin in the Treatment of Mycobacterial Infections

| Study # | Trial design | Dosage, Route of Administration and Duration | Study Subjects (N=number) | Mean Age (Range) |
|---------|-------------------------|--|--|------------------|
| 500 | Randomized double-blind | 500 mg BID 1000 mg BID 2000 mg BID | CDC-defined AIDS and CD ₄ counts <100 cells/mCL (n=154) | Adult |
| 577 | Open-label* | 500 mg BID 1000 mg BID | CDC-defined AIDS and CD ₄ counts <100 cells/mCL (n=469) | Adult |

* Compassionate use.

Legend: BID = twice daily

The results of the study 500 are described below. The study 577 results were similar to the results of the study 500.

MAC Bacteremia:

Decreases in MAC bacteremia or negative blood cultures were seen in the majority of patients in all dose groups. Mean reductions in colony forming units (CFU) are shown below. Included

in the table are results from a separate study with a 4-drug regimen (ciprofloxacin, ethambutol, rifampicin, and clofazimine). Since patient populations and study procedures may vary between these 2 studies, comparisons between the clarithromycin results and the combination therapy results should be interpreted cautiously (Table 21).

Table 21 - Mean Reductions in Log CFU from Baseline (After 4 Weeks of Therapy)

| 500 mg BID | 1000 mg BID | 2000 mg BID | Four Drug Regimen |
|-------------------|--------------------|--------------------|--------------------------|
| (N=35) | (N=32) | (N=26) | (N=24) |
| 1.5 | 2.3 | 2.3 | 1.4 |

Legend: BID = twice daily

Although the 1000 mg and 2000 mg twice daily doses showed significantly better control of bacteremia during the first 4 weeks during therapy, no significant differences were seen beyond that point. The percent of patients whose blood was sterilized as shown by 1 or more negative cultures at any time during acute therapy was 61% (30/49) for the 500 mg twice daily group and 59% (29/49) and 52% (25/28) for the 1000 and 2000 mg twice daily groups, respectively. The percent of patients who had 2 or more negative cultures during acute therapy that were sustained through study Day 84 was 25% (12/49) in both the 500 and 1000 mg twice daily groups and 8% (4/48) for the 2000 mg twice daily group. By Day 84, 23% (11/49), 37% (18/49), and 56% (27/48) of patients had died or discontinued from the study, and 14% (7/49), 12% (6/49), and 13% (6/48) of patients had relapsed in the 500, 1000, and 2000 mg twice daily dose groups, respectively. All of the isolates had a minimum inhibitory concentration (MIC) < 8 mcg/mL at pretreatment. Relapse was almost always accompanied by an increase in MIC. The median time to first negative culture was 54, 41, and 29 days for the 500, 1000, and 2000 mg twice daily groups, respectively.

Clinically Significant Disseminated MAC Disease

Among patients experiencing night sweats prior to therapy, 84% showed resolution or improvement at some point during the 12 weeks of clarithromycin at 500 to 2000 mg twice daily doses. Similarly, 77% of patients reported resolution or improvement in fevers at some point. Response rates for clinical signs of MAC are given in Table 22.

Table 22 - Response Rates for Clinical Signs of MAC

| Resolution of Fever | | | Resolution of Night Sweats | | |
|---------------------|-----------------|---------------------------|----------------------------|-------------------|-----------------------------|
| BID dose (mg) | % ever afebrile | % afebrile \geq 6 weeks | BID dose (mg) | % ever resolving | % resolving \geq 6 weeks |
| 500 | 67 | 23 | 500 | 85 | 42 |
| 1000 | 67 | 12 | 1000 | 70 | 33 |
| 2000 | 62 | 22 | 2000 | 72 | 36 |
| Weight Gain > 3% | | | Hemoglobin Increase > 1 g | | |
| BID dose (mg) | % ever gaining | % gaining \geq 6 weeks | BID dose (mg) | % ever increasing | % increasing \geq 6 weeks |
| 500 | 33 | 14 | 500 | 58 | 26 |
| 1000 | 26 | 17 | 1000 | 37 | 6 |
| 2000 | 26 | 12 | 2000 | 62 | 18 |

Legend: BID = twice daily

The median duration of response, defined as improvement of resolution of clinical signs and symptoms, was 2 to 6 weeks.

Since the study was not designed to determine the benefit of monotherapy beyond 12 weeks, the duration of response may be underestimated for the 25 to 33% of patients who continued to show clinical response after 12 weeks.

Survival

Median survival time from study entry (Study 500) was 249 days at the 500 mg twice daily dose compared to 215 days with the 1000 mg twice daily dose. However, during the first 12 weeks of therapy, there were 2 deaths in 53 patients in the 500 mg twice daily group *versus* 13 deaths in 51 patients in the 1000 mg twice daily group. The reason for this apparent mortality difference is not known. Survival in the 2 groups was similar beyond 12 weeks. The median survival times for these dosages were similar to recent historical controls with MAC when treated with combination therapies.

Median survival time from study entry in Study 577 was 199 days for the 500 mg twice daily dose and 179 days for the 1000 mg twice daily dose. During the first 4 weeks of therapy, while patients were maintained on their originally assigned dose, there were 11 deaths in 255 patients taking 500 mg twice daily and 18 deaths in 214 patients taking 1000 mg twice daily.

Eradication of *Helicobacter pylori*

Triple Therapy: clarithromycin/omeprazole/amoxicillin

In a well-controlled double-blind study, *Helicobacter pylori* (*H. pylori*) infected duodenal ulcer patients received triple therapy with clarithromycin 500 mg twice daily, omeprazole 20 mg daily and amoxicillin 1000 mg twice daily for 10 days or dual therapy with clarithromycin 500 mg three times daily and omeprazole 40 mg daily for 14 days. *H. pylori* was eradicated in 90% of the patients receiving clarithromycin triple therapy and in 60% of the patients receiving dual therapy.

A summary of the Trial Design is presented in Table 23.

Table 23 - Summary of the Trial Design Efficacy of Clarithromycin in the Eradication of *Helicobacter pylori* – Triple Therapy

| Study # | Trial Design | Dosage, Route of Administration and Duration | Study Subjects (N=number) | Mean Age (Range) |
|---------|--|--|---------------------------|------------------|
| 183 | Phase III, randomized, double blind, multicenter | Treatment 1 Clarithromycin 500 mg BID with Omeprazole 20 mg QD and Amoxicillin 1000 mg BID Treatment 2 Clarithromycin 500 mg BID with Omeprazole 40 mg QD Oral Treatment 1: 10 days Treatment 2: 14 days | 267 patients | 18 to 75 years |

Legend: BID = twice daily; QD= once daily

The ulcer healing rates and corresponding 95% confidence intervals are presented in Table 24.

Table 24 - Ulcer Healing [95% C.I.] at 4- to 6-Week Follow-up

| Patient Subset | Clarithromycin + Omeprazole + Amoxicillin | Clarithromycin + Omeprazole | p-value |
|----------------------|---|-------------------------------|---------|
| Clinically evaluable | 93% (118/127) [87.0, 96.7] | 91% (104/114) [84.5, 95.7] | 0.641 |
| Intent-to-treat #1 | 93% (122/131) [87.4, 96.8] | 92% (111/121) [85.3, 96.0] | 0.812 |
| Intent-to-treat #2 | 90% (122/136) [83.3, 94.3] | 85% (111/130) [78.1, 91.0] | 0.353 |

- An ulcer was defined as a circumscribed break in the duodenal mucosa that measured 5 to 25 mm in the longest diameter with apparent depth and was covered with an exudate.

- Duodenal ulcer was identified by endoscopy and *H. pylori* infection at baseline was defined as at least two of three positive tests from ¹³C UBT, CLOtest[®], histology and culture.
- *H. pylori* eradication at 4 to 6 weeks posttreatment was defined as at least two of three negative tests from ¹³C UBT gastric biopsy for culture, histology and CLOtest[®].

Intent-to-treat #1: excluded patients with no confirmed evidence of *H. pylori* pretreatment, patients who had no duodenal ulcer pretreatment, and patients who did not return for a particular visit or did not have a particular procedure performed (e.g. endoscopy).

Intent-to-treat #2: excluded patients with no confirmed evidence of *H. pylori* pretreatment and patients with no duodenal ulcer pretreatment, but included as failures patients who did not return for a particular visit or did not have a particular procedure performed (e.g. endoscopy).

The *H. pylori* eradication rates and corresponding 95% confidence intervals are summarized in Table 25.

For all patient subsets, triple therapy with clarithromycin, omeprazole, and amoxicillin achieved a statistically higher eradication rate than dual therapy ($p < 0.001$). These differences were also observed when the eradication rates were adjusted for potentially influential factors such as ulcer characteristics, age, and smoking. In addition, the eradication rates within each treatment group were similar for smokers and non-smokers.

Table 25 - Global Eradication [95% C.I.] at 4- to 6-Week Follow-up

| | Clarithromycin + Omeprazole + Amoxicillin | Clarithromycin + Omeprazole | p-value |
|----------------------------------|--|--|----------------|
| Bacteriologically evaluatable | 91% (115/127) [84.1, 95.0] | 59% (68/115) [49.6, 68.2] | <0.001 |
| Intent-to-treat #1 | 90% (120/133) [83.9, 94.7] | 60% (72/120) [50.7, 68.8] | <0.001 |
| Intent-to-treat #2 | 88% (120/136) [81.6, 93.1] | 55% (72/130) [46.4, 64.1] | <0.001 |

- An ulcer was defined as a circumscribed break in the duodenal mucosa that measured 5 to 25 mm in the longest diameter with apparent depth and was covered with an exudate.
- Duodenal ulcer was identified by endoscopy and *H. pylori* infection at baseline was defined as at least two of three positive tests from ¹³C UBT, CLOtest[®], histology and culture.
- *H. pylori* eradication at 4 to 6 weeks posttreatment was defined as at least two of three negative tests from ¹³C UBT gastric biopsy for culture, histology and CLOtest[®].

Intent-to-treat #1: excluded patients with no confirmed evidence of *H. pylori* pretreatment, patients who had no duodenal ulcer pretreatment, and patients who did not

return for a particular visit or did not have a particular procedure performed (e.g. endoscopy).

Intent-to-treat #2: excluded patients with no confirmed evidence of *H. pylori* pretreatment and patients with no duodenal ulcer pretreatment, but included as failures patients who did not return for a particular visit or did not have a particular procedure performed (e.g. endoscopy).

International, Randomized, Double-blind, Placebo-controlled Study

In an international, randomized, double-blind, placebo-controlled study involving more than 100 patients in each of 6 treatment groups, patients with proven duodenal ulcer disease were randomized to treatment twice daily for 1 week with omeprazole, 20 mg (O), plus either placebo (P) or combinations of 2 of the following antimicrobials: amoxicillin, 1 g (A), clarithromycin, 250 mg or 500 mg (C250, C500), or metronidazole, 400 mg (M). *H. pylori* eradication rates for the “all-patients-treated” analysis were 96% (OAC500), 95% (OMC250), 90% (OMC500), 84% (OAC250), 79% (OAM), and 1% (OP).

Independent, Open, and Non-randomized Study

In an independent, open, and non-randomized study, *H. pylori* infected patients received eradication therapy with clarithromycin 500 mg twice daily in conjunction with amoxicillin 1000 mg twice daily and omeprazole 20 mg once daily (Group A) or omeprazole 20 mg twice daily (Group B) for 7 days. In those patients not previously treated with anti-*H. pylori* therapy, *H. pylori* was eradicated in 86% (95% CI=69-95) of patients in Group A and 75% (95% CI=62-85) of patients in Group B, the difference was not statistically significant.

14.3 Comparative Bioavailability Studies

Fasting Study

A randomized, two-way, single dose crossover comparative bioavailability study was performed to assess the bioequivalence of CLARITHROMYCIN tablets, 500 mg (Sivem Pharmaceuticals ULC) and BIAXIN BID® tablets, 500 mg (Abbott Laboratories Inc.) in healthy subjects under fasting conditions. The summary of the comparative bioavailability data from the 18 subjects that were included in the statistical analysis are presented in the following tables:

SUMMARY TABLE OF THE COMPARATIVE BIOAVAILABILITY DATA

| Clarithromycin (1 x 500 mg) Geometric Mean Arithmetic Mean (CV %) | | | | |
|--|------------------------------|------------------------------|----------------------------|-------------------------|
| Parameter | Test ¹ | Reference ² | % Ratio of Geometric Means | 90% Confidence Interval |
| AUC _T (ng·h/mL) | 16506.84 17700.08 (39.48) | 16874.19 18169.15 (42.08) | 97.8 | 89.3-107.2 |
| AUC _I (ng·h/mL) | 17465.88 18670.50 (38.23) | 17663.38 18939.63 (41.08) | 98.9 | 90.9-107.6 |
| C _{max} (ng/mL) | 2011.07 2138.23 (36.61) | 2189.42 2316.08 (32.56) | 91.8 | 80.1-105.4 |
| T _{max} ³ (h) | 2.11 (0.667-6.00) | 2.25 (0.667-6.00) | | |
| T _{1/2} ⁴ (h) | 4.86 (28.84) | 4.85 (31.86) | | |

¹ CLARITHROMYCIN (clarithromycin) tablets, 500 mg (Sivem Pharmaceuticals ULC)

² BIAXIN BID® (clarithromycin) tablets, 500 mg (Abbott Laboratories Inc., purchased in Canada)

³ Expressed as the median (range) only.

⁴ Expressed as arithmetic mean (CV%) only.

Fed Study

A randomized, two-way, single dose crossover comparative bioavailability study was performed to assess the bioequivalence of CLARITHROMYCIN tablets, 500 mg (Sivem Pharmaceuticals ULC) and BIAXIN BID® tablets, 500 mg (Abbott Laboratories Inc.) in healthy subjects under high fat, high calorie fed conditions. The summary of the comparative bioavailability data from the 18 subjects that were included in the statistical analysis are presented in the following tables:

SUMMARY TABLE OF THE COMPARATIVE BIOAVAILABILITY DATA

| Clarithromycin (1 x 500 mg) Geometric Mean Arithmetic Mean (CV %) | | | | |
|--|------------------------------|------------------------------|----------------------------|-------------------------|
| Parameter | Test ¹ | Reference ² | % Ratio of Geometric Means | 90% Confidence Interval |
| AUC _T (ng·h/mL) | 15203.85 16468.68 (47.15) | 16096.90 17201.76 (46.88) | 94.4 | 83.8-106.5 |
| AUC _I (ng·h/mL) | 15771.06 17040.85 (46.53) | 16689.50 17798.87 (46.64) | 94.5 | 84.1-106.1 |
| C _{max} (ng/mL) | 2969.57 3143.25 (36.46) | 3194.37 3368.89 (36.46) | 93.0 | 78.2-110.6 |
| T _{max} ³ (h) | 1.82 (0.667-4.00) | 1.89 (1.00-4.50) | | |
| T _{1/2} ⁴ (h) | 3.97 (27.84) | 3.98 (31.70) | | |

¹ CLARITHROMYCIN (clarithromycin) tablets, 500 mg (Sivem Pharmaceuticals ULC)

² BIAXIN BID® (clarithromycin) tablets, 500 mg (Abbott Laboratories Inc., purchased in Canada)

³ Expressed as the median (range) only.

⁴ Expressed as arithmetic mean (CV%) only.

15 MICROBIOLOGY

Clarithromycin exerts its antimicrobial action by binding to the 50S ribosomal subunit of susceptible microorganisms resulting in inhibition of protein synthesis.

Clarithromycin is active *in vitro* against various aerobic and anaerobic gram-positive and gram-negative organisms as well as most MAC microorganisms. The *in vitro* activity of clarithromycin is presented in Table 28.

Additionally, the 14-OH clarithromycin metabolite also has significant antimicrobial activity which may be additive to the activity of the parent compound. Against *Hæmophilus influenzae*, 14-OH clarithromycin is twice as active as the parent compound *in vitro*. However, for MAC isolates, the 14-OH metabolite was 4 to 7 times less active than clarithromycin. The clinical significance of this activity against MAC is unknown.

Clarithromycin is bactericidal to *H. pylori*; this activity is greater at neutral pH than at acid pH.

The ranges of MICs of clarithromycin, 14-OH-clarithromycin metabolite and the MICs required to inhibit 50% (MIC₅₀) and 90% (MIC₉₀) of bacteria are presented in Tables 29 and 30. Beta-lactamase production should not have any effect on clarithromycin activity.

Cross-resistance to azithromycin has been documented. Attention should be paid to the possibility of cross-resistance between clarithromycin and other macrolide drugs, as well as lincomycin and clindamycin.

The *in vitro* data indicate enterobacteriaceæ, pseudomonas species and other non-lactose fermenting gram-negative bacilli are not sensitive to clarithromycin.

Clarithromycin has been shown to be active against most strains of the following microorganisms both *in vitro* and in clinical infections (see [1 INDICATIONS](#)).

| Aerobic Gram-Positive microorganisms | Aerobic Gram-negative microorganisms | Other microorganisms | Mycobacteria |
|---|---|------------------------------------|---|
| <i>Staphylococcus aureus</i> | <i>Haemophilus influenzae</i> | <i>Mycoplasma pneumoniae</i> | <i>Mycobacterium avium</i> complex (MAC) consisting of: |
| <i>Streptococcus pneumoniae</i> | <i>Haemophilus parainfluenzae</i> | <i>Chlamydia pneumoniae</i> (TWAR) | <i>Mycobacterium avium</i> |
| <i>Streptococcus pyogenes</i> | <i>Moraxella catarrhalis</i> | | <i>Mycobacterium Intracellulare</i> |

The following *in vitro* data are available, but their clinical significance is unknown. Clarithromycin exhibits *in vitro* activity against most strains of the following microorganisms; however, the safety and effectiveness of clarithromycin in treating clinical infections due to these microorganisms have not been established in adequate and well-controlled clinical trials (See [15 MICROBIOLOGY, Tables 28-30 below](#)):

| Aerobic Gram-positive microorganisms | Aerobic Gram-negative microorganisms | Anaerobic Gram-positive microorganisms | Anaerobic Gram-negative microorganisms | <i>Campylobacter</i> |
|---|---|---|---|-----------------------------|
| <i>Streptococcus agalactiae</i> | <i>Bordetella pertussis</i> | <i>Clostridium perfringens</i> | <i>Bacteroides melaninogenicus</i> | <i>Campylobacter jejuni</i> |
| <i>Viridans group streptococci</i> | <i>Pasteurella multocida</i> | <i>Propionibacterium acnes</i> | | |

Table 28 - In Vitro Susceptibility of Strains of Gram-Positive and Gram-Negative Bacteria to Clarithromycin

| Microorganisms | Number of Strains | Cumulative % of Strains Inhibited at MIC (mg/L) | | | | | | | | | | | |
|--|-------------------|---|------|------|------|------|------|------|------|------|------|------|------|
| | | .031 | .062 | .125 | .250 | .500 | 1.00 | 2.00 | 4.00 | 8.00 | 16.0 | 32.0 | 64.0 |
| Gram-Positive | | | | | | | | | | | | | |
| <i>Staphylococcus aureus</i> methicillin-resistant | 25 | - | 4 | 4 | 8 | 8 | 12 | 12 | 12 | 12 | 12 | 12 | 100 |
| <i>Staphylococcus aureus</i> methicillin-susceptible | 126 | - | 20 | 75 | 84 | 86 | 87 | 87 | 87 | 88 | 88 | 88 | 100 |
| All <i>Staphylococcus aureus</i> | 151 | - | 17 | 63 | 72 | 73 | 74 | 74 | 74 | 75 | 75 | 75 | 100 |
| <i>Staphylococcus epidermidis</i> | 59 | - | 18 | 37 | 42 | 44 | 45 | 47 | 50 | 50 | 54 | 54 | 100 |
| Other coagulase negative <i>staphylococcus</i> | 27 | - | 14 | 44 | 44 | 48 | 48 | 48 | 55 | 55 | 59 | 59 | 100 |
| <i>Streptococcus pyogenes</i> (GrA) | 48 | 89 | 91 | 93 | 97 | 97 | 97 | 100 | - | - | - | - | - |
| <i>Enterococcus</i> | 97 | 1 | 4 | 8 | 25 | 59 | 61 | 63 | 63 | 64 | 64 | 68 | 100 |
| <i>Streptococcus pneumoniae</i> | 26 | 38 | 84 | 84 | 84 | 100 | - | - | - | - | - | - | - |
| <i>Streptococcus agalactiae</i> (GrB) | 41 | 95 | 95 | 95 | 95 | 95 | 97 | 100 | - | - | - | - | - |
| <i>Streptococcus viridans</i> | 15 | 86 | 86 | 86 | 93 | 93 | 93 | 93 | 93 | 93 | 93 | 93 | 100 |
| Other β -hemolytic <i>Streptococcus</i> | 19 | 78 | 78 | 78 | 84 | 84 | 84 | 89 | 89 | 94 | 94 | 94 | 100 |
| <i>Corynebacterium</i> species | 11 | 27 | 45 | 54 | 63 | 63 | 63 | 81 | 81 | 90 | 100 | - | - |
| <i>Listeria monocytogenes</i> | 7 | 28 | 100 | - | - | - | - | - | - | - | - | - | - |
| Gram-Negative | | | | | | | | | | | | | |
| <i>Neisseria gonorrhoeae</i> | 39 | 23 | 35 | 64 | 100 | - | - | - | - | - | - | - | - |
| <i>Hæmophilus influenzae</i> | 56 | 3 | 3 | 3 | 7 | 16 | 37 | 80 | 100 | - | - | - | - |
| <i>Neisseria meningitides</i> | 6 | - | 33 | 50 | 83 | 100 | - | - | - | - | - | - | - |
| <i>Campylobacter</i> species | 30 | - | 10 | 10 | 43 | 80 | 93 | 100 | - | - | - | - | - |

* MICs do not take into account the antimicrobial activity of the 14-OH clarithromycin metabolite.

Table 29 - *In vitro* Susceptibility of Different Bacteria to Clarithromycin

| Microorganisms | Number of Strains | MIC (mg/L) | | |
|--|-------------------|--------------|--------|--------|
| | | Range | 50% | 90% |
| <i>Mycoplasma pneumoniae</i> | 30 | ≤0.004-0.125 | ≤0.004 | ≤0.031 |
| <i>Bordetella pertussis</i> | 18 | ≤0.008-0.06 | ≤0.008 | 0.03 |
| <i>Legionella pneumophila</i> | 14 | 0.12-0.25 | 0.12 | 0.25 |
| <i>Hæmophilus influenzae</i> | 22 | 2-8 | 4 | 8 |
| <i>Moraxella catarrhalis</i> | 17 | 0.03-0.25 | 0.06 | 0.25 |
| <i>Chlamydia trachomatis</i> | 11 | 0.002-0.008 | 0.004 | 0.008 |
| <i>Neisseria gonorrhoeae</i> | 26 | 0.0625-4 | 0.125 | 0.5 |
| <i>Mycobacterium avium</i> | 30 | 4-32 | 8 | 16 |
| <i>Mycobacterium avium-intracellulare</i> | 124 | <0.25-4 | 1 | 2 |
| <i>Mycobacterium chelonae</i> | 137 | - | - | 0.25 |
| <i>Mycobacterium fortuitum</i> | 86 | - | 2.0 | >8.0 |
| <i>Mycobacterium kansasii</i> | 24 | ≤0.125-0.25 | ≤0.125 | 0.25 |
| <i>Pasteurella multocida</i> | 10 | 1.0-4 | 1.0 | 2.0 |
| <i>Bacteriodes melaninogenicus</i> | 12 | ≤0.125-0.2 | ≤0.125 | ≤0.125 |
| <i>Clostridium perfringens</i> | 10 | 0.25-0.5 | 0.5 | 0.5 |
| <i>Staphylococcus aureus</i> (methicillin sensitive) | 20 | 0.06-0.25 | 0.17 | 0.24 |
| <i>Streptococcus pyogenes</i> | 10 | ≤0.06 | ≤0.06 | ≤0.06 |
| <i>Chlamydia pneumoniae</i> | 49 | 0.004-0.025 | 0.016 | 0.031 |
| <i>Helicobacter pylori</i> ‡ | 13 | 0.03-0.06 | 0.03 | 0.03 |

‡ Hardy DJ, Hanson CW, Hensey DM, Beyer, JM, Fernandes PB, Susceptibility of *Campylobacter pylori* to macrolides and fluoroquinolones. J. Antimicrob Chemother 1988;22:631-636.

Table 30 - *In vitro* Susceptibility of Different Bacteria to 14-OH-Clarithromycin

| Microorganisms | Number of Strains | MIC (mg/L) | | |
|---------------------------------|-------------------|--------------|-------|-------|
| | | Range | 50% | 90% |
| <i>Streptococcus pyogenes</i> | 15 | 0.015-0.03 | 0.015 | 0.03 |
| <i>Streptococcus pneumoniae</i> | 13 | ≤0.004-0.015 | 0.008 | 0.015 |
| <i>Streptococcus agalactiae</i> | 15 | 0.03-0.06 | 0.06 | 0.06 |
| <i>Listeria monocytogenes</i> | 14 | 0.25-0.5 | 0.5 | 0.5 |
| <i>Moraxella catarrhalis</i> | 17 | 0.03-0.12 | 0.06 | 0.12 |
| <i>Neisseria gonorrhoeae</i> | 15 | 0.06-1 | 0.25 | 0.5 |
| <i>Campylobacter jejuni</i> | 12 | 0.25-2 | 0.5 | 2 |
| <i>Legionella pneumophila</i> | 14 | 0.12-0.5 | 0.25 | 0.5 |
| <i>Hæmophilus influenzae</i> | 22 | 1-4 | 2 | 4 |

| Microorganisms | Number of Strains | MIC (mg/L) | | |
|--------------------------------|-------------------|-------------|-------|------|
| | | Range | 50% | 90% |
| <i>Bordetella pertussis</i> | 18 | ≤0.008-0.06 | 0.015 | 0.06 |
| <i>Bacteroides fragilis</i> | 10 | 0.5->128 | 1 | 1 |
| <i>Clostridium perfringens</i> | 10 | 0.5-0.5 | 0.5 | 0.5 |
| <i>Propionibacterium acnes</i> | 12 | 0.03->128 | 0.03 | 0.06 |

Clarithromycin Kill Kinetics Against *Helicobacter pylori*

Figure 5 illustrates the kill kinetics of clarithromycin and 14-OH clarithromycin against *H. pylori* at 8 x MIC and at pH 8.0; and Figure 6 illustrates the kill kinetics of clarithromycin and amoxicillin against *H. pylori* at pH 6.5.




Figure 5: Kill kinetics of clarithromycin and 14-OH clarithromycin against *H. pylori* strain 2597 at 8 x MIC and at pH 8.0. A flask was inoculated to produce a starting inoculum of approximately 10⁶ cfu/mL. The flask was then incubated in an anaerobe jar with CampyPak[®] and shaken gently at 37°C. Counts were done at 0, 2, 4, 8, 24, and 48 h in physiological saline after 72 h incubation. ●, No antimicrobial; ○, clarithromycin (0.12 mg/L); x, 14-OH clarithromycin (0.24 mg/L).




Figure 6: Kill kinetics of clarithromycin and amoxicillin against *H. pylori* strain 2597 at pH 6.5. Counts were done at 0, 2, 4, 8, 24, and 48 h in physiological saline after 72 h incubation. ●, No antimicrobial; ○, clarithromycin (3 mg/L); amoxicillin ■, (3 mg/L).

Susceptibility Testing excluding *Mycobacteria* and *Helicobacter*

Dilution Techniques

Quantitative methods are used to determine antimicrobial minimal inhibitory concentrations (MICs). These MICs provide estimates of the susceptibility of bacteria to antimicrobial compounds. The MICs should be determined using a standardized procedure. Standardized procedures are based on a dilution method⁴³ (broth or agar) or equivalent with standardized inoculum concentrations and standardized concentrations of clarithromycin powder.

The standard single disc susceptibility test (using the 15 mcg clarithromycin disc) and the dilution susceptibility test should be interpreted according to the criteria in Table 31.

Table 31 - Criteria for the Interpretation of Standard Single Disc and Dilution Susceptibility Tests except for *H. influenzae* and *H. pylori*

| | Zone Diameter (mm) | Appropriate MIC Correlate (mg/L) |
|---------------|--------------------|----------------------------------|
| Susceptible | ≥ 18 | ≤ 2 |
| Intermediate* | 14 to 17 | 4 |
| Resistant | ≤ 13 | ≥ 8 |

* Indicates that the test results are equivocal; therefore, dilution tests may be indicated. N.B. These criteria and the definition are in agreement with NCCLS. Documents M2-A6⁴² and M100-S8⁴³.

The standard single disc susceptibility test (using the 15 mcg clarithromycin disc) for *H. influenzae* should be interpreted according to the criteria in Table 32.

Table 32 - Criteria for the Interpretation of Standard Single Disc and Dilution Susceptibility Tests for *H. influenzae*

| | Zone Diameter (mm) | Appropriate MIC Correlate (mg/L) |
|---------------|--------------------|----------------------------------|
| Susceptible | ≥ 13 | ≤ 8 |
| Intermediate* | 11 to 12 | 16 |
| Resistant | ≤ 10 | ≥ 32 |

* Indicates that the test results are equivocal; therefore, dilution tests may be indicated.

N.B. According to the revised NCCLS 1997 and 1998 Guidelines, the zone diameter and MIC values reflect both the activities of the parent compound and 14-OH metabolite.

A report of “Susceptible” indicates that the pathogen is likely to respond to monotherapy with clarithromycin.

A report of “Intermediate” indicates that the result be considered equivocal, and if the microorganism is not fully susceptible to alternative clinically feasible drugs, the test should be repeated. This category implies possible clinical applicability in body sites where clarithromycin is physiologically concentrated or in situations where high clarithromycin dosages can be used. This category provides a buffer zone which prevents small uncontrolled technical factors from causing major discrepancies in interpretations.

A report of “Resistant” indicates that achievable drug concentrations are unlikely to be inhibitory, and other therapy should be selected.

Diffusion Techniques

Quantitative methods that require measurement of zone diameters also provide reproducible estimates of the susceptibility of bacteria to antimicrobial compounds. One such standardized procedure⁴⁴ requires the use of standardized inoculum concentrations. This procedure uses paper disks impregnated with 15 mcg clarithromycin to test the susceptibility of microorganisms to clarithromycin.

Reports from the laboratory providing results of the standard single-disk susceptibility test with a 15 mcg clarithromycin disk should be interpreted according to the criteria in Table 31.

Standardized Dilution Techniques

Standardized susceptibility test procedures require the use of laboratory control microorganisms to control the technical aspects of the laboratory procedures. Standard clarithromycin powder should provide the following MIC values for *S. aureus* and *H. influenzae*

(Table 33).

Table 33 - Standard Clarithromycin Powder MIC Values

| Microorganisms | MIC (mcg/mL) |
|--------------------------------|--------------|
| <i>S. aureus</i> ATCC 29213 | 0.12 to 0.5 |
| <i>H influenzae</i> ATCC 49247 | 4 to 16 |

Standardized Diffusion Techniques

As with standardized dilution techniques, diffusion methods require the use of laboratory control microorganisms that are used to control the technical aspects of the laboratory procedures. For the diffusion technique, the 15 mcg clarithromycin disk should provide the following zone diameters for *S. aureus* and *H. influenzae* (Table 34).

Table 34 - Zone Diameter for the 15 mcg Clarithromycin Disc

| Microorganisms | Zone Diameter (mm) |
|--------------------------------|--------------------|
| <i>S. aureus</i> ATCC 25923 | 26 to 32 |
| <i>H influenzae</i> ATCC 49247 | 11 to 17 |

In vitro Activity of Clarithromycin against Mycobacteria

Clarithromycin has demonstrated *in vitro* activity against MAC microorganisms isolated from both AIDS and non-AIDS patients. While gene probe techniques may be used to distinguish *M. avium* species from *M. intracellulare*, many studies only reported results on MAC isolates.

Various *in vitro* methodologies employing broth or solid media at different pH's, with and without oleic acid-albumin-dextrose-catalase (OADC), have been used to determine clarithromycin MIC values for mycobacterial species. In general, MIC values decrease more than 16-fold as the pH of Middlebrook 7H12 broth increases from 5.0 to 7.4. At pH 7.4, MIC values determined with Mueller-Hinton agar were 4- to 8-fold higher than those observed with Middlebrook 7H12 media. Utilization of OADC in these assays has been shown to further alter MIC values.

Clarithromycin activity against 80 MAC isolates from AIDS patients and 211 MAC isolates from non-AIDS patients was evaluated using a microdilution method with Middlebrook 7H9 broth. Results showed MIC values of ≤ 4.0 mcg/mL in 81% and 89% of the AIDS and non-AIDS MAC isolates, respectively. Twelve percent of the non-AIDS isolates had an MIC value ≤ 0.5 mcg/mL. Clarithromycin activity was evaluated against phagocytized MAC in mouse and human macrophage cell cultures as well as in the beige mouse infection model.

Clarithromycin activity was evaluated against *Mycobacterium tuberculosis* microorganisms. In 1 study utilizing the agar dilution method with Middlebrook 7H10 media, 3 of 30 clinical isolates had an MIC of 2.5 mcg/mL. Clarithromycin inhibited all isolates at > 10.0 mcg/mL.

Susceptibility Testing for Mycobacterium avium Complex

The disk diffusion and dilution techniques for susceptibility testing against gram-positive and gram-negative bacteria should not be used for determining clarithromycin MIC values against mycobacteria. *In vitro* susceptibility testing methods and diagnostic products currently available for determining MIC values against MAC organisms have not been standardized nor validated. Clarithromycin MIC values will vary depending on the susceptibility testing method employed, composition and pH of the media, and the utilization of nutritional supplements. Breakpoints to determine whether clinical isolates of *M. avium* or *M. intracellulare* are susceptible or resistant to clarithromycin have not been established.

In vitro Activity of Clarithromycin against Helicobacter pylori

Clarithromycin has demonstrated *in vitro* activity against *H. pylori* isolated from patients with duodenal ulcers. *In vitro* susceptibility testing methods (broth microdilution, agar dilution, E-test, and disk diffusion) and diagnostic products currently available for determining MICs and zone sizes have not been standardized, validated, or approved for testing *H. pylori*. The clarithromycin MIC values and zone sizes will vary depending on the susceptibility testing methodology employed, media, growth additives, pH, inoculum concentration tested, growth phase, incubation atmosphere, and time.

Susceptibility Test for Helicobacter pylori

In vitro susceptibility testing methods and diagnostic products currently available for determining MICs and zone sizes have not been standardized, validated, or approved for testing *H. pylori* microorganisms. MIC values for *H. pylori* isolates collected during 2 U.S. clinical trials evaluating clarithromycin plus omeprazole, were determined by broth microdilution MIC methodology (Hachem CY et al., 1996). Results obtained during the clarithromycin plus omeprazole clinical trials fell into a distinct bimodal distribution of susceptible and resistant clarithromycin MICs.

If the broth microdilution MIC methodology published in Hachem CY et al., 1996 is used and the following tentative breakpoints are employed, there should be reasonable correlation between MIC results and clinical and microbiological outcomes for patients treated with clarithromycin plus omeprazole (Table 35).

Table 35 - Susceptibility Testing for *Helicobacter pylori* in Patients Treated with Clarithromycin and Omeprazole

| MIC (mcg/mL) | Interpretation |
|--------------|------------------|
| ≤ 0.06 | Susceptible (S) |
| 0.12 to 2.0 | Intermediate (I) |
| ≥ 4 | Resistant (R) |

These breakpoints should not be used to interpret results obtained using alternative methods.

16 NON-CLINICAL TOXICOLOGY

General Toxicology:

Acute Toxicity

The acute toxicity of clarithromycin administered by a variety of routes, was studied in mice and rats. The median lethal dose by the oral route ranged from 2.7 to >5.0 g/kg. Acute toxicity did not differ markedly between sexes (Table 36).

Table 36 - Acute LD₅₀ values of Clarithromycin

| Species | Sex | Route | LD ₅₀ value (g/kg) |
|---------|-----|-------|-------------------------------|
| Mice | M | PO | 2.74 |
| | F | PO | 2.7 |
| | M | SC | >5.0 |
| | F | SC | >5.0 |
| | M | IP | 1.03 |
| | F | IP | 0.85 |
| | M | IV | 0.17 |
| | F | IV | 0.2 |
| Rats | M | PO | 3.47 |
| | F | PO | 2.7 |
| | M | SC | >5.0 |
| | F | SC | >5.0 |
| | M | IP | 6.69 |
| | F | IP | 7.58 |

Legend: IP = intraperitoneal; IV = intravenous; PO = oral; SC = subcutaneous

The primary signs of toxicity included reduction in activities, behaviours, weight gains, respiration rates and sedation. The emetic activity of clarithromycin prevented the determination of the lethal dose in dog.

The acute oral toxicity of clarithromycin in very young mice and rats was determined. The median lethal dose (1.2 g/kg) was about 2-fold that seen in the older rodents.

Subchronic Toxicity

Studies were conducted in rats, dogs and monkeys with clarithromycin administered orally. The duration of administration ranged from 14 days to 42 days.

Rats

One study in rats (with oral doses up to 800 mg/kg/day) failed to show adverse effects in rats exposed to 50 mg/kg/day for 4 weeks. The clinical signs observed at toxic doses were reduced motility, piloerection, hypothermia and perineal urine staining. Changes occurred in biochemical parameters at 200 and 800 mg/kg/day indicative of hepatotoxicity which was confirmed by histopathologic findings of hepatocyte necrosis.

Other pathologic findings at the top 2 dose levels included swelling of the renal cortical tubular epithelia and atrophic changes to the lymphatic and genital systems. The same toxicity profile was observed in immature rats following the daily administration of oral doses up to 150 mg/kg/day of clarithromycin for 6 weeks. At 150 mg/kg/day, there was an increase in relative weights of liver and kidneys.

Dogs

Dogs were dosed orally with 0, 6.25, 25, 100 or 400 mg/kg/day of clarithromycin daily for 28 days. Emesis occurred sporadically in the treated dogs. No other adverse effects were seen in dogs exposed to 6.25 mg/kg/day. The clinical signs at higher dosages included loose stools, lacrimation and conjunctivitis.

Slight anorexia was noted in dogs receiving 100 mg/kg/day or more. Dogs at 400 mg/kg/day exhibited reduced red blood cell count, hematocrit, hemoglobin concentration, serum albumin, and mean urine pH and specific gravity. Increases were seen in serum transaminase, alkaline phosphatase, and total bilirubin concentrations.

Bilirubin was detected in the urine. Other pathologic changes at 400 mg/kg/day included biliary hyperplasia, gastric glandular atrophy, renal tubule epithelial atrophy, edema of the iris, ciliary body and choroid, capillary proliferation in the cornea, suppression of spermatogenesis, and adrenal medullary degeneration.

Monkeys

Monkeys were treated daily for 1 month with oral doses of 0, 25, 100 or 400 mg/kg/day. Two animals out of 10 receiving 400 mg/kg/day died. Salivation was recorded at all dosage levels. No other adverse effects were seen in animals treated daily with 25 mg/kg/day.

The clinical signs observed at higher doses and most frequently at 400 mg/kg/day were vomiting, emesis, sunken eyes, dehydration, emaciation, low rectal temperature, body weight loss, reduced food consumption, cloudiness of the cornea and reduction in intra-ocular pressure. Yellow discoloured feces were passed on a few isolated occasions by some animals given a dose of 400 mg/kg/day. As with the other species, the liver was the primary target at toxic doses as shown by early elevation of serum concentration of glucose, BUN, creatinine, ALT, AST, LDH, amylase and/or triglyceride; an electrolyte imbalance and low levels of protein, cholesterol, phospholipid; elevated leucine aminopeptidase (LAP).

Principal histopathologic changes were seen mainly in high-dose monkeys, but some mid-dose monkeys exhibited similar alterations. Changes included, necrosis and vacuolation of hepatocytes, vacuolation of renal cortical tubules, no spermatogenesis, thymic regression and single cell necrosis of the stomach. In man the recommended dose is 500 to 1000 mg/day or 7.1 to 14.3 mg/kg/day (70 kg person).

Chronic Toxicity

Rats (20/sex/group) were treated daily with oral doses of 0, 15, 37.5, 75 or 150 mg/kg/day for 3 months. There were 8 incidental deaths, but none of them were considered treatment related. Clinical signs included increased salivation, dehydration, hyperactivity and were observed in a dose-related manner. The only toxic effect noted was some variation in body weight gain. No toxicologically significant changes occurred in hematology, biochemistry or urinalysis results.

Post-mortem, there was an increase in mean relative liver and kidney weights at the top dose level. No microscopic changes were detected in the kidneys, but in the liver, there was a sex/dose-related increase in multinucleated hepatocytes. Effects were only seen in females at 150 mg/kg/day but in males occurred as low as 37.5 mg/kg/day.

A 6-month oral study was performed in rats (20 to 27/sex/group) at dosages of 0, 1 to 6, 8, 40 or 200 mg/kg/day. Seven male and female rats from the control group and the 40 and 200 mg/kg/day groups were allowed a 63 day non-dosed recovery period. No mortalities occurred. Body weight and food intake were reduced at high doses during the dosing phase but normalized during recovery.

Water intake and urine volume increased in males and females of the 40 and 200 mg/kg/day groups. Dose-related hematological changes included reduced erythrocytes and HCT with increased MCV, MCH and MCHC and relative eosinophil counts. Biochemical changes were mainly restricted to the high-dose group and included increased ALP and decreased phospholipids; decreased total cholesterol and triglycerides, and increased AST and ALT in males only and decreased albumin in females only.

Organ weight increases were found to include cecum, adrenals, liver, and spleen. Histopathological examinations showed drug-related, recovery-reversible, increases in

multinucleated hepatocytes associated with minimal and focal necrosis in livers of both sexes at the top 2 dose levels. No relevant pathology was found in the cecum, adrenals or spleen to account for the increased weights. After recovery only the 200 mg/kg/day group had increased multinucleated hepatocytes.

Dogs (7/sex/group) were administered daily with oral doses of 0, 10, 30, or 100 mg/kg/day of clarithromycin for 3 months. Emesis occurred at levels of 30 mg/kg and above. One male high-dose dog was killed *in extremis* on day 69. Drug-related lesions were seen in the liver, gallbladder, thymus and stomach.

Hematological and biochemical changes at the high-dose level included, decreased RBC and HCT, increased ALT, ALP, GGT, and decreased total protein and albumin. No significant organ weight changes were recorded, but treatment-related microscopic alterations in the liver and stomach of mild- and high-dose dogs were seen, as well as changes in gallbladder, spleen and thymus of high-dose animals.

A 6-month oral study was also performed in dogs (4 to 5/sex/group) at dosages of 0, 0.8, 4, 20 or 100 mg/kg/day. At the 0 and 100 mg/kg levels, 1 male and 1 female dog were allowed a 1-month, non-dosed, recovery period. One male high-dose dog died on day 174. This death was considered to be as a direct result of clarithromycin administration. Histopathologic examination revealed hepatic parenchymal damage, identifying the cause of clinical jaundice. Clinical signs during the dosing phase of the study were restricted to the top 2 dose levels and included emesis and ocular signs. Food consumption and water intake were reduced at 20 and 100 mg/kg/day.

Hematologic changes at 100 mg/kg were indicative of subclinical anemia. Biochemical alterations at the same level were associated with liver damage. Ocular changes were only apparent at the top dose level.

Increase in the weights of lung, liver, spleen, adrenals and kidneys were found at 100 mg/kg/day. Histopathologic examination of these organs showed degeneration of liver parenchyma, and toxic effects in adrenals. The thymus weight was reduced at 100 mg/kg/day. At the end of the recovery period all findings had regressed or reduced.

Monkeys (5 to 6/sex/group) were similarly administered clarithromycin at levels of 0, 25, 50 or 100 mg/kg/day for 6 months. At the 0 and 100 mg/kg/day levels, 1 male and 1 female monkey were allowed a one-month recovery period. One high-dose female died in week 25. Inhalation of vomit was considered to be the cause of death. Clinical signs were restricted to a dose-related incidence of emesis and salivation. No treatment-related effects were found in food consumption, ophthalmoscopy or hematology. Weight loss was restricted to 1 high-dose female. Minor serum chemistry changes were seen at the 100 mg/kg level, particularly in plasma proteins. Urinalysis revealed a dose-related lowering of pH and SG at 13 weeks only. Organ weight increases in liver, adrenal and kidneys were seen at high doses, but pathology was

restricted to minimal liver changes consisting of cytoplasmic rarefaction of centrilobular hepatocytes. All changes were reversed during the recovery period.

Carcinogenicity:

Long-term studies in animals have not been performed to evaluate the carcinogenic potential of clarithromycin.

Mutagenicity

The following *in vitro* mutagenicity tests have been conducted with clarithromycin: *Salmonella*/mammalian microsome test, bacterial-induced mutation frequency test, *in vitro* chromosome aberration test, rat hepatocyte DNA synthesis assay, mouse lymphoma assay, mouse dominant lethal study, mouse micronucleus test.

All tests had negative results except the *in vitro* chromosome aberration test which was weakly positive in one test and negative in another. In addition, a Bacterial Reverse-Mutation Test (Ames Test) has been performed on clarithromycin metabolites with negative results.

Reproduction and Developmental Teratology

Fertility and reproduction studies have shown that daily doses of 150 to 160 mg/kg/day to male and female rats caused no adverse effects on the estrous cycle, fertility, parturition, or number and viability of offspring. Plasma levels in rats after 150 mg/kg/day were 2 times the human serum levels.

In the 150 mg/kg/day monkey studies, plasma levels were 3 times the human serum levels. When given orally after 150 mg/kg/day, clarithromycin was shown to produce embryonic loss in monkeys. This effect has been attributed to marked maternal toxicity of the drug at this high dose.

In rabbits, *in utero* fetal loss occurred at an intravenous dose of 33 mg/m², which is 17 times less than the maximum proposed human oral daily dose of 618 mg/m².

Special Toxicology

Acute Renal Toxicity

There was no evidence of nephrotoxicity of clarithromycin in the rat at doses up to 500 mg/kg/day.

Hepatotoxicity

In the *in vitro* and *in vivo* hepatotoxicity studies comparing clarithromycin with erythromycin, it was found that clarithromycin caused no greater cytotoxicity than erythromycin stearate and

much less toxicity than erythromycin estolate. Hepatic enzyme induction was not found in doses below 500 mg/kg/day. In cynomolgus monkeys, the closest metabolic model for humans, elevations of ALT and LDH were identified at 200 mg/kg/day.

In dogs, a rise of ALT has been seen at 100 mg/kg/day, and in Wistar rats, a similar elevation of enzymes was seen at 200 mg/kg/day. Morphologic lesions related to prolonged exposure to clarithromycin (up to 6 months) have been consistent with reportedly reversible changes in rat, dog, and monkey studies. Such doses are many times beyond the therapeutic range in humans, which is within 8 to 10 mg/kg/day.

Ocular Toxicity:

Ocular lesions appear confined to dogs and monkeys receiving lethal doses, which were large multiples of the human therapeutic dose. Radiolabelled clarithromycin studies indicate the eye is not selectively burdened by drug deposits and that clearance from this tissue follows that seen in other tissues. Opacities occur in the cornea following widespread extraocular tissue changes which are detectable *via* numerous diagnostic methods. Reduced intraocular pressure precedes corneal opacity in a relatively predictive manner. Some evidence for transient opacity and at least partial resolution was noted in animal studies, but most animals succumbed to other organ dysfunctions shortly after opacities were observed.

Animals given doses close to the therapeutic dose had no ocular changes. No ophthalmologic effects were noted in rabbits treated at doses of 40 and 160 mg/kg/day for 28 days.

Ototoxicity

No effects on pinna reflex were seen in guinea pigs at a dose of 400 mg/kg/day but inner and outer hair cells disappeared suggesting toxic damage. No evidence of damage was reported at 200 mg/kg/day.

17 SUPPORTING PRODUCT MONOGRAPHS

Biaxin BID® (clarithromycin tablets, film-coated, 250 mg and 500 mg, USP), submission control: 288756; Product Monograph, BGP Pharma ULC, Date of Revision: December 17,, 2024.

PATIENT MEDICATION INFORMATION

READ THIS FOR SAFE AND EFFECTIVE USE OF YOUR MEDICINE

Pr **CLARITHROMYCIN**

Clarithromycin Tablets

Read this carefully before you start taking **CLARITHROMYCIN** and each time you get a refill. This leaflet is a summary and will not tell you everything about this drug. Talk to your healthcare professional about your medical condition and treatment and ask if there is any new information about **CLARITHROMYCIN**.

Serious Warnings and Precautions

- CLARITHROMYCIN should not be used in pregnancy, especially during the first 3 months. If there are no other medicines you can take for your infection, your healthcare professional may give you CLARITHROMYCIN. If this happens, they will discuss the risks to your baby with you. Talk to your healthcare professional before taking CLARITHROMYCIN if you are pregnant or think you might be pregnant.
- Taking CLARITHROMYCIN along with certain other drugs may lead to serious safety issues. Talk to your healthcare professional about all the medicines you take.

What is CLARITHROMYCIN used for?

- CLARITHROMYCIN is used to treat certain infections like pneumonia, bronchitis and infections of the sinuses, skin, and throat that are caused by bacteria.
- It is used with other medicines to kill bacteria called *Helicobacter pylori* (*H. pylori*). This may prevent duodenal ulcers from coming back. Duodenal ulcers are sores on the upper part of the small intestine.
- It is used to prevent and to treat MAC disease in patients with HIV. MAC is a short word for *Mycobacterium avium complex*, the bacteria that cause MAC disease.

Antibacterial drugs like CLARITHROMYCIN treat only bacterial infections. They do not treat viral infections such as the common cold. Although you may feel better early in treatment, CLARITHROMYCIN should be taken exactly as directed. Misuse or overuse of CLARITHROMYCIN could lead to the growth of bacteria that will not be killed by CLARITHROMYCIN (resistance). This means that CLARITHROMYCIN may not work for you in the future. Do not share your medicine.

How does CLARITHROMYCIN work?

CLARITHROMYCIN is an antibiotic that kills bacteria in your body.

What are the ingredients in CLARITHROMYCIN?

Medicinal ingredient: Clarithromycin

Non-medicinal ingredients: Colloidal silica hydrated, croscarmellose sodium, magnesium stearate, microcrystalline cellulose, opadry yellow (D&C Yellow #10 aluminum lake, hydroxypropyl methylcellulose, lactose monohydrate, polyethylene glycol 4000, titanium dioxide) and powdered cellulose.

CLARITHROMYCIN comes in the following dosage forms:

250 mg and 500 mg tablets.

Do not use CLARITHROMYCIN if:

- You are allergic to clarithromycin or any of the other ingredients in CLARITHROMYCIN.
- You are allergic to another medicine called erythromycin or any other medicines from a class of antibiotics called macrolides (such as azithromycin or telithromycin).
- You are taking any of the following medications:
 - Ergotamine, dihydroergotamine (for migraine); Lovastatin, simvastatin, lomitapide (for high cholesterol); Ticagrelor (for cardiovascular disease); Saquinavir (treatment for HIV); Oral midazolam (for trouble sleeping or agitation); Pimozide (for schizophrenia); Colchicine (for gout); Domperidone (for gastrointestinal disorders); ivabradine (for treatment of stable chronic heart failure).
 - Pimozide, ergotamine, dihydroergotamine and colchicine can interact with CLARITHROMYCIN, possibly leading to an irregular heartbeat. Deaths have occurred.
- You had liver problems after taking CLARITHROMYCIN, or any other medicine containing clarithromycin, in the past.
- You have severe liver failure in combination with kidney problems.
- You have a history of heart disturbance or irregular heartbeat such as arrhythmias, QT prolongation or torsades de pointes.
- You have low levels of potassium in the blood (hypokalemia) or low levels of magnesium in the blood (hypomagnesaemia).

To help avoid side effects and ensure proper use, talk to your healthcare professional before you take CLARITHROMYCIN. Talk about any health conditions or problems you may have, including if you:

- Have now or have had health problems in the past.
- Have or develop severe diarrhea as this may be a sign of a more serious condition.
- Have kidney problems.
- Have liver problems.

- Are taking medicines called digoxin (for heart failure); atorvastatin or pravastatin (for high cholesterol); or midazolam (a sedative).
- Are taking a medicine called quetiapine (for schizophrenia, bipolar depression). Serious and life-threatening side effects have occurred in people taking clarithromycin and quetiapine, including malignant neuroleptic syndrome (fever, rigid muscles, dizziness, fainting, and altered mental state). Your healthcare professional will decide if you should take this medication.
- Are allergic to other medicines, foods, dyes, or preservatives.
- Are pregnant, trying to get pregnant or think you might be pregnant.
- Are breastfeeding or planning to breastfeed. CLARITHROMYCIN passes into your breastmilk and can harm your baby.
- Have a condition called myasthenia gravis which is a chronic disease that causes muscle weakness. CLARITHROMYCIN may make your myasthenia gravis worse.
- Are taking CLARITHROMYCIN and oral medicines for diabetes (such as gliclazide, glyburide) and / or with insulin as this can result in serious low blood sugar levels (hypoglycemia). Discuss with your healthcare professional how you should monitor your blood sugar levels.
- Are taking warfarin, as there is a risk of serious bleeding with CLARITHROMYCIN.
- Are taking triazolam, alprazolam or other benzodiazepines (midazolam). These should be used cautiously with CLARITHROMYCIN due to the serious risk of effects on your brain and spinal cord.
- Are taking CLARITHROMYCIN and medicines used to prevent blood clots such as dabigatran, rivaroxaban, apixaban and edoxaban, particularly if your healthcare professional has told you that you are at high risk of bleeding.

Other warnings you should know about:

Serious heart problems:

Use of antibiotics like CLARITHROMYCIN have resulted in heart problems such as irregular heartbeat, torsades de pointes and QT prolongation sometimes leading to death. Talk to your healthcare professional if you are elderly or have risk factors such as:

- Heart disease, heart problems or slow heartbeat.
- If you are taking other medicines which are known to cause serious disturbances in heart rhythm.
- If you have disturbances in the levels of salts (electrolytes) in your blood, such as low levels of magnesium (hypomagnesemia).

Antibiotic resistance and HIV:

Development of antibiotic resistance (where the medicine no longer works to kill bacteria) has been seen in patients with HIV taking clarithromycin. To avoid this, you should always take your medicine as advised by your healthcare professional.

Driving and using machines:

If you feel dizzy, confused or disorientated while taking CLARITHROMYCIN, do not drive or operate machines.

Tell your healthcare professional about all the medicines you take, including any drugs, vitamins, minerals, natural supplements or alternative medicines.

The following may interact with CLARITHROMYCIN:

- Alfentanil (used during surgery).
- Alprazolam, hexobarbital, phenobarbital, midazolam, triazolam (sedative medications).
- Amlodipine, diltiazem, verapamil (calcium channel blockers often used for high blood pressure).
- Aripiprazole, pimozide, quetiapine, risperidone (for schizophrenia, bipolar depression).
- Atazanavir, indinavir, ritonavir, saquinavir, nevirapine, efavirenz, etravirine, zidovudine (treatments for HIV).
- Atorvastatin, lovastatin, pravastatin, rosuvastatin, simvastatin, lomitapide (for high cholesterol).
- Bromocriptine (used for problems with your pituitary gland and Parkinson's disease).
- Carbamazepine (for seizures, nerve pain or bipolar depression).
- Cilostazol, digoxin, quinidine, disopyramide, warfarin/acenocoumarol, ticagrelor (diseases of your blood vessels and heart).
- Colchicine (treatment for gout).
- Cyclosporine (used for psoriasis, rheumatoid arthritis and after organ transplant).
- Domperidone (used for gastrointestinal disorders).
- Ergotamine, dihydroergotamine (often used for migraine headaches).
- Fluconazole, itraconazole (for fungal infections).
- Hydroxychloroquine, chloroquine (used to treat conditions including rheumatoid arthritis, or to treat or prevent malaria)
- Insulin, nateglinide, pioglitazone, repaglinide, rosiglitazone (for diabetes).
- Ivabradine (used for treatment of stable chronic heart failure).
- Lansoprazole, omeprazole (proton pump inhibitors for heart burn and reflux).
- Corticosteroids (eg. methylprednisolone, an anti-inflammatory).
- Phenytoin, valproic acid (treatment of seizures and epilepsy).
- Rifabutin, rifampin (treatments for infections).
- Rivaroxaban, apixaban, edoxaban (to prevent blood clots).
- Sildenafil, tadalafil, vardenafil (treatments for erectile dysfunction).
- St. John's Wort (for depression).
- Tacrolimus (used after organ transplant).
- Theophylline (asthma and other lung problems).
- Tolterodine (treatment for overactive bladder).
- Vinblastine, ibrutinib (cancer treatment).

How to take CLARITHROMYCIN:

- Always take it exactly how your healthcare professional has told you.
- Your healthcare professional will tell you how much CLARITHROMYCIN to take and when to take it.
- How much you are prescribed will depend on the condition you have.
- You can take CLARITHROMYCIN with or without meals.

Usual dose:

For respiratory tract infections (like pneumonia, bronchitis and infections of the sinuses and throat) and skin infections:

The usual dose of CLARITHROMYCIN is 250 mg to 500 mg every 12 hours for 7 to 14 days.

For infections with *H. pylori* and treatment of duodenal ulcers (a sore in your intestine):

The usual dose of CLARITHROMYCIN is 500 mg every 12 hours for 10 days. You will take CLARITHROMYCIN together with omeprazole (20 mg once a day) and amoxicillin (1 g every 12 hours).

For prevention and treatment of MAC disease in patients with HIV:

The usual dose of CLARITHROMYCIN is 500 mg every 12 hours. Your healthcare professional will tell you how long you should continue taking CLARITHROMYCIN for.

Overdose:

Symptoms of CLARITHROMYCIN overdose are abdominal pain, vomiting, nausea and diarrhea

If you think you, or a person you are caring for, have taken too much CLARITHROMYCIN, contact a healthcare professional, hospital emergency department, or regional poison control centre immediately, even if there are no symptoms.

Missed Dose:

- If you miss a dose, take it as soon as you remember.
- If it is almost time for your next dose, do not take the missed dose.
- Take your next dose when you would normally take it.
- Never take a double dose to make up for a missed dose.

What are possible side effects from using CLARITHROMYCIN?

These are not all the possible side effects you may have when taking CLARITHROMYCIN. If you experience any side effects not listed here, tell your healthcare professional.

Side effects may include:

- abdominal pain
- abnormal taste
- diarrhea
- ear disorder (trouble hearing and ringing in your ears)

- flatulence
- indigestion
- headache
- nausea
- rash
- vomiting

| Serious side effects and what to do about them | | | |
|---|--------------------------------------|--------------|---|
| Symptom/effect | Talk to your healthcare professional | | Stop taking drug and get immediate medical help |
| | Only if severe | In all cases | |
| UNCOMMON | | | |
| Allergic reactions: itching, hives, rash, sore throat, fever, swelling, difficulty breathing, lightheadedness/ dizziness, swelling of your tongue or throat, warm red skin or wheezing. | | | ✓ |
| Clostridium difficile colitis (bowel inflammation): severe diarrhea (bloody or watery) with or without fever, abdominal pain, or tenderness. | | ✓ | |
| Irregular heartbeat | | | ✓ |
| Myasthenia gravis: muscle weakness, drooping eyelid, vision changes, difficulty chewing and swallowing, trouble breathing. | | | ✓ |
| Hepatitis (liver inflammation): abdominal pain, nausea, vomiting, yellowing of skin and eyes, dark urine. | | | ✓ |

If you have a troublesome symptom or side effect that is not listed here or becomes bad enough to interfere with your daily activities, tell your healthcare professional.

Reporting Side Effects

You can report any suspected side effects associated with the use of health products to Health Canada by:

- Visiting the Web page on Adverse Reaction Reporting (<https://www.canada.ca/en/health-canada/services/drugs-health-products/medeffect-canada/adverse-reaction-reporting.html>) for information on how to report online, by mail or by fax; or
- Calling toll-free at 1-866-234-2345.

NOTE: Contact your health professional if you need information about how to manage your side effects. The Canada Vigilance Program does not provide medical advice.

Storage:

CLARITHROMYCIN should be stored at controlled room temperature between 15 and 30°C in a tightly closed container. Protect from light. Do not use beyond the expiration date.

Keep CLARITHROMYCIN out of reach and sight of children.

If you want more information about CLARITHROMYCIN:

- Talk to your healthcare professional.
- Find the full Product Monograph that is prepared for healthcare professionals and includes this Patient Medication Information by visiting the Health Canada website <https://www.canada.ca/en/health-canada/services/drugs-health-products/drug-products/drug-product-database.html> or the manufacturer's website www.sivem.ca, or by calling 1-855-788-3153.

This leaflet was prepared by Sivem Pharmaceuticals ULC.

Last Revised: JUL 10, 2025