

PRODUCT MONOGRAPH

PrMiconazole in 0.9% Sodium Chloride Injection

Sterile Solution,
1 mg / mL (50 mg / 50 mL, 100 mg / 100 mL and 150 mg / 150 mL) miconazole
(as miconazole sodium),
Intravenous

Manufacturer's Standard

Antifungal Agent

Baxter Corporation
7125 Mississauga Road,
Mississauga, ON
L5N 0C2

Date of Initial Authorization:
SEP 17, 2025

Submission Control Number: 268188

TABLE OF CONTENTS

PART I: HEALTH PROFESSIONAL INFORMATION.....	3
SUMMARY PRODUCT INFORMATION	3
INDICATIONS AND CLINICAL USE.....	3
CONTRAINDICATIONS.....	4
WARNINGS AND PRECAUTIONS	4
ADVERSE REACTIONS	6
DRUG INTERACTIONS.....	12
DOSAGE AND ADMINISTRATION.....	13
OVERDOSAGE.....	15
ACTION AND CLINICAL PHARMACOLOGY	15
STORAGE AND STABILITY	18
DOSAGE FORMS, COMPOSITION AND PACKAGING	19
PART II: SCIENTIFIC INFORMATION.....	20
PHARMACEUTICAL INFORMATION.....	20
CLINICAL TRIALS	21
DETAILED PHARMACOLOGY.....	30
MICROBIOLOGY	30
TOXICOLOGY.....	32
REFERENCES.....	33
PART III: CONSUMER INFORMATION	34

PrMiconazole in 0.9% Sodium Chloride Injection**PART I: HEALTH PROFESSIONAL INFORMATION****SUMMARY PRODUCT INFORMATION**

Route of Administration	Dosage Form / Strength	All Nonmedicinal Ingredients
Intravenous infusion	Solution / 50 mg per 50 mL, 100 mg per 100 mL and 150 mg per 150 mL micafungin (as micafungin sodium)	Citric acid, sodium chloride, sodium citrate, water for injection.

INDICATIONS AND CLINICAL USE

Micafungin in 0.9% Sodium Chloride Injection (micafungin sodium) is indicated for use in adults and children 4 months and older for:

- Treatment of patients with Candidemia, Acute Disseminated Candidiasis, *Candida* Peritonitis and Abscesses infections. Micafungin sodium has not been adequately studied in patients with endocarditis, osteomyelitis and meningitis due to *Candida* infections.
- Treatment of patients with esophageal candidiasis.
- Prophylaxis of *Candida* infections in patients undergoing hematopoietic stem cell transplantation.

Geriatrics (≥ 65 years of age): No overall differences in safety or effectiveness have been observed between geriatric subjects (≥ 65 years) and younger subjects in clinical studies.

Pediatrics (4 months - 16 years): The safety and effectiveness of micafungin sodium in pediatric patients aged four (4) months to 16 years of age have been supported by evidence from adequate and well- controlled studies in adult and pediatric patients as well as pharmacokinetic data in pediatric patients.

The safety and efficacy of micafungin sodium in pediatric patients below the age of 4 months has not been established.

NOTE: The efficacy of micafungin sodium against infections caused by fungi other than *Candida* has not been established.

CONTRAINDICATIONS

Micafungin in 0.9% Sodium Chloride Injection is contraindicated in patients with hypersensitivity to micafungin, echinocandins or any component of this drug product (For a complete listing, see the **DOSAGE FORMS, COMPOSITION AND PACKAGING** section of the product monograph).

WARNINGS AND PRECAUTIONS

General

Isolated cases of serious hypersensitivity (anaphylaxis and anaphylactoid) reactions (including shock), Stevens-Johnson syndrome, and toxic epidermal necrolysis have been reported in patients receiving micafungin sodium. If these reactions occur, Micafungin in 0.9% Sodium Chloride Injection infusion should be discontinued and appropriate treatment administered (See **ADVERSE REACTIONS**).

Carcinogenesis, Mutagenesis, Impairment of Fertility

Hepatic carcinomas and adenomas were observed in a 6-month intravenous toxicology study with an 18-month recovery period of micafungin sodium in rats designed to assess the reversibility of hepatocellular lesions. Micafungin in 0.9% Sodium Chloride Injection should not be used in higher than recommended doses and for prolonged duration because the potential risk of liver tumor formation cannot be fully excluded.

Rats administered micafungin sodium for 3 months at 32 mg/kg/day (corresponding to 8 times the highest recommended human dose [150 mg/day], based on AUC comparisons), exhibited coloured patches/zones, multinucleated hepatocytes and altered hepatocellular foci after 1- or 3-month recovery periods, and adenomas were observed after a 20-month recovery period. Rats administered micafungin sodium at the same dose for 6 months exhibited adenomas after a 12-month recovery period; after an 18-month recovery period, an increased incidence of adenomas was observed, and additionally, carcinomas were detected. A lower dose of micafungin sodium (equivalent to 5 times the human AUC) in the 6-month rat study resulted in a lower incidence of adenomas and carcinomas following 18 months recovery. The duration of micafungin dosing in these rat studies (3 or 6 months) exceeds the usual duration of micafungin sodium dosing in patients, which is typically less than 1 month for treatment of esophageal candidiasis, but dosing may exceed 1 month for *Candida* prophylaxis.

Although the increase in carcinomas in the 6-month rat study did not reach statistical significance, the persistence of altered hepatocellular foci subsequent to micafungin dosing, and the presence of adenomas and carcinomas in the recovery periods suggest a causal relationship between micafungin sodium, altered hepatocellular foci, and hepatic neoplasms. Whole-life carcinogenicity studies of micafungin sodium in animals have not been conducted, and it is not known whether the hepatic neoplasms observed in treated rats also occur in other species, or if there is a dose threshold for this effect (See **TOXICOLOGY**).

Hepatic

Laboratory abnormalities in liver function tests have been seen in healthy volunteers and patients treated with micafungin sodium. In some patients with serious underlying conditions who were receiving micafungin sodium along with multiple concomitant medications, clinical hepatic abnormalities have occurred, and isolated cases of significant hepatic impairment, hepatitis, or worsening hepatic failure have been reported. Patients with hepatic impairment or who develop

abnormal liver function tests during Micafungin in 0.9% Sodium Chloride Injection therapy should be monitored for evidence of worsening hepatic function and evaluated for the risk/benefit of continuing Micafungin in 0.9% Sodium Chloride Injection therapy (See **ADVERSE REACTIONS**).

Renal

Elevations in BUN and creatinine, and isolated cases of significant renal impairment or acute renal failure have been reported in patients who received micafungin sodium. In fluconazole-controlled studies, the incidence of drug-related renal adverse events was 0.4% for micafungin sodium treated patients and 0.5% for fluconazole treated patients. Patients who develop abnormal renal function tests during Micafungin in 0.9% Sodium Chloride Injection therapy should be monitored for evidence of worsening renal function (See **ADVERSE REACTIONS**).

Hematological Effects

Acute intravascular hemolysis and hemoglobinuria were seen in a healthy volunteer during infusion of micafungin sodium (200 mg) and oral prednisolone (20 mg). This event was transient, and the subject did not develop significant anemia. Isolated cases of significant hemolysis and hemolytic anemia have also been reported in patients treated with micafungin sodium. Patients who develop clinical or laboratory evidence of hemolysis or hemolytic anemia during Micafungin in 0.9% Sodium Chloride Injection therapy should be monitored closely for evidence of worsening of these conditions and evaluated for the risk/benefit of continuing Micafungin in 0.9% Sodium Chloride Injection therapy (See **ADVERSE REACTIONS**).

Sexual Function/Reproduction

Male rats treated intravenously with micafungin sodium for 9 weeks showed vacuolation of the epididymal ductal epithelial cells at or above 10 mg/kg (about 0.6 times the recommended clinical dose for esophageal candidiasis, based on body surface area comparisons). Higher doses (about twice the recommended clinical dose, based on body surface area comparisons) resulted in higher epididymis weights and reduced numbers of sperm cells. In a 39-week intravenous study in dogs, seminiferous tubular atrophy and decreased sperm in the epididymis were observed at 10 and 32 mg/kg, doses equal to about 2 and 7 times the recommended clinical dose, based on body surface area comparisons. There was no impairment of fertility in animal studies with micafungin sodium (See **TOXICOLOGY**).

Special Populations

Pregnant Women: No adequate, well-controlled studies have been conducted in pregnant women. Micafungin in 0.9% Sodium Chloride Injection should be used during pregnancy only if the benefits outweigh the potential risks.

Micafungin sodium administration to pregnant rabbits (intravenous dosing on days 6 to 18 of gestation) resulted in visceral abnormalities and abortion at 32 mg/kg, a dose equivalent to about four times the recommended dose based on body surface area comparisons. Visceral abnormalities included abnormal lobation of the lung, levocardia, retrocaval ureter, anomalous right subclavian artery, and dilatation of the ureter. However, animal studies are not always predictive of human response (See **TOXICOLOGY**).

Nursing Women: It is not known whether micafungin is excreted in human milk. Micafungin was found in the milk of lactating, drug-treated rats. Caution should be exercised when Micafungin in 0.9% Sodium Chloride Injection is administered to a nursing woman.

Pediatrics (< 4 months): The safety and efficacy of micafungin sodium in pediatric patients below the age of 4 months has not been established. Micafungin in 0.9% Sodium Chloride Injection should not be used in children less than 4 months of age.

Geriatrics (≥ 65 years of age): A total of 418 subjects in clinical studies of micafungin sodium were 65 years of age and older, and 124 subjects were 75 years of age and older. No overall differences in safety or effectiveness were observed between these subjects and younger subjects. Other reported clinical experience has not identified differences in responses between the elderly and younger patients, but greater sensitivity of some older individuals cannot be ruled out.

ADVERSE REACTIONS

Adverse Drug Reaction Overview

Possible histamine-mediated symptoms have been reported with micafungin sodium, including rash, pruritus, facial swelling, and vasodilatation. Serious hypersensitivity (anaphylaxis and anaphylactoid) reactions (including shock) have been reported during administration of micafungin sodium.

Injection site reactions, including phlebitis and thrombophlebitis have been reported, at micafungin sodium doses of 50-150 mg/day. These reactions tended to occur more often in patients receiving micafungin sodium via peripheral intravenous administration.

Clinical Trial Adverse Drug Reactions

Because clinical trials are conducted under very specific conditions, the adverse reaction rates observed in the clinical trials may not reflect the rates observed in practice and should not be compared to the rates in the clinical trials of another drug. Adverse drug reaction information from clinical trials is useful for identifying drug-related adverse events and for approximating rates.

Adults

Candidemia and Other *Candida* Infections:

In a phase III, randomized, double-blind study for treatment of candidemia and other *Candida* infections, treatment-emergent adverse events occurred in 183/200 (92%), 187/202 (93%) and 171/193 (89%) adult patients in the micafungin sodium 100 mg/day, micafungin sodium 150 mg/day, and caspofungin (70/50 mg/day) treatment groups, respectively. Selected treatment-emergent adverse events occurring in ≥ 5% of the patients in any treatment study groups are shown in Table 1.

Table 1: Selected Treatment-Emergent Adverse Events* in Adult Patients with Candidemia and Other <i>Candida</i> Infections			
System Organ Class† (Preferred Term)‡	Micafungin sodium 100 mg n (%)	Micafungin sodium 150 mg n (%)	Comparator§ n (%)
Number of Patients	200	202	193
Gastrointestinal Disorders	81 (41)	89 (44)	76 (39)
Diarrhea	15 (8)	26 (13)	14 (7)
Nausea	19 (10)	15 (7)	20 (10)
Vomiting	18 (9)	15 (7)	16 (8)
Metabolism and Nutrition Disorders	77 (39)	83 (41)	73 (38)
Hypoglycemia	12 (6)	14 (7)	9 (5)
Hypernatremia	8 (4)	13 (6)	8 (4)
Hyperkalemia	10 (5)	8 (4)	5 (3)
General Disorders/Administration Site Conditions	59 (30)	56 (28)	51 (26)
Pyrexia	14 (7)	22 (11)	15 (8)
Investigations	36 (18)	49 (24)	37 (19)
Blood Alkaline Phosphatase Increased	11 (6)	16 (8)	8 (4)
Cardiac Disorders	35 (18)	48 (24)	36 (19)
Atrial Fibrillation	5 (3)	10 (5)	0

Patient base: all randomized patients who received at least 1 dose of study drug

* During IV treatment + 3 days

† MedDRA v5.0

‡ Within a system organ class patients may experience more than 1 adverse event.

§ 70 mg loading dose on day 1 followed by 50 mg/day thereafter (caspofungin)

In a second phase III, randomized, double-blind study for treatment of candidemia and other *Candida* infections, treatment-emergent adverse events occurred in 245/264 (93%) and 250/265 (94%) patients in the micafungin sodium (100 mg/day) and amphotericin B (3 mg/kg/day) treatment groups, respectively. The following treatment-emergent adverse events in patients at least 16 years of age were notable: nausea (10% vs. 8%), diarrhea (11% vs. 11%), vomiting (13% vs. 9%), abnormal liver function tests (4% vs. 3%), increased aspartate aminotransferase (3% vs. 2%) and increased blood alkaline phosphatase (3% vs. 2%) in the micafungin sodium and amphotericin B treatment groups, respectively.

Esophageal Candidiasis: In a phase III, randomized, double-blind study for treatment of esophageal candidiasis, a total of 202/260 (78%) adult patients who received micafungin sodium 150 mg/day and 186/258 (72%) patients who received intravenous fluconazole 200 mg/day experienced an adverse event. Treatment-emergent adverse events resulting in discontinuation were reported in 17 (7%) micafungin sodium treated patients; and in 12 (5%) fluconazole treated patients. Selected treatment-emergent adverse events occurring in $\geq 5\%$ of the patients in either treatment group are shown in Table 2.

Table 2: Selected Treatment-Emergent Adverse Events* in Adult Patients with Esophageal Candidiasis		
System Organ Class[†] (Preferred Term) [‡]	Micafungin sodium 150 mg/day n (%)	Comparator[§] n (%)
Number of Patients	260	258
Gastrointestinal Disorders	84 (32)	93 (36)
Diarrhea	27 (10)	29 (11)
Nausea	20 (8)	23 (9)
Vomiting	17 (7)	17 (7)
Abdominal Pain NOS	10 (3.8)	15 (5.8)
General Disorders / Administration Site Conditions	52 (20)	45 (17)
Pyrexia	34 (13)	21 (8)
Nervous System Disorders	42 (16)	40 (16)
Headache	22 (9)	20 (8)
Vascular Disorders	54 (21)	21 (8)
Phlebitis	49 (19)	13 (5)
Skin and Subcutaneous Tissue Disorders	36 (14)	26 (10)
Rash	14 (5)	6 (2)

Patient base: all randomized patients who received at least 1 dose of study drug

*During treatment + 3 days

[†] MedDRA v5.0

[‡] Within a system organ class patients may experience more than 1 adverse event

[§] Fluconazole 200 mg/day

Prophylaxis of *Candida* Infections in Hematopoietic Stem Cell Transplant Recipients: A double-blind, Phase III study was conducted in a total of 791 adult patients scheduled to undergo an autologous or allogeneic hematopoietic stem cell transplant. The median duration of treatment was 18 days (range 1 to 46 days) in both treatment arms.

All patients who received micafungin sodium (382) and all patients who received fluconazole (409) experienced at least one adverse event during the study. Treatment-emergent adverse events resulting in micafungin sodium discontinuation were reported in 15 (4%) patients while those resulting in fluconazole discontinuation were reported in 32 (8%) patients. Selected treatment-emergent adverse events occurring in $\geq 15\%$ of the patients in either treatment group are shown in Table 3.

Table 3: Selected Treatment-Emergent Adverse Events* in Adult Patients During Prophylaxis of <i>Candida</i> Infection in Hematopoietic Stem Cell Transplant Recipients		
System Organ Class[‡] (Preferred Term) [†]	Micafungin sodium 50 mg/day n (%)	Comparator[‡] n (%)
Number of Patients	382	409
Gastrointestinal Disorders	377 (99)	404 (99)
Diarrhea	294 (77)	327 (80)
Nausea	270 (71)	290 (71)
Vomiting	252 (66)	274 (67)
Abdominal Pain	100 (26)	93 (23)
Blood and Lymphatic System Disorders	368 (96)	385 (94)
Neutropenia	288 (75)	297 (73)
Thrombocytopenia	286 (75)	280 (69)
Skin and Subcutaneous Tissue Disorders	257 (67)	275 (67)
Rash	95 (25)	91 (22)
Nervous System Disorders	250 (65)	254 (62)
Headache	169 (44)	154 (38)
Psychiatric Disorders	233 (61)	235 (58)
Insomnia	142 (37)	140 (34)
Anxiety	84 (22)	87 (21)
Cardiac Disorders	133 (35)	138 (34)
Tachycardia	99 (26)	91 (22)

Patient base: all randomized adult patients who received at least 1 dose of study drug

*During treatment + 3 days

[‡] MedDRA v12.0

[†] Within a system organ class patients may experience more than 1 adverse reaction.

[‡] Fluconazole 400 mg/day

Selected Treatment-Emergent Adverse Events < 5% in All Adult Clinical Studies

The overall safety of micafungin sodium was assessed in 2748 adult patients and 520 volunteers in 40 clinical studies, including the invasive candidiasis, esophageal candidiasis and prophylaxis studies, who received single or multiple doses of micafungin sodium, ranging from 12.5 mg to ≥ 150 mg/day in adult patients. The mean duration of treatment was 18.9 days.

Overall, 2497 (91%) adult patients who received micafungin sodium experienced an adverse event.

The most common serious adverse events in adult patients treated with micafungin sodium were sepsis 63 (2%), respiratory failure 86 (3%), septic shock 50 (2%), pneumonia 39 (1%), hypotension 36 (1%), multi-organ failure 32 (1%), renal failure 32 (1%), cardiac arrest 24 (1%), pyrexia 23 (1%), and bacterial sepsis 21 (1%).

Other selected treatment-adverse events reported in <5% in all adult patients treated with micafungin sodium are listed below:

- *Blood and lymphatic system disorders*: coagulopathy, pancytopenia, thrombotic thrombocytopenic purpura
- *Cardiac disorders*: cardiac arrest, myocardial infarction, pericardial effusion
- *General disorders and administration site conditions*: infusion reaction, injection site thrombosis
- *Hepatobiliary disorders*: hepatocellular damage, including fatal cases, hepatomegaly, jaundice, hepatic failure
- *Immune system disorders*: hypersensitivity, anaphylactic reaction
- *Nervous system disorders*: convulsions, encephalopathy, intracranial hemorrhage

- *Psychiatric disorders*: delirium
- *Skin and subcutaneous tissue disorders*: urticarial

Pediatrics

The overall safety of micafungin sodium was assessed in 479 patients, aged 3 days through 16 years of age, who received at least one dose of micafungin sodium in 11 clinical studies. The mean treatment duration was 24.8 days. A total of 246 patients received at least one dose of micafungin sodium 2 mg/kg or higher.

Of these pediatric patients, 264 (55%) were male and 319 (67%) were Caucasian. The age distribution was as follows: 116 (24%) less than 2 years, 108 (23%) between 2 and 5 years, 140 (29%) between 6 and 11 years, and 115 (24%) between 12 and 16 years of age.

In all pediatric studies with micafungin sodium, 439/479 (92%) patients experienced at least one treatment-emergent adverse event.

Two studies that included pediatric patients were randomized, double-blind, and active-controlled: The invasive candidiasis and candidemia study investigated the efficacy and safety of micafungin sodium (2 mg/kg/day for patients weighing \leq 40 kg and 100 mg/day for patients weighing $>$ 40 kg) compared to amphotericin B (3 mg/kg/day) in 112 pediatric patients. Treatment-emergent adverse events occurred in 51/56 (91%) of patients in the micafungin sodium group and 52/56 (93%) of patients in the amphotericin B group. Treatment-emergent adverse events resulting in micafungin sodium discontinuation were reported in 2 (4%) pediatric patients; these events were acute renal failure and cardiac arrest.

The prophylaxis study in patients undergoing HSCT investigated the efficacy of micafungin sodium (1 mg/kg/day for patients weighing \leq 50 kg and 50 mg/day for patients weighing $>$ 50 kg) as compared to fluconazole (8 mg/kg/day for patients weighing \leq 50 kg and 400 mg/day for patients weighing $>$ 50 kg). All 91 pediatric patients experienced at least one treatment-emergent adverse event. Three (7%) pediatric patients discontinued micafungin sodium due to adverse events; these events were arthralgia, intracranial hemorrhage and meningitis.

The incidence of the following treatment-emergent adverse events was higher in pediatric patients than in adult patients: rash, pruritis, urticaria, infusion related reactions, and alanine aminotransferase increased.

The selected treatment-emergent adverse events (\geq 15%) occurring in pediatric patients and more frequently in a micafungin sodium group, for all micafungin sodium pediatric studies and for the two comparative studies (candidemia and prophylaxis) described in the above section are shown in Table 4.

Table 4: Selected Treatment-Emergent Adverse Events ≥ 15% in Pediatric Patients from All Clinical Studies, in Pediatric Patients with Candidemia and Other <i>Candida</i> Infections (C/IC), and in Hematopoietic Stem-Cell Transplant Recipients During Prophylaxis of <i>Candida</i> Infections					
System Organ Class[¥] (Preferred Term) [†]	All Micafungin sodium- treated Patients n = 479 n (%)	C/IC		Prophylaxis	
		Micafungin sodium n = 56 n (%)	Comparator[‡] n = 56 n (%)	Micafungin sodium n = 43 n (%)	Comparator[§] n = 48 n (%)
Gastrointestinal disorders	285 (60)	22 (40)	18 (32)	43 (100)	45 (94)
Vomiting	146 (31)	10 (18)	8 (14)	28 (65)	32 (67)
Diarrhea	106 (22)	4 (7)	5 (9)	22 (51)	31 (65)
Nausea	91 (19)	4 (7)	4 (7)	30 (70)	25 (52)
Abdominal pain	76 (16)	2 (4)	2 (4)	15 (35)	12 (25)
Abdominal distension	29 (6)	1 (2)	1 (2)	8 (19)	6 (13)
General disorders and administration site conditions	256 (53)	14 (25)	14 (25)	41 (95)	46 (96)
Pyrexia	103 (22)	5 (9)	9 (16)	26 (61)	31 (65)
Infusion related reaction	24 (5)	0	3 (5)	7 (16)	4 (8)
Skin and subcutaneous tissue disorders	197 (41)	11 (20)	8 (14)	33 (77)	38 (79)
Pruritus	54 (11)	0	1 (2)	14 (33)	15 (31)
Rash	55 (12)	1 (2)	1 (2)	13 (30)	13 (27)
Urticaria	24 (5)	0	1 (2)	8 (19)	4 (8)
Respiratory, thoracic and mediastinal disorders	194 (41)	9 (16)	13 (23)	30 (70)	33 (69)
Epistaxis	45 (9)	0	0	4 (9)	8 (17)
Blood and lymphatic system disorders	161 (34)	17 (30)	13 (23)	40 (93)	44 (92)
Thrombocytopenia	70 (15)	5 (9)	3 (5)	31 (72)	37 (77)
Neutropenia	61 (13)	3 (5)	4 (7)	33 (77)	34 (71)
Anemia	63 (13)	10 (18)	6 (11)	22 (51)	24 (50)
Febrile neutropenia	23 (5)	0	0	7 (16)	7 (15)
Investigations	191 (40)	12 (21)	8 (14)	24 (56)	25 (52)
Alanine aminotransferase increased	45 (10)	0	0	7 (16)	1 (2)
Urine output decreased	18 (4)	0	0	10 (23)	8 (17)
Cardiac disorders	97 (20)	7 (13)	3 (5)	10 (23)	17 (35)
Tachycardia	47 (10)	2 (4)	1 (2)	7 (16)	12 (25)
Renal and urinary disorders	78 (16)	4 (7)	4 (7)	16 (37)	15 (31)
Hematuria	18 (4)	0	0	10 (23)	7 (15)
Psychiatric disorders	80 (17)	3 (5)	1 (2)	20 (47)	9 (19)
Anxiety	35 (7)	0	0	10 (23)	3 (6)

Patient base: all randomized patients who received at least one dose of study drug.

[¥]MedDRA v12.0

[†]Within a system organ class, patients may experience more than 1 adverse reaction.

[‡]Amphotericin B

[§]Fluconazole

Other selected treatment-emergent adverse events reported in < 15% of pediatric patients receiving micafungin sodium in all clinical studies were:

- *Hepatobiliary disorders:* hyperbilirubinemia, hepatomegaly
- *Immune system disorders:* anaphylactoid reaction
- *Investigations:* aspartate aminotransferase increased
- *Renal disorders:* renal failure

Post-Market Adverse Drug Reactions

The following adverse events have been identified during the post-approval use of micafungin sodium. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency.

- *Blood and lymphatic system disorders:* hemolytic anemia, disseminated intravascular coagulation
- *Hepatobiliary disorders:* hyperbilirubinemia, hepatic disorder, hepatocellular damage
- *Investigations:* hepatic function abnormal, white blood cell count decreased
- *Renal and urinary disorders:* acute renal failure and renal impairment
- *Skin and subcutaneous tissue disorders:* Stevens-Johnson syndrome, toxic epidermal necrolysis, erythema multiforme
- *Vascular disorders:* shock

DRUG INTERACTIONS

Drug-Drug Interactions

A total of 14 clinical drug-drug interaction studies were conducted in healthy volunteers to evaluate the potential for interaction between micafungin sodium and amphotericin B, mycophenolate mofetil, cyclosporine, tacrolimus, prednisolone, sirolimus, nifedipine, fluconazole, itraconazole, voriconazole, ritonavir, or rifampin. In these studies, no interaction that altered the pharmacokinetics of micafungin was observed.

Co-administration of micafungin sodium and amphotericin B deoxycholate was associated with a 30% increase in amphotericin B deoxycholate exposure. The clinical significance of this increase is unclear however this co-administration should only be used when the benefits clearly outweigh the risks, with close monitoring of amphotericin B deoxycholate toxicities.

Co-administration of voriconazole with micafungin sodium was associated with approximately 20% reduction of voriconazole exposure on average. In the same study co-administration of placebo and voriconazole was associated with a similar reduction of voriconazole exposure. When the effect of micafungin sodium was corrected for the effect of placebo, there was no significant effect of micafungin sodium on the pharmacokinetics of voriconazole.

There was no effect of a single dose or multiple doses of micafungin sodium on mycophenolate mofetil, cyclosporine, tacrolimus, prednisolone, voriconazole and fluconazole pharmacokinetics.

Sirolimus AUC was increased by 21% with no effect on C_{max} in the presence of steady-state micafungin sodium compared with sirolimus alone. Nifedipine AUC and C_{max} were increased by 18% and 42%, respectively, in the presence of steady-state micafungin sodium compared with nifedipine alone. Itraconazole AUC and C_{max} were increased by 22% and 11%, respectively, in the presence of steady-state micafungin sodium compared with itraconazole alone. Patients receiving sirolimus, itraconazole or nifedipine in combination with Micafungin in 0.9% Sodium Chloride Injection should be monitored for sirolimus, itraconazole or nifedipine toxicity. The sirolimus, itraconazole or nifedipine dosage should be reduced if necessary.

Micafungin is not an inhibitor of P-glycoprotein and, therefore, would not be expected to alter P-glycoprotein-mediated drug transport activity.

Drug-Food Interactions

Interactions with food have not been established.

Drug-Herb Interactions

Interactions with herbs have not been established.

Drug-Laboratory Test Interactions

Interactions with laboratory tests have not been established.

DOSAGE AND ADMINISTRATION

Dosing Considerations

Micafungin in 0.9% Sodium Chloride Injection in GALAXY Container (PL 2501 Plastic) is a premixed solution intended only for intravenous infusion.

- No further dilution is necessary.
- Check for minute leaks by squeezing container firmly. If leaks are detected, discard solution as sterility may be impaired.

Do not mix or co-infuse Micafungin in 0.9% Sodium Chloride Injection with other medications. Micafungin has been shown to precipitate when mixed directly with a number of other commonly used medications.

Micafungin in 0.9% Sodium Chloride Injection should be administered by a slow intravenous infusion over the period of 1 hour. More rapid infusions may result in more frequent histamine mediated reactions.

NOTE: An existing intravenous line should be flushed with 0.9% Sodium Chloride Injection, USP, prior to infusion of Micafungin in 0.9% Sodium Chloride Injection.

Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration, whenever solution and container permit. Do not use if there is any evidence of precipitation or foreign matter.

- Micafungin in 0.9% Sodium Chloride Injection is preservative-free. Discard partially used bags. Do not freeze.
- Solution is stable for up to 10 days in the original carton at room temperature up to 77°F (25°C).
- Product does not require light protection during administration.

Recommended Dose and Dosage Adjustment

Table 5: Micafungin in 0.9% Sodium Chloride Injection Dosage in Adults	
Indication	Recommended Dose
	Adults
Treatment of Candidemia, Acute Disseminated Candidiasis, <i>Candida</i> Peritonitis and Abscesses infections	100 mg/day ¹
Prophylaxis of <i>Candida</i> Infections in HSCT Recipients	50 mg/day ²
Treatment of Esophageal Candidiasis	150 mg/day ³

¹ In patients treated successfully for candidemia and other *Candida* infections, the mean duration of treatment was 15 days (range 10-47 days)

² In hematopoietic stem cell transplant (HSCT) recipients who experienced success of prophylactic therapy, the mean duration of prophylaxis was 19 days (range 6-51 days)

³ In patients treated successfully for esophageal candidiasis, the mean duration of treatment was 15 days (range 10- 30 days)

Micafungin in 0.9% Sodium Chloride Injection Dose and Schedule for Pediatric Patients

The recommended doses for pediatric patients based on indication and weight are shown in Table 6.

Table 6: Micafungin in 0.9% Sodium Chloride Injection Dosage in Pediatric Patients 4 Months or Older		
Indication	Pediatrics (aged ≥ 4 months) Dose Given Once Daily	
	30 kg or less	Greater than 30 kg
Treatment of Candidemia, Acute Disseminated Candidiasis, <i>Candida</i> Peritonitis and Abscesses infections	2 mg/kg maximum daily dose 100 mg	
Prophylaxis of <i>Candida</i> Infections in HSCT Recipients	1 mg/kg maximum daily dose 50 mg	
Treatment of Esophageal Candidiasis	3 mg/kg	2.5 mg/kg maximum daily dose 150 mg

A loading dose is not required; typically, 85% of the steady-state concentration is achieved after three daily Micafungin in 0.9% Sodium Chloride Injection doses.

Gender/Race: No dose adjustment is required based on gender or race (see **ACTION AND CLINICAL PHARMACOLOGY – Special Populations**).

Use in Patients with Renal Impairment: No dose adjustment is required in patients with renal impairment (see **ACTION AND CLINICAL PHARMACOLOGY – Special Populations**). Supplementary dosing is not required following hemodialysis (see **ACTION AND CLINICAL**

PHARMACOLOGY – Special Populations).

Use in Patients with Hepatic Impairment: No dose adjustment is required in patients with mild, moderate or severe hepatic impairment (see **ACTION AND CLINICAL PHARMACOLOGY – Special Populations**).

Use with Other Drugs: No dose adjustment for Micafungin in 0.9% Sodium Chloride Injection is required with concomitant use of mycophenolate mofetil, cyclosporine, tacrolimus, prednisolone, sirolimus, nifedipine, fluconazole, voriconazole, itraconazole, amphotericin B, ritonavir, or rifampin (see **DRUG INTERACTIONS**).

Administration

DIRECTIONS FOR USE OF MICAFUNIN IN 0.9% SODIUM CHLORIDE INJECTION IN GALAXY CONTAINER (PL 2501 PLASTIC)

Preparation for Administration

- Suspend container from support.
- Remove protector from outlet port at bottom of container.
- Attach Intravenous administration set to outlet port.

OVERDOSAGE

For the most recent information in the management of a suspected drug overdose, contact your regional poison control centre or Health Canada's toll-free number, 1-844-POISON-X (1-844-764-7669).

Micafungin sodium is highly protein bound and, therefore, is not dialyzable. No cases of micafungin sodium overdosage have been reported. Repeated daily doses of up to 8 mg/kg (maximum total dose of 896 mg) in adult patients, up to 6 mg/kg in pediatric patients \geq 4 months of age, and up to 10 mg/kg in pediatric patients < 4 months of age have been administered in clinical studies with no reported dose-limiting toxicity. The minimum lethal dose of micafungin sodium is 125 mg/kg in rats, equivalent to 8 times the highest recommended adult clinical dose (150 mg) and approximately 7 times the highest pediatric clinical dose (3 mg/kg) based on body surface area comparisons.

ACTION AND CLINICAL PHARMACOLOGY**Mechanism of Action**

Micafungin is a semisynthetic lipopeptide (echinocandin) compound synthesized by a chemical modification of a fermentation product of *Coleophoma empetri* F-11899. Micafungin is a member of a class of antifungal drugs known as echinocandins. Micafungin inhibits the synthesis of 1,3- beta-D glucan, an essential component of fungal cell walls, which is not present in mammalian cells.

Drug Resistance

There have been reports of clinical failures in patients receiving micafungin therapy due to the development of drug resistance. Some of these reports have identified specific mutations in the *FKS* protein component of the glucan synthase enzyme that are associated with higher MICs and breakthrough infection.

Pharmacokinetics

The pharmacokinetics of micafungin were determined in healthy subjects, hematopoietic stem cell transplant recipients, and patients with esophageal candidiasis or invasive candidiasis up to a maximum daily dose of 8 mg/kg body weight.

The relationship of area under the concentration-time curve (AUC) to micafungin dose was linear over the daily dose range of 50 mg to 150 mg and 3 mg/kg to 8 mg/kg body weight.

Steady-state pharmacokinetic parameters in relevant patient populations after repeated daily administration are presented in Table 7.

Population	N	Dose (mg)	Pharmacokinetic Parameters (Mean ± Standard Deviation)				
			C _{max} (µg/mL)	AUC ₀₋₂₄ (µg·h/mL)	T _{1/2} (h)	Cl (mL/min/kg)	
Patients with IC [Day 1]	20	100	5.7±2.2	83±51	14.5±7.0	0.359 ±0.179	
	20	100	10.1±4.4	97±29	13.4±2.0	0.298 ±0.115	
HIV-Positive Patients with EC [Day 1]	20	50	4.1±1.4	36±9	14.9±4.3	0.321 ±0.098	
	20	100	8.0±2.4	108±31	13.8±3.0	0.327 ±0.093	
	14	150	11.6±3.1	151±45	14.1±2.6	0.340 ±0.092	
	[Day 14 or 21]	20	50	5.1±1.0	54±13	15.6±2.8	0.300±0.063
		20	100	10.1±2.6	115±25	16.9±4.4	0.301±0.086
		14	150	16.4±6.5	167±40	15.2±2.2	0.297±0.081
HSCT Recipients [Day 7]	8	Per kg 3	21.1±2.84	234±34	14.0±1.4	0.214±0.031	
	10	4	29.2±6.2	339±72	14.2±3.2	0.204±0.036	
	8	6	38.4±6.9	479±157	14.9±2.6	0.224±0.064	
	8	8	60.8±26.9	663±212	17.2±2.3	0.223±0.081	

HIV = human immunodeficiency virus; EC = esophageal candidiasis; HSCT = hematopoietic stem cell transplant; IC = Candidemia and other *Candida* infections; *AUC_{0-infinity} is presented for day 1 and AUC₀₋₂₄ is presented for steady state.

Distribution: The mean ± standard deviation volume of distribution of micafungin at terminal phase was 0.39 ± 0.11 L/kg body weight when determined in adult patients with esophageal candidiasis at the dose range of 50 mg to 150 mg. Micafungin is highly (> 99%) protein bound *in vitro*, independent of plasma concentrations over the range of 10 to 100 mcg/mL. The primary binding protein is albumin; however, micafungin, at therapeutically relevant concentrations, does not competitively displace bilirubin binding to albumin. Micafungin also binds to a lesser extent to α₁-acid-glycoprotein.

Metabolism: Micafungin is metabolized to M-1 (catechol form) by arylsulfatase, with further metabolism to M-2 (methoxy form) by catechol-O-methyltransferase. M-5 is formed by hydroxylation at the side chain (ω -1 position) of micafungin catalyzed by cytochrome P450 (CYP) isozymes. Even though micafungin is a substrate for and a weak inhibitor of CYP3A *in vitro*, hydroxylation by CYP3A is not a major pathway for micafungin metabolism *in vivo*. Micafungin is neither a P-glycoprotein substrate nor inhibitor *in vitro*.

In four healthy volunteer studies, the ratio of metabolite to parent exposure (AUC) at a dose of 150 mg/day was 6% for M-1, 1% for M-2, and 6% for M-5. In patients with esophageal candidiasis, the ratio of metabolite to parent exposure (AUC) at a dose of 150 mg/day was 11% for M-1, 2% for M-2, and 12% for M-5.

Excretion: The excretion of radioactivity following a single intravenous dose of ^{14}C -micafungin sodium for injection (25 mg) was evaluated in healthy volunteers. At 28 days after administration, mean urinary and fecal recovery of total radioactivity accounted for 82.5% (76.4% to 87.9%) of the administered dose. Fecal excretion is the major route of elimination (total radioactivity at 28 days was 71.0% of the administered dose).

Special Populations

Pediatrics (4 months – 16 years): Micafungin pharmacokinetics in 229 pediatric patients ≥ 4 months through 16 years of age were characterized using population pharmacokinetics. Population pharmacokinetic analysis in pediatric patients identified body weight as the major determinant of total systemic clearance. As a result, weight-based (mg/kg) dosing was used to reduce between-subject variability in exposure and to equate pediatric exposure with adult exposure. Micafungin exposure was dose proportional across the dose and age range studied (Table 8).

Body weight group	N	Dose [§] mg/kg	C _{max,ss} [†] (mcg/mL)	AUC _{ss} [†] (mcg·h /mL)	t _{1/2} [‡] (h)	CL [‡] (mL/min/kg)
30 kg or less	149	1.0	7.1 +/- 4.7	55 +/- 16	12.5 +/- 4.6	0.328 +/- 0.091
		2.0	14.2 +/- 9.3	109 +/- 31		
		3.0	21.3 +/- 14.0	164 +/- 47		
Greater than 30 kg	80	1.0	8.7 +/- 5.6	67 +/- 17	13.6 +/- 8.8	0.241 +/- 0.061
		2.0	17.5 +/- 11.2	134 +/- 33		
		2.5	23.0 +/- 14.5	176 +/- 42		

[§] Or the equivalent if receiving the adult dose (50, 100, or 150 mg)

[†] Derived from simulations from the population PK model

[‡] Derived from the population PK model

Geriatric (≥ 65 years): The exposure and disposition of a 50 mg micafungin sodium dose administered as a single 1-hour infusion to 10 healthy subjects aged 66-78 years were not significantly different from those in 10 healthy subjects aged 20-24 years. No dose adjustment is necessary for the elderly.

Race and Gender: No dose adjustment of Micafungin in 0.9% Sodium Chloride Injection is required based on gender or race. After 14 daily doses of 150 mg to healthy subjects, micafungin AUC in women was greater by approximately 23% compared with men, due to smaller body weight. No notable differences among white, black, and Hispanic subjects were seen. The micafungin AUC was greater by 26% in Japanese subjects compared to blacks, due to smaller body weight.

Renal Impairment: Micafungin in 0.9% Sodium Chloride Injection does not require dose adjustment in patients with renal impairment. A single 1-hour infusion of 100 mg micafungin sodium was administered to 9 adult subjects with severe renal impairment (creatinine clearance < 30 mL/min) and to 9 age-, gender-, and weight-matched subjects with normal renal function (creatinine clearance > 80 mL/min). The maximum concentration (C_{max}) and AUC were not significantly altered by severe renal impairment.

Since micafungin is highly protein bound, it is not dialyzable. Supplementary dosing should not be required following hemodialysis.

Hepatic Impairment: A single 1-hour infusion of 100 mg micafungin sodium was administered to 8 adult subjects with moderate hepatic impairment (Child-Pugh score 7-9) and 8 age-, gender-, and weight-matched subjects with normal hepatic function. The C_{max} and AUC values of micafungin were lower by approximately 22% in subjects with moderate hepatic impairment compared to normal subjects. This difference in micafungin exposure does not require dose adjustment of Micafungin in 0.9% Sodium Chloride in patients with moderate hepatic impairment.

A single 1-hour infusion of 100 mg micafungin sodium was administered to 8 adult subjects with severe hepatic impairment (Child-Pugh score 10-12) and 8 age-, gender-, ethnic- and weight-matched subjects with normal hepatic function. The mean C_{max} and AUC values of micafungin were lower by approximately 30% in subjects with severe hepatic impairment compared to normal subjects. The mean C_{max} and AUC values of M-5 metabolite were approximately 2.3-fold higher in subjects with severe hepatic impairment compared to normal subjects; however, this exposure (parent and metabolite) was comparable to that in patients with systemic *Candida* infection. Therefore, no micafungin dose adjustment is necessary in patients with severe hepatic impairment.

Studies in pediatric subjects with hepatic impairment have not been conducted.

STORAGE AND STABILITY

Store Micafungin in 0.9% Sodium Chloride Injection in refrigerator at 2°C to 8°C in the original carton. The drug is sensitive to light, keep in original container until ready to use. Do not use after the expiration date printed on the carton and container label.

The drug product should be stored under prescribed conditions in the refrigerator. In rare cases, if needed, the unopened/sterile product in the original carton may be stored at room temperature of 25°C

up to maximum 10 days. Any product stored a room temperature exceeding 10 days period should be discarded.

Do not freeze and do not use Micafungin in 0.9% Sodium Chloride Injection if it has been frozen.

Note: Micafungin in 0.9% Sodium Chloride Injection is preservative-free.

SPECIAL HANDLING INSTRUCTIONS

Not applicable.

DOSAGE FORMS, COMPOSITION AND PACKAGING

Micafungin in 0.9% Sodium Chloride Injection is supplied as a refrigerated 50 mL, 100 mL, or 150 mL single-dose GALAXY container in the following presentations:

50 mg/50 mL (1 mg/mL): containing 50 mg of micafungin (equivalent to 50.87 mg of micafungin sodium); 36 mg citric acid, anhydrous, USP; 450 mg sodium chloride, USP; 92 mg sodium citrate, dihydrate, USP;

100 mg/100 mL (1 mg/mL): containing 100 mg of micafungin (equivalent to 101.73 mg of micafungin sodium); 72 mg citric acid, anhydrous, USP; 900 mg sodium chloride, USP; 184 mg sodium citrate, dihydrate, USP;

150 mg/150 mL (1 mg/mL): containing 150 mg of micafungin (equivalent to 152.60 mg of micafungin sodium); 108 mg citric acid, anhydrous, USP; 1350 mg sodium chloride, USP; 276 mg sodium citrate, dihydrate, USP.

The premixed solution is clear and colorless, with a pH range of 4.5 to 5.1.

PART II: SCIENTIFIC INFORMATION

PHARMACEUTICAL INFORMATION

Drug Substance


Proper name: micafungin sodium

Chemical name: Pneumocandin A0, 1-[(4R,5R)-4,5-dihydroxy-N²-[4-[5-[4-(pentyloxy)phenyl]-3-isoxazolyl]benzoyl]-L-ornithine]-4-[(4S)-4-hydroxy-4-[4-hydroxy-3-(sulfooxy)phenyl]-L-threonine]-, monosodium salt.

Molecular formula: C₅₆H₇₀N₉NaO₃S

Molecular mass: 1292.26

Structural formula:



Physicochemical properties:

Physical Form: Micafungin sodium is a light-sensitive, hygroscopic white powder that is freely soluble in water, isotonic sodium chloride solution, N,N-dimethylformamide and dimethylsulfoxide, slightly soluble in methyl alcohol, and practically insoluble in acetonitrile, ethyl alcohol (95%), acetone, diethyl ether and n-hexane.

pH: The pH of the reconstituted solution, with 0.9% Sodium Chloride Injection, USP, is between 5.0 - 7.0.

CLINICAL TRIALS

The results from clinical studies by indication are presented below.

The safety and effectiveness of micafungin sodium in pediatric patients aged four (4) months to 16 years of age have been supported by evidence from 2 studies that enrolled adult and pediatric patients, as well as pharmacokinetic data in pediatric patients.

Treatment of Candidemia and Other *Candida* Infections

Study demographics and trial design

Table 9: Summary of Patient Demographics for Micafungin sodium Clinical Studies in the Treatment of Candidemia and Other <i>Candida</i> Infections				
Dosage, Route of Administration and Duration	Number of Study Subjects	Mean Age (Range)	Gender	Race
Study 03-0-192: Phase III, randomized (1:1:1), double-blind, parallel group non-inferiority study (in Adult Patients Only)				
Micafungin sodium 100 mg/day, intravenous infusion, minimum 14 days to a maximum of 4 weeks*.	191	56.61 (18 - 92)	M = 107 (56%) F = 84 (44%)	Caucasian = 134 (70%) Black = 21 (11%) Other = 36 (19%)
Micafungin sodium 150 mg/day, intravenous infusion, minimum of 14 days to a maximum of 4 weeks*.	199	55.41 (18 - 90)	M = 117 (59%) F = 82 (41%)	Caucasian = 129 (65%) Black = 36 (18%) Other = 34 (17%)
Caspofungin 70 mg on day 1 and 50 mg/day thereafter, intravenous infusion, minimum 14 days to a maximum of 4 weeks*.	188	55.84 (19 - 95)	M = 112 (60%) F = 76 (40%)	Caucasian = 129 (69%) Black = 26 (14%) Other = 33 (18%)
Study FG-463-21-08: Phase III, randomized, double-blind, parallel group study (Adult and Pediatric Patients)				
Once daily IV infusion of Micafungin sodium 100 mg/day [†] in patients less than 40 kg and 2 mg/kg in patients weighing 40 kg or more for a minimum of 14 days to a maximum of 4 weeks*.	Adults			
	247 [†]	52.7 (18 - 89)	M = 155 (63%) F = 92 (37%)	Caucasian = 149 (60%) Black = 13 (5%) Other = 85 (34%)
	Pediatrics			
	48	4.0 (0 - 15)	M = 31 (65%) F = 17 (35%)	Caucasian = 27 (56%) Black = 5 (10%) Other = 16 (33%)
Once daily IV infusion of AmBisome 3 mg/kg for a minimum of 14 days to a maximum of 4 weeks*.	Adults			
	247 [†]	53.6 (16 - 97)	M = 147 (60%) F = 100 (40%)	Caucasian = 153 (62%) Black = 10 (4%) Other = 84 (34%)
	Pediatrics			
	50	2.2 (0 - 15)	M = 30 (60%) F = 20 (40%)	Caucasian = 25 (50%) Black = 9 (18%) Other = 16 (32%)

* Administration of study drug could be prolonged up to a maximum of 8 weeks for patients with chronic disseminated candidiasis, *Candida* osteomyelitis or *Candida* endocarditis.

† Micafungin dose increases to 200 mg/day (patients weighing > 40 kg) or 4 mg/kg/day (patients weighing ≤ 40 kg) were permitted. Amphotericin B dose increases up to 5 mg/kg were permitted.

‡ In the presentation of the Efficacy Analysis for study FG-463-21-08 (Table 11), the Independent Data Review Board modified Full Analysis Set (IDRB-mFAS) was used. The IDRB-mFAS population was N=248 for Micafungin sodium and N = 246 for amphotericin B.

Study Results

Adult Only Study

Two dose levels of micafungin sodium were evaluated in a phase III, randomized, double-blind study to determine the efficacy and safety versus caspofungin in adult patients with invasive candidiasis and candidemia. Patients were randomized to receive once daily intravenous infusions (IV) of micafungin sodium, either 100 mg/day or 150 mg/day or caspofungin (70 mg loading dose followed by 50 mg maintenance dose). Patients were stratified by APACHE II score (≤ 20 or > 20) and by geographic region. Patients with *Candida* endocarditis were excluded from this analysis. Outcome was assessed by overall treatment success based on clinical (complete resolution or improvement in attributable signs and symptoms and radiographic abnormalities of the *Candida* infection and no additional antifungal therapy) and mycological (eradication or presumed eradication) response at the end of IV therapy. Deaths that occurred during IV study drug therapy were treated as failures.

In this study, 111/578 (19.2%) of the patients had baseline APACHE II scores of > 20 , and 50/578 (8.7%) were neutropenic at baseline (absolute neutrophil count less than 500 cells/mm³). Outcome, relapse and mortality data are shown for the recommended dose of micafungin sodium (100 mg/day) and caspofungin in Table 10.

Table 10: Efficacy Analysis: Treatment Success in Patients with Candidemia and other <i>Candida</i> Infection in Study 03-0-192 (Adult Patients Only)		
	Micafungin sodium 100 mg/day n (%) % treatment difference (95% CI)	Comparator¹ n (%)
Treatment Success at End of IV Therapy²	135/191 (70.7) 7.4 (-2.0, 16.3)	119/188 (63.3)
Success in Patients with Neutropenia at Baseline	14/22 (63.6)	5/11 (45.5)
Success by Site of Infection		
Candidemia	116/163 (71.2)	103/161 (64)
Abscess	4/5 (80)	5/9 (55.6)
Acute disseminated³	6/13 (46.2)	5/9 (55.6)
Endophthalmitis	1/3	1/1
Chorioretinitis	0/3	0
Skin	1/1	0
Kidney	2/2	1/1
Pancreas	1/1	0
Peritoneum	1/1	0
Lung/Skin	0/1	0
Lung/Spleen	0/1	0
Liver	0	0/2
Intraabdominal abscess	0	3/5
Chronic disseminated	0/1	0
Peritonitis	4/6 (66.7)	2/5 (40.0)

Table 10: Efficacy Analysis: Treatment Success in Patients with Candidemia and other *Candida* Infection in Study 03-0-192 (Adult Patients Only)

	Micafungin sodium 100 mg/day n (%) % treatment difference (95% CI)	Comparator¹ n (%)
Success by Organism⁴		
<i>C. albicans</i>	57/81 (70.4)	45/73 (61.6)
<i>C. glabrata</i>	16/23 (69.6)	19/31 (61.3)
<i>C. tropicalis</i>	17/27 (63)	22/29 (75.9)
<i>C. parapsilosis</i>	21/28 (75)	22/39 (56.4)
<i>C. krusei</i>	5/8 (62.5)	2/3 (66.7)
<i>C. guilliermondii</i>	1/2	0/1
<i>C. lusitaniae</i>	2/3 (66.7)	2/2
Relapse through 6 Weeks⁵		
Overall	49/135 (36.3)	44/119 (37)
Culture confirmed relapse	5	4
Required systemic antifungal therapy	11	5
Died during follow-up	17	16
Not assessed	16	19
Overall study mortality		
Mortality during IV therapy	58/200 (29)	51/193 (26.4)
Mortality attributed to fungal infection	28/200 (14) 7/200 (3.5)	27/193 (14) 7/193 (3.6)

¹ 70 mg loading dose on day 1 followed by 50 mg/day thereafter (casposfungin)

² All patients who received at least one dose of study medication and had documented invasive candidiasis or candidemia. Patients with *Candida* endocarditis were excluded from the analyses.

³ A patient may have had > 1 organ of dissemination

⁴ A patient may have had > 1 baseline infection species

⁵ All patients who had a culture confirmed relapse or required systemic antifungal therapy in the post treatment period for a suspected or proven *Candida* infection. Also includes patients who died or were not assessed in follow-up.

Adult and Pediatric Study

Adults

In a second phase III, randomized, double-blind, parallel group study in 494 adult and 98 pediatric patients, the efficacy and safety of micafungin sodium versus amphotericin B were determined in non- neutropenic and neutropenic (absolute neutrophil count [ANC] < 500 cells/mcL) patients with invasive candidiasis or candidemia. Patients received an initial dose of IV micafungin 100 mg qd (2.0 mg/kg for patients weighing ≤ 40 kg) or IV amphotericin B 3 mg/kg by 1-hour infusion in a blinded manner.

The primary endpoint was success based on clinical (complete or partial) and mycological (eradication or presumed eradication) response at the end of IV therapy and no additional systemic antifungal therapy. Deaths that occurred during IV study drug therapy were treated as failures.

In this study a total of 32/248 (12.9%) of adult patients in the micafungin group and 25/246 (10%) of adult patients in the amphotericin B group were neutropenic at baseline (absolute neutrophil count less than 500 cells/mcL). A baseline APACHE II score of > 20 was recorded for 61/226 (27%) of patients in the micafungin group and 53/219 (24.2%) in the amphotericin B group. Outcome, relapse and mortality data from adults are shown for the recommended dose of micafungin sodium (100 mg/day) and amphotericin B in Table 11.

Table 11: Efficacy Analysis: Treatment Success in Patients with Candidemia and other <i>Candida</i> Infection in Study FG-463-21-08 (Adult Patients)		
	Micafungin sodium 100 mg/day n (%)	Comparator[†] n (%)
Treatment Success at End of IV Therapy¹ % treatment difference (95% CI)²	156/248 (62.9) 2.3 (-5.7, 11.4)	149/246 (60.6)
Success in Patients with Neutropenia at Baseline³	15/32 (46.9)	11/25 (44.0)
Success by Type of Infection		
Candidemia	131/206 (63.6)	129/209 (61.7)
Invasive Candidiasis	24/40 (60.0)	20/37 (54.1)
Success by Site of Infection		
Abscess	4/5 (80)	4/6 (66.7)
Blood	127/203 (62.6)	129/207 (62.3)
Bone	0/0	0/1 (0.0)
Central Line/Catheter Associated	4/4 (100.0)	0/2 (0.0)
Disseminated	5/10 (50.0)	5/10 (50.0)
Endocardium	1/2 (50.0)	3/5 (60.0)
Kidney	0/0	0/1 (0.0)
Peritonitis	13/20 (65.0)	8/14 (57.1)
Success by Organism⁴		
<i>C. albicans</i>	67/103 (65.0)	65/109 (59.6)
<i>C. dubliniensis</i>	1/1 (100.0)	1/1 (100.0)
<i>C. famata</i>	3/3 (100.0)	1/1 (100.0)
<i>C. glabrata</i>	14/30 (46.7)	8/19 (42.1)
<i>C. guilliermondii</i>	4/5 (80.0)	4/5 (80.0)
<i>C. inconspicua</i>	1/1 (100.0)	0/0
<i>C. intermedia</i>	0/0	2/3 (66.7)
<i>C. kefyr</i>	0/0	1/2 (50.0)
<i>C. krusei</i>	4/9 (44.4)	5/10 (50.0)
<i>C. lipolytica</i>	0/0	1/1 (100.0)
<i>C. lusitaniae</i>	1/1 (100.0)	1/2 (50.0)
<i>C. non-albicans</i>	0/0	0/1 (0.0)
<i>C. palmiophila</i>	0/0	0/1 (0.0)
<i>C. parapsilosis</i>	30/42 (71.4)	24/38 (63.2)
<i>C. pelliculosa</i>	0/0	0/4 (0.0)
<i>C. rugosa</i>	0/2 (0.0)	1/1 (100.0)
<i>C. sake</i>	0/0	2/3 (66.7)
<i>C. sp. nos</i>	3/3 (100.0)	4/7 (57.1)
<i>C. tropicalis</i>	39/66 (59.1)	39/62 (62.9)
<i>C. utilis</i>	1/1 (100.0)	0/1 (0.0)
<i>Trichosporon asahii</i>	0/1 (0.0)	0/0
Yeast <i>sp. nos</i>	1/1 (100.0)	3/3 (100.0)
Relapse through end of study		
Overall	57/156 (36.5)	60/149 (40.3)
Culture confirmed relapse	6/156 (3.8)	5/149 (3.4)
Systemic antifungal therapy during post-treatment	18/156 (11.5)	19/149 (12.8)
Died during follow-up ⁵	33/156 (21.2)	36/149 (24.2)
Overall study mortality	106/264 (40.2)	108/267 (40.4)
Mortality during IV therapy ⁶	50/264 (18.9)	53/267 (19.9)
Mortality attributed to fungal infection	34/264 (12.9)	25/267 (9.4)

[†] Amphotericin B 3 mg/day

¹ Success is defined as a positive clinical response (complete or partial) and a positive microbiological response (eradication or presumed eradication) at the end of blinded therapy. Patients that died during treatment (first dose through last dose day +1), missed evaluation, or met one of following 3 criteria of systemic antifungal therapy were considered a failure: i. pre-treatment (within 72 hours of study drug administration) systemic (IV, PO, CIV) antifungal use for therapeutic, non-prophylaxis purposes for > 2 days; ii. any on study treatment systemic antifungal use for > 1 day; or iii. post-treatment systemic antifungal use for therapeutic purposes initiated within 48 hours of discontinuation of study therapy.

² Micafungin 100 –Amphotericin B. 95% CI for difference based on the Cochran-Mantel-Haenszel method controlling for baseline neutropenia status.

³ Neutropenia was assessed by Investigator

⁴ A patient may have had more than one baseline fungal infection species.

⁵ Death day > Last dose day + 1

⁶ IV therapy duration is from first dose day through last dose day + 1

Pediatrics

The response rate at the end of therapy for micafungin sodium treated pediatric patients was 32/48 (67%) compared to 32/50 (64%) for amphotericin B treated patients. In the micafungin treated group, 3 pediatric patients with treatment success had proven recurrent infection. In the amphotericin B group, none of the successfully treated pediatric patients had a recurrent infection.

Treatment of Esophageal Candidiasis

Study demographics and trial design

Table 12: Summary of Patient Demographics for Micafungin Sodium Clinical Studies for Esophageal Candidiasis (Adult Patients Only)				
Dosage, Route of Administration and Duration	Number of Study Subjects	Mean Age (Range)	Gender	Race
Study 03-7-005: Phase III, multicenter, randomized, multinational, double-blind, parallel, non-inferiority trial				
Micafungin sodium: 150 mg/day once daily as an intravenous infusion for a minimum of 14 days or 7 days after resolution of clinical symptoms	260	37.2 ± 10.59 (17.0 - 80.0)	M = 131 (50%) F = 129 (50%)	Black = 176 (68%) Caucasian = 38 (15%) Mestizo = 32 (12%) Other = 14 (5%)
Fluconazole: 200 mg/day once daily as an intravenous infusion for a minimum of 14 days or 7 days after resolution of clinical symptoms	258	37.5 ± 11.16 (17.0 - 87.0)	M = 116 (45%) F = 142 (55%)	Black = 178 (69%) Caucasian = 35 (14%) Mestizo = 29 (11%) Other = 16 (6%)

Table 12: Summary of Patient Demographics for Micafungin Sodium Clinical Studies for Esophageal Candidiasis (Adult Patients Only)				
Dosage, Route of Administration and Duration	Number of Study Subjects	Mean Age (Range)	Gender	Race
Study FG463-21-09: Phase II, multicenter, prospective randomized, reference therapy controlled, double-blind, parallel, group study				
Micafungin sodium: 50 mg/day once daily as an intravenous infusion for 14 days	64	33.9 ± 7.5 (19 - 54)	M = 30 F = 34	Black = 31 (48%) Caucasian = 25 (39%) Other = 8 (13%)
Micafungin sodium: 100 mg/day once daily as an intravenous infusion for 14 days	62	36.8 ± 8.1 (24 - 68)	M = 26 F = 36	Black = 33 (53%) Caucasian = 26 (42%) Other = 3 (5%)
Micafungin sodium: 150 mg/day once daily as an intravenous infusion for 14 days	59	36.7 ± 8.8 (23 - 68)	M = 33 F = 26	Black = 30 (51%) Caucasian = 25 (42%) Other = 4 (7%)
Fluconazole: 200 mg/day once daily as an intravenous infusion for 14 days	60	35.5 ± 8.1 (19 - 56)	M = 28 F = 32	Black = 32 (53%) Caucasian = 22 (37%) Other = 6 (10%)

Study Results

In two controlled studies involving 763 patients with esophageal candidiasis, 445 adults with endoscopically-proven candidiasis received micafungin sodium, and 318 received fluconazole for a median duration of 14 days (range 1-33 days).

Micafungin sodium was evaluated in a phase III, randomized, double-blind study which compared micafungin sodium 150 mg/day (n = 260) to intravenous fluconazole 200 mg/day (n = 258) in adults with endoscopically-proven esophageal candidiasis. Most patients in this study had HIV infection, with CD4 cell counts < 100 cells/mm³.

Outcome was assessed by endoscopy and by clinical response at the end of treatment. Endoscopic cure was defined as endoscopic grade 0, based on a scale of 0-3. Clinical cure was defined as complete resolution in clinical symptoms of esophageal candidiasis (dysphagia, odynophagia, and retrosternal pain). Overall therapeutic cure was defined as both clinical and endoscopic cure. Mycological eradication was determined by culture, and by histological or cytological evaluation of esophageal biopsy or brushings obtained endoscopically at the end of treatment. As shown in Table 13 below, endoscopic cure, clinical cure, overall therapeutic cure, and mycological eradication were comparable for patients in the micafungin sodium and fluconazole treatment groups.

Treatment Outcome*	Micafungin sodium 150 mg/day N=260	Comparator[§] N=258	% Difference[†] (95% CI)
Endoscopic Cure	228 (87.7%)	227 (88.0%)	-0.3% (-5.9, +5.3)
Clinical Cure	239 (91.9%)	237 (91.9%)	0.06% (-4.6, +4.8)
Overall Therapeutic Cure	223 (85.8%)	220 (85.3%)	0.5% (-5.6, +6.6)
Mycological Eradication	141/189 (74.6%)	149/192 (77.6%)	-3.0% (-11.6, +5.6)

*Endoscopic and clinical outcome were measured in modified intent-to-treat population, including all randomized patients who received ≥ 1 dose of study treatment. Mycological outcome was determined in the per protocol (evaluable) population, including patients with confirmed esophageal candidiasis who received at least 10 doses of study drug, and had no major protocol violations.

[§]Fluconazole 200 mg/day

[†]Calculated as micafungin sodium – fluconazole

Most patients (96%) in this study had *Candida albicans* isolated at baseline. The efficacy of micafungin sodium was evaluated in less than 10 patients with *Candida* species other than *C. albicans*, most of which were isolated concurrently with *C. albicans*.

Relapse was assessed at 2 and 4 weeks post-treatment in patients with overall therapeutic cure at end of treatment. Relapse was defined as a recurrence of clinical symptoms or endoscopic lesions (endoscopic grade > 0). There was no statistically significant difference in relapse rates at either 2 weeks or through 4 weeks post-treatment for patients in the micafungin sodium and fluconazole treatment groups, as shown in Table 14.

Relapse	Micafungin sodium 150 mg/day N=223	Comparator[§] N=220	% Difference* (95% CI)
Relapse [†] at Week 2	40 (17.9%)	30 (13.6%)	4.3% (-2.5, 11.1)
Relapse [†] Through Week 4 (cumulative)	73 (32.7%)	62 (28.2%)	4.6% (-4.0, 13.1)

[§]Fluconazole 200 mg/day

*Calculated as micafungin sodium – fluconazole; N=number of patients with overall therapeutic cure (both clinical and endoscopic cure at end-of-treatment)

[†]Relapse included patients who died or were lost to follow-up, and those who received systemic anti-fungal therapy in the post-treatment period

In this study, 459 of 518 (88.6%) patients had oropharyngeal candidiasis in addition to esophageal candidiasis at baseline. At the end of treatment, 192/230 (83.5%) micafungin sodium treated patients and 188/229 (82.1%) of fluconazole treated patients experienced resolution of signs and symptoms of oropharyngeal candidiasis. Of these, 32.3% in the micafungin sodium group, and 18.1% in the fluconazole group (treatment difference = 14.2%; 95% confidence interval [5.6, 22.8]) had symptomatic relapse at 2 weeks post-treatment. Relapse included patients who died or were lost to follow-up, and those who received systemic antifungal therapy during the post-treatment period. Cumulative relapse at 4 weeks post-treatment was 52.1% in the micafungin sodium group and 39.4% in the fluconazole group (treatment difference 12.7%, 95% confidence

interval [2.8, 22.7]).

Effectiveness in pediatric patients was determined based on an open-label clinical study and pharmacokinetic data.

Prophylaxis of *Candida* Infections in Hematopoietic Stem Cell Transplant Recipients

Study demographics and trial design

Table 15: Summary of Patient Demographics for Micafungin sodium Clinical Trial for Prophylaxis of Fungal Infections (Adult and Pediatric Patients)				
Dosage, Route of Administration and Duration	Number of Study Subjects	Mean Age	Gender	Race
Study 98-0-050: Phase III, randomized, double-blind, comparative study				
Once daily IV infusion of micafungin sodium: 50 mg in patients greater than 50 kg and 1 mg/kg in patients weighing 50 kg or less for a maximum of 42 days.	Adults			
	382	47.7 (17.7 - 73.2)	M = 230 (60.2%) F = 152 (39.8%)	Caucasian = 350 (91.6%) Black = 26 (6.8%) Other = 6 (1.6%)
	Pediatrics			
	43	8.7 (0.6 - 16.9)	M = 23 (53.5%) F = 20 (46.5%)	Caucasian = 37 (86.0%) Black = 4 (9.3%) Other = 2 (4.7%)
Fluconazole: Intravenous infusion of 400 mg in patients greater than 50 kg of weight and 8 mg/kg in patients weighing 50 kg or less for a maximum of 42 days.	Adults			
	409	46.4 (17 - 72)	M = 245 (59.9%) F = 164 (40.1%)	Caucasian = 368 (90.0%) Black = 33 (8.1%) Other = 8 (2.0%)
	Pediatrics			
	48	8.2 (0.6 - 16.7)	M = 29 (60.4%) F = 19 (39.6%)	Caucasian = 43 (89.6%) Black = 4 (8.3%) Other = 1 (2.1%)

Study Results

Adults

In a randomized, double-blind study, micafungin sodium (50 mg IV once daily) was compared to fluconazole (400 mg IV once daily) in 882 (791 adults and 91 pediatric) patients undergoing an autologous or syngeneic (46%) or allogeneic (54%) stem cell transplant. The status of the patients' underlying malignancy at the time of randomization was: 365 (41%) patients with active disease, 326 (37%) patients in remission, and 195 (22%) patients in relapse. The more common baseline underlying diseases in the 476 allogeneic transplant recipients were: chronic myelogenous leukemia (22%), acute myelogenous leukemia (21%), acute lymphocytic leukemia (13%), and non-Hodgkin's lymphoma (13%). In the 404 autologous and syngeneic transplant recipients the more common baseline underlying diseases were: multiple myeloma (37.1%), non-Hodgkin's lymphoma (36.4%), and Hodgkin's disease (15.6%). During the study, 198 of 882 (22.4%) transplant recipients had proven graft-versus-host disease; and 475 of 882 (53.9%) recipients received immunosuppressive medications for treatment or prophylaxis of graft-versus-host disease.

Study drug was continued until the patient had neutrophil recovery to an absolute neutrophil count (ANC) of ≥ 500 cells/mm³ or up to a maximum of 42 days after transplant. The average duration of drug administration was 18 days (range 1 to 51 days).

Successful prophylaxis was defined as the absence of a proven, probable, or suspected systemic fungal infection through the end of therapy (usually 18 days), and the absence of a proven or probable systemic fungal infection through the end of the 4-week post-therapy period. A suspected systemic fungal infection was diagnosed in patients with neutropenia (ANC < 500 cells/mm³); persistent or recurrent fever (while ANC < 500 cells/mm³) of no known etiology; and failure to respond to at least 96 hours of broad spectrum antibacterial therapy. A persistent fever was defined as four consecutive days of fever greater than 38°C. A recurrent fever was defined as having at least one day with temperatures $\geq 38.5^\circ\text{C}$ after having at least one prior temperature

1. 38°C; or having two days of temperatures $> 38^\circ\text{C}$ after having at least one prior temperature
2. 38°C. Transplant recipients who died or were lost to follow-up during the study were considered failures of prophylactic therapy.

Successful prophylaxis was documented in 80.7% of recipients who received micafungin sodium, and in 73.7% of recipients who received fluconazole (7.0% difference [95% CI = 1.5, 12.5]), as shown in Table 16, along with other study endpoints. The use of systemic antifungal therapy post-treatment was 42% in both groups.

The number of proven breakthrough *Candida* infections was 4 (0.9%) in the micafungin sodium and 2 (0.4%) in the fluconazole group.

The efficacy of micafungin sodium against infections caused by fungi other than *Candida* has not been established.

Table 16: Results from Clinical Study of Prophylaxis of <i>Candida</i> Infections in Hematopoietic Stem Cell Transplant Recipients (Adult and Pediatric Patients)		
Outcome of Prophylaxis	Micafungin sodium 50 mg/day (n=425)	Comparator[§] (n=457)
	Probable/Proven	Probable/Proven
Success*	343 (80.7%)	337 (73.7%)
Failure:	82 (19.3%)	120 (26.3%)
All Deaths ¹	18 (4.2%)	26 (5.7%)
Proven/probable fungal infection prior to death	1 (0.2%)	3 (0.7%)
Proven/probable fungal infection (not resulting in death) ¹	6 (1.4%)	8 (1.8%)
Suspected fungal infection ²	53 (12.5%)	83 (18.2%)
Lost to follow-up	5 (1.2%)	3 (0.7%)

[§] Fluconazole 400 mg/day

* Defined as the absence of proven, probable or suspected fungal infections through the end of therapy and the absence of a proven or probable infection at the end of the 4-week post-treatment period. Difference (micafungin

sodium – Fluconazole): +7.0% [95% CI=1.5, 12.5]

¹ Through end-of-study (4 weeks post-therapy)

² Through end-of-therapy

Pediatrics

Clinical success was reported in 72% (n=31) pediatric patients treated with micafungin sodium as compared to 54% (N=26) in comparator arms. Treatment failure was reported in 28% (n=12) and 46% (n=22) patients in micafungin sodium and comparator arms respectively. Mortality through the end- of-study (4 weeks post-therapy) was reported in 12% (n=5) and 13% (n=6) patients in micafungin sodium and comparator arms, respectively.

DETAILED PHARMACOLOGY

Human pharmacokinetics of micafungin sodium were determined in healthy subjects, hematopoietic stem cell transplant recipients and patients with esophageal candidiasis up to a maximum of daily dose of 8 mg/kg body weight. The relationship of area under the curve (AUC) to micafungin dose was linear over the daily dose range of 50 mg to 150 mg and 3 mg/kg to 8 mg/kg body weight.

Detailed pharmacokinetics in relevant patient groups and special populations are described previously in the Action and Clinical Pharmacology section.

MICROBIOLOGY

Activity In Vitro

Micafungin exhibited *in vitro* activity against *C. albicans*, *C. glabrata*, *C. guilliermondii*, *C. kefyr*, *C. krusei*, *C. lusitanae*, *C. parapsilosis*, *C. pelliculosa* and *C. tropicalis*.

Susceptibility Testing Methods

The interpretive standards for micafungin against *Candida* species are applicable only to tests performed using Clinical Laboratory and Standards Institute (CLSI) microbroth dilution (MIC; based on partial inhibition endpoint) and CLSI disk diffusion reference method; both MIC and zone diameter results are read at 24 hours.

When available, the clinical microbiology laboratory should provide the results of *in vitro* susceptibility test results for antimicrobial drug products used in resident hospitals to the physician as periodic reports that describe the susceptibility profile of pathogens. These reports should aid the physician in selecting an antifungal drug product for treatment. The techniques for Broth Microdilution and Disk Diffusion are described below.

Broth Microdilution Technique

Quantitative methods are used to determine antifungal MICs. These MICs provide estimates of the susceptibility of *Candida* species to antifungal agents. MICs should be determined using the standardized CLSI procedure. Standardized procedures are based on a microdilution method (broth) with standardized inoculum concentrations and standardized concentrations of micafungin powder. The MIC values should be interpreted according to the criteria provided in Table 17.

Disk Diffusion Technique

Qualitative methods that require measurement of zone diameters also provide reproducible estimates of the susceptibility of *Candida* species to antifungal agents. The CLSI procedure uses

standardized inoculum concentrations and paper disks impregnated with 10 µg of micafungin should be used to test the susceptibility of *Candida* species to micafungin at 24 hours. Disk diffusion interpretive criteria are provided in Table 17.

Table 17: Susceptibility Interpretive Criteria for Micafungin						
	Broth Microdilution MIC (µg/mL) at 24 hours			Disk Diffusion at 24 hours (Zone diameters in mm)		
	Susceptible (S)	Intermediate (I)	Resistant (R)	Susceptible (S)	Intermediate (I)	Resistant (R)
<i>Candida albicans</i>	≤ 0.25	0.5	≥ 1	≥ 22	20-21	≤ 19
<i>Candida tropicalis</i>	≤ 0.25	0.5	≥ 1	≥ 22	20-21	≤ 19
<i>Candida krusei</i>	≤ 0.25	0.5	≥ 1	≥ 22	20-21	≤ 19
<i>Candida parapsilosis</i>	≤ 2	4	≥ 8	≥ 16	14-15	≤ 13
<i>Candida guilliermondii</i>	≤ 2	4	≥ 8	≥ 16	14-15	≤ 13
<i>Candida glabrata</i>	≤ 0.06	0.12	≥ 0.25	Not applicable [†]	Not applicable [†]	Not applicable [†]

MIC: minimum inhibitory concentration

[†] Disk diffusion zone diameters have not been established for this strain/antifungal agent combination.

A report of “Susceptible” indicates that the isolate is likely to be inhibited if the antimicrobial compound in the blood reaches the concentrations usually achievable.

The “Intermediate” category implies that an infection due to the isolate may be appropriately treated in body sites where the drug is physiologically concentrated or when a high dosage of drug is used. The “Resistant” category implies that the isolates are not inhibited by the concentrations of the drug usually achievable with normal dosage schedules and clinical efficacy of the drug against the pathogen has not been reliably shown in treatment studies.

Quality Control

Standardized susceptibility test procedures require the use of quality control organisms to monitor and ensure the accuracy and precision of supplies and reagents used in the assay, and the technique of the individual performing the test. Standard micafungin powder and 10 mcg disks should provide the following range of values noted in Table 18.

Table 18: Acceptable Quality Control Ranges for Micafungin to be Used in Validation of Susceptibility Test Results		
QC strains	Broth microdilution (MIC in mcg/mL) at 24-hour	Disk Diffusion (Zone diameter in mm) at 24-hour
<i>Candida parapsilosis</i> ATCC [†] 22019	0.5 – 2.0	14-23
<i>Candida krusei</i> ATCC 6258	0.12 – 0.5	23-29
<i>Candida tropicalis</i> ATCC 750	Not applicable [§]	24-30
<i>Candida albicans</i> ATCC 90028	Not applicable [§]	24-31

MIC: minimum inhibitory concentration

[†] ATCC is a registered trademark of the American Type Culture Collection.

[§] Quality control ranges have not been established for this strain/antifungal agent combination.

Activity In Animal Models

Micafungin sodium has shown activity in both mucosal and disseminated murine models of candidiasis. Micafungin sodium, administered to immunosuppressed mice in models of

disseminated candidiasis prolonged survival and/or decreased the mycological burden.

TOXICOLOGY

High doses of micafungin sodium (5 to 8 times the highest recommended human dose, based on AUC comparisons) have been associated with irreversible changes to the liver when administered for 3 or 6 months, and these changes may be indicative of pre-malignant processes.

Carcinogenesis, Mutagenesis and Impairment of Fertility

Hepatic carcinomas and adenomas were observed in a 6-month intravenous toxicology study with an 18-month recovery period of micafungin sodium in rats designed to assess the reversibility of hepatocellular lesions.

Rats administered micafungin sodium for 3 months at 32 mg/kg/day (corresponding to 8 times the highest recommended human dose [150 mg/day], based on AUC comparisons), exhibited colored patches/zones, multinucleated hepatocytes and altered hepatocellular foci after 1- or 3-month recovery periods, and adenomas were observed after 20-month recovery period. Rats administered micafungin sodium at the same dose for 6 months exhibited adenomas after a 12-month recovery period; after an 18-month recovery period, an increased incidence of adenomas was observed, and additionally, carcinomas were detected. A lower dose of micafungin sodium (equivalent to 5 times the human AUC) in the 6-month rat study resulted in a lower incidence of adenomas and carcinomas following 18 months recovery. The duration of micafungin dosing in these rat studies (3 or 6 months) exceeds the usual duration of micafungin sodium dosing in patients, which is typically less than 1 month for treatment of esophageal candidiasis, but dosing may exceed 1 month for *Candida* prophylaxis.

Although the increase in carcinomas in the 6-month rat study did not reach statistical significance, the persistence of altered hepatocellular foci subsequent to micafungin dosing, and the presence of adenomas and carcinomas in the recovery periods suggest a causal relationship between micafungin sodium, altered hepatocellular foci, and hepatic neoplasms. Whole-life carcinogenicity studies of micafungin sodium in animals have not been conducted, and it is not known whether the hepatic neoplasms observed in treated rats also occur in other species, or if there is a dose threshold for this effect.

Micafungin sodium was not mutagenic or clastogenic when evaluated in a standard battery of *in vitro* and *in vivo* tests (i.e., bacterial reversion - *S. typhimurium*, *E. coli*; chromosomal aberration; intravenous mouse micronucleus).

Male rats treated intravenously with micafungin sodium for 9 weeks showed vacuolation of the epididymal ductal epithelial cells at or above 10 mg/kg (about 0.6 times the recommended clinical dose for esophageal candidiasis, based on body surface area comparisons). Higher doses (about twice the recommended clinical dose, based on body surface area comparisons) resulted in higher epididymis weights and reduced numbers of sperm cells. In a 39-week intravenous study in dogs, seminiferous tubular atrophy and decreased sperm in the epididymis were observed at 10 and 32 mg/kg, doses equal to about 2 and 7 times the recommended clinical dose, based on body surface area comparisons. There was no impairment of fertility in animal studies with micafungin sodium.

REFERENCES

1. De Wet NT, Bester AJ, Viljoen JJ. *et al.*, A randomized, double blind, comparative trial of micafungin (FK463) vs. fluconazole for the treatment of oesophageal candidiasis. *Aliment Pharmacol Ther* 2005; 21:899–907.
2. De Wet N, Llanos-Cuentas A, Suleiman J. *et al.*, A Randomized, Double-Blind, Parallel- Group, Dose-Response Study of Micafungin Compared with Fluconazole for the Treatment of Esophageal Candidiasis in HIV-Positive Patients. *Clinical Infectious Diseases* 2004; 39:842–9.
3. Kuse ER, Chetchotisakd P, da Cunha CA, *et al.* Micafungin versus liposomal amphotericin B for candidaemia and invasive candidosis: a phase III randomized double-blind trial. *Lancet*. 2007 May 5;369(9572):1519-27.
4. Pappas PG, Rotstein CMF, Betts RF, *et al.*, Micafungin versus Caspofungin for Treatment of Candidemia and Other Forms of Invasive Candidiasis. *Clinical Infectious Diseases* 2007 Oct 1;45(7):883-93.
5. van Burik JA, Ratanatharathorn V, Stepan D, *et al.*, Micafungin versus Fluconazole for Prophylaxis against Invasive Fungal Infections during Neutropenia in Patients Undergoing Hematopoietic Stem Cell Transplantation. *Clinical Infectious Diseases* 2004; 39:1407–16.
6. Clinical and Laboratory Standards Institute (CLSI). Reference Method for Broth Dilution Antifungal Susceptibility Testing of Yeasts – Approved Standard – Third Edition. CLSI document M27-A3. Clinical and Laboratory Standards Institute, Wayne, PA, USA, 2008.
7. Clinical and Laboratory Standards Institute (CLSI). Reference Method for Broth Dilution Antifungal Susceptibility Testing of Yeasts; Fourth Informational Supplement. CLSI document M27-S4. Clinical and Laboratory Standards Institute, Wayne, PA, USA, 2012.
8. Clinical and Laboratory Standards Institute (CLSI). Method for Antifungal Disk Diffusion Susceptibility Testing of Yeasts; Approved Guideline - Second Edition. CLSI document M44-A2. Clinical and Laboratory Standards Institute, Wayne, PA, USA, 2009.
9. ^{Pf}Mycamine[®] (micafungin sodium for injection, 50 mg and 100 mg micafungin sodium per vial), submission control 294524, Product Monograph, Sandoz Canada, Inc. (March 20, 2025)

PART III: CONSUMER INFORMATION

PrMiconazole in 0.9% Sodium Chloride Injection

This leaflet is part III of a three-part "Product Monograph" published when Miconazole in 0.9% Sodium Chloride Injection was approved for sale in Canada and is designed specifically for Consumers. This leaflet is a summary and will not tell you everything about Miconazole in 0.9% Sodium Chloride Injection. Contact your doctor or pharmacist if you have any questions about the drug.

ABOUT THIS MEDICATION

What the medication is used for:

Your doctor has prescribed Miconazole in 0.9% Sodium Chloride Injection to treat one of several types of fungal infections described below.

- Miconazole in 0.9% Sodium Chloride Injection is used to treat certain fungal infections of the esophagus (food tube connecting the mouth to the stomach). These infections are called esophageal candidiasis and are caused by *Candida* (fungus). Healthy individuals usually have *Candida* in their mouth and throat without any ill effects. An infection occurs when the body's resistance is lowered.
- Miconazole in 0.9% Sodium Chloride Injection is used to treat patients with certain fungal infections caused by *Candida* including Candidemia and other *Candida* infections as determined by the doctor.
- Miconazole in 0.9% Sodium Chloride Injection is used to help prevent fungal infections caused by *Candida* in patients who are undergoing a stem cell transplant.

Miconazole in 0.9% Sodium Chloride Injection has not been studied for the treatment of other types of fungal infections.

What it does:

Miconazole in 0.9% Sodium Chloride Injection is an antifungal drug that belongs to a class of drugs called echinocandins. Miconazole in 0.9% Sodium Chloride Injection interferes with the production of a component (glucan polysaccharide) of the fungal cell wall that is necessary if the fungus is to

continue living and growing. Fungal cells exposed to Miconazole in 0.9% Sodium Chloride Injection have incomplete or defective cell walls, making them fragile and unable to grow.

When it should not be used:

Do not use Miconazole in 0.9% Sodium Chloride Injection if you are allergic to it, another echinocandin (eg. caspofungin acetate (Cancidas) or anidulafungin (Eraxis)) or any of the ingredients in Miconazole in 0.9% Sodium Chloride Injection (see *What the non-medicinal ingredients are*).

Use in children:

- Miconazole in 0.9% Sodium Chloride Injection should not be used in patients under 4 months of age.

What the medicinal ingredient is:

Miconazole sodium

What the non-medicinal ingredients are:

Citric acid, sodium chloride, sodium citrate and water for injection

This is a complete listing of all non-medicinal ingredients.

What dosage forms it comes in:

Miconazole in 0.9% Sodium Chloride Injection is supplied as a refrigerated sterile solution in the following presentations:

50 mg / 50 mL: contains 50 mg miconazole (as miconazole sodium)

100 mg / 100 mL: contains 100 mg miconazole (as miconazole sodium)

150 mg / 150 mL: contains 150 mg miconazole (as miconazole sodium)

WARNINGS AND PRECAUTIONS

Serious Warnings and Precautions:

- The use of Miconazole in 0.9% Sodium Chloride Injection may sometimes cause severe allergic reactions including shock (see *Side Effects*).
- Serious liver problems including liver inflammation or worsening of liver failure (see *Side Effects*).
- Miconazole in 0.9% Sodium Chloride Injection may cause kidney problems, kidney failure, and abnormal kidney function tests.

- Micafungin in 0.9% Sodium Chloride Injection may cause destruction of red blood cells called hemolysis or hemolytic anemia.

Use in pregnancy and breast-feeding:

Micafungin in 0.9% Sodium Chloride Injection has not been studied in pregnant women. Micafungin in 0.9% Sodium Chloride Injection should not be used in pregnancy unless the doctor decides the potential benefit justifies the potential risk to the fetus.

It is not known if micafungin sodium is excreted in breast milk. You and your doctor will discuss this.

Use in patients with liver problems:

Patients with liver problems may require extra vigilance by their doctor to monitor liver function. Be sure to tell your doctor if you have had or now have liver problems.

BEFORE you use Micafungin in 0.9% Sodium Chloride Injection talk to your doctor or pharmacist if:

You are taking or plan to take other medications, including those obtained without a prescription.

- You have liver problems
- You are pregnant
- You are breast feeding
- You are allergic to any component of Micafungin in 0.9% Sodium Chloride Injection.

INTERACTIONS WITH THIS MEDICATION

Micafungin in 0.9% Sodium Chloride Injection and other medicines may interact with each other. Tell your healthcare professional about all the medicines you take including prescription and non-prescription medicines, vitamins, and herbal supplements.

Especially, tell your healthcare professional if you take:

- Sirolimus
- nifedipine
- itraconazole
- amphotericin B

The doses of these medicines may need to be

reduced while you are receiving Micafungin in 0.9% Sodium Chloride Injection.

PROPER USE OF THIS MEDICATION

Usual dose (adults and children over 4 months of age):

The treatment schedule and dosage will be set by your doctor, who will monitor your response and condition. Micafungin in 0.9% Sodium Chloride Injection should be administered once daily by slow intravenous infusion of approximately 1 hour.

Overdose

If you think you, or a person you are caring for, have taken too much Micafungin in 0.9% Sodium Chloride Injection, contact a healthcare professional, hospital emergency department, regional poison control centre or Health Canada's toll-free number, 1-844 POISON-X (1-844-764-7669) immediately, even if there are no signs or symptoms.

Serious side effects and what to do about them			
Symptom / effect	Talk to your healthcare professional		Stop taking the drug and get immediate medical help
	Only if severe	In all cases	
Common			
Swollen veins (phlebitis, thrombophlebitis)		✓	
Liver problems (eg. yellowing of the skin and eyes, abdominal pain, nausea, vomiting and fatigue)		✓	
Uncommon			
Serious allergic reaction and symptoms such as severe rash, itching, swelling of hands and feet, trouble breathing			✓

Any medicine may have unintended or undesirable effects, so-called side effects.

Common side effects of Micafungin in 0.9% Sodium Chloride Injection include rash, mental confusion, nausea, vomiting, itching, facial swelling, diarrhea, fever, fatigue and relaxing of blood vessels (vasodilation). Micafungin in 0.9% Sodium Chloride Injection may also cause injection site reactions such as inflammation of the veins.

Other reported medication-related undesirable effects include: anemia, low white blood cells count, abdominal pain, injection site pain, itching, trouble breathing, swelling of the hands, ankles, or feet, impaired liver function, sleep problems, and alterations in some laboratory blood tests. Life-threatening allergic reactions have been reported rarely during administration of Micafungin in 0.9%

Sodium Chloride Injection.

Other side effects may also occur rarely; and, as with any prescription medication, some side effects may be serious. Ask your doctor or pharmacist for more information. Tell your doctor promptly about these or any other unusual symptoms.

This is not a complete list of side effects. For any unexpected effects while taking Micafungin in 0.9% Sodium Chloride Injection contact your doctor.

HOW TO STORE IT

Store in refrigerator at 2°C to 8°C in the original carton. The drug is sensitive to light, keep in original container until ready to use.

The drug product should be stored under prescribed conditions in the refrigerator. In rare cases, if needed, the unopened/sterile product in the original carton may be stored at room temperature of 25 °C up to a maximum of 10 days. Any product stored at room temperature exceeding 10 days should be discarded.

Do not freeze and do not use Micafungin in 0.9% Sodium Chloride Injection if it has been frozen.

Reporting side effects

You can report any suspected side effects associated with the use of health products to Health Canada by:

- Visiting the Web page on Adverse Reaction Reporting (canada.ca/drug-device-reporting) for information on how to report online, by mail or by fax; or
- Calling toll-free at 1-866-234-2345.

NOTE: Contact your healthcare professional if you need information about how to manage your side effects. The Canada Vigilance Program does not provide medical advice.

MORE INFORMATION**If you want more information about
Micafungin in 0.9% Sodium Chloride
Injection:**

- Talk to your healthcare professional
- Find the full product monograph that is prepared for healthcare professionals and includes this Consumer Information by visiting the Health Canada Drug Product Database website ([Drug Product Database: Access the database](#)); the manufacturer's website www.baxter.ca; or by calling 1-888-719-9955.

This leaflet was prepared by Baxter Corporation, Mississauga, Ontario, L5N 0C2, Canada

BAXTER AND GALAXY ARE REGISTERED TRADEMARKS OF BAXTER INTERNATIONAL INC.

Date of Revision: SEP 17, 2025