

PRODUCT MONOGRAPH

MICRO Zn

Zinc Sulfate Injection USP

(Zn²⁺ 1 mg/mL and 5 mg/mL)

Trace Element

Sandoz Canada Inc.
4600 Armand-Frappier St.
Saint-Hubert, QC, Canada
J3Z 1G5

Date of Preparation: December 23, 1991
Date of Revision: SEP 19, 2025

Submission Control No.: 274131

MICRO Zn

Zinc Sulfate Injection USP (Zn²⁺ 1 mg/mL and 5 mg/mL)

Trace Element

CLINICAL PHARMACOLOGY

Zinc is an essential nutritional element that is important in many enzyme systems either as a metalloenzyme or as an enzyme activator. More than 70 different zinc metalloenzymes have been characterized including carbonic anhydrase, alkaline phosphatase, alcohol dehydrogenase, procarboxypeptidase, superoxide dismutase, glyceraldehydes-3-P dehydrogenase and retinene reductase.

A zinc metalloenzyme is also involved in the synthesis of RNA and DNA, making it important in the normal growth and development process. Zinc facilitates wound healing and helps maintain the senses of taste and smell and normal skin hydration.

The distribution of zinc is wide and nonuniform with the highest concentrations found in the eye, prostate, kidney, liver, muscle, bones, teeth (dental enamel), hair, nails and skin. Zinc is also present in the blood with 75-88% of the total zinc of normal human blood in the red cells, 12-22% in the plasma and 3% in the leukocytes. Normal zinc levels are 8.8 mcg/mL in whole blood, 1.21 mcg/mL in plasma, 14.4 mcg/mL in erythrocytes and 1.0 mcg/mL in serum.

In a study with 99 healthy young men, a mean serum zinc concentration of 102 mcg/100 mL (range 68-136) was reported. Thirty to forty percent of plasma zinc is bound to alpha 2-macroglobulin and sixty to seventy percent is loosely bound to albumin.

Profound changes in zinc blood levels are seen in various disease states and under stress conditions. Subnormal plasma zinc levels have been reported in patients with malignant tumours, atherosclerosis, postalcoholic cirrhosis of the liver and other liver diseases, tuberculosis and leprosy, untreated pernicious anemia, chronic and acute infections, and after acute tissue injury, regardless of origin.

Zinc is absorbed primarily from the small intestine. The main route of zinc excretion is in the feces, which contains the total endogenously excreted zinc (pancreatic and intestinal secretions) and zinc not absorbed from the diet. Small amounts of zinc are lost in urine (0.3 to 0.6 mg/day). However, accumulative zincuria has been observed following major operations, severe burns, nephrosis, postalcoholic hepatic cirrhosis, hepatic porphyria and starvation. Zinc is also lost through sweat, in hair and sloughing skin.

In patients with gastrointestinal disease receiving total parental nutrition (TPN), abnormal zinc excretion occurred from the gastrointestinal tract in diarrheal stools and intestinal fluid lost through suction and fistulous discharge.

Zinc deficiency occurs during long term TPN and, in some cases, during short term TPN, particularly in patients with long standing enteropathies. TPN patients with zinc deficiency are characteristically apathetic, depressed, and develop diarrhea, alopecia and a moist eczematous rash in the nasolabial fold, followed by bullous or pustular lesions on other parts of the face, in the groin, and on the hands and feet. These conditions are reversed or relieved by zinc administration. All or some of these zinc deficiency symptoms have been reported in adults, children and premature infants, the most predominant clinical manifestations reported being skin lesions and diarrhea resembling symptoms of acrodermatitis enteropathica.

Plasma zinc levels also declined in premature infants maintained on TPN without supplementation. During the last 10 to 12 weeks of pregnancy, two-thirds of the infant's zinc stores are transferred from the mother. This patient population is at high risk of developing zinc deficiency because they are born with low body stores, need zinc for growth and may be in negative zinc balance up to 60 days after birth.

Therefore, providing zinc during TPN prevents development of the following deficiency symptoms: parakeratosis, hypogenusia, anorexia, dysosmia, geophagia, hypogonadism, growth retardation and hepatosplenomegaly.

INDICATIONS AND CLINICAL USE

Micro Zn (Zinc Sulfate Injection USP) is indicated as a supplement to intravenous solutions given for TPN. Its administration in TPN solutions helps to maintain plasma zinc levels and to prevent depletion of endogenous stores of zinc and subsequent deficiency symptoms.

WARNINGS

Micro Zn is a hypotonic solution which should be administered in admixtures only.

If toxicity symptoms occur due to zinc, discontinue supplementation of TPN solutions immediately.

Do not give undiluted Micro Zn by direct injection into a peripheral vein because of the potential of infusion phlebitis.

PRECAUTIONS

The possibility of zinc retention should be a consideration in patients with renal dysfunction and caution should be exercised since zinc is excreted *via* the kidneys.

ADVERSE REACTIONS

No adverse reactions have been reported for the amount of zinc present in this product. The amount is small and toxicity symptoms are not likely to occur at the suggested dosage level.

OVERDOSAGE

Zinc is a relatively nontoxic compound, but toxicity can occur by oral administration, inhalation and hemodialysis. Ingestion of excess zinc has usually resulted from storage of food or beverages in galvanized containers, which results in diarrhea, vomiting and fever. One report of intoxication following inhalation of zinc oxide fumes causing fever, headache and vomiting has been reported in the literature. In 1972, a case of zinc poisoning was reported in a patient on hemodialysis with zinc-contaminated water. The patient developed nausea, fever and severe anemia.

Infusions of 40 to 80 mg/day of zinc have been used with no apparent ill effects. No adverse effects were reported when a group of 22 patients received a 20 mg infusion before and after surgery. One case of ill effects was reported when a daily 10 mg dose of zinc was infused over one hour for 5 days. The ill effects were tachycardia, hypothermia, profuse sweating and blurred vision.

One death resulted from an overdose of intravenous zinc which was due to a local prescribing error. A 72 year old woman with a high output enterocutaneous fistula inadvertently received a 46 mmol (7.4 g) of zinc sulfate infused over a 60-hour period. Analysis of her serum zinc showed a zinc level of 4184 mcg/100 mL. Clinical manifestations were edema, jaundice, vomiting, diarrhea and oliguria.

Seven patients who received an accidental overdose (25 mg zinc/litre TPN solution; equivalent to 50 to 70 mg zinc/day) exhibited hyperamylasemia (557 to 1850 Klein Units; normal 130 to 310).

For management of a suspected drug overdose, contact your regional Poison Control Centre immediately.

DOSAGE AND ADMINISTRATION

Dosage

The suggested dosage ranges are:

Adults

For the metabolically stable adult receiving TPN, the suggested intravenous dosage level is 2.5 to 4 mg of zinc per day.

For acute catabolic states an additional 2 mg of zinc per day is suggested.

For the stable adult with fluid loss from the small bowel, an additional 12.2 mg of zinc per litre of TPN solution; or an additional 17.1 mg of zinc per kg of stool or ileostomy output is recommended.

Pediatrics

For full-term infants and children up to 5 years of age, 100 mcg zinc/kg/day is recommended.

For premature infants weighing up to 3 kg in body weight, 300 mcg zinc/kg/day is recommended.

Administration

Routine monitoring of zinc plasma levels is suggested as a guideline for administration.

Normal plasma levels for zinc vary from approximately 68 to 136 mcg per 100 mL. Frequently monitor the blood zinc levels for those patients receiving more than the usual maintenance dosage level of zinc.

Micro Zn is for intravenous use after dilution only.

PHARMACEUTICAL INFORMATION

Proper Name:	Zinc Sulfate
Chemical Name:	Zinc Sulfate Heptahydrate
Molecular Formula:	$\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$
Molecular Weight:	287.5
Description:	Zinc sulfate is an odourless, colourless, transparent, efflorescent crystals or white crystalline powder with an astringent metallic taste freely soluble in water. Each gram of zinc sulfate represents 3.50 mmol (7 mEq) of zinc. Zinc sulfate 220 mg is approximately equivalent to 50 mg of zinc.

Dilution for Intravenous Use

Aseptic addition of Micro Zn to the amino acid/dextrose component of a TPN solution under a laminar flow hood is recommended. After dilution, the solution must be used within 24 hours.

Visually inspect parenteral drug products for particulate matter and discoloration prior to administration whenever container and solution permit.

DOSAGE FORMS, COMPOSITION AND PACKAGING

Each mL of Micro Zn contains:

	Micro Zn 1 mg/mL	Micro Zn 5 mg/mL
Zinc (as zinc sulfate heptahydrate)	1 mg	5 mg
Water for injection	q.s.	q.s.

Sulfuric acid is used to adjust the pH.

Micro Zn (Zinc Sulfate Injection USP) is available in two strengths, 1 mg/mL and 5 mg/mL of zinc, in single use vials of 10 mL, boxes of 10.

STORAGE AND STABILITY

Store between 15 and 28°C. Protect from freezing.

The chlorobutyl rubber stopper is not made with natural rubber latex.

REFERENCES

1. Boddapati S, Yang K and Murty R. Intravenous solution compatibility and filter-retention characteristics of trace element preparations. *Am J Hosp Pharm* 1981; 38:1731-1736.
2. Ellenhorn MJ and Barceloux, eds. *Medical Toxicology. Diagnosis and Treatment of Human Poisoning*. Elsevier Science Publishing Co. Inc., New York, 1988; 1064-1065.
3. Expert Panel, AMA Department of Food and Nutrition. Guidelines for essential trace element preparations for parenteral use. *JAMA* 1979; 241: 2051-2054.
4. Phillips GD and Odgers CL. Parenteral Nutrition: Current status and concepts. *Drug* 1982; 23: 276-323.
5. Report of a WHO expert group on trace elements in human nutrition. 3 Zinc Tech. Rep. Ser. *Wed. Hlth. Org. No. 532*, 1973; 9-14.
6. Reynolds JE ed. *Martindale. The Extra Pharmacopeia*. The Pharmaceutical Press, London 1982; 943-946.
7. Tasman-Jones C, Kay RG and Lee SP. Zinc and copper deficiency, with particular reference to parenteral nutrition. *Surg. Annu* 1978, 10: 23-52.
8. Underwood EJ. *Trace Elements in Human and Animal Nutrition*. 4th ed. New Academic Press 1977; 196-242.
9. Wolman SL, Anderson GH, Marliss EB and Jeejeebhoy KN. Zinc in total parenteral nutrition: requirements and metabolic effects. *Gastroenterology* 19.
10. Zlotkin SH. Total parenteral nutrition in children. *Pediatr Clin North Am*, 1985; 32(2):381-400.
11. Brocks A, Reid H, Glazer G. Acute intravenous zinc poisoning. *Br Med J*, 1977; May 28;1(6073):1390-1.