

PRODUCT MONOGRAPH
INCLUDING PATIENT MEDICATION INFORMATION

Pr **TUKYSA**[®]

Tucatinib Tablets

Tablets, 50 mg and 150 mg, Oral

Protein Kinase Inhibitor (XXX)

Pfizer Canada ULC
17,300 Trans-Canada Hwy,
Kirkland, Québec
H9J 2M5

Date of Initial Authorization:
JUN 06, 2020

Date of Revision:
NOV 24, 2025

Submission Control No.: 299882

TUKYSA is a registered trademark of Seagen Inc. in Canada

RECENT MAJOR LABEL CHANGES

Not applicable

TABLE OF CONTENTS

RECENT MAJOR LABEL CHANGES	1
TABLE OF CONTENTS	2
PART I: HEALTH PROFESSIONAL INFORMATION	4
1 INDICATIONS	4
1.1 Pediatrics	4
1.2 Geriatrics	4
2 CONTRAINDICATIONS	4
3 SERIOUS WARNINGS AND PRECAUTIONS BOX	5
4 DOSAGE AND ADMINISTRATION	5
4.1 Dosing Considerations	5
4.2 Recommended Dose and Dosage Adjustment	5
4.3 Missed Dose	8
5 OVERDOSAGE	8
6 DOSAGE FORMS, STRENGTHS, COMPOSITION AND PACKAGING	8
7 WARNINGS AND PRECAUTIONS	8
7.1 Special Populations	11
7.1.1 Pregnant Women	11
7.1.2 Breast-feeding	11
7.1.3 Pediatrics	11
7.1.4 Geriatrics	11
8 ADVERSE REACTIONS	12
8.1 Adverse Reaction Overview	12
8.2 Clinical Trial Adverse Reactions	12
8.3 Less Common Clinical Trial Adverse Reactions	14
8.4 Abnormal Laboratory Findings: Hematologic, Clinical Chemistry and Other Quantitative Data	15
9 DRUG INTERACTIONS	15
9.1 Overview	15
9.2 Drug-Drug Interactions	16
9.3 Drug-Food Interactions	18
9.4 Drug-Herb Interactions	18
9.5 Drug-Laboratory Test Interactions	18
10 ACTION AND CLINICAL PHARMACOLOGY	18
10.1 Mechanism of Action	18
10.2 Pharmacodynamics	18
10.3 Pharmacokinetics	19

11	STORAGE, STABILITY AND DISPOSAL.....	20
12	SPECIAL HANDLING INSTRUCTIONS.....	20
	PART II SCIENTIFIC INFORMATION	21
13	PHARMACEUTICAL INFORMATION.....	21
14	CLINICAL TRIALS.....	22
	14.1 Trial Design and Study Demographics	22
	14.2 Study Results	24
15	MICROBIOLOGY.....	26
16	NON-CLINICAL TOXICOLOGY.....	26
	PATIENT MEDICATION INFORMATION	28

PART I: HEALTH PROFESSIONAL INFORMATION

1 INDICATIONS

TUKYSA (tucatinib) is indicated in combination with trastuzumab and capecitabine for treatment of patients with locally advanced unresectable or metastatic HER2-positive breast cancer, including patients with brain metastases, who have received prior treatment with trastuzumab, pertuzumab, and trastuzumab emtansine, separately or in combination.

Clinical trial data supporting the effectiveness of TUKYSA in combination with trastuzumab and capecitabine are limited to patients who had received at least one prior HER2-directed therapy in the metastatic setting.

1.1 Pediatrics

Pediatrics (<18 years of age):

No data are available to Health Canada; therefore, Health Canada has not authorized an indication for pediatric use.

1.2 Geriatrics

Geriatrics (≥65 years of age):

In the HER2CLIMB trial, 116/612 (19%) patients were 65 years or older. No overall differences in effectiveness were observed between these patients and younger patients. Differences in safety were observed between patients 65 years or older and younger patients in the TUKYSA arm; patients 65 years or older were more likely to experience a serious adverse event and more likely to discontinue TUKYSA compared to younger patients <65 years (see Warnings and Precautions Special Populations, Geriatrics).

2 CONTRAINDICATIONS

TUKYSA is contraindicated in patients who are hypersensitive to this drug or to any ingredient in the formulation, including any non-medicinal ingredient, or component of the container. For a complete listing, see Dosage Forms, Strengths, Composition and Packaging.

Refer to the Product Monographs of capecitabine and trastuzumab for further information on the contraindications of these drugs.

3 SERIOUS WARNINGS AND PRECAUTIONS BOX

Serious Warnings and Precautions

- TUKYSA in combination with trastuzumab and capecitabine can cause severe diarrhea, leading to dehydration, acute kidney injury and death. Antidiarrheal treatment may be indicated (see Warnings and Precautions, Diarrhea).
- TUKYSA can cause severe hepatotoxicity; monitor transaminase and bilirubin (see Warnings and Precautions, Hepatotoxicity).
- TUKYSA may cause fetal harm and developmental malformation when administered to a pregnant woman (see Warnings and Precautions, Sexual Health and Special Populations, Pregnant Women).

4 DOSAGE AND ADMINISTRATION

4.1 Dosing Considerations

TUKYSA tablets should be swallowed whole. Tablets should not be chewed, crushed, or split prior to swallowing.

4.2 Recommended Dose and Dosage Adjustment

The recommended dose of TUKYSA is 300 mg (two 150 mg tablets) taken orally twice daily in combination with trastuzumab and capecitabine until disease progression or unacceptable toxicity (see Table 1). For additional information on dosing and administration of the co-administered trastuzumab and capecitabine, consult the respective Product Monographs.

Table 1: Recommended dosing

Treatment	Dose	Treatment Days	Timing Relative to Food Intake
TUKYSA	300 mg orally twice daily	Continuously	Take with or without a meal
Capecitabine	1000 mg/m ² orally twice daily	Days 1 to 14 every 21 days	Take within 30 minutes after a meal
Trastuzumab <u>Intravenous dosing</u> Initial dose Subsequent doses OR <u>Subcutaneous dosing</u>	8 mg/kg intravenously 6 mg/kg intravenously 600 mg subcutaneously	Day 1 Every 21 days Every 21 days	Not applicable

TUKYSA should be taken approximately 12 hours apart, at the same time every day, with or without a meal. TUKYSA may be taken at the same time with capecitabine.

Patients with Hepatic Impairment

For patients with baseline severe hepatic impairment (Child-Pugh C), reduce the starting dose of TUKYSA to 200 mg orally twice daily (see ACTION AND CLINICAL PHARMACOLOGY, Pharmacokinetics).

No starting dose adjustment is required in patients with baseline mild (Child-Pugh A) or moderate (Child-Pugh B) hepatic impairment.

Patients with Renal Impairment

TUKYSA in combination with capecitabine and trastuzumab cannot be used in patients with severe renal impairment (creatinine clearance [CrCL] < 30 mL/min), because capecitabine is contraindicated in patients with severe renal impairment.

No starting dose adjustment is required in patients with baseline mild to moderate renal impairment (CrCL ≥ 30 mL/min) (see ACTION AND CLINICAL PHARMACOLOGY, Pharmacokinetics).

Concomitant Use with Strong CYP2C8 Inhibitors

Avoid concomitant use of strong CYP2C8 inhibitors with TUKYSA. If concomitant use with a strong CYP2C8 inhibitor cannot be avoided, reduce the TUKYSA starting dose to 100 mg twice daily and increase monitoring for tucatinib-related toxicity. After discontinuation of the strong CYP2C8 inhibitor for 3 elimination half-lives, resume the TUKYSA dose taken prior to initiating the inhibitor (see DRUG INTERACTIONS).

Pediatrics (<18 years of age)

Health Canada has not authorized an indication for pediatric use.

Geriatrics (≥ 65 years old)

No dose adjustment is required in patients ≥ 65 years of age (see ACTION AND CLINICAL PHARMACOLOGY, Pharmacokinetics).

Dose Modifications for Adverse Reactions

The recommended TUKYSA dose modifications for patients with adverse reactions are provided in Tables 2 to 3. Refer to the Product Monographs for co-administered trastuzumab and capecitabine for dose modifications for toxicities suspected to be caused by those therapies.

Table 2: TUKYSA Dose Reduction Schedule

Dose Level	TUKYSA Dose
Starting dose	300 mg twice daily
First dose reduction	250 mg twice daily
Second dose reduction	200 mg twice daily
Third dose reduction	150 mg twice daily ¹

1. Permanently discontinue TUKYSA in patients unable to tolerate 150 mg twice daily.

Table 3: TUKYSA Dose Modifications for Adverse Reactions

Adverse Reaction¹	Severity	TUKYSA Dose Modification
Diarrhea	Grade 3 without anti-diarrheal treatment	Initiate or intensify appropriate medical therapy. Hold TUKYSA until recovery to ≤ Grade 1, then resume TUKYSA at the same dose level.
	Grade 3 with anti-diarrheal treatment	Initiate or intensify appropriate medical therapy. Hold TUKYSA until recovery to ≤ Grade 1, then resume TUKYSA at the next lower dose level.
	Grade 4	Permanently discontinue TUKYSA.
Hepatotoxicity	Grade 2 elevation of bilirubin (>1.5 – ≤ 3 x ULN)	Hold TUKYSA until recovery to ≤ Grade 1, then resume TUKYSA at the same dose level.
	Grade 3 elevation of ALT or AST (> 5 – ≤ 20 x ULN) OR Grade 3 elevation of bilirubin (> 3 ≤ 10 x ULN)	Hold TUKYSA until severity ≤ Grade 1. Then resume TUKYSA at the next lower dose level.
	Grade 4 elevation of ALT or AST (> 20 x ULN) OR Grade 4 elevation of bilirubin (> 10 x ULN)	Permanently discontinue TUKYSA.
	ALT or AST > 3 x ULN AND Bilirubin > 2 x ULN	Permanently discontinue TUKYSA.
Other Adverse Reactions	Grade 3	Hold TUKYSA until severity ≤ Grade 1. Then resume TUKYSA at the next lower dose level.
	Grade 4	Permanently discontinue TUKYSA.

ULN: upper limit of normal; ALT: alanine aminotransferase; AST: aspartate aminotransferase

1. Grading per CTCAE v4.03

4.3 Missed Dose

If a patient misses or vomits a dose, they should take their next dose at the regularly scheduled time.

5 OVERDOSAGE

There is no specific antidote, and the benefit of hemodialysis in the treatment of TUKYSA overdose is unknown. In the event of an overdose, withhold TUKYSA and apply general supportive measures.

For management of a suspected drug overdose, contact your regional poison control centre.

6 DOSAGE FORMS, STRENGTHS, COMPOSITION AND PACKAGING

Table 4: Dosage Forms, Strengths, Composition and Packaging

Route of Administration	Dosage Form / Strength/Composition	Non-medicinal Ingredients
Oral	50 mg tablets: round, convex, yellow, film-coated, debossed with "TUC" on one side and "50" on the other side. 150 mg tablets: oval-shaped, yellow, film-coated, debossed with "TUC" on one side and "150" on the other side.	<u>Tablet core</u> Colloidal silicon dioxide Copovidone Crospovidone Potassium chloride Sodium bicarbonate Sodium chloride Magnesium stearate Microcrystalline cellulose <u>Coating</u> Polyvinyl alcohol Titanium dioxide Macrogol/polyethylene glycol Talc Yellow iron oxide non-irradiated

7 WARNINGS AND PRECAUTIONS

General

Patients with brain metastases requiring immediate local therapy were excluded from the pivotal HER2CLIMB study. These patients should undergo local CNS directed therapy prior to being treated with TUKYSA, if appropriate (see CLINICAL TRIALS).

Driving and Operating Machinery

Due caution should be exercised when driving or operating a vehicle or potentially dangerous

machinery.

Gastrointestinal

Diarrhea

Diarrhea, including severe events, has been reported during treatment with TUKYSA in combination with trastuzumab and capecitabine (see ADVERSE REACTIONS). In HER2CLIMB, 81% of patients who received TUKYSA experienced diarrhea, including 12% with Grade 3 diarrhea and 0.5% with Grade 4 diarrhea. Both patients who developed Grade 4 diarrhea subsequently died, with diarrhea as a contributor to death. The median time to onset of the first episode of diarrhea was 12 days; 80% of diarrhea events resolved with a median time to resolution of 8 days. Diarrhea led to dose reductions of TUKYSA in 6% of patients and discontinuation of TUKYSA in 1% of patients. Prophylactic use of antidiarrheals was not mandated in the HER2CLIMB trial.

If diarrhea occurs, administer antidiarrheals as clinically indicated. Based on the severity of the diarrhea, interrupt dose, then dose reduce or permanently discontinue TUKYSA (see DOSAGE AND ADMINISTRATION) and refer to the trastuzumab and capecitabine Product Monographs for relevant safety information and dose modifications. Perform diagnostic tests as clinically indicated to exclude infectious causes of Grade 3 or 4 diarrhea or diarrhea of any grade with complicating features (dehydration, fever, neutropenia).

Stomatitis

Stomatitis has been reported during treatment with TUKYSA in combination with trastuzumab and capecitabine. In HER2CLIMB, 32% of patients who received TUKYSA experienced stomatitis, including 2.5% with Grade 3.

If stomatitis occurs, follow the dose modification guidelines for TUKYSA in Table 3 (see DOSAGE AND ADMINISTRATION) and refer to the trastuzumab and capecitabine Product Monographs for relevant safety information and dose modifications.

Hepatic/Biliary/Pancreatic

Hepatotoxicity

Hepatotoxicity, including severe hepatotoxicity, has been reported during treatment with TUKYSA. In HER2CLIMB, the median time to onset of any grade increased ALT, AST, or bilirubin was 36 days; 84% of events resolved, with a median time to resolution of 22 days. In HER2CLIMB, 8% of patients who received TUKYSA had an ALT increase $> 5 \times$ ULN, 6% had an AST increase $> 5 \times$ ULN, and 1.5% had a bilirubin increase $> 3 \times$ ULN (Grade ≥ 3); 2% of patients had AST and/or ALT increased $> 10 \times$ ULN. Hepatotoxicity led to dose reduction of TUKYSA in 8% of patients and discontinuation of TUKYSA in 1.5% of patients.

If hepatotoxicity occurs, based on its severity, interrupt dose, then dose reduce or permanently discontinue TUKYSA (see DOSAGE AND ADMINISTRATION).

Patients with known chronic liver disease, or carriers of hepatitis B or C were excluded from the pivotal study. Patients with pre-existing liver function test abnormalities (total bilirubin $> 1.5 \times$ ULN, or AST/ALT $> 2.5 \times$ ULN, or AST/ALT $> 5 \times$ ULN if liver metastases were present) were excluded from clinical trials with TUKYSA.

Monitoring and Laboratory Tests

Liver Function Monitoring

Monitor ALT, AST, and bilirubin prior to starting TUKYSA, every three weeks, and as clinically indicated.

Creatinine Increased

Although not an adverse reaction, increased serum creatinine was observed in 14% of patients treated with TUKYSA due to inhibition of renal tubular transport of creatinine without affecting glomerular function. In clinical studies, increases in serum creatinine (30% mean increase) occurred within the first 21 days of treatment with TUKYSA remained elevated but stable throughout treatment and were reversible upon treatment discontinuation. Alternative markers such as BUN, cystatin C, or calculated GFR (if not based on creatinine), may be considered to determine whether renal function is impaired.

Sexual Health

Reproduction

Pregnancy testing

Verify the pregnancy status of female patients of reproductive potential prior to initiating TUKYSA (see Special Populations).

Contraception

Advise female patients of reproductive potential of potential risk to a fetus. Advise females of reproductive potential to use effective contraception during treatment and for at least 1 week after the last dose (see Special Populations).

Advise male patients with female partners who are pregnant, possibly pregnant, or who could become pregnant to use effective barrier contraception during treatment with TUKYSA and for at least 1 week after the last dose of TUKYSA. Male patients are also advised not to donate or store semen during treatment and at least 1 month after the last dose of TUKYSA.

Fertility

No fertility studies in women or men have been conducted.

Based on findings from animal studies, TUKYSA may impair male and female fertility (see NON-CLINICAL TOXICOLOGY).

Skin

Palmar-plantar erythrodysesthesia syndrome

Palmar-plantar erythrodysesthesia syndrome has been reported during treatment with TUKYSA in combination with trastuzumab and capecitabine. In HER2CLIMB, 63% of patients who received TUKYSA experienced palmar-plantar erythrodysesthesia, including 13% with Grade 3.

If palmar-plantar erythrodysesthesia syndrome occurs, follow the dose modification guidelines for TUKYSA in Table 3 (see DOSAGE AND ADMINISTRATION) and refer to the trastuzumab and capecitabine Product Monographs for relevant safety information and dose modifications.

7.1 Special Populations

7.1.1 Pregnant Women

Pregnancy

TUKYSA may cause fetal harm based on findings from animal studies. There are no data from the use of TUKYSA in pregnant women. TUKYSA should not be used in pregnant women. If the patient becomes pregnant while taking TUKYSA, the patient should be apprised of potential hazard to the fetus.

In animal studies, administration of TUKYSA to pregnant rabbits during organogenesis, at exposures similar to those in patients, resulted in embryo-fetal malformations (see NON-CLINICAL TOXICOLOGY).

7.1.2 Breast-feeding

No data are available regarding the presence of tucatinib or its metabolites in human or animal milk or its effects on the breastfed child or on milk production.

Because many drugs are excreted into human milk and because of the potential for serious adverse reactions in a breastfed child from TUKYSA advise lactating women not to breastfeed while taking TUKYSA and for at least 1 week after the last dose.

7.1.3 Pediatrics

No data are available to Health Canada; therefore, Health Canada has not authorized an indication for pediatric use.

7.1.4 Geriatrics

In the HER2CLIMB trial, 116/612 (19%) patients were 65 years or older. No overall differences in effectiveness were observed between these patients and younger patients. Differences in safety were observed between patients 65 years or older and younger patients in both the TUKYSA and control treatment arms. In patients ≥ 65 years, the incidence of serious adverse events was 34% and 36% in patients receiving TUKYSA and placebo, respectively; in patients < 65 , the incidence of serious adverse events was 24% and 25%, respectively. In the TUKYSA arm, diarrhea and vomiting were the most frequent serious adverse events, which occurred in 9% and 6% of patients ≥ 65 years compared to 3% and 2% of patients < 65 years, respectively. Higher incidence rates for Grade ≥ 3 diarrhea and vomiting were observed in patients ≥ 65 years (17% and 7%, respectively) compared to patients < 65 years (12% and 2%, respectively) in the TUKYSA arm of HER2CLIMB. Higher rates of treatment discontinuation due to adverse events were observed in both treatment arms for patients ≥ 65 years (9.8% in the TUKYSA arm and 9.1% in the control arm) compared to patients < 65 years (4.7% in the TUKYSA arm and 1.8% in the control arm).

8 ADVERSE REACTIONS

8.1 Adverse Reaction Overview

The data described in this section reflect exposure to TUKYSA in combination with trastuzumab and capecitabine from HER2CLIMB, a randomized, double-blind, placebo-controlled, active comparator, global trial in patients with locally advanced unresectable or metastatic HER2-positive breast cancer, who received at least one dose of study drug.

The median duration of exposure to TUKYSA or placebo was 5.8 months (range, <0.1, 35.1) compared to 4.4 months (range, <0.1 to 24.0) on the control arm.

Patients treated with TUKYSA experienced higher incidences of treatment-emergent adverse events (99.3% vs 97.0), grade 3-4 adverse events (53.2% vs 45.7%), and treatment-related adverse events (85% vs 73%) than those treated with placebo.

The most common treatment emergent adverse events in patients who received TUKYSA ($\geq 20\%$) were diarrhea, palmar-plantar erythrodysesthesia, nausea, fatigue, hepatotoxicity, vomiting, stomatitis, decreased appetite, abdominal pain, headache, anemia, and rash.

Serious adverse events occurred in 26% of patients treated with TUKYSA compared to 27% of patients treated with placebo + trastuzumab + capecitabine (control arm). The most common serious adverse events ($\geq 2\%$) in patients treated with TUKYSA were diarrhea (4%), vomiting (2%), nausea (2%), abdominal pain (2%), and seizure (2%).

Fatal adverse events occurred in 2% of patients who received TUKYSA compared to 3% of patients on the control arm. Events leading to death in the TUKYSA arm included sudden death, sepsis, dehydration, and cardiogenic shock.

Adverse events leading to treatment discontinuation of TUKYSA/placebo occurred in 6% of patients on the TUKYSA arm compared to 3% of patients in the control arm; the most common adverse events leading to treatment discontinuation of TUKYSA were diarrhea (1%) and ALT increased (1%). Adverse events leading to dose reduction occurred in 21% of patients treated with TUKYSA compared to 11% of patients in the control arm; the most common adverse events leading to dose reduction of TUKYSA were diarrhea (6%), ALT increased (5%), and AST increased (4%).

8.2 Clinical Trial Adverse Reactions

Because clinical trials are conducted under very specific conditions, the adverse reaction rates observed in the clinical trials may not reflect the rates observed in practice and should not be compared to the rates in the clinical trials of another drug. Adverse reaction information from clinical trials is useful for identifying drug-related adverse events and for approximating rates.

Table 5 summarizes treatment emergent adverse events reported in patients in HER2CLIMB.

Table 5: Treatment Emergent Adverse Events (≥10%) in Patients Who Received TUKYSA and with a Difference Between Arms of ≥ 5% Compared to Placebo in HER2CLIMB (All Grades)

SOC Preferred Term	TUKYSA + Trastuzumab + Capecitabine N = 404			Placebo + Trastuzumab + Capecitabine N = 197		
	All Grade n (%)	Grade 3 n (%)	Grade 4 n (%)	All Grade n (%)	Grade 3 n (%)	Grade 4 n (%)
Blood and lymphatic system disorders						
Anemia ¹	85 (21)	15 (3.7)	0 (0)	25 (13)	5 (2.5)	0 (0)
Gastrointestinal disorders						
Diarrhea	327 (81)	50 (12)	2 (0.5)	105 (53)	17 (9)	0 (0)
Nausea	236 (58)	15 (3.7)	0 (0)	86 (44)	6 (3)	0 (0)
Vomiting	145 (36)	12 (3)	0 (0)	50 (25)	7(3.6)	0 (0)
Stomatitis ²	131 (32)	10 (2.5)	0 (0)	42 (21)	1(0.5)	0 (0)
Hepatobiliary disorders						
Hepatotoxicity ³	169 (42)	37 (9)	1 (0.2)	47 (24)	7 (3.6)	0 (0)
Investigations						
Creatinine Increased ⁴	56 (14)	0 (0)	0 (0)	3 (1.5)	0 (0)	0 (0)
Weight decreased	54 (13)	4 (1)	0 (0)	11 (6)	1 (0.5)	(0)
Metabolism and nutrition disorders						
Decreased appetite	100 (25)	2 (0.5)	0 (0)	39 (20)	0 (0)	0 (0)
Musculoskeletal and connective tissue disorders						
Arthralgia	59 (15)	2 (0.5)	0 (0)	9 (4.6)	1 (0.5)	0 (0)
Nervous System Disorders						
Peripheral Neuropathy ⁵	54 (13)	2 (0.5)	(0)	13 (7)	2 (1)	0 (0)
Respiratory, thoracic and mediastinal disorders						
Epistaxis	47 (12)	0 (0)	0 (0)	9 (5)	0 (0)	0 (0)
Skin and subcutaneous tissue disorders						
Palmar-plantar erythrodysesthesia syndrome	256 (63)	53 (13)	0 (0)	104 (53)	18 (9)	0 (0)
Rash ⁶	82 (20)	3 (0.7)	0 (0)	29 (15)	1 (0.5)	0 (0)

1. Anemia includes anemia, hemoglobin decreased, and normocytic anemia

2. Stomatitis includes stomatitis, oropharyngeal pain, oropharyngeal discomfort mouth ulceration, oral pain, lip ulceration, glossodynia, tongue blistering, lip blister, oral dysaesthesia, tongue ulceration, aphthous ulcer

3. Hepatotoxicity includes hyperbilirubinemia, blood bilirubin increased, bilirubin conjugated increased, alanine aminotransferase increased, transaminases increased, hepatotoxicity, aspartate aminotransferase increased, liver function test increased, liver injury, and hepatocellular injury
4. Due to inhibition of renal tubular transport of creatinine without affecting glomerular function
5. Peripheral neuropathy includes peripheral sensory neuropathy, neuropathy peripheral, peripheral motor neuropathy, and peripheral sensorimotor neuropathy
6. Rash includes rash maculo-papular, rash, dermatitis acneiform, erythema, rash macular, rash papular, rash pustular, rash pruritic, rash erythematous, skin exfoliation, urticaria, dermatitis allergic, palmar erythema, plantar erythema, skin toxicity, and dermatitis

8.3 Less Common Clinical Trial Adverse Reactions

Additional clinically important adverse events, regardless of relationship to TUKYSA, that occurred in <10% of patients treated with TUKYSA in combination with trastuzumab and capecitabine include:

Cardiac disorders: cardiac failure, palpitations

Gastrointestinal disorders: dysphagia, rectal hemorrhage

General disorders and administration site conditions: chest discomfort, influenza-like illness, non-cardiac chest pain, peripheral swelling, pyrexia

Infections and infestations: septic shock

Investigations: glomerular filtration rate decreased, neutrophil count decreased, platelet count decreased

Metabolism and nutrition disorders: dehydration, hyperglycemia, hypernatremia, hypoglycemia

Musculoskeletal and connective tissue disorders: muscular weakness

Nervous system disorders: lethargy, seizure

Psychiatric disorders: depression

Renal and urinary disorders: dysuria, urinary incontinence

Reproductive system and breast disorders: vaginal hemorrhage

Skin and subcutaneous tissue disorders: alopecia, night sweats, skin ulcer

Vascular disorders: hypotension

8.4 Abnormal Laboratory Findings: Hematologic, Clinical Chemistry and Other Quantitative Data

Table 6: Laboratory Abnormalities (≥20%) Worsening from Baseline in Patients Who Received TUKYSA and with a Difference of ≥5% Compared to Placebo in HER2CLIMB

	TUKYSA + Trastuzumab +Capecitabine ¹		Placebo + Trastuzumab +Capecitabine ¹	
	All Grades n (%)	Grades ≥3 n (%)	All Grades n (%)	Grades ≥3 n (%)
Hematology				
Decreased hemoglobin	234 (59)	13 (3.3)	100 (51)	3 (1.5)
Chemistry				
Decreased phosphate	221 (57)	30 (8)	87 (45)	14 (7)
Increased bilirubin	190 (47)	6 (1.5)	60 (30)	6 (3.1)
Increased ALT	185 (46)	31 (8)	54 (27)	1 (0.5)
Increased AST	171 (43)	23 (6)	49 (25)	2 (1)
Decreased magnesium	156 (40)	3 (0.8)	48 (25)	1 (0.5)
Decreased potassium ²	146 (36)	25 (6)	61 (31)	10 (5)
Increased creatinine ³	132 (33)	0 (0)	12 (6)	0 (0)
Decreased sodium ⁴	114 (28)	10 (2.5)	46 (23)	4 (2)
Increased alkaline phosphatase	103 (26)	2 (0.5)	34 (17)	0 (0)

1. The denominator used to calculate the rate varied from 351 to 400 in the TUKYSA arm and 173 to 197 in the control arm based on the number of patients with a baseline value and at least one post-treatment value. Grading was based on NCI-CTCAE v.4.03 for laboratory abnormalities, except for increased creatinine which only includes patients with a creatinine increase based on the upper limit of normal definition for grade 1 events (NCI CTCAE v5.0).
2. Laboratory criteria for Grade 1 is identical to laboratory criteria for Grade 2.
3. Due to inhibition of renal tubular transport of creatinine without affecting glomerular function.
4. There is no definition for Grade 2 in CTCAE v.4.03.

9 DRUG INTERACTIONS

9.1 Overview

Tucatinib is metabolized primarily by CYP2C8 and to a lesser extent via CYP3A (See ACTION AND CLINICAL PHARMACOLOGY, Pharmacokinetics). Concomitant use of TUKYSA with a moderate or strong inducer of CYP2C8 or a strong inducer of CYP3A can decrease tucatinib exposure. Concomitant use of TUKYSA with a moderate or strong inhibitor of CYP2C8 can increase tucatinib exposure. Co-administration of tucatinib with itraconazole, a strong CYP3A inhibitor, did not result in a clinically significant drug interaction.

9.2 Drug-Drug Interactions

The drugs listed in this table are based on either drug interaction studies, or potential interactions due to the expected magnitude and seriousness of the interaction.

Table 7: Drug-Drug Interactions

Common Name	Source of Evidence	Effect	Clinical Comment
Drug Interactions that Affect TUKYSA			
Moderate or Strong CYP2C8 Inducers (e.g. rifampin)	CT, P	Concomitant use with a moderate or strong CYP2C8 inducer decreases tucatinib plasma concentrations (see Table 8), which may reduce TUKYSA efficacy.	Avoid concomitant use of TUKYSA with moderate or strong CYP2C8 inducers.
Strong CYP3A Inducers (e.g. phenytoin, rifampin)	CT	Concomitant use with a strong CYP3A inducer decreases tucatinib plasma concentrations (see Table 8), which may reduce TUKYSA efficacy.	Avoid concomitant use of TUKYSA with strong CYP3A inducers.
Moderate or Strong CYP2C8 Inhibitors (e.g. clopidogrel, gemfibrozil)	CT, P	Concomitant use with a moderate or strong CYP2C8 inhibitor increases tucatinib plasma concentrations (see Table 8), which may increase the risk of TUKYSA toxicity.	Avoid concomitant use of TUKYSA with strong CYP2C8 inhibitors (see DOSAGE AND ADMINISTRATION). Increase monitoring for tucatinib-related toxicity if co-administered with moderate CYP2C8 inhibitors.
TUKYSA Drug Interactions that Affect Other Drugs			
CYP3A Substrates (e.g. midazolam, triazolam)	CT	Concomitant use with CYP3A substrates increases the plasma concentrations of CYP3A substrates (see Table 9), which may lead to increased toxicity of the CYP3A substrates.	Avoid concomitant use of TUKYSA with sensitive CYP3A substrates. If the use of sensitive CYP3A substrates is unavoidable, consider dose modification of CYP3A substrates with narrow therapeutic indices and/or increased monitoring for potential adverse reactions as described in the medication's prescribing information.
P-glycoprotein (P-gp) Substrates (e.g. dabigatran etexilate, digoxin, fexofenadine)	CT	Concomitant use with P-gp substrates increases the plasma concentrations of P-gp substrates (see Table 9), which	P-gp substrates with narrow therapeutic indices, such as digoxin, should be used with caution when

		may lead to increased toxicity of the P-gp substrates.	coadministered with TUKYSA. Refer to the prescribing information of sensitive P-gp substrates for dose adjustment recommendations due to drug interactions.
--	--	--	---

CT = Clinical Trial; P = Predicted

Clinical Studies

Table 8: Effect of Other Drugs on TUKYSA

Concomitant Drug (Dose)	TUKYSA Dose	Ratio (90% CI) of Exposure Measures of Tucatinib With and Without Concomitant drug	
		C _{max}	AUC
<u>Strong CYP3A Inhibitor</u> Itraconazole (200 mg BID)	300 mg single dose	1.32 (1.23, 1.42)	1.34 (1.26, 1.43)
<u>Strong CYP3A/Moderate CYP2C8 Inducer</u> Rifampin (600 mg once daily)		0.63 (0.53, 0.75)	0.52 (0.45, 0.60)
<u>Strong CYP2C8 Inhibitor</u> Gemfibrozil (600 mg BID)		1.62 (1.47, 1.79)	3.04 (2.66, 3.46)

Table 9: Effect of TUKYSA on Other Drugs

Concomitant Drug (Dose)	TUKYSA Dose	Ratio (90% CI) of Exposure Measures of Concomitant Drug With and Without Tucatinib	
		C _{max}	AUC
<u>CYP2C8 Substrate</u> Repaglinide (0.5 mg single dose)	300 mg twice daily	1.69 (1.37, 2.10)	1.69 (1.51, 1.90)
<u>CYP3A Substrate</u> Midazolam (2 mg single dose)		3.01 (2.63, 3.45)	5.74 (5.05, 6.53)
<u>P-gp Substrate</u> Digoxin (0.5 mg single dose)		2.35 (1.90, 2.90)	1.46 (1.29, 1.66)
<u>MATE1/2-K Substrate</u> Metformin ¹ (850 mg single dose)		1.08 (0.95, 1.23)	1.39 (1.25, 1.54)

1. Tucatinib reduced the renal clearance of metformin without any effect on glomerular filtration rate (GFR) as measured by iothexol clearance and serum cystatin C.

Based on drug interaction studies conducted with TUKYSA, no clinically significant drug interactions have been observed when TUKYSA is combined with omeprazole (a proton pump inhibitor) or tolbutamide (a sensitive CYP2C9 substrate).

In Vitro Studies

Tucatinib is a reversible inhibitor of CYP2C8 and CYP3A and a time-dependent inhibitor of CYP3A, but has low potential to inhibit CYP1A2, CYP2B6, CYP2C9, CYP2C19, CYP2D6, and UGT1A1 at clinically relevant concentrations.

Tucatinib is a substrate of P-gp and BCRP, but not a substrate of OAT1, OAT3, OCT1, OCT2, OATP1B1, OATP1B3, MATE1, MATE2-K, and BSEP.

Tucatinib inhibits MATE1/MATE2-K mediated transport of metformin and OCT2/MATE1-mediated transport of creatinine. The observed serum creatinine increase in clinical studies with tucatinib is due to inhibition of tubular secretion of creatinine via OCT2 and MATE1.

9.3 Drug-Food Interactions

The effect of food on the pharmacokinetics of tucatinib was not clinically meaningful, thus tucatinib may be administered without regard to food (see ACTION AND CLINICAL PHARMACOLOGY, Pharmacokinetics).

9.4 Drug-Herb Interactions

Interactions with herbal products have not been established.

9.5 Drug-Laboratory Test Interactions

Interactions with laboratory tests have not been established.

10 ACTION AND CLINICAL PHARMACOLOGY

10.1 Mechanism of Action

Tucatinib is a reversible and selective tyrosine kinase inhibitor of HER2. In cellular signaling assays, tucatinib is >1000-fold more selective for HER2 compared to epidermal growth factor receptor. In vitro, tucatinib inhibits phosphorylation of HER2 and HER3, resulting in inhibition of downstream cell signaling and cell proliferation, and induces death in HER2 driven tumor cells. In vivo, tucatinib inhibits the growth of HER2 driven tumors and the combination of tucatinib and trastuzumab showed enhanced anti-tumor activity in vitro and in vivo compared to either drug alone. In an intracranial mouse tumor model, tucatinib demonstrated increased distribution to tumor tissue compared with brain parenchyma and resulted in increased survival.

10.2 Pharmacodynamics

Cardiac Electrophysiology

In a randomised, partially double-blind, placebo- and positive-controlled 3-way crossover ECG assessment study in healthy subjects (N=50), tucatinib, administered as 300 mg twice-daily for four days and then as a single 300 mg dose on the morning of the fifth day, did not have any clinically relevant effect on the QTcF interval, the QRS duration, the PR interval, or ventricular heart rate. In this study, the geometric mean (CV%) C_{max} value of tucatinib was 519 ng/mL (26.7%).

10.3 Pharmacokinetics

Table 10 presents population pharmacokinetic model-predicted pharmacokinetic parameters in patients with metastatic breast cancer after treatment with TUKYSA in combination with trastuzumab and capecitabine.

Table 10: Summary of Tucatinib Pharmacokinetic Parameters in Metastatic Breast Cancer Patients

	C_{max} (ng/mL)	T_{max} (h)	Effective t_{1/2} (h)	AUC₀₋₁₂ (ng*h/mL)	CL/F (L/h)	Vss/F (L)
Steady State (300 mg BID)	630	2	9.55	5234	57.3	730

C_{max}: maximum serum concentration; T_{max}: time present at maximum concentration; t_{1/2}: half life; AUC: area under the curve; CL/F: apparent clearance; Vss/F: apparent steady-state volume of distribution.

Plasma tucatinib exposure (AUC_{0-INF} and C_{max}) demonstrated approximately dose proportional increases at oral doses from 50 to 300 mg twice daily (0.17 to 1 times the recommended dose) administered alone in healthy volunteers. Tucatinib exhibited 1.7-fold accumulation for AUC and 1.5-fold accumulation for C_{max} following administration of TUKYSA 300 mg twice daily for 14 days in patients. Time to steady state was approximately 4 days.

Absorption: Following a single oral dose of 300 mg tucatinib, the median time to peak plasma concentration was approximately 2.0 hours (range 1.0 to 4.0 hours).

Effects of Food

Administration of a single dose of tucatinib (2 x 150 mg) in 11 subjects after a high-fat, high-calorie meal (153 calories from protein, 257 calories from carbohydrates and 571 calories from fat), resulted in a 49% increase in AUC_{0-INF} and a delay in T_{max} from 1.5 hours to 4.0 hours when compared with administration of tucatinib (2 x 150 mg) under fasting conditions. Food did not significantly affect C_{max}. The effect of food on the pharmacokinetics of tucatinib was not clinically meaningful.

Distribution: The apparent volume of distribution of tucatinib was approximately 1670 L in healthy volunteers. The plasma protein binding was 97.1% at clinically relevant concentrations.

Metabolism: Tucatinib is metabolized primarily by CYP2C8 and to a lesser extent via CYP3A. In plasma, approximately 75.6% of the plasma radioactivity was unchanged tucatinib, 19.3% was attributed to identified metabolites including 9.2% as an active metabolite ONT-993, and 5.1% was unassigned.

Elimination: Following a single oral dose of 300 mg, tucatinib is cleared from plasma with a mean half-life of approximately 8.7 hours and apparent clearance of 148 L/h in healthy volunteers.

Tucatinib is predominantly eliminated by the hepatobiliary route and is not appreciably renally eliminated. Following a single oral dose of 300 mg radiolabeled tucatinib, approximately 86% of the total radiolabeled dose was recovered in feces (16% of the administered dose as unchanged tucatinib) and 4% in urine with an overall total recovery of 90% within 13 days post-

dose.

Special Populations and Conditions

Based on population pharmacokinetic analyses, age (19 to 77 years), albumin (2.5 to 5.2 g/dL), body weight (41 kg to 138 kg), and race (White, Black, and Asian) did not have a clinically meaningful effect on tucatinib exposure. The effect of gender on the pharmacokinetics of tucatinib has not been studied.

Pediatrics: Pharmacokinetics of tucatinib have not been evaluated in children and adolescents <18 years of age.

Hepatic Insufficiency: The effect of hepatic impairment on tucatinib pharmacokinetics was studied in non-cancer subjects with mild (Child-Pugh Class A), moderate (Child-Pugh Class B), or severe (Child-Pugh C) hepatic impairment following a single oral administration of 300 mg tucatinib. Mild (Child-Pugh A) and moderate (Child-Pugh B) hepatic impairment had no clinically relevant effect on tucatinib exposure. The mean AUC_{0-INF} of tucatinib was approximately 1.6-fold higher in subjects with severe hepatic impairment compared to subjects with normal hepatic function.

Renal Insufficiency: Renal excretion is a minor route of tucatinib elimination, accounting for approximately 4% of the administered dose. The pharmacokinetics of tucatinib have not been evaluated in a dedicated renal impairment study. In a population pharmacokinetic analysis, mild renal impairment (CrCL 60 to 89 mL/min) had no effect on tucatinib pharmacokinetics. There was insufficient data to evaluate the effect of moderate (CrCL 30 to 59 mL/min) and severe renal impairment (CrCL < 30 mL/min) on tucatinib pharmacokinetics in the population pharmacokinetic analysis.

11 STORAGE, STABILITY AND DISPOSAL

Store at 20°C to 25°C.

Keep out of reach and sight of children.

60 count bottle presentation for 50 mg and 150 mg tablets:

White, 75cc Wide Mouth Round Bottle, made of polyethylene with child-resistant cap, polypropylene closure, and foil induction inner seal. A 2g desiccant canister with silica gel is enclosed with the tablets in each bottle.

120 count bottle presentation for 150 mg tablets:

White, 150cc Wide Mouth Round Bottle, made of polyethylene with child-resistant cap, polypropylene closure, and foil induction inner seal. A 2g desiccant canister with silica gel is enclosed with the tablets in each bottle.

Dispense only in original container. Do not discard desiccant. Replace cap securely each time after opening. Discard any unused tablets 3 months after opening the bottle.


12 SPECIAL HANDLING INSTRUCTIONS

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

PART II SCIENTIFIC INFORMATION

13 PHARMACEUTICAL INFORMATION

Drug Substance

Proper/common name:	Tucatinib
Chemical name:	Ethanol, compd. With N6-(4,5-dihydro-4,4-dimethyl-2-oxazolyl)-N4-[3-methyl-4-([1,2,4]triazolo[1,5-a] 21yridine-7-yloxy)phenyl]-4,6-quinazolinediamine (1:2)
Molecular formula and molecular mass:	$C_{26}H_{24}N_8O_2 - \frac{1}{2} C_2H_5OH$; 503.57 g/mol
Structural formula:	

Physicochemical properties: Tucatinib hemi-ethanolate is an off-white to yellow powder. In aqueous media, tucatinib hemi-ethanolate is soluble at pH <2.9, sparingly soluble at pH 4.0, and insoluble at pH ≥4.6.

14 CLINICAL TRIALS

14.1 Trial Design and Study Demographics

Table 11: Summary of trial design of HER2CLIMB, tucatinib versus Placebo in Combination with Trastuzumab and Capecitabine in Patients with Previously Treated Locally Advanced Unresectable or Metastatic HER2+ Breast Carcinoma

Trial design	Dosage, route of administration and duration	Study subjects (n)	Median age (Range)
Randomized, double-blind, placebo-controlled, active comparator Subjects randomized in a 2:1 ratio to receive tucatinib or placebo in combination with trastuzumab and capecitabine	TUKYSA or placebo, 300 mg orally twice per day, was administered until disease progression or unacceptable toxicity. Trastuzumab was administered intravenously as a loading dose of 8 mg/kg on Day 1 of Cycle 1, followed by a maintenance dose of 6 mg/kg on Day 1 of each subsequent 21-day cycle. An alternate dosing option for trastuzumab was a fixed dose of 600 mg administered subcutaneously on Day 1 of each 21-day cycle. Capecitabine, 1000 mg/m ² orally twice per day, was administered on Days 1 through 14 of each 21-day cycle.	TUKYSA in combination with trastuzumab and capecitabine (N=410) Placebo in combination with trastuzumab and capecitabine (N=202). Total N=612 Randomization was stratified by the presence or history of brain metastases (yes vs. no), Eastern Cooperative Oncology Group (ECOG) performance status (0 vs. 1), and region (U.S., Canada, or rest of world).	TUKYSA + Trastuzumab + Capecitabine 55 (22-80) Placebo + Trastuzumab + Capecitabine 54 (25-82)

The efficacy of TUKYSA in combination with trastuzumab and capecitabine was evaluated in the global HER2CLIMB trial. Enrolled patients were required to have HER2-positive unresectable locally advanced or metastatic breast cancer, with or without brain metastases, and had prior treatment with trastuzumab, pertuzumab, and ado-trastuzumab emtansine (T-DM1) separately or in combination, in the neoadjuvant, adjuvant or metastatic setting. All patients had received at least one prior HER2-directed agent in the metastatic setting, with 94% of patients having prior trastuzumab, 91% of patients with prior pertuzumab, and 99% of patients with prior T-DM1 in the metastatic setting. HER2 overexpression or amplification was confirmed by central laboratory analysis prior to patient enrollment.

Patients with brain metastases, including those with untreated or progressing lesions, were eligible to enroll provided they were neurologically stable and did not require immediate radiation or surgery. Patients who received systemic corticosteroids for control of symptoms of CNS metastases <28 days prior to the first dose of study treatment were excluded from the trial. The trial also excluded patients with leptomeningeal disease.

Table 12: Summary of patient demographics and baseline disease characteristics in HER2CLIMB (Full Analysis Set)

	TUKYSA + trastuzumab + capecitabine (N=410)	Placebo + trastuzumab + capecitabine (N=202)	Total (N=612)
Median Age, years (range)	55 (22, 80)	54 (25, 82)	54 (22, 82)
Age Category, n (%)			
< 65 years	328 (80)	168 (83)	496 (81)
≥ 65 years	82 (20)	34 (17)	116 (19)
Gender n (%)			
Male	3 (0.7)	2 (1)	5 (0.8)
Female	407 (99)	200 (99)	607 (99)
Race			
Asian	18 (4)	5 (2)	23 (4)
Black	41 (10)	14 (7)	55 (9)
White	287 (70)	157 (78)	444 (73)
Baseline ECOG, n (%)			
0	204 (50)	94 (47)	298 (49)
1	206 (50)	108 (53)	314 (51)
Estrogen/progesterone receptor status, n (%)			
Positive for either or both	243 (59)	127 (63)	370 (60)
Negative for both	161 (39)	75 (37)	236 (39)
Sites of metastatic disease			
Lung	200 (49)	100 (50)	300 (49)
Liver	137 (33)	78 (39)	215 (35)
Skin	58 (14)	28 (14)	86 (14)
Patients with history or presence of brain metastases, n (%)	198 (48)	93 (46)	291 (48)
Treated stable	80/198 (40)	37/93 (40)	117/291 (40)
Treated progressing	74/198 (37)	34/93 (37)	108/291 (37)
Untreated	44/198 (22)	22/93 (24)	66/291 (23)
Median number of prior lines of systemic therapy (range)	4 (2, 14)	4 (2, 17)	4 (2, 17)
Median number of prior lines of systemic therapy in the metastatic setting (range)	3 (1, 14)	3 (1, 13)	3 (1,14)

The primary efficacy endpoint was progression-free survival (PFS) in the first 480 randomized patients, as assessed by blinded independent central review (BICR) according to Response Evaluation Criteria in Solid Tumours (RECIST) v1.1.

Secondary efficacy endpoints were evaluated in all randomized patients (N=612) and included overall survival (OS), PFS among patients with a history or presence of brain metastases (PFS_{BrainMets}), and confirmed objective response rate (ORR).

14.2 Study Results

The median duration of exposure to TUKYSA was 7.3 months (range <0.1, 35.1) for patients on the TUKYSA + trastuzumab and capecitabine arm compared to 4.4 months (range <0.1, 24.0) of placebo for patients on the placebo + trastuzumab and capecitabine arm.

Efficacy results are summarized in Table 13 and Figures 1 and 2.

Table 13: Summary of primary and key secondary efficacy results based on RECIST 1.1 criteria in HER2CLIMB, per Blinded Independent Central Review (BICR)

	TUKYSA + Trastuzumab + Capecitabine	Placebo + Trastuzumab + Capecitabine
Primary endpoint¹		
Progression-free survival	N=320	N=160
Number of events (%)	178 (56)	97 (61)
Hazard ratio (95% CI) ²	0.54 (0.42, 0.70)	
P-value ³	<0.00001	
Median PFS, months (95% CI)	7.8 (7.5, 9.6)	5.6 (4.2, 7.1)
Secondary endpoints		
Overall survival	N=410	N=202
Number of deaths	130 (32)	85 (42)
Hazard ratio (95% CI) ²	0.66 (0.50, 0.87)	
P-value ⁴	0.00480	
Median OS, months (95% CI)	21.9 (18.3, 31.0)	17.4 (13.6, 19.9)
PFS_{BrainMets}⁵	N=198	N=93
Number of events (%)	106 (53.5)	51 (54.8)
Hazard ratio (95% CI) ⁶	0.483 (0.339, 0.689)	
P-value ⁶⁷	<0.00001	
Median, months (95% CI)	7.6 (6.2, 9.5)	5.4 (4.1, 5.7)
Confirmed Objective Response Rate for Patients with Measurable Disease	N=340	N=171
ORR (95% CI) ⁸	40.6 (35.3, 46.0)	22.8 (16.7, 29.8)
P-value ⁹	0.00008	

CR	3 (0.9)	2 (1.2)
PR	135 (39.7)	37 (21.6)

CI=confidence interval; PFS=progression-free survival; OS=overall survival; ORR=objective response rate; CR=complete response; PR=partial response.

1. Primary PFS analysis conducted in first 480 randomized patients.
2. Hazard ratio and 95% confidence intervals are based on stratified Cox proportional hazards regression model controlling for stratification factors (presence or history of brain metastases, ECOG status, and region of world)
3. Two-sided p-value based on re-randomization procedure (Rosenberger and Lachin 2002) controlling for stratification factors, compared with the allocated alpha of 0.05
4. Two-sided p-value based on re-randomization procedure (Rosenberger and Lachin 2002) controlling for stratification factors, compared with the allocated alpha of 0.0074 for this interim analysis (with 60% of the planned number of events for final analysis)
5. Analysis includes patients with history or presence of parenchymal brain metastases at baseline, including target and non-target lesions. Any site of disease progression (not only in the brain) was considered an event. Does not include patients with dural lesions only.
6. Hazard ratio and 95% confidence intervals are based on stratified Cox proportional hazards regression model controlling for stratification factors (ECOG status and region of world)
7. Two-sided p-value based on re-randomization procedure (Rosenberger and Lachin 2002) controlling for stratification factors, compared with the allocated alpha of 0.0080 for this interim analysis (with 71% of the planned number of events for final analysis)
8. Two-sided 95% exact confidence interval, computed using the Clopper-Pearson method (1934)
9. Two-sided p-value based on Cochran-Mantel-Haenszel test controlling for stratification factors (presence or history of brain metastases, ECOG status, and region of world), compared with the allocated alpha of 0.05

Figure 1: PFS per BICR





Figure 2: Overall Survival



No. at Risk	0	3	6	9	12	15	18	21	24	27	30	33	36
TUC+Tras+Cape 410	410	388	322	245	178	123	80	51	34	20	10	4	0
Pbo+Tras+Cape 202	202	191	160	119	77	48	32	19	7	5	2	1	0

Efficacy results were consistent across patient subgroups including stratification factors and hormone receptor status.

15 MICROBIOLOGY

Not applicable.

16 NON-CLINICAL TOXICOLOGY

General Toxicology

Rat

Tucatinib was administered orally to rats at doses of 6, 20, and 60 mg/kg/day for 13 weeks. At ≥ 6 mg/kg/day in females, there were higher alanine aminotransferase and alkaline phosphatase, centrilobular hepatocyte hypertrophy, increased interstitial cells in the ovary, atrophy of the uterus, mucification of the vagina, and lobular atrophy of the mammary gland in males. Additional changes at ≥ 20 mg/kg/day included minimally higher liver weights. Additional changes at 60 mg/kg/day included lower uterus/cervix (females) and prostate gland (males) weights. There were no test article-related findings at the end of recovery with the exception of the ovary of females administered ≥ 20 mg/kg/day. The exposure achieved at a dose of 6 mg/kg/day is approximately 2% of the exposure achieved at the maximum recommended human dose of 300 mg twice daily based on AUC comparison.

Cynomolgus monkeys

Tucatinib was administered orally to cynomolgus monkeys at doses of 5, 20, and 40 mg/kg/day for 13 weeks. At ≥ 5 mg/kg/day, there were fecal abnormalities and increased heart rates (at vital signs measurement only). Additional findings at 40 mg/kg/day, included transient body weight loss, lean body condition, and apparent blood in the stool which was successfully treated with antibiotic therapy, and increased liver weights. All changes were reversible. The dose of 20 mg/kg/day is approximately 24% of the maximum recommended human dose of 300 mg twice daily based on AUC comparison.

Carcinogenicity and Mutagenicity

Carcinogenicity studies have not been conducted with tucatinib.

Tucatinib was not mutagenic in an *in vitro* bacterial reverse-mutation study (Ames test) or clastogenic in either an *in vitro* mouse bone marrow chromosomal aberration assay or an *in vivo* mouse bone marrow erythrocyte micronucleus assay.

Reproductive and Developmental Toxicology

Impairment of Fertility

In repeat-dose toxicity studies in rats, decreased corpora lutea/corpus luteum cyst, increased interstitial cells of the ovary, atrophy of the uterus, and mucification of the vagina were observed at doses of ≥ 6 mg/kg/day administered twice daily, which resulted in exposures based on AUC₀₋₁₂ of approximately 4% of the human exposure at the recommended dose. Changes in male rats included lobular atrophy of the male mammary gland and decreased prostate gland weights. No histological effects were observed on male or female reproductive tracts in cynomolgus monkeys or on male reproductive tracts in rats at doses resulting in exposures up to 8 times (in monkey) or 18 times (in rat) the human exposure at the recommended dose based on AUC₀₋₁₂.

Developmental Toxicity

Embryo-fetal development studies were conducted in rabbits and rats. Tucatinib administered to pregnant dams caused teratogenicity in rabbits and embryo-fetal toxicity in rats. In pregnant rabbits, increased resorptions, decreased percentages of live fetuses, and skeletal, visceral, and external malformations were observed in fetuses at ≥ 90 mg/kg/day; at this dose, maternal exposure is approximately equivalent to the human exposure at the recommended dose based on AUC. In pregnant rats, decreased maternal body weight and body weight gain were observed at doses of ≥ 90 mg/kg/day. Fetal effects of decreased weights and delayed ossification were observed at ≥ 120 mg/kg/day.

Juvenile Animal Studies

No juvenile toxicity studies have been conducted.

PATIENT MEDICATION INFORMATION

READ THIS FOR SAFE AND EFFECTIVE USE OF YOUR MEDICINE

Pr **TUKYSA**[®] tucatinib tablets

Read this carefully before you start taking **TUKYSA** and each time you get a refill. This leaflet is a summary and will not tell you everything about this drug. Talk to your healthcare professional about your medical condition and treatment and ask if there is any new information about **TUKYSA**.

Your breast cancer will be treated with **TUKYSA** in combination with other medications (trastuzumab and capecitabine). Read the Consumer Information leaflets for those medications as well as this one.

Serious Warnings and Precautions

TUKYSA can cause severe side effects, including:

Severe diarrhea. This is when you have frequent loose or liquid bowel movements. It can cause you to become dehydrated and have severe kidney problems that could be life threatening. Your healthcare professional may give you medications to help treat diarrhea. Tell your healthcare professional right away at the first sign of diarrhea.

Hepatotoxicity. These are severe liver problems. Your healthcare professional will do blood tests to check if your liver is working properly.

Potential harm to your unborn baby. If you are a female patient who is pregnant or plans to become pregnant, be aware that TUKYSA may harm your unborn baby. Females should avoid becoming pregnant while taking TUKYSA. Use effective birth control methods while taking TUKYSA and for at least 1 week after your last dose. Your healthcare professional can tell you about ways to prevent pregnancy while you are taking TUKYSA. If you become pregnant, or think you might be pregnant while taking TUKYSA, tell your healthcare professional right away.

What is TUKYSA used for?

TUKYSA is used with the medications trastuzumab and capecitabine. It is used to treat adults with breast cancer that:

- is positive for human epidermal growth factor receptor 2 (HER2 positive),
- cannot be removed by surgery,
- has spread outside the breast to other parts of the body such as the brain. This is called locally advanced or metastatic disease, and
- has been treated previously with the medications trastuzumab, pertuzumab, and trastuzumab emtansine.

How does TUKYSA work?

Tucatinib, the active ingredient in TUKYSA, belongs to a family of medications called kinase inhibitors. These medications work by blocking the HER2 receptors on the cancer cells. This may slow or stop cancer cells from growing or may kill them.

What are the ingredients in TUKYSA?

Medicinal ingredients: tucatinib

Non-medicinal ingredients: Colloidal silicon dioxide, copovidone, crospovidone, macrogol/polyethylene glycol, magnesium stearate, microcrystalline cellulose, polyvinyl alcohol, potassium chloride, sodium bicarbonate, sodium chloride, talc, titanium dioxide, yellow iron oxide non-irradiated.

TUKYSA comes in the following dosage forms:

Tablets: 50 mg and 150 mg

Do not use TUKYSA if:

- you are allergic to tucatinib or any of the other ingredients in this medication.

Remember to read the Consumer Information leaflets for capecitabine and trastuzumab. These will help you to determine if there are other reasons why you should not be treated with this combination of medications.

To help avoid side effects and ensure proper use, talk to your healthcare professional before you take TUKYSA. Talk about any health conditions or problems you may have, including if you:

- have liver problems.
- have brain metastases. This is when your cancer has spread to your brain. Some brain metastases may need other treatment before TUKYSA can be started to ensure proper use.

Other warnings you should know about:**Female patients:**

- If you are able to get pregnant, your doctor will do a pregnancy test before you start taking TUKYSA.
- If you are breastfeeding or planning to breastfeed: it is not known if TUKYSA passes into breast milk. You should not breastfeed while you are taking TUKYSA and for at least 1 week after your last dose. Talk to your healthcare professional about the best way to feed your baby during this time.

Male patients with female sexual partners who are pregnant, might be pregnant or could become pregnant:

- Use condoms each time you have sex while taking TUKYSA and for at least 1 week after your last dose.
- Avoid fathering a child during your treatment. If your partner becomes pregnant, tell your healthcare professional right away.
- It is not known if TUKYSA is present in semen.
- Do not donate or store semen during your treatment and for at least 1 month after your last dose of TUKYSA.

Fertility: Taking TUKYSA may affect your fertility. This means that it may be difficult for you to have a child. Talk to your healthcare professional if you have questions about this.

Palmar-Plantar Erythrodysesthesia (Hand-Foot Syndrome): Some patients taking TUKYSA have experienced hand-foot syndrome. Talk to your healthcare professional if you feel any tingling, tenderness, redness or swelling on the palm of the hand or the sole of the foot. These are signs of hand-foot syndrome.

Stomatitis: Some patients taking TUKYSA have experienced stomatitis, which is a condition that affects the inside of your mouth. Talk to your healthcare professional if you develop sores in your mouth or redness and swelling of the lining of your mouth. These are signs of stomatitis.

Blood tests:

During your treatment with TUKYSA, you will need to have blood tests. These blood tests will help your healthcare professional to know how well your liver and kidneys are working. Depending on the test results, your healthcare professional may need to lower your dose or stop your treatment with TUKYSA.

Driving and using machines: Before you do tasks that require special attention, wait until you know how you respond to TUKYSA.

Tell your healthcare professional about all the medicines you take, including any drugs, vitamins, minerals, natural supplements or alternative medicines.

The following may interact with TUKYSA:

- A medicine used to treat bacterial and fungal infections called rifampicin
- A medicine used to treat seizures called phenytoin
- Medicines used to treat high blood pressure or heart problems such as digoxin, clopidogrel
- A medicine used to treat abnormal levels of fat in the blood called gemfibrozil
- Medicines used to treat alcohol dependence, seizures, anxiety disorders, panic, agitation, and insomnia such as midazolam, triazolam
- Medicines used to treat blood clots in your body such as dabigatran etexilate
- A medicine used to prevent and treat allergy symptoms called fexofenadine

How to take TUKYSA:

- exactly as your healthcare professional tells you.
- twice a day by mouth, with or without food. Take your doses about 12 hours apart at about the same times each day.
- swallow the tablets whole. Do not chew, crush or split the tablets before swallowing them.
- You may take your TUKYSA tablets at the same time as your capecitabine.

You will take TUKYSA with the medications trastuzumab and capecitabine. Your healthcare professional will tell you how much of these medications you will take and how you will receive them.

Usual dose: 300 mg (two 150 mg tablets) twice a day

If you have severe liver or kidney problems or are taking certain medications, you may start at a lower dose of TUKYSA.

Your healthcare professional may interrupt, stop or lower your dose of TUKYSA. This can happen if you experience certain side effects or if your disease gets worse.

Overdose:

If you think you have taken too much TUKYSA, contact your healthcare professional, hospital emergency department or regional poison control centre immediately, even if there are no symptoms.

Missed Dose:

If you miss a dose of TUKYSA, take your next dose at the regularly scheduled time. Do not double your dose or take extra tablets to make up the missed dose.

If you vomit after taking your TUKYSA, do not take an extra dose. Take your next dose at the usual time.

What are possible side effects from using TUKYSA?

These are not all the possible side effects you may feel when taking TUKYSA. You may also get side effects from the other medications taken with TUKYSA. If you experience any side effects not listed here, contact your healthcare professional.

- Weight loss
- Decreased appetite
- Trouble swallowing
- Abdominal pain
- Joint pain
- Chest discomfort or pain
- Flu-like illness
- Fever
- Leg swelling
- Muscle weakness
- Low energy
- Feeling sad
- Painful urination
- Accidental peeing
- Nosebleed
- Rash
- Hair loss
- Night sweats
- Open wound or sore
- Low blood pressure

TUKYSA can cause abnormal blood test results. Your healthcare professional will decide when to perform blood tests and will interpret the results.

Serious side effects and what to do about them

Symptom / effect	Talk to your healthcare professional		Stop taking drug and get immediate medical help
	Only if severe	In all cases	
VERY COMMON			
Diarrhea: loose, watery stools		x	
Nausea: feeling sick		x	
Vomiting: being sick		x	
Hepatotoxicity (liver problems): itching, yellowing of the skin or eyes (jaundice), dark urine, and pain or discomfort in the right upper stomach area.		x	
Stomatitis: Mouth ulcers/sores, redness and swelling of the lining of the mouth		x	
Anemia (low level of red blood cells): shortness of breath, feeling very tired, pale skin, fast heartbeat, loss of energy, weakness		x	
Peripheral neuropathy (damage to nerves): weakness, numbness and pain in hands and feet		x	
Palmar-Plantar Erythrodysesthesia (Hand-Foot Syndrome): tingling, tenderness, redness and swelling usually on the palm of the hand or the sole of the foot		x	
COMMON			
Dehydration (loss of too much fluid from the body): thirst, dry mouth, headache, decreased amount of urine, dark yellow urine		x	
Palpitations: fast heartbeat, heartbeat fluttering rapidly		x	
Rectal hemorrhage: bleeding from the rectum		x	
Seizure (convulsion): muscle twitching, changes in emotions, confusion, loss of consciousness with uncontrollable shaking			x
Vaginal hemorrhage: bleeding from the vagina		x	
UNCOMMON			

Cardiac Failure (heart does not pump blood as well as it should): shortness of breath, fatigue and weakness, swelling in the ankles and feet, cough nausea, rapid or irregular heartbeat			X
Septic Shock (severely low blood pressure from blood infection): fever, dizziness, chills, rapid breathing, rapid heartbeat			X

If you have a troublesome symptom or side effect that is not listed here or becomes bad enough to interfere with your daily activities, talk to your healthcare professional.

Reporting Side Effects

You can report any suspected side effects associated with the use of health products to Health Canada by:

- Visiting the Web page on Adverse Reaction Reporting (<https://www.canada.ca/en/health-canada/services/drugs-health-products/medeffect-canada/adverse-reaction-reporting.html>) for information on how to report online, by mail or by fax; or
- Calling toll-free at 1-866-234-2345.

NOTE: Contact your health professional if you need information about how to manage your side effects. The Canada Vigilance Program does not provide medical advice.

Storage:

- Store TUKYSA at 20°C to 25°C.
- Replace the cap on the bottle after each time you open it.
- 3 months after opening the bottle, any unused tablets should be discarded according to your local rules. If you are not sure, ask your pharmacist about what to do with unused medications that you no longer need.

Keep out of reach and sight of children.

If you want more information about TUKYSA:

- Talk to your healthcare professional
- Find the full product monograph that is prepared for healthcare professionals and includes this Patient Medication Information by visiting the Health Canada website (<https://www.canada.ca/en/health-canada.html>); the manufacturer's website www.pfizer.ca, or by calling 1-800-463-6001.

This leaflet was prepared by Pfizer Canada ULC.

TUKYSA is a registered trademark of Seagen Inc. in Canada

Last Revised: NOV 24, 2025