

PRODUCT MONOGRAPH
INCLUDING PATIENT MEDICATION INFORMATION

Pr **PLENDIL**®

Felodipine

Extended Release Tablets, 2.5 mg, 5 mg and 10 mg, oral

Antihypertensive Agent

Glenwood GmbH Pharmazeutische Erzeugnisse
Arabellastr. 17
81925 Munich
Germany

Date of Authorization:
2025-12-09

Imported by:
McKesson Specialized Distribution Inc.
8449 Lawson Rd, Unit 102
Milton, Ontario L9T 9L1

Submission Control Number: 297826

RECENT MAJOR LABEL CHANGES

Not applicable.

TABLE OF CONTENTS

Sections or subsections that are not applicable at the time of authorization are not listed.

RECENT MAJOR LABEL CHANGES	2
TABLE OF CONTENTS	2
PART I: HEALTH PROFESSIONAL INFORMATION	4
1 INDICATIONS	4
1.1 Pediatrics.....	4
2 CONTRAINDICATIONS	4
4 DOSAGE AND ADMINISTRATION	4
4.1 Dosing Considerations.....	4
4.2 Recommended Dose and Dosage Adjustment	4
4.4 Administration	5
4.5 Missed Dose	5
5 OVERDOSAGE	5
6 DOSAGE FORMS, STRENGTHS, COMPOSITION AND PACKAGING	6
7 WARNINGS AND PRECAUTIONS	6
7.1 Special Populations	8
7.1.1 Pregnancy	8
7.1.2 Breast-feeding.....	8
7.1.3 Pediatrics	8
7.1.4 Geriatrics.....	8
8 ADVERSE REACTIONS	9
8.1 Adverse Reaction Overview	9
8.2 Clinical Trial Adverse Reactions	9
8.3 Less Common Clinical Trial Adverse Reactions	10
8.4 Abnormal Laboratory Findings: Hematologic, Clinical Chemistry and Other Quantitative Data	11
9 DRUG INTERACTIONS	11
9.1 Serious Drug Interactions.....	11

9.2	Drug Interactions Overview	11
9.3	Drug-Behaviour Interactions	12
9.4	Drug-Drug Interactions	12
9.5	Drug-Food Interactions	15
9.6	Drug-Herb Interactions	16
9.7	Drug-Laboratory Test Interactions	16
10	CLINICAL PHARMACOLOGY	16
10.1	Mechanism of Action	16
10.2	Pharmacodynamics	16
10.3	Pharmacokinetics	18
11	STORAGE, STABILITY AND DISPOSAL	20
	PART II: SCIENTIFIC INFORMATION	20
13	PHARMACEUTICAL INFORMATION	20
14	CLINICAL TRIALS	20
15	MICROBIOLOGY	21
16	NON-CLINICAL TOXICOLOGY.....	21
	PATIENT MEDICATION INFORMATION	33

PART I: HEALTH PROFESSIONAL INFORMATION

1 INDICATIONS

PLENDIL (felodipine) is indicated for:

- the treatment of mild to moderate essential hypertension.

The combination of PLENDIL with a thiazide diuretic or a beta-blocker has been found to be compatible and showed an additive antihypertensive effect. The safety and efficacy of concurrent use of PLENDIL with other antihypertensive agents have not been established.

1.1 Pediatrics

Pediatrics (≤ 18 years of age): No data are available to Health Canada; therefore, Health Canada has not authorized an indication for pediatric use.

2 CONTRAINDICATIONS

PLENDIL (felodipine) is contraindicated:

- In patients with a known hypersensitivity to felodipine or any other components of PLENDIL.
- In patients with a known hypersensitivity to other dihydropyridines.
- In women of childbearing potential, in pregnancy, and during lactation. Fetal malformations and adverse effects on pregnancy have been reported in animals.

4 DOSAGE AND ADMINISTRATION

4.1 Dosing Considerations

In clinical trials, doses above 10 mg daily showed an increased blood pressure response but a disproportionately higher incidence of peripheral edema and other vasodilatory adverse events.

4.2 Recommended Dose and Dosage Adjustment

The usual recommended initial dose is 5 mg once daily.

Depending on the patient's response, the dosage should be adjusted accordingly. Dose adjustment, if necessary, should be done at intervals of not less than two weeks.

The maintenance dosage range is 2.5 to 10 mg once daily.

Use in the Elderly

Patients over 65 years of age may develop elevated plasma concentrations of felodipine. A starting dose no higher than 2.5 mg once daily is recommended. A dosage of 10 mg daily should not be exceeded (see 7.1.4 WARNINGS and PRECAUTIONS, Special Populations, Geriatrics).

Use in Children

PLENDIL is not recommended in children (see 7.1.3 WARNINGS and PRECAUTIONS, Special Populations, Pediatrics).

Use in Patients with Impaired Liver Function

Patients with impaired liver function may develop elevated plasma concentrations of felodipine. A starting dose no higher than 2.5 mg once daily is recommended. A dosage of 10 mg daily should not be exceeded (see 7 WARNINGS and PRECAUTIONS, Hepatic/Biliary/Pancreatic).

Use in Patients with Renal Impairment

Modification of the recommended dosage is usually not required in patients with renal impairment.

4.4 Administration

PLENDIL should be swallowed whole and not crushed or chewed. The tablets should not be administered with a meal rich in carbohydrates or fat (see 10.3 CLINICAL PHARMACOLOGY, Pharmacokinetics, Special Populations and Conditions).

4.5 Missed Dose

If the patient misses a dose of PLENDIL and remembers within 12 hours, then the usual dose should be taken as soon as possible. The patient should then return to the regular schedule. If the patient remembers more than 12 hours later, then the missed dose should not be taken, but the next dose taken on time.

A double dose of PLENDIL should never be taken to make up for missed tablets.

5 OVERDOSAGE

Symptoms

Overdosage can cause excessive peripheral vasodilation with marked hypotension and possibly bradycardia.

Treatment

In the case of known overdosage, activated charcoal may be used. If severe hypotension occurs, symptomatic treatment should be instituted. The patient should be placed supine with the legs elevated. The intravenous administration of fluids may be used to treat hypotension. Plasma volume may be increased by infusion of a plasma volume expander. When accompanied by bradycardia, atropine 0.5-1 mg should be administered intravenously. Sympathomimetic drugs predominantly affecting the α 1-adrenoceptor may be given if the above-mentioned measures are considered insufficient. Removal of felodipine from the circulation by hemodialysis has not been established.

For the most recent information in the management of a suspected drug overdose, contact your regional poison control centre or Health Canada's toll-free number, 1-844 POISON-X (1-844-764-7669).

6 DOSAGE FORMS, STRENGTHS, COMPOSITION AND PACKAGING

Table 1 – Dosage Forms, Strengths, Composition and Packaging

Route of Administration	Dosage Form / Strength/Composition	Non-medicinal Ingredients
oral	Extended release tablet, 2.5 mg, 5 mg, 10 mg felodipine	aluminum silicate, carnauba wax, hydrogen peroxide, hydroxypropyl cellulose, hydroxypropyl methylcellulose, iron oxide yellow, iron oxide red-brown (absent in 2.5 mg tablet), lactose anhydrous, microcrystalline cellulose, polyethylene glycol, polyoxyl 40 hydrogenated castor oil, propyl gallate, sodium stearyl fumarate, titanium dioxide.

PLENDIL is available as follows:

PLENDIL 2.5 mg Tablet: A yellow, circular, biconvex film-coated tablet, engraved $\frac{A}{FL}$ on one side and 2.5 on the other.

PLENDIL 5 mg Tablet: A pink, circular, biconvex film-coated tablet, engraved $\frac{A}{FM}$ on one side and 5 on the other.

PLENDIL 10 mg Tablet: A red-brown, circular, biconvex film-coated tablet, engraved $\frac{A}{FE}$ on one side and 10 on the other.

Each tablet strength is available in blister packages containing 30 tablets.

7 WARNINGS AND PRECAUTIONS

Cardiovascular

Congestive Heart Failure

The safety and efficacy of PLENDIL in patients with heart failure have not been established. Caution should, therefore, be exercised when using PLENDIL in hypertensive patients with compromised ventricular function, particularly in combination with a beta-blocker. Acute hemodynamic studies in a small number of patients with New York Heart Association Class II or III heart failure treated with felodipine have not demonstrated negative inotropic effects.

Hypotension, Myocardial Ischemia

PLENDIL may, occasionally, precipitate symptomatic hypotension and rarely syncope. It may lead to reflex tachycardia which, particularly in patients with severe obstructive coronary artery disease, may result in myocardial ischemia. Careful monitoring of blood pressure during the initial administration and titration of felodipine is recommended.

Care should be taken to avoid hypotension especially in patients with a history of cerebrovascular insufficiency, and in those taking medications known to lower blood pressure.

Beta-Blocker Withdrawal

PLENDIL gives no protection against the dangers of abrupt beta-blocker withdrawal; any such withdrawal should be a gradual reduction of the dose of beta-blockers.

Outflow Obstruction

PLENDIL should be used with caution in the presence of left ventricular outflow obstruction.

Peripheral Edema

Mild to moderate peripheral edema was the most common adverse event in the clinical trials. The incidence of peripheral edema was dose-dependent. Frequency of peripheral edema ranged from about 10 percent in patients under 50 years of age taking 5 mg daily to about 30 percent in those over 60 years of age taking 20 mg daily. This adverse effect generally occurs within 2-3 weeks of the initiation of treatment. Care should be taken to differentiate this peripheral edema from the effects of increasing left ventricular dysfunction.

Dependence, Tolerance and/or Abuse Liability

The potential for PLENDIL to cause dependence, tolerance and/or abuse has not been studied.

Driving and Operating Machinery

PLENDIL can induce hypotension and visual disturbances that may impact the ability to drive or use machines. Exercise caution when driving or operating a vehicle or potentially dangerous machinery.

Gastrointestinal

Gingival Hyperplasia

PLENDIL can induce gingival enlargement in patients with pronounced gingivitis and periodontitis. However, such changes may be avoided or reversed by measures of good oral hygiene and mechanical debridement of the teeth. In very rare instances, PLENDIL has also caused gingivitis.

Hepatic/Biliary/Pancreatic

Patients with impaired liver function may have elevated plasma concentrations of felodipine and, therefore, may require lower doses of PLENDIL (see 10.3 CLINICAL PHARMACOLOGY, Pharmacokinetics, Special Populations and Conditions). These patients should have their blood pressure monitored closely during initial administration and after dosage adjustment of PLENDIL. A dosage of 10 mg daily should not be exceeded (see 4.2 DOSAGE AND ADMINISTRATION, Recommended Dose and Dosage Adjustment).

Reproductive Health

PLENDIL is contraindicated in women of childbearing potential and in pregnancy (see 2 CONTRAINDICATIONS).

Sensitivity/Resistance

Lactose

PLENDIL contains lactose and should not be given to patients with hereditary galactose intolerance or glucose-galactose malabsorption.

Skin

Dermatologic Lesion

Along with leucocytoclastic vasculitis, other dermatologic events have been observed. These include rash and flush. All cases of dermatologic lesions should be carefully diagnosed and monitored.

7.1 Special Populations

7.1.1 Pregnancy

PLENDIL is contraindicated in women of childbearing potential and in pregnancy (see 2 CONTRAINDICATIONS).

Fetal malformations and adverse effects on pregnancy have been reported in animals. See 16 NON-CLINICAL TOXICOLOGY, Reproductive and Developmental Toxicology, for additional information on teratology observed in animals.

7.1.2 Breast-feeding

PLENDIL is contraindicated during lactation (see 2 CONTRAINDICATIONS).

7.1.3 Pediatrics

Pediatrics (≤ 18 years of age): No data are available to Health Canada; therefore, Health Canada has not authorized an indication for pediatric use.

7.1.4 Geriatrics

Patients over 65 years of age may have elevated plasma concentrations of felodipine and, therefore, may require lower doses of PLENDIL (see 10.3 CLINICAL PHARMACOLOGY, Pharmacokinetics, Special Populations and Conditions). These patients should have their blood pressure monitored closely during initial administration and after dosage adjustment of PLENDIL. A dosage of 10 mg daily should not be exceeded (see 4.2 DOSAGE AND ADMINISTRATION, Recommended Dose and Dosage Adjustment).

8 ADVERSE REACTIONS

8.1 Adverse Reaction Overview

Mild to moderate peripheral edema was the most common adverse event in the clinical trials. The incidence of peripheral edema was dose-dependent.

8.2 Clinical Trial Adverse Reactions

Clinical trials are conducted under very specific conditions. The adverse reaction rates observed in the clinical trials; therefore, may not reflect the rates observed in practice and should not be compared to the rates in the clinical trials of another drug. Adverse reaction information from clinical trials may be useful in identifying and approximating rates of adverse drug reactions in real-world use.

In 861 patients with essential hypertension treated once daily with 2.5 to 10 mg PLENDIL (felodipine) as monotherapy in controlled clinical trials, the most common clinical adverse events were peripheral edema and headache.

Adverse events that occurred with an incidence of 1.5% or greater at any of the recommended doses of 2.5 mg to 10 mg once a day, without regard to causality, are listed by dose in Table 2 below. These events are reported from controlled clinical trials with patients who were randomized to either a fixed dose of PLENDIL or titrated from an initial dose of 2.5 mg or 5 mg once a day. **A dose of 20 mg once a day has been evaluated in some clinical studies. Although the antihypertensive effect of PLENDIL is increased at 20 mg once a day, there is a disproportionate increase in adverse events, especially those associated with vasodilatory effects (see 4 DOSAGE AND ADMINISTRATION).**

Table 2 Percent Of Patients With Adverse Events In Controlled Trials Of PLENDIL (N=861)* As Monotherapy Without Regard To Causality (Incidence Of Discontinuations Shown In Parentheses)

Body System Adverse Events	Placebo n=334	2.5 mg n=255	5 mg n=581	10 mg n=408
Body as a Whole				
Peripheral Edema	3.3 (0.0)	2.0 (0.0)	8.8 (2.2)	17.4 (2.5)
Asthenia	3.3 (0.0)	3.9 (0.0)	3.3 (0.0)	2.2 (0.0)
Cardiovascular				
Palpitation	2.4 (0.0)	0.4 (0.0)	1.4 (0.3)	2.5 (0.5)
Warm Sensation/Flushing	0.9 (0.3)	3.9 (0.0)	6.2 (0.9)	8.4 (1.2)
Digestive				
Nausea	1.5 (0.9)	1.2 (0.0)	1.7 (0.3)	1.0 (0.7)
Dyspepsia	1.2 (0.0)	3.9 (0.0)	0.7 (0.0)	0.5 (0.0)
Constipation	0.9 (0.0)	1.2 (0.0)	0.3 (0.0)	1.5 (0.2)
Nervous				

Body System	Placebo	2.5 mg	5 mg	10 mg
Adverse Events	n=334	n=255	n=581	n=408
Headache	10.2 (0.9)	10.6 (0.4)	11.0 (1.7)	14.7 (2.0)
Dizziness	2.7 (0.3)	2.7 (0.0)	3.6 (0.5)	3.7 (0.5)
Paresthesia	1.5 (0.3)	1.6 (0.0)	1.2 (0.0)	1.2 (0.2)
Respiratory				
Upper Respiratory Infection	1.8 (0.0)	3.9 (0.0)	1.9 (0.0)	0.7 (0.0)
Cough	0.3 (0.0)	0.8 (0.0)	1.2 (0.0)	1.7 (0.0)
Skin				
Rash	0.9 (0.0)	2.0 (0.0)	0.2 (0.0)	0.2 (0.0)

* Some patients have been exposed to more than one dose level of PLENDIL.

8.3 Less Common Clinical Trial Adverse Reactions

Adverse events that occurred in 0.5 up to 1.5 percent of patients who received PLENDIL in all controlled clinical trials at the recommended dosage range of 2.5 to 10 mg once a day or during post-marketing experience are listed below. These events are listed in order of decreasing severity within each category regardless of relationship to PLENDIL therapy:

Body as a Whole: Chest pain, facial edema, flu-like illness, fever;

Cardiovascular: Tachycardia, premature beats, postural hypotension, bradycardia;

Dermatologic: Pruritus, erythema multiforme, erythema nodosum, leucocytoclastic vasculitis, urticaria, photosensitivity reactions;

Gastrointestinal: Abdominal pain, diarrhea, vomiting, dry mouth, flatulence, acid regurgitation, cholestatic hepatitis, gingival hyperplasia, salivary gland enlargement;

Metabolic: ALT (SGPT) increased;

Musculoskeletal: Arthralgia, muscle cramps, myalgia;

Nervous/Psychiatric: Insomnia, depression, anxiety disorders, irritability, nervousness, somnolence, decrease in libido, tremor, confusion;

Respiratory: Dyspnea, epistaxis;

Special Senses: Visual disturbances;

Urogenital: Impotence/sexual dysfunction, urinary frequency, urinary urgency, dysuria, polyuria.

Serious adverse events reported from controlled clinical trials and during marketing experience (incidence <0.5 percent) were myocardial infarction, hypotension, syncope, angina pectoris, arrhythmia and anemia.

Isolated cases of angioedema have been reported. Angioedema may be accompanied by breathing difficulty.

8.4 Abnormal Laboratory Findings: Hematologic, Clinical Chemistry and Other Quantitative Data

Clinical Trial Findings

For the following laboratory values, statistically significant decreases were observed; bilirubin, red blood count, hemoglobin, and urate. Statistically significant increases were found in erythrocyte sedimentation rate and thrombocyte count. In isolated cases, there were increased liver enzymes. None of the changes were considered to be of clinical significance.

9 DRUG INTERACTIONS

9.1 Serious Drug Interactions

Serious Drug Interactions

Strong Inhibitors of CYP3A4: Use of PLENDIL with drugs that result in strong inhibition of CYP3A4, such as ketoconazole, clarithromycin, ritonavir, may lead to increased plasma levels of felodipine and associated serious adverse events. Such concomitant use should be avoided.

9.2 Drug Interactions Overview

As with all drugs, care should be exercised when treating patients with multiple medications. Dihydropyridine calcium channel blockers undergo biotransformation by the cytochrome P-450 system, mainly via the CYP 3A4 isoenzyme. Coadministration of felodipine with other drugs which follow the same route of biotransformation may result in altered bioavailability of felodipine or these drugs. Dosages of similarly metabolized drugs, particularly those of low therapeutic ratio, and especially in patients with renal and/or hepatic impairment, may require adjustment when starting or stopping concomitantly administered felodipine to maintain optimum therapeutic blood levels.

Cytochrome P-450 Enzyme Substrates

Drugs known to be substrates of the cytochrome P-450 system include: benzodiazepines, flecainide, imipramine, propafenone, terfenadine, theophylline (see Table 3).

Cytochrome P-450 Enzyme Inhibitors: Drugs known to be inhibitors of the cytochrome P-450 system include: azole antifungals (ketoconazole and itraconazole), cimetidine, cyclosporine, erythromycin, quinidine, warfarin. Enzyme inhibitors of the cytochrome P450 3A4 system have been shown to cause an increase in felodipine plasma concentrations (see Table 3).

CYP 3A4 Enzyme Inducers: Enzyme inducers of the cytochrome P450 3A4 system have been shown to cause a decrease in plasma concentrations of felodipine, e.g. Hypericum perforatum (Saint John's Wort) (see Drug-Herb Interactions). Drugs known to be inducers of the cytochrome P-450 3A4 isoenzyme include: phenobarbital, phenytoin, rifampin (see Table 3).

Concomitant Use With Strong Inhibitors of CYP3A4

Use of PLENDIL with drugs that result in strong inhibition of CYP3A4, such as ketoconazole, clarithromycin, ritonavir, may lead to increased plasma levels of felodipine and associated serious adverse events. Such concomitant use should be avoided.

An observational study demonstrated an increased risk of hospitalisation with acute kidney injury when felodipine was used concomitantly with clarithromycin in elderly patients (> 65 years of age) compared to when it was used concomitantly with azithromycin, odds ratio [felodipine: 2.97 (95% C.I. 1.09-8.06)].

Other Concomitant Therapy

In healthy subjects there were no clinically significant interactions when felodipine was given concomitantly with indomethacin or spironolactone.

9.3 Drug-Behaviour Interactions

The interaction of PLENDIL with individual behavioural risks (e.g. cigarette smoking, cannabis use, and/or alcohol consumption) has not been studied.

9.4 Drug-Drug Interactions

The drugs listed in Table 3 are based on either drug interaction case reports or studies, or potential interactions due to the expected magnitude and seriousness of the interaction (i.e., those identified as contraindicated).

Table 3 - Established or Potential Drug-Drug Interactions

Proper/Common name	Source of Evidence	Effect	Clinical comment
Alcohol	T	Alcohol induces vasodilation.	Alcohol can enhance the hemodynamic effects of felodipine.

Proper/Common name	Source of Evidence	Effect	Clinical comment
Beta-adrenoceptor Blocking Agents (such as Metoprolol)	CT	A pharmacokinetic study of felodipine in conjunction with metoprolol demonstrated no significant effects on the pharmacokinetics of felodipine. The AUC and C _{max} of metoprolol, however, were increased approximately 31 and 36 percent, respectively.	In controlled clinical trials, however, beta-blockers including metoprolol were concurrently administered with felodipine and were well tolerated.
Cytochrome P-450 Enzyme substrates: benzodiazepines, flecainide, imipramine, propafenone, terfenadine, theophylline	CT	Enzyme substrates of the cytochrome P450 3A4, when coadministered with felodipine, may act like P450 3A4 inhibitors and cause an increase in felodipine plasma concentrations.	Dose adjustment and monitoring may be required.
Cytochrome P-450 Enzyme Inhibitors:			
Cimetidine	CT	In healthy volunteers pharmacokinetic studies showed an approximately 50 percent increase in the area under the plasma concentration time curve (AUC) as well as the C _{max} of felodipine when given concomitantly with cimetidine.	It is anticipated that a clinically significant interaction may occur. Therefore, it is recommended that low doses of PLENDIL be used when given concomitantly with cimetidine.

Proper/Common name	Source of Evidence	Effect	Clinical comment
Clarithromycin	CT	In elderly patients (> 65 years of age), concomitant use of felodipine with clarithromycin was associated with increased risk of hospitalization with acute kidney injury.	Avoid concomitant use.
Erythromycin	CT	Concomitant treatment with erythromycin has been shown to cause an increase in felodipine plasma levels. Coadministration of felodipine extended-release tablets with erythromycin resulted in approximately 2.5-fold increase in the AUC and C _{max} , and about 2-fold prolongation in the half-life of felodipine.	It is expected that a clinically significant interaction may occur. Therefore, low doses of PLENDIL are recommended to be used when given concomitantly with erythromycin.
Itraconazole Ketoconazole	CT	Coadministration of another extended release formulation of felodipine with itraconazole resulted in approximately 8-fold increase in the AUC, more than 6-fold increase in the C _{max} , and 2-fold prolongation in the half-life of felodipine.	Caution should be used when CYP3A4 inhibitors are coadministered with felodipine and a conservative approach to dosing felodipine should be taken.

Proper/Common name	Source of Evidence	Effect	Clinical comment
CYP 3A4 Enzyme Inducers: Phenytoin, Carbamazepine and Phenobarbital	CT	In a pharmacokinetic study, maximum plasma concentrations of felodipine were considerably lower in epileptic patients on long term anticonvulsant therapy (phenytoin, carbamazepine, phenobarbital) than in healthy volunteers. The mean area under the felodipine plasma concentration-time curve was also reduced in epileptic patients to approximately 6% of that observed in healthy volunteers.	Since a clinically significant interaction may be anticipated, alternative antihypertensive therapy should be considered in these patients.
Digoxin	CT	When given concomitantly with felodipine as conventional tablets the peak plasma concentration (C _{max}) of digoxin was significantly increased.	With the extended release formulation of felodipine there was no significant change in C _{max} or AUC of digoxin.
Tacrolimus	C	Felodipine may increase the concentration of tacrolimus.	When used together, the tacrolimus serum concentration should be monitored and the tacrolimus dose may need to be adjusted.

Legend: C = Case Study; CT = Clinical Trial; T = Theoretical

9.4 Drug-Food Interactions

Grapefruit juice - Coadministration of felodipine with grapefruit juice resulted in more than 2-fold increase in the AUC and C_{max}, but no prolongation in the half-life of felodipine (see 10.3 CLINICAL PHARMACOLOGY, Pharmacokinetics, Special Populations and Conditions).

Published data show that through inhibition of cytochrome P-450, grapefruit juice can increase plasma levels and augment pharmacodynamic effects of dihydropyridine calcium channel blockers. In

view of the absolute bioavailability of PLENDIL, the potential for a significant increase in pharmacodynamic effects exists (see 10.3 CLINICAL PHARMACOLOGY, Pharmacokinetics). Therefore, the consumption of grapefruit juice prior to or during treatment with PLENDIL should be avoided.

Food – The bioavailability of PLENDIL is influenced by the presence of food. When administered with either a high fat or carbohydrate diet, C_{max} is significantly increased by approximately 1.5 to 2.0 fold; AUC is unchanged. This may cause a clinically significant fall in blood pressure. Therefore PLENDIL should not be administered with meals rich in carbohydrate or fat (see 10.3 CLINICAL PHARMACOLOGY, Pharmacokinetics).

9.5 Drug-Herb Interactions

Hypericum perforatum – St-John's Wort is an inducer of CYP3A4 and has been shown to cause a decrease in plasma concentrations of felodipine. Therefore, dosage of felodipine may have to be increased.

9.6 Drug-Laboratory Test Interactions

Interactions with laboratory tests have not been established.

10 CLINICAL PHARMACOLOGY

10.1 Mechanism of Action

PLENDIL (felodipine) is a calcium ion influx inhibitor (calcium channel blocker). Felodipine is a member of the dihydropyridine class of calcium channel blockers.

The therapeutic effect of this group of drugs is believed to be related to their specific cellular action of selectively inhibiting transmembrane influx of calcium ions into cardiac muscle and vascular smooth muscle. The contractile processes of these tissues are dependent upon the movement of extracellular calcium into the cells through specific ion channels. Felodipine blocks transmembrane influx of calcium through the slow channel without affecting to any significant degree the transmembrane influx of sodium through the fast channel. This results in a reduction of free calcium ions available within cells of the above tissues.

Felodipine does not alter total serum calcium. *In vitro* studies show that the effects of felodipine on contractile mechanisms are selective, with greater effects on vascular smooth muscle than on cardiac muscle. Negative inotropic effects can be detected *in vitro*, but such effects have not been seen in intact animals.

The effect of felodipine on blood pressure in man is principally a consequence of a dose-related decrease in peripheral vascular resistance, with a modest reflex increase in heart rate (see 10.2 CLINICAL PHARMACOLOGY, Pharmacodynamics).

10.2 Pharmacodynamics

Animal Pharmacodynamics

Felodipine is a selective vasodilating antihypertensive dihydropyridine which has been shown to lower arterial blood pressure in normotensive rats, cats and dogs and in rats with primary or secondary forms of hypertension.

The plasma concentration needed for 20% reduction of mean arterial pressure, in normotensive dogs and spontaneously hypertensive rats (SHR), was of the order of 15-30 nmol/L.

The oral dose of felodipine required to obtain a 20% reduction of mean arterial blood pressure in dogs (3 studies, 17 males in total) and in rats (1 study, 11 males) when given by gavage or in capsules was 1-5 µmol/kg. When felodipine was mixed in with the diet of SHR (3 studies, treatment duration ranging from 2 weeks to 6 months) daily doses of up to 85 µmol/kg had to be administered to achieve the same reduction in mean arterial pressure.

The primary effects of felodipine at plasma concentrations of 1-30 nmol/L were examined in several studies in rats and dogs given felodipine intravenously and orally. The only primary effect observed, in addition to arterial vasodilation, was diuresis-natriuresis due to a tubular action.

Felodipine exhibits a hundredfold vascular versus myocardial selectivity, based on the inhibitory potency ratios for vascular and cardiac tissues, studied *in vitro*.

Acutely, there is reflex tachycardia which may be blocked by beta-adrenoceptor antagonists and which moderates during maintained antihypertensive treatment due to baroreflex resetting.

Two different studies, in which felodipine was given to 8 male rats in a dose of 247 nmol/kg administered intravenously over 90 minutes and to 7 female rats as an oral dose of 78 µmol/kg with food for 6 months, have demonstrated that the reabsorption of filtered sodium is reduced in the distal tubules and collecting ducts in the kidney. Potassium excretion and absorption were unaffected by felodipine.

Human Pharmacodynamics

The acute hemodynamic effect of felodipine is a reduction in total peripheral resistance which leads to a decrease in blood pressure associated with a modest reflex increase in heart rate. This reflex increase in heart rate frequently occurs during the first week of therapy and generally attenuates over time. Heart rate increases of 5-10 beats per minute may be seen during chronic administration. The effect on the heart rate is inhibited by beta-blocking agents. Following administration of PLENDIL a reduction in blood pressure generally occurs within two to five hours.

During chronic administration, substantial blood pressure control lasts for approximately 24 hours; reductions in diastolic blood pressure at trough plasma levels were 40-60% of those at peak plasma levels. The antihypertensive effect is dose-dependent and correlates with the plasma concentration of felodipine.

Felodipine in therapeutic doses has no effect on conduction in the conducting system of the heart and no effect on the A-V nodal refractoriness. No direct additional effects to those registered after beta-blockade are observed when PLENDIL is given concomitantly.

Renal vascular resistance is decreased by felodipine while glomerular filtration rate remains unchanged. Mild diuresis, natriuresis and kaliuresis have been observed during the first week of therapy. No significant effects on serum electrolytes have been observed during short and long-term therapy. No general salt and water retention occurs during long-term therapy. In clinical trials increases in noradrenaline plasma levels have been observed.

10.3 Pharmacokinetics

Animal Pharmacokinetics

Felodipine is rapidly and completely absorbed after oral administration in rats and dogs. First-pass elimination reduces oral bioavailability to 20 to 30% for a dose of 5 µmol/kg. This is comparable with the 15% availability in man. Saturation of the first-pass elimination at high felodipine doses to the rat, 150 µmol/kg, results in almost 100% bioavailability.

Felodipine exhibits multiexponentially declining plasma concentration-time curves after i.v. doses. In the terminal phase distribution volumes are 10 to 20 L/kg. The binding to plasma proteins is >99.5% in rat, dog and man. An autoradiography study with (¹⁴C) felodipine in mice did not indicate any irreversible tissue binding. It was also found that the drug and/or its metabolites pass the blood-brain and placental barriers.

Elimination of felodipine was studied in single dose studies in rats (7 males) and dogs (3 females) administered drug in doses of 0.2 µmol/kg i.v. and 5 µmol/kg oral. The studies indicate that felodipine is almost exclusively eliminated as metabolites in almost equal amounts in the urine and feces of the rat and dog.

The primary step in the biotransformation of felodipine is oxidation to the corresponding pyridine analogue, which lacks vasodilating properties. Sequential metabolic processes involve ester hydrolysis and hydroxylation of pyridine methyl groups.

Eight different inactive metabolites have been identified in rat urine. The urinary metabolic pattern in mouse, rabbit, dog and man resembles in all essentials that found in the rat.

Human Pharmacokinetics

Absorption

Felodipine is completely absorbed from the gastrointestinal tract after oral administration.

Distribution:

Due to rapid biotransformation of felodipine during its first pass through the portal circulation the systemic availability is approximately 15% and is independent of the dose in the range of 5-20 mg per day. The plasma protein binding of felodipine is approximately 99%. It is bound predominately to the albumin fraction.

Metabolism:

Felodipine is extensively metabolized in the liver, predominantly by cytochrome P-450 CYP 3A4. Six metabolites, which account for 23% of the oral dose, have been identified: none has significant vasodilating activity.

Elimination:

After 72 hours, approximately 70% of a given dose is excreted as metabolites in the urine and 10% is secreted in the feces. Less than 0.5% of a dose is recovered unchanged in the urine.

Felodipine has been observed to have a mean blood clearance of 914±355 mL/min in hypertensive patients, 606±245 mL/min in elderly hypertensive patients and 1337±413 mL/min in young healthy

volunteers. Its mean terminal half-life was 24.5 ± 7.0 hours in hypertensive patients, 27.5 ± 8.4 hours in elderly hypertensive patients and 14.1 ± 5.6 hours in young healthy volunteers.

Effect of Extended Release Formulation on Pharmacokinetics: The extended release formulation prolongs the absorption phase of felodipine resulting in an increased time to reach peak plasma concentrations (t_{max}), and a reduced maximum plasma concentration (C_{max}). The mean t_{max} ranges from 2.5 to 5 hours. The area under the plasma concentration versus time curve and C_{max} are linearly related to the dose in the 10 to 40 mg range. Following administration of PLENDIL to hypertensive patients, mean C_{max} at steady state is approximately 20% higher after multiple doses than after a single dose. No increase in the AUC is found during multiple dosing. The inter-individual variation in C_{max} and AUC after repeated dosing is approximately threefold and indicates a need for individualized dosing.

Effect of Food on Pharmacokinetics: The bioavailability of PLENDIL is not influenced by the presence of food in the gastrointestinal tract. However, the peak plasma concentration of felodipine (C_{max}) is significantly increased by 1.5 to 2 fold when PLENDIL is taken after a high fat or high carbohydrate meal versus fasting. Because the effects of felodipine on blood pressure are related to plasma levels, this increase in C_{max} may cause a clinically significant fall in blood pressure. Therefore PLENDIL should not be administered with meals rich in carbohydrate or fat. However, the absorption characteristics of felodipine are not affected when PLENDIL is administered with a light meal low in fat and carbohydrates (i.e. 2 slices of toast with cheese, 150 mL milk with cornflakes, and 150 mL orange juice).

Studies in healthy male volunteers showed significant alterations in the pharmacokinetics of felodipine when felodipine was administered concomitantly with grapefruit juice. Following the administration of a single dose of plain felodipine 5 mg tablets with 200 mL grapefruit juice or 200 mL water, AUC and C_{max} of felodipine increased about threefold as compared to administration with water. When felodipine extended release tablets were administered as PLENDIL 10 mg with 250 mL grapefruit juice, felodipine AUC and C_{max} values doubled as compared to those observed with water. When grapefruit juice was taken for up to 24 hours prior to PLENDIL administration, a significant pharmacokinetic interaction was observed (see 9.4 DRUG INTERACTIONS, Drug-Food Interactions).

Special Populations and Conditions

- **Geriatrics:** Plasma concentrations of felodipine, after a single dose and at steady state, increase with age. Mean clearance of felodipine in elderly hypertensives (mean age 74 years) was only 45 percent of that in young volunteers (mean age 26 years). At steady state mean AUC for young patients was 39 percent of that for the elderly patients.
- **Pregnancy and Breast-feeding:** Animal studies have demonstrated that felodipine crosses the blood-brain barrier and the placenta. PLENDIL is contraindicated during pregnancy and while breast-feeding (see 2 CONTRAINDICATIONS).
- **Hepatic Insufficiency:** In patients with hepatic disease, the clearance of felodipine was reduced to about 60 percent of that seen in normal young volunteers.
- **Renal Insufficiency:** Renal impairment does not alter the plasma concentration profile of felodipine. Although higher concentrations of the metabolites are present in the plasma due to decreased urinary excretion, these are hemodynamically inactive.

11 STORAGE, STABILITY AND DISPOSAL

Store PLENDIL at 15-30°C.

PART II: SCIENTIFIC INFORMATION

13 PHARMACEUTICAL INFORMATION


Drug Substance

Proper name: felodipine

Chemical name: 3,5-pyridinedicarboxylic acid, 4-(2,3-dichlorophenyl)-1,4-dihydro 2,6-dimethyl-,ethyl methyl ester

Molecular formula and molecular mass: C₁₈H₁₉Cl₂NO₄, 384.26

Structural formula:



Physicochemical properties: Felodipine is a slightly yellowish, crystalline powder. It is not hygroscopic. It is soluble in acetone, dichloromethane, ethanol, heptane, methanol, n-octanol, 2-propanol, and practically insoluble in water.

14 CLINICAL TRIALS

No clinical trial information is available for this drug product.

15 MICROBIOLOGY

No microbiological information is required for this drug product.

16 NON-CLINICAL TOXICOLOGY

General Toxicology:

Acute Toxicity

Table 4 Acute Toxicity Studies

SPECIES	ROUTE	SEX	NO. OF ANIMALS	DOSE LEVELS	LD50 VALUES	LD50 VALUES
				mg/kg	mg/kg	µmol/kg
Mice	p.o.	Male	10	100-700	240 (185-279)	630 (480-730)
	p.o.	Female	10	150-750	264 (189-330)	690 (490-860)
Mice	i.v.	Male	10	6-12	8.6 (8.0-9.5)	22 (21-25)
	i.v.	Female	10	5-12.5	10.4 (9.2-11.8)	27 (24-31)
Rats	p.o.	Male	6	1000-4500	2390 (1710-2840)	6230 (4450-7380)
	p.o.	Female	6	125-4500	2250 (1300-3850)	5900 (3400-10000)
Rats	i.v.	Male	6	5-10	6.8 (5.9-7.5)	18 (15-19)
	i.v.	Female	6	3-8	6.4 (4.9-7.1)	17 (13-19)

Signs of acute toxicity were sedation, convulsions, diarrhea and body weight loss.

In a single study, Beagle dogs (2 male, 2 female) were studied after single doses of 20, 40 and 80 µmol/kg felodipine given orally, and after repeated doses of 10 µmol/kg felodipine b.i.d. given orally for 7 days followed by 20 µmol/kg felodipine b.i.d. given orally for 3 days. After single administration of felodipine 80 µmol/kg, food consumption was decreased for 1 to 3 days. At all dose levels the ECGs revealed tachycardia which persisted for more than 6 hours. When the dogs received the repeated doses of felodipine there was a dose-dependent hyperemia of the mucus membranes and skin, and dose-dependent reduced food consumption. Tiredness was reported at the 20 µmol/kg b.i.d. dose level. Two dogs died at the dose level 20 µmol/kg (7.7 mg/kg) b.i.d. It is concluded that after a single oral dose of felodipine, the minimum lethal dose in dogs is higher than 80 µmol/kg. With repeated daily doses it has been shown that 20 µmol/kg (7.7 mg/kg) twice daily with four hour intervals, may be lethal to dogs.

General Toxicity After Repeated Administration

The general toxicity of felodipine has been studied after repeated oral administration in rats and dogs and repeated i.v. administration in rats.

The studies performed are summarized in Table 5.

Table 5 Repeat Dose Toxicity Studies

Species	No. of Animals/ Sex/Group	Route of Administration	Frequency Duration of Dosing	Dose Groups		Deaths/ Group		Toxic Effects
				($\mu\text{mol/kg}$)	(mg/kg)	M	F	
Rats	6	p.o.	Once daily for one month.	0	0			At the 150 $\mu\text{mol/kg}$ dose, hyperemia manifested in redness of mucous membranes nose and ears. Decrease in mean food intake and body weight in females during week 1. Decrease in blood glucose in females and serum potassium concentrations in males. Insignificant increase in serum protein and albumin concentrations in females. Increase in serum urea in females. Reduction in testicular weight without signs of morphological change. Elevated kidney weight values in females.
				5	2			
				15	5.8			
				50	19.2			
				150	57.6			
Rats	10	p.o.	Once daily for 5 weeks.	0	0			The 130 $\mu\text{mol/kg}$ group showed decreased food consumption (most notable on earlier treatment days), lowered levels of plasma neutral fat, increases in liver weight and decreases in submaxillary gland weight. Male weight gain was suppressed. The 470 $\mu\text{mol/kg}$ group showed decreased spontaneous locomotor activity, ptosis, loss of hair gloss, abdominal swelling, nasal bleeding, anemia and deaths from debility in cases with severe toxic symptoms. Decreases in female blood lymphocyte ratio and increase in neutrophil ratio. Increase in liver weight and decreases in submaxillary gland weight. In the dead animals, atrophy of the spleen and thymus, degenerative hematopoiesis of the bone marrow were found.
				5	2			
				26	10			
				130	50			
				470	180	5	9	
Rats	25	p.o.	Once daily for 6 months.	0	0	2		Distinct hyperemia of the ears, lasting several hours after treatment in all 3 active groups. In mid and high dose groups, males became slightly tense and irritated after 6 weeks of treatment. The high dose group showed a clear reduction in food consumption during the first few weeks with a corresponding lag in body weight gain, most pronounced in females. Water consumption was increased in high dose males and doubled in high dose females after 3rd week. Blood glucose concentrations were decreased in all groups but most pronounced in the high dose group. The high dose group also showed a decrease in serum potassium and chloride concentrations and some decrease in osmolality of the urine. Females showed increased serum urea concentrations. Mid and high dose animals showed low-degree weight increases of several organs. High dose females had enlarged colons, often doubled, in both sexes increased ileum weight. Both mid and high dose animals showed increased activity of the zona glomerulosa of the adrenal glands. Death of the low dose female was attributed to myocarditis and the 4 deaths in the high dose group to acute circulatory insufficiency. Only the high dose deaths were attributed to felodipine.
				5	2		1	
				25	9.6			
				125	48.0	3	1	

Table 5 Repeat Dose Toxicity Studies (continued)

Species	No. of Animals/ Sex/Group	Route of Administration	Frequency Duration of Dosing	Dose Groups		Deaths/ Group		Toxic Effects
				(μ mol/kg)	(mg/kg)	M	F	
Rats	10	i.v.	Once daily for two weeks.	0	saline			Dose levels 0.3 and 1.0 μ mol/kg produced peripheral vasodilation and apparent hyperthermia 1-3 hours after dosing. Higher liver weight gain in males. Males given 1 μ mol/kg dose showed inferior body weight gain during first 4 days of treatment.
				0	saline			
				0.1	0.04			
				0.3	0.12			
				1.0	0.38			
Dogs	2	p.o.	Once daily for 1 month.	0	0			Dose dependent hyperemia of the mucous membrane and abdominal skin in mid and high doses. Dose dependent tachycardia noted in all groups. High dose groups showed depression of ST-j or ST-segment. High dose group males and females showed non-significant increase of heart and kidney weights.
				5	2			
				10	3.8			
				20	9.6			
Dogs	5 grp 1 5 grp 2 5 grp 3 5 grp 4	p.o.	Twice daily for 12 months in grps 1-4.	0	0			Increased heart rate throughout the study. Mid and high dose animals developed decreased heart rate before the first daily dose. Hyperemia of the mucous membranes post-dose noted in mid and high dose animals (more pronounced in the high dose). A decrease in packed cell volume hemoglobin concentration and number of red blood cells, with preponderance in males, noted in mid and high dose groups.
				1.0 b.i.d.	0.38 b.i.d.			
				3.0 b.i.d.	1.2 b.i.d.			
				6.0 b.i.d.	2.3 b.i.d.			
Dogs	3 grp 5 3 grp 6 3 grp 7 3 grp 8	p.o.	Twice daily for 6 months in grps 5-8.	0	0			Decrease in osmolality of the urine in mid dose females and both sexes of the high dose. Insignificant increase in serum glucose concentration in mid dose females. Enlargement of the gingiva observed clinically in both mid and high dose groups; pathologically, a non-inflammatory gingival hyperplasia with male high dose preponderance. Increased activity of the zona glomerulosa in mid dose animals.
				1.0 b.i.d.	0.38 b.i.d.			
				3.0 b.i.d.	1.2 b.i.d.			
				6.0 b.i.d.	2.3 b.i.d.	1*	1*	
<p>* The initial high dose was 10 μmol/kg. At this dose 2 dogs died after 3 days. Death was attributed to acute circulatory failure. All dogs showed tiredness post dose. After the dose was reduced, this sign was reduced.</p>								

Carcinogenicity:

The carcinogenic effect of felodipine has been studied in mice (50 males and 50 females/group) at doses of 40, 120 and 360 µmol/kg over 99 weeks and rats (50 males and 50 females/group) at doses of 20, 60 and 180 µmol/kg over 112 weeks. In the mouse study, the high dose group (360 µmol/kg) showed an increased incidence of hepatocellular neoplasms (carcinomas and adenomas) compared with the animals terminally sacrificed from the other groups (see Table 6).

Table 6 Incidence Of Hepatocellular Neoplasms In Mice

Group	Sex	No. of Animals ¹ / Group ²	Dose Level µmol/L	Total No. of Mice with Adenomas ¹	Total No. of Mice with Carcinomas ¹
1	M	93 (14)	0	3	18 (2)
2	M	41 (9)	40	3	8 (1)
3	M	42 (9)	120	1	7
4	M	39 (25)	360	0	11 (3)
1	F	94 (36)	0	1 (1)	4 (1)
2	F	46 (14)	40	1	2
3	F	48 (25)	120	2	1 (1)
4	F	42 (28)	360	2 (1)	3

1 Number of animals entering week 54 and continuing to termination of the study.

2 Numbers in brackets are the number of mice that died between week 54 and termination of the study.

Few neoplasms (10 in total; 6 in the treatment groups, 4 in controls) were found in animals dying before termination of the study. Because of the difference in mortality between the groups, it was necessary to analyze pre-terminal and terminal deaths together in order to evaluate the carcinogenic potential of felodipine in mice. In doing this analysis, there was no significant difference between treated and untreated groups. Although there was an increased incidence of neoplasms in the high dose group compared with animals terminally sacrificed from the other groups, the incidence of hepatic carcinomas in all groups were within the historical laboratory control range for this strain of mice.

In view of the differing survival patterns in the different dose groups, the mouse carcinogenicity study was subjected to a further statistical analysis which indicated that no significant increase in hepatic neoplasms had occurred in the felodipine treated groups. The liver slides from this study were also examined "blindly" by another specialized pathologist who concluded that "comparisons of the incidence or mean grades or any hepatocellular proliferative lesions, or combination of lesions fail to suggest a compound-related effect".

A repeat study in male mice with felodipine in doses of 40, 90 and 180 µg/kg over 76 weeks was performed. The results of the histopathological examination of the livers showed that there was no significant increase in hepatic tumours in the felodipine treated animals at any of the dose levels.

In the rat carcinogenicity study discussed above, an increased incidence of benign interstitial cell tumours (Leydig, cell tumours) was found in the testes of the low (26%=13/50), mid (38%=19/49), and high (40%=20/50) dose males when compared to controls (9%=9/100). It is considered that these tumours have an endocrinological basis in the rat as they can be induced by various experimental

interventions that increase the level of luteinizing hormone (LH) in the blood. *In vivo* and *in vitro* studies have shown effects of felodipine which indicate that there is an endocrinological mechanism behind the increased incidence of Leydig cell tumours in rats. In human studies felodipine (given as a single doses of 5, 10 or 20 mg to 12 young healthy volunteers and as 10-20 mg daily for 8 weeks followed by 10 mg daily for one week to 10 hypertensive patients) has not revealed effects on testosterone levels. LH levels were also determined in the hypertensive patients and showed no abnormalities.

Treated and untreated rats in the above carcinogenicity study also differed with respect to the occurrence of bile-duct hyperplasia and squamous-cell hyperplasia in the esophageal groove of the forestomach. Bile-duct hyperplasia was found in 30% (males) and 23% (females) of the control groups and at increased incidence in medium (42% males, 24% females) and high (64% males, 46% females) dose animals. Squamous-cell hyperplasia of the esophageal groove of the forestomach was found in 0% (males) and 6% (females) of the controlled groups, and at an increased incidence in the low (40 % males, 16% females), medium (46% males, 34% females) and high (56% males, 38% females) dose animals.

Genotoxicity:

Felodipine has not shown any mutagenic potential.

Reproductive and Developmental Toxicology:

Teratogenic Effects

Studies in pregnant rabbits administered doses of 0.46, 1.2, 2.3 and 4.6 mg/kg/day (from 0.4 to 4 times the maximum recommended human dose on a mg/m² basis) showed digital anomalies consisting of reduction in size and degree of ossification of the terminal phalanges in the fetuses. The frequency and severity of the changes appeared dose-related and were noted even at the lowest dose. These changes have been shown to occur with other members of the dihydropyridine class. Similar fetal anomalies were not observed in rats given felodipine.

In a teratology study in Cynomolgus monkeys, no reduction in the size of the terminal phalanges was observed but an abnormal position of the distal phalanges was noted in about 40 percent of the fetuses.

Non-Teratogenic Effects

In a study on fertility and general reproductive performance in rats, prolongation of parturition with difficult labour and an increased frequency of fetal and early postnatal deaths were observed in the groups treated with doses of 9.6 mg/kg/day and above.

Significant enlargement of the mammary glands in excess of the normal enlargement for pregnant rabbits was found with doses greater than or equal to 1.2 mg/kg/day. This effect occurred only in pregnant rabbits and regressed during lactation. Similar changes in the mammary glands were not observed in rats or monkeys.

Please also refer to Table 7.

Table 7 Reproductive and Developmental Toxicity Studies

Species and Strain	No. of Animals/ Group	Period of Administration	Route of Administration			Dams	Litters	
				µmol/kg	mg/kg			
Rats, Sprague Dawley	15 M	Apr. 11 weeks ¹ Apr. 10 weeks ¹ (Seg I)	p.o.	0		<u>Parents</u> Dose dependent prolongation of parturition and hard labour in the animals receiving 25 µmol/kg or 70 µmol/kg.	Dose dependently increased frequencies of late fetal deaths and early postnatal deaths in animals receiving 25 µmol/kg or 70 µmol/kg.	
	30 F			10	3.8			
				25	9.6			
				70	26.9			
Rats, Sprague Dawley	20 F	Days 6-15 of gestation (Seg II)	p.o.	0		<u>70 µmol/kg</u> Slightly lower food consumption during the dosing period and slightly reduced body weight gain towards the end of the dosing period.	No signs of embryotoxic, teratogenic or fetotoxic effects related to the treatment could be detected.	
					10			3.8
					25			9.6
					70			26.9
Rats, Sprague Dawley	20 F	From day 15 of gestation to day 20 post part. (Seg III)	p.o.	0		<u>30 µmol/kg</u> Slight prolongation of the gestation period, prolonged parturition and hard labour.	<u>30 µmol/kg</u> Increased frequencies of stillborn fetuses and early postnatal deaths.	
					3			1.2
					10			3.8
					30			11.5
Rabbit, New Zealand White	5 F	Days 6-19 of gestation (seg II, pilot study)	p.o.	0		One control animal and 2 high dose animals were found dead during the study. Dose dependent decrease in body weight changes and food consumption values.	In a dose dependent manner, the mean resorption values were higher than control and the mean fetal viability values were lower than control in the treated groups. The mean fetal body weights and the mean fetal lengths were less than control in all of the treated groups. These effects had an insignificant dose relation.	
					10			3.8
					25			9.6
					50			19.2
Rabbit, New Zealand White	15 F	Days 6-19 of gestation (Seg II)	p.o.	03	1.2	<u>3 µmol/kg</u> Depression noted more frequently in all treated groups than in the control group. A dose-related incidence of thickening of mammary tissue was observed in all treated groups (1/15, 10/15, & 9/14, resp.) at sacrifice day 29 of gestation. <u>6 µmol/kg</u> Depression, thickening of mammary tissue. Incidence of premature deliveries slightly higher than control & wt losses during days 20-29 higher than control; these effects were also observed in the animals receiving 12 µmol/kg. <u>12 µmol/kg</u> Depression, body wt loss, thickening of mammary tissue, premature deliveries as above.	Dose-related skeletal anomalies were observed in the extremities of fetuses in all treated groups. <u>3 µmol/kg</u> Small distal phalanges in the 4th digits of the hind feet. <u>6 µmol/kg</u> Small or missing distal phalanges in the 4th digits of the hind feet. Small distal phalanges in the first digits of the fore feet. One mid dose fetus had one fore foot with the entire first digit missing. <u>12 µmol/kg</u> Small or missing distal phalanges in the 4th digits of the hind feet. Small or missing distal phalanges in the first digits of the fore feet.	
					6			2.3
					12			4.6

¹Males were dosed for 9 weeks prior to mating and through the mating period (maximum 12 days). Females were dosed for 2 weeks prior to mating and then up to 21 days post partum. Half of the females were killed on Day 14 of pregnancy.

Table 7 Reproductive and Developmental Toxicity Studies (continued)

Species and Strain	No. of Animals/ Group	Period of Administration	Route of Administration	$\mu\text{mol/kg}$	mg/kg	Dams	Litters
Rabbit, New Zealand White	15 F	Days 6-18 of gestation (Seg II)	p.o.	0 6 12	 2.3 4.6	<p><u>6 $\mu\text{mol/kg}$</u> Reduced food intake during first few days of dosing; this was also seen in the animals receiving 12 $\mu\text{mol/kg}$ (dose related). Suppression of body wt gain during the first few days of dosing, also seen in the animals receiving 12 $\mu\text{mol/kg}$. A dose related enlargement of mammary glands; also macroscopic changes in the glands including colour darkening and no milk leakage on dissection and microscopic changes, including increased number of small acini with abundant eosinophilic secretion.</p> <p><u>12 $\mu\text{mol/kg}$</u> Reduced food intake and suppression of body weight gain as described above; wt loss on cessation of dosing. Cold ears (more persistent and consistent observation). Enlargement of mammary glands.</p> <p>Other differences from control values possibly but less certainly related to treatment:</p> <ul style="list-style-type: none"> - The more persistent occurrence of nonspecific signs after initiation of dosing, particularly in animals receiving 12 $\mu\text{mol/kg}$. - A higher incidence of nonaccidental deaths and abortions after this initiation of dosing. 	<p><u>6 $\mu\text{mol/kg}$</u> Digital anomalies, with a dose related trend in terms of the numbers affected and the degree of effect. The most noticeable pattern was for the greater effect on the more distal parts of the paw, particularly the hind paws, and the proportions of fetuses showing subtle reductions in the degree of claw ossification were 0, 71.4 and 100% respectively at 0, 6 and 12 $\mu\text{mol/kg}$.</p> <p><u>12 $\mu\text{mol/kg}$</u> Digital anomalies as observed above. Increased preimplantation loss and slight increase in early post-implantation Reduced litter size and litter wt.</p>

Table 7 Reproductive and Developmental Toxicity Studies (continued)

Species and Strain	No. of Animals/ Group	Period of Administration	Route of Administration				
				μmol/kg	mg/kg	Dams	Litters
Rabbit,	21 F	Days 6-28	p.o.	0		<p><u>Days 6-12</u> In all treatment groups treatment was associated with an initial decline in general condition indicated by reduced food intake, low faecal output, weight loss or suppression of weight gain. Recovery was evident within days of cessation of treatment. Palpable thickening of mammary glands with a corresponding significantly increased wt at termination was associated with initiation of treatment Day 6 of pregnancy, the response being less marked for animals dosed to Day 12 than to Day 28.</p> <p><u>Days 13-18</u> Initial decline in general condition as described above. No significant effect on mammary glands with regard to palpable thickening or increased wt.</p> <p><u>Days 6-28</u> See Days 6-12. Also, increased mammary weight gain.</p>	<p><u>Days 6-12</u> Litter size was reduced in all treatment groups compared to the controls. In groups treated from Day 6, the reduced litter size was mainly attributed to non-significant increases in post-implantation loss in combination with slightly higher values for preimplantation loss. Mean fetal wt lower in all treatment groups, the decrease was minimal for animals dosed Days 6-12. Lower values for litter wt in all treatment groups. Increased incidence of fetuses with an extra rib associated with initiation of treatment on Day 6.</p> <p><u>Days 13-18</u> Lower corpora lutea count contribution to a reduced litter size. Decreased mean fetal wt and lower values for litter wt. Treatment after Day 12 was associated with anomalies of the claws of almost all fetuses. The degree of effect appeared to be greater for this period of treatment than for the longer period (Days 6-28).</p> <p><u>Days 6-28</u> Reduced litter size, decrease in mean fetal wt and litter wt. Increased incidence of fetuses with an extra rib and anomalies of the claws occurred.</p>
New Zealand	21 F	Days 6-12		12	4.6		
White	20 F	Days 13-18		12	4.6		
	20 F	Days 6-28 of gestation (Seg II)		12	4.6		

Table 7 Reproductive and Developmental Toxicity Studies (continued)

Species and Strain	No. of Animals/ Group	Period of Administration	Route of Administration	µmol/kg	mg/kg	Dams	Litters
Rabbit, New Zealand White	10 F 10 F 20 F 19 F	Days 6 - 18 of gestation (Seg II)	p.o.	0 0 12 12	 4.6 4.6	Reduced body wt gain from start of dosing, compensated on Day 14 and reduced food consumption during the dosing period in treated animals. Increased wts due to hyperplasia of the mammary glands, in excess of the normal gestational one, in treated animals killed on Day 29. In animals killed on Day 35, no differences were observed with respect to wt or histomorphological appearance of the mammary glands.	Increased incidence of minor skeletal anomalies: disturbed development of distal phalanges, extra center of ossification between middle and distal phalanges in one or more digits in the fetuses of treated animals.
Rabbit, New Zealand White	9 F 9 F	Day 13 Days 13-18 of gestation	p.o.	12 12	4.6	The clinical signs observed and the effects on body wt gain, food consumption and mammary glands were essentially the same as those seen in earlier performed studies in the rabbit.	N/A
Rabbit, New Zealand White	8 F not pregnant	13 days (Days 0-12)	p.o.	0 1.2 12	 0.46 4.6	Reduced food consumption and decreased body wt during the dosing period in animals receiving 12 µmol/kg. No dose or compound related lesions were seen in the mammary glands which were all within the normal variation of mammary glands in non-pregnant rabbits. The results from this study and results of the above study in pregnant rabbits indicate that pregnancy has an insignificant effect on the pharmacokinetics of felodipine.	N/A

Table 7 Reproductive and Developmental Toxicity Studies (continued)

Species and Strain	No. of Animals/ Group	Period of Administration	Route of Administration	µmol/kg	mg/kg	Dams	Litters
Rabbit, New Zealand White	9 F 15 F 11 F	Days 6 - 18 of gestation (Seg II)	p.o.	0 1.2 122	0 0.46 4.6	<p><u>1.2 µmol/kg Plasma Concentration</u> Mean plasma levels and time course were very similar to those previously seen in non-pregnant rabbits, at the same dose level. The similarity of results on Days 13 and 18 indicated that there was no accumulation of the test compound.</p> <p><u>12 µmol/kg Signs of Maternal Toxicity</u> Body wt gain and food consumption were decreased during the first part of the dosing period.</p> <p><u>Mammary Glands</u> Treatment induced enlarged mammary glands. The mammary gland wts were significantly increased. The microscopic examination showed that this hyperplasia consisted of an increased volume of the glandular parenchyma due to an increased lobular size. The histological architecture, however, did not differ from that of the control animals.</p>	<p><u>12 µmol/kg (Day 29 of Gestation)</u> Fetal loss was slightly increased compared to the control group.</p> <p><u>1.2 and 12 µmol/kg Effects on the Phalanges</u> A dose related effect on both size and structure (reduced ossification) of the distal phalanges was seen. The most pronounced effect was on the 4th digit of the hind paws. This hypoplasia of the phalanges could be detected by external examination, but at the dose level of 12 µmol/kg only. In addition, 6 pups at 12 µmol/kg showed an extra center of ossification between the middle and distal phalanx of the pollex.</p>
Rabbit, New Zealand White	28 F3 32 F3	Days 6-18 of gestation (Seg II)	p.o.	0 12	0 4.6	<p>The effects on the treated dams sacrificed on Day 29 or allowed to litter were of the same type and magnitudes as in earlier studies and consisted of reduced body wt, reduced food consumption, and enlargement of the mammary glands, histologically characterized as a hyperplastic thickening due to an increased volume of the glandular parenchyma, in excess of the normal gestational hyperplasia. Changes in mammary glands were still present, although less marked in the dose group on Day 32 post parturition. The mean wts of the mammary glands were significantly increased for the treated dams on Days 29 and 32 post parturition. Since no disturbance of the post natal group of the pups was observed, the conclusion was drawn that there was no functional impairment of the mammary glands.</p>	<p>The findings in the fetuses were mainly disturbances of the development of the distal parts of the digits. The effects may be characterized as disturbances of the differentiation of the distal phalanges observed as decreased size of the distal phalanges and fusion of the outer phalanges in some digits. These anomalies of the digits could still be seen on Day 32 post parturition, and in addition, extra centres of ossification were also observed in some pups. The histological examination of the distal phalanges indicated a disturbed differentiation of the cartilaginous rudiments as a possible explanation of the observed effects.</p> <p>A slightly increased incidence of wavy ribs, an increased incidence of fetuses with an extra rib, and a decreased frequency of fetuses with variant sternebra among fetuses with normal number of ribs were observed in the dose group fetuses examined on Day 29. A few pups with gross malformations and visceral anomalies were also found in the treated group.</p>

²The dose level of 12 µmol/kg had been used as the highest dose in several previous segment II studies in the rabbit, and was known to cause maternal toxicity, enlargement of the mammary glands and minor skeletal effects in the offspring.

³Each group was divided into two subgroups. The animals in one of the subgroups were killed on Day 29 of pregnancy. The animals in the other one were allowed to litter normally and to rear their pups up to Day 32 post parturition, when all pups and dams were killed.

Table 7 Reproductive and Developmental Toxicity Studies (continued)

Species and Strain	No. of Animals/Group	Period of Administration	Route of Administration	µmol/kg	mg/kg	Dams	Litters
Rabbit, New Zealand White	1/I	3 F4	13	p.o.	12	4.6	<p><u>Clinical Observations</u> One dam dosed on Day 16 aborted on Day 20 and was killed on Day 23 for humanitarian reasons. One dam dosed on Day 17 was found dead on Day 29 of pregnancy. Two dams dosed on Days 15 and 18, respectively, aborted and one dam dosed on Day 15 gave premature birth. Retardation in body wt gain, reduced food consumption and reduced water intake were observed after the dosing day. The effects were most pronounced and lasted for the longest period of time in dams treated on Day 16 or 17 of gestation.</p> <p><u>Terminal Autopsy</u> Two dams dosed on Day 16 showed pale livers and distended caecum. Also two dams dosed on Day 17 showed pale livers.</p> <p><u>Effects on Distal Phalanges</u> No abnormalities of the fetal phalanges were observed after treatment of the dam on Day 13 or 18, while all fetuses showed reduced degree of ossification of one or more of the distal phalanges if treatment had occurred on Day 14, 15, 16 or 17 of gestation. The most pronounced effects were observed on Day 16, when all examined fetuses showed either reduced or very reduced size of one more of the distal phalanges. The hypoplasia of the phalanges induced on Day 16 was in general, in contrast to other days of treatment, visible at external examination before alizarin staining of the skeletons.</p>
	2/I	3 F	14		12	4.6	
	3/I	3 F	15		12	4.6	
	4/I	3 F	16		12	4.6	
	1/II	3 F	15		12	4.6	
	2/II	3 F	16		12	4.6	
	3/II	3 F	17		12	4.6	
	4/II	3 F	18		12	4.6	
Rabbit, New Zealand White	1	4 F	single, Day 16 of gestation	p.o.	-	-	THIS STUDY DID NOT DIFFERENTIATE BETWEEN DAMS AND LITTERS
	2	4 F			12	4.6	<u>Concentration of Felodipine in Maternal Plasma, Fetal Tissue and Amniotic Fluid</u>
	3	4 F			12	4.6	Plasma Concentrations: The highest felodipine concentrations (~350 nmol/L) were recorded at 4 hours after treatment; concentrations were still high at 12 hours (≥200 nmol/L) and 24 hours (≥100 nmol/L).
	4	3 F			12	4.6	Concentrations in Fetal Tissue: Mean concentrations in fetal tissue were 4-6 times lower than those seen in maternal plasma. The levels were virtually constant between 4 and 12 hours after treatment (~50 nmol/kg) and at 24 hours they were around 20 nmol/kg.
	5	4 F			12	4.6	Concentration in Amniotic Fluid: These were about 2 to 4 times lower than those in fetal tissue.
	6	6 F			125	4.6	<p><u>Histological Examination of the Limb Plates</u> Fetuses from dams treated with felodipine:</p> <p>2 and 4 Hours After Dose: Marked expansion of limb plates due to extensive mesenchymal edema. Widening of the thin-walled embryonal vessels, and the border between the cartilaginous primordium of phalanges and surrounding, undifferentiated mesenchymal less well-defined.</p> <p>8 Hours After Dose: Marked edema of limb plates and occasionally ruptures of marginal blood vessels, with hemorrhage.</p> <p>12 Hours After Dose: Mesenchymal edema of limb plates somewhat less pronounced. In addition to hemorrhages, occasional digital blister caused by cleavage vesicles between the mesenchymal and ectoderm.</p> <p>24 Hours After Dose: As at 12 hours after dose. Also occasional small necroses in the apex of digits, most often at the site of the 3rd phalange.</p>

⁴ The study was divided into two experiments, I and II, in which the dams were treated on different days of gestation.

⁵ The dose level of 12 µmol/kg had been used as the highest dose in several previous segment II studies in the rabbit, and was known to cause maternal toxicity, enlargement of the mammary glands and minor skeletal effects in the offspring.

PATIENT MEDICATION INFORMATION

READ THIS FOR SAFE AND EFFECTIVE USE OF YOUR MEDICINE

PrPLENDIL®

Felodipine Extended Release Tablets

Read this carefully before you start taking **Plendil** and each time you get a refill. This leaflet is a summary and will not tell you everything about this drug. Talk to your healthcare professional about your medical condition and treatment and ask if there is any new information about **Plendil**.

What is Plendil used for?

Plendil is used in adults to treat hypertension (high blood pressure).

It can be used together with other medicines that treat hypertension such as thiazide diuretics or beta-blockers.

How does Plendil work?

Plendil belongs to a group of medicines called “calcium channel blockers”. It helps to relax and widen blood vessels, improving blood flow and lowering blood pressure.

What are the ingredients in Plendil?

Medicinal ingredient: felodipine.

Non-medicinal ingredients: aluminum silicate, carnauba wax, hydrogen peroxide, hydroxypropyl cellulose, hydroxypropyl methylcellulose, iron oxide yellow, iron oxide red-brown (5 mg or 10 mg tablets), lactose anhydrous, microcrystalline cellulose, polyethylene glycol, polyoxyl 40 hydrogenated castor oil, propyl gallate, sodium stearyl fumarate, and titanium dioxide.

Plendil comes in the following dosage forms:

Extended release tablets: 2.5 mg, 5 mg, and 10 mg of felodipine.

Do not use Plendil if:

- you are allergic to felodipine or any of the other ingredients in Plendil.
- you allergic to other dihydropyridines (medicines used to treat hypertension and angina).
- you are pregnant, plan to become pregnant, or are able to become pregnant. You should tell your healthcare professional as soon as possible if you become pregnant while using Plendil.
- you are breast-feeding or plan to breast-feed.
- you are age 18 or younger.

To help avoid side effects and ensure proper use, talk to your healthcare professional before you take Plendil. Talk about any health conditions or problems you may have, including if you:

- have heart or blood vessel problems.
- have liver problems.
- have gum problems (e.g., gingivitis or periodontitis).
- have problems digesting certain milk sugars (e.g., galactose intolerance or glucose-galactose malabsorption). Plendil capsules contain lactose, a milk sugar. If you are unsure ask your healthcare professional.

Other warnings you should know about:

- **Driving and using machines:** Plendil can lower your blood pressure and affect your vision. Before you drive or do tasks that require special attention, wait until you know how you respond to Plendil.
- **Hypotension (low blood pressure):** Plendil may occasionally cause low blood pressure. Your healthcare professional will monitor your blood pressure, especially when you first start taking Plendil or if you are older than 65 years of age.
- **Oral hygiene:** Plendil can cause mild tenderness or swelling of the gums. This effect can be prevented or reversed with good dental care. Brush your teeth carefully and often with a soft toothbrush, and use dental floss daily. Massaging your gums regularly with a soft toothbrush will also help. If your gums do become tender, red or swollen, tell your healthcare professional.

Tell your healthcare professional about all the medicines you take, including any drugs, vitamins, minerals, natural supplements or alternative medicines.

Serious Drug Interactions

Do **not** take Plendil if you are already taking medications known as “strong inhibitors of CYP3A4”. These include:

- clarithromycin and erythromycin, medicines used to treat bacterial infections.
- “azole” drugs, medicines used to treat fungal infections (e.g., ketoconazole and itraconazole).
- ritonavir, a medicine used to treat human immunodeficiency virus (HIV).

Taking Plendil with any of these medicines may cause serious drug interactions. If you are unsure if you are taking any of these medicines, ask your healthcare professional.

The following may also interact with Plendil:

- alcohol, including alcohol containing products.
- antiarrhythmics, medicines used to treat or prevent irregular heartbeats (e.g., quinidine, flecainide and propafenone).
- other antibacterials, medicines used to treat bacterial infections (e.g., erythromycin and rifampin).
- medicines used to prevent the body from rejecting a transplanted organ, including the kidney or liver (e.g., cyclosporine and tacrolimus).
- warfarin, a medicine used to prevent blood clotting.
- beta-blockers, medicines used to treat high blood pressure (e.g., metoprolol).

- digoxin, a medicine used to treat heart failure and certain heartbeats problems.
- benzodiazepines, medicines used to treat anxiety, insomnia, and seizures.
- theophylline, a medicine used to relieve symptoms of asthma.
- cimetidine, a medicine used to treat stomach ulcers and reduce stomach acid.
- anticonvulsants, medicines used to prevent epilepsy or seizures (e.g., phenytoin, carbamazepine and phenobarbital).
- antidepressants, medicines used to treat depression (e.g., imipramine).
- terfenadine, a medicine used to treat allergies.
- St. John's Wort, a herbal medicine used to treat depression.
- grapefruit juice.

How to take Plendil:

- Take exactly as directed by your healthcare professional. If you are unsure, ask your healthcare professional.
- Swallow the Plendil tablets whole with a glass of water. Do **not** crush or chew on the tablets.
- Avoid taking Plendil with a meal rich in fat or carbohydrates. Breakfast foods which are rich in fat and/or carbohydrates include bacon, sausage, hash brown potatoes, and sugared cereals. Plendil may be taken with a light meal which is low in fat and carbohydrates (i.e., two slices of toast with cheese, cereal with milk, and orange juice).
- To help remember each dose, try to take Plendil with something you do regularly each day. For example, upon waking or at breakfast.
- Check with your healthcare professional if you want to drink alcohol (including wine with your meals) while you are taking Plendil. Drinking alcohol while on Plendil may make you feel dizzier than usual. Alcohol may also cause an uncomfortable drop in blood pressure.
- Do **not** eat grapefruit or drink grapefruit juice while on Plendil.
- It is important to take Plendil even when you feel well. A constant amount of drug is needed in your body to control your blood pressure.

Usual dose:

Your healthcare professional will decide the right dose of Plendil for you. This will depend on your health, age, and how you respond to Plendil.

The usual dose is 2.5 mg to 10 mg once a day. Even if your healthcare professional has prescribed 2 tablets a day, both should be taken at the same time, unless otherwise indicated. Do **not** adjust or stop your dose without first consulting your healthcare professional.

Overdose:

An overdose with Plendil may cause the following:

- feeling faint,
- feeling light-headed,
- feeling dizzy,
- very low blood pressure,
- slow heart rate.

If you think you, or a person you are caring for, have taken too much Plendil, contact a healthcare professional, hospital emergency department, regional poison control centre, or Health Canada's toll-free number, 1-844 POISON-X (1-844-764-7669) immediately, even if there are no signs or symptoms.

Missed Dose:

- If you miss a dose of Plendil and remember within 12 hours, take the missed dose as soon as possible. Then go back to your regular schedule.
- If you miss a dose of Plendil and it has been more than 12 hours when you remember, do not take the missed dose. Just take the next dose at your regular scheduled time.

Never take a double dose of Plendil to make up for missed tablets. If you are still unsure, check with your healthcare professional to see what you should do.

What are possible side effects from using Plendil?

These are not all the possible side effects you may have when taking Plendil. If you experience any side effects not listed here, tell your healthcare professional.

The side effects of Plendil may include:

- cold or flu-like symptoms;
- constipation;
- cough;
- dizziness;
- feeling of pins and needles;
- feeling tired;
- flushing or warm sensation;
- headache;
- joint stiffness;
- muscle cramps or pain;
- upset stomach, nausea, or indigestion.

Serious side effects and what to do about them			
Symptom / effect	Talk to your healthcare professional		Stop taking drug and get immediate medical help
	Only if severe	In all cases	
VERY COMMON			
Peripheral edema (swelling of the limbs): swelling in the legs, ankles or feet, feeling of heaviness, or skin appears tight or shiny.		X	
COMMON			
Bradycardia or tachycardia (abnormally slow or fast heart beat)		X	
Gastrointestinal (GI) problems: abdominal pain, diarrhea, vomiting, dry mouth, passing gas, acid reflux, fatigue, fever, itchiness, light-coloured stool, or yellowing of the skin or whites of eyes.		X	

Gum problems: swollen gums, gum tenderness and soreness, discomfort in the mouth, problems with oral hygiene, bleeding gums, bad breath, or overgrowth or receding gums.		X	
Hypotension (low blood pressure): dizziness, fainting, light-headedness, blurred vision, nausea, vomiting, or fatigue (may occur when you go from lying or sitting to standing up).		X	
Skin reactions: itchiness, dry skin, redness, rashes, blisters, raised red bumps, red spots, swelling, or increased sensitivity to sunlight.		X	
Visual disturbances	X		
UNCOMMON			
Anemia (decreased number of red blood cells): fatigue, loss of energy, pale complexion, shortness of breath, or weakness.		X	
Angioedema with breathing problems (swelling of tissue under the skin and difficulty breathing): difficulty breathing, swollen face, hands, feet, genitals tongue, or throat, diarrhea, nausea, or vomiting.			X
Heart problems: chest pain, chest pressure, shortness of breath, sweating, nausea, dizziness, fatigue, palpitations, irregular heartbeat, light-headedness, or fainting.		X	
RARE			
Allergic reaction: difficulty swallowing, difficulty breathing, wheezing, drop in blood pressure, nausea, vomiting, hives, rash, or swelling of the face, lips, tongue or throat.			X

If you have a troublesome symptom or side effect that is not listed here or becomes bad enough to interfere with your daily activities, tell your healthcare professional.

Reporting Side Effects

You can report any suspected side effects associated with the use of health products to Health Canada by:

- Visiting the Web page on Adverse Reaction Reporting (canada.ca/drug-device-reporting) for information on how to report online, by mail or by fax; or
- Calling toll-free at 1-866-234-2345.

NOTE: Contact your healthcare professional if you need information about how to manage your side effects. The Canada Vigilance Program does not provide medical advice.

Storage:

Store Plendil tablets at room temperature between 15°C to 30°C. To protect your Plendil tablets, keep them in the original blister packaging.

Keep out of reach and sight of children.

If you want more information about Plendil:

- Talk to your healthcare professional.
- Find the full Product Monograph that is prepared for healthcare professionals and includes the Patient Medication Information by visiting the Health Canada Drug Product Database website (<https://www.canada.ca/en/health-canada/services/drugs-health-products/drug-products/drugproduct-database.html>) or by calling the manufacturer at 1-833-905-2937.

This leaflet was prepared by Glenwood GmbH Pharmazeutische Erzeugnisse.

Last Revised: 2025-12-09