

Product Monograph
Including Patient Medication Information

^{Pr}CEFAZOLIN FOR INJECTION

Cefazolin sodium

Powder for Solution

For Intravenous, Intramuscular use

500 mg, 1 g and 10 g Cefazolin (as cefazolin sodium) per vial
USP

Antibiotic

Teva Canada Limited
30 Novopharm Court
Toronto, Ontario
M1B 2K9
Canada
www.tevacanada.com

Date of Authorization: 2025-12-29

Control Number: 297943

Recent Major Label Changes

<i>None at time of the most recent authorization</i>	NA
--	----

Table of Contents

Certain sections or subsections that are not applicable at the time of the preparation of the most recent authorized product monograph are not listed.

Recent Major Label Changes	2
Table of Contents	2
Part 1: Healthcare Professional Information	4
1 Indications	4
1.1 Pediatrics	5
1.2 Geriatrics	5
2 Contraindications	5
4 Dosage and Administration	5
4.2 Recommended Dose and Dosage Adjustment.....	5
4.3 Reconstitution	7
4.4 Administration.....	8
5 Overdose	9
6 Dosage Forms, Strengths, Composition, and Packaging	9
7 Warnings and Precautions	10
General.....	10
Gastrointestinal	10
Immune	10
Monitoring and Laboratory Tests	10
Overgrowth of Non-Susceptible Organisms	10
Renal	10
Sensitivity/Resistance.....	11
Skin	11
7.1 Special Populations.....	11
7.1.1 Pregnancy	11
7.1.2 Breastfeeding	11
7.1.3 Pediatrics	11
8 Adverse Reactions	12
9 Drug Interactions	12
9.4 Drug-Drug Interactions.....	12
9.7 Drug-Laboratory Test Interactions.....	12
10 Clinical Pharmacology	13
10.1 Mechanism of Action.....	13
10.3 Pharmacokinetics	13
11 Storage, Stability, and Disposal	14

Part 2: Scientific Information	15
13 Pharmaceutical Information.....	15
14 Clinical Trials	15
15 Microbiology.....	16
16 Non-Clinical Toxicology	17
Patient Medication Information	19

Part 1: Healthcare Professional Information

1 Indications

CEFAZOLIN FOR INJECTION (cefazolin sodium) is indicated in the treatment of the following infections when caused by susceptible strains of the listed organisms:

Respiratory Tract Infections caused by *Streptococcus pneumoniae*, *Klebsiella pneumoniae*, *Hemophilus influenzae*, *Staphylococcus aureus* (penicillin-sensitive and penicillin-resistant) and group A *beta-haemolytic streptococci*.

Urinary Tract Infections caused by *Escherichia coli*, *Proteus mirabilis*, *Klebsiella pneumoniae* and some strains of enterobacter, and enterococci. See note below.

Skin and Soft Tissue Infections caused by *Staphylococcus aureus* (penicillin-sensitive and penicillin-resistant), group A *beta-haemolytic streptococci* and other strains of streptococci.

Bone and Joint Infections caused by *Staphylococcus aureus*.

Septicemia caused by *Streptococcus pneumoniae*, *Staphylococcus aureus* (penicillin-sensitive and penicillin resistant), *Proteus Mirabilis*, *Escherichia coli* and *Klebsiella pneumoniae*. See note below.

Endocarditis caused by *Staphylococcus aureus* (penicillin-sensitive and penicillin-resistant) and group A *beta haemolytic streptococci*.

Determine susceptibility of the causative organism to cefazolin sodium, by performing appropriate culture and susceptibility studies should be performed. (See [15 Microbiology](#) for disc susceptibility tests and dilution techniques).

Note: Most strains of *Enterococci*, indole positive *Proteus (P. vulgaris)*, *Enterobacter cloacae*, *Morganella morganii*, *Providencia rettgeri* and methicillin-resistant *Staphylococci* are resistant. *Serratia*, *Pseudomonas*, and *Acinetobacter calcoaceticus* (formerly *Mima* and *Herellea* species) are almost uniformly resistant to cefazolin. (See [15 Microbiology](#)).

Perioperative Prophylaxis: In patients undergoing potentially contaminated surgical procedures, and in patients in whom infection would pose a serious risk (e.g., during open-heart surgery and prosthetic arthroplasty), the preoperative, intraoperative and postoperative administration of CEFAZOLIN FOR INJECTION may reduce the incidence of certain post-operative infections.

Identification of the causative organisms should be made by culture should signs of infection occur, so that appropriate therapy may be instituted.

To reduce the development of drug-resistant bacteria and maintain the effectiveness of CEFAZOLIN FOR INJECTION and other antibacterial drugs, CEFAZOLIN FOR INJECTION should be used only to treat infections that are proven or strongly suspected to be caused by susceptible bacteria. When culture and susceptibility information are available, they should be considered in selecting or modifying antibacterial therapy. In the absence of such data, local epidemiology and susceptibility patterns may contribute to the empiric selection of therapy.

1.1 Pediatrics

Pediatrics (< 1 month): The safety and efficacy of CEFAZOLIN FOR INJECTION in pretermatures and infants under one month of age has not been established (see [4.2 Recommended Dose and Dosage Adjustment, Pediatric Use](#)).

1.2 Geriatrics

Geriatrics: No data are available to Health Canada; therefore, Health Canada has not authorized an indication for geriatric use.

2 Contraindications

CEFAZOLIN FOR INJECTION is contraindicated in patients who are hypersensitive to this drug or to any ingredient in the formulation, including any non-medicinal ingredient, or component of the container, or to the cephalosporin group of antibiotics. For a complete listing, see [6 Dosage Forms, Strengths, Composition, and Packaging](#).

4 Dosage and Administration

4.2 Recommended Dose and Dosage Adjustment

After reconstitution CEFAZOLIN FOR INJECTION may be administered either intramuscularly or intravenously. In both cases total daily dosages are the same.

Adults:

Adult Dosage Guide

Type of Infection	Dose	Frequency
Mild infections caused by susceptible Gram-positive cocci	250 mg to 500 mg	Every 8 hours
Acute, uncomplicated urinary tract infections*	1 g	Every 12 hours
Moderate to severe infections	500 mg to 1 g	Every 6 to 8 hours

*This dosage recommendation applies to intramuscular use. The efficacy of cefazolin sodium when administered intravenously at 12-hour intervals has not been established.

CEFAZOLIN FOR INJECTION has been administered in dosages of 6 g per day in serious infections such as endocarditis.

Treatment should be continued for at least 10 days in beta-haemolytic streptococcal infections to minimize possible complications associated with the disease.

Dosage in Patients with Reduced Renal Function:

After an initial loading dose appropriate to the severity of the infection, the following reduced dosage schedule is recommended:

Dosage Guide for Patients with Renal Impairment

Creatinine Clearance (mL/s)	Serum Creatinine (mMol/L)	Dosage
≤0.91	≥140	250 mg to 1 g every 6-12 hours
0.58-0.90	141-273	250 mg to 1 g every 8-12 hours
0.18-0.57	274-406	125 mg to 500 mg every 12 hours
≤0.17	≥407	125 mg to 500 mg every 18 hours

Perioperative Prophylactic Use:

The recommended dosage regimen to prevent postoperative infection in contaminated or potentially contaminated surgery is:

- a. One gram intravenously or intramuscularly administered ½ hour to 1 hour prior to the start of surgery so that at the time of the initial surgical incision adequate antibiotic levels are present in the serum and tissues.
- b. For lengthy operative procedures (e.g., 2 hours or more) 0.5-1.0 g administered intravenously or intramuscularly during surgery. (Administration should be modified according to the duration of the operative procedure and the time of greatest exposure to infective organisms.)
- c. Postoperatively, 0.5-1.0 gram intravenously or intramuscularly every 6 to 8 hours for 24 hours postoperatively. The prophylactic administration of CEFAZOLIN FOR INJECTION maybe continued for 3 to 5 days following the completion of surgery in which the occurrence of infection may be particularly devastating (e.g., open-heart surgery and prosthetic arthroplasty).

Pediatric Use:

A total daily dosage of 25 to 50 mg per kg (approximately 10 to 20 mg per pound) of body weight, divided into three or four equal doses, is effective for most mild to moderately severe infections in children.

For severe infections total daily dosage maybe increased to 100 mg per kg (45 mg per pound) of body weight. The use of CEFAZOLIN FOR INJECTION in prematures and in infants under one month is not recommended since the safety for use in these patients has not been established (see [1.1 Pediatrics](#)).

Pediatric Dosage Guide – 25 mg/kg/day

Weight		25 mg/kg/day Divided Into 3 Doses		25 mg/kg/day Divided into 4 doses	
lb	kg	Approximate Single Dose mg/q8h	Volume Needed of 125 mg/mL* Solution	Approximate Single Dose mg/q6h	Volume Needed of 125 mg/mL* Solution
10	4.5	40 mg	0.35 mL	30 mg	0.25 mL
20	9	75 mg	0.60 mL	55 mg	0.45 mL
30	13.6	115 mg	0.90 mL	85 mg	0.70 mL
40	18.1	150 mg	1.20 mL	115 mg	0.90 mL
50	22.7	190 mg	1.50 mL	140 mg	1.10 mL

* 125 mg/mL concentration may be obtained by reconstituting the 500 mg vial with 3.8 mL of diluent.

Pediatric Dosage Guide-50 mg/kg/day

Weight		50 mg/kg/day Divided Into 3 Doses		50 mg/kg/day Divided into 4 doses	
lb	kg	Approximate Single Dose mg/q8h	Volume Needed of 225 mg/mL* Solution	Approximate Single Dose mg/q6h	Volume Needed of 225 mg/mL* Solution
10	4.5	75 mg	0.35 mL	55 mg	0.25 mL
20	9	150 mg	0.70 mL	110 mg	0.50 mL
30	13.6	225 mg	1.00 mL	170 mg	0.75 mL
40	18.1	300 mg	1.35 mL	225 mg	1.00 mL
50	22.7	375 mg	1.70 mL	285 mg	1.25 mL

* 225 mg/mL concentration may be obtained by reconstituted the 500 mg vial with 2.0 mL of diluent.

Treatment with 60 percent of the normal daily dose may be administered in divided doses every 12 hours to children with mild to moderate renal impairment (Ccr 0.67-1.17 mL/s). Children with moderate to severe renal impairment (Ccr 0.33- 0.87 mL/s) should be given 25 percent of the normal daily dose in equally divided doses every 12 hours, and children with severe renal impairment (Ccr 0.08-0.33 mL/s) should receive 10 percent of the normal daily dose every 24 hours.

All dosage recommendations apply after an initial loading dose.

4.3 Reconstitution

Reconstituted Solutions

Parenteral drug products should be SHAKEN TO DISSOLVE ALL POWDER when reconstituted, and inspected visually for particulate matter prior to administration. The drug solutions should be discarded if particulate matter is evident in reconstituted fluids.

Reconstituted solutions may range in colour from pale yellow to yellow without a change in potency.

(1) For Intramuscular Injection:

Single Dose Vials:

Reconstitute according to the table which follows. SHAKE TO DISSOLVE ALL POWDER.

Single Dose Vial Reconstitution Table

Strength (mg)	Diluent	Volume to be Added to Vial (mL)	Approximate Available Volume (mL)	Nominal Concentration (mg/mL)
500	Sodium Chloride Injection	2	2.2	225
	or Sterile Water for Injection	3.8	4	125
1000	Sterile Water for Injection	2.5	3	334

(2) For Direct Intravenous (bolus) Injection:

Single Dose Vial:

Reconstitute as directed above. SHAKE TO DISSOLVE ALL POWDER. A minimum of 10 mL of Sterile Water for Injection should be used to dilute the reconstituted solution.

Pharmacy Bulk Vial:

Pharmacy Bulk Vials should be used for intravenous use only. Add, according to the table below, Sterile Water for Injection or Sodium Chloride Injection. SHAKE TO DISSOLVE ALL POWDER.

Pharmacy Bulk Vial Reconstitution Table

Strength	Amount of Diluent	Approximate Available Volume	Approximate Concentration
10 grams	45 mL	50 mL	200 mg/mL
	96 mL	100 mL	100 mg/mL

The vial is intended for single puncture and multiple dispensing, and the vial contents should be used within 8 hours.

(3) For intermittent or continuous intravenous infusion, reconstituted CEFAZOLIN FOR INJECTION may be further diluted as follows:

Single Dose Vials:

Reconstitute according to the Single Dose Vial Reconstitution Table above. SHAKE TO DISSOLVE ALL POWDER. Further dilute the reconstituted CEFAZOLIN FOR INJECTION to 50 to 100 mL in one of the following solutions:

Sodium Chloride Injection 0.9%

Dextrose Injection 5% or 10%

Dextrose 5% in Lactated Ringer's Injection

Dextrose 5% and Sodium Chloride Injection 0.9% (also may be used with Dextrose 5% and Sodium Chloride Injection 0.45% or 0.2%)

Lactated Ringer's Injection

Ringer's Injection

Sodium Bicarbonate 5% in Sterile Water for Injection

Pharmacy Bulk Vial:

Reconstitute according to the Pharmacy Bulk Vial Reconstitution Table. SHAKE TO DISSOLVE ALL POWDER. Further dilute aliquots in 50 to 100 mL of Sterile Water for Injection or one of the solutions listed above.

The further diluted solutions above should be used within 24 hours at room temperature or 72 hours under refrigeration from the time of initial puncture.

4.4 Administration

Note: See [4.3 Reconstitution](#) for reconstitution and dilution directions.

Intramuscular Administration:

Inject the reconstituted solution into a large muscle mass. Pain on injection of CEFAZOLIN FOR INJECTION occurs infrequently.

Intravenous Administration:

Direct (bolus) injection: Inject the appropriately diluted reconstituted solution slowly over 3 to 5 minutes directly into a vein or through tubing for patients receiving parenteral fluids. (See list of solutions for intravenous infusion in [4.3 Reconstitution](#).)

Intermittent or Continuous Infusion: The reconstituted solution can be administered along with primary intravenous fluid management programs in a volume control set or in a separate secondary i.v. bottle. (See list of solutions for intravenous infusion in [4.3 Reconstitution](#).)

5 Overdose

There is a lack of experience with acute CEFAZOLIN FOR INJECTION overdose. Supportive therapy should be instituted according to symptoms in cases of suspected overdose.

For the most recent information in the management of a suspected drug overdose, contact your regional poison control centre or Health Canada’s toll-free number, 1-844 POISON-X (1-844-764-7669).

6 Dosage Forms, Strengths, Composition, and Packaging

Table 1: Dosage Forms, Strengths, and Composition

Route of Administration	Dosage Form/Strength/Composition	Non-Medicinal Ingredients
Intravenous	Powder For Solution 500 mg, 1 g and 10 g cefazolin (as cefazolin sodium) per vial	None
Intramuscular	Powder For Solution 500 mg and 1 g cefazolin (as cefazolin sodium) per vial	None

Description

CEFAZOLIN FOR INJECTION is supplied as a powder in clear glass vials equivalent to 500 mg or 1 g of cefazolin. CEFAZOLIN FOR INJECTION is also available as a Pharmacy Bulk Vial in vials equivalent to 10 g of cefazolin.

The availability of the pharmacy bulk vial is intended for hospitals with a recognized IV admixture program.

Composition

Each vial contains cefazolin sodium. Each gram of cefazolin sodium contains 48 mg of sodium. CEFAZOLIN FOR INJECTION does not contain any preservative.

7 Warnings and Precautions

General

In beta-haemolytic streptococcal infections, treatment should be continued for at least 10 days, to minimize possible complications associated with the disease.

Gastrointestinal

In patients with a history of lower gastrointestinal disease, particularly colitis, CEFAZOLIN FOR INJECTION should be prescribed with caution.

Immune

CEFAZOLIN FOR INJECTION should be used with caution in penicillin-allergic patients. There is clinical evidence of partial cross-allergenicity of the penicillins and the cephalosporins. There are instances of patients who have had reactions to both penicillins and cephalosporins (including fatal anaphylaxis after parenteral use). Clinical and laboratory evidence of partial cross-allergenicity of the two drug classes exists.

CEFAZOLIN FOR INJECTION should be administered cautiously and then only when absolutely necessary to any patient who has demonstrated allergy, particularly to drugs. Immediate emergency treatment with epinephrine is indicated for serious anaphylactoid reactions. As indicated, oxygen, intravenous steroids, and airway management, including intubation, should also be employed.

Monitoring and Laboratory Tests

Clinitest^R tablets solution, but not enzyme-based tests such as Clinistix^R and Tes-Tape^R, may falsely indicate glucose in the urine of patients on cefazolin.

Positive direct and indirect Coombs' tests have been reported during treatment with cefazolin. These may also occur in neonates whose mothers received cephalosporins before delivery. The clinical significance of this effect has not been established.

Overgrowth of Non-Susceptible Organisms

The overgrowth of non-susceptible organisms may result from the prolonged use of CEFAZOLIN FOR INJECTION. It is essential that the patient be carefully observed.

Renal

Caution should be exercised in treating patients with pre-existing renal damage although CEFAZOLIN FOR INJECTION has not shown evidence of nephrotoxicity.

Patients with low urinary output due to impaired renal function should be administered reduced daily dosages of CEFAZOLIN FOR INJECTION (see [4.2 Recommended Dose and Dosage Adjustment, Dosage in Patients with Reduced Renal Function](#)). Blood levels of CEFAZOLIN FOR INJECTION in dialysis patients remain fairly high and should be monitored.

Probenecid may decrease renal tubular secretion of cefazolin when used concurrently with CEFAZOLIN FOR INJECTION, resulting in increased and prolonged cefazolin blood levels (see [9.4 Drug-Drug Interactions](#)).

Seizures may occur with the administration of CEFAZOLIN FOR INJECTION, particularly in patients with renal impairment when the dosage is not reduced appropriately. Discontinue CEFAZOLIN FOR INJECTION if seizures occur or make appropriate dosage adjustments in patients with renal impairment. Anticonvulsant therapy should be continued in patients with known seizure disorders.

Sensitivity/Resistance

Development of Drug Resistant Bacteria

Prescribing CEFAZOLIN FOR INJECTION in the absence of a proven or strongly suspected bacterial infection is unlikely to provide benefit to the patient and risks the development of drug-resistant bacteria.

Skin

Severe Cutaneous Adverse Reactions

Severe cutaneous adverse reactions (SCAR) such as acute generalized exanthematous pustulosis (AGEP), drug reaction with eosinophilia and systemic symptoms (DRESS), Stevens-Johnson syndrome (SJS), and toxic epidermal necrolysis (TEN) have been reported in association with beta-lactam treatment. When SCAR is suspected, CEFAZOLIN FOR INJECTION should be discontinued and appropriate therapy and/or measures should be taken.

7.1 Special Populations

7.1.1 Pregnancy

The safety of the use of CEFAZOLIN FOR INJECTION during pregnancy has not been established.

7.1.2 Breastfeeding

Very low concentrations of cefazolin are found in the milk of nursing mothers. CEFAZOLIN FOR INJECTION should be administered with caution to a nursing woman.

7.1.3 Pediatrics

The safety of the use of CEFAZOLIN FOR INJECTION in prematures and infants under one month of age has not been established (see [4.2. Recommended Dose and Dosage Adjustment, Pediatric Use](#)).

8 Adverse Reactions

The following adverse reactions have been reported:

Gastrointestinal: Diarrhea, oral candidiasis (oral thrush), vomiting, nausea, stomach cramps, anorexia. During antibiotic treatment symptoms of pseudo membranous colitis can appear. There have been rare reports of nausea and vomiting. There have been reports of pseudo membranous colitis with the use of cephalosporins. It is therefore important to consider its diagnosis in patients who develop diarrhea in association with antibiotic use.

Haematologic: Cases of impaired blood coagulation have been reported during cefazolin treatment. Patients with risk factors causing vitamin K deficiency or affecting other clotting mechanisms (parenteral alimentation, dietary deficiencies, reduced hepatic and renal function) may be at increased risk of bleeding during cefazolin treatment. Monitoring of patients and vitamin K supplementation should be considered as indicated.

Blood clotting disturbances including prolongation of the prothrombin time, increased INR and bleeding. Neutropenia, anemia, leukopenia, thrombocytopenia, positive direct and indirect antiglobulin (Coombs') tests.

Hepatobiliary and Renal: Without clinical evidence of renal or hepatic impairment transient increases in AST (SGOT), ALT (SGPT), BUN and alkaline phosphatase levels have been observed. Transient hepatitis and cholestatic jaundice have been reported rarely, as with some penicillins and some other cephalosporins.

Immune system disorders: Allergic reactions occur infrequently and include: anaphylaxis, eosinophilia, itching, drug fever, skin rash.

Local Reactions: Phlebitis at the site of injection has occurred rarely. Infrequently there is pain at the site of injection following intramuscular injection. Some induration has been reported.

Other: Vulvar pruritus, genital moniliasis, vaginitis and anal pruritus.

9 Drug Interactions

9.4 Drug-Drug Interactions

The renal tubular secretion of CEFAZOLIN FOR INJECTION may be decreased when probenecid is used concurrently, resulting in increased and prolonged CEFAZOLIN FOR INJECTION blood levels.

9.7 Drug-Laboratory Test Interactions

Clinitest^R tablets solution, but not enzyme-based tests such as Clinistix^R and Tes-Tape^R, may falsely indicate glucose in the urine of patients on cefazolin.

Positive direct and indirect Coombs' tests have been reported during treatment with cefazolin. These may also occur in neonates whose mothers received cephalosporins before delivery. The clinical significance of this effect has not been established.

10 Clinical Pharmacology

10.1 Mechanism of Action

CEFAZOLIN FOR INJECTION is a cephalosporin antibiotic for parenteral administration. Cefazolin exerts its bactericidal effect by inhibiting bacterial cell wall synthesis.

10.3 Pharmacokinetics

Cefazolin is about 85% bound to serum protein. The peak level in serum is approximately 32-42 mg/mL after an intramuscular (i.m.) injection of 500 mg. Over 80% of injected cefazolin is excreted in the urine during the first 24 hours after i.m. injection; most is excreted during the first 4-6 hours.

The blood levels of cefazolin listed on the following tables were determined following intramuscular and intravenous administration.

Serum Concentration (mg/mL) Following Administration:

(Time After Intravenous Injection in Minutes)

	5	15	30	60	120	240
Cefazolin 1 g	188.4	135.8	106.8	73.7	45.6	16.5

(Time After Intramuscular Injection in Hours)

Cefazolin	½	1	2	4	6	8
1 g	65.8	68.3	60.6	29.3	11.2	6.5
500 mg	36.2	36.8	37.9	15.5	6.3	3
250 mg	15.5	17	13	5.1	2.5	<1.5

The serum half-life is approximately 1.8 hours following intravenous administration and 2.0 hours after intramuscular administration.

The mean peak serum levels of cefazolin in hospitalized patients are approximately equivalent to those seen in normal volunteers.

Healthy volunteers received a continuous intravenous infusion of 3.5 mg/kg for 1 hour (approximately 250 mg) and 1.5 mg/kg hourly for the next two hours (approximately 100 mg). A steady serum level of 28 mg/mL was attained at the third hour.

Cefazolin levels in synovial fluid and serum are similar four hours after drug administration.

Levels in cord blood are equivalent to 40% of those found in maternal blood.

In patients without obstructive biliary disease, serum levels of cefazolin can be up to five times lower than bile levels of cefazolin. However, bile levels of cefazolin are considerably lower than serum levels in patients with obstructive biliary disease.

Cefazolin is excreted unchanged in the urine. Approximately 60% of the drug is excreted in the first six hours, and this increases to 70%-80% within 24 hours. Peak urine concentrations of approximately 2400 mcg/mL and 4000 mcg/mL are achieved following intramuscular doses of 500 mg and 1 gram, respectively.

11 Storage, Stability, and Disposal

CEFAZOLIN FOR INJECTION should be stored between 15° - 30°C, protected from light.

Reconstituted CEFAZOLIN FOR INJECTION may be stored for 24 hours at controlled room temperature not exceeding 25°C, or for 72 hours under refrigeration (2 to 8°C), protected from light.

CEFAZOLIN FOR INJECTION solution reconstituted with bacteriostatic diluent and used for intramuscular administration as multiple-dose containers should be used within 7 days when stored under refrigeration.

The Pharmacy Bulk Vial is intended for multiple dispensing for intravenous use only, employing a single puncture. Following reconstitution, the solution should be dispensed and diluted for use within eight hours. Any unused reconstituted solution should be discarded after eight hours.

Part 2: Scientific Information

13 Pharmaceutical Information

Drug Substance:


Non-proprietary name of the drug substance: Cefazolin sodium

Chemical name: Monosodium (6R, 7R)-3-[[5-methyl-1,3,4-thiadiazol-2-yl]thio]methyl]-8-oxo-7-[2-(1H-tetrazol-1-yl)acetamido]-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylate.

Molecular formula: $C_{14}H_{13}N_8NaO_4S_3$

Molecular weight: 476.5

Structural formula:



Physicochemical properties:

Cefazolin sodium is a white to off-white, practically odourless, solid. The drug is freely soluble in water and very slightly soluble in alcohol. It is practically insoluble in chloroform and ether. The pH is between 4.5 and 6.0, in an aqueous solution containing 100 mg of cefazolin per millilitre.

14 Clinical Trials

Clinical trial information is not available.

15 Microbiology

Cefazolin Activity Against Clinical Isolates

	No. of Strains		Cumulative Percentage Susceptible to Strains Indicated Concentration (µg/mL)				
		<0.05	<0.1 - 0.78	1.56 - 3.13	6.25 - 12.5	25 – 50	100
<i>S. aureus</i>	700	0.14	59.1	90.6 – 92.4*	97.3	99.7	99.9
<i>S. pyogenes</i>	5	80+	100				
<i>S. faecalis</i>	2				50	100	
<i>S. pneumoniae</i>	6	100+					
<i>E. coli</i>	484		8.7	67.9	92.1	95.9	97.7
<i>P. mirabilis</i>	30			50	86.7	90	90
<i>K. pneumoniae</i>	138		2.9	53.6	73.2	91.3	93.5
<i>Enterobacter</i>	31			6.5	29.0	64.5	77.4
<i>H. influenzae</i>	30			13.3	70.0	100	
<i>N. gonorrhoeae</i>	13		38.5	100			
<i>Shigella</i> spp.	2			50	50	100	
<i>Salmonella</i> spp.	8			100			
<i>Staphylococci</i> (coagulase - negative)	295		66	82	90	93	100

* Reported as 3.13-6.25 µ.g/mL

+ Reported as ≤ 0.1 µ.g /mL

Disc Susceptibility Tests:

The following criteria should be used to interpret tests using a standardized 30 µg cephalosporin-class disc:

Zones of 18 mm or greater indicate that the tested organisms are susceptible and are likely to respond to therapy. Zones of 15 to 17 mm indicate organisms of intermediate susceptibility which may be susceptible if high dosage is used or if the infection is confined to tissues and fluids (e.g., urine) in which high antibiotic levels are attained. Zones of 14 mm or less are produced by resistant organisms.

The cephalothin disc should not be used for testing susceptibility to other cephalosporins.

Dilution Techniques:

If the minimal inhibitory concentration (MIC) for cefazolin is not more than 16 mg/mL, then a bacterial isolate may be considered susceptible. If the MIC is equal to or greater than 64 mg/mL, organisms are considered to be resistant.

The ranges of MIC for the control strains were:

E. coli ATCC 25922 1.0-4.0 mg/mL
S. aureus ATCC 25923 0.25-1.0 mg/mL

16 Non-Clinical Toxicology

General toxicology

Acute Toxicity:

Parenteral and oral cefazolin demonstrated low toxicity in rodents, canines and rabbits tested in acute toxicity studies.

Species	Route Of Administration	LD ₅₀ LD (g/kg)
mice	intravenous	≥3.9
	intraperitoneal	≥4.0
	subcutaneous	7.6
	oral	>11.0
rats	intravenous	≥3.0
	intraperitoneal	7.4
	subcutaneous	>10
	oral	>11.0
rabbits	intravenous	>2.0
dogs	intravenous	>2.0

Subacute and Chronic Toxicity:

Rats and dogs were studied in subacute and chronic parenteral toxicity of cefazolin. Rats were treated for 3 and 6 months subcutaneously and for one month intraperitoneally. The highest doses ranged from 2000 mg/kg per day in the 6 month study to 4000 mg/kg per day in the 1 and 3 month studies. Anemia was the only significant abnormality attributable to s.c. drug administration. In all experiments there was a definite dose-related depression of SGPT levels.

Leukocytosis and hypererythropoiesis accompanied the anemia, which was probably related to hemorrhaging at the injection site.

The lowering of the SGPT was dependent upon both the dose and the duration of treatment. This was not statistically significant at the low doses and was reversible upon withdrawal of the drug.

Equivalent chronic studies in dogs produced similar results: at the higher doses there was a fall in SGPT and frank anemia resulted from high subcutaneous doses. Dogs treated intravenously did not develop the anemia indicating that it was probably associated with hemorrhaging at the site of injection.

Reproductive and developmental toxicology

Rabbits and mice were administered cefazolin in doses of 240 mg/kg/day and 2400 mg/kg/day. No teratologic effects were observed. No adverse effects on mating, fertility, gestation, delivery and

lactation were observed in rats administered 2000 mg/kg per day. Baby rats whose mothers were injected with 1200 mg/kg/day of cefazolin prior to delivery and throughout lactation were observed and there was no effect on the birth, or peri- and postnatal development.

Special toxicology

Nephrotoxicity:

The nephrotoxicity of cefazolin was studied following intravenous injections of rabbits and subcutaneous injections of mice and rats. The mean nephrotoxic intravenous dose in rabbits was between 300 and 400 mg/kg/day. No evidence of renal damage was produced when cefazolin was injected subcutaneously into mice at a dose of 8 g/kg/day for up to 3 days and into rats at a dose of 4 g/kg/day for up to 7 days.

Patient Medication Information

READ THIS FOR SAFE AND EFFECTIVE USE OF YOUR MEDICINE

PrCEFAZOLIN FOR INJECTION

This Patient Medication Information is written for the person who will be receiving **CEFAZOLIN FOR INJECTION**. This may be you or a person you are caring for. Read this information carefully. Keep it as you may need to read it again.

This Patient Medication Information is a summary. It will not tell you everything about this medication. If you have more questions about this medication or want more information about **CEFAZOLIN FOR INJECTION**, talk to a healthcare professional.

What CEFAZOLIN FOR INJECTION is used for:

CEFAZOLIN FOR INJECTION is used to treat certain bacterial infections in the:

- respiratory tract;
- urinary tract;
- bones and joints;
- skin and soft tissues;
- bloodstream (septicemia);
- inner lining of the heart's chambers and valves.

CEFAZOLIN FOR INJECTION can also be used to prevent infections, before and after surgery.

Antibacterial drugs like CEFAZOLIN FOR INJECTION treat **only** bacterial infections. They do not treat viral infections, such as the common cold.

How CEFAZOLIN FOR INJECTION works:

CEFAZOLIN FOR INJECTION is an antibiotic that belongs to a group of medicines called cephalosporins. It kills bacteria by preventing them from forming their cell walls.

The ingredients in CEFAZOLIN FOR INJECTION are:

Medicinal ingredient: cefazolin sodium

Non-medicinal ingredients: none

CEFAZOLIN FOR INJECTION comes in the following dosage form:

Sterile powder for solution: 500 mg, 1 g and 10 g cefazolin per vial.

Do not receive CEFAZOLIN FOR INJECTION if:

- you have had an allergic reaction to:
 - cefazolin sodium, any other ingredient used to prepare CEFAZOLIN FOR INJECTION, or components of the container; or
 - other cephalosporin antibiotics.

To help avoid side effects and ensure proper use, talk to your healthcare professional before you receive CEFAZOLIN FOR INJECTION. Talk about any health conditions or problems you may have,

including if you:

- are allergic to or react badly to penicillins or other antibiotics.
- have a history of bowel disease, particularly colitis (inflammation of the colon).
- have kidney problems.
- are pregnant, think you might be pregnant or plan to become pregnant.
- are breast-feeding or planning to breastfeed. CEFAZOLIN FOR INJECTION may pass into breast milk.
- have previously had issues with blood clotting (bleeding), dietary deficiency (such as vitamin k deficiency); or have had liver / kidney issues or dietary deficiencies that have caused low levels of vitamin K.

Other warnings you should know about:

- CEFAZOLIN FOR INJECTION may interfere with some blood and urine test results. If you are given a blood or urine test, tell your healthcare professional that you are receiving treatment
- with CEFAZOLIN FOR INJECTION.

Tell your healthcare professional about all the medicines you take, including any drugs, vitamins, minerals, natural supplements or alternative medicines.

The following may interact with CEFAZOLIN FOR INJECTION:

- Probenecid, used to treat gout.

How CEFAZOLIN FOR INJECTION is given:

- CEFAZOLIN FOR INJECTION will be given to you by a healthcare professional as an injection into either a vein or a muscle.
- Although you may feel better early in treatment, CEFAZOLIN FOR INJECTION should be used exactly as directed.
- Misuse or overuse of CEFAZOLIN FOR INJECTION could lead to the growth of bacteria that will not be killed by CEFAZOLIN FOR INJECTION (resistance). This means that CEFAZOLIN FOR INJECTION may not work for you in the future.
- Do not share your medicine.

Usual dose:

Your healthcare professional will decide how much CEFAZOLIN FOR INJECTION to give you and how often.

Overdose:

If you think you, or a person you are caring for, have been given too much CEFAZOLIN FOR INJECTION, contact a healthcare professional, hospital emergency department, or regional poison control centre or Health Canada's toll-free number, 1-844 POISON-X (1-844-764-7669) immediately, even if there are no signs or symptoms.

Missed dose:

If you miss a scheduled treatment, talk to your healthcare professional as soon as possible to find out what to do.

Possible side effects from using CEFAZOLIN FOR INJECTION:

These are not all the possible side effects you may have when receiving CEFAZOLIN FOR INJECTION. If you experience any side effects not listed here, tell your healthcare professional.

Side effects with CEFAZOLIN FOR INJECTION may include:

- diarrhea, nausea, vomiting
- stomach cramps, loss of appetite
- rash, itching
- pain, tenderness or a hardened mass at the injection site
- vaginal and anal itching

Serious side effects and what to do about them

Frequency/Side Effect/Symptom	Talk to your healthcare professional		Stop taking this drug and get immediate medical help
	Only if severe	In all cases	
Common			
Thrombocytopenia (high levels of platelets): burning, redness, throbbing, numbness and/or tingling in the hands and feet, headache, dizziness, weakness, fainting, chest pain, vision changes		√	
Pseudomembranous colitis (advanced inflammation of the colon): watery, bloody diarrhea, mucus in the stool, abdominal cramps and pain, fever			√
Phlebitis (swelling of a vein near the injection site): pain, tenderness, redness		√	
Rare			
Severe Cutaneous Adverse Reactions (SCAR) (severe skin reactions that may also affect other organs): <ul style="list-style-type: none"> • Skin peeling, scaling, or blistering (with or without pus) which may also affect your eyes, mouth, nose or genitals, itching, severe rash, bumps under the skin, skin pain, skin colour changes (redness, yellowing, purplish) • Swelling and redness of eyes or face 			√

Frequency/Side Effect/Symptom	Talk to your healthcare professional		Stop taking this drug and get immediate medical help
	Only if severe	In all cases	
<ul style="list-style-type: none"> • Flu-like feeling, fever, chills, body aches, swollen glands, cough • Shortness of breath, chest pain or discomfort 			
Unknown			
Allergic reaction: rash, hives, swelling of the face, lips, tongue or throat, difficulty swallowing or breathing			√
Anemia (low levels of red blood cells): fatigue, pale skin, loss of energy, weakness, shortness of breath		√	
Bleeding disorders: bruising or bleeding longer than usual, black stool or blood in stool, bleeding in surgical wounds or in your catheter, severe headache, nausea and vomiting, confusion, dizziness, seizures, loss of consciousness.		√	
Seizures (fits): uncontrollable shaking with or without loss of consciousness. You are more likely to experience this if you have kidney problems.			√
Liver disorder: yellowing of the skin or eyes, dark urine, abdominal pain, nausea, vomiting, loss of appetite		√	
Neutropenia/leukopenia (low levels of white blood cells): infection, fatigue, fever, aches, pain, flu-like symptoms		√	
Oral candidiasis (yeast infection): creamy white bumps on the tongue, cheeks, gums or throat that bleed when scraped, pain, trouble swallowing, bad taste in the mouth		√	
Vaginal yeast infection: vaginal itching, burning during intercourse or urination, pain, redness, swelling, discharge		√	

If you have a troublesome symptom or side effect that is not listed here or becomes bad enough to interfere with your daily activities, tell your healthcare professional.

Reporting side effects

You can report any suspected side effects associated with the use of health products to Health Canada by:

- Visiting the Web page on Adverse Reaction Reporting (canada.ca/drug-device-reporting) for information on how to report online, by mail or by fax; or
- Calling toll-free at 1-866-234-2345.

NOTE: Contact your healthcare professional if you need information about how to manage your side effects. The Canada Vigilance Program does not provide medical advice.

Storage:

- CEFAZOLIN FOR INJECTION will be stored by your healthcare professional at room temperature (15°C – 30°C) and protected from light.
- Keep out of reach and sight of children.

If you want more information about CEFAZOLIN FOR INJECTION:

- Talk to your healthcare professional
- Find the full product monograph that is prepared for healthcare professionals and includes the Patient Medication Information by visiting the Health Canada Drug Product Database website: (<https://www.canada.ca/en/health-canada/services/drugs-health-products/drug-products/drug-product-database.html>); the manufacturer's website <http://www.tevacanada.com> ; or by calling 1-800-268-4127 ext. 3; or emailing druginfo@tevacanada.com.

This leaflet was prepared by Teva Canada Limited.

Date of Authorization: 2025-12-29