

PRODUCT MONOGRAPH
INCLUDING PATIENT MEDICATION INFORMATION

^{Pr} **OLANZAPINE-ODT**

Olanzapine Orally Disintegrating Tablets

Oral, 5 mg, 10 mg, 15 mg and 20 mg

USP

N05AH03, Antipsychotic Agent

Jubilant Generics Limited
1A, Sector 16A,
Institutional Area
Noida, Uttar Pradesh, India - 201301

Date of Initial Authorization:
AUG 22, 2014

Canadian Importer/distributor:
Mint Pharmaceuticals Inc.
1093 Meyerside Drive
Mississauga, Ontario
L5T 1J6

Date of Revision: JAN 27, 2026

Submission Control Number: 301000

RECENT MAJOR LABEL CHANGES

None at the time of the most recent authorization.

TABLE OF CONTENTS

Certain sections or subsections that are not applicable at the time of the preparation of the most recent authorized product monograph are not listed.

RECENT MAJOR LABEL CHANGES	2
PART I: HEALTH PROFESSIONAL INFORMATION.....	4
1 INDICATIONS	4
1.1 Pediatrics	4
1.2 Geriatrics.....	4
2 CONTRAINDICATIONS	4
3 SERIOUS WARNINGS AND PRECAUTIONS BOX	5
4 DOSAGE AND ADMINISTRATION	5
4.1 Dosing Considerations.....	5
4.2 Recommended Dose and Dosage Adjustment	5
4.4. Administration	6
4.5. Missed Dose.....	7
5 OVERDOSAGE	7
6 DOSAGE FORMS, STRENGTHS, COMPOSITION AND PACKAGING	7
7 WARNINGS AND PRECAUTIONS	8
7.1 Special Populations.....	15
7.1.1 Pregnant Women	15
7.1.2 Breast-feeding	16
7.1.3 Pediatrics	16
7.1.4 Geriatrics	16
8 ADVERSE REACTIONS	17
8.1 Adverse Reaction Overview	17
8.2 Clinical Trial Adverse Reactions.....	18

8.3	Less Common Clinical Trial Adverse Reactions	22
8.4	Abnormal Laboratory Findings: Hematologic, Clinical Chemistry and Other Quantitative Data	28
8.5	Post-Market Adverse Reactions	32
9	DRUG INTERACTIONS.....	34
9.2	Drug Interactions Overview	34
9.3	Drug-Behaviour Interactions	34
9.4	Drug-Drug Interactions.....	34
9.5	Drug-Food Interactions	37
9.6	Drug-Herb Interactions.....	37
9.7	Drug-Laboratory Test Interactions	37
10	CLINICAL PHARMACOLOGY	38
10.1	Mechanism of Action	38
10.2	Pharmacodynamics	38
10.3	Pharmacokinetics	41
11	STORAGE AND STABILITY.....	43
12	SPECIAL HANDLING INSTRUCTIONS	43
	PART II: SCIENTIFIC INFORMATION	44
13	PHARMACEUTICAL INFORMATION.....	44
14	CLINICAL TRIALS	44
14.1	Efficacy and Safety Studies	44
14.2	Study Results	46
14.3	Comparative Bioavailability Studies.....	56
15	MICROBIOLOGY	56
16	NON-CLINICAL TOXICOLOGY	56
17	SUPPORTING PRODUCT MONOGRAPHS.....	59
	PATIENT MEDICATION INFORMATION.....	60

PART I: HEALTH PROFESSIONAL INFORMATION

1. INDICATIONS

OLANZAPINE-ODT (olanzapine orally disintegrating tablets) is indicated for

- the acute treatment of schizophrenia and related psychotic disorders
- maintenance treatment of schizophrenia and related psychotic disorders.

In controlled clinical trials, olanzapine was found to improve both positive and negative symptoms.

Olanzapine has been shown to be effective in maintaining clinical improvement during 1-year of continuation therapy in patients who had shown an initial treatment response.

OLANZAPINE-ODT (olanzapine orally disintegrating tablets) is indicated for the acute treatment of

- manic or mixed episodes in bipolar I disorder.

Olanzapine may be used as monotherapy or cotherapy with agents commonly used in the treatment of acute bipolar disorder (e.g., lithium or divalproex sodium).

The efficacy of olanzapine as monotherapy maintenance treatment in bipolar patients with manic or mixed episodes who responded to acute treatment with OLANZAPINE-ODT was demonstrated in two 1-year “time to relapse” trials (see [14 CLINICAL TRIALS](#)).

The physician who elects to use OLANZAPINE-ODT for extended periods should periodically re-evaluate the long-term usefulness of the drug for the individual patient (see [4 DOSAGE AND ADMINISTRATION](#) section).

1.1 Pediatrics

Pediatrics: Based on the data submitted and reviewed by Health Canada, the safety and efficacy of olanzapine in pediatric patients has not been established. Therefore, Health Canada has not authorized an indication for pediatric use (see [7.1.3 Pediatrics](#)).

1.2 Geriatrics

Geriatrics (≥65 years old): OLANZAPINE-ODT is not indicated for treatment of elderly patients with dementia (see [3 SERIOUS WARNINGS AND PRECAUTIONS BOX](#) and [7.1.4. Geriatrics](#)). Caution should be used when treating geriatric patients with OLANZAPINE-ODT (see [10.3 Pharmacokinetics](#), [7.1.4 Geriatrics](#), and [4.1 Dosing Considerations](#)).

2. CONTRAINDICATIONS

OLANZAPINE-ODT is contraindicated in those patients who are hypersensitive to this drug or to any ingredient in the formulation, including any non-medicinal ingredient, or component of the container. For a complete listing, see [6 DOSAGE FORMS, STRENGTHS, COMPOSITION AND PACKAGING](#) section.

3. SERIOUS WARNINGS AND PRECAUTIONS BOX

Serious Warnings and Precautions

Increased Mortality in Elderly Patients with Dementia:

Elderly patients with dementia treated with atypical antipsychotic drugs are at an increased risk of death compared to placebo. Analyses of thirteen placebo controlled trials with various atypical antipsychotics (modal duration of 10 weeks) in these patients showed a mean 1.6-fold increase in death rate in the drug-treated patients. Although the causes of death were varied, most of the deaths appeared to be either cardiovascular (e.g., heart failure, sudden death) or infectious (e.g., pneumonia) in nature. (See [7 WARNINGS AND PRECAUTIONS, 7.1 Special Populations, Use in Geriatric Patients with Dementia](#)). OLANZAPINE-ODT (olanzapine) is not approved for use in elderly patients with dementia.

4. DOSAGE AND ADMINISTRATION

4.1 Dosing Considerations

General Considerations for Oral Dosing in Special Populations

Elderly or Debilitated Patients:

In clinical trials, 44 patients with schizophrenia or related disorders who were 65 years of age or over were treated with OLANZAPINE-ODT (5 to 20 mg daily) (see [7 WARNINGS AND PRECAUTIONS, 7.1. Special Populations](#)). Given the limited experience with olanzapine in the elderly, and the higher incidence of concomitant illness and concomitant medication in this population, OLANZAPINE-ODT should be used with caution.

The recommended starting dose is 5 mg in patients who are elderly, debilitated, who have a predisposition to hypotensive reactions, who otherwise exhibit a combination of factors that may result in slower metabolism of olanzapine (e.g., nonsmoking female patients), or who may be pharmacodynamically more sensitive to olanzapine. When indicated, dose escalation should be performed with caution in these patients.

Patients with Hepatic and/or Renal Impairment:

As clinical experience is lacking in these patients, the lower initial starting dose and slower titration to initial target dose should be considered. Further dose escalation, when indicated, should be conservative (see [7 WARNINGS AND PRECAUTIONS, 7.1 Special Populations](#)).

4.2 Recommended Dose and Dosage Adjustment

Schizophrenia and Related Disorders

Adults: OLANZAPINE-ODT should be administered on a once-a-day schedule without regard to meals, generally beginning with 5 to 10 mg, with a target dose of 10 mg/day within several days. Further dosage adjustments, if indicated, should generally occur at intervals of not less than 1 week, since steady state for olanzapine would not be achieved for approximately 1 week in the typical patient. When dosage adjustments are necessary, dose increments/ decrements of 5 mg

per day are recommended. An increase to a dose greater than target dose of 10 mg/day (i.e., to a dose of 15 mg/day or greater) is normally recommended only after clinical assessment.

In clinical trials a dose range of 5 to 20 mg/day was studied (see [14 CLINICAL TRIALS](#)).

Doses above 20 mg/day have been evaluated from a safety perspective (see Table 6 in Adverse Events, Dose- Dependent Adverse Events subsection); however, efficacy at doses above 20 mg/day has not been systematically evaluated.

Maintenance Therapy in Schizophrenia:

It is recommended that responding patients with schizophrenia be continued on OLANZAPINE-ODT at the lowest dose needed to maintain remission. Patients should be reassessed periodically to determine the need for maintenance treatment. While there is no body of evidence available to answer the question of how long the patient should be treated with olanzapine, the effectiveness of maintenance treatment is well established for many other antipsychotic drugs.

Bipolar Disorder

Bipolar Mania

Adults: The recommended starting dose for olanzapine is 15 mg administered once a day in monotherapy and 10 mg daily in combination therapy.

It may be given without regard to meals as its absorption is not affected by food. The dosage range of olanzapine is from 5 mg to 20 mg per day. Daily dosage should be adjusted in response to clinical assessment.

Maintenance Therapy in Bipolar Disorder:

Patients who have been receiving and responding to OLANZAPINE-ODT for the treatment of acute manic or mixed episodes of bipolar disorder should initially continue maintenance therapy at the same dose (see [14 CLINICAL TRIALS](#)). Subsequent daily dosage should be adjusted on the basis of clinical status within a range of 5 to 20 mg per day.

Patients should be periodically reassessed to determine the need for maintenance treatment and the appropriate dose for such treatment.

Health Canada has not authorized an indication for pediatric use.

4.4. Administration

OLANZAPINE-ODT (orally disintegrating tablet) is intended for oral administration only. It begins disintegrating in the mouth within seconds, allowing its contents to be subsequently swallowed with or without liquid.

The orally disintegrating tablet breaks easily and should be handled carefully, with dry hands. Direct contact with hands should be avoided if possible. One blister cell must be separated from the strip prior to peeling the blister backing. The blister backing should be carefully peeled back and the orally disintegrating tablet pushed out and placed directly in the mouth. The orally disintegrating tablet may also be stirred into 125 mL (4 ounces) of water, milk, coffee, orange juice or apple juice and the contents promptly consumed.

4.5. Missed Dose

If a patient misses a dose by a few hours, advise patient to take as soon as he/she remembers. If most of the day has passed, advise patient to wait until the next scheduled dose. Advise patients to not take 2 doses of OLANZAPINE-ODT at once.

5. OVERDOSAGE

Signs and Symptoms

Very common symptoms reported in olanzapine overdose ($\geq 10\%$ incidence) include tachycardia, agitation/aggressiveness, dysarthria, various extrapyramidal symptoms, and reduced level of consciousness ranging from sedation to coma.

Other medically significant sequelae of olanzapine overdose include delirium, convulsion, possible neuroleptic malignant syndrome, respiratory depression, aspiration, hypertension or hypotension, cardiac arrhythmias ($< 2\%$ of overdose cases) and cardiopulmonary arrest. Fatal outcomes have been reported for acute overdoses as low as 450 mg of oral olanzapine but survival has also been reported following acute overdose of approximately 2,000 mg of oral olanzapine.

Management of Overdose

There is no specific antidote for olanzapine. Induction of emesis is not recommended. Standard procedures for management of overdose may be indicated (i.e., gastric lavage, administration of activated charcoal). The concomitant administration of activated charcoal was shown to reduce the oral bioavailability of olanzapine by 50 to 60%.

Symptomatic treatment and monitoring of vital organ function should be instituted according to clinical presentation, including treatment of hypotension and circulatory collapse and support of respiratory function. Do not use epinephrine, dopamine, or other sympathomimetic agents with beta-agonist activity since beta stimulation may worsen hypotension.

For the most recent information in the management of a suspected drug overdose, contact your regional poison control centre or Health Canada's toll-free number, 1-844 POISON-X (1-844-764-7669).

6. DOSAGE FORMS, STRENGTHS, COMPOSITION AND PACKAGING

Table 1: Dosage Forms, Strengths, Composition and Packaging

Route of Administration	Dosage Form / Strength	Non-medicinal Ingredients
Oral	orally disintegrating tablets / 5 mg, 10 mg, 15 mg, 20 mg	Aspartame, Low substituted hydroxypropyl cellulose, Magnesium stearate, Mannitol, Silica, Strawberry flavour

Availability of dosage forms:

The 5 mg tablets are yellow coloured, round, flat faced bevelled edge tablets with characteristic flavour, debossed with 'D5' on one side and 'CO' on the other.

The 10 mg tablets are yellow coloured, round, flat faced bevelled edge tablets with characteristic flavour, debossed with 'D10' on one side and 'CO' on the other.

The 15 mg tablets are yellow coloured, round, flat faced bevelled edge tablets with characteristic flavour, debossed with 'D15' on one side and 'CO' on the other.

The 20 mg tablets are yellow coloured, round, flat faced bevelled edge tablets with characteristic flavour, debossed with 'D20' on one side and 'CO' on the other.

The 5 mg, 10 mg, 15 mg and 20 mg tablets are supplied in aluminum blister strips of 10 tablets and 30 tablets per carton.

Phenylketonurics: OLANZAPINE-ODT contains phenylalanine (1.12, 2.24, 3.36, or 4.8 mg per 5, 10, 15, or 20 mg oral lyophilisate, respectively).

7. WARNINGS AND PRECAUTIONS

Please see [3 SERIOUS WARNINGS AND PRECAUTIONS BOX](#)

General

Clinical experience with olanzapine in patients with concomitant illness is limited. Caution is thus advised when using OLANZAPINE-ODT in patients with diseases or conditions that could affect the metabolism or the pharmacodynamic activity of olanzapine (see [4 DOSAGE AND ADMINISTRATION](#) section and [10 CLINICAL PHARMACOLOGY](#)).

Body Temperature Regulation:

Disruption of the body's ability to reduce core body temperature has been attributed to antipsychotic agents. Appropriate care is advised when prescribing OLANZAPINE-ODT for patients who will be experiencing conditions which may contribute to an elevation of core temperature, e.g., exercising strenuously, exposure to extreme heat, receiving concomitant medication with anticholinergic activity, or being subject to dehydration.

Falls:

Olanzapine may cause somnolence, postural (orthostatic) hypotension, motor and sensory instability, which may lead to falls and, consequently, fractures or other injuries. For patients with diseases, conditions, or medications that could exacerbate these effects, complete fall risk assessments when initiating antipsychotic treatment and recurrently for patients on long-term antipsychotic therapy.

Anticholinergic effects

As olanzapine demonstrated anticholinergic activity *in vitro*, caution is advised when prescribing for patients with symptomatic prostatic enlargement, narrow-angle glaucoma or paralytic ileus and

related conditions.

Carcinogenesis and Mutagenesis

For animal data, see [16 NON-CLINICAL TOXICOLOGY](#) section.

Cardiovascular

Olanzapine has not been evaluated in patients with a recent history of myocardial infarction or unstable heart disease. Patients with these conditions were excluded from pre-marketing clinical trials.

Hypotension and Syncope:

As with other drugs that have high alpha-1 adrenergic receptor blocking activity, olanzapine may induce orthostatic hypotension, tachycardia, dizziness, and sometimes syncope, especially at the initiation of treatment. In a clinical trial database of 2500 patients treated with oral olanzapine, syncope was reported in 0.6% (15/2500). The risk of orthostatic hypotension and syncope may be minimized by initiating therapy with 5 mg QD (see [4 DOSAGE AND ADMINISTRATION](#) section). A more gradual titration to the target dose should be considered if hypotension occurs.

OLANZAPINE-ODT should be used with particular caution in patients with known cardiovascular disease (history of myocardial infarction or ischemia, heart failure, or conduction abnormalities), cerebrovascular disease, and conditions which would predispose patients to hypotension (dehydration, hypovolemia, and treatment with antihypertensive medications).

QT Interval:

In clinical trials, clinically meaningful QTc prolongations (Fridericia QT correction [QTcF] $\geq$ 500 milliseconds [msec] at any time post baseline in patients with baseline QTcF < 500 msec) were uncommon (0.1% to 1%) in patients treated with olanzapine, with no significant differences in associated cardiac events compared to placebo. However, as with other antipsychotics, caution should be exercised when olanzapine is prescribed with medicines known to increase QTc interval, especially in the elderly, in patients with congenital long QT syndrome, congestive heart failure, heart hypertrophy, hypokalaemia or hypomagnesaemia (see [8 ADVERSE REACTIONS, 8.5 Post-Market Adverse Reactions](#)).

Cardiac Death:

In a retrospective observational study, patients treated with atypical antipsychotics (including olanzapine) or typical antipsychotics had a similar dose-related increase of presumed sudden cardiac death (SCD) compared to non-users of antipsychotics (almost twice the risk than that for non-users). In postmarketing reports with olanzapine, the event of SCD has been reported very rarely.

Driving and Operating Machinery

Potential Effect on Cognitive and Motor Performance:

Because olanzapine may cause somnolence, patients should be cautioned about operating hazardous machinery, including motor vehicles, until they are reasonably certain that OLANZAPINE-ODT therapy does not affect them adversely.

Exercise caution when driving or operating a vehicle or potentially dangerous machinery.

Endocrine and Metabolism

Weight Gain:

Olanzapine was associated with weight gain during clinical trials. Clinically significant weight gain was observed across all baseline body mass index (BMI) categories (see [8 ADVERSE REACTIONS, Other Adverse Events Observed During Clinical Trials with Olanzapine Across All Indications, Weight Changes](#)). Using pooled data from patients treated with olanzapine over the dosage range of 5 mg to 20 mg per day mean gain was 5.4 kg. The mean change in weight was comparable for patients with schizophrenia and bipolar mania. A retrospective analysis of 573 patients receiving olanzapine for up to 3 years found that dose was not a significant predictor of greater long-term changes in weight.

In long-term studies (at least 48 weeks), both the magnitude of weight gain and the proportion of olanzapine-treated patients who had clinically significant weight gain were greater than in short-term studies. The percentage of patients who gained $\geq 25\%$ of their baseline body weight with long-term exposure was very common ($\geq 10\%$).

Hyperglycaemia:

As with some other antipsychotics, exacerbation of pre-existing diabetes and hyperglycaemia have been reported rarely and diabetic ketoacidosis and diabetic coma including some fatal cases have been reported very rarely during the use of olanzapine sometimes in patients with no reported history of hyperglycaemia (see [8 ADVERSE REACTIONS, 8.5 Post-Market Adverse Reactions](#) section). In some cases, a prior increase in body weight has been reported which may be a predisposing factor. Patients should have baseline and periodic monitoring of blood glucose and body weight.

In clinical trials (up to 52 weeks) olanzapine was associated with a greater mean change in glucose relative to placebo. Treatment-emergent clinically significant changes in fasting glucose were observed in patients with or without evidence of glucose dysregulation at baseline (see [8 ADVERSE REACTIONS, Other Adverse Events Observed During Clinical Trials with Olanzapine Across All Indications, Glucose Changes](#)).

Assessment of the relationship between atypical antipsychotic use and glucose abnormalities is complicated by the possibility of an increased background risk of diabetes mellitus in patients with schizophrenia and the increasing incidence of diabetes mellitus in the general population. Given these confounders, the relationship between atypical antipsychotic use and hyperglycemia-related adverse events is not completely understood. However, epidemiological studies suggest an increased risk of treatment-emergent hyperglycemia-related adverse events in patients treated with the atypical antipsychotics. Precise risk estimates for hyperglycemia-related adverse events in patients treated with atypical antipsychotics are not available.

Any patient treated with atypical antipsychotics should be monitored for symptoms of hyperglycemia including polydipsia, polyuria, polyphagia, and weakness. Patients who develop symptoms of hyperglycemia during treatment with atypical antipsychotics should undergo fasting blood glucose testing. In some cases, hyperglycemia has resolved when the atypical antipsychotic was discontinued; however, some patients required continuation of anti-diabetic treatment despite discontinuation of the suspect drug. Patients with risk factors for diabetes mellitus (e.g., obesity, family history of diabetes) who are starting treatment with atypical antipsychotics should undergo fasting blood glucose testing at the beginning of treatment and periodically during treatment. Patients with an established diagnosis of diabetes mellitus who are started on atypical antipsychotics should be monitored regularly for worsening of glucose control.

Hyperprolactinemia:

As with other drugs that block dopamine D₂ and/or serotonin 5-HT₂ receptors, olanzapine may elevate prolactin levels. Elevations associated with olanzapine treatment are generally mild, and may decline during continued administration.

Since tissue culture experiments indicate that approximately one third of human breast cancers are prolactin-dependent *in vitro*, OLANZAPINE-ODT should only be administered to patients with previously detected breast cancer if the benefits outweigh the potential risks. Caution should also be exercised when considering OLANZAPINE-ODT treatment in patients with pituitary tumours. Possible manifestations associated with elevated prolactin levels are amenorrhea, galactorrhea, and menorrhagia.

As is common with compounds which stimulate prolactin release, the administration of olanzapine resulted in an increase in the incidence of mammary neoplasms in both rats and mice. The physiological differences between rats and humans with regard to prolactin make the clinical significance of these findings unclear. To date, neither clinical nor epidemiological studies have shown an association between chronic administration of these drugs and mammary tumorigenesis.

Long-standing hyperprolactinemia when associated with hypogonadism may lead to decreased bone mineral density in both female and male subjects.

Lipids:

Increases in lipids have been observed in olanzapine-treated patients in placebo-controlled clinical trials. Treatment-emergent clinically significant changes in fasting lipids were observed in patients with or without evidence of dyslipidemia at baseline (see [8 ADVERSE REACTIONS, Other Adverse Events Observed During Clinical Trials with Olanzapine Across All Indications, Lipids subsection](#)). Appropriate clinical monitoring is recommended, including baseline and follow-up lipid evaluations.

Gastrointestinal

Antiemetic Effect:

Consistent with its dopamine antagonist effects, olanzapine may have an antiemetic effect. Such an effect may mask signs of toxicity due to overdosage of other drugs, or may mask symptoms of disease such as brain tumour or intestinal obstruction.

Genitourinary

Priapism:

Rare cases of priapism have been reported with antipsychotic use, such as olanzapine. This adverse reaction, as with other psychotropic drugs, did not appear to be dose-dependent and did not correlate with the duration of treatment. The most likely mechanism of action of priapism is a relative decrease in sympathetic tone.

Urinary Retention:

Olanzapine possesses anticholinergic properties, which can lead to adverse drug reactions such as urinary retention. There have been several serious post-marketing reports of urinary retention in OLANZAPINE-ODT-treated patients and in some cases, catheterization was required. OLANZAPINE-ODT should be prescribed with caution in patients with a current diagnosis or prior history of urinary retention and in patients with other risk factors for urinary retention (e.g. benign prostatic hyperplasia). OLANZAPINE-ODT should also be prescribed with caution in patients receiving medications with anticholinergic activity that can affect voiding.

Hematologic

Venous Thromboembolism:

Venous thromboembolism (VTE), including fatal pulmonary embolism, has been reported in temporal association with antipsychotic drugs, including olanzapine, in case reports and/or observational studies. When prescribing OLANZAPINE-ODT, all potential risk factors for VTE should be identified and preventative measures undertaken, particularly since patients with schizophrenia often present with risk factors for VTE. Very rare cases of VTE have been reported in olanzapine-treated patients during the post-marketing period.

Hematologic Indices:

In oral olanzapine clinical trials, there were no data to suggest olanzapine adversely affected bone marrow function, even in patients with a history of clozapine-associated neutropenia or leukopenia. Olanzapine was associated with a 5.7% incidence of mainly transient treatment-emergent elevations of eosinophil counts above the normal range. Elevations were not associated with any symptoms, identifiable allergic phenomena, or changes in other hematologic indices. Rare cases of leukopenia have been reported with olanzapine. In case of symptoms of infection, WBC count and differential count should be considered.

Neutropenia, granulocytopenia and agranulocytosis have been reported during antipsychotic use. Therefore, it is recommended that patients have their complete blood count (CBC) tested prior to starting OLANZAPINE-ODT and then periodically throughout treatment.

Hepatic

Small single-dose clinical pharmacology studies (see [10 CLINICAL PHARMACOLOGY](#) section) did not reveal any major alterations in olanzapine pharmacokinetics in subjects with hepatic impairment. Given the limited clinical experience with olanzapine in patients with these conditions, caution should be exercised (see [4 DOSAGE AND ADMINISTRATION](#) section).

Aminotransferase Elevations:

During pre-marketing clinical trials, therapy with oral olanzapine was associated with elevation of hepatic aminotransferases, primarily ALT (SGPT). Within a clinical trial database of 2280 olanzapine-treated patients, with baseline ALT (SGPT) levels ≤ 60 IU/L, 5.9% (134/2280) had treatment-emergent ALT (SGPT) elevations to > 120 IU/L, 1.9% (44/2280) had elevations to > 200 IU/L, and 0.2% (5/2280) had elevations to > 400 IU/L. No patients had values in excess of 700 IU/L. None of the olanzapine-treated patients who had elevated aminotransferase values manifested clinical symptomatology associated with liver impairment. The majority of aminotransferase elevations were seen during the first six weeks of treatment. Most elevations were transient (66%) while patients continued on olanzapine therapy, or falling (11%) at the last available measurement. Of the 134 olanzapine-treated patients whose enzyme levels increased to > 120 IU/L, 20 discontinued treatment (6 for hepatic, 14 for other reasons) while their ALT (SGPT) values were still rising. In 38 olanzapine-treated patients with baseline ALT (SGPT) > 90 IU/L, none experienced an elevation to > 400 IU/L.

Rare post-marketing reports of hepatitis have been received. Very rare cases of cholestatic or mixed liver injury have also been reported in the post-marketing period. Hepatic failure, including fatalities has also been reported very rarely in the post-marketing period. See [8.5 Post-Market Adverse Reactions](#).

Precaution should be exercised when using OLANZAPINE-ODT in patients with pre-existing hepatic disorders, in patients who are being treated with potentially hepatotoxic drugs, or if treatment-emergent signs or symptoms of hepatic impairment appear.

For patients who have known or suspected abnormal hepatic function prior to starting OLANZAPINE-ODT, standard clinical assessment including measurement of aminotransferase levels is recommended. Periodic clinical reassessment with aminotransferase levels is recommended for such patients, as well as for patients who develop any signs and symptoms suggestive of a new onset liver disorder during OLANZAPINE-ODT therapy.

Neurologic

Neuroleptic Malignant Syndrome:

Neuroleptic malignant syndrome (NMS) is a potentially fatal symptom complex that has been reported in association with antipsychotic drugs, including olanzapine.

Clinical manifestations of NMS are hyperpyrexia, muscle rigidity, altered mental status, and evidence of autonomic instability (irregular pulse or blood pressure, tachycardia, diaphoresis, and cardiac dysrhythmia). Additional signs may include elevated creatine phosphokinase, myoglobinuria (rhabdomyolysis), and acute renal failure.

In arriving at a diagnosis, it is important to identify cases where the clinical presentation includes both serious medical illness (e.g., pneumonia, systemic infection, etc.), and untreated or inadequately treated extrapyramidal signs and symptoms (EPS). Other important considerations in the differential diagnosis include central anticholinergic toxicity, heat stroke, drug fever, and primary central nervous system pathology.

The management of NMS should include 1) immediate discontinuation of all antipsychotic drugs including OLANZAPINE-ODT and other drugs not essential to therapy; 2) intensive symptomatic treatment and medical monitoring; and 3) treatment of any concomitant serious medical problems for which specific treatments are available. There is no general agreement about specific pharmacological treatment for uncomplicated NMS.

If a patient requires antipsychotic drug treatment after recovery from NMS, the potential re-introduction of therapy should be very carefully considered. The patient should be carefully monitored since recurrences of NMS have been reported.

Tardive Dyskinesia:

Tardive dyskinesia (TD), a syndrome consisting of potentially irreversible involuntary dyskinetic movements, is associated with the use of antipsychotic drugs. Tardive dyskinesia occurs more frequently in elderly patients; however, patients of any age can be affected. It is unknown whether antipsychotic drugs may differ in their potential to cause TD. However, during long-term, double-blind, extension schizophrenia maintenance trials (894 olanzapine-treated patients; median olanzapine treatment, 237 days), olanzapine was associated with a statistically significantly lower incidence of treatment-emergent dyskinesia compared to haloperidol. During long-term, double-blind, monotherapy extension bipolar maintenance trials (567 olanzapine-treated patients, for up to 1 year), there were no cases of TD in the olanzapine arms, as determined by either reported adverse events or the Abnormal Involuntary Movement Scale (AIMS). TD has been reported very rarely (≤

0.0025%) in post-market surveillance.

The risk of developing tardive dyskinesia and the chance of it becoming irreversible are believed to increase as the duration of treatment and the cumulative dose of antipsychotic drug increase. However, the syndrome can develop, although less commonly, after relatively brief periods of treatment at low doses. There is no known treatment for established cases of TD. The syndrome may remit, partially or completely, if antipsychotic drug treatment is withdrawn. Antipsychotic drug treatment itself, however, may suppress the signs and symptoms of tardive dyskinesia, thereby masking the underlying process.

Given these considerations, OLANZAPINE-ODT should be prescribed in a manner that is most likely to minimize the risk of tardive dyskinesia. As with any antipsychotic drug, OLANZAPINE-ODT should be reserved for patients who appear to be receiving substantial benefit from the drug. In such patients the lowest effective dose and the shortest duration of treatment should be sought. The need for continued treatment should be reassessed periodically.

If signs or symptoms of tardive dyskinesia appear in a patient on OLANZAPINE-ODT, drug discontinuation should be considered. However, some patients may benefit from continued treatment with OLANZAPINE-ODT despite the presence of the syndrome.

In clinical trials, a single case of pre-existing intracranial hypertension was exacerbated.

Seizures:

Conventional neuroleptics are known to lower seizure threshold. In clinical trials, seizures have occurred in a small number (0.9%, 22/2500) of olanzapine-treated patients. There were confounding factors that may have contributed to the occurrence of seizures in many of these cases. OLANZAPINE-ODT should be used cautiously in patients who have a history of seizures or have conditions associated with seizures or have a lowered seizures threshold.

Psychiatric

Suicide:

The possibility of suicide or attempted suicide is inherent in psychosis, and thus close supervision and appropriate clinical management of high-risk patients should accompany drug therapy.

Renal

Small single-dose clinical pharmacology studies (see [10 CLINICAL PHARMACOLOGY](#) section) did not reveal any major alterations in olanzapine pharmacokinetics in subjects with renal impairment. Given the limited clinical experience with olanzapine in patients with these conditions, caution should be exercised (see [4 DOSAGE AND ADMINISTRATION](#) section).

Uric Acid:

In the pre-marketing clinical trial database, oral olanzapine was associated with mild elevations of uric acid in some patients. However, only one olanzapine-treated patient experienced treatment-emergent gout, and the baseline uric acid concentration for this patient was at least as large as all concentrations observed while the patient was receiving olanzapine.

Skin

Drug Reaction with Eosinophilia and Systemic Symptoms (DRESS):

Drug Reaction with Eosinophilia and Systemic Symptoms (DRESS) has been reported with olanzapine

exposure. DRESS consists of a combination of three or more of the following: cutaneous reaction (such as rash or exfoliative dermatitis), eosinophilia, fever, lymphadenopathy and one or more systemic complications such as hepatitis, nephritis, pneumonitis, myocarditis, and pericarditis. Severe cutaneous adverse reactions are sometimes fatal. Discontinue olanzapine if DRESS is suspected.

7.1. Special Populations

7.1.1. Pregnant Women

There are no adequate and well-controlled studies in pregnant women. Patients should be advised to notify their physician if they become pregnant or intend to become pregnant during treatment with OLANZAPINE-ODT. Because human experience in pregnant females is limited, this drug should be used in pregnancy only if the potential benefit justifies the potential risk to the fetus.

Non-Teratogenic Effects:

Neonates exposed to antipsychotic drugs (including OLANZAPINE-ODT) during the third trimester of pregnancy are at risk for extrapyramidal and/or withdrawal symptoms following delivery. There have been reports of agitation, hypertonia, hypotonia, tremor, somnolence, respiratory distress and feeding disorder in these neonates. These complications have varied in severity; while in some cases symptoms have been self-limited, in other cases neonates have required intensive care unit support and prolonged hospitalization.

OLANZAPINE-ODT should not be used during pregnancy unless the expected benefits to the mother markedly outweigh the potential risks to the fetus.

Labour and Delivery:

Parturition in rats was not affected by olanzapine. The effect of olanzapine on labour and delivery in humans is not known.

Reproductive and Developmental toxicology in animal models

Fertility studies in male and female rats and teratology studies in rats and rabbits have been conducted by the oral route. Mating performance was affected by administration of very high olanzapine doses due to sedation in male rats, but the effect was quickly reversed when treatment stopped. Estrous cycles were affected and reproduction parameters were influenced in rats given high doses. No adverse effects were observed on numbers of corpora lutea, implantations, fetal viability, or fetal weight, and there were no effects on litter size or on the survival, growth, or development of the offspring from parents given up to 18 times the maximum daily human dose (5 mg/kg/day in rats). Although the reproductive process in female rats from mating through fertilization was not adversely affected by treatment, this evidence does not exclude a possible interference with maintenance of pregnancy at high doses of olanzapine. The reproduction studies conducted did not reveal evidence of harm to the fetus. Maternal toxicity, developmental toxicity (indicated by fetal growth retardation and slightly delayed ossification at birth), and increased numbers of nonviable offspring occurred at higher doses. However, fetal malformations were not increased. Transient modest decreases in activity levels of the progeny from females given 0.25 mg/kg/day and skeletal changes indicative of growth retardation in fetuses from females given 5 mg/kg/day were observed. Transient decreases in offspring activity have occurred at all doses; however, there were no effects on body weight, growth, mating, fertility, or live births in second-generation animals. Placental transfer of olanzapine occurs in rat fetuses (see [16 NON-CLINICAL](#)

TOXICOLOGY).**7.1.2. Breast-feeding**

In a study in lactating, healthy women, olanzapine was excreted in breast milk. Mean infant exposure (mg/kg) at steady state was estimated to be 1.8% of the maternal olanzapine dose (mg/kg). Patients should be advised not to breast feed an infant if they are taking OLANZAPINE-ODT.

Data from animal model

Olanzapine was also detected in the milk of rats at concentrations up to three-fold higher than those in the plasma (see [16 NON-CLINICAL TOXICOLOGY](#)).

7.1.3. Pediatrics

Pediatrics (< 18 years of age): The safety and efficacy of OLANZAPINE-ODT in children under the age of 18 years have not been established and its use is not recommended.

Weight gain has been observed with atypical antipsychotic use in pediatric and adolescent patient populations. Independent of any changes during treatment with atypical antipsychotics, weight gain can be associated with adverse changes in other metabolic parameters (e.g., glucose and lipid metabolism). Abnormal childhood weight and metabolic status have been associated with adverse cardiovascular outcomes in adulthood. Weight gain and changes in other metabolic parameters associated with atypical antipsychotics can be more frequent or more severe in pediatric and adolescent patients than in the adult patients.

The long-term safety of atypical antipsychotics, including potential cardiometabolic effects and effects on growth, maturation and behavioural development in patients under 18 years of age has not been systematically evaluated.

A greater magnitude of weight gain and lipid alterations has been reported in adolescents compared with adults. Adolescents treated with olanzapine experienced a significantly higher incidence of elevated prolactin levels and significantly higher mean increases in prolactin levels compared with adults. Hepatic aminotransferase elevations are more common in adolescents as compared to adults. Sedation-related events are more common in adolescents as compared to adults.

See also [8 ADVERSE REACTIONS/Other Investigational Trials/Adverse Events in Adolescent Patients \(ages 13 to 17 years\)](#).

No data are available to Health Canada; therefore, Health Canada has not authorized an indication for pediatric use.

7.1.4. Geriatrics

Geriatrics (≥ 65 years of age):

The number of patients 65 years of age or over with schizophrenia or related disorders exposed to oral olanzapine during clinical trials was limited (N = 44). Caution should thus be exercised with the use of OLANZAPINE-ODT in the elderly patient, recognizing the more frequent hepatic, renal,

central nervous system, and cardiovascular dysfunctions, and more frequent use of concomitant medication in this population (see [4 DOSAGE AND ADMINISTRATION](#) section).

Use in Geriatric Patients with Dementia

Overall Mortality:

Elderly patients with dementia treated with atypical antipsychotic drugs showed increased mortality compared to placebo in a meta-analysis of 13 controlled trials of various atypical antipsychotic drugs. In five placebo-controlled trials with oral olanzapine in this population, the incidence of mortality was 3.5% for olanzapine-treated patients compared to 1.5% for placebo-treated patients. OLANZAPINE-ODT is not indicated in elderly patients with dementia (see [3 SERIOUS WARNINGS AND PRECAUTIONS BOX](#)).

Dysphagia:

Esophageal dysmotility and aspiration have been associated with antipsychotic drug use. Aspiration pneumonia is a common cause of morbidity and mortality in patients with advanced Alzheimer's disease. Olanzapine and other antipsychotic drugs should be used cautiously in patients at risk for aspiration pneumonia.

Cerebrovascular Adverse Events (CVAEs), Including Stroke, in Elderly Patients with Dementia:

Cerebrovascular adverse events (e.g., stroke, transient ischemic attack), including fatalities, were reported in trials of olanzapine in elderly patients with dementia-related psychosis. In placebo-controlled studies, there was a higher incidence of CVAEs in patients treated with olanzapine compared to patients treated with placebo (1.3% vs. 0.4%, respectively; see [8 ADVERSE REACTIONS](#) section). OLANZAPINE-ODT is not approved for the treatment of elderly patients with dementia.

There is insufficient evidence to determine whether CVAEs in elderly patients with dementia are associated specifically with olanzapine or all antipsychotic agents. Clinical trial data appear to suggest that patients with a dementia diagnosis of vascular or mixed type had a higher likelihood of experiencing CVAEs than other types of dementia.

The risks and benefits of the use of OLANZAPINE-ODT in elderly patients with dementia should be assessed taking into account the risk predictors for CVAEs in the individual patient. Patients/caregivers should be advised to immediately report signs and symptoms of potential CVAEs, such as sudden weakness or numbness in the face, arms, or legs, and speech or vision problems.

8. ADVERSE REACTIONS

8.1. Adverse Reaction Overview

The stated frequencies of adverse events represent the proportion of individuals who experienced at least once, a treatment-emergent adverse event of the type listed. An event was considered treatment-emergent if it occurred for the first time or worsened while receiving therapy following baseline evaluation. It is important to emphasize that although the events were reported during therapy, they were not necessarily caused by the therapy.

The most commonly reported adverse reactions in patients with schizophrenia that were reported more frequently with olanzapine oral therapy than with placebo were discontinuation due to ALT (SGPT) elevation, dizziness, constipation, personality disorder, weight gain, akathisia, and postural hypotension (see [7 WARNINGS AND PRECAUTIONS](#)).

The most commonly reported adverse reactions in patients with bipolar disorder that were reported more frequently with olanzapine oral therapy than with placebo were somnolence, dry mouth, dizziness, asthenia, constipation, dyspepsia, increased appetite, weight gain, back pain, speech disorder, increased salivation, amnesia, paresthesia, and tremor (see [7 WARNINGS AND PRECAUTIONS](#)).

8.2. Clinical Trial Adverse Reactions

The prescriber should be aware that the figures in the tables and tabulations cannot be used to predict the incidence of side effects in the course of usual medical practice where patient characteristics and other factors differ from those that prevailed in the clinical trials. Similarly, the cited frequencies cannot be compared with figures obtained from other clinical investigations involving different treatments, uses, and investigators. The figures cited, however, do provide the prescribing physician with some basis for estimating the relative contribution of drug and nondrug factors to the side effect incidence in the populations studied. Information in tabular format can be found in Table 2. The information included in these tables is also outlined in the following text sections. Adverse events associated with discontinuation are considered separately.

Incidence of Adverse Events Associated with Discontinuation:

Schizophrenia and Related Disorders:

In short-term, placebo-controlled trials, there was no statistically significant difference in rates of discontinuation of olanzapine or placebo attributed to adverse events. Overall, 5% of olanzapine-treated patients discontinued treatment for adverse events compared with 6% of placebo-treated patients. Discontinuations due to ALT (SGPT) elevations, however, were considered to be drug related (2% for olanzapine versus 0% for placebo) (see [7 WARNINGS AND PRECAUTIONS, Renal subsection](#)).

Bipolar Disorder:

Bipolar Mania

In short-term, placebo-controlled clinical trials, there was no difference overall in the incidence of discontinuation due to adverse events (2% for olanzapine versus 2% for placebo).

Bipolar Maintenance

In the long-term (1-year), placebo-controlled clinical trial, of the 225 olanzapine-treated patients, 16% (n = 35) discontinued due to an adverse event, compared with 9% (n = 12) of 136 placebo-treated patients.

In the long-term (1-year), active-controlled clinical trial, of the 217 olanzapine-treated patients, 19% (n = 41) discontinued due to an adverse event, compared with 26% (n = 55) of 214 lithium-treated patients.

All Short-Term Trials - Schizophrenia and Bipolar Mania Trials:

In short-term, active-controlled clinical trials, of the 1796 oral olanzapine-treated patients in

comparative clinical trials with haloperidol, 98 (5%) discontinued treatment for adverse events compared with 66 of 810 (8%) haloperidol-treated patients.

All Short-Term Trials - Overall Integrated Safety Database:

In a pre-marketing clinical trial database of 2500 olanzapine-treated patients, 14.9% (372/2500) discontinued due to an adverse event. About half (183/372) of these discontinuations were associated with the underlying psychopathology. Other adverse events most commonly (incidence of 0.5% to 0.6%) reported as the reason for discontinuation among olanzapine-treated patients were: ALT (SGPT) increased, unintended pregnancy, creatine phosphokinase increased, and convulsion.

Incidence of Commonly Observed Adverse Events:

Schizophrenia and Related Disorders:

In the schizophrenia placebo-controlled trials, the most commonly observed adverse events associated with the use of olanzapine (incidence of $\geq 5\%$ and at least twice placebo) were: dizziness (11% for olanzapine vs 4% for placebo), constipation (9% vs 3%), ALT (SGPT) increased (8% vs 3%), personality disorder (8% vs 4%), weight gain (6% vs 1%), akathisia (5% vs 1%), and postural hypotension (5% vs 2%).

Bipolar Disorder:

Bipolar Mania

In the bipolar mania monotherapy placebo-controlled trials, the most commonly observed adverse events associated with the use of olanzapine (incidence of $\geq 5\%$ and at least twice placebo) were: somnolence (35% vs 13%), dry mouth (22% vs 7%), dizziness (18% vs 6%), asthenia (15% vs 6%), constipation (11% vs 5%), dyspepsia (11% vs 5%), increased appetite (6% vs 3%), and tremor (6% vs 3%).

In bipolar mania combination placebo-controlled trials, the most commonly observed adverse events associated with the combination of olanzapine and lithium or valproate (incidence of $\geq 5\%$ and at least twice placebo) were: dry mouth (32% for olanzapine combination vs 9% for placebo), weight gain (26% vs 7%), increased appetite (24% vs 8%), dizziness (14% vs 7%), back pain (8% vs 4%), constipation (8% vs 4%), speech disorder (7% vs 1%), increased salivation (6% vs 2%), amnesia (5% vs 2%), and paresthesia (5% vs 2%). In addition to the latter list of adverse events identified during bipolar mania combination clinical trials tremor ($\geq 10\%$) has also been identified.

Bipolar Maintenance

In the one-year 'time to relapse' placebo-controlled clinical trial in bipolar disorder, the most commonly observed adverse events associated with olanzapine (incidence of $\geq 5\%$ and at least twice placebo) were: weight increased (8% for olanzapine vs 1.5% for placebo), headache NOS (6.7% vs 2.9%), fatigue (6.2% vs 1.5%), depression (5.8% vs 2.9%).

Other Indication Trials:

Abnormal gait and falls have been observed very commonly ($\geq 10\%$) in clinical trials with elderly patients with dementia-related psychosis. Also, urinary incontinence and pneumonia were commonly reported ($\geq 1\%$ and $< 10\%$) in these patients.

In clinical trials in patients with drug-induced (dopamine agonist) psychosis associated with Parkinson's disease, worsening of Parkinsonian symptomatology and hallucinations were reported very commonly and more frequently than with placebo.

Adverse Events Occurring at an Incidence of 1% or More Among Oral Olanzapine-Treated**Patients:**

Certain portions of the discussion below relating to objective or numeric safety parameters are derived from studies in patients with schizophrenia and have not been duplicated for bipolar disorder trials. However, this information is also generally applicable to bipolar disorder. Table 2 enumerates the incidence of treatment-emergent adverse events, rounded to the nearest percent, that occurred during acute therapy (up to 6 weeks) of schizophrenia in 1% or more of patients treated with oral OLANZAPINE-ODT (doses ≥ 2.5 mg/day) where the incidence in patients treated with OLANZAPINE-ODT was greater than the incidence in placebo-treated patients.

Table 2: Schizophrenia Trials: Treatment-Emergent Adverse Events Incidence in Placebo- Controlled Clinical Trials with Oral Olanzapine – Acute Phase¹

Body System/Adverse Event	Percentage of Patients Reporting Event	
	Olanzapine (N = 248)	Placebo (N = 118)
<i>Body As a Whole</i>		
Headache	17%	15%
Pain	10%	9%
Fever	5%	3%
Abdominal pain	4%	2%
Back pain	4%	3%
Chest pain	4%	2%
Neck rigidity	2%	1%
Intentional injury	1%	0%
<i>Cardiovascular System</i>		
Postural hypotension	5%	2%
Tachycardia	4%	1%
Hypotension	2%	1%
<i>Digestive System</i>		
Constipation	9%	3%
Dry mouth	7%	4%
Gamma glutamyl transpeptidase increased	2%	1%
Increased appetite	2%	1%
<i>Hemic and Lymphatic</i>		
Leukopenia	1%	0%
<i>Metabolic and Nutritional Disorders</i>		
SGPT increased	8%	3%
Weight gain ²	6%	1%
Edema	2%	0%
Peripheral edema	2%	0%
SGOT increased	2%	0%
Creatine phosphokinase increased	1%	0%
	Percentage of Patients Reporting Event	

Body System/Adverse Event	Olanzapine (N = 248)	Placebo (N = 118)
<i>Musculoskeletal System</i>		
Arthralgia	3%	2%
Joint disorder	2%	1%
Twitching	2%	1%
<i>Nervous System</i>		
Somnolence ²	26%	15%
Agitation	23%	17%
Insomnia	20%	19%
Nervousness	16%	14%
Hostility	15%	14%
Dizziness ²	11%	4%
Anxiety	9%	8%
Personality disorder	8%	4%
Akathisia ²	5%	1%
Hypertonia	4%	3%
Speech disorder	4%	1%
Tremor	4%	3%
Amnesia	2%	0%
Drug dependence	2%	0%
Euphoria	2%	0%
Neurosis	1%	0%
<i>Respiratory System</i>		
Rhinitis	10%	6%
Cough increased	5%	3%
Pharyngitis	5%	3%
<i>Skin and Appendages</i>		
Fungal dermatitis	2%	0%
Vesiculobullous rash	2%	1%
<i>Special Senses</i>		
Amblyopia	5%	4%
Blepharitis	2%	1%
Eye disorder	2%	1%
Corneal lesion	1%	0%
<i>Urogenital System</i>		
Menstrual disorder ³	2%	0%

¹ The following events had an incidence equal to or less than placebo: abnormal dreams, accidental injury, anorexia, apathy, asthenia, cogwheel rigidity, confusion, conjunctivitis, depression, diarrhea, dysmenorrhea³, dyspepsia, ecchymosis, emotional lability, hallucinations, hyperkinesia, hypertension, hypokinesia, libido increased, myalgia, nausea, paranoid reaction, paresthesia, pruritus, rash, schizophrenic reaction, sweating, thinking abnormal, tooth caries, vaginitis³, vomiting.

² Statistically significantly more frequent in patients treated with oral olanzapine than in patients treated with placebo.

³ Denominator used was for females only (N = 41 olanzapine; N = 23 Placebo).

Weight Changes:

During acute therapy (up to 6 weeks) in controlled clinical trials comparing OLANZAPINE-ODT with placebo in the treatment of schizophrenia, the percentages of patients with weight gain $\geq 7\%$ of baseline body weight at any time were 29% for olanzapine and 3% for placebo, which was a statistically significant difference. The average weight gain during acute therapy in patients treated with olanzapine was 2.8 kg. Clinically significant weight gain was observed across all baseline body mass index (BMI) categories. In long-term extension schizophrenia trials, there was an average gain of 5.4 kg, and 56% of olanzapine-treated patients with weight gain $> 7\%$ of baseline body weight. In long-term extension bipolar maintenance trials, there was a mean weight gain of 3.8 kg, and with 31% of olanzapine-treated patients with weight gain $> 7\%$ of baseline body weight (see [7 WARNINGS AND PRECAUTIONS, Endocrine and Metabolism](#) subsection).

8.3. Less Common Clinical Trial Adverse Reactions***Other Adverse Events from Schizophrenia Trials:***

Certain portions of the discussion below relating to objective or numeric safety parameters, namely, vital sign changes, weight gain, and ECG changes are derived from studies in patients with schizophrenia and have not been duplicated for bipolar mania. However, this information is also generally applicable to bipolar mania.

Vital Sign Changes:

In placebo-controlled clinical trials, orthostatic hypotension (greater than 30 mm decrease in systolic blood pressure) occurred with an incidence of 5% in oral OLANZAPINE-ODT treated patients compared to 2% in placebo-treated patients (vital sign measurements collected only after 3-7 days of olanzapine treatment). Oral olanzapine was associated with a mean baseline to endpoint increase in heart rate of 2.4 beats per minute compared to no change among placebo-treated patients (see [7 WARNINGS AND PRECAUTIONS, Cardiovascular](#) subsection).

ECG Changes:

Between-group comparisons for pooled placebo-controlled trials revealed no statistically significant olanzapine/placebo differences in the proportions of patients experiencing potentially important changes in ECG parameters, including QT, QTc, and PR intervals.

Other Adverse Events Observed During Clinical Trials with Olanzapine Across All Indications

The following discussion relates primarily to weight gain changes observed during clinical trials across all indications.

Weight Changes:

Weight gain has been very commonly observed in olanzapine-treated patients during clinical trials. In 13 placebo-controlled olanzapine monotherapy studies, 22.2% of olanzapine-treated patients gained $\geq 7\%$ of their baseline body weight versus 3% of placebo-treated patients, with a median exposure to event of 8 weeks; 4.2% of olanzapine-treated patients gained $\geq 15\%$ of their baseline weight versus 0.3% of placebo-treated patients, with a median exposure to event of 12 weeks. Clinically significant weight gain was observed across all baseline body mass index (BMI) categories.

In long-term studies (at least 48 weeks), both the magnitude of weight gain and the proportion of olanzapine-treated patients who had clinically significant weight gain were greater than in short-term studies. The percentage of patients who gained $\geq 25\%$ of their baseline body weight with long-term exposure was very common ($\geq 10\%$).

Prolactin:

In controlled clinical trials (up to 12 weeks), elevations in prolactin were observed in 30% of olanzapine-treated patients as compared to 10.5% of placebo-treated patients. In the majority of these patients, the elevations were mild. In patients with schizophrenia, menstrual-related adverse events potentially associated with prolactin elevations¹ were common ($< 10\%$ to $\geq 1\%$), whereas sexual function-related and breast-related adverse events were infrequent ($< 1\%$ to $\geq 0.1\%$). In patients treated for other mental illnesses², sexual function-related adverse events (erectile dysfunction, libido decreased, loss of libido, orgasm abnormal) potentially associated with prolactin elevations were common ($< 10\%$ to $\geq 1\%$), whereas breast-related and menstrual-related adverse events were infrequent ($< 1\%$ to $\geq 0.1\%$).

Vital Sign Changes:

Bradycardia was uncommonly observed in clinical trials.

Photosensitivity Reactions:

Photosensitivity reactions were uncommonly observed in clinical trials. Table 3 summarizes core adverse drug reaction terms and their frequencies identified from an integrated database of 42 completed olanzapine clinical studies in adults, consisting of 7787 patients exposed to olanzapine in placebo- or comparator-controlled clinical studies.

Table 3: Core Adverse Drug Reactions from Clinical Trials of Olanzapine

Body System/Adverse Reaction Term	Frequency				
	$\geq 10\%$	$< 10\%$ and $\geq 1\%$	$< 1\%$ and $\geq 0.1\%$	$< 0.1\%$ and $\geq 0.01\%$	$< 0.01\%$
<i>Body as a Whole</i>					
Pyrexia		X			
<i>Cardiovascular</i>					
¹ Orthostatic Hypotension	X				
<i>Digestive System</i>					
Abdominal Distension			X		
<i>Musculoskeletal System</i>					
Arthralgia		X			

¹ TEAEs analysis up to 52 weeks of treatment

² Bipolar Depression, Psychotic Depression, Borderline Personality Disorder and Bipolar Mania

Body System/Adverse Reaction Term	Frequency				
	≥ 10%	< 10% and ≥ 1%	< 1% and ≥ 0.1%	< 0.1% and ≥ 0.01%	< 0.01%
Nervous System					
Amnesia			X		
Respiratory, Thoracic and Mediastinal Disorders					
Epistaxis			X		
Laboratory Analytes					
Clinical Chemistry					
¹ Alkaline phosphatase –Increased		X			
¹ Gamma Glutamyltransferase (GGT) (U/L) - High		X			
¹ Uric Acid (μmol/L) – High		X			
Hematology					
¹ Leukopenia, including Neutropenia		X			

¹ As assessed by measured values within the clinical trial database.

Dose-Dependent Adverse Events:

Dose-relatedness of adverse events was assessed using data from a clinical trial with a fixed dosage range. Table 4 enumerates the treatment-emergent adverse events in which there was a statistically significantly increasing dose response in this clinical trial.

Table 4: Schizophrenia Trials: Dose-Dependent Adverse Events in a Fixed Dosage Range, Placebo-Controlled Clinical Trial¹ of Oral Olanzapine

Body System/ Adverse Event	Percentage of Patients Reporting Event			
	Placebo (N = 68)	Olanzapine 5 ± 2.5 mg/day (N = 65)	Olanzapine 10 ± 2.5 mg/day (N = 64)	Olanzapine 15 ± 2.5 mg/day (N = 69)
Digestive System				
Constipation	0%	6.2%	9.4%	14.5%
Nervous System				
Abnormal dreams	0%	0%	1.6%	4.3%
Dizziness	2.9%	7.7%	9.4%	17.4%
Somnolence	16.2%	20.0%	29.7%	39.1%
Respiratory System				
Pharyngitis	1.5%	3.1%	1.6%	10.1%

¹ Fungal dermatitis was also reported with a statistically significantly increasing dose response, but is not included as a drug cause was remote.

Table 5 enumerates the treatment-emergent adverse events from one 8-week, randomized, double-

blind, fixed- dose trial comparing 10 (N = 199), 20 (N = 200) and 40 (N = 200) mg/day of olanzapine in patients with schizophrenia or schizoaffective disorder. Statistically significant differences among the 3 dose groups were observed for the following safety outcomes: fatigue, dizziness, prolactin elevation, and weight gain (mean change).

Table 5: Schizophrenia Trial: Dose-Dependent Adverse Events in a Fixed Dose, Placebo-Controlled Clinical Trial of Oral Olanzapine¹

Adverse Event	Olanzapine 10 mg/day (N = 195)	Olanzapine 20 mg/day (N = 191)	Olanzapine 40 mg/day (N = 197)
Fatigue ^{2,3} (% reporting event)	1.5%	2.1%	6.6%
Dizziness ³ (% reporting event)	2.6%	1.6%	6.6%
Prolactin Elevation ^{2,3,4} (% reporting event)	31.2%	42.7%	61.1%
Prolactin Elevation ^{2,3} (mean change from baseline to endpoint)	-10.5 ng/mL	-1.7 ng/mL	4.9 ng/mL
Weight Gain $\geq$ 7% at any time (% reporting event)	14.0%	18.4%	20.5%
Weight Gain ² (mean change from baseline to endpoint)	1.9 kg	2.3 kg	3.0 kg

¹ Study HGLF: 8-week, Phase IV, parallel, randomized, double-blind, fixed-dose study in patients with schizophrenia and schizoaffective disorder evaluating the dose-response efficacy and safety of olanzapine 10, 20, and 40 mg/day. Patients were titrated up to their randomized dose over 2 weeks.

² significant difference between 10 vs. 40 mg/day

³ significant difference between 20 vs. 40 mg/day

⁴ > 24.2 ng/mL (female) or > 18.77 ng/mL (male) at any time during the trial

Incidence of Treatment-Emergent Extrapyramidal Symptoms:

Table 6 enumerates the percentage of patients with treatment-emergent extrapyramidal symptoms as assessed by categorical analyses of formal rating scales during acute therapy in a controlled clinical trial comparing oral olanzapine at 3 fixed dosage ranges with placebo in the treatment of schizophrenia.

Table 6: Schizophrenia Trials: Treatment-Emergent Extrapyramidal Symptoms Assessed By Rating Scales Incidence In A Fixed Dosage Range, Placebo-Controlled Clinical Trial -- Acute Phase¹

	Percentage of Patients			
	Placebo	Olanzapine 5 $\pm$ 2.5 mg/day	Olanzapine 10 $\pm$ 2.5 mg/day	Olanzapine 15 $\pm$ 2.5 mg/day
Parkinsonism ²	15%	14%	12%	14%
Akathisia ³	23%	16%	19%	27%

¹ No statistically significant differences.

² Percentage of patients with a Simpson-Angus Scale total score $\geq$ 3.

³ Percentage of patients with a Barnes Akathisia Scale global score ≥ 2 .

Table 7 enumerates the percentage of patients with treatment-emergent extrapyramidal symptoms as assessed by spontaneously reported adverse events during acute therapy in the same controlled clinical trial comparing oral olanzapine at 3 fixed dosage ranges with placebo in the treatment of schizophrenia. Similar results were found during the long-term (up to 1-year) double-blind monotherapy extension bipolar maintenance trial comparing olanzapine with placebo; there was a higher statistical incidence of akathisia for combined doses of olanzapine versus placebo.

Table 7: Schizophrenia Trials: Treatment-Emergent Extrapyramidal Symptoms Assessed by Adverse Events Incidence in an Oral Fixed Dosage Range, Placebo-Controlled Clinical Trial -- Acute Phase¹

Extrapyramidal Symptoms	Percentage of Patients Reporting Event			
	Placebo (N = 68)	Olanzapine 5 ± 2.5 mg/day (N = 65)	Olanzapine 10 ± 2.5 mg/day (N = 64)	Olanzapine 15 ± 2.5 mg/day (N = 69)
Dystonic events ²	1%	3%	2%	3%
Parkinsonism events ³	10%	8%	14%	20%
Akathisia events ⁴	1%	5%	11% ¹	10% ¹
Dyskinetic events ⁵	4%	0%	2%	1%
Residual events ⁶	1%	2%	5%	1%
Any extrapyramidal event	16%	15%	25%	32% ¹

¹ Statistically significantly different from placebo.

² Patients with the following COSTART terms were counted in this category: dystonia, generalized spasm, neck rigidity, oculogyric crisis, opisthotonos, torticollis.

³ Patients with the following COSTART terms were counted in this category: akinesia, cogwheel rigidity, extrapyramidal syndrome, hypertonia, hypokinesia, masked facies, tremor.

⁴ Patients with the following COSTART terms were counted in this category: akathisia, hyperkinesia.

⁵ Patients with the following COSTART terms were counted in this category: buccoglossal syndrome, choreoathetosis, dyskinesia, tardive dyskinesia.

⁶ Patients with the following COSTART terms were counted in this category: movement disorder, myoclonus, twitching.

Other Investigational Trials

Cerebrovascular Adverse Events (CVAEs) in Elderly Patients with Dementia

Data from 5 placebo-controlled trials in elderly patients with dementia-related psychosis (Alzheimer's, vascular, and mixed; olanzapine n = 1178 and placebo n = 478) suggest that there was a higher incidence of CVAE in patients treated with olanzapine compared to patients treated with placebo (1.3% vs. 0.4%, respectively). Although the incidence of CVAE was not significantly different when analyzed with Fisher's Exact Test ($p = 0.177$), the difference was found to be significant when simultaneously controlling for age, gender, and type of dementia using Poisson Regression ($p = 0.0428$). Four patients died in the olanzapine group versus 1 patient in the placebo group. In open-label safety trials studied for up to 59 weeks in dementia patients (N = 231), 7 cases of CVAEs, including 2 fatalities, were reported

(see [7 WARNINGS AND PRECAUTIONS](#) section).

Data from these trials suggest that patients with a dementia diagnosis of vascular or mixed type had a 5-fold higher likelihood of experiencing CVAEs than patients with a diagnosis of Alzheimer's. There is insufficient information to determine whether CVAEs in elderly patients with dementia are associated specifically with olanzapine or all antipsychotic agents.

OLANZAPINE-ODT is not approved for use in elderly patients with dementia.

Overall Mortality:

Elderly patients with dementia treated with atypical antipsychotic drugs showed increased mortality compared to placebo in a meta-analysis of 13 controlled trials of various atypical antipsychotic drugs. In five placebo-controlled trials with oral olanzapine in this population, the incidence of mortality was 3.5 % for olanzapine-treated patients compared to 1.5 % for placebo-treated patients. OLANZAPINE-ODT is not indicated in elderly patients with dementia.

Adverse Events in Adolescent Patients (ages 13 to 17 years)

OLANZAPINE-ODT is not indicated for pediatric use, including adolescent patients (ages 13 to 17).

The types of adverse events observed in adolescent patients treated with olanzapine were similar to those seen in adult patients. Although no clinical trials designed to compare adolescents to adults were conducted, the data from the adolescent trials were compared to those of the adult trials.

Mean increase in weight in adolescents (4.6 kg over 3 weeks median duration of exposure) was greater than in adults (2.6 kg over 7 weeks median duration of exposure).

In long-term studies (at least 24 weeks), both the magnitude of weight gain and the proportion of adolescent olanzapine-treated patients who had clinically significant weight gain were greater than in short-term studies and were greater than in adult patients with comparable exposures. With long-term exposure, approximately half of adolescent patients gained $\geq 15\%$ and almost a third gained $\geq 25\%$ of their baseline body weight. Among adolescent patients, mean weight gain was greatest in patients who were overweight or obese at baseline.

Sedation-related events are more common in adolescents (44%) as compared to adults (29%).

Core Adverse Drug Reactions and Frequencies from Clinical Trials in Adolescent Patients (ages 13 to 17 years)

Table 8 summarizes core adverse drug reaction terms for olanzapine compared to placebo and their frequencies identified only during clinical trials in adolescent patients (ages 13 to 17 years).

Table 8: Core Adverse Drug Reactions and Frequencies from Clinical Trials in Adolescent Patients (ages 13 to 17 years)

Body System/Adverse Reaction Term	Olanzapine		Placebo	
	Frequency	N	Frequency	N
Body as a Whole				
Weight gain $\geq$ 7% of baseline body weight (kg) ²	40.6%	197	9.8%	112
Weight gain $\geq$ 15% of baseline body weight (kg) ³	7.1%	197	2.7%	112
Digestive System				
Dry Mouth	6.15%	179	0%	89
Increased Appetite	24%	179	6%	89
Nervous System				
Sedation ¹	44.1%	179	9%	89

¹ Represented cluster of MedDRA terms including: hypersomnia, lethargy, sedation, somnolence.

² Median duration of exposure to event = 4 weeks

³ Median duration of exposure to event = 19 weeks

8.4. Abnormal Laboratory Findings: Hematologic, Clinical Chemistry and Other Quantitative Data

Lipids:

In placebo-controlled clinical trials of up to 12 weeks in duration, olanzapine-treated patients had a greater mean increase in fasting total cholesterol, LDL cholesterol, and triglycerides compared to placebo-treated patients.

Mean increases in fasting lipid values (total cholesterol, LDL cholesterol, and triglycerides) were greater in patients without evidence of lipid dysregulation at baseline. However, for mean changes in fasting triglycerides, the difference between olanzapine and placebo was greater in patients with evidence of lipid dysregulation at baseline. Elevations in fasting triglyceride levels $\geq$ 11.3 mmol/L were uncommonly observed with olanzapine use (8 week median duration of exposure).

Table 9: Changes in Fasting Lipids Values from Adult Placebo-Controlled Olanzapine Monotherapy Studies with Treatment Duration up to 12 Weeks

Laboratory Analyte	Olanzapine*	Placebo
Triglycerides: fasting normal to high (< 1.70 mmol/L to $\geq$ 2.26 mmol/L)	9.2% (N = 457)	4.4% (N = 251)

Laboratory Analyte	Olanzapine*	Placebo
Triglycerides: fasting borderline to high (≥ 1.70 mmol/L and < 2.26 mmol/L to ≥ 2.26 mmol/L)	39.3% (N = 135)	20.0% (N = 65)
Cholesterol-Total: fasting normal to high (< 5.18 mmol/L to ≥ 6.22 mmol/L)	2.8% (N = 392)	2.4% (N = 207)
Cholesterol-Total: fasting borderline to high (≥ 5.18 mmol/L and < 6.22 mmol/L to ≥ 6.22 mmol/L)	23.0% (N = 222)	12.5% (N = 112)
LDL cholesterol: fasting normal to high (< 2.59 mmol/L to ≥ 4.14 mmol/L)	0% (N = 154)	1.2% (N = 82)
LDL cholesterol: fasting borderline to high (≥ 2.59 mmol/L and < 4.14 mmol/L to ≥ 4.14 mmol/L)	10.6% (N = 302)	8.1% (N = 173)

* Median duration of exposure 8 weeks.

For fasting HDL cholesterol, no statistically significant differences were observed between olanzapine-treated patients and placebo-treated patients (see [7 WARNINGS AND PRECAUTIONS, Endocrine and Metabolism](#) subsection).

The proportion of patients who had changes in total cholesterol, LDL cholesterol or triglycerides from normal or borderline to high, or changes in HDL cholesterol from normal or borderline to low, was greater in long-term studies (at least 48 weeks) as compared with short-term studies. Treatment-emergent clinically significant changes in fasting lipids were observed in patients with or without evidence of dyslipidemia at baseline.

Glucose

In clinical trials (up to 52 weeks) olanzapine was associated with a greater mean change in glucose relative to placebo.

The difference in mean changes between olanzapine and placebo was greater in patients with evidence of glucose dysregulation at baseline (including those patients diagnosed with diabetes mellitus or who met criteria suggestive of hyperglycemia), and these patients had a greater increase in HbA1c compared to placebo.

In patients with baseline normal fasting glucose levels (< 5.5 mmol/L), 2.2% (N = 543) of those treated with olanzapine (median exposure duration of 8 weeks) were found to have high glucose levels (≥ 6.99 mmol/L) during olanzapine treatment versus 3.4% (N = 293) of those treated with placebo. In patients with baseline borderline fasting glucose levels (≥ 5.5 mmol/L and < 6.99 mmol/L), 17.4% (N = 178) of those treated with olanzapine (median exposure duration of 5 weeks) were found to have high glucose levels (≥ 6.99 mmol/L) during olanzapine treatment versus 11.5% (N = 96) of those treated with placebo.

The proportion of patients who had a change in glucose level from normal or borderline at baseline to high increased over time. Treatment-emergent clinically significant changes in fasting glucose

were observed in patients with or without evidence of glucose dysregulation at baseline.

Glycosuria was commonly reported in olanzapine-treated patients during clinical trials.

Other Laboratory Changes

Olanzapine is associated with asymptomatic increases in SGPT, SGOT, and GGT (see [7 WARNINGS AND PRECAUTIONS, Hepatic](#) subsection). Olanzapine is also associated with generally mild increases in serum prolactin, which usually decreases with continued drug treatment. Olanzapine is also associated with asymptomatic elevations of eosinophils and uric acid (see [7 WARNINGS AND PRECAUTIONS, Renal](#) subsection), and with decreases in serum bicarbonate.

In 5 double-blind, placebo controlled clinical trials of elderly patients with dementia-related psychosis (n = 1184 total in the olanzapine arms and n = 478 total in placebo arms), olanzapine-treated patients showed significantly greater incidence rates compared to placebo-treated patients of low albumin (10.4% vs 5.5%, respectively), low hemoglobin (4.2% vs 1.8%) and low hematocrit (4.6% vs 2.4%). Of patients who had low albumin values, 3.6% in the olanzapine-treated group vs 1.4% in the placebo-treated also experienced a treatment-emergent respiratory infection. A causal relationship between the two adverse events has not been determined.

Mean increases in fasting glucose were similar in adolescents and adults treated with olanzapine; however, the difference between olanzapine and placebo groups was greater in adolescents compared to adults.

In long-term studies (at least 24 weeks), changes in glucose from normal at baseline to high were uncommon (< 1% and ≥ 0.1%).

Mean increases in fasting total cholesterol, LDL cholesterol, and triglycerides were generally greater in adolescents than in adults treated with olanzapine. However, in short term trials, the differences between olanzapine and placebo were similar for adolescents and adults.

The proportion of treatment-emergent clinically significant changes in normal-to-high or borderline-to-high fasting total cholesterol, LDL cholesterol and triglycerides was greater in adolescents compared to adults, and the differences between olanzapine and placebo in these categories of laboratory values were also generally greater in adolescents. In long-term studies, treatment-emergent clinically significant changes in total cholesterol, LDL cholesterol, and triglycerides were observed in adolescents with or without evidence of dyslipidemia at baseline.

Compared with adults, adolescents treated with olanzapine experienced a significantly higher incidence of elevated prolactin levels (47% in olanzapine-treated adolescents vs 30% in olanzapine-treated adults) and significantly higher mean increases in prolactin levels.

Hepatic aminotransferase elevations (3 times the Upper limit of Normal) are more common in adolescents (12.1%) as compared to adults (5.4%).

Table 10: Adverse Drug Reactions related to Clinical Chemistry and their Frequencies from Clinical Trials in Adolescent Patients (ages 13 to 17 years)

Clinical Chemistry				
ALT/SGPT > 3X ULN all randomized patients with ALT baseline ≤ 3X ULN ¹	12.1%	174	2.3%	87
Clinical Chemistry				
AST/SGOT – Increased ²	27.6%	163	3.8%	79
Total bilirubin – Decreased ³	22.1%	154	6.7%	75
GGT – Increased ⁴	10.1%	169	1.2%	83
Prolactin – Increased ⁵	47.4%	116	6.8%	59
Cholesterol – total, fasting normal to high (< 4.40 mmol/L to ≥ 5.18 mmol/L) ⁶	6.9%	87	2.3%	43
Cholesterol – total, fasting borderline to high (≥ 4.40 mmol/L and < 5.18 mmol/L to ≥ 5.18 mmol/L) ⁶	38.9 %	36	7.7%	13
LDL cholesterol: fasting normal to high (< 2.85 mmol/L to ≥ 3.37 mmol/L)	5.1%	98	4.5%	44
LDL cholesterol: fasting borderline to high (≥ 2.85 mmol/L and < 3.37 mmol/L to ≥ 3.37 mmol/L)	48.3%	29	0%	9
Triglycerides, fasting normal to high (< 1.02 mmol/L to > 1.47 mmol/L) ⁶	26.9%	67	10.7%	28
Triglycerides, fasting borderline to high (≥ 1.02 mmol/L and ≤ 1.47 mmol/L to > 1.47 mmol/L) ⁶	59.5%	37	35.3%	17
Glucose, fasting normal to high (< 5.55 mmol/L to ≥ 6.99 mmol/L) ⁶	0%	124	1.9%	53
Glucose, fasting borderline to high (≥ 5.55 mmol/L and < 6.99 mmol/L to ≥ 6.99 mmol/L) ⁷	14.3%	14	0%	13

- ¹ Covance reference ranges:
- | | | | |
|-------|----------------------|-----|------|
| (U/L) | Female 13 - ≤ 17.999 | Low | High |
| (U/L) | Male 13 - ≤ 17.99 | 6 | 34 |
- ² Covance reference ranges:
- | | | | |
|-------|----------------------|-----|------|
| (U/L) | Female 13 - ≤ 17.999 | Low | High |
| (U/L) | Male 13 - ≤ 17.99 | 6 | 43 |
- ³ Covance reference ranges:
- | | | | |
|----------|----------------------|-----|------|
| (mmol/L) | Female 13 - ≤ 17.999 | Low | High |
| (mmol/L) | Male 13 - ≤ 17.99 | 10 | 40 |
- ⁴ Covance reference ranges:
- | | | | |
|-------|----------------------|-----|------|
| (U/L) | Female 13 - ≤ 17.999 | Low | High |
| (U/L) | Male 13 - ≤ 17.99 | 10 | 40 |
- ⁵ Covance reference ranges for prolactin as published by Wiedemann and Jonetz-Mentzel (1993)
- | | | |
|---------|-----------------|--------------|
| Female: | 12 to 14 years: | 2.52 – 16.90 |
| | | ng/mL |
| | 14 to 19 years: | 4.20 – 39.00 |
| | | ng/mL |
| Male: | 12 to 14 years: | 2.84 – 24.00 |
| | | ng/mL |

14 to 19 years:

2.76 – 16.10
ng/mL⁶ Median duration of exposure was 3 weeks⁷ Median duration of exposure was 5 weeks

Health Canada has not authorized an indication for pediatric use.

8.5 Post-Market Adverse Reactions

Table 11 summarizes core adverse drug reaction terms and their frequencies identified from global post-marketing surveillance in addition to what was reported in clinical trials (see preceding section [8 ADVERSE REACTIONS, 8.2 Clinical Trial Adverse Reactions](#)). A causal relationship between olanzapine and the emergence of these events has not been established.

Table 11: Core Adverse Drug Reactions Seen with Olanzapine Formulations¹

Body System/Adverse Reaction Term	Frequency				
	≥ 10%	< 10% and ≥ 1%	< 1% and ≥ 0.1%	< 0.1% and ≥ 0.01%	< 0.01%
Body as a Whole					
Allergic reaction ²					X
Discontinuation reaction ³					X
Cardiovascular					
Venous Thromboembolism, including Pulmonary Embolism and Deep Vein Thrombosis					X
Digestive System					
Pancreatitis					X
Salivary Hypersecretion ⁸			X		
Hematologic					
Thrombocytopenia ⁴					X
Hepatobiliary disorders					
Hepatitis				X	
Jaundice					X
Hepatic failure					X
Metabolic					
Diabetic Coma					X
Diabetic Ketoacidosis ⁵					X
Hypercholesterolemia ⁷					X
Hyperglycaemia				X	
Hypertriglyceridemia ^{6,7}					X
Exacerbation of pre-existing diabetes				X	
Musculoskeletal System					
Rhabdomyolysis					X
Nervous System					
Restless Legs Syndrome (RLS) ⁸			X		

Body System/Adverse Reaction Term	Frequency				
	≥ 10%	< 10% and ≥ 1%	< 1% and ≥ 0.1%	< 0.1% and ≥ 0.01%	< 0.01%
Seizures				X	
Stuttering ^{1,9}			X		
<i>Skin and Appendages</i>					
Alopecia					X
Rash				X	
Drug Reaction with Eosinophilia and Systemic Symptoms (DRESS)					X
<i>Urogenital System</i>					
Priapism					X
Urinary Incontinence					X
Urinary Retention					X
<i>Laboratory Analytes</i>					
<i>Clinical Chemistry</i>					
Total bilirubin –Increased					X

¹ Adverse event identified from spontaneous post-marketing surveillance.

² e.g., maculopapular rash, anaphylactoid reaction, angioedema, pruritis, or urticaria.

³ i.e., diaphoresis, nausea or vomiting.

⁴ Including a case of thrombocytopenic purpura.

⁵ COSTART term is diabetic acidosis.

⁶ COSTART term is hyperlipemia.

⁷ Random cholesterol levels of ≥ 6.22 mmol/L and random triglyceride levels of ≥ 11.30 mmol/L have been very rarely reported.

⁸ Adverse event identified from spontaneous post-marketing reporting with frequency determined using the olanzapine clinical trial database.

⁹ Stuttering was only studied in oral formulations and the review did not include details about the rapid IM formulation.

As with other atypical anti-psychotics, there have been isolated post-market reports with olanzapine of serious cardiovascular-related adverse events, including fatalities (see [7 WARNINGS AND PRECAUTIONS, Cardiovascular](#)).

Neutropenia, granulocytopenia and agranulocytosis have been reported during antipsychotic use. Therefore, it is recommended that patients have their complete blood count (CBC) tested prior to starting OLANZAPINE-ODT and then periodically throughout treatment.

Venous Thromboembolism (VTE), including fatal pulmonary embolism, has been reported with antipsychotic drugs, including olanzapine. However, since patients who require treatment with antipsychotics often present with acquired risk factors for VTE all possible risk factors of VTE e.g. immobilization, should be identified and preventative measures undertaken.

Patients should be advised of the risk of severe constipation during OLANZAPINE-ODT treatment, and that they should tell their doctor if constipation occurs or worsens, as they may need laxatives.

Atypical antipsychotic drugs, including olanzapine, have been associated with cases of sleep apnea, with or without concomitant weight gain. In patients who have a history of or are at risk for sleep apnea, OLANZAPINE-ODT should be prescribed with caution.

Risks of somnambulism (sleep walking) and sleep-related eating disorder have been associated with the use of atypical antipsychotics including olanzapine.

Cases of syndrome of inappropriate secretion of antidiuretic hormone (SIADH) and hyponatremia have been identified during the post-marketing period.

9. DRUG INTERACTIONS

9.2 Drug Interactions Overview

The metabolism of olanzapine may be affected by inhibitors or inducers of the P450 cytochrome isoforms, specifically CYP1A2 activity. Olanzapine clearance was increased by smoking or carbamazepine coadministration. Smoking and carbamazepine therapy are known to induce CYP1A2 activity. Known potent inhibitors of CYP1A2 activity may decrease olanzapine clearance. Olanzapine is not a potent inhibitor of CYP1A2 activity. The pharmacokinetics of theophylline, a drug principally metabolized by CYP1A2, are not altered by olanzapine.

9.3 Drug-Behaviour Interactions

Alcohol: Given the primary CNS effects of olanzapine, caution should be used when it is taken in combination with other centrally acting drugs and alcohol since additive pharmacological effects such as increased sedation may occur.

Smoking: Concomitant smoking may induce the metabolism of olanzapine.

9.4 Drug-Drug Interactions

The drugs listed in this Table 12 are based on either drug interaction case reports or studies, or potential interactions due to the expected magnitude and seriousness of the interaction (i.e., those identified as contraindicated).

Table 12: Established or Potential Drug-Drug Interactions

Proper/Common name	Source of Evidence	Effect	Clinical comment
Activated Charcoal	CT	The concomitant administration of activated charcoal reduced the oral bioavailability of olanzapine by 50% to 60%.	Activated charcoal may be a useful agent to reduce olanzapine absorption in situations of overdose (see 5 OVERDOSE).

Proper/Common name	Source of Evidence	Effect	Clinical comment
Antacids, e.g. aluminium, magnesium	CT	No effect	Single doses of antacid (aluminium, magnesium) did not affect the oral bioavailability of olanzapine.
Antihypertensive Agents	T	N/A	Because of its potential for inducing hypotension, olanzapine may enhance the effects of certain antihypertensive agents. Caution should be exercised in patients who receive medicinal products that can induce hypotension, bradycardia, or respiratory depression.
Carbamazepine	CT	Olanzapine C _{max} , AUC, and t _{1/2} were consistently lower when given after carbamazepine.	Concomitant carbamazepine therapy may induce the metabolism of olanzapine.
Cimetidine	CT	No effect	Single doses of cimetidine did not affect the oral bioavailability of olanzapine.
CYP1A2 Inducers	In vitro	Population modeling has suggested that clearance of olanzapine correlates with the activity of CYP1A2.	Agents that induce CYP1A2 such as omeprazole may increase clearance of olanzapine.
CYP1A2 Inhibitors, e.g. fluvoxamine, ciprofloxacin, ketoconazole	CT	Fluvoxamine, a specific CYP1A2 inhibitor, has been shown to significantly inhibit the metabolism of olanzapine. The mean increase in olanzapine C _{max} following fluvoxamine was 54% in female nonsmokers and 77% in male smokers. The mean increase in olanzapine AUC was 52% and 108%, respectively.	A lower starting dose of olanzapine should be considered in patients who are using fluvoxamine or any other CYP1A2 inhibitors, such as ciprofloxacin or ketoconazole. A decrease in the dose of olanzapine should be considered if treatment with an inhibitor of CYP1A2 is initiated.

Proper/Common name	Source of Evidence	Effect	Clinical comment
Diazepam	CT	No effect	In clinical trials with single doses of oral olanzapine, no inhibition of the metabolism of diazepam (P450 CYP3A4) was evident.
Dopamine Agonists, Levodopa Agonists	T		As it exhibits <i>in vitro</i> dopamine antagonism, olanzapine may antagonize the effects of levodopa and dopamine agonists.
Drugs Metabolized via P450-CYP1A2, -CYP2C9, - CYP2C19, -CYP2D6, and - CYP3A	In vitro	In <i>in vitro</i> studies with human microsomes, olanzapine showed little potential to inhibit cytochromes P450-CYP1A2, -CYP2C9, - CYP2C19, - CYP2D6, and - CYP3A (see 10 CLINICAL PHARMACOLOGY).	Olanzapine is thus unlikely to cause clinically important drug-drug interactions mediated through the metabolic routes described. However, the possibility that olanzapine may alter the metabolism of other drugs, or that other drugs may alter the metabolism of olanzapine, should be considered when prescribing OLANZAPINE-ODT.
Fluoxetine	CT	Fluoxetine (60 mg single dose or 60 mg daily for 8 days) causes a mean 16% increase in the maximum concentration of olanzapine and a mean 16% decrease in olanzapine clearance.	The magnitude of the impact of this factor is small in comparison to the overall variability between individuals, and therefore dose modification is not routinely recommended
Imipramine/Desipramine	CT	No effect	In clinical trials with single doses of oral olanzapine, no inhibition of the metabolism of imipramine/desipramine (P450-CYP2D6) was evident.
Lithium or Biperiden	CT	No effect	Oral olanzapine showed no interaction when co-administered with lithium or biperiden.

Proper/Common name	Source of Evidence	Effect	Clinical comment
Theophylline	CT	No effect	The pharmacokinetics of theophylline, a drug principally metabolized by CYP1A2, were not altered by olanzapine in a clinical trial with single doses of IV theophylline.
Valproate	In vitro	Studies <i>in vitro</i> using human liver microsomes showed that olanzapine has little potential to inhibit the glucuronidation of valproate, which is the major metabolic pathway. Furthermore, valproate was found to have little effect on the metabolism of olanzapine <i>in vitro</i> . Daily concomitant <i>in vivo</i> administration of 10 mg olanzapine for 2 weeks did not affect steady state plasma concentrations of valproate.	Therefore, concomitant olanzapine administration does not require dosage adjustment of valproate.
Warfarin	CT	No effect	In clinical trials with single doses of oral olanzapine, no inhibition of the metabolism of warfarin (P450 CYP2C9) was evident.

Legend: CT = Clinical Trial; T = Theoretical

9.5. Drug-Food Interactions

Absorption of olanzapine is not affected by food.

9.6. Drug-Herb Interactions

Interactions with herbal products have not been identified.

9.7. Drug-Laboratory Test Interactions

Interactions with laboratory tests have not been identified.

10. CLINICAL PHARMACOLOGY

10.1. Mechanism of Action

Olanzapine, a thienobenzodiazepine, is an antipsychotic agent that demonstrates a broad pharmacologic profile across a number of receptor systems. In preclinical studies, olanzapine exhibited affinities for serotonin 5HT_{2A/C}, 5HT₃, 5HT₆; dopamine D₁, D₂, D₃, D₄, D₅; muscarinic M₁₋₅; adrenergic α_1 ; and histamine H₁ receptors.

10.2. Pharmacodynamics

Pharmacodynamic Properties:

Olanzapine displays high receptor affinity binding *in vitro* at dopamine D₂, D₃, D₄ (K_i = 11-31 nM), 5-HT_{2A/C} (K_i = 4 and 11 nM, respectively), 5-HT₃, 5-HT₆, muscarinic M₁₋₅ (K_i = 1.9-2.5 nM), adrenergic α_1 (K_i = 19 nM), and histamine H₁ (K_i = 7 nM) receptor subtypes, while displaying a lower affinity at dopamine D₁ and D₅ receptor subtypes (K_i = 51 – 119 nM). In a behavioural paradigm predictive of antipsychotic activity, olanzapine reduced conditioned avoidance response in rats at doses lower than 4 times those required to produce catalepsy. In a single dose (10 mg) PET study in healthy subjects, olanzapine produced higher 5-HT_{2A} than dopamine D₂ receptor occupancy. The percent of D₂ occupancy was less than the threshold value predictive of extrapyramidal events.

In animals olanzapine has been observed to produce a significant reduction in the firing of A10 dopaminergic cells. The number of spontaneously active A9 neurons either remained constant or was increased. This may explain the low incidence of extrapyramidal side effects with olanzapine usually associated with the typical antipsychotics.

Olanzapine also increases extracellular levels of dopamine in a regionally specific manner in the prefrontal cortex, similar to mood stabilizers, lithium and valproate.

In Vitro Receptor Binding Affinities:

The binding affinities of olanzapine versus clozapine and haloperidol are summarized in *Table 13*. The binding profile of olanzapine has similarities to that produced by clozapine, although the affinity of olanzapine is somewhat greater for dopamine D₁ and D₂ receptors and lower at 2 receptors. With respect to 5-HT receptor subtypes, both agents show greatest affinity for 5-HT_{2A} and 5-HT_{2C} receptors. The ratio of activity between 5-HT_{2A} and D₂ receptors is slightly less for olanzapine than for clozapine, although olanzapine is still about twice as active at 5-HT_{2A} receptors compared with D₂ receptors. Both compounds also have a high affinity for muscarinic receptor subtypes, particularly the m₁ site. The affinity constants (K_i , nM) for olanzapine, clozapine, and haloperidol are shown below:

Table 13: Affinity constants for olanzapine, clozapine, and haloperidol

Compound	Dopamine D ₁	Dopamine D ₂	α_1	α_2	Histamine H ₁
Olanzapine	31 ± 0.7	11 ± 2	19 ± 1	230 ± 40	7 ± 0.3
Clozapine	85 ± 0.7	125 ± 20	7 ± 4	8 ± 3	6 ± 2
Haloperidol	25 ± 7	1 ± 0.04	46 ± 6	360 ± 100	3630 ± 85

Compound	5-HT _{1A}	5-HT _{1B}	5-HT _{1D}	5-HT _{2A}	5-HT _{2C}	5-HT ₃
Olanzapine	>10,000	1355 ± 380	800 ± 190	4 ± 0.4	11 ± 1	57
Clozapine	770 ± 220	1200 ± 170	980 ± 115	12 ± 3	8 ± 0.8	69
Haloperidol	7930 ± 500	>10,000	6950 ± 950	78 ± 22	3085	>1000

Compound	m ₁	m ₂	m ₃	m ₄	m ₅
Olanzapine	1.9 ± 0.1	18 ± 5	25 ± 2	13 ± 2	6 ± 0.8
Clozapine	1.9 ± 0.4	10 ± 1	14 ± 1	18 ± 5	5 ± 1.2
Haloperidol	1475 ± 300	1200 ± 180	1600 ± 305	>10,000	Not Tested

Olanzapine has no significant activity at GABA_A, benzodiazepine, or β receptors. Olanzapine also interacts with dopamine D₄ receptors (K_i 27 nM).

In vivo biochemical studies were conducted to confirm the binding data and investigate the functional consequences of interacting with these neurotransmitter receptor sites.

In Vivo Neuroendocrine Studies:

It has been shown that corticosterone concentrations in rats can be elevated by 5-HT-mediated or dopamine-mediated mechanisms. Olanzapine antagonizes the 5-HT (quipazine-induced) (ED₅₀ 0.57 mg/kg) and D₂ dopamine receptor-mediated (pergolide-induced) (ED₅₀ 3 mg/kg) increases in corticosterone. These results show that olanzapine has greater activity at 5-HT compared with D₂ dopamine receptors *in vivo*. These results complement the behavioural studies showing that olanzapine preferentially antagonizes a 5-HT-induced response.

In Vivo Behavioural Pharmacology:

In behavioural studies, olanzapine exhibits a broad pharmacologic profile, as predicted from the biochemical data.

Olanzapine blocks apomorphine induced climbing behaviour with an ED₅₀ of approximately 5 mg/kg.

The climbing response has previously been shown to require both D1 and D2 receptor activation. These results therefore indicate that olanzapine possesses dopamine antagonist activity in vivo.

A second study in mice looked at the ability of olanzapine to block 5 hydroxytryptophan (5 HTP) induced head twitches, a test probably mediated by 5 HT₂ receptors. Olanzapine produced dose related reductions in the head twitch response with approximate ED₅₀s of 2 mg/kg. Olanzapine preferentially blocks the head twitch, compared with the climbing response, demonstrating that this agent exhibits greater activity at the 5 HT receptor compared with dopamine receptors in vivo. These results agree with those reported in rats, showing that olanzapine preferentially antagonizes 5 HT mediated rather than dopamine mediated elevations in corticosterone (Moore et al. 1993).

Olanzapine doses of 2.5 to 10 mg/kg produced a significant reduction in oxotremorine induced tremor in mice, with an ED₅₀ of 3 mg/kg. These results demonstrate that olanzapine possesses anticholinergic activity in vivo at doses which also antagonize dopamine mediated effects.

Inhibition of a conditioned avoidance response has been widely used as a test to predict the antipsychotic potential of a compound, while the induction of catalepsy in rats is associated with the occurrence of extrapyramidal symptoms in the clinic. ED₅₀s for the various compounds in blocking a conditioned avoidance response or inducing catalepsy in rats are given in Table 14.

Table 14 Effect of Olanzapine and Haloperidol on Conditioned Avoidance Responding (CAR) and the Induction of Catalepsy (CAT) in Lister Hooded Rats

Compound	CAR	CAT	Ratio
Olanzapine	5.6 (4.6-6.8)	23 (18.7-29)	4.1
Haloperidol	0.28 (0.24-0.33)	0.74 (0.6-0.9)	2.6

Note: The results are expressed as ED₅₀ values (mg/kg p.o.) with 95% confidence intervals stated in parentheses. The ratio is the ED₅₀ CAT / ED₅₀ CAR.

Although olanzapine induces catalepsy, this only occurs at doses higher than those required to block the conditioned avoidance response.

A number of reports have shown that the "atypical" agent, clozapine, differs from "typical" antipsychotics in its effects on schedule controlled behaviour. In a rat or pigeon conflict test, olanzapine, clozapine, and chlordiazepoxide produced the characteristic changes in rates of responding associated with anxiolytics, although the effect of olanzapine and clozapine was smaller than that seen with chlordiazepoxide. All three compounds decreased or had no effect on the high rates of responding produced in the reward component, whereas the rates in time out and particularly the conflict period were increased. This type of profile was not seen with the "typical" antipsychotic, haloperidol, which only decreased the rates in all the components. These data further emphasize the "atypical" profile of olanzapine.

In Vivo Electrophysiology:

"Typical" antipsychotic agents, such as haloperidol, reduce the spontaneous firing of both A9 and A10 dopaminergic neurons in the CNS following chronic dosing. The A9 (nigrostriatal system) is

thought to mediate extrapyramidal motor disturbances, while the A10 (mesolimbic system) has been associated with the antipsychotic activity of compounds. Olanzapine (10 and 20 mg/kg subcutaneously for 21 days) produced a significant reduction in the firing of A10 dopaminergic cells. The number of spontaneously active A9 neurons either remained constant or was increased. These results are very similar to those reported previously for clozapine and further emphasize the “atypical” pharmacologic profile of olanzapine.

Human Versus Animal Metabolism:

In animal species (mice, rats, and dogs) used for toxicologic evaluation, olanzapine was metabolized through aromatic hydroxylation (forming phenolic metabolites and/or their glucuronide conjugates), allylic (alkyl) oxidation, N dealkylation, and N oxidation reactions.

Although similarities in the metabolic fate of olanzapine in animals (mice, rats, and dogs) and humans include the 2 alkyl hydroxylation, N dealkylation, and N oxidation pathways, two significant differences can be noted. First, direct glucuronidation, producing mainly 10 N glucuronide and to a lesser extent 4' N glucuronide, was a significant metabolic pathway in humans. These N glucuronides were absent in animal species except for a trace amount of 10 N glucuronide in dog urine. Second, metabolites resulting from aromatic oxidation were not found in any human biological fluids. The monkey also did not appear to form 10 N glucuronide, but was similar to humans in apparently not forming metabolites resulting from the oxidative attack of the benzene ring of olanzapine.

10.3 Pharmacokinetics

In the following Table 15, a summary of clinically relevant pharmacokinetic parameters for Olanzapine can be found.

Table 15: Summary of Olanzapine Pharmacokinetic Parameters in Healthy Subjects

Statistic	$t_{1/2}$ (h)	CL (L/h)	Vd (L)
Number of observations	491	491	479
Mean	33.1	26.1	1,148
Median	30.5	23.6	1,091
5th Percentile	20.7	12.0	660
95th Percentile	54.1	46.9	1,792
Minimum	14.5	7.1	400
Maximum	79.5	142.0	2,438

CL, Plasma Clearance; $t_{1/2}$, Elimination Half-Life; Vd, Volume of Distribution.

Estimates of Olanzapine pharmacokinetic parameters were pooled selectively across 15 clinical pharmacology studies (single dose studies only) comprising a total of 193 subjects.

Oral Administration:

Table 16: Olanzapine Key Pharmacokinetics

Patient Characteristics	Half-Life (hours)	Plasma Clearance (L/hr)
Nonsmoking	38.6	18.6
Smoking	30.4	27.7
Female	36.7	18.9
Male	32.3	27.3
Elderly (65 and older)	51.8	17.5
Non-elderly	33.8	18.2

Although smoking status, gender, and, to a lesser extent, age may affect olanzapine clearance and half-life, the magnitude of the impact of these single factors is small in comparison to the overall variability between individuals.

Absorption:

Olanzapine is well absorbed after oral administration, reaching peak plasma concentrations within 5 to 8 hours. The absorption is not affected by food.

Distribution:

Plasma concentrations of orally administered olanzapine were linear and dose proportional in trials studying doses from 1 to 20 mg. The maximum plasma concentrations (C_{max}) of olanzapine after single oral doses of 5, 10 and 15 mg averaged 7, 14, and 21 ng/mL, respectively (20 ng/mL = 0.064 mcM). In young healthy volunteers, after once-a-day repeated dosing, steady-state C_{max} was approximately twice that achieved after a single dose (e.g., 23 ng/mL versus 12 ng/mL for a 10-mg dose). In the elderly, the steady state plasma concentration was approximately 3-fold higher than that achieved after a single dose (e.g., 16 ng/mL versus 5 ng/mL for a 5-mg dose). In both, young and elderly, steady-state concentrations of olanzapine were obtained after seven days of once daily dosing.

Over time and dosage range, pharmacokinetic parameters within an individual are very consistent. However, plasma concentrations, half-life and clearance of olanzapine may vary between individuals on the basis of smoking status, gender, and age (see [7 WARNINGS AND PRECAUTIONS, 7.1 Special Populations](#)). Data from pooled, single dose pharmacokinetic studies showed the half-life of olanzapine to range from 21 to 54 hours (5th to 95th percentile), and the apparent plasma clearance to range from 12 to 47 L/hr (5th to 95th percentile).

The plasma protein binding of olanzapine was about 93% over the concentration range of about 7 to about 1000 ng/mL. Olanzapine is bound predominantly to albumin and α 1-acid glycoprotein.

Metabolism:

Olanzapine is metabolized in the liver by conjugative and oxidative pathways. The major circulating metabolite is the 10-N-glucuronide, which is pharmacologically inactive and does not pass the blood brain barrier. Cytochrome P450 isoforms CYP1A2 and CYP2D6 contribute to the formation of the N-desmethyl and 2-hydroxymethyl metabolites. Both metabolites exhibited significantly less *in vivo* pharmacological activity than olanzapine in animal studies. The predominant pharmacologic activity is from the parent olanzapine.

In vitro microsomal studies show that olanzapine is a weak inhibitor of CYP1A2 (K_i = 36 mcM),

CYP2D6 ($K_i = 89 \text{ mcM}$), and CYP3A4 ($K_i = 490 \text{ mcM}$). Based upon these K_i values, little inhibition of these cytochrome P-450 enzymes is expected *in vivo* at concentrations below 5 mcM (roughly 1500 ng/mL) because the olanzapine concentration will be less than 10% of its K_i value. In clinical studies, observed steady-state plasma concentrations of olanzapine are rarely > 150 ng/mL (approximately 0.5 mcM). Olanzapine is thus not likely to cause clinically important pharmacokinetic drug-drug interactions mediated through the metabolic routes outlined above. (See [9 DRUG INTERACTIONS](#) section).

Elimination:

After oral administration to healthy subjects, the mean terminal elimination half-life was 33 hours (21 to 54 hours for 5th to 95th percentile) and the mean olanzapine plasma clearance was 26 L/hr (12 to 47 L/hr for the 5th to 95th percentile). Olanzapine pharmacokinetics varied on the basis of smoking status, gender, and age.

Special Populations and Conditions

Geriatrics: In a study involving 24 healthy subjects, the mean elimination half-life of olanzapine was about 1.5 times greater in elderly (> 65 years) than in non-elderly subjects (≤ 65 years). Caution should be used in dosing the elderly, especially if there are other factors that might additively influence drug metabolism and/or pharmacodynamic sensitivity (see [4 DOSAGE AND ADMINISTRATION](#) section).

Sex: Clearance of olanzapine is approximately 30% lower in women than in men. There were, however, no apparent differences between men and women in effectiveness or adverse effects. Dosage modifications based on gender should not be needed.

Ethnic origin: In a study of Caucasians, Japanese, and Chinese subjects, there were no differences in olanzapine pharmacokinetics among the three populations. Cytochrome P450 isoform CYP2D6 status does not affect the metabolism of olanzapine.

Hepatic Insufficiency: No differences in the single-dose pharmacokinetics of oral olanzapine were noted in subjects with clinically significant cirrhosis (who were mostly smokers) when compared to healthy subjects (all non-smokers). Multiple-dose studies in patients with hepatic impairment, however, have not been performed.

Renal Insufficiency: There was no significant difference in mean elimination half-life or olanzapine plasma clearance between subjects with severely impaired renal function compared to individuals with normal renal function. Approximately 57% of radio-labelled olanzapine is excreted in urine, principally as metabolites.

11. STORAGE AND STABILITY

Store at 15° to 30°C. Protect from light and moisture.

12. SPECIAL HANDLING INSTRUCTIONS

OLANZAPINE-ODT should be handled carefully with dry hands.


PART II: SCIENTIFIC INFORMATION**13. PHARMACEUTICAL INFORMATION****Drug Substance**

Proper name: Olanzapine

Chemical name: 2-Methyl-4-(4-methyl-1-piperazinyl)-10H-thieno[2,3-b] [1,5] benzodiazepine

Molecular formula and molecular mass: C₁₇H₂₀N₄S (312.44 g/mol)

Structural formula:



Physicochemical properties:

Description:	Olanzapine is an antipsychotic agent of the thienobenzodiazepine class. It is a yellow crystalline powder, which is practically insoluble in water, freely soluble methylene chloride, slightly soluble in methanol and in ethanol (96%).
pKa:	5.0 and 7.4
Melting Point:	190°C - 193°C

14. CLINICAL TRIALS**14.1. Efficacy and Safety Studies*****Schizophrenia and Related Disorders Trials***

The efficacy of oral olanzapine in the reduction of and maintenance of the reduction of the manifestations of schizophrenia and related psychotic disorders was established in 3 well-controlled clinical trials of psychotic inpatients who, at entry met the DSM-III-R criteria for schizophrenia (most with a course at entry of 'chronic with acute exacerbation') and 1 well-controlled clinical trial of psychotic inpatients and outpatients who, at entry, met the DSM-III-R criteria for schizophrenia, schizophreniform disorder, or schizoaffective disorder. Study designs for the studies mentioned above and in section 4814.2 can be found in Table 17.

Table 17: Schizophrenia and Related disorder trials characteristics: oral administration

Parameter	1	2	3	4	5
Number of subjects	335	152	431	1996	894
Duration of study	6 weeks	6 weeks	6 weeks	6 weeks	Extension maintenance trials with median treatment length of 237 days
Treatment arms	3 fixed dosage ranges: 5 ± 2.5 mg/day 10 ± 2.5 mg/day 15 ± 2.5 mg/day	2 fixed dose ranges: 1 mg/day 10 mg/day	3 fixed dose ranges: 1 mg/day 5 ± 2.5 mg/day 10 ± 2.5 mg/day 15 ± 2.5 mg/day	1 dosage range: 5 to 20 mg/day	See Studies 1, 3, and 4
Type of control	Haloperidol 15 ± 5 mg/day BID, Placebo	Placebo	Haloperidol 15 ± 5 mg/day BID	Haloperidol 5 to 20 mg/day	Controlled
Further characteristics	-	-	-	2:1 randomization	Double-blinded; extension maintenance trials of studies 1, 3, and 4

Bipolar Disorder Trials

Bipolar Mania:

The efficacy of oral olanzapine in treating acute bipolar mania was demonstrated in 5 controlled studies, including 2 placebo-controlled studies, 2 active comparator studies and 1 cotherapy study. All patients enrolled in these studies had a diagnosis of bipolar I disorder and displayed an acute manic or mixed episode (with or without psychotic features) according to the DSM-IV criteria based on clinical assessment and confirmed by the structured clinical interview for the diagnostic and statistical manual, SCID-P. Study designs for the studies mentioned above and in section 14.2 can be found in Table 18.

Table 18: Bipolar Mania Trials characteristics: oral administration

	Placebo-controlled trials		Active Comparator trials		Cootherapy trial
Parameter	1	2	3	4	5
Number of subjects	Y-MRS: 70 LOCF: 69	Y-MRS: 60 LOCF: 55	Olanzapine arm: 125 Divalproex arm: 126	Olanzapine arm: 234 Haloperidol arm: 219	Olanzapine + Valproate or Lithium: 229 Valproate or Lithium alone: 115
Duration of study	3 weeks	4 weeks	3 weeks (continuation phase 11 months)	6 weeks (6 weeks maintenance of response phase in the absence of a placebo arm)	6 weeks (18 months re- randomized phase)
Treatment arms	Dose range: 5 to 20 mg/day	Dose range: 5 to 20 mg/day	Dose-range Olanzapine: 5 to 20 mg/day Dose-range Divalproex: 500 to 2500 mg/day	Dose-range Olanzapine: 5 to 20 mg/day Dose-range Haloperidol: 3 to 15 mg/day	Cootherapy vs. Monotherapy
Type of control	Placebo	Placebo	Divalproex	Haloperidol	Monotherapy
Further characteristics	-	-	Double-blinded, Randomized	Double-blinded	Double-blinded

14.2 Study Results

Schizophrenia and Related Disorders Trials Results

(1) A 6-week, placebo-controlled trial (N = 335) compared 3 fixed dosage ranges of olanzapine (5 ± 2.5 , 10 ± 2.5 , and 15 ± 2.5 mg/day QD), 1 dosage range of haloperidol (15 ± 5 mg/day on a BID schedule), and placebo. The 2 higher dosage ranges of olanzapine were statistically significantly superior to placebo on the Brief Psychiatric Rating Scale (BPRS) total, the Clinical Global Impressions - Severity of Illness (CGI - S) scale, and the BPRS positive psychosis cluster. The highest dosage range of olanzapine was statistically significantly superior to placebo and to haloperidol on the Scale for the Assessment of Negative Symptoms (SANS). Efficacy of olanzapine generally increased with dose. The 5 ± 2.5 mg/day dosage range of olanzapine was numerically, but not statistically, significantly superior to placebo on BPRS total and other assessments of overall psychopathology.

(2) A 6-week, placebo-controlled trial (N = 152) compared 2 fixed doses of olanzapine (1 or 10 mg/day QD) and placebo. Olanzapine, 10 mg/day, was statistically significantly superior to placebo on the BPRS total, the BPRS positive psychosis cluster, the CGI-S scale, the Positive and Negative Syndrome Scale (PANSS) total, the PANSS positive subscale, and the PANSS negative subscale. Olanzapine, 1 mg/day, appeared to be a no- effect dose with no difference, clinically or statistically, from placebo on any assessment of psychopathology.

(3) A 6-week, dose comparison trial (N = 431) compared 3 fixed dosage ranges of olanzapine (5 ± 2.5 , 10 ± 2.5 and 15 ± 2.5 mg/day QD), olanzapine (1 mg/day QD), and haloperidol (15 ± 5 mg/day on

a BID schedule). There were no statistically significant differences between groups on efficacy measures except for the highest dosage range of olanzapine, which was statistically significantly superior to olanzapine, 1 mg, on the BPRS positive psychosis cluster, PANSS positive subscale, and the CGI-S scale.

(4) A 6-week comparator-controlled trial (N = 1996, 2:1 randomization, olanzapine: haloperidol) compared 1 dosage range of olanzapine (5 to 20 mg/day QD) and 1 dosage range of haloperidol (5 to 20 mg/day QD). The acute mean modal doses (for those patients with at least 3 weeks of treatment) were 13.2 mg/day for olanzapine and 11.8 mg/day for haloperidol. Olanzapine was statistically significantly superior to haloperidol on the BPRS total, the BPRS negative psychosis cluster, the PANSS negative subscale, and the CGI-S scale. Olanzapine was also statistically significantly superior to haloperidol on the Montgomery-Asberg Depression Rating Scale (MADRS). The validity of this scale in patients with schizophrenia, however, is not established.

(5) The effectiveness of olanzapine in long-term therapy, ie, > 6 weeks, was evaluated in 3 double-blind, controlled, extension maintenance trials (of acute trials 1, 3, and 4 above). Patients who showed adequate clinical improvement following double-blind acute therapy were allowed to continue on in a double-blind, long-term extension maintenance phase on their acute dosage regimen. Long-term maintenance of treatment response (as defined by continued reduction in signs and symptoms sufficient to not require hospitalization for psychosis) was compared over time (894 olanzapine-treated patients; median length of treatment was 237 days). The percentage of patients maintaining treatment response over one year was compared. Olanzapine was statistically significantly superior to placebo in the one placebo-controlled trial and was comparable or statistically significantly superior to the active comparator in 3 of 3 active comparator-controlled trials.

Summary of Schizophrenia and Related Disorders Trials

While the efficacy of olanzapine at a dose of 5 mg/day was not statistically superior to placebo (see (1 above)), some individual patients receiving this dose had a good acute response, and were well maintained during a 1-year extension phase.

The above trials (including open-label extension) and an additional trial in geriatric patients with primary degenerative dementia of the Alzheimer's type constitute the primary database (N = 2500 patients treated with olanzapine, corresponding to 1122.2 patient-years; N = 810 patients treated with haloperidol, corresponding to 193.0 patient-years; N = 236 patients treated with placebo, corresponding to 27.1 patient-years).

Bipolar Disorder Trials Results

Bipolar Mania:

Placebo-Controlled Trials: The 2 placebo-controlled trials evaluated the efficacy of olanzapine versus placebo in treating bipolar manic or bipolar mixed episodes as measured by the Y-MRS total score LOCF mean change from baseline to endpoint over 3 weeks (n = 70 and n = 69, respectively) and 4 weeks (n = 60 and n = 55, respectively). These trials demonstrated superiority in the efficacy of olanzapine compared with placebo. The key findings were as follows:

- Olanzapine, at a dose range of 5-20 mg/day, was statistically superior to placebo in improving manic symptoms in each study (p = 0.019 and p < 0.001, respectively).

- In each study, a statistically significantly greater percentage of olanzapine-treated patients (48.6% and 64.8%, respectively) compared with placebo-treated patients (24.2% and 42.9%, respectively) responded to treatment ($\geq 50\%$ reduction in Y-MRS total score) ($p = 0.004$ and $p = 0.023$, respectively).
- In each study, the percentage of patients that were in clinical remission (endpoint Y-MRS total score ≤ 12) were significantly greater among olanzapine patients (45.7% and 61.1%, respectively) compared with placebo patients (25.8% and 35.7%, respectively) ($p = 0.020$ and $p = 0.013$, respectively).
- Olanzapine efficacy did not differ significantly among the main subtypes of bipolar mania, for example patients with a history of rapid cycling, with or without psychotic features, and bipolar mixed or bipolar manic.

Active Comparator Trials: Two active comparator trials were conducted.

(a) The first active comparator study evaluated the efficacy of olanzapine versus divalproex in treating bipolar manic and bipolar mixed episodes by using the Y-MRS total score LOCF mean change from baseline to endpoint. This study was a 3-week, double-blind study with a double-blind continuation phase of 11 months. The primary objective of the study was to demonstrate non-inferiority in the efficacy of olanzapine compared with divalproex at 3 weeks. Patients were randomized to either olanzapine (5-20 mg/day, $n = 125$) or divalproex (500-2500 mg/day, $n = 126$). The key findings were as follows:

- Olanzapine was statistically superior to divalproex in improving manic symptoms as measured by Y-MRS change score at 3 weeks (mean improvements of 13.4 and 10.4 points, respectively, $p = 0.028$).
- The proportion of patients meeting the criteria for response was not statistically significantly different between olanzapine and divalproex groups (54.4% and 42.3%, respectively) ($p = 0.059$).
- The proportion of patients that was in clinical remission was significantly greater among olanzapine patients (47.2%) compared with divalproex patients (34.1%) ($p = 0.039$).

The second active comparator study evaluated the efficacy of olanzapine versus haloperidol in treating bipolar manic or mixed episodes by assessing the proportion of patients in protocol-defined remission from manic and depressive symptoms at 6 weeks. Remission was defined as: 1) achieving improvement in clinical symptomatology in manic and depressive symptoms; 2) having achieved specific reductions in Y-MRS and HAMD-21 total scores; and 3) continuing to take study medication at Week 6. This trial consisted of a 6-week double-blind phase followed by a 6-week double-blind maintenance of response phase in the absence of a placebo arm. Patients were randomly assigned to treatment with olanzapine 5-20 mg/day ($n = 234$) or haloperidol 3-15 mg/day ($n = 219$). The key findings were as follows:

- Olanzapine and haloperidol were similarly effective in improving manic symptoms.
- A clinical response to treatment was defined as a $\geq 50\%$ improvement in Y-MRS total score from baseline to endpoint. In both treatment groups, a large proportion of patients responded to treatment. At the end of the acute phase 72.3% and 74.2% of olanzapine and haloperidol patients, respectively met the response criteria, and at the end of the continuation phase almost all patients were classified as responders (96.3% of 160 olanzapine patients and 94.1% of 136 haloperidol patients).

- The proportion of patients in symptomatic remission at the end of the acute phase (6 weeks) was similar for olanzapine and haloperidol patients (52.1% versus 46.1%, respectively ($p = 0.152$)). Among patients who entered the continuation phase and were not in symptomatic remission at 6-weeks, significantly more olanzapine patients (68.3%) than haloperidol patients (41.0%) were in remission by the end of the continuation period ($p = 0.014$).
- Manic symptoms continued to improve among olanzapine patients to a statistically significant extent.
- Olanzapine was statistically significantly more efficacious than haloperidol in patients without psychotic features (acute phase remission rates were 56.7% in 104 olanzapine patients and 41.6% in 89 haloperidol patients, respectively) ($p = 0.043$).

Cotherapy Trial: This trial evaluated the efficacy of olanzapine plus either valproate or lithium (cotherapy, $n = 229$) versus valproate or lithium alone (monotherapy, $n = 115$) in treating bipolar manic or mixed episodes as measured by the Y-MRS total score LOCF mean change from baseline to endpoint. This study was a 6-week, double-blind study with a re-randomized double-blind phase of 18 months. The key findings were as follows:

- Olanzapine in combination with either valproate or lithium was significantly more efficacious than monotherapy (valproate or lithium) in improving manic symptoms (mean improvements of 13.1 and 9.1 points, respectively) ($p = 0.003$).
- The proportion of patients that clinically responded to treatment was statistically significantly greater among patients receiving olanzapine cotherapy (67.7%) than lithium or valproate monotherapy (44.7%, $p < 0.001$).
- The percentage of patients that were in clinical remission was significantly greater in the olanzapine cotherapy group (78.6%) compared with the lithium or valproate monotherapy group (65.8%, $p = 0.012$).
- The difference in time to remission was also statistically significantly different ($p = 0.002$). The median estimated remission time was 14 days for olanzapine cotherapy-treated patients and 22 days for monotherapy-treated patients.

Bipolar Maintenance:

The efficacy of oral olanzapine as monotherapy for maintenance treatment of bipolar disorder in patients who responded to acute treatment with olanzapine for a manic or mixed episode was demonstrated in two 1-year 'time to event' controlled trials: one placebo-controlled and one active comparator trial against lithium monotherapy.

All patients enrolled in these studies had a diagnosis of bipolar I disorder and displayed an acute manic or mixed episode (with or without psychotic features) according to the DSM-IV criteria.

For both studies: Patients had to meet study-defined response criteria (YMRS total score of ≤ 12 and a HAM-D-21 total score ≤ 8) during open-label treatment with olanzapine (or olanzapine plus lithium in the active comparator study) in order to be randomized into the double-blind maintenance period for observation of study-defined relapse. Dosing was flexible (5 - 20 mg/day for olanzapine; serum levels 0.6 - 1.2 mEq/L for lithium).

The exit criteria was symptomatic relapse of bipolar disorder, either mania or depression. Symptomatic relapse of mania was defined as reaching a YMRS total score ≥ 15 , and symptomatic

relapse of depression as reaching a HAMD-21 total score ≥ 15 ; for the placebo- controlled study only, the definitions also included being hospitalized for mania or depression. Thus, the primary efficacy variable was time to, and incidence of, the exit symptomatic relapse of bipolar disorder, based on analysis of Kaplan-Meier time-to-relapse curves. The key characteristics for the bipolar maintenance trials can be found in Table 19.

Table 19: Bipolar Maintenance Trials characteristics: oral administration

Parameter	1	2
Number of subjects	Olanzapine arm: 225 Placebo arm: 136	Olanzapine + Placebo arm: 217 Lithium + Placebo arm: 214
Duration of study	1 year	1 year
Treatment arms	Dose range: 5 to 20 mg/day	Dose range Olanzapine arm: 5 to 20 mg/day Dose range Lithium arm: 0.6–1.2 mEq/L serum level
Type of control	Placebo	Lithium Monotherapy

1) **Placebo-controlled trial:**

The study evaluated the efficacy of olanzapine vs placebo in maintenance treatment of manic or mixed bipolar episodes by using survival curve analysis to assess the time to, and incidence of, relapse of bipolar disorder. In this trial, 361 patients who had demonstrated response criteria for an average of 16 days were randomized to either continuation of olanzapine at their same dose (n = 225) or to placebo (n = 136), for observation of relapse for up to one year.

The key findings were as follows:

- Figure 1 shows the 1-year time-to-relapse curves for total discontinuations from the study in each arm over time, whether exiting due to relapse, or withdrawal due to adverse events or other reasons. The percentage of patients remaining in the study (i.e., relapse-free, and have not withdrawn for any reason) can be seen at each of the 3, 6, 9 and 12 month points; at study-end, this is 24% (n = 53) for olanzapine and 10% (n = 13) for placebo. The time-point at which 50% of the patients in a specific arm had withdrawn for any reason was Day 59 for the olanzapine group compared to Day 23 for the placebo group.
- Figure 2 shows the 1-year time-to-relapse curves for specifically the exit criterion of bipolar relapse (i.e., patients who withdrew for other reasons were censored and excluded from the calculated numerators and denominators). Olanzapine was superior to placebo, for both incidence of bipolar relapse (46.7% vs 80.1%, respectively), and median time to relapse (174 days vs 22 days, respectively). Note that a high relapse incidence for the placebo arm is not unexpected given the limited time that patients had been demonstrating response criteria prior to randomization.
- Figures 2a and 2b show the efficacy time-to-relapse curves for each of manic and depressive relapse, respectively. Olanzapine showed a statistically significant advantage over placebo in terms of each of mania and depression, although a greater advantage was seen in mania.




Figure 1: Time to Event (Relapse or Discontinuation) Study HGHL; Double-Blind Treatment Phase




Figure 2: Time to Symptomatic Relapse of Bipolar Disorder, Including Hospitalization Study HGHL; Double-Blind Treatment Phase




Figure 2a: Time to Symptomatic Relapse of Mania, Including




Figure 2b: Time to Symptomatic Relapse of Depression, Including Hospitalization Study HGHL; Double-Blind Treatment Phase

2) Active comparator trial:

The study evaluated the efficacy of olanzapine vs lithium in maintenance treatment of manic or mixed bipolar episodes in a non-inferiority design to assess the incidence of relapse of bipolar disorder and further, by using survival curve analysis to assess time to relapse. In this trial, 543 patients who had demonstrated response criteria for an average of 20 days were randomized to either olanzapine plus placebo (n = 217) or lithium plus placebo (n = 214) for observation of relapse for up to one year. The first month of the double blind period was a taper period to allow for non-abrupt lithium discontinuation. The non-inferiority margin used in this study was: $\pm 20\%$ of the efficacy seen for the reference population.

The key findings were as follows:

- Figure 3 shows the 1-year time-to-relapse curves for total discontinuations from the study in each arm over time, whether exiting due to relapse, or withdrawal due to adverse events or other reasons. The percentage of patients remaining in the study (i.e., relapse-free, and have not withdrawn for any reason) can be seen at each of the 3, 6, 9 and 12 month points; at study-end, this is 42% (n = 94) for olanzapine and 28% (n = 61) for lithium. The time-point at which 50% of the patients in a specific arm had withdrawn for any reason was Day 255 for the olanzapine group compared to Day 192 for the lithium group.
- Figure 4 shows the 1-year time-to-relapse curves for specifically the exit criterion of bipolar relapse (i.e., patients who withdrew for other reasons were censored and excluded from the calculations of numerators and denominators). Olanzapine was non-inferior to lithium for both incidence of bipolar relapse (30.0% vs 38.8%, respectively), and time to 25% of patients experiencing relapse (122 days vs 143 days, respectively).

- It can be seen from Figure 4 that for approximately the first five months of the 1-year trial, relapse rate was higher in olanzapine-treated patients; thereafter, the rate of relapse for lithium increases, while that for olanzapine flattens out.
- Figures 4a and 4b are the 1-year time-to-relapse curves for each of the exit criterion of manic and depressive relapse respectively. Olanzapine showed a statistically significant advantage over lithium in rate of mania relapse, and was non-inferior for depressive relapse.




Figure 3: Time to Event (Relapse or Discontinuation) Study HGHT; Double-Blind Treatment Phase




Figure 4: Time to Symptomatic Relapse of Bipolar Disorder Study HGHT; Double-Blind Treatment Phase




Figure 4a: Time to Symptomatic Relapse of Mania Study HGHT; Double-Blind Treatment Phase




Figure 4b: Time to Symptomatic Relapse of Depression Study HGHT; Double-Blind Treatment Phase

Summary of Bipolar Disorder Trials

Bipolar Mania: Olanzapine was more efficacious than placebo and divalproex and as effective as haloperidol in improving overall manic symptomatology in patients with acute bipolar I disorder, manic or mixed episode, with or without psychotic symptoms, and with or without a history of rapid cycling. Olanzapine is associated with a faster onset of action (based on median time to remission estimated with Kaplan-Meier analysis) compared to divalproex and similar to that of haloperidol. The addition of olanzapine also improved patients not responding to lithium or valproate. Olanzapine was not associated with inducing or worsening symptoms of depression.

Bipolar Maintenance: Two 1-year controlled studies support the use of olanzapine monotherapy in maintenance treatment of bipolar patients who responded to acute olanzapine treatment for a manic or mixed episode. Based on analysis of one-year Kaplan-Meier survival curves, olanzapine was superior to placebo, and non-inferior to lithium, in both time to, and incidence of, bipolar relapse over one year.

14.3 Comparative Bioavailability Studies

A randomized, blinded, two-way, single-dose crossover, bioequivalence study comparing OLANZAPINE-ODT orally disintegrating tablets, 5 mg (Jubilant Generics Limited) and ZYPREXA® ZYDIS® orally disintegrating tablets, 5 mg (Eli Lilly Canada Inc.) in healthy adult subjects, under fasting conditions. Comparative bioavailability data from the 26 subjects who completed the study and were included in the statistical analysis are presented in the following table:

Table 20 - SUMMARY TABLE OF THE COMPARATIVE BIOAVAILABILITY DATA

Olanzapine (1 x 5 mg) Geometric Mean Arithmetic Mean (CV %)				
Parameter	Test ¹	Reference ²	% Ratio of geometric means	90% Confidence Interval
AUC _T (ng·hr/mL)	176.25 179.73 (18.91)	173.00 178.49 (23.90)	101.9	93.0 - 111.6
AUC _I (ng·hr/mL)	256.02 263.21 (23.08)	251.52 260.94 (25.72)	101.8	92.7 - 111.8
C _{MAX} (ng/mL)	8.48 8.70 (21.83)	8.60 8.96 (27.67)	98.7	89.7 - 108.5
T _{MAX} ³ (h)	4.90 (29.83)	5.06 (24.43)		
T _½ ³ (h)	29.70 (15.06)	29.68 (17.93)		

¹ OLANZAPINE-ODT (olanzapine) orally disintegrating tablets, 5 mg (Jubilant Generics Limited)

² Zyprexa® Zydys® (olanzapine) orally disintegrating tablets, 5 mg (Eli Lilly Canada Inc.)

³ Expressed as the arithmetic mean (CV %) only.

15. MICROBIOLOGY

No microbiological information is required for this drug product.

16. NON-CLINICAL TOXICOLOGY

An extensive series of acute, subchronic, chronic, reproduction, and genetic toxicity as well as oncogenicity studies have been conducted to support clinical trials with olanzapine. In most of these studies, olanzapine was given by the oral route to rodents, rabbits, and monkeys in an aqueous suspension with 5% to 10% acacia and to dogs as neat material in capsules.

General Toxicology:

The findings of toxicology studies support the safety of olanzapine for oral use in humans as an

antipsychotic agent.

Acute Toxicity:

The acute toxicity of olanzapine was studied in mice, rats, dogs, and monkeys. The estimated median lethal dose for each species is shown below in Table 26.

Table 21: Acute Toxicity Summary

Estimated Median Lethal Dose (mg/kg/day)			
Species	Route	Males	Females
Mouse	Oral	211	208
Rat	Oral	174	177
Dog	Oral	Both sexes > 100 mg/kg	
Monkey	Nasogastric	Both sexes > 100 mg/kg	
Rat	Intraperitoneal	112	107

Signs of toxicity in rodents included hypoactivity, lethargy, leg weakness, coma, tremors, clonic convulsions, salivation, poor grooming, and depressed body weight gain.

Long-term toxicity:

Subchronic administration studies of up to 3 months in duration have been conducted by the oral route in mice, rats, and dogs and chronic toxicity studies of up to 1 year were conducted by the oral route in rats and dogs.

The predominant effects in laboratory animals given olanzapine were CNS depression and anticholinergic effects related to the pharmacology of the drug. Tolerance to the CNS depression developed in repeated-dose studies. Depressed body weight gain was a consistent finding in mice given 30 mg/kg/day and in rats given 4 mg/kg/day. Effects on hematology parameters were found in each species studied in repeated-dose studies. Rats given 16 mg/kg/day had decreased lymphocyte and neutrophil counts and atrophy of bone marrow consistent with the marked reduction in body weight gain. Mice given 3 mg/kg/day developed leukopenia, due primarily to lymphocytopenia, but also associated with neutropenia. Lymphoid necrosis of thymus and spleen was seen in mice given ≥ 10 mg/kg/day. Instances of reversible neutropenia, with or without thrombocytopenia, or anemia developed in a low number of individual dogs treated with 8 or 10 mg/kg/day. Bone marrow from some dogs with olanzapine-induced neutropenia responded to olanzapine with lower than expected numbers of maturing granulocytic cells; however, progenitor and proliferating cells were present in adequate numbers. No olanzapine-related hematologic effects were seen in dogs receiving olanzapine at either 2 or 5 mg/kg/day.

Effects observed in rats consistent with increased plasma concentrations of prolactin in rats included decreased weights of ovaries and uterus. Histopathologic tissue alterations in mammary gland morphology and vaginal epithelium and increased prominence of ovarian follicles were also consistent with elevated prolactin concentrations. Prolactin-induced histopathologic tissue alterations found in rats regressed after treatment cessation. No unexpected toxicologically important findings unrelated to pharmacologic activity were found in the 1-year studies in rats given 4 mg/kg/day or in dogs given 5 mg/kg/day.

Genotoxicity:

Olanzapine was not mutagenic or clastogenic in a full range of standard tests which included bacterial mutation tests and *in vitro* and *in vivo* mammalian tests. Appropriate positive controls were used in each test to verify the sensitivity of the test systems.

Carcinogenicity:

The oncogenic potential of olanzapine was evaluated in studies in rats and mice. Carcinogenicity studies were conducted in CD-1 mice and Fischer 344 rats. Olanzapine was administered orally to mice at doses of 3, 10, or 20 mg/kg for 19 months (males) or 21 months (females) in an initial study, and in a subsequent study at doses of 0.5, 2, or 8 mg/kg for 21 months (males and females). Rats received oral doses of 0.25, 1, 2.5, or 4 mg/kg (males) or 0.25, 1, 2.5, 4, or 8 mg/kg (females) for 24 months. Initial dose levels 2.5 and 4 mg/kg/day in females were increased to 4 and 8 mg/kg/day, respectively, on Day 211. These doses are equivalent to 2 to 70 times the maximum daily human dose (mouse studies) or 0.9 to 28 times the maximum daily human dose (rats). A maximum tolerated dose was achieved in both mouse and rat studies. Increased mortality was seen in mice at doses of 10 and 20 mg/kg and decreases in circulating lymphocytes and neutrophils were seen at doses 0.5 mg/kg. In female mice treated with olanzapine, the incidence of mammary tumours was increased at doses ≥ 2 mg/kg. Female rats treated with 4 or 8 mg/kg had an increase in malignant mammary tumours, but the overall incidence of mammary gland neoplasia was unchanged. An increase in total incidence of mammary gland tumours in female mice given 10 or 20 mg/kg/day (the high dose was decreased from 30 mg/kg/day due to excess mortality) was also noted.

Antipsychotic drugs, including olanzapine, have been shown to chronically elevate prolactin concentrations in rodents. An increase in mammary neoplasms has been found in rodents after chronic administration of other antipsychotic drugs and is considered to be prolactin mediated. The role of prolactin in human breast cancer has not been defined conclusively, and there are presently no epidemiologic data indicating increased risk for breast cancer for humans using antipsychotic drugs.

Reproductive and Developmental toxicology:

Fertility studies in male and female rats and teratology studies in rats and rabbits have been conducted by the oral route. Mating performance was affected by administration of olanzapine due to sedation in male rats given doses greater than 18 times the maximum daily human dose, (5 mg/kg/day in rats), but the effect was quickly reversed when treatment stopped. Estrous cycles were affected and reproduction parameters were influenced in rats given doses greater than 4 times the maximum daily human dose (≥ 1 mg/kg/day in rats). No adverse effects were observed on numbers of corpora lutea, implantations, fetal viability, or fetal weight, and there were no effects on litter size or on the survival, growth, or development of the offspring from parents given up to 18 times the maximum daily human dose (5 mg/kg/day in rats). Although the reproductive process in female rats from mating through fertilization was not adversely affected by treatment, this evidence does not exclude a possible interference with maintenance of pregnancy at high doses of olanzapine. Reproduction studies, performed in rats and rabbits at doses of olanzapine 3.5 and 7 times the maximum daily human dose (20 mg), respectively, have revealed no evidence of harm to the fetus. Maternal toxicity, developmental toxicity (indicated by fetal growth retardation and slightly delayed ossification at birth), and increased numbers of nonviable offspring occurred at higher doses (in rats

at 14 and 63 times the maximum daily human dose and in rabbits at 28 and 105 times the maximum daily human dose). However, fetal malformations were not increased. Transient modest decreases in activity levels of the progeny from females given 0.25 mg/kg/day and skeletal changes indicative of growth retardation in fetuses from females given 5 mg/kg/day were observed. Transient decreases in offspring activity have occurred at all doses; however, there were no effects on body weight, growth, mating, fertility, or live births in second-generation animals. Placental transfer of olanzapine occurs in rat fetuses. Olanzapine was also detected in the milk of rats at concentrations up to three-fold higher than those in the plasma.

Hematologic Indices:

In animal studies with olanzapine, the principal hematologic findings were reversible peripheral cytopenias in individual dogs given high doses of olanzapine (24 to 30 times the maximum daily human dose), dose-related decreases in lymphocytes and neutrophils in mice, and lymphopenia secondary to compromised nutritional status in rats. A few dogs treated with 24 to 30 times the maximum daily human dose developed reversible neutropenia or reversible hemolytic anemia between 1 and 10 months of treatment. Effects on hematology parameters in each species involved circulating blood cells, and no evidence of bone marrow cytotoxicity was found in any of the species examined.

17. SUPPORTING PRODUCT MONOGRAPHS

1. ZYPREXA® (Tablets, 2.5 mg, 5 mg, 7.5 mg, 10 mg and 20 mg), ZYPREXA® ZYDIS® (Orally Disintegrating Tablets, 5 mg, 10 mg, 15 mg and 20 mg), ZYPREXA® IntraMuscular (powder, 10 mg / vial), control 281906, product monograph, CHEPLAPHARM Registration GmbH. (2025-01-30)

PATIENT MEDICATION INFORMATION

READ THIS FOR SAFE AND EFFECTIVE USE OF YOUR MEDICINE

Pr **OLANZAPINE-ODT**

Olanzapine Orally Disintegrating Tablets

Read this carefully before you start taking **OLANZAPINE-ODT** and each time you get a refill. This leaflet is a summary and will not tell you everything about this drug. Talk to your healthcare professional about your medical condition and treatment and ask if there is any new information about **OLANZAPINE-ODT**.

Serious Warnings and Precautions

Higher Death Rate in Elderly Patients with Dementia:

- OLANZAPINE-ODT is not for use in elderly patients with dementia.
- Elderly patients with dementia have a higher death rate when taking OLANZAPINE-ODT and other medicines. Most deaths are caused by serious heart problems, stroke or infections.

Tell your healthcare professional right away if you have sudden weakness or numbness in the face, arms, or legs, and speech or vision problems.

What is OLANZAPINE-ODT used for?

OLANZAPINE-ODT (olanzapine orally disintegrating tablets) is used in adults to treat symptoms of:

- schizophrenia and related psychotic disorders
- bipolar disorder

How does OLANZAPINE-ODT work?

OLANZAPINE-ODT is an antipsychotic medicine. It helps control certain natural chemicals in the brain. This will help control the symptoms of schizophrenia and bipolar mania.

What are the ingredients in OLANZAPINE-ODT?

Medicinal ingredient: olanzapine.

Non-medicinal ingredients: Aspartame, Low substituted hydroxypropyl cellulose, Mannitol, Magnesium stearate, Strawberry flavour, Silica.

OLANZAPINE-ODT comes in the following dosage form:

Orally disintegrating tablets; 5 mg, 10 mg, 15 mg, and 20 mg.

Do not use OLANZAPINE-ODT if:

- you are allergic to olanzapine or any of the ingredients in OLANZAPINE-ODT or the container.

To help avoid side effects and ensure proper use, talk to your healthcare professional before you take OLANZAPINE-ODT. Talk about any health conditions or problems you may have, including if you:

- are pregnant or plan to become pregnant

- are breast feeding or plan on breast feeding
- have had an allergic reaction to any medicine which you have taken previously to treat your current condition
- have diabetes or a family history of diabetes
- have a history of any problems with the way your heart beats or have any heart problems
- have a history of stroke or high blood pressure
- have risk factors for developing blood clots such as: a family history of blood clots, age over 65, smoking, obesity, recent major surgery (such as hip or knee replacement), immobility due to air travel or other reason, or take oral birth control.
- are a smoker
- have ever had blackouts or seizures
- drink alcoholic beverages or use drugs
- exercise vigorously or work in hot or sunny places
- have a history of liver problems, hepatitis, or yellowing of the eyes and skin (jaundice)
- have a history of kidney problems
- have prostate or bladder problems
- have intestinal or bowel problems (paralytic ileus)
- have raised pressure within the eye (glaucoma)
- cannot take phenylalanine because OLANZAPINE-ODT contains aspartame, a source of phenylalanine
- have breast cancer
- have pituitary tumours
- have or have a history of sleep apnea
- are 65 years or older

Other warnings you should know about:

Falls: OLANZAPINE-ODT may cause tiredness, low blood pressure and unstable muscle coordination. This may lead to falls and injuries.

Hyperprolactinemia: OLANZAPINE-ODT may cause higher levels of prolactin. You may be at higher risk of breaking a bone if you have high levels of prolactin and a condition called hypogonadism.

Pregnancy, breast-feeding and newborns:

- If you plan to get pregnant, tell your healthcare professional. Your healthcare professional will decide if you should take OLANZAPINE-ODT.
- If you get pregnant during treatment, tell your healthcare professional immediately. Your healthcare professional will decide if you should continue to take OLANZAPINE-ODT.
- If you take OLANZAPINE-ODT during pregnancy your baby might have the following:
 - difficulty breathing, are overly sleepy, have muscle stiffness, or floppy muscles (like a rag doll), are shaking, or are having difficulty feeding.

Tell your healthcare professional if your newborn baby has these symptoms. Seek emergency hospital care right away if the symptoms are serious. The symptoms may resolve on their own.

Check-ups and monitoring:

You will have regular visits with your healthcare professional, before, during and at the end of your

treatment. They will:

- Check that your heart is working properly.
- blood tests to check your blood sugar levels, blood fat levels, blood and liver health.
- Monitor your blood pressure.
- Check your body weight

Tell your healthcare professional about all the medicines you take, including any drugs, vitamins, minerals, natural supplements or alternative medicines.

The following may interact with OLANZAPINE-ODT:


- Medicines used to treat anxiety or to help you sleep such as lorazepam
- Medicines used to treat depression or obsessive-compulsive disorder like fluvoxamine
- Medicines used to treat high blood pressure such as metoprolol
- Medicines used to treat stomach problems like omeprazole
- Medicines typically used to treat epilepsy (seizures) such as carbamazepine
- Medicines used to treat fungal infections like ketoconazole
- Medicines used to treat bacterial infections like ciprofloxacin
- Medicines used to treat symptoms in Parkinson's disease like pramipexole or levodopa
- Activated charcoal
- Drinking alcohol
- Smoking

How to take OLANZAPINE-ODT:

- Always take OLANZAPINE-ODT exactly as your healthcare professional has told you. Check with your healthcare professional if you are not sure.
- Take OLANZAPINE-ODT for as long as your healthcare professional prescribes it. Do not stop taking this medicine unless your healthcare professional tells you to.
- Take your prescribed dose at the same time each day.

Proper Handling Instructions

- Handle OLANZAPINE-ODT carefully with dry hands.



Follow the instructions below:

1. Separate one blister cell from the strip by tearing along the dotted line.
2. Carefully peel off the backing foil.
3. Gently push the tablet out from the bottom of the blister.
4. Avoid touching the tablet with your hands. Put the tablet directly into your mouth. It will begin to dissolve in your mouth within a few seconds. You can also place the tablet directly into a full glass (125 mL or 4 ounces) of water, milk, coffee, orange juice or apple juice. Stir and drink all of the contents right away.

Usual dose:**OLANZAPINE-ODT (Adults):**

- Your healthcare professional will decide the best dose depending on the disease.
- Your healthcare professional will monitor your health. They may interrupt, reduce, increase or stop your dose. This may occur based on your current health, if you take certain other medications or if you have certain side effects.

Overdose:

If you think you, or a person you are caring for, have taken too much OLANZAPINE-ODT, contact a healthcare professional, hospital emergency department, regional poison control centre or Health Canada's toll-free number, 1-844 POISON-X (1-844-764-7669) immediately, even if there are no signs or symptoms.

Show the healthcare professional your bottle of tablets. Do this even if there are no signs of discomfort or poisoning.

Missed Dose:

- If you miss a dose of OLANZAPINE-ODT by a few hours, take the dose when you remember.
- If most of the day has passed, wait until your next scheduled dose and try not to miss any more.
- **Do not take 2 doses at once.**

What are possible side effects from using OLANZAPINE-ODT?

These are not all the possible side effects you may have when taking OLANZAPINE-ODT. If you experience any side effects not listed here, tell your healthcare professional:

- drowsiness
- weight gain
- dizziness
- anxiety
- increased appetite
- fluid retention
- constipation
- dry mouth
- a feeling of restlessness (akathisia)
- low blood pressure upon rising from a lying or sitting position
- back pain
- stuttering (disruptive speech)
- increased salivation (salivary hypersecretion)
- memory loss
- breast enlargement in men
- milk secretion, changes in period in women

Serious side effects and what to do about them			
Symptom / effect	Talk to your healthcare professional		Stop taking drug and get immediate medical help
	Only if severe	In all cases	
COMMON			
Constipation (new or worsening)		✓	
Hypotension (low blood pressure): dizziness, fainting, light-headedness, blurred vision, nausea, vomiting, fatigue (may occur when you go from lying or sitting to standing up)		✓	
Tachycardia (abnormally fast heartbeat): dizziness, light headedness, shortness of breath, racing heart		✓	
UNCOMMON			
Bradycardia (slow heartbeat)		✓	
RARE			
Hyperglycaemia, Glycosuria (high sugar in blood or urine): Extreme thirst, increased urination, dry mouth, tiredness, visual disturbances, nausea and vomiting, abdominal pain, increased hunger		✓	
Liver problems: fever, yellow skin or eyes, dark urine, weakness, abdominal pain, nausea, vomiting, loss of appetite, itching, light coloured stool, trouble thinking clearly			✓
Low white blood cell count: infection, such as cold, flu-like symptoms, fever, sore throat, as well as weakness or general feeling of unwellness		✓	
Seizures: loss of consciousness with uncontrollable shaking ("fit")			✓
VERY RARE			
Allergic reactions: skin rash, hives, swelling, difficulty breathing			✓
Blood clots: swelling, pain and redness in an arm or leg that can be warm to touch, sudden chest pain, difficulty breathing and heart palpitations		✓	
Neuroleptic Malignant Syndrome (NMS, a nervous system disorder): High fever, muscle rigidity, rapid heartbeat, profuse sweating, irregular pulse			✓
Pancreatitis (pancreas inflammation): severe abdominal pain, fever, nausea, vomiting			✓
Priapism: long-lasting (greater than 4 hours in duration) and painful erection of the penis			✓

Serious skin reactions: (Drug Reaction with Eosinophilia and Systemic Symptoms [DRESS]): skin rash or redness developing into widespread rash with blisters and peeling skin, swollen lymph nodes and fever.			✓
Sudden cardiac death (SCD, heart failure): chest pain, shortness of breath, fainting, fast heartbeats			✓
Cerebrovascular adverse events (disturbance of blood flow to the brain), including stroke : Sudden weakness or numbness in the face, arms, or legs, and speech or vision problems			✓
Tardive dyskinesia : Muscle twitching or abnormal movements of the face or tongue			✓
Thrombocytopenia (low blood platelets): bruise easily, excessive bleeding		✓	
Rhabdomyolysis (skeletal muscle break down): Very dark (“tea coloured”) urine, muscle tenderness and/or aching			✓
UNKNOWN			
Hyponatremia (low level of sodium in the blood): lethargy, confusion, muscular twitching, achy, stiff or uncoordinated muscles, seizures, coma		✓	
SIADH—syndrome of inappropriate antidiuretic hormone secretion : concentrated urine (dark in colour), feel or are sick, have muscle cramps, confusion and fits (seizures) which may be due to inappropriate secretion of ADH (antidiuretic hormone).		✓	

If you have a troublesome symptom or side effect that is not listed here or becomes bad enough to interfere with your daily activities, tell your healthcare professional.

Reporting side effects

You can report any suspected side effects associated with the use of health products to Health Canada by:

- Visiting the Web page on Adverse Reaction Reporting (canada.ca/drug-device-reporting) for information on how to report online, by mail or by fax; or
- Calling toll-free at 1-866-234-2345.

NOTE: Contact your healthcare professional if you need information about how to manage your side effects. The Canada Vigilance Program does not provide medical advice.

Storage:

- Store OLANZAPINE-ODT at 15°C-30°C, in original package, in a dry place. Protect from light.

- Keep out of reach and sight of children.

If you want more information about OLANZAPINE-ODT:

- Talk to your healthcare professional
- Find the full product monograph that is prepared for healthcare professionals and includes the Patient Medication Information by visiting the Health Canada Drug Product Database website: (<https://www.canada.ca/en/health-canada/services/drugs-health-products/drug-products/drug-product-database.html>); or by contacting the distributor, Mint Pharmaceuticals Inc., at 1-877-398-9696.

This leaflet was prepared by Jubilant Generics Limited.

1A- Sector 16A, Institutional Area,
Noida, Uttar Pradesh, India, 201301.

Importer/Distributor:

Mint Pharmaceuticals Inc.

1093 Meyerside Drive

Mississauga, Ontario

L5T 1J6

Last Revised: JAN 27, 2026