

**Product Monograph
Including Patient Medication Information**

PrSAIZEN®

Somatropin for Injection

Produced in genetic engineered mouse C127 cells by recombinant DNA technology

Lyophilized Powder for reconstitution

5 mg/vial

Somatropin

Produced in genetic engineered mouse C127 cells by recombinant DNA technology

Solution for Injection in a Cartridge

6 mg (5.83 mg/mL)

12 mg (8 mg/mL)

20 mg (8 mg/mL)

Human Growth Hormone

EMD Serono, A Division of EMD Inc., Canada

2695 North Sheridan Way, Suite 200

Mississauga, Ontario, L5K 2N6

EMD Serono is the Canadian healthcare business of Merck KGaA,

Darmstadt, Germany

www.emdserono.ca

Date of Authorization:

2026-08-28

Control Number: 304422

® Registered trademark of Merck KGaA, Darmstadt, Germany or its affiliates

Recent Major Label Changes

[7 Warnings and Precautions, Musculoskeletal](#)

2026-06

Table of Contents

Certain sections or subsections that are not applicable at the time of the preparation of the most recent authorized product monograph are not listed.

Recent Major Label Changes	<u>2</u>
Table of Contents	<u>2</u>
Part 1: Healthcare Professional Information	<u>5</u>
1. Indications	<u>5</u>
1.1. Pediatrics	<u>5</u>
1.2. Geriatrics	<u>5</u>
2. Contraindications	<u>5</u>
3. Serious Warnings and Precautions Box	<u>6</u>
4. Dosage and Administration	<u>7</u>
4.1. Dosing Considerations	<u>7</u>
4.2. Recommended Dose and Dosage Adjustment	<u>7</u>
4.3. Reconstitution	<u>8</u>
4.4. Administration	<u>8</u>
4.5. Missed Dose	<u>10</u>
5. Overdose	<u>10</u>
6. Dosage Forms, Strengths, Composition, and Packaging	<u>10</u>
7. Warnings and Precautions	<u>11</u>
General	<u>11</u>
Carcinogenesis and Genotoxicity	<u>12</u>
Congenital Disorders	<u>12</u>
Dependence, Tolerance and/or Abuse Liability	<u>13</u>
Driving and Operating Machinery	<u>13</u>
Endocrine and Metabolism	<u>13</u>
Hematologic	<u>14</u>
Hepatic/Biliary/Pancreatic	<u>14</u>

Immune	<u>14</u>
Monitoring and Laboratory Tests	<u>15</u>
Musculoskeletal	<u>16</u>
Neurologic.....	<u>17</u>
Reproductive Health	<u>17</u>
7.1. ____Special Populations.....	<u>18</u>
7.1.1. ____Pregnancy	<u>18</u>
7.1.2. ____Breastfeeding.....	<u>18</u>
7.1.3. ____Pediatrics	<u>18</u>
7.1.4. ____Geriatrics.....	<u>19</u>
8. ____Adverse Reactions	<u>19</u>
8.1. ____Adverse Reaction Overview	<u>19</u>
8.2. ____Clinical Trial Adverse Reactions	<u>20</u>
8.2.1. ____Clinical Trial Adverse Reactions – Pediatrics	<u>23</u>
8.3. ____Less Common Clinical Trial Adverse Reactions.....	<u>40</u>
8.3.1. ____Less Common Clinical Trial Adverse Reactions – Pediatrics.....	<u>41</u>
8.5. ____Post-Market Adverse Reactions.....	<u>44</u>
9. ____Drug Interactions.....	<u>44</u>
9.2. ____Drug Interactions Overview	<u>44</u>
9.3. ____Drug-Behaviour Interactions	<u>44</u>
9.4. ____Drug-Drug Interactions	<u>44</u>
9.5. ____Drug-Food Interactions	<u>46</u>
9.6. ____Drug-Herb Interactions	<u>46</u>
9.7. ____Drug-Laboratory Test Interactions	<u>46</u>
10. ____Clinical Pharmacology	<u>46</u>
10.1. ____Mechanism of Action	<u>46</u>
10.2. ____Pharmacodynamics	<u>47</u>
10.3. ____Pharmacokinetics.....	<u>51</u>
10.4. ____Immunogenicity	<u>53</u>
11. ____Storage, Stability, and Disposal.....	<u>54</u>
Part 2: Scientific Information	<u>55</u>

13._Pharmaceutical Information	<u>55</u>
14._Clinical Trials.....	<u>55</u>
14.1. __Clinical Trials by Indication.....	56
Pediatric Indications	56
Adult Indications.....	65
16._Non-Clinical Toxicology	<u>68</u>
Patient Medication Information.....	<u>71</u>
Patient Medication Information.....	<u>80</u>

Part 1: Healthcare Professional Information

1. Indications

SAIZEN (somatropin) is indicated for:

Pediatric Indications

Growth Hormone Insufficiency or Deficiency:

SAIZEN is indicated for the long-term treatment of children with growth failure due to inadequate secretion of normal endogenous growth hormone. Other causes for growth failure should be ruled out.

Turner Syndrome:

SAIZEN is indicated for the treatment of short stature in girls with gonadal dysgenesis (Turner syndrome) when epiphyses are not closed.

Chronic Renal Failure:

SAIZEN is indicated for the treatment of growth failure in children due to Chronic Renal Failure.

Small for Gestational Age (SGA):

SAIZEN is indicated for growth disturbance (current height Standard Deviation Score (SDS) <-2) in short children born small for gestational age (SGA) with a birth weight and/or length below -2 standard deviations (SD), who failed to show catch-up growth (Height Velocity SDS < 0 during the last year) by 2 years of age or later.

Adult Indication

Adult Growth Hormone Deficiency:

SAIZEN is indicated for the replacement therapy in adult patients with acquired or idiopathic growth hormone deficiency (GHD) as diagnosed by a single dynamic test for growth hormone deficiency (peak GH ≤ 5 $\mu\text{g/L}$). Patients with a growth hormone deficiency with onset in childhood should be retested before treatment starts.

1.1. Pediatrics

Pediatrics (<18 years of age): Based on the data submitted and reviewed by Health Canada, the safety and efficacy of SAIZEN in pediatric patients has been established. Therefore, Health Canada has authorized an indication for pediatric use (see [1 Indications](#)).

1.2. Geriatrics

Geriatrics (≥ 65 years of age): No data are available to Health Canada; therefore, Health Canada has not authorized an indication for geriatric use.

2. Contraindications

SAIZEN is contraindicated and should not be administered in the following cases:

- Acute critical illness with complications following cardiac surgery, abdominal surgery, multiple accidental trauma or acute respiratory failure. Clinical studies demonstrated that high doses of

somatropin were associated with a significantly increased morbidity and mortality in those patients (see [7 Warnings and Precautions](#)).

- In patients with closed epiphyses, SAIZEN has no effect on cartilaginous growth areas of the long bone. Treatment of pediatric growth disorders with SAIZEN should be discontinued when the patient has reached satisfactory adult height, or the epiphyses are fused.
- In the presence of progression of an underlying intracranial tumour. An intracranial tumour should be inactive, with evidence of remission prior to instituting therapy, and SAIZEN should be discontinued if there is evidence of recurrent activity. Patients should be examined frequently for progression or recurrence of the underlying disease process.
- Patients known to be hypersensitive to somatropin and to any of the excipients in the two formulations: lyophilized powder for reconstitution (and solvent) and solution for injection in a cartridge.
- Active neoplasia (either newly diagnosed or recurrent). Any pre-existing neoplasia should be inactive. Somatropin should be discontinued if there is evidence of recurrent tumor growth
- Proliferative or preproliferative diabetic retinopathy.

Somatropin is contraindicated in patients with Prader-Willi syndrome who are severely obese, have a history of upper airway obstruction or sleep apnea, or have severe respiratory impairment. There have been reports of sudden death when somatropin was used in such patients. SAIZEN is not indicated for the treatment of pediatric patients who have growth failure due to genetically confirmed Prader-Willi syndrome (see [3 Serious Warnings and Precautions Box](#)).

SAIZEN treatment should be discontinued in critically ill patients.

SAIZEN is not recommended for use during pregnancy and lactation.

In children with chronic renal disease, treatment with somatropin must be discontinued at the time of renal transplantation.

SAIZEN reconstituted with bacteriostatic diluent should not be administered to neonates (infants 28 days old or younger), or to patients sensitive to benzyl alcohol, found in the diluent.

3. Serious Warnings and Precautions Box

- SAIZEN treatment should be carried out under regular guidance of a physician experienced in the diagnosis and management of growth disorders: Growth Hormone Insufficiency or Deficiency, Turner syndrome, Chronic Renal Failure, Small for Gestational Age or adult patients with either childhood-onset or adult-onset growth hormone deficiency
- SAIZEN shall only be used if, once reconstituted, the resulting solution is water-clear and devoid of particulate matter (See [Patient Medication Information](#), *How to take SAIZEN*)
- Benzyl alcohol, used as a preservative in Bacteriostatic Water for Injection, USP has been associated with toxicity in newborns*. When administering SAIZEN to newborns, reconstitute with sterile water for injection, USP. Only use one reconstituted dose per growth hormone vial and discard the unused portion.
- There have been reports of fatalities associated with the use of growth hormone in pediatric

patients with Prader-Willi syndrome who have one or more of the following risk factors: severe obesity, history of respiratory impairment or sleep apnea or unidentified [i.e. previously undiagnosed/mildly symptomatic] respiratory infections (see [2 Contraindications](#); [7 Warnings and Precautions, Congenital Disorders](#)).

* A newborn infant, or neonate, is a child under 28 days of age.

4. Dosage and Administration

4.1. Dosing Considerations

- Before initiating a patient on SAIZEN therapy, please review completely the [2 Contraindications](#) and [7 Warnings and Precautions](#) sections.
- SAIZEN dosage should be individualized for each patient according to body weight.
- SAIZEN treatment should be carried out under regular guidance of a physician experienced in the diagnosis and management of growth disorders.
- The patient's medical history for hypersensitivity reactions should be carefully evaluated prior to performing the injection (see [7 Warnings and Precautions, Immune](#)).
- For SAIZEN 5 mg, once the appropriate dose for a patient has been determined, reconstitute each vial of SAIZEN with the diluent supplied. Do not use in patients sensitive to benzyl alcohol, see [2 Contraindications](#).
- Oral estrogens may reduce the serum IGF-I response to SAIZEN treatment. Female patients receiving oral estrogen replacement may require higher SAIZEN dosages. However, the maximum recommended weekly dose should not be exceeded (see [9.4 Drug-Drug Interactions](#)).

4.2. Recommended Dose and Dosage Adjustment

Growth failure due to inadequate endogenous growth hormone secretion:

It is recommended that SAIZEN be administered subcutaneously at a dose of 0.2 mg/kg body weight per week. The dosage can be increased to 0.27 mg/kg per week if there is insufficient response to treatment.

Growth failure in girls due to gonadal dysgenesis (Turner syndrome):

It is recommended that SAIZEN be administered subcutaneously at a dose of 0.375 mg/kg body weight per week (optimal dosing 0.32 – 0.375 mg/kg/week).

Concomitant therapy with non-androgenic anabolic steroids in patients with Turner syndrome can enhance the growth response.

Growth failure in children with Chronic Renal Failure:

It is recommended that SAIZEN be administered subcutaneously at a dose of 0.35 mg/kg body weight per week.

Growth disturbance in short children born small for gestational age (SGA):

It is recommended that SAIZEN be administered subcutaneously at a dose of 0.47 mg/kg body weight/week.

Adult Growth Hormone Deficiency:

It is recommended that SAIZEN be administered subcutaneously at a dose of 0.005 mg/kg/day at the start of therapy. This dose may be increased after 4 weeks to 0.01 mg/kg/day if well tolerated. The minimum effective dose should be used and dose requirements may decline with age.

4.3. Reconstitution

SAIZEN (somatropin for injection) 5 mg Vial:

The recommended diluents for reconstitution are:

Water for Injection, USP and Bacteriostatic Water for Injection, USP.

When administering SAIZEN to newborns, reconstitute with sterile water for injection, USP. Only use one reconstituted dose per growth hormone vial and discard the unused portion (see [2 Contraindications](#)).

Incompatibility: SAIZEN should not be mixed with other drugs.

Preparation of Solution

To prevent possible contamination of the vial, wipe the rubber stopper with an antiseptic solution before puncturing it with the needle.

After determining appropriate patient dose, reconstitute each 5 mg vial of SAIZEN with 1 to 3 mL of Bacteriostatic Water for Injection, USP (Benzyl Alcohol preserved).

To reconstitute SAIZEN, inject the diluent into the vial of SAIZEN aiming the liquid against the glass vial wall. Swirl the vial with a GENTLE rotary motion until contents are dissolved completely. DO NOT SHAKE. If shaken, the solution will appear opalescent; however, this opalescence does not indicate any decrease in potency. Parenteral drug products should be inspected visually prior to administration. Do not inject if the reconstituted product contains particulate matter or is discoloured. For use in patients sensitive to the diluent see [7 Warnings and Precautions](#).

Table 1 – Reconstitution

Vial Size	Volume of Diluent To Be Added to Vial	Approximate Available Volume	Concentration Per mL
Nominal capacity: 2 mL*	1 to 3 mL	1 to 3 mL	1.67 to 5.0 mg/mL depending on volume of diluent used.

* According to ISO 8362-1, brimful capacity: 4 mL ± 0.5 mL

4.4. Administration

SAIZEN solution should not be administered if it contains particles or is not clear.

SAIZEN 5 mg Vial:

SAIZEN should be administered using sterile, disposable syringes and needles. The syringe used should be of appropriately small volume to ensure the accurate dose withdrawal. The calculated dose should be withdrawn for either subcutaneous or intramuscular injection.

SAIZEN 6 mg (5.83 mg/mL), 12 mg (8 mg/mL), 20 mg (8 mg/mL) Cartridges:

The cartridge containing the solution of SAIZEN is for use with the easypod® electromechanical auto-injector or the Aluetta® pen injector.

The Aluetta pen injectors and SAIZEN cartridges are available in several presentations. Each Aluetta pen injector is colour coded and must only be used with the matching colour coded SAIZEN cartridge to give the correct dose:

- The Aluetta pen injector 6 (blue) must be used with the SAIZEN 6 mg (5.83 mg/mL) cartridge (blue).
- The Aluetta pen injector 12 (red) must be used with the SAIZEN 12 mg (8 mg/mL) cartridge (red).
- The Aluetta pen injector 20 (yellow) must be used with the SAIZEN 20 mg (8 mg/mL) cartridge (yellow).

The route of injection is subcutaneous.

Pediatric Indications

Growth failure due to inadequate endogenous growth hormone secretion:

a) Subcutaneous injection:

The weekly dose can be divided into 3 single doses (corresponding to 0.067 mg/kg per injection) or into 6 or 7 single daily doses (corresponding to 0.033 or 0.028 mg/kg per injection, respectively). The injection site should be altered to prevent lipoatrophy. For subcutaneous injections, the use of a needle which is 1.25 cm long is recommended.

b) Intramuscular injection:

The weekly dose should be divided into 3 single injections (corresponding to 0.067 mg/kg). For intramuscular injections, the use of a needle which is at least 2.5 cm long is recommended to ensure the injection reaches the intramuscular layer.

Growth failure in girls due to gonadal dysgenesis (Turner syndrome):

The weekly dose can be divided into 3 single doses (corresponding to 0.137 - 0.161 mg/kg per injection) or into 7 single daily subcutaneous doses (corresponding to 0.045 - 0.054 mg/kg per injection).

Growth failure in children with Chronic Renal Failure:

The daily subcutaneous injection consists of a single injection of 0.05 mg/kg body weight. The injection site should be altered to prevent lipoatrophy. A needle 1.25 cm long should be used for subcutaneous injections.

Growth disturbance in short children born small for gestational age (SGA):

For SGA patients, SAIZEN should be administered as a daily subcutaneous injection consisting of a single injection of 0.067 mg/kg body weight. The injection site should be altered to prevent lipoatrophy. A needle 1.25 cm long should be used for subcutaneous injections.

Adult Indication

Adult Growth Hormone Deficiency

At the start of somatropin therapy, a low dose of 0.005 mg/kg/day is recommended, given as a daily subcutaneous injection. The dose should be adjusted stepwise, controlled by IGF-1 age-adjusted normal values, to 0.01 mg/kg/day if well tolerated. The recommended final GH dose seldom exceeds 1.0 mg/day. In general, the lowest efficacious dose should be administered. In older or overweight patients, lower doses may be necessary.

4.5. Missed Dose

For patients who miss a dose, it is not recommended to double the next dose. Administer the regular dose at the next scheduled dosage time.

5. Overdose

No cases of acute overdosage have been reported. However, exceeding the recommended doses can cause side effects. Overdosage can lead to hypoglycemia followed by hyperglycemia. Long-term overdosage could result in signs and symptoms of gigantism and/or acromegaly consistent with the known effects of excess human growth hormone. If any signs of overdosage occur, treatment should be discontinued. Moreover, somatropin overdose is likely to cause manifestations of fluid retention.

For the most recent information in the management of a suspected drug overdose, contact your regional poison control centre or Health Canada's toll-free number, 1-844 POISON-X (1-844-764-7669).

6. Dosage Forms, Strengths, Composition, and Packaging

To help ensure the traceability of biologic products, healthcare professionals should record both the brand name and the non-proprietary (active ingredient) name as well as other product-specific identifiers such as the Drug Identification Number (DIN) and the batch/lot number of the product supplied.

Table 2 – Dosage Forms, Strengths, and Composition

Route of Administration	Dosage Form/ Strength/Composition	Non-Medicinal Ingredients
Subcutaneous injection, Intramuscular injection	Lyophilized powder for reconstitution/5 mg/vial	Phosphoric acid, sodium hydroxide, sucrose.
Subcutaneous injection	Solution for Injection in a Cartridge/ 6 mg (5.83 mg/mL), 12 mg (8 mg/mL), 20 mg	Citric acid, phenol, Poloxamer 188, sucrose.

	(8 mg/mL)	
--	-----------	--

Description

SAIZEN 5 mg/vial:

SAIZEN is a sterile, non-pyrogenic, lyophilized powder.

Each vial contains 5 mg somatropin, 1.2 mg phosphoric acid, 0.7 mg of sodium hydroxide and 34.2 mg sucrose.

SAIZEN 6 mg (5.83 mg/mL), 12 mg (8 mg/mL), 20 mg (8 mg/mL) cartridges:

6 mg (5.83 mg/mL) Cartridge: Each cartridge contains 6 mg of somatropin, 77.3 mg sucrose, 2.1 mg Poloxamer 188, 3.8 mg phenol and citric acid (pH 6.1 ± 0.1).

12 mg (8 mg/mL) Cartridge: Each cartridge contains 12 mg somatropin, 112.5 mg sucrose, 3.0 mg Poloxamer-188, 5.6 mg phenol and citric acid (pH 6.1 ± 0.1).

20 mg (8 mg/mL) Cartridge: Each cartridge contains 20 mg somatropin, 187.5 mg sucrose, 5.0 mg Poloxamer 188, 9.3 mg phenol and citric acid (pH 6.1 ± 0.1).

Packaging

SAIZEN 5 mg/vial:

SAIZEN is available as a sterile, non-pyrogenic, lyophilized powder.

Each carton contains 1 vial of 5 mg somatropin for injection together with 1 vial of diluent (3 mL bacteriostatic Water for Injection, USP).

The recommended route of administration is subcutaneous or intramuscular.

SAIZEN 6 mg (5.83 mg/mL), 12 mg (8 mg/mL) and 20 mg (8 mg/mL) Cartridges:

SAIZEN, solution for injection in a cartridge is supplied in a standard 3 mL nominal capacity glass cartridge (Type I).

They are available in a single (1) cartridge pack size.

The SAIZEN cartridges are colour coded: SAIZEN 6 mg (5.83 mg/mL) in blue, 12 mg (8 mg/mL) in red, and 20 mg (8 mg/mL) in yellow.

The recommended route of administration is subcutaneous.

7. Warnings and Precautions

General

Fluid retention is expected during growth hormone replacement therapy in adults. Growth hormone increases sodium retention with expansion of the extracellular volume and this anti-natriuretic effect appears to be mediated through the rennin-angiotensin-aldosterone system. Adverse events such as peripheral edema, joint swelling, myalgia, arthralgia, paresthesia, carpal tunnel syndrome and benign intracranial hypertension may be clinical manifestation of fluid retention and appear to be more frequent in elderly patients with adult-onset disease (see [8 Adverse Reactions](#)). However, these

symptoms/signs are usually transient and dose dependent therefore reduce dose as necessary.

Concomitant glucocorticoid therapy may inhibit the response to SAIZEN and should not exceed 10-15 mg hydrocortisone equivalent/m² body surface area during SAIZEN treatment (see [9.4 Drug-Drug Interactions](#)).

To avoid transmission of disease, cartridge and prefilled syringe shall not be used by more than one person. Instructions on appropriate use should be given, (see [Patient Medication Information](#)).

Information for patients

Patients and/or their parents should be informed about potential advantages and disadvantages of growth hormone therapy including the possible side effects. If home use is determined to be desirable by the physician, patients should also be offered instruction for use of injection devices, storage, travelling and other pertinent information. (see [Patient Medication Information](#)).

Female patients should be advised to inform their doctor if they are pregnant or are contemplating pregnancy. Careful monitoring, as well as general health is essential in pregnant patients. (see [7.1 Special Populations](#) and [Patient Medication Information](#)).

Carcinogenesis and Genotoxicity

Carcinogenicity studies have not been conducted. Mutagenicity studies showed no mutagenic activity with SAIZEN.

Leukemia has been reported in a small number of growth hormone deficient patients, treated with growth hormone. Based on the current evidence, experts cannot conclude that growth hormone therapy is responsible for these occurrences.

Treatment with growth hormone may have an increased risk of developing neoplasm.

Secondary Neoplasm in Survivors of Childhood Cancer:

In childhood cancer survivors, an increased risk of a second neoplasm (benign and malignant) has been reported in patients treated with growth hormones. Intracranial tumours, in particular meningiomas in patients treated with radiation to the head for their first neoplasm, were the most common of these second neoplasms.

Congenital Disorders

Prader-Willi Syndrome:

SAIZEN is not indicated for the long-term treatment of pediatric patients who have growth failure due to genetically confirmed Prader-Willi Syndrome, unless they also have a diagnosis of growth hormone deficiency. There have been reports of fatalities after initiating therapy with growth hormone in pediatric patients with Prader-Willi syndrome who had one or more of the following risk factors: severe obesity, history of upper airway obstruction or sleep apnea, or unidentified (i.e. previously undiagnosed/mildly symptomatic) respiratory infection (see [3 Serious Warnings and Precautions Box](#)).

Male patients with one or more of these factors may be at greater risk than females. Patients with Prader-Willi syndrome should be evaluated for signs of upper airway obstruction and sleep apnea before initiation of treatment with growth hormone. If during treatment with growth hormone, patients show signs of upper airway obstruction (including onset of or increased snoring) and/or new onset of sleep apnea, treatment should also be interrupted and the patients should be treated as indicated. All patients with Prader-Willi syndrome treated growth hormone should also have effective weight control and be monitored for signs of respiratory infection, which should be diagnosed as early as possible and treated aggressively. Unless patients with Prader-Willi syndrome also have a diagnosis of growth hormone deficiency, SAIZEN is not indicated for the long-term treatment of pediatric patients who have growth failure due to genetically-confirmed Prader-Willi syndrome.

Turner Syndrome:

Patients with Turner syndrome may be at increased risk for development of intracranial hypertension therefore these patients should be evaluated for signs and symptoms of intracranial hypertension and treated aggressively before initiation of treatment with somatropin.

Patients with Turner syndrome should be evaluated carefully for otitis media and other ear disorders before and during treatment with growth hormones since these patients have an increased risk of ear or hearing disorders. In the presence of ear infection or hearing disorders, these patients should be treated as indicated.

Patients with Turner syndrome are at risk for cardiovascular disorders (e.g. stroke, aortic aneurysm, hypertension) and these conditions should be monitored closely before and during treatment with somatropin.

Patients with Turner syndrome have an inherently increased risk of developing autoimmune thyroid disease. Therefore, patients should have periodic thyroid function tests and be treated as indicated. Skeletal abnormalities including scoliosis are commonly seen in untreated Turner syndrome patients.

Dependence, Tolerance and/or Abuse Liability

Inappropriate use of somatropin by individuals who do not have indications for which growth hormone is approved, may result in clinically significant negative health consequences.

Driving and Operating Machinery

SAIZEN does not interfere with the patient's ability to drive or use machinery.

Endocrine and Metabolism

Because human growth hormone may induce a state of insulin-resistance, SAIZEN patients should be monitored for evidence of glucose intolerance. SAIZEN should be used with caution in patients with diabetes mellitus (adjustment of their antidiabetic therapy may be required) or a family history of diabetes mellitus (see [Monitoring and Laboratory Tests](#)). SGA patients are a subgroup of patients at higher risk of developing diabetes in whom fasting insulin and blood glucose should be closely monitored before initiating and during treatment with SAIZEN. SAIZEN administration is followed by a transient phase of hypoglycemia of approximately 2 hours, then from 2-4 hours onward by an increase

in blood glucose levels despite high insulin concentrations. To detect insulin resistance, patients should be monitored for evidence of glucose intolerance.

In adults with risk factors for insulin resistance or glucose intolerance, such as obesity, a family history of diabetes mellitus and those on high dose corticosteroid therapy, growth hormone therapy may induce Type II Diabetes Mellitus if the insulin secretory capacity is impaired.

Growth hormone can affect the metabolism of thyroid hormones by increasing the extrathyroidal conversion of T4 to T3 and this lowering effect on T4 may unmask incipient central hypothyroidism in hypopituitary patients. Therefore, thyroid function should be evaluated before starting SAIZEN therapy and regularly assessed during treatment, not less frequently than annually. If hypothyroidism is diagnosed in the course of SAIZEN therapy, it should be corrected because untreated hypothyroidism will jeopardize the response to growth hormone.

In patients with hypopituitarism (multiple hormonal deficiencies) standard hormonal replacement therapy should be monitored closely when human growth hormone therapy is administered (see [Monitoring and Laboratory Tests](#)).

Hematologic

Serum levels of inorganic phosphorus, alkaline phosphatase, and Insulin-like Growth Factor 1 (IGF-1) may increase with SAIZEN therapy.

Hepatic/Biliary/Pancreatic

SAIZEN is indicated for the treatment of growth failure in children due to Chronic Renal Failure. The safety of SAIZEN has not been established in patients with hepatic, biliary or pancreatic impairments. Growth hormone requirements may need to be adjusted in patients with renal and/or hepatic and/or biliary and/or pancreatic impairments.

Pancreatitis

Cases of pancreatitis have been reported rarely in children and adults receiving somatropin treatment. Children treated with growth hormone may have an increased risk of developing pancreatitis compared to adults. Published literature indicates that girls who have Turner syndrome may be at a greater risk than other somatropin-treated children. Pancreatitis should be considered in growth hormone-treated patients, especially children, who develop abdominal pain.

Immune

Local allergic reactions:

With growth hormone therapies, patients may experience redness, swelling, pain, inflammation, or itching at the site of injection. (see [8 Adverse Reactions](#)). Most of these minor reactions usually resolve in a few days to a few weeks. They may occur if the injection is not properly made (irritants in the skin cleansing agent or poor injection technique), or if the patient is allergic to the growth hormone or any excipients. (see [2 Contraindications](#)).

Injection sites should be varied to prevent localized lipoatrophy, in particular in the case of long-term, subcutaneous administrations of SAIZEN. On rare occasion, injection site reactions may require discontinuation of SAIZEN therapy.

Systemic allergic reactions:

There is a potential risk with somatropin that severe cases of generalized allergy including anaphylactic reaction may be life threatening.

Antibody Production:

As with all protein pharmaceuticals, a small percentage of patients may develop antibodies during treatment with growth hormones. The clinical significance of these antibodies is unknown, though to date the antibodies have been of low binding capacity and have not been associated with growth deceleration except in patients with gene deletions. If growth deceleration is observed that is not attributable to another cause, testing for antibodies to somatropin should be considered for any patient who fails to respond to therapy.

Monitoring and Laboratory Tests

With SAIZEN the need for regular IGF-1 monitoring shall be considered to maintain IGF-1 within the normal range for age and sex (see [Patient Medication Information](#)). Because human growth hormone may induce a state of insulin resistance, patients should be observed for evidence of glucose intolerance. Patients with diabetes or glucose intolerance should be monitored closely during therapy with human growth hormone.

Patients with Turner syndrome should be evaluated carefully for otitis media and other ear disorders since these patients have an increased risk of ear or hearing disorders.

Patients with Turner syndrome are at risk for cardiovascular disorders (e.g. stroke, aortic aneurysm, and hypertension) and these conditions should be monitored closely.

Patients with Turner syndrome have an inherently increased risk of developing autoimmune thyroid disease. Therefore, patients should have periodic thyroid function tests and be treated as indicated (see [Endocrine and Metabolism](#)).

In patients with hypopituitarism (multiple hormonal deficiencies) standard hormonal replacement therapy should be monitored closely when human growth hormone therapy is administered.

Hypothyroidism may develop during treatment with human growth hormone. Inadequate treatment of hypothyroidism may prevent optimal response to human growth hormone. Thyroid function should be evaluated before starting growth hormone therapy and regularly assessed during treatment and should be treated with thyroid hormone when indicated.

Serum levels of inorganic phosphorus, alkaline phosphatase, and parathyroid hormone (PTH) may increase with growth hormone therapy.

Patients on growth hormone therapy should be monitored for the emergence of any new malignancy and the treatment discontinued if a new tumor or signs of relapse are detected.

Patients with growth failure due to chronic renal failure should be regularly examined and monitored for progression of renal osteodystrophy. Slipped capital femoral epiphysis or avascular necrosis of the femoral head may occur in children with advanced renal osteodystrophy and it is uncertain whether these complications are affected by growth hormone therapy. In these patients, radiologic imaging of the hips and laboratory exams (serum calcium, phosphorus, alkaline phosphatase and PTH) should be made prior to initiating growth hormone therapy and regularly followed subsequently. Physicians and parents should be alert to the development of a limp or complaints of hip or knee pain (as knee pain may be referred hip pain) in patients treated with growth hormone therapy.

Bone age should be monitored periodically during SAIZEN administration especially in patients who are pubertal and/or receiving concomitant thyroid replacement therapy. Under these circumstances, epiphyseal maturation may progress rapidly.

Patients with growth hormone deficiency secondary to an intracranial lesion should be examined frequently for progression or recurrence of the underlying disease process.

Patients with an intra or extracranial neoplasia in remission who are receiving treatment with growth hormone should be examined carefully and at regular intervals by the physician. Patients developing neoplasia should be reported to Health Canada by the treating physician.

In short children born SGA other medical reasons or treatments that could explain growth disturbance should be ruled out before starting treatment.

For SGA patients, it is recommended to measure IGF-I level before starting treatment and twice a year thereafter. If on repeated measurements IGF-I levels exceed +2 SD compared to references for age and pubertal status, the IGF-I/IGF Binding Protein-3 (BP-3) ratio could be taken into account to consider dose adjustment.

Experience in initiating treatment in SGA patients near onset of puberty is limited. It is therefore not recommended to initiate treatment near onset of puberty. Experience with SGA patients with Silver-Russell syndrome is limited.

Some of the height gain obtained by treating short children born SGA with somatropin may be lost if treatment is stopped before final height is reached.

Growth hormone administration is followed by a transient phase of hypoglycemia of approximately 2 hours, then from 2-4 hours onward by an increase in blood glucose levels despite high insulin concentrations. To detect insulin resistance, patients should be monitored for evidence of glucose intolerance.

Musculoskeletal

Increased tissue turgor and musculoskeletal discomfort may occur during treatment with growth hormones (see [8 Adverse Reactions](#)). These symptoms may resolve spontaneously, with analgesic therapy, or after reducing the dosage.

Carpal tunnel syndrome may occur during treatment with growth hormone, SAIZEN (see [8 Adverse Reactions](#)). If the symptoms of carpal tunnel syndrome do not resolve by decreasing the dosage of growth hormone, it is recommended that treatment be discontinued.

Patients with endocrine disorders, including growth hormone deficiency (GHD), as well as those with rapid growth, are at increased risk of epiphysiolyis of the hip compared to the general population. Cases of epiphysiolyis capitis femoris including slipped capital femoral epiphysis (SCFE) and/or avascular necrosis (osteonecrosis) of the femoral head (including Legg-Calve-Perthes disease) have been reported in children treated with SAIZEN. Physicians and parents should be alert to the development of a limp or complaints of hip or knee pain in children being treated or who have been treated with SAIZEN.

In case of persistent edema or severe paraesthesia the dosage should be decreased in order to avoid the development of carpal tunnel syndrome. Growth hormone deficiency in the adult is a lifelong condition and should be treated accordingly, however experience with patients over sixty years and experience with prolonged treatment is limited.

SAIZEN has not been shown to increase the incidence of scoliosis. Progression of scoliosis can occur in pediatric patients who experience rapid growth. Because somatropin increases growth rate, patients with a history of scoliosis who are treated with growth hormone should be monitored for progression of scoliosis.

Neurologic

Intracranial Hypertension

Intracranial hypertension (IH) with papilloedema, visual changes, headache, nausea and/or vomiting has been reported in a small number of patients treated with growth hormone products. Symptoms usually occurred within the first eight weeks of initiation of growth hormone therapy. In all reported cases, IH-associated signs and symptoms resolved after discontinuation of therapy or a reduction of growth hormone dose. Physicians and parents should be attentive to relevant symptoms such as headache and visual problems in patients under SAIZEN therapy. Fundoscopic examination should be performed routinely before initiating treatment with SAIZEN to exclude pre-existent papilloedema and repeated if there is any clinical suspicion. If papilloedema is confirmed by fundoscopy, SAIZEN treatment should be stopped. It can be restarted at a lower dose after idiopathic-intracranial hypertension has resolved which occurs rapidly when treatment is withdrawn. If growth hormone treatment is restarted, careful monitoring for symptoms of intracranial hypertension is necessary, and treatment should be discontinued if intracranial hypertension reoccurs.

Perioperative Considerations and Acute Critical Illness

In clinical studies, a significant increase in morbidity and mortality has been reported among somatropin treated patients with acute critical illness in intensive care units due to complications following cardiac surgery, abdominal surgery, multiple accident trauma or acute respiratory failure (see [2 Contraindications](#)). Mortality was higher in patients treated with 5.3 mg or 8 mg somatropin daily (41.9 %) compared to patients receiving placebo (19.3 %). Based on this information, these patients must not be treated with somatropin.

Reproductive Health

No adequate and well controlled studies in pregnant women have been performed. From the reproductive studies performed in animals with SAIZEN, there is no evidence of an increased risk of adverse reactions for the embryo or foetus (see [16 Non-Clinical Toxicology](#)).

Skin

SC administration of SAIZEN can result in lipoatrophy (depression in the skin). Patients should be advised to consult their doctor if they notice any of these conditions.

Injection sites should be varied to prevent localized lipoatrophy, in particular with long term subcutaneous (SC) administration of SAIZEN.

7.1. Special Populations

Adult Patients:

Patients with epiphyseal closure who were treated with growth hormone replacement therapy in childhood should be re-evaluated according to the criteria in [1 Indications](#) before continuation of SAIZEN at the reduced dose level required for growth hormone-deficient adults.

Experience with prolonged treatment in adults is limited.

7.1.1. Pregnancy

There are no adequate and well controlled studies in pregnant women. Therefore, SAIZEN should be used during pregnancy only if it is clearly indicated and under medical supervision. Animal studies do not indicate direct or indirect harmful effects with respect to pregnancy, embryo-fetal development, parturition or postnatal development (reproductive toxicology studies do not indicate any adverse effect on fertility and reproduction, despite administration of doses sufficiently high to produce some pharmacological effects on growth) (see [16 Non-Clinical Toxicology](#)).

Patients shall inform their doctor if they are pregnant or are contemplating pregnancy. Caution should be exercised when prescribing to pregnant women.

7.1.2. Breastfeeding

It is not known whether this drug is excreted in human milk. Due to the large molecular weight, it is unlikely that it would be passed intact into the maternal milk and absorption of intact protein from the gastrointestinal tract of the infant is also unlikely. However, secretion of breakdown products of the drug in breast milk has not been studied. Because many drugs are excreted in human milk, caution should be exercised when somatropin is administered to a nursing mother.

7.1.3. Pediatrics

Pediatrics (<18 years of age): SAIZEN is indicated for use in children (see [1 Indications](#)). Patients with endocrine disorders, including growth hormone deficiency, may develop slipped capital epiphyses more frequently. Any pediatric patient with an unexplained limp should be evaluated by a healthcare professional (see [7 Warnings and Precautions, Musculoskeletal](#)).

7.1.4. Geriatrics

Geriatrics (≥ 65 years of age): No data are available to Health Canada; therefore, Health Canada has not authorized an indication for geriatric use.

8. Adverse Reactions

8.1. Adverse Reaction Overview

Carcinogenesis and Genotoxicity

Frequency not known

Some cases of acute leukemia have been reported in growth hormone deficient children, untreated as well as treated with growth hormone, and might possibly represent a slightly increased incidence compared with non-growth hormone deficient children. A causal relationship to growth hormone therapy has not yet been established.

General Disorders and Administration Site Conditions

Common

Injection site reactions: Some patients may experience redness and itching at the site of injection, particularly when the subcutaneous route is used.

Localized lipoatrophy, which can be avoided by varying the site of injection

Common (in adults), Uncommon (in children)

Fluid retention: peripheral edema, stiffness, arthralgia, myalgia, paresthesia

Immune System Disorders

Frequency not known

Localized and generalized hypersensitivity reactions

Nervous System Disorders

Common

Headache

Uncommon

Benign intracranial hypertension, which can be resolved after discontinuation of therapy or a reduction of growth hormone dose

Endocrine Disorders

Very Rare

Hypothyroidism

Gastrointestinal Disorders

Frequency not known

Pancreatitis

Musculoskeletal disorders

Very Rare

Slipped capital femoral epiphysis (epiphysiolysis) at the site of the hip joint or avascular necrosis (osteonecrosis) of the femoral head including Legg-Calve-Perthes disease may occur. A child with an unexplained limp should be examined.

Metabolism disorders

Common

Hyperglycemia

Frequency not known

Hyperinsulinism, insulin resistance

Intermittent dosage has been associated with the appearance of hypoglycemia.

Toxicity in newborns has been associated with benzyl alcohol as a preservative (see [7 Warnings and Precautions](#)).

8.2. Clinical Trial Adverse Reactions

Clinical trials are conducted under very specific conditions. Therefore, the frequencies of adverse reactions observed in the clinical trials may not reflect frequencies observed in clinical practice and should not be compared to frequencies reported in clinical trials of another drug.

Adult Growth Hormone Deficiency

The safety of SAIZEN replacement therapy in adults was evaluated in a randomized, double-blind, placebo-controlled (DBPC) study involving 115 patients (see [14.1 Clinical Trials by Indication, Adult Indications](#)). During the 6-month study, adverse reactions were reported in 56 patients (93.3%) in the somatropin-treated group and 42 patients (76.4%) in the placebo-treated group.

Withdrawals for this study during both the double-blind, placebo controlled phase and the open-label phase were due to patient decision (7%), protocol violation (0.9%), adverse events (12.2%), lost to follow-up (0.9%) and other (2.6%).

Edema and edema peripheral (>10%), myalgia (<10%), arthralgia (up to 35.5%), and other joint disorders (<10%) were reported to occur in adult patients receiving growth hormone replacement therapy. These side effects occurred primarily early in therapy and tended to be transient.

Adult patients with growth hormone deficiency, following diagnosis of growth hormone deficiency in childhood, reported side effects less frequently than those with adult onset growth hormone deficiency.

Table 3 - Percentage of Patients with Adverse Events by System Organ Class, Preferred Term (whether on or off treatment with SAIZEN)

System Organ Class	Preferred Term	On treatment (N=107)	Off treatment (N=44)
Musculoskeletal and connective tissue disorders	Arthralgia	35.5%	18.2%
	Back pain	13.1%	6.8%
	Myalgia	9.3%	2.3%
	Pain in extremity	9.3%	2.3%
	Joint swelling	6.5%	6.8%
	Joint stiffness	6.5%	
	Musculoskeletal stiffness	4.7%	2.3%
	Tendonitis	4.7%	
	Groin pain	1.9%	4.5%
	Shoulder pain	3.7%	
	Muscle spasms	1.9%	2.3%
	Pain in jaw	1.9%	2.3%
	Chest wall pain	1.9%	
	Dupuytren's contracture	1.9%	
	Musculoskeletal discomfort	1.9%	
Nervous system disorders	Headache	20.6%	18.2%
	Paraesthesia	9.3%	2.3%
	Dizziness	7.5%	6.8%
	Carpal tunnel syndrome	7.5%	4.5%
	Hypoaesthesia	6.5%	
	Sinus headache	3.7%	2.3%
	Sciatica	1.9%	4.5%
	Memory impairment	2.8%	
	Loss of consciousness	1.9%	
Infections and infestations	Influenza	17.8%	4.5%
	Nasopharyngitis	14.0%	2.3%
	Lower respiratory tract infection	9.3%	6.8%
	Urinary tract infection	4.7%	2.3%
	Bronchitis	3.7%	4.5%
	Upper respiratory tract infection	4.7%	
	Tooth abscess	4.7%	
	Ear infection	1.9%	2.3%

Table 3 - Percentage of Patients with Adverse Events by System Organ Class, Preferred Term (whether on or off treatment with SAIZEN)

System Organ Class	Preferred Term	On treatment (N=107)	Off treatment (N=44)
	Gastroenteritis	2.8%	
	Otitis externa	1.9%	
	Tonsillitis	1.9%	
General disorders and administration site conditions	Edema peripheral	16.8%	9.1%
	Fatigue	10.3%	4.5%
	Influenza like illness	6.5%	2.3%
	Injection site bruising	1.9%	6.8%
	Asthenia	2.8%	4.5%
	Chest pain	2.8%	
	Edema	1.9%	2.3%
	Pyrexia	2.8%	
	Chills	1.9%	
	Pain	1.9%	
Investigations	Free fatty acids increased	10.3%	4.5%
	Insulin-like growth factor increased	7.5%	
	Blood cholesterol increased	2.8%	2.3%
	Thyroxine free decreased	3.7%	
	Alanine aminotransferase increased	2.8%	
	Glycosylated hemoglobin increased	2.8%	
	Weight decreased	2.8%	
	Neutrophil count increased	1.9%	
	White blood cell count increased	1.9%	
	Blood urine present	1.9%	
	Lymphocyte count decreased	1.9%	
Gastrointestinal disorders	Nausea	6.5%	6.8%
	Diarrhea	7.5%	2.3%
	Abdominal pain upper	7.5%	
	Vomiting	4.7%	
	Abdominal distension	2.8%	
	Abdominal pain	2.8%	
	Stomach discomfort	2.8%	
	Abdominal discomfort	1.9%	
	Frequent bowel movements	1.9%	
	Gastroesophageal reflux disease	1.9%	
Metabolism and nutrition disorders	Fluid retention	8.4%	
	Hyperglycemia	3.7%	2.3%
	Dehydration	2.8%	

Table 3 - Percentage of Patients with Adverse Events by System Organ Class, Preferred Term (whether on or off treatment with SAIZEN)

System Organ Class	Preferred Term	On treatment (N=107)	Off treatment (N=44)
	Dyslipidemia	2.8%	
Skin and subcutaneous tissue disorders	Hyperhidrosis	4.7%	
	Rash	2.8%	2.3%
	Skin disorder	2.8%	
	Pruritus	1.9%	
	Nail pigmentation	1.9%	
Respiratory, thoracic and mediastinal disorders	Pharyngolaryngeal pain	4.7%	4.5%
	Cough	2.8%	6.8%
	Dyspnea	1.9%	4.5%
	Nasal congestion	2.8%	2.3%
Psychiatric disorders	Insomnia	7.5%	
	Depression	4.7%	
	Depressive delusion	2.8%	
	Anxiety	1.9%	
	Depressed mood	1.9%	
Eye disorders	Conjunctivitis	2.8%	
	Vision blurred	2.8%	
	Eye pain	1.9%	2.3%
Renal and urinary disorders	Hematuria	8.4%	4.5%
	Nephrolithiasis	1.9%	
Injury, poisoning and procedural complications	Fall	1.9%	
	Joint dislocation	1.9%	
	Joint sprain	1.9%	
Vascular disorders	Hypertension	3.7%	6.8%
	Hypotension	1.9%	
Endocrine disorders	Hypothyroidism	6.5%	
	Hyperthyroidism	1.9%	
Reproductive system and breast disorders	Metrorrhagia	2.8%	
	Dysmenorrhea	1.9%	
Blood and lymphatic system disorders	Anaemia	1.9%	
	Lymphadenopathy	1.9%	
Neoplasms benign, malignant and unspecified (incl cysts and polyps)	Pituitary tumour	1.9%	
Ear and labyrinth disorders	Ear discomfort	1.9%	2.3%
	Ear pain	1.9%	

8.2.1. Clinical Trial Adverse Reactions – Pediatrics

Growth Hormone Deficiency

In clinical studies in which SAIZEN was administered to growth hormone deficient children (see [14.1 Clinical Trials by Indication, Pediatric Indications](#)), the following reactions were infrequently seen: local reactions at the injection site (such as pain, numbness, redness and swelling), hypothyroidism, hypoglycemia, seizures, exacerbation of preexisting psoriasis and disturbances in fluid balance.

The most frequently reported AEs were those commonly reported in any pediatric patient population, with pharyngolaryngeal pain (32.6%), pyrexia (38.4%), cough (30.4%), headache (37.5%), nasopharyngitis (21.4%), upper respiratory tract infection (17.9%) and influenza (17.0%) being the most frequently reported. These events were well tolerated without the need for hospitalisation. In addition to the treatment related adverse events reported above, two patients developed anti-hGH antibodies. In both cases, the antibodies did not have any growth inhibiting effect. None of the patients developed antibodies to host cell protein. Three transfer patients who had anti-hGH antibodies prior to treatment became negative within 6 months of treatment with SAIZEN. Hypothyroidism (5.8%) and decreased thyroxine (19.2%) were seen in several patients. One patient died of recurrent craniopharyngioma and one patient experienced lipoatrophy.

Table 4 - Percentage of Patients with Adverse Events by System Organ Class, Preferred Term (whether on or off treatment with SAIZEN)

System Organ Class	Preferred Term	On treatment (N=224)	Off treatment (N=15)
Infections and infestations	Nasopharyngitis	21.4%	
	Upper respiratory tract infection	17.9%	
	Influenza	17.0%	
	Bronchitis	12.5%	
	Otitis media	9.4%	
	Ear infection	8.9%	
	Varicella	7.6%	6.7%
	Pharyngitis streptococcal	7.1%	
	Pharyngitis	6.7%	
	Tonsillitis	6.7%	
	Viral infection	6.7%	
	Rhinitis	6.3%	
	Sinusitis	5.8%	6.7%
	Gastroenteritis viral	6.3%	
	Urinary tract infection	5.4%	
	Gastroenteritis	5.4%	
	Acute tonsillitis	4.0%	6.7%
	Scarlet fever	4.5%	
	Pneumonia	4.0%	
	Herpes simplex	1.8%	
	Rubella	1.8%	

Table 4 - Percentage of Patients with Adverse Events by System Organ Class, Preferred Term (whether on or off treatment with SAIZEN)

System Organ Class	Preferred Term	On treatment (N=224)	Off treatment (N=15)
	Mumps	1.3%	
	Impetigo	1.3%	
	Respiratory tract infection	1.3%	
	Viral pharyngitis	1.3%	
General disorders and administration site conditions	Pyrexia	38.4%	6.7%
	Injection site pain	6.7%	
	Fatigue	6.3%	
	Asthenia	1.8%	
	Chest pain	1.3%	
	Injection site bruising	1.3%	
	Injection site reaction	1.3%	
Respiratory, thoracic and mediastinal disorders	Pharyngolaryngeal pain	32.6%	
	Cough	30.4%	
	Nasal congestion	7.6%	
	Epistaxis	6.7%	
	Rhinorrhea	4.9%	
	Asthma	3.1%	
	Rhinitis allergic	3.1%	
	Paranasal sinus hypersecretion	1.3%	
	Wheezing	1.3%	
	Dysphonia	1.3%	
	Sinus congestion	1.3%	
Nervous system disorders	Headache	37.5%	
	Convulsion	3.6%	
	Dizziness	2.7%	
	Epilepsy	2.2%	
	Disturbance in attention	1.8%	
	Lethargy	1.3%	
Gastrointestinal disorders	Vomiting	17.4%	
	Diarrhea	11.6%	
	Abdominal pain upper	9.8%	
	Gastrointestinal disorder	7.6%	
	Abdominal pain	5.4%	
	Nausea	4.5%	
	Stomach discomfort	4.0%	
	Constipation	2.7%	
	Toothache	1.3%	
	Dental discomfort	1.3%	
Investigations	Thyroxine decreased	19.2%	

Table 4 - Percentage of Patients with Adverse Events by System Organ Class, Preferred Term (whether on or off treatment with SAIZEN)

System Organ Class	Preferred Term	On treatment (N=224)	Off treatment (N=15)
	Hormone level abnormal	2.2%	
	Blood triglycerides increased	1.8%	6.7%
	Body temperature increased	1.8%	
	Weight increased	1.3%	
	Cardiac murmur	1.3%	
	Drug specific antibody present	1.3%	
Surgical and medical procedures	Substitution therapy	12.9%	
	Tooth extraction	2.7%	
	Appendicectomy	1.8%	
	Myringotomy	1.8%	
	Tonsillectomy	1.3%	
Musculoskeletal and connective tissue disorders	Arthralgia	12.9%	
	Pain in extremity	9.4%	
	Muscle spasms	2.7%	
	Back pain	1.8%	
	Bone pain	1.3%	
	Myalgia	1.3%	
Skin and subcutaneous tissue disorders	Rash	7.1%	
	Eczema	2.7%	
	Pruritus	2.7%	
	Psoriasis	1.8%	13.3%
	Erythema	2.2%	
	Urticaria	1.8%	
	Acne	1.3%	
Injury, poisoning and procedural complications	Treatment noncompliance	3.6%	
	Joint injury	2.2%	
	Hand fracture	2.2%	
	Road traffic accident	1.8%	6.7%
	Fall	1.8%	
	Skin laceration	1.8%	
	Arthropod bite	1.3%	
	Joint sprain	1.3%	
	Foot fracture	1.3%	
Endocrine disorders	Delayed puberty	6.3%	
	Hypothyroidism	5.8%	6.7%
	Hypogonadism	2.2%	
	Hypopituitarism	1.8%	
	Diabetes insipidus	1.3%	

Table 4 - Percentage of Patients with Adverse Events by System Organ Class, Preferred Term (whether on or off treatment with SAIZEN)

System Organ Class	Preferred Term	On treatment (N=224)	Off treatment (N=15)
	Secondary hypogonadism	1.3%	
Ear and labyrinth disorders	Ear pain	11.6%	
	Hypoacusis	1.3%	
	Middle ear effusion	1.3%	
	Otorrhea	1.3%	
Metabolism and nutrition disorders	Hypoglycemia	3.1%	
	Iron deficiency	2.2%	
	Obesity	2.2%	
	Insulin resistance	1.3%	
Reproductive system and breast disorders	Varicocele	2.2%	
	Gynaecomastia	1.8%	
Immune system disorders	Hypersensitivity	4.5%	
	Seasonal allergy	4.5%	
Psychiatric disorders	Mental disorder	1.8%	
	Nervousness	1.8%	
	Depression	1.3%	
Blood and lymphatic system disorders	Lymphadenopathy	2.7%	
	Anaemia	2.2%	
Renal and urinary disorders	Enuresis	2.7%	
Eye disorders	Conjunctivitis	2.2%	
Congenital, familial and genetic disorders	Cryptorchism	2.7%	13.3%
Neoplasms benign, malignant and unspecified (incl cysts and polyps)	Craniopharyngioma	1.8%	

Turner Syndrome

A clinical study was conducted in 91 girls with Turner syndrome to receive either SAIZEN alone or in conjunction with oxandrolone (see Dosage Regimen Table under [14 Clinical Trials](#)).

In girls treated with SAIZEN alone the percentage of patients who experienced specific adverse events were: skin reaction at injection site (13%), pain at injection site (7%), deepening/hoarseness of voice (7%), pain in limbs (7%), pigmented naevi (4%), clitorimegaly (3%), hypercholesterolaemia (3%), and 1% each for edema, hair loss, increased ephelides and seborrhea.

For the group treated with SAIZEN and oxandrolone the percentage of patients who experienced adverse events were: clitoromegaly (30%), pain in limbs (11%), deepening/hoarseness of voice (9%), pain at injection site (9%), skin reaction at injection site (8%), elevated creatinine kinase (4%), hypercholesterolaemia (4%), and 2% each for virilization, exanthem, hyperlipidemia, pigmented naevi, edema, lipodystrophy, hematoma, muscle cramps, increased freckles and hair loss. Thus, the addition of

oxandrolone was associated with some virilizing effects, especially at doses of more than 0.05 mg/kg daily.

A total of 18 (20%) patients exhibited a treatment emergent abnormality in the response to glucose loading at some time during the study, of whom only 7 patients (7.7%) had detectable glucose intolerance on two or more occasions. Four patients discontinued treatment in association with these abnormalities. It should be noted that impaired glucose tolerance is commonly found in Turner syndrome patients.

Table 5 - Percentage of Patients with Adverse Events by System Organ Class, Preferred Term (whether on or off treatment with SAIZEN)

System Organ Class	Preferred Term	On treatment (N=81)	Off treatment (N=0)
Respiratory, thoracic and mediastinal disorders	Cough	51.9%	
	Pharyngolaryngeal pain	45.7%	
	Epistaxis	14.8%	
	Dysphonia	11.1%	
	Rhinorrhea	8.6%	
	Vocal cord thickening	1.2%	
General disorders and administration site conditions	Pyrexia	46.9%	
	Injection site reaction	16.0%	
	Injection site pain	12.3%	
	Edema	2.5%	
	Localised Edema	1.2%	
Infections and infestations	Rhinitis	17.3%	
	Influenza	9.9%	
	Ear infection	6.2%	
	Otitis media	6.2%	
	Bronchitis	4.9%	
	Sinusitis	4.9%	
	Fungal infection	3.7%	
	Nasopharyngitis	3.7%	
	Urinary tract infection	3.7%	
	Varicella	3.7%	
	Fungal skin infection	2.5%	
	Scarlet fever	2.5%	
	Tonsillitis	2.5%	
	Pneumonia	1.2%	
	Candidiasis	1.2%	
	Acute tonsillitis	1.2%	
	Gastroenteritis	1.2%	
	Helminthic infection	1.2%	
	Herpes zoster	1.2%	

Table 5 - Percentage of Patients with Adverse Events by System Organ Class, Preferred Term (whether on or off treatment with SAIZEN)

System Organ Class	Preferred Term	On treatment (N=81)	Off treatment (N=0)
	Measles	1.2%	
	Meningitis viral	1.2%	
	Mumps	1.2%	
	Otitis media chronic	1.2%	
	Paronychia	1.2%	
	Pertussis	1.2%	
	Pharyngitis	1.2%	
	Respiratory tract infection	1.2%	
	Viral infection	1.2%	
	Vulvitis	1.2%	
Nervous system disorders	Headache	44.4%	
	Petit mal epilepsy	1.2%	
	Convulsion	1.2%	
	Dizziness	1.2%	
	Epilepsy	1.2%	
	Febrile convulsion	1.2%	
	Hypertonia	1.2%	
Ear and labyrinth disorders	Ear pain	28.4%	
	Hearing impaired	2.5%	

Chronic Renal Failure

In clinical studies with SAIZEN in Chronic Renal Failure (see [14 Clinical Trials](#)), the following adverse events were considered possibly related to treatment by the investigator: pseudotumor cerebri, deterioration of renal function, hyperthyroidism, injection site infection, renal transplant rejection, papilloedema, hypothyroidism, impaired OGTT and abnormal SGOT.

Table 6 - Percentage of Patients with Adverse Events by System Organ Class, Preferred Term (whether on or off treatment with SAIZEN)

System Organ Class	Preferred Term	On treatment (N=65)	Off treatment (N=11)
Infections and infestations	Upper respiratory tract infection	26.2%	
	Otitis media	20.0%	
	Viral infection	20.0%	
	Catheter related infection	18.5%	9.1%
	Rhinitis	16.9%	
	Urinary tract infection	13.8%	
	Influenza	13.8%	
	Gastroenteritis	10.8%	
	Nasopharyngitis	9.2%	
	Herpes simplex	7.7%	
	Bronchitis	7.7%	
	Pharyngitis	7.7%	

Table 6 - Percentage of Patients with Adverse Events by System Organ Class, Preferred Term (whether on or off treatment with SAIZEN)

System Organ Class	Preferred Term	On treatment (N=65)	Off treatment (N=11)
	Tonsillitis	7.7%	
	Varicella	7.7%	
	Staphylococcal infection	6.2%	
	Pyelonephritis	4.6%	
	Sinusitis	4.6%	
	Tonsillitis streptococcal	4.6%	
	Sepsis	3.1%	
	Cytomegalovirus infection	3.1%	
	Infection parasitic	3.1%	
	Hepatitis C	3.1%	
	Localised infection	3.1%	
	Pharyngitis streptococcal	3.1%	
	Respiratory tract infection viral	3.1%	
	Streptococcal infection	3.1%	
	Ear infection	1.5%	
	Fungal infection	1.5%	
	Abscess	1.5%	
	Appendicitis	1.5%	
	Cystitis	1.5%	
	Dental caries	1.5%	
	Endotoxic shock	1.5%	
	Gastroenteritis viral	1.5%	
	Hepatitis B	1.5%	
	Herpes zoster	1.5%	
	Infection	1.5%	
	Injection site abscess	1.5%	
	Laryngitis	1.5%	
	Onychomycosis	1.5%	
	Paronychia	1.5%	
	Pseudomonas infection	1.5%	
	Scarlet fever	1.5%	
	Skin infection	1.5%	
	Staphylococcal sepsis	1.5%	
	Vaginal candidiasis	1.5%	
	Vaginal infection	1.5%	
	Viral rash	1.5%	
Gastrointestinal disorders	Peritonitis	20.0%	
	Vomiting	20.0%	
	Diarrhea	15.4%	
	Abdominal pain	7.7%	

Table 6 - Percentage of Patients with Adverse Events by System Organ Class, Preferred Term (whether on or off treatment with SAIZEN)

System Organ Class	Preferred Term	On treatment (N=65)	Off treatment (N=11)
	Inguinal hernia	7.7%	
	Gingival hyperplasia	7.7%	
	Constipation	6.2%	
	Nausea	6.2%	
	Abdominal pain upper	4.6%	
	Tooth disorder	3.1%	
	Dysphagia	3.1%	
	Abdominal hernia	1.5%	
	Breath odour	1.5%	
	Colonic polyp	1.5%	
	Dyspepsia	1.5%	
	Enteritis	1.5%	
	Faecal incontinence	1.5%	
	Food poisoning	1.5%	
	Frequent bowel movements	1.5%	
	Gastritis	1.5%	
	Gingival hypertrophy	1.5%	
	Intestinal obstruction	1.5%	
	Salivary gland enlargement	1.5%	
	Stomach discomfort	1.5%	
	Umbilical hernia	1.5%	
General disorders and administration site conditions	Pyrexia	23.1%	9.1%
	Fatigue	4.6%	
	Face edema	4.6%	
	Edema peripheral	3.1%	
	Chest pain	3.1%	
	Gait disturbance	3.1%	
	Injection site hemorrhage	3.1%	
	Injection site pain	3.1%	
	Local swelling	1.5%	
	Asthenia	1.5%	
	Catheter site inflammation	1.5%	
	Chills	1.5%	
	Difficulty in walking	1.5%	
	Generalised edema	1.5%	
	Inflammation	1.5%	
	Influenza like illness	1.5%	
	Injection site reaction	1.5%	
	Edema	1.5%	
Nervous system disorders	Headache	23.1%	
	Dizziness	4.6%	
	Convulsion	3.1%	

Table 6 - Percentage of Patients with Adverse Events by System Organ Class, Preferred Term (whether on or off treatment with SAIZEN)

System Organ Class	Preferred Term	On treatment (N=65)	Off treatment (N=11)
	Psychomotor hyperactivity	1.5%	
	Benign intracranial hypertension	1.5%	
	Paraesthesia	1.5%	
	Petit mal epilepsy	1.5%	
	Balance disorder	1.5%	
	Brain edema	1.5%	
	Cerebral infarction	1.5%	
	Coordination abnormal	1.5%	
	Hypertonia	1.5%	
	Mental retardation severity unspecified	1.5%	
	Nervous system disorder	1.5%	
	Optic neuritis	1.5%	
Surgical and medical procedures	Renal transplant	27.7%	45.5%
Renal and urinary disorders	Renal failure	12.3%	9.1%
	Dysuria	4.6%	
	Renal impairment	4.6%	
	Proteinuria	3.1%	
	Enuresis	3.1%	
	Hydronephrosis	1.5%	
	Renal disorder	1.5%	
	Bladder disorder	1.5%	
	Hematuria	1.5%	
	Micturition disorder	1.5%	
	Neurogenic bladder	1.5%	
	Pyuria	1.5%	
	Urethral disorder	1.5%	
	Urinary retention	1.5%	
Respiratory, thoracic and mediastinal disorders	Pharyngolaryngeal pain	15.4%	
	Cough	13.8%	
	Rhinitis allergic	6.2%	9.1%
	Rhinorrhea	4.6%	
	Epistaxis	3.1%	
	Adenoidal hypertrophy	3.1%	
	Pulmonary edema	3.1%	
	Wheezing	3.1%	
	Asthma	1.5%	
	Pharyngeal erythema	1.5%	
	Respiratory tract congestion	1.5%	
	Atelectasis	1.5%	

Table 6 - Percentage of Patients with Adverse Events by System Organ Class, Preferred Term (whether on or off treatment with SAIZEN)

System Organ Class	Preferred Term	On treatment (N=65)	Off treatment (N=11)
	Dyspnoea	1.5%	
	Mediastinal disorder	1.5%	
	Nasal congestion	1.5%	
	Nasal discomfort	1.5%	
	Rales	1.5%	
	Sneezing	1.5%	
	Throat irritation	1.5%	
Musculoskeletal and connective tissue disorders	Pain in extremity	13.8%	
	Arthralgia	10.8%	
	Neck pain	4.6%	
	Back pain	4.6%	
	Renal osteodystrophy	3.1%	
	Bone pain	3.1%	
	Groin pain	1.5%	
	Aseptic necrosis bone	1.5%	
	Muscle spasms	1.5%	
	Musculoskeletal discomfort	1.5%	
	Osteochondrosis	1.5%	
	Arthropathy	1.5%	
	Epiphysiolysis	1.5%	
	Knee deformity	1.5%	
	Lower limb deformity	1.5%	
	Myalgia	1.5%	
	Rickets	1.5%	
	Shoulder pain	1.5%	
Metabolism and nutrition disorders	Hypercalcaemia	7.7%	
	Hyperkalaemia	6.2%	
	Glucose tolerance impaired	3.1%	
	Fluid overload	3.1%	
	Hyperphosphataemia	3.1%	
	Decreased appetite	3.1%	
	Hyperuricaemia	1.5%	9.1%
	Anorexia	1.5%	
	Diabetes mellitus	1.5%	
	Fluid retention	1.5%	
	Hyperglycemia	1.5%	
	Hyperlipidaemia	1.5%	
	Hypermagnesaemia	1.5%	
	Hypervolaemia	1.5%	
	Hypokalaemia	1.5%	
	Malnutrition	1.5%	
	Metabolic acidosis	1.5%	

Table 6 - Percentage of Patients with Adverse Events by System Organ Class, Preferred Term (whether on or off treatment with SAIZEN)

System Organ Class	Preferred Term	On treatment (N=65)	Off treatment (N=11)
Injury, poisoning and procedural complications	Injury	13.8%	
	Accidental overdose	3.1%	
	Fall	3.1%	
	Arthropod bite	1.5%	
	Excoriation	1.5%	
	Post procedural vomiting	1.5%	
Investigations	Blood creatinine increased	7.7%	
	Weight decreased	3.1%	
	Aspartate aminotransferase increased	1.5%	
	Blood parathyroid hormone increased	1.5%	
	Blood pressure decreased	1.5%	
	Blood urea increased	1.5%	
	Blood urine present	1.5%	
	Body temperature increased	1.5%	
	Liver function test abnormal	1.5%	
	Metabolic function test	1.5%	
	Urine output decreased	1.5%	
Skin and subcutaneous tissue disorders	Acne	4.6%	
	Rash	1.5%	
	Onychorrhexis	1.5%	
	Rash pruritic	1.5%	
	Skin depigmentation	1.5%	
	Skin hypertrophy	1.5%	
	Skin nodule	1.5%	
	Urticaria	1.5%	
	Alopecia	1.5%	
	Angioneurotic edema	1.5%	
	Dermatitis contact	1.5%	
	Ecchymosis	1.5%	
	Eczema	1.5%	
	Nail dystrophy	1.5%	
	Psoriasis	1.5%	
	Rash papular	1.5%	
	Skin discolouration	1.5%	
	Skin inflammation	1.5%	
	Skin reaction	1.5%	
Immune system disorders	Transplant rejection	6.2%	9.1%
	Hypersensitivity	4.6%	
	Kidney transplant rejection	4.6%	

Table 6 - Percentage of Patients with Adverse Events by System Organ Class, Preferred Term (whether on or off treatment with SAIZEN)

System Organ Class	Preferred Term	On treatment (N=65)	Off treatment (N=11)
	Drug hypersensitivity	1.5%	
Blood and lymphatic system disorders	Anaemia	9.2%	
	Lymphadenopathy	7.7%	
	Nephrogenic anaemia	4.6%	
	Neutropenia	1.5%	
	Coagulopathy	1.5%	
	Leukocytosis	1.5%	
Vascular disorders	Hypertension	9.2%	
	Hypotension	6.2%	
	Hemorrhage	3.1%	
	Hypertensive crisis	1.5%	
	Hematoma	1.5%	
	Hot flush	1.5%	
	Peripheral coldness	1.5%	
	Vasculitis	1.5%	
	Vasospasm	1.5%	
Reproductive system and breast disorders	Gynaecomastia	3.1%	
	Testicular torsion	3.1%	
	Balanitis	1.5%	
	Breast disorder	1.5%	
	Epididymitis	1.5%	
	Testicular disorder	1.5%	
Eye disorders	Eyelid edema	3.1%	
	Eye pain	1.5%	
	Optic atrophy	1.5%	
	Optic discs blurred	1.5%	
	Papilloedema	1.5%	
	Vision blurred	1.5%	
Ear and labyrinth disorders	Ear pain	4.6%	
	Ear disorder	1.5%	
	Hypoacusis	1.5%	
	Ear discomfort	1.5%	
Endocrine disorders	Hyperparathyroidism	4.6%	
	Hypoparathyroidism	1.5%	
	Hypothyroidism	1.5%	
Cardiac disorders	Cyanosis	1.5%	
	Cardiac disorder	1.5%	
	Cardiac failure	1.5%	
	Pericarditis	1.5%	
Congenital, familial and genetic disorders	Hydrocele	3.1%	
	Congenital foot malformation	1.5%	
	Skin papilloma	3.1%	

Table 6 - Percentage of Patients with Adverse Events by System Organ Class, Preferred Term (whether on or off treatment with SAIZEN)

System Organ Class	Preferred Term	On treatment (N=65)	Off treatment (N=11)
Neoplasms benign, malignant and unspecified (incl cysts and polyps)	Parathyroid tumour benign	1.5%	
Psychiatric disorders	Attention deficit/hyperactivity disorder	1.5%	
	Insomnia	1.5%	

Small for Gestational Age

SAIZEN was evaluated in two randomized, controlled Phase III clinical studies in children born SGA: GF 4001 (n=101) and GF 6283 (n=58) (see [14 Clinical Trials](#)). The most common (incidence > 5%) adverse events observed in clinical trials with SGA patients were mild to moderate in severity. The most frequently reported AEs were those commonly reported in any pediatric patient population, with ear infection, bronchitis, nasopharyngitis, gastroenteritis, varicella and pharyngolaryngeal pain being the most frequently reported. These events were well tolerated without the need for drug discontinuation. The number, type and severity of events did not differ between periods with r-hGH treatment and periods with observation without treatment, or between the first and second and third year of r-hGH treatment.

Oral glucose tolerance tests (OGTT) were used during the treatment and observation periods in studies GF 4001 and GF 6283. Increased insulin levels were observed after 18 months of r-hGH treatment in study GF 4001 and this increase was sustained during the 3-year treatment period but were normalised during the follow-up period. Similar results were observed for the children who received continuous treatment (Group TTOO), but were less apparent for children who received intermittent treatment (Group TOTO) in study GF 6283. Abnormal glucose levels during the OGTT indicative of impaired glucose tolerance were observed in a few patients in both studies, and more often during continuous treatment (study GF 4001 and Group TTOO in study GF 6283) than during intermittent treatment (Group TOTO in study GF 6283). There was no report on diabetes mellitus in any of the studies but in one child (patient no. 6283102003) in study GF 6283, who had received r-hGH treatment continuously for 2 years, a fasting glucose value of 11.2 mmol/L was observed after 2 years of observation without treatment. There was no withdrawal in any of the studies due to change in glucose tolerance. These results are similar to those reported in the literature.

Common and very common adverse reactions (frequency ≥ 1%) are tabulated below.

Table 7 - Percentage of Patients with Adverse Events by System Organ Class, Preferred Term (whether on or off treatment with SAIZEN - pooling of GF 4001 and GF 6283)

System Organ Class	Preferred Term	On treatment (N=100)	Off treatment (N=34)
Infections and infestations	Ear infection	19.0%	14.7%
	Bronchitis	19.0%	8.8%

Table 7 - Percentage of Patients with Adverse Events by System Organ Class, Preferred Term (whether on or off treatment with SAIZEN - pooling of GF 4001 and GF 6283)

System Organ Class	Preferred Term	On treatment (N=100)	Off treatment (N=34)
	Nasopharyngitis	14.0%	20.6%
	Gastroenteritis	14.0%	5.9%
	Varicella	13.0%	11.8%
	Otitis media	7.0%	11.8%
	Otitis media acute	4.0%	2.9%
	Tonsillitis	4.0%	
	Laryngitis	4.0%	
	Lung infection	3.0%	2.9%
	Pharyngitis	2.0%	5.9%
	Influenza	2.0%	2.9%
	Urinary tract infection	2.0%	2.9%
	Upper respiratory tract infection	2.0%	
	Viral infection	2.0%	
	Acute tonsillitis		2.9%
	Tooth abscess	1.0%	2.9%
	Cystitis	1.0%	
	Furuncle	1.0%	
	Helicobacter gastritis		2.9%
	Infectious mononucleosis	1.0%	
	Paronychia	1.0%	
	Pneumonia viral	1.0%	
	Respiratory tract infection	1.0%	
	Rubella	1.0%	
	Sepsis	1.0%	
	Sinusitis	1.0%	
	Tracheobronchitis	1.0%	
	Viral rash	1.0%	
Respiratory, thoracic and mediastinal disorders	Pharyngolaryngeal pain	17.0%	8.8%
	Rhinitis	5.0%	5.9%
	Asthma	5.0%	
	Maxillary sinusitis	3.0%	
	Epistaxis	2.0%	
	Lung disorder	1.0%	
	Nasal congestion	1.0%	
	Rales	1.0%	
Gastrointestinal disorders	Diarrhea	2.0%	5.9%
	Constipation	2.0%	2.9%
	Abdominal pain	1.0%	2.9%

Table 7 - Percentage of Patients with Adverse Events by System Organ Class, Preferred Term (whether on or off treatment with SAIZEN - pooling of GF 4001 and GF 6283)

System Organ Class	Preferred Term	On treatment (N=100)	Off treatment (N=34)
	Vomiting	1.0%	2.9%
	Food poisoning	1.0%	
	Gastroesophageal reflux disease	1.0%	
	Inguinal hernia	1.0%	
	Irritable bowel syndrome	1.0%	
	Peritonitis	1.0%	
	Toothache		2.9%
Injury, poisoning and procedural complications	Head injury	2.0%	
	Upper limb fracture	1.0%	2.9%
	Arthropod sting	1.0%	
	Burns second degree		2.9%
	Cervical vertebral fracture	1.0%	
	Contusion	1.0%	
	Fall	1.0%	
	Foreign body trauma	1.0%	
	Joint sprain	1.0%	
	Thermal burn	1.0%	
	Traumatic hematoma	1.0%	
	Wrist fracture	1.0%	
Surgical and medical procedures	Tonsillectomy	4.0%	
	Appendicectomy	3.0%	
	Ear tube insertion	3.0%	
	Myringoplasty	1.0%	
	Gastric operation	1.0%	
	Otorhinolaryngological surgery	1.0%	
	Skin neoplasm excision	1.0%	
Nervous system disorders	Headache	5.0%	5.9%
	Coma	1.0%	
	Dizziness	1.0%	
	Febrile convulsion	1.0%	
	Tremor	1.0%	
Blood and lymphatic system disorders	Anaemia	3.0%	
	Hypochromic anaemia	2.0%	2.9%
	Eosinophilia	2.0%	2.9%
	Thrombocytopenia	2.0%	2.9%
	Pancytopenia	1.0%	
	Granulocytopenia	1.0%	

Table 7 - Percentage of Patients with Adverse Events by System Organ Class, Preferred Term (whether on or off treatment with SAIZEN - pooling of GF 4001 and GF 6283)

System Organ Class	Preferred Term	On treatment (N=100)	Off treatment (N=34)
	Iron deficiency anaemia	1.0%	
Musculoskeletal and connective tissue disorders	Arthralgia	3.0%	2.9%
	Muscle hypertrophy	1.0%	
	Myalgia	1.0%	
	Osteoarthritis	1.0%	
	Osteochondrosis	1.0%	
General disorders and administration site conditions	Pyrexia	4.0%	
	Injection site inflammation	1.0%	
	Difficulty in walking	1.0%	
Skin and subcutaneous tissue disorders	Urticaria	1.0%	2.9%
	Acne	1.0%	
	Henoch-Schonlein purpura	1.0%	
	Pruritus generalised	1.0%	
	Rash	1.0%	
Investigations	Aspartate aminotransferase increased	1.0%	
	Blood glucose increased	1.0%	
	Body temperature increased	1.0%	
	Glycosylated hemoglobin increased	1.0%	
	Hemoglobin decreased	1.0%	
Renal and urinary disorders	Hematuria	1.0%	
	Polyuria		2.9%
	Renal insufficiency	1.0%	
	Ureteric stenosis	1.0%	
Psychiatric disorders	Sleep disorder	2.0%	
	Aggression	1.0%	
	Polydipsia psychogenic	1.0%	
Congenital, familial and genetic disorders	Cryptorchism		2.9%
	Eyelid ptosis congenital	1.0%	
	Pigmented naevus		2.9%
Cardiac disorders	Cardiac disorder	1.0%	
	Cardiac failure	1.0%	
Endocrine disorders	Autoimmune thyroiditis	1.0%	
	Hypothyroidism		2.9%
Immune system disorders	Allergy to animal	1.0%	
	Drug hypersensitivity	1.0%	
Eye disorders	Conjunctivitis		2.9%

Table 7 - Percentage of Patients with Adverse Events by System Organ Class, Preferred Term (whether on or off treatment with SAIZEN - pooling of GF 4001 and GF 6283)

System Organ Class	Preferred Term	On treatment (N=100)	Off treatment (N=34)
Metabolism and nutrition disorders	Glucose tolerance impaired	1.0%	
Neoplasms benign, malignant and unspecified (incl cysts and polyps)	Cyst	1.0%	
Reproductive system and breast disorders	Hypertrophy breast		2.9%

8.3. Less Common Clinical Trial Adverse Reactions

Clinical trial adverse drug reactions in adult patients with a frequency of less than 1% are presented in the following listing.

GHD

Blood and lymphatic system disorders: Microcytic anaemia, Neutropenia, Neutrophilia

Cardiac disorders: Atrial fibrillation, Coronary artery disease, Left ventricular failure

Endocrine disorders: Adrenocortical insufficiency acute, Pituitary cyst, Toxic nodular goitre

Eye disorders: Accommodation disorder, Asthenopia, Cataract, Conjunctivitis allergic, Dry eye, Eyelid cyst, Eyelid edema, Optic discs blurred

Gastrointestinal disorders: Anal fissure, Colitis ulcerative, Constipation, Diverticulum, Dysphagia, Food poisoning, Gastritis, Gastrointestinal hemorrhage, Hemorrhoids, Intestinal polyp, Tooth disorder, Toothache

General disorders and administration site conditions: Application site reaction, Chest discomfort, Discomfort, Facial pain, Injection site pain, Injection site reaction, Malaise

Hepatobiliary disorders: Cholelithiasis, Gallbladder polyp, Hepatic function abnormal

Immune system disorders: Hypersensitivity, Seasonal allergy

Infections and infestations: Breast infection, Cystitis, Eye infection, Gastroenteritis viral, Gingival infection, Helicobacter infection, Herpes zoster oticus, Hordeolum, Infected sebaceous cyst, Labyrinthitis, Pharyngitis, Sinusitis, Viral infection, Vulvovaginal mycotic infection

Injury, poisoning and procedural complications: Ankle fracture, Back injury, Contusion, Laceration, Muscle strain, Procedural pain, Wrist fracture

Investigations: Blood calcium increased, Blood creatinine increased, Blood glucose decreased, Blood potassium decreased, Body temperature decreased, Gamma-glutamyltransferase increased, Hemoglobin increased, Heart rate increased, Hepatic enzyme increased, High density lipoprotein decreased, Liver function test abnormal, Low density lipoprotein increased, Lymph node palpable, Monocyte count decreased, Prostate examination abnormal, Semen volume decreased, Weight increased

Metabolism and nutrition disorders: Decreased appetite, Diabetes mellitus, Hypercholesterolaemia, Hyperkalaemia, Hypertriglyceridaemia, Hypoglycemia, Increased appetite, Iron deficiency, Polydipsia

Musculoskeletal and connective tissue disorders: Arthritis, Arthropathy, Axillary mass, Bone pain, Bursitis, Ganglion, Limb deformity, Muscle fatigue, Muscle tightness, Musculoskeletal chest pain, Musculoskeletal pain, Myopathy, Sacral pain, Synovitis, Tendon disorder

Neoplasms benign, malignant and unspecified (incl. cysts and polyps): Acrochordon, Basal cell carcinoma, Glioma, Hemangioma, Uterine leiomyoma

Nervous system disorders: Cerebellar infarction, Cerebrovascular accident, Disturbance in attention, Drooling, Hyperaesthesia, Lethargy, Syncope, Syncope vasovagal, Tremor, Trigeminal neuralgia, Visual field defect

Psychiatric disorders: Aggression, Early morning awakening, Mood altered, Mood swings, Stress

Renal and urinary disorders: Micturition urgency, Proteinuria, Renal colic, Urinary incontinence

Reproductive system and breast disorders: Breast pain, Breast tenderness, Gynaecomastia, Menorrhagia, Menstruation irregular, Withdrawal bleed, Withdrawal bleeding irregular

Respiratory, thoracic and mediastinal disorders: Asthma, Nasal dryness, Rhinitis allergic, Sinus disorder, Sleep apnoea syndrome

Skin and subcutaneous tissue disorders: Acne, Alopecia, Dermatitis, Dermatitis contact, Erythema, Hypertrichosis, Parapsoriasis, Petechiae, Photosensitivity reaction, Rash generalised, Rash pruritic, Scar, Seborrhoeic dermatitis, Skin inflammation, Skin nodule, Sweat gland disorder, Urticaria

Vascular disorders: Hot flush, Lymphoedema, Orthostatic hypotension

8.3.1. Less Common Clinical Trial Adverse Reactions – Pediatrics

GHD

Blood and lymphatic system disorders: Iron deficiency anaemia, Neutrophilia, Plasmacytosis

Cardiac disorders: Angina pectoris, Aortic valve stenosis, Palpitations

Congenital, familial and genetic disorders: Atrial septal defect, Epidermal naevus, Facial dysmorphism, Foetal alcohol syndrome, Lymphangioma, Pigmented naevus, Turner syndrome

Ear and labyrinth disorders: Deafness, Deafness bilateral, Ear congestion, Ear discomfort, Motion sickness, Tinnitus, Tympanic membrane hyperaemia, Tympanic membrane perforation

Endocrine disorders: Adrenocortical insufficiency chronic, Empty sella syndrome, Hyperthyroidism

Eye disorders: Blepharitis, Blindness, Diplopia, Eye hemorrhage, Eye irritation, Eyelid edema, Eyelid ptosis, Lacrimal cyst, Optic atrophy, Papilloedema, Vision blurred, Visual acuity reduced, Visual disturbance

Gastrointestinal disorders: Abdominal pain lower, Aphthous stomatitis, Dyspepsia, Faeces hard, Flatulence, Gingival bleeding, Gingival hypertrophy, Irritable bowel syndrome, Mouth ulceration, Oral mucosal blistering, Pancreatitis, Salivary gland hypertrophy, Tooth disorder

General disorders and administration site conditions: Adverse drug reaction, Adverse event, Application site pain, Chills, Cyst rupture, Feeling cold, Feeling hot, Hernia, Influenza like illness, Infusion site bruising, Injection site atrophy, Injection site hemorrhage, Injection site hypertrophy, Injection site induration, Injection site irritation, Injection site mass, Injection site rash, Injection site scar, Injection site swelling, Instillation site pruritus, Irritability, Local swelling, Malaise, Mucosal ulceration, No adverse effect, edema peripheral, Pain

Hepatobiliary disorders: Liver disorder

Immune system disorders: Drug hypersensitivity, Multiple allergies, Selective IgG subclass deficiency

Infections and infestations: Acarodermatitis, Appendicitis, Bacteriuria, Body tinea, Conjunctivitis infective, Cystitis, Enterobiasis, Erythema infectiosum, Eye infection, Febrile infection, Fungal infection, Furuncle, Gastric infection, Genital infection, Hepatitis B, Herpes zoster, Herpetic stomatitis, Infectious mononucleosis, Injection site infection, Kidney infection, Lice infestation, Localised infection, Lower respiratory tract infection, Lymph gland infection, Measles, Meningitis viral, Molluscum contagiosum, Nail infection, Oral candidiasis, Orchitis, Otitis externa, Paronychia, Pertussis, Pharyngotonsillitis, Postoperative wound infection, Pyelonephritis, Skin infection, Staphylococcal infection, Streptococcal infection, Tinea capitis, Tinea infection, Tooth abscess, Tooth infection, Vaginal infection, Viraemia, Viral upper respiratory tract infection, Vulvitis, Vulvovaginal mycotic infection, Wound infection

Injury, poisoning and procedural complications: Animal scratch, Arthropod sting, Concussion, Confusion postoperative, Contusion, Exposure to toxic agent, Fibula fracture, Head injury, Heat exhaustion, Humerus fracture, Joint ligament rupture, Limb injury, Lower limb fracture, Medical device discomfort, Mouth injury, Multiple fractures, Muscle strain, Soft tissue injury, Splinter, Sunburn, Superficial injury of eye, Thermal burn, Tibia fracture, Vaccination complication, Wound

Investigations: Blood corticotrophin decreased, Blood cortisol decreased, Blood gonadotrophin decreased, Blood iron decreased, Blood sodium decreased, Catheterisation cardiac, Diagnostic procedure, Glucose tolerance test abnormal, Glucose urine, Head circumference abnormal, Hepatic enzyme increased, Iodine uptake abnormal, Lipids increased, Platelet count decreased, Transaminases increased, Weight decreased, White blood cell count decreased, White blood cells urine positive

Metabolism and nutrition disorders: Anorexia, Decreased appetite, Dehydration, Fluid overload
Fluid retention, Glucose tolerance impaired, Hypercholesterolaemia, Hyperinsulinaemia,
Hyponatraemia, Increased appetite, Polydipsia, Weight gain poor

Musculoskeletal and connective tissue disorders: Arthropathy, Chondropathy
Flank pain, Groin pain, Growth retardation, Joint effusion, Joint range of motion decreased, Juvenile
arthritis, Limb discomfort, Muscle contracture, Musculoskeletal pain, Myopathy, Neck pain,
Osteochondrosis, Rickets, Scoliosis, Shoulder pain, Spinal disorder, Temporomandibular joint syndrome,
Torticollis

Neoplasms benign, malignant and unspecified (incl. cysts and polyps): Astrocytoma, Glioma,
Metastases to spine, Neoplasm progression, Pinealoma, Pituitary tumour, Skin papilloma, Tumour flare

Nervous system disorders: Amnesia, Brain edema, Carpal tunnel syndrome, Cerebral atrophy, Clonus,
Coordination abnormal, Depressed level of consciousness, Encephalitis, Grand mal convulsion,
Hemiparesis, Hyperreflexia, Hypoaesthesia, Hipotonia, Intracranial pressure increased, Memory
impairment, Mental retardation severity unspecified, Migraine, Paraesthesia, Psychomotor
hyperactivity, Reflexes abnormal, Somnolence, Status epilepticus, Syncope, Tonic convulsion, Transient
ischemic attack, Tremor

Psychiatric disorders: Abnormal behaviour, Aggression, Attention deficit/hyperactivity disorder, Crying,
Emotional disorder, Personality change, Personality disorder, Restlessness, Stress, Tic

Renal and urinary disorders: Bladder spasm, Dysuria, Glycosuria, Hematuria, Leukocyturia, Nocturia,
Polyuria, Proteinuria

Reproductive system and breast disorders: Bilateral breast buds, Breast pain, Breast swelling,
Dysmenorrhea, Epididymal cyst, Female genital-digestive tract fistula, Menstruation irregular,
Metrorrhagia, Priapism, Pruritus genital, Testicular retraction, Testicular swelling, Vaginal erythema,
Vaginal hemorrhage, Vaginal ulceration

Respiratory, thoracic and mediastinal disorders: Dyspnoea, Dyspnoea exertional, Hyperventilation,
Pharyngeal erythema, Pharyngeal ulceration, Rhinalgia, Tonsillar hypertrophy

Skin and subcutaneous tissue disorders: Alopecia, Angioneurotic edema, Dandruff, Dermatitis
Dermatitis allergic, Dermatitis contact, Dry skin, Hair growth abnormal, Hyperhidrosis, Hyperkeratosis,
Keloid scar, Lipoatrophy, Lipodystrophy acquired, Neurodermatitis, Periorbital edema, Pigmentation
disorder, Rash erythematous, Rash maculo-papular, Rash papular, Rash pruritic, Scar, Seborrhea, Skin
discolouration, Skin hyperpigmentation, Skin hypopigmentation, Skin lesion

Social circumstances: Family stress

Surgical and medical procedures: Abscess drainage, Adenoidectomy, Adenotonsillectomy, Adhesiolysis,
Allergenic desensitisation procedure, Astrocytoma surgery, Brain tumour operation, Dental disorder
prophylaxis, Dental treatment, Drug therapy, Ear operation, Ear tube insertion, Explorative laparotomy,
Eye muscle tenotomy, Foot operation, Hernia repair, Hormone replacement therapy, Intra-aortic

balloon placement, Medical device implantation, Mineral supplementation, Mole excision, Orchidopexy, Surgery, Testicular operation, Urethral operation, Urethral repair, Wart excision

Vascular disorders: Flushing, Poor peripheral circulation

TS, CRF, SGA

No patients experienced AEs on treatment with a frequency of less than 1%.*

*CRF, TS, SGA: clinical trials included a subject number inferior or equal to 100 patients.

8.5. Post-Market Adverse Reactions

The majority of the adverse reactions of SAIZEN have been identified from the clinical trials and are summarized in the aforementioned studies.

The adverse reactions reported with marketed use of SAIZEN that are already mentioned in the clinical study tables are not repeated here unless there is a difference in frequency. These reactions have been identified during post-marketing surveillance and their exact frequency is unknown, except if specified.

General disorders and administration site conditions: localized lipoatrophy

Metabolism disorders: Hyperglycemia

Musculo-skeletal disorders: Avascular necrosis (osteonecrosis) of the femoral head including Legg-Calve-Perthes disease (Very rare).

Epiphysiolysis capitis femoris including slipped capital femoral epiphysis (SCFE).

Reproductive system and breast disorders: Gynaecomastia (uncommon)

9. Drug Interactions

9.2. Drug Interactions Overview

SAIZEN may interact with insulin, glucocorticoids, oral estrogens, thyroid hormones, and drugs known to be metabolized by CYP P450 or CYP3A4 hepatic enzymes (see Section [9.4 Drug-Drug Interactions](#)).

9.3. Drug-Behaviour Interactions

The interaction of SAIZEN with individual behavioural risks (e.g. cigarette smoking, cannabis use, and/or alcohol consumption) has not been studied.

9.4. Drug-Drug Interactions

Antidiabetic:

It is well documented that GH in the basal state increases plasma free fatty acid levels, even in the presence of insulin stimulation, thus counteracting the antilipolytic action of insulin. Given the anti-

insulin effects of GH, SAIZEN patients should be monitored for evidence of glucose intolerance in patients with diabetes mellitus or with a family history of diabetes mellitus. The care of diabetes in GH-replaced adults should follow standard guidelines, but intensified monitoring of metabolic control is advocated in the early phase of GH replacement of such patients. Patients with diabetes mellitus may require adjustment of their antidiabetic therapy.

Glucocorticoid effects on growth hormone response:

Concomitant glucocorticoid therapy may reduce the growth promoting effect of somatropin. If glucocorticoid replacement is required, the dose should be carefully adjusted to avoid either adrenal insufficiency or inhibition of growth promoting effects.

Growth hormone affecting the metabolism of glucocorticoids:

Several clinical studies report an impact of GH administration on glucocorticoid secretion, even if its mode of action remains unclear. There is evidence to suggest that GH lowers the levels of cortisol binding globulin and increases the net conversion of cortisol to cortisone, thus reducing bioactivity of glucocorticoids during GH replacement.

Initiation of GH replacement may unmask secondary adrenal insufficiency in some patients by reducing the activity of 11 β -hydroxysteroid dehydrogenase, type 1 (11 β -HSD1), an enzyme converting inactive cortisone to cortisol. Initiation of somatropin in patients receiving glucocorticoid replacement therapy may lead to manifestation of cortisol deficiency. Adjustment of glucocorticoid dose may be required. In a patient with central adrenal failure, initiation of GH treatment may require an increase in hydrocortisone dose. Careful monitoring of patients' symptoms such as weight, appetite, and mood are required to assess the need for glucocorticoid dose modification.

Growth hormone as inducer of cytochrome P450 3A4:

Published in vitro data indicate that growth hormone may be an inducer of CYP3A4, a cytochrome P450 enzyme involved in glucocorticoid catabolism. Therefore, growth hormone therapy may both unmask unsuspected adrenocorticotrophic hormone (ACTH) deficiencies and negate low replacement glucocorticoid doses used in secondary adrenal insufficiency (AI) by decreasing the availability of cortisol. Patients starting growth hormone therapy may require adjustments in their glucocorticoid replacement doses, and stress doses. Caution is recommended when administering SAIZEN with compounds that are metabolized by the CYP450 or CYP3A4 liver enzymes (e.g., corticosteroids, sex steroids, anticonvulsants, cyclosporine and others). When SAIZEN is administered in combination with drugs known to be metabolized by CYP 450 or CYP3A4 hepatic enzymes, it is advisable to monitor clinical effectiveness of such drugs.

Oral Estrogen:

Because oral estrogens may reduce the serum IGF-1 response to somatropin treatment, girls and women receiving oral estrogen replacement may require greater somatropin dosages. However, the maximum recommended weekly dose should not be exceeded.

Thyroid hormones:

Growth hormone administration in healthy normal adult subjects acutely increases serum T3, leading to reciprocal decreases in Free Thyroxine (FT4) and Thyroid-Stimulating Hormone (TSH) with a consequent increase in the T3 to the T4 ratio. This GH effect may as such unmask incipient hypothyroidism.

Therefore, the importance of evaluating thyroid function in GHD children prior to commencing therapy is emphasized.

9.5. Drug-Food Interactions

Interactions with food have not been established.

9.6. Drug-Herb Interactions

Interactions with herbal products have not been established.

9.7. Drug-Laboratory Test Interactions

Interactions with laboratory tests have not been established.

10. Clinical Pharmacology

10.1. Mechanism of Action

Human growth hormone is physiologically released in response to pulses of the hypothalamic growth hormone releasing factor (GRF) at night during sleep. Endogenously liberated hGH has a short half-life. At least part of the actual growth-promoting effect takes place through stimulation of the production of insulin-like growth factors or somatomedins, a group of peptide hormones with a long half-life which, via a negative feedback mechanism, influence the synthesis and release of hGH.

Somatropin is a polypeptide hormone consisting of 191 amino acid residues and its structure is identical to that of growth hormone extracted from human pituitary glands. It is produced by recombinant (rDNA) technology in a mammalian cell expression system. Somatropin is also therapeutically equivalent to human growth hormone of pituitary origin.

SAIZEN provides an exogenous supply of human growth hormone for those patients lacking the ability to produce adequate endogenous supplies.

Linear growth: Somatropin stimulates linear growth in patients with pituitary growth hormone deficiency, Turner syndrome and chronic renal failure. Treatment of these patients with SAIZEN results in increased growth rates and IGF-1 levels similar to those seen for children treated with growth hormone of pituitary origin.

Skeletal growth: The measurable increase in growth (body length) after somatropin treatment results from its effect on cartilaginous growth areas of the long bones. It is known that somatropin's effect is mediated by a sulfation factor, IGF-1, which permits the incorporation of sulfate into cartilage. IGF-1 is present in low concentration in the serum of growth hormone deficient patients and increases during somatropin therapy.

Cell growth: Somatropin brings about cellular growth as demonstrated by an increase in the muscular, visceral and red cell mass. In muscle tissue, the increase in mass is associated with a corresponding increase in both number and dimension of muscular fibre cells.

10.2. Pharmacodynamics

A single-blind, randomized, placebo-controlled study in 16 normal-weight, healthy male volunteers was conducted to evaluate the pharmacology of SAIZEN. SAIZEN 0.067 mg/kg or placebo was administered over a period of 14 days. There was no difference between the treatment groups regarding blood pressure, headache, queasiness, and itching and local side effects (redness, swelling, soreness, and sensitivity to touch which occurred in both groups) which was caused by the high osmolality of the administered solution.

A single-blind, placebo-controlled, partly randomized cross-over phase I study was performed in six healthy male volunteers, under standard diet. Single doses of r-hGH were administered: 5 mg as a 6-hour iv infusion, 6 mg and 20 mg by sc route.

During the iv infusion, GH concentrations rapidly reached a plateau followed by a rapid decay, while the GH concentration curve rose and decreased smoothly after the sc injection, the absorption representing the rate limiting step. A moderate degree of non-linearity was observed between the 6 and 20 mg sc doses, the latter displaying a greater bioavailability and a prolonged mean absorption time. The 3 doses of r-hGH elicited similar side effects, though in varying magnitudes (transient weight gain, a rise in IGF-1, IGFBP-3, non-esterified fatty acids and glycerol, and a decrease in blood urea, urinary nitrogen, sodium and potassium excretions). These effects were more striking after the 6 mg sc dose than after the 5 mg iv infusion, although both doses produced similar AUCs of GH. The biological changes induced by the 20 mg dose were not much higher than after the 6 mg dose, suggesting a relative saturation of the effects of r-hGH at high doses. This was not the case for the glycemia and the C-peptide urinary excretion, which were significantly elevated only after the 20 mg dose.

Table 8 - Effects of 5 mg as a 6-hour iv infusion, 6 mg and 20 mg by sc route

Variable	Treatment vs time interaction	Effect of 5mg iv vs placebo	Effect of 6mg sc vs placebo	Effect of 20 mg sc vs placebo	Effect of 6mg sc vs 20 mg sc
Weight gain	0.08	24h-72 h	24h-96h	24h-72h	72h-96h
IGF-1	<0.0001	8h-48h	8h-72h	8h-96h	16h-48h
IGFBP-3	0.014	12h-48h	16h-48h	16h-72h	72h-96h
NEFA	0.0002	4h-12h	4h-12h	4h-24h	--
Glycerol	0.01	--	24h	8h, 24h, 48h	8h
Blood urea	<0.0001	8h-24h	8h-36h	12h-48h	48h
Urinary nitrogen	0.05	8h-24h	4h-24h	12h-24h	4h-8h
Urinary sodium	0.006	4h-24h	4h-24h	0h-96h	4h-12h
Urinary potassium	<0.0001	4h-24h	4h-24h	0h-96h	4h-12h

The above table lists the time periods (hours) over which the variable differs significantly ($p < 0.5$) from the placebo (or 6 mg sc). The values are calculated from a 3-way analysis of variance for repeated measures and randomized block design.

Refer to [14.1 Clinical Trials by Indication, Adult Indications, Cardiac function](#) subsection.

Apart from its growth-promoting effect, growth hormone influences the carbohydrate, the lipid, and the protein metabolism.

Carbohydrate Metabolism:

Somatropin has an effect on carbohydrate metabolism. The diabetogenic effect of somatropin is well-known in clinical medicine. Acromegalic patients often suffer from diabetes mellitus while hypopituitary children experience hypoglycemia. In healthy patients, very large doses of somatropin may interfere with glucose tolerance. A simultaneous increase in the plasma insulin level is observed upon somatropin administration.

The diabetogenic activity of somatropin is perhaps due to several concomitant factors:

- Reduced transport of glucose into peripheral tissues.
- Increased release of glucose from the liver.
- Reduced concentration of insulin at the muscular level.
- Reduced glycolysis from the block of the enzyme triose phosphate dehydrogenase, mediated by non-esterified fatty acids.

By inhibiting glucose uptake in the tissue, hGH has an anti-insulinogenic effect. At the same time, hGH increases the release of pancreatic insulin. SAIZEN (0.067 mg/kg) administration to 8 male volunteers every 48 hours over a period of 14 days resulted in a greater increase in glucose and a higher insulin secretion during the OGTT than before treatment whereas a single intramuscular or subcutaneous injection (1.33 mg SOMATROPIN/m²) had no significant effect on the blood sugar and C-peptide levels within the first 4 hours. The mean basal insulin level was increased after 4 hours (p=0.0738). Results of these two studies are summarized:

Table 9 - Glucose and Insulin levels in male patients receiving SC injections of r-hGH every 48hr for 14 days [mean (se)]

	Occasion	Glucose (mg/dL)		Insulin (IU/mL)	
		HGH-rDNA	Placebo	HGH-rDNA	Placebo
0 min	premedication	80.19 (1.70)	80.86 (3.92)	6.35 (1.27)	9.28 (2.26)
	stage II Day 7	92.58 (3.23)	84.02 (1.67)	17.28 (2.96)	9.26 (1.52)
	stage II Day 15	89.87 (2.98)	80.86 (2.00)	16.03 (1.54)	10.26 (1.48)
30 min	premedication	126.82 (6.08)	130.64 (9.74)	83.16 (15.82)	91.14 (23.64)
	stage II Day 7	168.49(12.50)	151.59 (8.38)	100.64 (28.45)	97.7 (21.96)
	stage II Day 15	168.71(12.00)	144.61 (8.69)	124.46 (20.15)	102.87 (26.19)
60 min	premedication	88.3 (6.67)	98.66 (11.38)	47.78 (11.7)	49.81 (8.36)
	stage II Day 7	196.64 (11.75)	145.74 (14.08)	176 (33.26)	123.14 (36.59)
	stage II Day 15	173.67 (18.31)	133.8 (12.39)	169.99 (23.78)	114.45 (17.08)
90 min	premedication	73.21 (3.94)	89.2 (10.30)	22.45 (5.31)	42.42 (9.99)
	stage II Day 7	187.86 (15.30)	113.3 (8.15)	216.91 (26.80)	94.61 (26.57)
	stage II Day 15	152.94 (15.30)	110.37 (8.41)	154.5 (30.96)	101.38 (17.60)
120 min	premedication	65.10 (4.18)	77.94 (10.44)	11.71 (2.23)	32.81 (10.58)
	stage II Day 7	149.12 (12.99)	105.19 (7.62)	177.94 (33.99))	81.62 (24.20)
	stage II Day 15	118.03 (10.97)	88.75 (8.37)	138.92 (24.24)	53.41 (6.57)
150 min	premedication	67.12 (6.26)	68.25 (6.74)	6.4 (1.95)	13 (4.91)
	stage II Day 7	116.9 (13.81)	88.97 (7.10)	95.51 (21.69)	38.17 (8.49)
	stage II Day 15	86.5 (10.22)	69.6 (8.77)	76.45 (21.10)	34.06 (8.93)
180 min	premedication	70.5 (5.18)	72.53 (6.65)	4.86 (0.91)	9.19 (2.25)
	stage II Day 7	98.43 (11.26)	76.36 (4.50)	59.04 (17.99)	22.89 (5.08)
	stage II Day 15	70.05 (5.62)	77.04 (2.90)	29.45 (7.50)	15.5 (2.01)

Table 10 - Glucose, C-peptide and Insulin levels in patients receiving a single SC or IM injection of r-hGH [mean (sd)]

Time (h)	male after IM injection	female after IM injection	male after SC injection	female after SC injection
Glucose (m/dL)				
0	85.33 (5.57)	84.67 (4.80)	84.20 (4.82)	82.33 (4.80)
1	83.17 (5.12)	82.17 (8.66)	84.80 (5.54)	80.67 (4.84)
2	83.00 (8.53)	82.33 (7.92)	81.80 (6.42)	81.50 (5.32)
3	87.50 (7.37)	86.83 (5.19)	86.00 (6.28)	79.33 (7.42)

4	91.40 (6.02)	84.67 (7.37)	83.40 (4.28)	81.67 (8.52)
C-Peptide (IU/mL)				
0	83.17 (26.55)	66.83 (6.37)	65.40 (18.31)	68.17 (13.75)
1	62.83 (16.83)	61.83 (8.47)	61.20 (15.27)	65.83 (17.75)
2	61.50 (20.50)	63.17 (5.98)	70.50 (18.31)	65.83 (11.13)
3	60.83 (22.05)	64.50 (13.38)	68.60 (15.27)	61.00 (13.36)
4	71.00 (20.25)	68.00 (10.24)	68.80 (17.48)	65.00 (14.21)
Insulin (IU/mL)				
0	12.23 (2.77)	12.62 (2.12)	11.00 (2.61)	12.44 (2.20)
1	10.33 (2.95)	11.08 (1.86)	11.56 (3.34)	11.32 (1.39)
2	10.25 (4.09)	11.80 (1.67)	9.78 (2.40)	11.22(3.37)
3	10.92 (2.41)	12.32(3.95)	11.20 (2.61)	9.66 (2.34)
4	12.94 (5.97)	13.36 (1.86)	14.74 (6.50)	11.30 (2.03)

Connective tissue metabolism:

Connective tissue metabolism is affected by somatropin's ability to stimulate the synthesis of chondroitin sulfate and collagen as well as the urinary excretion of hydroxyproline.

Lipid Metabolism:

Lipid metabolism is also affected by somatropin. This occurs when intracellular lipolysis is stimulated, thus increasing the plasma concentration of free fatty acids and stimulating the oxidation of fatty acids. In the diabetic patient, somatropin has been shown to accentuate ketogenesis.

Due to an activation of hormone-sensitive lipases in the fatty tissue, the parenteral application of hGH leads to an increase in FFA in the serum with a maximum between 3 - 6 hours, which lasts for several hours. Following subcutaneous and intramuscular injections of SAIZEN (1.33 mg Somatropin/m²), a significant increase in the mean FFA levels could be demonstrated after 4 hours. A drop of serum cholesterol after administration of high doses of SAIZEN was also found in 8 male volunteers.

Mineral metabolism:

Somatropin affects mineral metabolism by inducing the net retention of phosphorus and potassium and to a lesser degree sodium. Somatropin induces the increased intestinal absorption of calcium and the increased renal tubular reabsorption of phosphorus with increased serum and inorganic phosphate. Increased serum alkaline phosphatase may also be observed during somatropin therapy.

Protein Metabolism:

Somatropin has an effect on protein metabolism. Somatropin is an anabolic agent that stimulates intracellular transport of amino acids, net retention of nitrogen and protein synthesis which can be quantified by observing the decline in urinary nitrogen excretion and BUN.

In hGH deficient children, the administration of hGH produces a marked retention of nitrogen. This was the reason for Prader et. al. to develop the nitrogen retention test for the differential diagnosis of short stature. Such metabolic tests have subsequently been simplified and refined using the stable isotope ¹⁵N. Recently, GH-deficient patients have been shown to respond to a recombinant hGH preparation in the same way as to pit-hGH. The effect on protein metabolism is closely connected with the growth-

promoting effect. The growing organism requires more protein, and this requirement is then met by stimulation of protein synthesis. SAIZEN administered subcutaneously to 8 male volunteers every 48 hours over a period of 14 days (0.2 IU/kg [approximately 0.067 mg/kg]) resulted in a statistically significant drop of the mean urea blood level from 34.4 ± 2.5 mg/dL to 24.8 ± 1.5 mg/dL.

Insulin-like Growth Factor-I (IGF-1):

Reduced IGF-1 serum concentrations are often found in children with hypothalamic-hypophyseal hGH deficiency. Recent investigations show that plasma IGF-1 levels do not (although they generally increase on hGH treatment) directly correlate with the therapeutic success of hGH, because other factors (nutrition, binding protein concentration) modulate the plasma concentrations, and because IGF-1 formation in the tissues is not directly reflected in the plasma levels. In addition, some animal experiments show that not all growth-promoting effects are necessarily mediated through IGF-1.

SAIZEN (1.33 mg Somatropin/m²) administration to 12 healthy adults (6 males and 6 females), showed increased mean serum concentrations of IGF-1 within 24 hours. SAIZEN (0.067 mg/kg) administration to 8 healthy male volunteers every 48 hours over a period of 14 days, resulted in an increased mean serum concentration.

Contrary to earlier reports, other authors have found no correlation between the growth velocity and short-term or long-term changes in the IGF-1 serum levels. In the U.S. multicentre study for SAIZEN, the mean IGF-1 serum level (n = 50) rose from a pre-treatment level of 8.12 ± 6.36 nmol/L to 16.95 ± 9.82 nmol/L within 12 months of treatment, whereas the other SAIZEN multicentre studies confirmed that IGF-1 cannot be used as a parameter for evaluating the efficacy of growth hormone. The reason for this might be that IGF-1 is produced by different tissues and that the effect on the target cells is carried out in a paracrine or autocrine manner. Furthermore, IGF-1 (as determined by RIA), is only one of the growth factors responsible for the stimulation of growth.

10.3. Pharmacokinetics

The pharmacokinetics of SAIZEN are linear at least up to doses of 8 IU (2.67 mg). At higher doses (60 IU/20 mg) some degree of non-linearity cannot be ruled out, however with no clinical relevance.

Following IV administration in healthy volunteers the volume of distribution at steady-state is around 7 L, total metabolic clearance is around 15 L/h while the renal clearance is negligible, and the drug exhibits an elimination half-life of 20 to 35 min.

Following single-dose SC and IM administration of SAIZEN, the apparent terminal half-life is much longer, around 1 to 6 hours (median 2.7 hours). This is due to a rate limiting absorption process.


Maximum serum growth hormone (GH) concentrations following injection are reached after approximately 4 hours (range 2 to 7 hours) and serum GH levels return to baseline within 24 hours, indicating that no accumulation of injected GH will occur during repeated administrations.

The absolute bioavailability of both routes is 70-90 %.

Table 11 - Summary of Somatropin's Pharmaceutical Parameters

	Distribution Steady State	Metabolic Clearance	Terminal half-life (range)	Time to C _{max}	Bioavailability
Single dose	7L	15L/h	4 hours (1 to 6 hours)	4 hours	70-90%

An open, non-comparative study was conducted in 12 healthy adults (6 males and 6 females). SAIZEN (1.33 mg Somatropin/m²) was injected intramuscularly and subcutaneously in equal doses, with at least one week between each injection. To determine the C_{max} and T_{max}, blood samples were taken at 1, 2, 3, 4, 6, 9, 12, 15 (3 mg only) and 24 hours; hGH was determined by specific radioimmunoassay. While the mean peak serum concentration of circulating hGH (36.9 ng/mL) occurred 3 hours after intramuscular injection, subcutaneous injections resulted in a more sustained elevation at lower levels, reaching a mean peak serum concentration between 4 hours (mean 16.4 ng/mL) and 6 hours (mean 16.3 ng/mL). The areas under the curves (AUCs) were very similar after subcutaneous and intramuscular injections. Comparable results were obtained after subcutaneous administration of the same amount of SAIZEN in a higher concentrated solution (3.33 mg/mL instead of 1.33 mg/mL). The T_{max} values reported for intramuscular injection of pit-hGH in the literature are between 2 and 4 hours which correlates well with the values found for SAIZEN.

Figure 1 – hGH serum concentrations (mean ± SD) and resulting curves

Animal Pharmacology

Two pharmacokinetic studies in primates (*Cynomolgus* monkeys) were conducted with SAIZEN, consisting of one single dose study using two routes of administration, subcutaneous and intravenous, and one repeat dose study by subcutaneous administration.

In the single dose study, the animals were dosed with 0.16 mg (in 0.5 mL)/kg SAIZEN (r-hGH) and 0.5 IU (in 0.5 mL)/kg Asellacrin [natural hGH or pituitary extraction] for comparison, and the pharmacokinetic profiles were compared. Eight monkeys were treated in a cross-over design with all four treatments with at least 1 week between subsequent treatments. Blood samples were drawn from a forelimb vein of the fasted animals immediately before and up to 48 hours after administration.

For both SAIZEN and Asellacrin, the half-life was significantly longer after subcutaneous administration than after intravenous administration. The relative bioavailability of SAIZEN was not significantly different from that for Asellacrin.

Table 12 - Mean Half-Life ($T_{1/2}$ Values)

Subcutaneous vs Intravenous	
<u>Subcutaneous $T_{1/2}$ (hours)</u>	<u>Intravenous $T_{1/2}$ (hours)</u>
r-hGH 5.1	r-hGH 3.2
n-hGH 4.7	n-hGH 2.4

In the repeat dose study, the pharmacokinetic profiles of SAIZEN and Asellacrin were compared by subcutaneous administration alone. Animals were treated at 24 hour intervals for 8 consecutive days, with either 0.16 mg/kg of SAIZEN or 0.5 IU/kg of Asellacrin, after being fasted for 16 hours. Blood samples were taken from a forelimb vein of each animal immediately prior to and 1 hour after each administration (meantime for highest serum concentration after administration as shown in the single dose pharmacokinetic study), on study days 1-7, and prior to and up to 48 hours after treatment on day 8.

Neither increasing nor decreasing trends could be detected in the serum concentration values obtained prior to or one hour after each daily administration. No accumulation nor induction phenomena occurred in this study. No significant difference for the elimination half life was detected for Asellacrin when compared with the single dose study. However, a significant difference (near to the 5% significance level) was seen for SAIZEN. It was concluded that the serum kinetic profile of hGH is not significantly altered by repeated administration. No significant difference was seen between the values of the main pharmacokinetic parameters calculated after the last administration in monkeys.

Special Populations and Conditions

SAIZEN is indicated for children with growth failure due to chronic renal failure (see [1 Indications](#)). No other studies have been conducted with special populations and conditions.

There have been reports of fatalities associated with the use of growth hormone in pediatric patients with Prader-Willi syndrome who have one or more of the following risk factors: severe obesity, history of respiratory impairment or sleep apnea or unidentified [i.e. previously undiagnosed/mildly symptomatic] respiratory infections (see [2 Contraindications](#); [7 Warnings and Precautions, Congenital Disorders](#)). In adults with risk factors for insulin resistance or glucose intolerance, such as obesity, a family history of diabetes mellitus and those on high dose corticosteroid therapy, growth hormone therapy may induce Type II Diabetes Mellitus if the insulin secretory capacity is impaired (see [7 Warnings and Precautions, Endocrine and Metabolism](#)).

10.4. Immunogenicity

All therapeutic proteins have the potential for immunogenicity.

The detection of antibody formation is highly dependent on the sensitivity and specificity of the assay. Additionally, the observed incidence of antibody (including neutralizing antibody) positivity in an assay may be influenced by several factors including assay methodology, sample handling, timing of sample

collection, concomitant medications, and underlying disease. For these reasons, comparison of incidence of antibodies in the studies described below with the incidences of antibodies in other studies or to other products may be misleading.

For more detailed discussion of the clinical effect(s) and pertinent clinical recommendations, see [7 Warnings and Precautions, Immune](#), Antibody Production.

11. Storage, Stability, and Disposal

SAIZEN 5 mg/vial:

Store SAIZEN lyophilized product at room temperature (not to exceed 25°C) and in original package to protect from light.

Do not use SAIZEN after the expiry date shown on label.

Stability of Solution and Storage

SAIZEN 5 mg/vial:

When reconstituted with 1 mL to 3 mL Bacteriostatic Water for Injection, USP, store the reconstituted solution at 2-8 °C for up to 14 days.

When reconstituted with Water for Injection, USP, the reconstituted solution should be administered immediately (within 3 hours). Any unused solution should be discarded.

SAIZEN 6 mg (5.83 mg/mL), 12 mg (8 mg/mL), 20 mg (8 mg/mL) cartridges:

Store cartridges in a refrigerator (2-8°C) in the original package to protect from light. When using the easypod electromechanical auto-injector or the Aluetta pen injector, the cartridge is kept in the device and the device must be stored in the refrigerator.

Do not freeze. After first injection, the cartridges must be stored at 2-8°C for a period of up to 28 days, of which no more than 7 days may be outside of the refrigerator at room temperature (not to exceed 25°C). Cartridges must be discarded after 28 days, or if the total period outside of the refrigerator exceeds 7 days.

Do not use SAIZEN after the expiry date, which is stated on the cartridge.

Any unused product or waste material should be disposed of in accordance with local requirements.

Part 2: Scientific Information

13. Pharmaceutical Information

Drug Substance

Non-proprietary name of the drug substance(s): somatropin for injection

Chemical name: recombinant human Growth Hormone for Injection (r-hGH)

Molecular formula and molecular mass: C₉₉₀ H₁₅₂₈ N₂₆₂ O₃₀₀ S₇

Structure (for biologics)/Structural formula:

-PHE-PRO-THR-ILE-PRO-LEU-SER-ARG-LEU-PHE-ASP
-ASN-ALA-MET-LEU-ARG-ALA-HIS-ARG-LEU-HIS-GLN-LEU-ALA
-PHE-ASP-THR-TYR-GLN-GLU-PHE-GLU-GLU-ALA-TYR-ILE-PRO
-LYS-GLU-GLN-LYS-TYR-SER-PHE-LEU-GLN-ASN-PRO-GLN-THR
-SER-LEU-CYS-PHE-SER-GLU-SER-ILE-PRO-THR-PRO-SER-ASN
-ARG-GLU-GLU-THR-GLN-GLN-LYS-SER-ASN-LEU-GLU-LEU-LEU
-ARG-ILE-SER-LEU-LEU-LEU-ILE-GLN-SER-TRP-LEU-GLU-PRO
-VAL-GLN-PHE-LEU-ARG-SER-VAL-PHE-ALA-ASN-SER-LEU-VAL
-TYR-GLY-ALA-SER-ASP-SER-ASN-VAL-TYR-ASP-LEU-LEU-LYS
-ASP-LEU-GLU-GLU-GLY-ILE-GLN-THR-LEU-MET-GLY-ARG-LEU
-GLU-ASP-GLY-SER-PRO-ARG-THR-GLY-GLN-ILE-PHE-LYS-GLN
-THR-TYR-SER-LYS-PHE-ASP-THR-ASN-SER-HIS-ASN-ASP-ASP
-ALA-LEU-LEU-LYS-ASN-TYR-GLY-LEU-LEU-TYR-CYS-PHE-ARG
-LYS-ASP-MET-ASP-LYS-VAL-GLU-THR-PHE-LEU-ARG-ILE-VAL
-GLN-CYS-ARG-SER-VAL-GLU-GLY-SER-CYS-GLY-PHE

Physicochemical properties: Molecular weight = 22, 125 daltons

Pharmaceutical standard: Professed/House

Product Characteristics:

Somatropin is a polypeptide hormone consisting of 191 amino acid residues and its structure is identical to that of growth hormone extracted from human pituitary glands. A large loop is formed by a disulfide bond between Cys⁵³ and Cys¹⁶⁵. A second, smaller loop is formed by a disulfide bond near the carboxyl-terminal between Cys¹⁸² and Cys¹⁸⁹. The solution is a slightly opalescent liquid.

The biological activity of growth hormone is approximately 3.0 international units/1mg.

14. Clinical Trials

SAIZEN (somatropin for injection) is a polypeptide hormone of recombinant DNA origin which is composed of 191 amino acid residues in the identical sequence and configuration as human pituitary growth hormone. It is indicated as classical endocrine replacement therapy for the long-term treatment of patients with growth failure due to inadequate secretion of normal endogenous growth hormone. In

vitro, preclinical and clinical studies have demonstrated that SAIZEN is therapeutically equivalent to human growth hormone (hGH) of pituitary origin.

14.1. Clinical Trials by Indication

Pediatric Indications

Inadequate Endogenous Growth Hormone Secretion

Trial Design and Study Demographics

Table 13 – Summary of Patient Demographics for Clinical Trials in Pediatric GHD

Study #	Study design	Dosage, route of administration and duration	Study subjects (n)	Age (range)	Sex (m, f)
1. GF 2078 2. GF 2376 3. GF 2386 4. GF 2415 5. GF 2537	Each study: open-label, multicentre, Phase III	0.20 mg/kg/week, 3-6 injections Subcutaneous administration, 2 years	304	0.0-19.0	1. 50, 19. 2. 73, 28 3. *. 4. 22, 17 5. Naïve patients 19 (12, 7). Transfer patients: 10*.

*: information missing in the clinical study report

Efficacy and safety of SAIZEN have been studied in five pivotal studies using pre-treatment growth measurements compared with treatment growth measured as a method of control.

Of the total patients enrolled in these studies, 70.3% with at least 12 months treatment have been analyzed for efficacy. All patients were prepubertal or pubertal children with classic growth hormone deficiency with or without previous growth hormone treatment. The patients in the different studies treated with SAIZEN were assessed for the occurrence of adverse events and laboratory abnormalities and were tested regularly for the presence of antibodies against hGH and against proteins of the host cells (C-127 mouse cells).

Study Results

In addition to the treatment related adverse events reported above, two patients developed anti-hGH antibodies. In both cases, the antibodies did not have any growth inhibiting effect. None of the patients developed antibodies to host cell protein. Three transfer patients who had anti-hGH antibodies prior to treatment became negative within 6 months of treatment with SAIZEN. Hypothyroidism was seen in several patients. One patient died of recurrent craniopharyngioma and one patient experienced lipotrophy.

The effectiveness of growth hormone treatment on growth was assessed primarily by changes in height velocity:

- Height velocity in cm/year, as a change from baseline.
- Height velocity as a change in standard deviation with reference to mean value for chronological age [Standard Deviation Score for Chronological Age (SDS CA)].
- Height velocity as a change in the standard deviation with reference to mean values for bone age [Standard Deviation Score for Bone Age (SDS BA)].

In contrast to the absolute data in cm/year, the calculation in SDS takes the different normal height velocities of different age groups into account.

In the above multicentre studies, 54.7% of the subjects were naive patients treated for 12 months. The efficacy results are summarized in [Table 14](#).

Table 14 - Height Velocities in Naive Patients Treated with SAIZEN for 12 Months in Comparison with 12 Month Literature Data for pit-hGH

Growth velocity	SAIZEN		pit-hGH	
	Before	During 12 months	Before	During 12 months
cm/year				
Germany	3.60 ± 1.22 (n=27)	9.54 ± 2.76 (n=27)		
USA	3.49 ± 1.10 (n=50)	8.56 ± 1.65 (n=50)		
Italy	3.22 ± 1.39 (n=26)	8.54 ± 2.45 (n=26)	3.68 ± 1.07 (n=25)	7.66 ± 2.37 (n=36)
United Kingdom	3.77 ± 1.75 (n=12)	10.02 ± 2.08 (n=12)		
France	3.88 ± 1.07 (n=25)	8.03 ± 1.59 (n=25)		
SDS CA				
Germany	-2.69 ± 1.28 (n=26)	+3.44±2.81 (n=26)	-2.45 ± 0.84 (n=?)	+2.29 ± 2.49 (n=36)
USA	-2.82 ± 1.27 (n=50)	+3.26±2.38 (n=50)		
SDS BA				
Germany	-2.92 ± 1.14 (n=25)	+2.19 ± 1.25 (n=25)	-3.0 ± 0.99 (n=22)	+0.71 ± 1.93 (n=35)
USA	-3.08 ± 1.09 (n=50)	+2.48 ± 2.41 (n=50)		

Turner Syndrome

Trial Design and Study Demographics

Table 15 – Summary of Patient Demographics for Clinical Trials in Turner Syndrome

Study #	Study design	Dosage, route of administration and duration	Study subjects (n)	Age (range)	Sex
GF 3152	Open-label phase III randomised	<u>XO group</u> : r-hGH 18 IU/m ² /week, 1 year, then r-hGH 24 IU/m ² /week, 1 year. <u>XM group</u> : r-hGH 18 IU/m ² /week, 2 years and oxandrolone 0.100 mg/kg/day the 1 st year, then oxandrolone 0.05 mg/kg/day the 2 nd year. Subcutaneous administration.	0/91	5.7-15.4	0, 91
GF 5413 (continuation of study GF 3152)	Open-label phase III randomised	r-hGH 24 IU/m ² /week, with or without oxandrolone 0.05 mg/kg/day, up to a total of 6 years over studies GF 3152 & 5413. Subcutaneous administration	0/91	5.7-15.4	0, 91

Turner syndrome is caused by the apparent complete or partial absence of one of the X chromosomes or by other chromosomal abnormalities involving the second X chromosome. It occurs with an incidence of about 1 in 1000 neonates. The majority of Turner syndrome patients experience poor growth and gonadal dysfunction.

Short stature is nearly a constant clinical feature of girls with Turner syndrome. Although there is considerable variation in phenotypic expression, the observed height velocities and final height of girls with TS are usually significantly lower than the normal average. The effectiveness of any therapeutic agent in the treatment of short stature in girls with TS is judged by the change in height velocity and the effect on predicted and/or final height.

The study described here is a pivotal, Phase III trial in patients with Turner syndrome. The purpose of this study was to assess the long-term efficacy as well as the safety of SAIZEN alone and in combination with oxandrolone in the treatment of growth retarded girls with Turner syndrome.

This was an open, comparative, randomised multicentre study where a total of 91 girls (aged 10.3 ± 2.3 years) with Turner syndrome were randomly allocated to receive either SAIZEN alone (XO group) or SAIZEN in combination with oxandrolone (XM group). The study medication dosing regimen is presented in [Table 16](#).

Table 16 - Dosage Regimen

		XO (initial n = 47)	XM (initial n = 44)
1 st Year	SAIZEN	18 IU/m ² /week (~0.029mg/kg/day)	18 IU/m ² /week (~0.029 mg/kg/day)
	Oxandrolone	-	0.1 mg/kg/day
2 nd Year	SAIZEN	24 IU/m ² /week (~0.038 mg/kg/day)	18 IU/m ² /week (~0.029 mg/kg/day)
	Oxandrolone	-	0.05 mg/kg/day

		XO (n=21)	XOM (n=26)	XM (n=33)	XMO (n=11)
3 rd Year and subsequent	SAIZEN	24 IU/m ² /week (~0.038mg/kg/day)	24 IU/m ² /week (~0.038mg/kg/day)	24IU/m ² /week (~0.038mg/kg/day)	24 IU/m ² /week (~0.038 mg/kg/day)
	Oxandrolone	-	0.05 mg/kg/day	0.05 mg/kg/day	-

After the second year, oxandrolone 0.05 mg/kg/day was also recommended for the XO group if HV was less than +2SD above the respective mean HV for age of girls with TS (group XOM). If subjects of the XM group had to stop oxandrolone, but continue SAIZEN, the dose of SAIZEN was increased to 24 IU/m²/week (group XMO).

Study Results

Changes in Height Velocity (cm/year)

The mean HV increased significantly over baseline in both groups (XO and XM) during the first 12 months of treatment. The difference between the mean gains in HV of the two groups, 2.4 ± 1.3 (XO) vs. 4.6 ± 1.8 (XM) cm/year, was statistically significant ($P < 0.0001$). During the second year of treatment, the mean HV was maintained at a higher level compared with baseline in both groups. The TS girls in the XO group grew at a rate of 5.5 ± 1.1 cm/year ($+1.5 \pm 1.1$ cm/year over baseline) significantly more than before treatment ($P < 0.0001$). In the XM group, a mean HV of 6.5 ± 1.4 cm/year was achieved with a $+2.4 \pm 1.9$ cm/year gain over baseline ($P < 0.0001$). The difference in HV between the XO versus XM groups over the first two years of treatment was also statistically significant ($P < 0.05$).

Height

Heights at the start of treatment showed a normal distribution around the 50th centile curve for a population of untreated girls with TS. After following 2 to 7.5 years of treatment, the majority of the TS patients were more than 1 SD above the mean for age, and had achieved a height greater than 150 cm. The best responses were achieved in the youngest patients.

Predicted Final Height

Changes in final height predictions (FHP) were analysed in the 35 TS girls who completed 6 years of treatment using the Bayley-Pinneau method based on bone age. The mean change in predicted adult height after 6 years ranged between 9 - 11 cm for the 4 groups. This suggests that final height will have improved by approximately 1.7 SD since the start of therapy.

Final Height

Twenty-six of the original 91 TS girls have reached final height defined as a height velocity = 0.5 cm/year over the last year of observation or treatment. For the group as a whole, the mean ($\pm$ SD) final height was 150.6 ± 5.5 cm. The results are consistent with those reported in the literature for other TS girls treated with GH. When normalized for age, the improvement for the group of patients who reached final height as a whole was 1.1 SD. It should be noted that the age at attainment of final height was 18.2 years compared to 15.7 years for those who had not yet reached final height and on average were treated for a shorter period of time, 4.9 versus 5.3 years.

Factors influencing response to treatment

In order to assess the influence of certain factors on the response to SAIZEN therapy, we examined the effect of age at start of therapy and the duration of therapy on the change in height standard deviation

score (HSDS). There was a highly significant inverse relationship between age at start and response to therapy ($r=-0.49$, $p<0.0001$), suggesting that the earlier the treatment is initiated, the better final height is likely to be. There was also a highly significant positive relationship between the duration of treatment and the improvement in HSDS ($r=0.44$, $p<0.0001$), suggesting that the final height will indeed be influenced by the duration of treatment.

In order to separate the effects of age at start and duration of treatment on the response to therapy, the effect of age at start on HSDS was analyzed by holding the duration of treatment constant. Highly significant negative correlations were again evident: $r=-0.54$ ($p<0.0001$) for the 56 girls treated for 5 years, and $r=-0.56$ ($p<0.0005$) for the 35 girls treated for 6 years.

Chronic Renal Failure

Trial Design and Study Demographics

Table 17 – Summary of Patient Demographics for Clinical Trial in CRF

Study #	Study design	Dosage, route of administration and duration	Study subjects (n)	Age (range)	Sex
GF 4941	Open-label, Phase III	28 IU/m ² , 3 years, then 36IU/m ² if unsatisfactory growth. Subcutaneous administration	81	1.7, 16.4	58, 23

A primary intent to treat analysis is not available for the published literature in this area. Growth failure is common and serious sequelae of chronic renal failure in childhood with profound consequences for a child's psychological development and social integration. Even when managed optimally, patients with CRF continue to lose height over time, both relative to their normally growing peers and in terms of final height.

The study described here is a pivotal, multicenter study where a total of 81 children with chronic renal failure (17 post-transplant, 27 on dialysis and 37 compensated CRF) were evaluated to assess the safety and efficacy of SAIZEN in the treatment of growth failure in children with chronic renal failure (CRF). SAIZEN was administered subcutaneously at a weekly dose of 0.35 mg/kg per week (28 IU/m²/ week) for the first 3 years of treatment.

Study Results

For the group as a whole, laboratory changes were similar to those reported in patients receiving GH for growth hormone deficiency. This suggests that the dose used in this study is, in effect, an appropriately physiological one for children with CRF.

Changes in Height Velocity (HV)

After 12 months: Of the 63 patients who were available for analysis, 59 (94%) experienced an increase over baseline in HV. Mean HV (SD) increased by 4.4 ± 4.0 cm/yr. ($p < 0.001$).

After 24 months: In the 44 children available for analysis, 39 (89%) experienced a sustained increase over baseline HV and the mean HV for this cohort was 7.5 ± 2.9 cm/yr., an increase of 3.0 ± 3.6 cm/yr. over baseline ($p < 0.001$).

Changes in Height Standard Deviation Score (H SDS)

After 12 months: Of the 63 children available for analysis, 55 (87%) experienced an increase over baseline in H SDS. The percentage of children with a normal H SDS increased from 1% (1 of 81) to 17% (11 of 63). For the group as a whole, H SDS increased by $+0.7 \pm 0.7$ ($p < 0.001$).

After 24 months: In the 44 children available for analysis, 38 (86%) experienced a sustained increase over baseline in H SDS. The percentage of children achieving a normal H SDS increased to 43% (19 of 44). The mean HV SDS remained significantly higher than at baseline and was greater than zero, consistent with ongoing catch-up growth.

Table 18 - Mean Height Velocity (cm/yr) and Mean Height SDS during the First Two Years of the Study

	12 Months (n=63)	24 Months (n=44)
HV	9.0 ± 3.6	7.5 ± 2.9
HV from Baseline	$+4.4 \pm 4.0^*$	$+3.0 \pm 3.6^*$
H SDS	-3.0 ± 1.7	-2.5 ± 1.5
H SDS from Baseline	$+0.7 \pm 0.7^*$	$+1.2 \pm 1.2^*$

*Statistically significant at the 0.05 level

Final Height

The change in the HA/BA (Height Age/Bone Age) ratio was also quantified during the 2 years of the study. The change in linear growth relative to the change in skeletal maturation, a measure of the preservation or loss of potential final height, was examined by estimating the HA/ BA ratio. A ratio of 1.0 indicates proportional gains in height and skeletal maturation and preservation of potential final height: a value above 1.0 suggests potential improvement in final height while a value below 1.0 indicates a loss of final height.

At baseline, for the 80 patients in which they could both be obtained, the mean HA was 5.1 and the mean BA was 5.7. For the 56 children with data at one year, the HA/ BA was 1.6 ± 2.2 , significantly greater than unity ($p=0.040$) and suggestive of improvement in predicted final height. For the cohort of children treated for 2 years, HA/ BA remained ≥ 1.0 during the second year, consistent with ongoing preservation of final adult height.

Analysis of Efficacy Subgroups

A secondary analysis, stratifying the study participants according to whether they had entered the study with compensated CRF, end stage renal disease (on dialysis) or whether they had undergone transplantation, was also performed. HV and H SDS rose in all three groups at one year of treatment. In the second year HV remained elevated above baseline levels in both the compensated and transplant groups and H SDS improved in all three groups.

Small for Gestational Age

A child born SGA is defined as a neonate whose birth weight or birth crown-heel length is at least 2 standard deviations (SD) below the mean (≤ -2 SD is equivalent to the 2.3 percentile) for the infant's gestational age, based on data derived from a reference population. Most children born SGA catch up in growth during their first years of life, but between 8% and 15% of all children born SGA do not.

Two randomized, controlled Phase III clinical studies GF 4001 (n=101) and GF 6283 (n=58) were designed to assess efficacy and safety of treatment with SAIZEN in children born SGA who did not catch up in growth during their first years of life. Their design and patient demographics are summarized in the following table:

Study Demographics and Trial Design

Table 19 - Summary of Patient Demographics for Clinical Trials in SGA

Study Number	Trial Design	Dosage (kg/day)	Route of Administration	Administration Scheme	Duration (years)	Study Subjects Number	Mean Age (range) (years)	Sex
GF 4001	Open, Randomised (SAIZEN group + No treatment control group)	0.067 mg (0.2 IU)	Subcutaneous injection	Continuous	Study: 4 Treatment: 3	101	4.5 (2-8)	M 51 F 49
GF 6283	Open, Randomised (SAIZEN group + No treatment control group)	0.067 mg (0.2 IU)	Subcutaneous injection	Intermittent	Study: 4 Treatment: 2	58	3.3 (2-5)	M 28 F 30

Study GF 4001 was initiated in 1990 and a dose of 0.2 IU/kg/day (corresponding to 0.067 mg/kg/day) was selected for use based on the results from the EMD Serono study GF 2773 and on published information (Albertsson-Wikland et al., 1989). The aim of the treatment was to induce catch-up growth and, because failure to catch up is thought to be related to a relative resistance to GH, a dose higher than that used to treat children with GH deficiency was administered. The study included a group treated from study initiation (Group T), and a control group (Group C) that did not receive any treatment during the first year but did receive SAIZEN in subsequent years. During the course of the study all children received SAIZEN treatment for three years and were followed for five years after treatment, with the exception of the one patient who withdrew before the start of the treatment phase. Study GF 6283 was initiated in 1993 and examined the same SAIZEN dose as study GF 4001. The 4-year study used two different regimens, continuous treatment for two years with follow-up for two additional years (Treatment-Treatment-Observation-Observation, TTOO) or discontinuous treatment

(treatment during the first and third study years and observation during the second and fourth study years, TOTO). The TOTO regimen was chosen to investigate whether discontinuous treatment would result in a better effect on catch-up growth than continuous treatment (TTOO) since it was well known that the effect of SAIZEN on height velocity is greater during the first year of treatment than during subsequent years.

The two pivotal studies used similar inclusion and exclusion criteria: birth weight less than the 10th percentile of the gestational age-related standards, height SD score (H-SDS) ≤ -3.0 for chronological age and sex, and height velocity (HV) less than +0.5 SD for chronological age and sex. At study entry, children were aged between 2 and 8 years in study GF 4001 and between 2 and 5 years in study GF 6283. An additional inclusion criterion in study GF 6283 was a parental height ≥ 1.48 m for women and ≥ 1.60 m for men.

Similar numbers of boys and girls were included in each of two treatment groups for Study GF 4001. In Study GF 6283 distribution of boys and girls in the two treatment groups were slightly skewed (the two treatment groups contained 28 and 30 patients with respectively 11 males/17 females in TTOO and 17 males/13 females in TOTO). Taken together, the children enrolled in the two studies were representative of short children born SGA. They had not demonstrated any catch-up growth and no child had GH deficiency as determined by standard GH stimulation tests. Mean parental heights were somewhat below the French population average in both studies.

Study Results

The greatest effect on growth was observed during the first year of r-hGH treatment. However, during continued treatment, HV-SDS values demonstrated that growth rate remained consistently above the mean for age and sex and that catch-up growth was maintained (see table below). Mean HV values were 7.62 and 7.46 cm/year during the second year of SAIZEN therapy in the T and C groups, respectively, and 6.62 and 6.86 cm/year during the third year in study GF 4001 (compared with 5.83 and 6.16 cm/year at baseline in the two treatment groups, respectively). The corresponding mean HV-SDS values were 1.97 and 2.05 after 2 years of SAIZEN therapy and 1.07 and 1.33 after 3 years in study GF 4001.

Mean HV during the second year in the group that received continuous treatment in study GF 6283 was 7.85 cm/year (compared with 7.40 cm/year at baseline). This corresponded to HV-SDS of 1.55 after 2 years of continuous treatment in study GF 6283.

As a result of the accelerated growth, the children's mean H-SDS CA improved to -1.67 and -1.73 after 2 years, and -1.43 and -1.41 after 3 years in study GF 4001, and to -1.55 after 2 years of continuous treatment in study GF 6283. Thus, 2 years of treatment with SAIZEN yielded a net height increase of 1.5, 1.7 and 2.0 SD (studies GF 4001 and GF 6283) and 3 years of continuous treatment yielded an increase in height of 1.9 SD (study GF 4001).

Table 20 - Growth Results in Patients Treated with SAIZEN from Studies GF4001 and GF6283
Values presented as group means (SD)

Study	Treatment Arm	Number Enrolled/ Completed	Height velocity SDS (based on chronological age)					Height SDS (based on chronological age)					Height SDS (based on bone age)				
			At start of	1 year	2 years	3 years	4 years	At start of	1 year	2 years	3 years	4 years	At start of	1 year	2 years	3 years	4 years
GF 4001	Group T, 0.067 mg/kg/day	50/44	-1.42 (1.23)	4.00 (1.68)	1.97 (1.56)	1.07 (1.59)		-3.34 (0.64)	-2.21 (0.81)	-1.67 (0.91)	-1.43 (0.93)		-1.04 (2.02)	-0.54 (1.73)	-0.43 (1.56)	-0.50 (1.56)	
	Group C, 0.067 mg/kg/day	50/39	-0.40 (1.11)	4.01 (2.11)	2.05 (1.68)	1.33 (1.12)		-3.24 (0.85)	-2.22 (0.89)	-1.73 (1.03)	-1.41 (1.11)		-1.37 (1.40)	-0.89 (1.47)	-1.01 (1.52)	-1.02 (1.37)	
GF 6283	Group TTOO, 0.067 mg/kg/day	28/25	-0.97 (1.01)	3.76 (1.31)	1.55 (1.82)	-2.28 (1.67) *	-1.82 (1.25) *	-3.55 (0.60)	-2.18 (0.62)	-1.55 (0.82)	-1.80 (0.78) *	-1.99 (0.81) *	-0.66 (2.35)	-0.78 (1.28)	-0.61 (1.24)	-0.88 (1.35) *	-1.21 (1.28) *
	Group TOTO, 0.067 mg/kg/day	30/28	-1.31 (1.44)	3.15 (1.50)	-1.78 (1.06) *	2.56 (1.69)	-2.41 (1.19) *	-3.43 (0.74)	-2.16 (0.88)	-2.35 (0.90) *	-1.68 (1.05)	-2.00 (1.01) *	-1.04 (2.79)	-0.93 (1.93)	-1.51 (1.63) *	-1.22 (1.73)	-1.68 (1.61) *

* Denotes values obtained during observation;

Growth and bone maturation

Bone maturation was followed during the pivotal studies with SAIZEN by assessing the ratio of bone age to chronological age (BA/CA), the ratio of the rate of change in bone age to the rate of change in chronological age ($\Delta\text{BA}/\Delta\text{CA}$), and H-SDS based on bone age.

BA/CA ratios increased during r-hGH treatment but remained below unity at all times. The increase of the BA/CA ratio is consistent with the increased height velocity during treatment and suggests that adequate catch-up growth is achieved during treatment with SAIZEN.

Effects of discontinuation of treatment

Study GF 6283 provide follow-up information for 2 years after treatment and study GF 4001 provides such information for 5 years after treatment cessation.

In study GF 4001, mean H-SDS CA values of the combined treatment groups were -1.38 at the end of treatment, and -1.70 , -1.83 , -1.96 , -2.16 and -2.23 after 1, 2, 3, 4 and 5 years of follow-up, respectively (pre-treatment values were -3.34 and -3.24 for the two groups, respectively). This means that over 5 years patients may have lost an average improvement in height of 0.85 SDS. Thus, the net effect of treatment for 3 years over a total of 8 years was a gain in height of approximately 1 SDS. These data indicate clearly that a substantial part of the effect on growth was lost following discontinuation of treatment.

Similarly, after 2 years of follow-up in study GF 6283, mean H-SDS CA was -1.99 (compared to a pre-treatment value of -3.55) and in study GF 6018 it was -1.4 (pre-treatment value of -2.6). Thus, the net effect of treatment for 2 years during a total of 4 years was a gain in height of approximately 1.2 to 1.5 SD.

Hence, discontinuation of treatment is accompanied by a substantial loss of the benefit on growth. The data support the notion that treatment should be given continuously until final height.

The impact of ethnic factors has not been evaluated in SGA clinical trials.

Adult Indications***Adult Growth Hormone Deficiency***Trial Design and Study Demographics

Table 21 – Summary of Patient Demographics for Clinical Trials in Adult GHD

Study #	Study design	Dosage, route of administration and duration	Study subjects (n)	Age (range)	Sex
GF 7364	Double-blind, placebo-controlled, randomised, phase III	(1) 0.005 mg/kg/day, 4 weeks, then 0.010 mg/kg/day, 5 months or (2) placebo, 6 months. (1) and (2) followed by 12-30 months open-label: 0.005 mg/kg/day, 4 weeks, then 0.010 mg/kg/day. Subcutaneous administration.	115	20.2-69.7	67, 48

The pathogenesis and disease history leading to adult GHD differs depending on whether the deficiency has existed since childhood or has been acquired during adult life.

Hypopituitarism acquired in adult life is often caused by pituitary or peripituitary tumours and/or their associated therapy. It is estimated that acquired hypopituitarism with GHD annually affects 10 people per million. The evolution of GHD after radiation therapy of a pituitary adenoma or craniopharyngioma has been investigated during ten years following radiotherapy. The time taken for pronounced GHD to develop was between one and four years and was dependent upon the GH status before radiotherapy. The safety and efficacy of SAIZEN replacement therapy in adults was evaluated in a pivotal study. This was a randomized, double-blind, placebo-controlled (DBPC) study involving 115 patients. Sixty patients received 0.005 mg/kg/day for one month and then 0.01 mg/kg/day for five months, and 55 patients received corresponding placebo, administered by daily subcutaneous injection. This was followed by 12 to 30 months open-label treatment for all patients.

Table 22 - Number of patients still on treatment by visit since study start

Month from start of study	Total # of patients	Patient still on treatment	
		r-hGH	Placebo ^o
DBPC start	115	60 (100%)	55 (100%)
DBPC end	115	53 (88%)	51 (93%)
Month 12	115	49 (82%)	48 (87%)
Month 18	115	34 (57%)	34 (62%)
Month 24 *	42	15 (68%)	13 (65%)
Month 30 *	42	11 (50%)	11 (55%)
Month 36 *	42	6 (27%)	6 (30%)

* Only 2 of the 6 sites scheduled treatment beyond 18 months.

^o Treatment with SAIZEN started in month 6.

2 patients had their last visit before one of the presented month and the last after that month.

Study Results

Lean body mass (DEXA):

There was a statistically significant increase ($p < 0.0001$) after 6 months of r-hGH treatment compared to placebo (r-hGH: baseline 49.1 ± 11.7 kg, N = 59, 6 months 49.6 ± 12.0 kg, N = 52); (placebo: baseline

53.7 ± 12.2 kg, N = 54, 6 months 53.9 ± 11.9 kg, N = 52). After adjusting for centre and baseline lean body mass, the estimated overall treatment difference was an increase of 2.21 kg (95% CI 1.27–3.15) compared to placebo. This increase was sustained throughout the 30 months of follow-up treatment. A subsidiary, subgroup analysis by gender shows a treatment difference of 2.91 ± 0.47 kg compared to placebo, in males, while the difference is 0.80 ± 0.70 kg in females.

Treadmill exercise test (Weber protocol):

There was a slightly greater increase in VO₂max in the r-hGH group compared to placebo, but the difference was not statistically significant (r-hGH: baseline 21.21 ± 7.71 mL/kg/min N = 36, 6-months 25.50 ± 7.78 mL/kg/min, N = 26; placebo: baseline 23.36 ± 6.98 mL/kg/min, N = 35, 6 months 26.47 ± 8.58 mL/kg/min, N = 31). No statistically significant differences were noted for anaerobic threshold (r-hGH: baseline 13.13 ± 3.80, N = 35, 6-months 16.29 ± 4.41 mL/kg/min, N = 26; placebo: baseline 14.69 ± 4.29, N = 34; 6 months 16.38 ± 6.00 mL/kg/min, N = 31). The changes observed during the follow-up phase VO₂max and anaerobic threshold paralleled study duration and were thus related to the time on study rather than to time on r-hGH treatment.

Other body composition results:

DEXA assessments demonstrated a statistically significant reduction of total fat mass (p<0.0001) in the r-hGH group compared to placebo (r-hGH: baseline 27.73 ± 10.72 kg, N = 59, 6 months 23.82 ± 9.65 kg, N = 52; placebo: baseline 28.90 ± 14.83 kg, N = 54, 6 months 29.12 ± 15.33 kg, N = 52). The fat mass remained relatively stable throughout 30 months of follow-up treatment. The changes in total bone mineral content or density were not statistically different between the treatment groups or during the follow-up.

Of the BIA assessments, only the change in total body water was subject to statistical analysis, and no significant difference was found between treatment groups. Anthropometry demonstrated no statistically significant between-group differences for skinfolds, waist/hip ratio or body weight and was maintained throughout the 30 month follow-up. The sum of circumferences decreased significantly in the r-hGH group relative to placebo (p<0.017). Body weight and BMI were stable throughout r-hGH treatment.

Bone turnover markers:

Intact parathyroid hormone decreased significantly in the r-hGH treated group (p=0.0071), whereas bone specific alkaline phosphatase, C-terminal propeptide, osteocalcin levels (p<0.0001 for each parameter) and urinary excretion of deoxyproline (p<0.001) rose significantly during r-hGH treatment compared to placebo. The changes were noted during the first 6 months of treatment and were followed by relatively little further change.

Perceived well-being:

The results from the Nottingham Health Profile (NHP) questionnaire demonstrated a statistically significant difference between the treatment groups during the DBPC phase for the domain Emotional Reactions (p<0.017). The domains for Social Isolation, Energy and Sleep were also found to be statistically significant for the first 6 months of treatment for the follow-up phase of the study and were maintained or further improved during this phase. No other significant differences were found in NHP or the General Well-Being Index.

Handgrip strength:

No statistically significant differences were found in the assessments of dominant or non-dominant hand-grip strength.

Mid-thigh cross-sectional MRI:

There were no statistically significant differences in the assessments of percentages of fat, muscle or bone (carried out in only 1 centre); however the decrease in fat, increase in muscle and stability of bone content echo the trends in body composition.

Cardiac function:

Two-dimensional echocardiography showed statistically significant differences between the treatment groups for ejection fraction percentage (increase in the r-hGH group, $p < 0.048$; r-hGH: baseline $54.90 \pm 11.21\%$, $N = 52$, 6 months $60.89 \pm 9.47\%$, $N = 48$; placebo: baseline $54.41 \pm 12.91\%$, $N = 50$, 6 months $57.30 \pm 8.61\%$, $N = 49$) and remained significantly higher than baseline throughout 30 months. Left ventricular posterior wall thickness also increased significantly during the first 6 months of treatment and remained stable and greater than baseline values although significantly only after 12 months.

16. Non-Clinical Toxicology

General toxicology

A series of toxicology studies including acute, subacute, subchronic and long-term studies were conducted with SAIZEN. The animal species used for these studies included mice, rats and monkeys.

Single Dose Studies

Six acute toxicity studies were conducted in rats, mice, and monkeys.

Table 23 – Acute Toxicity Studies

Species	Route of Administration	Dose
Rats and mice	Subcutaneous	13.3 mg/kg
Rats and mice	Intravenous	13.3 mg/kg
Rats	Subcutaneous	83.33 mg/kg
	Oral	
Rats	Oral	1.67, 3.33, 6.67, 13.3 mg/kg
Monkeys	Subcutaneous	1.67, 3.33, 6.67 mg/kg
Mice	Oral	1.67, 3.33, 6.67, 13.3 mg/kg

No effects were observed except for minor histological changes (vacuole in hepatocytes and a hyaline droplet in renal epithelium) in one female monkey of the 6.67 mg/kg dose group.

Repeated Dose Studies

Six studies were conducted: 2 four-week subacute studies (one in rats and the other in monkeys), two thirteen-week studies (one in rats and one in monkeys) and 2 fifty-two-week studies (rats and monkeys) by subcutaneous route. Pituitary-derived hGH (p-hGH) was administered in parallel in the 4-week tox study in monkeys and in the 13-week tox study in rats.

Table 24 - Four-Week Studies

	Rats (4 week study)	Monkeys (4 week study)
# Animals	15 animals/sex/group	28/sex
Dosage	Daily injections of 0, 0.067, 0.33, 1.6 and 3.33 mg/kg SAIZEN in 0.9% NaCl	Daily injections of 0, 0.067, 0.33 and 1.6 mg/kg of SAIZEN or 0.2 or 0.5 IU/kg p-hGH in 0.9% NaCl
Observations	Clinical signs, mortality, ophthalmoscopic examinations, body weight, laboratory analyses and post-mortem examination (autopsy, organ weight, histology)	
Results	The drug was well tolerated up to 3.33 mg/kg. Small number of slight hematological, biochemical and morphological changes, most notably in 1.6 and 3.33 mg/kg groups. These were for the most part reversible, and none appeared detrimental to the health of the animals. Local tolerability was satisfactory.	The drug was well tolerated up to 1.6 mg/kg. None of the observations appeared to be treatment-related and no clearly identifiable antibodies were observed.

Table 25 - Thirteen-week Studies

	Rats (13 week study)	Monkeys (13 week study)
# Animals	15/sex/group	12/sex/group
Dosage	Daily injections of 0, 0.067, 0.33 and 3.33 mg/kg SAIZEN, or 0.2 and 10 IU/kg p-hGH in 0.9% NaCl	Daily injections of 0.067, 0.33 and 1.67 mg/kg/day SAIZEN
Observations	Clinical signs, mortality, body weight, ophthalmoscopic examinations, laboratory analyses and post-mortem examination (autopsy, organ weight, histology)	
Results	No overt signs of toxicity, number of slight changes were observed due mainly to the biological activity of a heterologous hormone administered for a prolonged period of time.	Higher values of GOT, GPT, GGT and LAP found in males of the 1.67 mg group were the only drug-related changes.

LAP=leucine aminopeptidase; GGT=gamma-glutamyl transferase; GOT=glutamic-oxaloacetic transaminase; GPT=glutamic-pyruvic transaminase

Table 26 - Fifty-two-week Studies

	Rats (52 week study)	Monkeys (52 week study)
# Animals	100 males and 100 females	16 males and 16 females
Dose	0, 0.067, 0.2 and 0.6 mg/kg SAIZEN	0, 0.067, 0.2 and 0.6 mg/kg SAIZEN
Observations	Clinical signs, mortality, body weight, ophthalmoscopic examinations, laboratory analyses and post-mortem examination (autopsy, organ weight, histology)	
Results	No treatment-related deaths. Ten rats died from incidental causes or spontaneous incidental pathology. No treatment-related clinical signs. No clinical changes at the injection site. Body weight and food consumption unaffected. No treatment-related eye abnormalities were seen. None of the hematology, blood chemistry or	No treatment-related deaths. One male (0.2 mg/kg) died on day 245 (accidental trauma). No clinical or laboratory modifications attributable to the drug. No gross alternations at injection sites. No anti-hGH detected. Post-mortem examination showed no changes

	urinalysis value modifications were related to the dose levels. All rats developed high levels of antibodies by week 12, still present 8 weeks after end of study. Slight tendency towards increase of adrenal gland and spleen mean absolute weights without dose-correlation possible partial correlation with the slight increase in body weight. No drug-related modifications in gross pathology and histology.	attributable to the drug.
--	--	---------------------------

Genotoxicity

A series of mutagenicity studies, consisting of Ames Test, Gene Conversion Test in *S. cerevisiae*, Unscheduled DNA Synthesis in Cultured HeLa Cells, Chromosome Aberration in Human Lymphocytes Cultured In Vitro, Micronucleus Test, were conducted.

No mutagenic activity was observed with the drug for any of the mutagenicity tests listed above.

Carcinogenicity

No long-term animal studies have been performed to evaluate the carcinogenic potential of somatropin.

Reproductive and developmental toxicology

In three independent pre-clinical reproductive toxicology (fertility, teratology, and pre- and post-natal) studies involving rats, SAIZEN was administered by SC route at the dosages of 0, 0.033, 0.33 and 3.33 mg/kg/day. SAIZEN was administered SC to males during pre-mating and mating, and females during pre-mating, mating, gestation and lactation periods. In these studies, no compound related embryotoxic or teratogenic effects were observed in any experimental group. In addition, no interference with the F₀ and F₁ reproductive performance was found. SAIZEN administration did have the effect of increasing body weight and food and water intakes at the dosage of 3.33 mg/kg/day and to a lesser extent at the dosage of 0.33 mg/kg/day. Since the increase in body weight noted in parents, offspring and fetuses is related to the pharmacological activity of SAIZEN (r-hGH), the non-toxic effect dose is considered to be 3.33 mg/kg/day for both parents and their progeny.

In a reproductive (teratology) study in female rabbits, SAIZEN was administered by SC route at the dosages of 0, 0.033, 0.33 and 3.33 mg/kg/day for 13 days during the organogenesis period. Administration of SAIZEN during pregnancy did not induce embryotoxic or teratogenic effects, and the autopsy of females at the scheduled killing did not reveal any pathological changes. The no-effect level is considered to be 3.33 mg/kg/day for does and fetuses.

Patient Medication Information

READ THIS FOR SAFE AND EFFECTIVE USE OF YOUR MEDICINE

PrSAIZEN® 5 mg (Somatropin for injection)

This patient medication information is written for the person who will be taking SAIZEN. This may be you or a person you are caring for. Read this information carefully. Keep it as you may need to read it again.

This patient medication information is a summary. It will not tell you everything about this medication. If you have more questions about this medication or want more information about **SAIZEN**, talk to a healthcare professional.

Serious warnings and precautions box

- SAIZEN therapy should be carried out under the regular guidance of a doctor who is experienced in the diagnosis and management of patients with growth disorders.
- SAIZEN shall only be used if, once reconstituted, the solution is water-clear and does not contain particles (See *How to take SAIZEN*)
- Benzyl alcohol is a preservative in Bacteriostatic Water for Injection, USP, and it can be toxic to newborn babies*. When giving SAIZEN to newborns, mix it with sterile water for injection, USP. Use only one prepared dose per growth hormone vial and throw away any leftover liquid.
- There have been reports of deaths in children with Prader-Willi syndrome who were treated with growth hormone and had one or more of the following risk factors: severely obese, breathing problems or colds and lung infections.

*A newborn infant, or neonate, is a child under 28 days of age

What SAIZEN is used for:

- the long term treatment of patients with growth failure due to inadequate secretion of growth hormone.
- the treatment of short stature in girls with gonadal dysgenesis (Turner syndrome) when epiphyses are not closed.
- the treatment of growth failure in children due to Chronic Renal Failure.
- growth disturbance (current height Standard Deviation Score (SDS) < -2) in short children born small for gestational age (SGA) with a birth weight and/or length below -2 standard deviations (SD), who failed to show catch-up growth (Height Velocity SDS < 0 during the last year) by 2 years of age or later.
- replacement therapy in adult patients with acquired or idiopathic growth hormone deficiency (GHD) as diagnosed by a single dynamic test for growth hormone deficiency (peak GH ≤ 5 µg/L).

How SAIZEN works:

SAIZEN contains somatropin which is identical to growth hormone found naturally in humans but is made in laboratories.

SAIZEN provides an external supply of human growth hormone for those patients lacking the ability to

produce adequate amounts naturally. SAIZEN has many effects on growth and metabolism in patients undergoing therapy;

- stimulates linear growth (growth rate)
- measurable increase in growth (body length) from effects on cartilaginous growth areas of the long bones.
- cellular growth as demonstrated by an increase in the muscular, visceral and red cell mass
- effect on carbohydrate metabolism, including glucose tolerance and insulin levels
- effect on protein metabolism. SAIZEN is an anabolic agent that stimulates intracellular transport of amino acids, net retention of nitrogen and protein synthesis.
- lipid metabolism is also affected when intracellular lipolysis is stimulated, thus increasing the plasma concentration of free fatty acids and stimulating the oxidation of fatty acids.
- connective tissue metabolism is affected by stimulating the synthesis of chondroitin sulfate and collagen as well as the urinary excretion of hydroxyproline
- affects mineral metabolism by inducing the retention of phosphorus and potassium and to a lesser degree sodium.
- also increased are the intestinal absorption of calcium, renal tubular reabsorption of phosphorus with increased serum and inorganic phosphate.

The ingredients in SAIZEN are:

Medicinal ingredient(s): somatropin

Non-medicinal ingredients: Phosphoric acid, sodium hydroxide, and sucrose.

SAIZEN comes in the following dosage form(s):

SAIZEN is available in a 5 mg vial, with a vial of bacteriostatic water for injection (BWFI) diluent, containing benzyl alcohol as a preservative.

Do not use SAIZEN if:

SAIZEN should not be used in the following cases:

- Acute critical illness with complications following cardiac surgery, abdominal surgery, multiple trauma or acute respiratory failure. Clinical studies demonstrated that high doses of another somatropin were associated with a significantly increased morbidity and mortality in those patients.
- In patients with closed epiphyses, SAIZEN is ineffective for growth. Treatment with SAIZEN should be discontinued when the patient has reached satisfactory adult height, or the epiphyses are fused.
- In the presence of progression of an underlying intracranial tumour. An intracranial tumour should be inactive prior to instituting therapy, and SAIZEN should be discontinued if there is evidence of recurrent activity. Patients should be examined frequently for progression or recurrence of the underlying disease process.
- Patients known to be hypersensitive to somatropin and any of the excipients in powder for solution for injection or the diluent.
- Active neoplasia (either newly diagnosed or recurrent). Any pre-existing neoplasia should be inactive. SAIZEN therapy should be stopped if there is recurrent tumour growth.
- Proliferative or preproliferative diabetic retinopathy.

- In patients with Prader-Willi syndrome who are severely obese or have severe respiratory impairment. Unless patients with Prader-Willi syndrome also have a diagnosis of growth hormone deficiency, SAIZEN is not indicated for long-term treatment of pediatric patients who have growth failure due to genetically confirmed Prader-Willi syndrome.

SAIZEN treatment should be discontinued in critically ill patients.

SAIZEN is not recommended for use during pregnancy and lactation.

In children with chronic renal disease, treatment with somatropin must be discontinued at the time of renal transplantation.

To help avoid side effects and ensure proper use, talk to your healthcare professional before you take SAIZEN. Talk about any health conditions or problems you may have, including if you:

- feel shaky or light-headed due to low blood sugar levels shortly after taking SAIZEN. The feelings will quickly disappear. Your blood sugar levels may then rise above normal 2-4 hours after administration. Since treatment with growth hormone (GH) can alter how your body handles sugar, your levels will be tested regularly by a health care professional.
- are diabetic or a member of the family has diabetes, the doctor will monitor closely the treatment with SAIZEN and may change the treatment for diabetes. The doctor may additionally prescribe another hormone if you are found to have developed a lack of thyroid hormone.
- in the past had a condition affecting the brain e.g. a tumour, the doctor will examine you regularly to check that this has not come back again.
- suffer from a bad or recurrent headache, or from problems with eyesight and vomiting or feeling sick, contact your doctor immediately. Very rarely, a swelling of the brain may develop and the doctor may want to examine your eyes to look for any sign of this. In this case it may be necessary to stop the growth hormone treatment, although the treatment may be re-started at a later date. If the symptoms of brain swelling recur, treatment with SAIZEN should be discontinued.

Other warnings you should know about:

- SAIZEN therapy should be carried out under the regular guidance of a doctor who is experienced in the diagnosis and management of patients with growth hormone deficiency.
- When the medicine is injected into the same place over a long time, it can cause damage to this area. It is therefore important to keep changing the injection site. The doctor or nurse can speak to you about which parts of the body should be used.
- Some children with growth hormone deficiency have developed leukemia, whether or not they have received treatment with growth hormone, and might be at a slightly higher risk of developing leukemia than non-growth hormone deficient children. No cause and effect relationship with growth hormone treatment have been proven.
- Hip problems may occur more commonly in children with hormone or kidney problems. If the patient has chronic renal failure (which can occur when kidneys are damaged) he or she should be examined periodically for evidence of bone disease. It is uncertain whether the bone disease in children with hormone or kidney problems is affected by growth hormone therapy. X-rays of the hip should be obtained prior to initiating therapy. If the patient develops a limp or complains of hip or knee pain while being treated with SAIZEN, notify your doctor. He/she

should make sure that these symptoms are not caused by a slip of the femur head (separation of the epiphysis), before continuing treatment with SAIZEN.

- Rarely, SAIZEN may cause an inflammation of the pancreas, which causes severe pain in the abdomen and back. Please consider this especially if your child suffers from stomach-ache and contact your doctor.
- SAIZEN may affect how your thyroid works. Your doctor may test your blood for levels of thyroid hormones and prescribe another hormone if you (or your child) are found to have developed a lack of thyroid hormone.
- An increase in sideways curvature of the spine (scoliosis) may progress in any child during rapid growth. During treatment with SAIZEN, your doctor will check you (or your child) for signs of scoliosis.
- In children with chronic renal failure, treatment should be discontinued at the time of renal transplant.
- Small for Gestational Age (SGA) patients: SGA means small for gestational age. Gestational age is the time a baby is in its mother's womb. SGA refers to a baby who is smaller than most babies of the same gestational age. Patients who are small for their gestational age have a greater chance of developing diabetes. It is important to have your fasting insulin and blood glucose levels checked before and during treatment with SAIZEN.
- Patients with Turner syndrome treated with SAIZEN should be closely examined, especially if bone pain occurs.
- SAIZEN may cause fluid retention in adult patients. This may appear as swelling and joint or muscle pain. If you feel these symptoms tell your doctor who may decide to adjust your dose of SAIZEN.
- Notify your doctor if the patient develops a limp, difficulty walking, or knee or hip pain. The doctor should make sure that these symptoms are not caused by a displacement of the femoral head in relation to the joint head (epiphysiolysis) or by a circulatory disorder of the femoral head (avascular necrosis [osteonecrosis] or Legg-Calve-Perthes disease).

Tell your healthcare professional about all the medicines you take, including any drugs, vitamins, minerals, natural supplements or alternative medicines.

The following may interact with SAIZEN:

- It is usually safe to take other medicines. However, if the patient is taking CORTICOSTEROIDS, it is necessary to tell the doctor or nurse. These medicines are used to treat several illnesses including asthma, allergies, kidney rejection and rheumatoid arthritis. These medicines might stop your growth treatment from working.
- Thyroid hormones
- Glucocorticoids
- Antidiabetic medications
- Medications metabolized by cytochrome P450 3A4 (e.g., corticosteroids, sex steroids, anticonvulsants, cyclosporine and others)
- Oral estrogens

How to take SAIZEN:

Preparing SAIZEN for Administration

Here are the things you will need before you inject SAIZEN:

- 3 alcohol swabs
- Cotton swab

- 3cc syringe & 23 gauge needle for mixing
- BD insulin syringe for injection
- 1 vial of SAIZEN
- Diluent vial (You need this sterile liquid - the diluent - to dissolve the SAIZEN powder and make it injectable.)
- Syringe safety disposal container for used vials and needles

Always use unopened, sterile needles and syringes and keep the needles capped until needed.

TIP: Your doctor or nurse will explain how much diluent to add to the vial of SAIZEN and how much SAIZEN to inject.

Getting Ready to use SAIZEN

1. Begin by choosing a clean flat surface (like a kitchen or bathroom counter).
2. Wash your hands thoroughly with soap and water. This helps prevent infection.
3. Check the expiration date of your SAIZEN.

Drawing Up the Diluent

1. Carefully twist the needle cover off the long needle syringe.
2. Pull out the plunger to the amount recommended by your doctor or nurse. This brings air into the syringe.
3. Remove the flip-off cap from the diluent vial and discard. Wipe the rubber stopper of the vial with an alcohol swab.
4. Hold the vial firmly on the countertop. Put the needle into the stopper of the SAIZEN diluent vial. Push the plunger of the syringe and inject the air into the vial.
5. Turn the vial upside down. Make sure the needle tip stays in the liquid. Pull back on the plunger until the marks on the barrel of the syringe show that the amount of the diluent suggested by your doctor or nurse has been drawn out.
6. If air bubbles appear in the syringe, gently push the plunger into the syringe to send the air into the vial. You may have to tap the syringe lightly so you can push the bubbles out. Draw up more diluent, if needed, until you have the amount your doctor has prescribed.
7. Pull out the needle from the diluent.

Mixing SAIZEN

1. Remove the flip-off cap from the SAIZEN vial and discard. Wipe the rubber stopper of the vial with an alcohol swab.
2. With the same syringe, put the long needle into the stopper of the SAIZEN vial. Gently place the needle tip against the vial wall. Slowly inject the diluent, aiming the stream of diluent at the glass wall of the vial. DO NOT AIM THE STREAM AT THE WHITE POWDER at the bottom of the vial.
3. Take out the needle and throw it away in the safety container.
4. Gently swirl (don't shake) the vial until the powder is completely dissolved. The SAIZEN mixture should be clear. If it stays hazy, cloudy or has pieces floating in it after mixing, do not use it.

TIP: If SAIZEN becomes cloudy after mixing, return it to your pharmacist or nurse.

Preparing SAIZEN For Injection

1. Re-wipe the rubber stopper of the SAIZEN vial with an alcohol swab.
2. Pick up the insulin syringe with the short needle and carefully take off the needle cover.


3. Pull out the plunger to the amount recommended by your doctor or nurse. This brings air into the syringe.
4. Slowly insert the needle straight through the center of the rubber stopper of the vial of newly mixed SAIZEN. Gently push the plunger to inject air into the vial.
5. Turn the vial upside down with the syringe needle still in it, holding the vial in one hand. Be sure the tip of the needle is in the solution. Using your other hand, slowly pull back on the plunger until the amount of SAIZEN prescribed is in the syringe.
6. Remove the needle from the vial.
7. Hold the syringe straight up and tap gently. Put the plastic needle guard back until injection time. The injection should be given as soon as possible after filling the syringe. Do not store SAIZEN in the syringe.

TIP: Be careful not to touch the uncapped needle with your fingers - or let the needle touch anything.

Picking an Injection Site

You should pick a different site to inject each day, rotating through arms, legs and abdomen. The buttocks can be used, as well (see Injection Site Diagram). Using a site too often can lead to infection, irritation, or fat loss.

Injection Site Diagram



TIP: Many parents find it's a good idea to practice giving injections to each other, so they know what it feels like and to understand their child's reaction better. You can use a small amount of normal saline from the doctor's office.

Injecting SAIZEN

1. Clean the skin at the injection site with an alcohol swab using a circular motion.

TIP: Let the skin dry after cleaning it with alcohol. This helps reduce stinging.

2. Remove the cap from the needle and, using the hand with which you write, pick up the syringe and hold it like a pencil.
3. Pinch up a generous fold of skin and hold it while quickly inserting the needle all the way in at a 90 degree angle to the skin. With your index finger, push the plunger in to inject the medication. Take as much time as you need to inject all the solution. You may wish to count to 5.

TIP: When inserting the needle, you need very little force, but quick action.

4. As you release the skin from your grip, withdraw the needle at the same angle at which it was inserted. Place the cotton swab on the injection site and apply a gentle pressure.

5. Do not put the needle back in the needle guard. Carefully throw away the needle guard and all used needles and syringes in the safety container after a single use.

TIP: NEVER reuse a needle.

Disposal containers must be made of thick, puncture-proof plastic with a lid that fits firmly, such as an empty pop bottle. Containers may be returned to the clinic for disposal or you may wish to contact your pharmacy for further information regarding the safe disposal of used syringes.

Things to remember

1. Make injections routine - give the injection at the same time each evening before bedtime.
2. Store vials at room temperature (not to exceed 25°C). However, once reconstituted with bacteriostatic diluent they must be stored in the fridge and may be used for up to 14 days.
3. Check the expiration date.
4. Do not use if it turns cloudy, lumpy or discoloured.
5. Make certain that the dosage is equal to the amount prescribed.
6. Rotate your injection sites each time, as discussed with your nurse.
7. Refrigerate any unused solution. Multi-dose only with SAIZEN reconstituted with bacteriostatic WFI (BWFI).
8. If you are unsure about the mixing of the medication or if you are having difficulty with the injection procedure, contact your nurse or doctor.

Usual dose:

SAIZEN should be injected preferably in the evening.

The dosage and administration schedule of SAIZEN will be adapted to the patient's body weight by the doctor according to the following scheme:

- Growth failure due to inadequate endogenous growth hormone secretion: It is recommended that SAIZEN be administered subcutaneously at a dose of 0.2 mg/kg body weight per week. The dosage can be increased to 0.27 mg/kg per week if there is insufficient response to treatment. For intramuscular injection, the weekly dose should be divided into 3 single injections (0.067 mg/kg). Use a needle which is at least 2.5 cm long to ensure the injection reaches the muscle.
- Growth failure in girls due to gonadal dysgenesis (Turner syndrome): It is recommended that SAIZEN be administered subcutaneously at a dose of 0.375 mg/kg body weight per week (optimal dosing 0.32 – 0.375 mg/kg/week). Concomitant therapy with non-androgenic anabolic steroids in patients with Turner syndrome can enhance the growth response.
- Growth failure in children with Chronic Renal Failure: It is recommended that SAIZEN be administered subcutaneously at a dose of 0.35 mg/kg body weight per week.
- Growth disturbance in short children born small for gestational age (SGA): It is recommended that SAIZEN be administered subcutaneously at a dose of 0.47 mg/kg body weight/week.
- Adult Growth Hormone Deficiency: It is recommended that SAIZEN be administered subcutaneously at a dose of 0.005 mg/kg/day at the start of therapy. This dose may be increased after 4 weeks to 0.01 mg/kg/day if well tolerated. The minimum effective dose should be used and dose requirements may decline with age.

Overdose:

If a dose is missed or too much SAIZEN injected it is important to tell your doctor as it may be necessary to change slightly the dose to make up for this. Injecting too much can lead to changes in blood sugar levels which could mean that the patient will feel shaky and light-headed. If this happens contact your doctor as soon as possible. If too much were taken over a period of time, this could cause an excessive

growth of some bones to occur, particularly the hands, feet and jaw. Also, taking too much SAIZEN for a long time can cause your body to keep extra fluids.

There have been no reports of the effects of acute overdose.

If you think you, or a person you are caring for, have taken too much SAIZEN, contact a healthcare professional, hospital emergency department, regional poison control centre or Health Canada's toll-free number, 1-844 POISON-X (1-844-764-7669) immediately, even if there are no signs or symptoms.

Missed dose:

If you missed a dose of this medication, you do not need to make up the missed dose. Skip the missed dose and continue with your next scheduled dose. Do not take two doses at the same time.

Possible side effects from using SAIZEN:

These are not all the possible side effects you may have when taking SAIZEN. If you experience any side effects not listed here, tell your healthcare professional.

As with any medicine, side effects can sometimes occur, however most people have no problems with their prescribed SAIZEN. Sometimes, however, redness and itching may appear at the injection site. If this appears to be particularly troublesome, you should discuss this with your doctor.

SAIZEN may bring about insulin resistance. Insulin resistance means your body cannot make good use of the insulin it produces. This causes higher levels of glucose in your blood. Your doctor will need to check your blood glucose on a regular basis. It is also important to check blood glucose if you have diabetes or a family history of diabetes.

Intracranial hypertension is pressure within the skull that is too high. This may be a complication of SAIZEN (growth hormone therapy). Call your doctor if you have a headache that does not go away or goes away and comes back, problems with your vision, a sick feeling in your stomach (nausea) or vomiting.

Very rarely a patient could develop antibodies to somatropin.

These are usually not associated with any side effects and do not usually interfere with growth.

If the patient shows an unexplained limp, please contact your doctor or nurse.

If the patient suffers from these or any other unwanted effects, please inform your doctor or nurse.

This is not a complete list of side effects. If the patient experiences any unusual symptom or side effects, you should report them to the doctor immediately. It is also wise to discuss the possibility of side effects with the doctor before beginning treatment.

Reporting side effects

You can report any suspected side effects associated with the use of health products to Health Canada by:

- Visiting the Web page on Adverse Reaction Reporting (canada.ca/drug-device-reporting) for information on how to report online, by mail or by fax; or
- Calling toll-free at 1-866-234-2345.

NOTE: Contact your healthcare professional if you need information about how to manage your side effects. The Canada Vigilance Program does not provide medical advice.

Storage:

Lyophilized product:

Store SAIZEN lyophilized product at room temperature (not to exceed 25°C).

Do not use SAIZEN after the expiry date shown on label.

Reconstituted product:

SAIZEN 5 mg/vial:

When reconstituted with 1 mL to 3.0 mL Bacteriostatic Water for Injection, USP, store the reconstituted solution at 2-8 °C for up to 14 days.

When reconstituted with Water for Injection, USP, the reconstituted solution should be administered immediately (within 3 hours). Any unused solution should be discarded.

Keep out of reach and sight of children.

If you want more information about SAIZEN:

- Talk to your healthcare professional
- Find the full product monograph that is prepared for healthcare professionals and includes the Patient Medication Information by visiting the Health Canada Drug Product Database website (<https://www.canada.ca/en/health-canada/services/drugs-health-products/drug-products/drug-product-database.html>); the manufacturer's website [<https://medinfo.emdserono.ca/en>]; or by calling MOMENTUM patient services program at 1-877-724-9361.

This leaflet was prepared by EMD Serono, A Division of EMD Inc., Canada, a business of Merck KGaA, Darmstadt, Germany

Date of Authorization: 2026-08-28

® Registered trademark of Merck KGaA, Darmstadt, Germany or its affiliates

Patient Medication Information

READ THIS FOR SAFE AND EFFECTIVE USE OF YOUR MEDICINE

PrSAIZEN® 6 mg (5.83 mg/mL), 12 mg (8 mg/mL), 20 mg (8 mg/mL) cartridges
(Somatropin, Solution for injection in a cartridge)

This patient medication information is written for the person who will be taking **SAIZEN**. This may be you or a person you are caring for. Read this information carefully. Keep it as you may need to read it again.

This patient medication information is a summary. It will not tell you everything about this medication. If you have more questions about this medication or want more information about **SAIZEN**, talk to a healthcare professional.

Serious warnings and precautions box

- SAIZEN therapy should be carried out under the regular guidance of a doctor who is experienced in the diagnosis and management of patients with growth disorders.
- SAIZEN shall only be used if, once reconstituted, the solution is water-clear and does not contain particles (See How to take SAIZEN)
- Benzyl alcohol is a preservative in Bacteriostatic Water for Injection, USP, and it can be toxic to newborn babies*. When giving SAIZEN to newborns, mix it with sterile water for injection, USP. Use only one prepared dose per growth hormone vial and throw away any leftover liquid.
- There have been reports of deaths in children with Prader-Willi syndrome who were treated with growth hormone and had one or more of the following risk factors: severely obese, breathing problems or colds and lung infections.

*A newborn infant, or neonate, is a child under 28 days of age

What SAIZEN is used for:

SAIZEN is indicated for

- the long term treatment of patients with growth failure due to inadequate secretion of growth hormone.
- the treatment of short stature in girls with gonadal dysgenesis (Turner syndrome) when epiphyses are not closed.
- the treatment of growth failure in children due to Chronic Renal Failure.
- growth disturbance (current height Standard Deviation Score (SDS) < -2) in short children born small for gestational age (SGA) with a birth weight and/or length below -2 standard deviations (SD), who failed to show catch-up growth (Height Velocity SDS < 0 during the last year) by 2 years of age or later.
- replacement therapy in adult patients with acquired or idiopathic growth hormone deficiency (GHD) as diagnosed by a single dynamic test for growth hormone deficiency (peak GH ≤ 5 µg/L).

How SAIZEN works:

SAIZEN contains somatropin which is identical to growth hormone found naturally in humans but is made in laboratories.

SAIZEN provides an external supply of human growth hormone for those patients lacking the ability to produce adequate amounts naturally. SAIZEN has many effects on growth and metabolism in patients undergoing therapy;

- stimulates linear growth (growth rate)
- measurable increase in growth (body length) from effects on cartilaginous growth areas of the long bones.
- cellular growth as demonstrated by an increase in the muscular, visceral and red cell mass
- effect on carbohydrate metabolism, including glucose tolerance and insulin levels
- effect on protein metabolism. SAIZEN is an anabolic agent that stimulates intracellular transport of amino acids, net retention of nitrogen and protein synthesis.
- lipid metabolism is also affected when intracellular lipolysis is stimulated, thus increasing the plasma concentration of free fatty acids and stimulating the oxidation of fatty acids.
- connective tissue metabolism is affected by stimulating the synthesis of chondroitin sulfate and collagen as well as the urinary excretion of hydroxyproline
- affects mineral metabolism by inducing the retention of phosphorus and potassium and to a lesser degree sodium.
- also increased are the intestinal absorption of calcium, renal tubular reabsorption of phosphorus with increased serum and inorganic phosphate.

The ingredients in SAIZEN are:

Medicinal ingredient(s): somatropin

Non-medicinal ingredients: Citric acid, phenol, Poloxamer 188, and sucrose.

SAIZEN comes in the following dosage form(s):

SAIZEN comes as a pre-filled cartridge. There are three presentations which are available:

1. SAIZEN 6 mg with a nominal volume of 1.03 mL of solution. Final concentration that will be injected is 5.83 mg/mL.
2. SAIZEN 12 mg with a nominal volume of 1.50 mL of solution. Final concentration that will be injected is 8 mg/mL.
3. SAIZEN 20 mg with a nominal volume of 2.50 mL of solution. Final concentration that will be injected is 8 mg/mL.

The SAIZEN cartridges are colour coded: SAIZEN 6 mg (5.83 mg/mL) in blue, 12 mg (8 mg/mL) in red, and 20 mg (8 mg/mL) in yellow.

Do not use SAIZEN if:

SAIZEN should not be used in the following cases:

- Acute critical illness with complications following cardiac surgery, abdominal surgery, multiple trauma or acute respiratory failure. Clinical studies demonstrated that high doses of another somatropin, were associated with a significantly increased morbidity and mortality in those patients.
- In patients with closed epiphyses, SAIZEN is ineffective for growth. Treatment with SAIZEN should be discontinued when the patient has reached satisfactory adult height, or the epiphyses are fused.
- In the presence of progression of an underlying intracranial tumour. An intracranial tumour should be inactive prior to instituting therapy, and SAIZEN should be discontinued if there is evidence of

recurrent activity. Patients should be examined frequently for progression or recurrence of the underlying disease process.

- Patients known to be hypersensitive to somatropin and any of the excipients in powder for solution for injection or the diluent.
- Active neoplasia (either newly diagnosed or recurrent). Any pre-existing neoplasia should be inactive. SAIZEN therapy should be stopped if there is recurrent tumour growth.
- Proliferative or preproliferative diabetic retinopathy.
- In patients with Prader-Willi syndrome who are severely obese or have severe respiratory impairment. Unless patients with Prader-Willi syndrome also have a diagnosis of growth hormone deficiency, SAIZEN is not indicated for long-term treatment of pediatric patients who have growth failure due to genetically confirmed Prader-Willi syndrome.

SAIZEN treatment should be discontinued in critically ill patients.

SAIZEN is not recommended for use during pregnancy and lactation.

In children with chronic renal disease, treatment with somatropin must be discontinued at the time of renal transplantation.

To help avoid side effects and ensure proper use, talk to your healthcare professional before you take SAIZEN. Talk about any health conditions or problems you may have, including if you:

- feel shaky or light-headed due to low blood sugar levels shortly after taking SAIZEN. The feelings will quickly disappear. Your blood sugar levels may then rise above normal 2-4 hours after administration. Since treatment with growth hormone (GH) can alter how your body handles sugar, your levels will be tested regularly by a health care professional.
- are diabetic or a member of the family has diabetes, the doctor will monitor closely the treatment with SAIZEN and may change the treatment for diabetes. The doctor may additionally prescribe another hormone if you are found to have developed a lack of thyroid hormone.
- If in the past you have had a condition affecting the brain e.g. a tumour, the doctor will examine the patient regularly to check that this has not come back again.
- suffer from a bad or recurrent headache, or from problems with eyesight and vomiting or feeling sick, contact your doctor immediately. Very rarely, a swelling of the brain may develop and the doctor may want to examine your eyes to look for any sign of this. In this case it may be necessary to stop the growth hormone treatment, although the treatment may be re-started at a later date. If the symptoms of brain swelling recur, treatment with SAIZEN should be discontinued.

Other warnings you should know about:

- SAIZEN therapy should be carried out under the regular guidance of a doctor who is experienced in the diagnosis and management of patients with growth hormone deficiency.
- When the medicine is injected into the same place over a long time, it can cause damage to this area. It is therefore important to keep changing the injection site. The doctor or nurse can speak to you about which parts of the body should be used.
- Some children with growth hormone deficiency have developed leukemia, whether or not they have received treatment with growth hormone, and might be at a slightly higher risk of

developing leukemia than non-growth hormone deficient children. No cause and effect relationship with growth hormone treatment have been proven.

- Hip problems may occur more commonly in children with hormone or kidney problems. If the patient has chronic renal failure (which can occur when kidneys are damaged) he or she should be examined periodically for evidence of bone disease. It is uncertain whether the bone disease in children with hormone or kidney problems is affected by growth hormone therapy. X-rays of the hip should be obtained prior to initiating therapy. If the patient develops a limp or complains of hip or knee pain while being treated with SAIZEN, notify your doctor. He/she should make sure that these symptoms are not caused by a slip of the femur head (separation of the epiphysis), before continuing treatment with SAIZEN.
- Rarely, SAIZEN may cause an inflammation of the pancreas, which causes severe pain in the abdomen and back. Please consider this especially if your child suffers from stomach-ache and contact your doctor.
- SAIZEN may affect how your thyroid works. Your doctor may test your blood for levels of thyroid hormones and prescribe another hormone if you (or your child) are found to have developed a lack of thyroid hormone.
- An increase in sideways curvature of the spine (scoliosis) may progress in any child during rapid growth. During treatment with SAIZEN, your doctor will check you (or your child) for signs of scoliosis.
- In children with chronic renal failure, treatment should be discontinued at the time of renal transplant.
- Small for Gestational Age (SGA) patients: SGA means small for gestational age. Gestational age is the time a baby is in its mother's womb. SGA refers to a baby who is smaller than most babies of the same gestational age. Patients who are small for their gestational age have a greater chance of developing diabetes. It is important to have your fasting insulin and blood glucose levels checked before and during treatment with SAIZEN.
- Patients with Turner syndrome treated with SAIZEN should be closely examined, especially if bone pain occurs.
- SAIZEN may cause fluid retention in adult patients. This may appear as swelling and joint or muscle pain. If you feel these symptoms tell your doctor who may decide to adjust your dose of SAIZEN.
- Notify your doctor if the patient develops a limp, difficulty walking, or knee or hip pain. The doctor should make sure that these symptoms are not caused by a displacement of the femoral head in relation to the joint head (epiphysiolysis) or by a circulatory disorder of the femoral head (avascular necrosis [osteonecrosis] or Legg-Calve-Perthes disease).

Tell your healthcare professional about all the medicines you take, including any drugs, vitamins, minerals, natural supplements or alternative medicines.

The following may interact with SAIZEN:

- It is usually safe to take other medicines. However, if the patient is taking CORTICOSTEROIDS, it is necessary to tell the doctor or nurse. These medicines are used to treat several illnesses including asthma, allergies, kidney rejection and rheumatoid arthritis. These medicines might stop your growth treatment from working.
- Thyroid hormones
- Glucocorticoids
- Antidiabetic medications

- Medications metabolized by cytochrome P450 3A4 (e.g., corticosteroids, sex steroids, anticonvulsants, cyclosporine and others)
- Oral estrogens

How to take SAIZEN:

Preparing SAIZEN for Administration

Place all elements needed for the injection of the medication on a clean surface and wash your hands with soap and water. The cartridge containing the medication of SAIZEN is ready to be used for administration with the easypod® electromechanical auto-injector or the Aluetta® pen injector. The solution should be clear to slightly opalescent with no particles. If the solution contains particles, it must not be injected.

The Aluetta pen injectors are available in several presentations. Each Aluetta pen injector is colour coded and must only be used with the matching colour coded SAIZEN cartridge to give the correct dose:


- The Aluetta pen injector 6 (blue) must be used with the SAIZEN 6 mg (5.83 mg/mL) cartridge (blue).
- The Aluetta pen injector 12 (red) must be used with the SAIZEN 12 mg (8 mg/mL) cartridge (red).
- The Aluetta pen injector 20 (yellow) must be used with the SAIZEN 20 mg (8 mg/mL) cartridge (yellow).

For instructions on how to load the cartridge into the easypod electromechanical auto-injector or the Aluetta pen injector, please carefully read the corresponding instruction manual provided with the device.

Picking an Injection Site

You should pick a different site to inject each day, rotating through arms, legs and abdomen. The buttocks can be used, as well (see Injection Site Diagram). Using a site too often can lead to infection, irritation, or fat loss.

Injection Site Diagram



Usual dose:

SAIZEN should be injected preferably in the evening.

The dosage and administration schedule of SAIZEN will be adapted to the patient's body weight by the

doctor according to the following scheme:

- Growth failure due to inadequate endogenous growth hormone secretion: It is recommended that SAIZEN be administered subcutaneously at a dose of 0.2 mg/kg body weight per week. The dosage can be increased to 0.27 mg/kg per week if there is insufficient response to treatment.
- Growth failure in girls due to gonadal dysgenesis (Turner syndrome): It is recommended that SAIZEN be administered subcutaneously at a dose of 0.375 mg/kg body weight per week (optimal dosing 0.32 – 0.375 mg/kg/week). Concomitant therapy with non-androgenic anabolic steroids in patients with Turner syndrome can enhance the growth response.
- Growth failure in children with Chronic Renal Failure: It is recommended that SAIZEN be administered subcutaneously at a dose of 0.35 mg/kg body weight per week.
- Growth disturbance in short children born small for gestational age (SGA): It is recommended that SAIZEN be administered subcutaneously at a dose of 0.47 mg/kg body weight/week.
- Adult Growth Hormone Deficiency: It is recommended that SAIZEN be administered subcutaneously at a dose of 0.005 mg/kg/day at the start of therapy. This dose may be increased after 4 weeks to 0.01 mg/kg/day if well tolerated. The minimum effective dose should be used and dose requirements may decline with age.

Overdose:

If a dose is missed or too much SAIZEN injected it is important to tell your doctor as it may be necessary to change slightly the dose to make up for this. Injecting too much can lead to changes in blood sugar levels which could mean that the patient will feel shaky and light-headed. If this happens contact your doctor as soon as possible. If too much were taken over a period in time, this could cause an excessive growth of some bones to occur, particularly the hands, feet and jaw. Also, taking too much somatropin for a long time can cause your body to keep extra fluids.

There have been no reports of the effects of acute overdose.

If you think you, or a person you are caring for, have taken too much SAIZEN, contact a healthcare professional, hospital emergency department, regional poison control centre or Health Canada's toll-free number, 1-844 POISON-X (1-844-764-7669) immediately, even if there are no signs or symptoms.

Missed dose:

If you missed a dose of this medication, you do not need to make up the missed dose. Skip the missed dose and continue with your next scheduled dose. Do not take two doses at the same time.

Possible side effects from using SAIZEN:

These are not all the possible side effects you may have when taking SAIZEN. If you experience any side effects not listed here, tell your healthcare professional.

As with any medicine, side effects can sometimes occur, however most people have no problems with their prescribed SAIZEN. Sometimes, however, redness and itching may appear at the injection site. If this appears to be particularly troublesome, you should discuss this with your doctor.

SAIZEN may bring about insulin resistance. Insulin resistance means your body cannot make good use of the insulin it produces. This causes higher levels of glucose in your blood. Your doctor will need to check your blood glucose on a regular basis. It is also important to check blood glucose if you have diabetes or a family history of diabetes.

Intracranial hypertension is pressure within the skull that is too high. This may be a complication of SAIZEN (growth hormone therapy). Call your doctor if you have a headache that does not go away or goes away and comes back, problems with your vision, a sick feeling in your stomach (nausea) or vomiting.

Very rarely a patient could develop antibodies to somatropin. These are usually not associated with any side effects and do not usually interfere with growth.

If the patient shows an unexplained limp, please contact your doctor or nurse.

If the patient suffers from these or any other unwanted effects, please inform your doctor or nurse. This is not a complete list of side effects. If the patient experiences any unusual symptoms or side effects, you should report them to the doctor immediately. It is also wise to discuss the possibility of side effects with the doctor before beginning treatment.

Reporting side effects

You can report any suspected side effects associated with the use of health products to Health Canada by:

- Visiting the Web page on Adverse Reaction Reporting (canada.ca/drug-device-reporting) for information on how to report online, by mail or by fax; or
- Calling toll-free at 1-866-234-2345.

NOTE: Contact your healthcare professional if you need information about how to manage your side effects. The Canada Vigilance Program does not provide medical advice.

Storage:

Do not use SAIZEN after the expiry date, which is stated on the cartridge.

Do not freeze. Store cartridge(s) in a refrigerator (2-8°C) in the original package.

When using the easypod electromechanical auto-injector or the Aluetta pen injector, the cartridge is kept in the device and the device must be stored in the refrigerator.

Do not freeze. After first injection, the cartridges must be stored at 2-8°C for a period of up to 28 days, of which no more than 7 days may be outside of the refrigerator at room temperature (not to exceed 25°C).

Cartridges must be discarded after 28 days, or if the total period outside of the refrigerator exceeds 7 days.

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required.

Keep out of reach and sight of children.

If you want more information about SAIZEN:

- Talk to your healthcare professional
- Find the full product monograph that is prepared for healthcare professionals and includes the Patient Medication Information by visiting the Health Canada Drug Product Database website (<https://www.canada.ca/en/health-canada/services/drugs-health-products/drug-products/drug->

product-database.html); the manufacturer's website [<https://medinfo.emdserono.ca/en>]; or by calling MOMENTUM patient services program at 1-877-724-9361.

This leaflet was prepared by EMD Serono, A Division of EMD Inc., Canada, a business of Merck KGaA, Darmstadt, Germany

Date of Authorization: 2026-08-28

® Registered trademark of Merck KGaA, Darmstadt, Germany or its affiliates